Title: Berberine is a novel mitochondrial calcium uniporter (MCU) inhibitor that disrupts MCU-EMRE assembly

Authors:

Haixin Zhao[1, 3, 5], Siqi Chen[1, 2, 5], Nian Cao[1, 4, 5], Wenjun Wu[1, 5], Guangqin Liu[1,2], Jun Gao[1], Jiayi Chen[1], Ting Li[1], Dingyi Lu[1], Lingmin Zeng[1], Haizhen Zhu[1], Weina Zhang[1], Qing Xia[1], Teng Li[1], Tao Zhou[1], Xue-Min Zhang[1,2], Ai-Ling Li[1, 2, *], Xin Pan[1, 2, *]

Affiliations:

1. Nanhu Laboratory, National Center of Biomedical Analysis, 27 Tai-Ping Road, Beijing 100039, China
2. School of Basic Medical Sciences, Fudan University, Shanghai 200032, China
3. State Key Laboratory of Experimental Haematology, Fifth Medical Center of Chinese PLA General Hospital, Beijing 100071, China
4. Department of Cardiology, the Sixth Medical Centre, Chinese PLA General Hospital, Beijing, P.R. China
5. These authors contributed equally

*Correspondence: xpan@ncba.ac.cn ; alli@ncba.ac.cn

Summary

The mitochondrial calcium uniporter (MCU) complex mediates Ca^{2+} entry into mitochondrial, which plays a crucial role in regulating cellular energy metabolism and apoptosis. Dysregulation of MCU is implicated in various diseases, such as neurodegenerative disorders, cardiac diseases and cancer. Despite its importance, developing specific and clinically viable MCU inhibitors has been challenging. Here, we identify Berberine, a well-established drug with a documented safety profile, as a potent MCU inhibitor through a virtual screening of an FDA-approved drug library. Berberine localizes within mitochondria and directly binds to the juxtamembrane loop domain of MCU. This binding disrupts the interaction of MCU with its essential regulator, EMRE, thereby inhibiting rapid Ca^{2+} entry into the mitochondria. Notably, Berberine pretreatment reduces mitochondrial Ca^{2+} overload and mitigate ischemia/reperfusion-induced myocardial injury in mice. Our findings establish Berberine as a potent MCU inhibitor, offering a safe therapeutic strategy for diseases associated with dysregulated mitochondrial calcium homeostasis.

Introduction

The entry of Ca^{2+} ions into mitochondria primarily occurs through a highly selective Ca^{2+} channels known as mitochondrial Ca^{2+} uniporter (MCU) (*1*). This process is crucial for regulating various cellular processes, including mitochondrial dynamics, energy metabolism and cell death (*2*). The MCU complex, along with its regulatory components MICU1, MICU2, and EMRE, ensures precise control of mitochondrial Ca^{2+} uptake, which is crucial for maintaining cellular homeostasis (*3*). However, under pathological conditions, dysregulation of mitochondrial Ca^{2+} homeostasis may lead to detrimental effects. Excessive Ca^{2+} accumulation in mitochondria, referred to as mitochondrial Ca^{2+} overload, triggers mitochondrial permeability transition pore opening, leading to mitochondrial membrane depolarization, release of pro-apoptotic factors, and ultimately leading to cell death. These processes contribute to various diseases, including neurodegenerative disorders, cardiac diseases, and ischemia/reperfusion injury (*4, 5*).

Despite the critical role of MCU in disease pathophysiology, developing effective therapeutic interventions targeting mitochondrial Ca^{2+} regulation remains challenging. Existing MCU inhibitors, such as ruthenium red and Ru360, exhibit poor delivery and toxicity issues (*6, 7*). More recent inhibitors, including DS16570511 and mitoxantrone, have shown promise in preclinical studies but are hampered by toxicity, limiting their clinical applicability (*8, 9*). Ideal MCU inhibitors that is suitable for clinical application must meet several criteria: 1) efficient penetration of cellular and mitochondrial membranes to effectively inhibit mitochondrial Ca^{2+} uptake; 2) selective modulatation of mitochondrial Ca^{2+} uptake without affecting cytosolic Ca^{2+} signaling, ensuring normal cellular functions; 3) preservation of mitochondrial membrane potential, which is vital for mitochondrial function; and 4) low toxicity at therapeutically relevant concentrations to ensure safety and minimal adverse effects.

In response to the demand for safe and effective MCU inhibitors, we employed virtual molecular docking followed by a functional screening of an FDA-approved drug library. This led to the identification of Berberine, a natural alkaloid with established safety profiles, as a novel MCU inhibitor. Berberine localizes to mitochondria, directly binds to MCU, and disrupts the assembly of the MCU-EMRE complex. Importantly, Berberine significantly reduces mitochondrial Ca^{2+} overload, providing cardioprotection against ischemia/reperfusion-induced myocardial injury in mice. This discovery positions Berberine as a promising candidate for further development in the broader context of diseases involving mitochondrial Ca^{2+} dysregulation.

Results

Screening for potential MCU inhibitors from an FDA-approved drug library

MCU functions as a major gate of Ca^{2+} flux from cytosol into the mitochondrial
matrix. Targeting MCU may serve as therapeutic strategies for many diseases, such as
pathological cardiac cell death (*10-12*). To closely emulate MCU-targeted clinical
treatment, we employed a screening approach using an FDA-approved drug library
comprising 2,816 compounds (Fig. 1A). In our initial step, we performed a virtual
screening via MCU structure-based molecular docking and identified 120 hits with a
docking score of < -7.5 kcal/mol, suggesting their potential MCU binding capacity
(fig. S1A). To assess the impact of these hits on mitochondrial Ca^{2+} uptake, we
established HeLa cells expressing 4mt-GCaMP6, a fluorescent indicator reflecting
mitochondrial Ca^{2+} levels (*13*). Subsequently, we generated a screening system for
mitochondrial Ca^{2+} uptake activity using permeabilized Hela cells. Exposure to 100
µM $CaCl_2$ led to a rapid increase of the fluorescent signal, indicating mitochondrial
Ca^{2+} uptake (fig. S1B). Ru360, a well-known MCU inhibitor (*6, 7*), effectively
blocked Ca^{2+} entry into mitochondria (fig. S1, B and C). In a small-scale screen using
the 120 hits, four small molecules demonstrated at least 50% inhibition of
mitochondrial Ca^{2+} uptake at a concentration of 10 µM (fig. S1, B to D). Among
them, Berberine exhibited the highest inhibitory effect, reaching up to 87% (fig. S1D).
To rule out the possibility that the observed effect was a secondary consequence of
mitochondrial membrane potential disruption, the major driving force for Ca^{2+} uptake
within the organelle (*14, 15*), we performed TMRM staining analysis. The
mitochondrial decoupler, CCCP, induced a decrease in TMRM intensity. All tested
drugs, except for Berberine, including Trifarotene, Eltrombopag, and Tipranavir,
significantly decreased the mitochondrial membrane potential (fig. S1E). Therefore,
this screening process identified Berberine, a natural compound derived from
medicinal herbs, as a promising candidate for further investigation as MCU inhibitor.

Berberine inhibits mitochondrial Ca^{2+} uptake

To assess Berberine's ability to traverse cellular membranes and inhibit
mitochondrial Ca^{2+} uptake, we evaluated its inhibitory effects in intact cells by
initiating intracellular Ca^{2+} signaling with histamine stimulation while simultaneously
monitoring cytosolic and mitochondrial Ca^{2+} dynamics (Fig. 1B). Berberine
specifically inhibited mitochondrial Ca^{2+} uptake, as indicated by the green fluorescent
indicator 4mt-GCaMP6 (Fig. 1, C and D), without affecting the histamine-induced
cytosolic Ca^{2+} signals, as evidenced by the red fluorescent dye Calbryte590 (Fig. 1, E
and F). Similar results were obtained using other Ca^{2+} indicators, cyto-GCaMP6, or
4mt-GCaMP8 in HeLa cells (fig. S2, A to D). These findings suggest that berberine

can freely traverse the cellular membrane, thereby inhibiting mitochondrial
Ca^{2+} uptake without impacting the cytosolic Ca^{2+} signals.

Next, to further quantify berberine's inhibitory effect on mitochondrial
Ca^{2+} uptake, we permeabilized the cells with digitonin, allowing equal doses of
calcium ions to be added outside the mitochondrial membrane, thus evaluating the
capacity of mitochondria to absorb Ca^{2+} at different concentrations of Berberine.
Berberine exhibited a potent inhibition of mitochondrial Ca^{2+} uptake in a dose-
dependent manner (Fig. 1G), with an IC_{50} value determined to be 2.202 µM (Fig. 1H).

To evaluate the safety of Berberine in cells, we treated the cells with increasing
amount of Berberine, and confirmed that up to 10 µM Berberine treatment did not
affect mitochondrial membrane potential by TMRM staining (fig. S2, E and F). We
further investigated the cytotoxicity of Berberine to understand its potential impact on
cell growth. It showed that at lower doses, specifically between 5-10 µM, which are
sufficient to inhibit MCU, Berberine exhibited minimal cytotoxic effects (fig. S2G).
This indicates that Berberine is an effective MCU inhibitor with low cytotoxicity,
making it a relative safe option at these concentrations.

Direct binding of Berberine to MCU

Molecular docking screenings indicate that BBR may directly bind to the MCU
monomer (Fig. 1A, and fig. S1A). This leads us to suggest that Berberine possesses
the unique ability to directly traverse cellular membranes, gaining entry into
mitochondria to target and subsequently inhibit MCU. Previous studies have
documented the subcellular localization of Berberine within mitochondria (*16, 17*).
Exploiting Berberine's autofluorescence property, we also observed its cellular
penetration at 10 µM, co-localizing with MitoTracker stain (fig. S3, A and B),
reinforcing our hypothesis.

To further validate our hypothesis, we synthesized biotin-conjugated Berberine
(Biotin-BBR) (Fig. 3A). The Biotin-BBR exhibited similar mitochondrial localization
(Fig. 2, B and C) and inhibition efficacy as unmodified Berberine (fig. S3, C and D),
suggesting that the Biotin-BBR retains the ability of interacting with the same
molecular targets as the unmodified Berberine. Consistent with these results, Biotin
pull-down assays demonstrated that Biotin-BBR could interact with both exogenous
and endogenous MCU in cell lysate (Fig. 2D, and fig. S3E). *In vitro* pull-down assays
further confirmed that Biotin-BBR directly binds to recombinant glutathione-S-
transferase (GST)-MCU (Fig. 2E). Notably, this binding was diminished when
unlabeled Berberine competitively interacted with GST-MCU, indicating a consistent
binding site and confirming the specificity of the binding (Fig. 2E). Microscale
thermophoresis (MST) analysis further confirmed that the affinity between GST-
MCU and Berberine was ~8 µM (Fig. 2F).

Berberine targets the juxtamembrane loop domain of MCU

Next, to explore the mechanism of Berberine's binding to MCU and its functional impact, we first mapped MCU regions interacting with Berberine. We *in vitro* translated MCU truncations based on its secondary structure (*18*), excluding the mitochondrial targeting sequence (residues 1-74) due to its non-conservative nature (*19*). We assessed their interactions with Berberine using Biotin pull-down assays. Berberine showed a strong affinity for the MCU juxtamembrane loop (JML) segment spanning residues 275-295 (Fig. 3, A and B), although minor binding to other areas was also observed (Fig. 3B). These observations were consistent with molecular docking simulations (Fig. 3C), which indicated that Berberine binds near JML domain, a region critical for EMRE-dependent MCU opening (*18, 20*). Docking analyses pinpointed key binding residues, including Tyr 281, Tyr 289, Tyr 291, and Ala 294 (Fig. 3C, and fig. S3F). Notably, Tyr 281 and Tyr 289 are likely pivotal for hydrogen bonds with Berberine as shown by molecular docking simulations (Fig. 3C, and fig. S3F).

To explore the interaction of Berberine with predicted binding sites within the MCU, we performed site-directed mutagenesis. Sequence alignment of the JML domain revealed four conserved residues (Y281, Y289, Y291, and A294) across various MCU homologs (fig. S3G), known to play a role in maintaining the open conformation of the MCU (*20*). It was reported that mutations of Y289 and Y291 to alanine, as well as A294 to phenylalanine, do not alter the open conformation of MCU and preserve robust mitochondrial Ca^{2+} uptake capacity. However, the critical residue Y281, when mutated to alanine, induces a closed conformation of the MCU (*20*). Under these circumstances, simultaneous mutations of Y281A, Y289A, Y291A, and A294F significantly disrupted the interaction with the Berberine-MCU JML domain (Fig. 3D), confirming Berberine's direct binding to these residues. To determine whether the ability of Berberine to bind was independent of the MCU's conformational state (open or closed), we introduced mutations Y281H and Y281R, which maintain the open conformation (*20*) but disrupt potential hydrogen bonding with Berberine. These mutations still abolished Berberine binding (Fig. 3D), negating the hypothesis that Berberine's binding is confined to the open conformation of MCU.

To further demonstrate that Berberine's inhibition of the calcium ion absorption function of the MCU is dependent on its binding to the JML domain, we engineered cell lines with stable reintroduction of either the wild-type MCU (MCU^{WT}) or the MCU^{HAAF} mutant (Fig. 3E). This experimental design enabled us to contrast the effects of Berberine on mitochondrial Ca^{2+} uptake across these scenarios. Berberine treatment led to a reduction in mitochondrial Ca^{2+} uptake in both control and MCU^{WT}-reconstituted cells (Fig. 3, F and G). In contrast, cells expressing the MCU^{HAAF} mutant, which does not bind Berberine due to alterations in the JML domain (Fig. 3D), exhibited no decrease in mitochondrial Ca^{2+} uptake following Berberine

187 treatment (Fig. 3, F and G). This divergence underscores the critical role of the JML
188 domain interaction in Berberine's mechanism of action on mitochondrial
189 Ca^{2+} regulation.

Berberine disrupts MCU-EMRE interaction

Ca^{2+} transfer through the mitochondrial inner membrane is mediated by the
mitochondrial calcium uniporter (MCU) complex, consisting of MCU, EMRE,
MICU1, and MICU2 (*3*). We investigated Berberine's influence on the MCU complex
assembly via co-immunoprecipitation (IP) experiments. The components of MCU
complex, including MCU, EMRE, MICU1 and MICU2, were SNAP-tagged and co-
expressed with Flag-tagged MCU in 293T cells. Following Berberine treatment, only
the interaction between SNAP-tagged EMRE and Flag-tagged MCU was significantly
diminished, while the interactions of MCU with other MCU components remained
unaltered (Fig. 4A). These results suggest that Berberine selectively disrupts the
EMRE-MCU interaction, which was further confirmed by a reverse IP assay (fig.
S4A).

To directly assess Berberine's effect on MCU-EMRE interaction, we employed
in vitro binding assays. These assays confirmed Berberine's ability to directly
interfere with MCU-EMRE assembly (Fig. 4B). To understand the mechanism behind
this disruption, we conducted molecular dynamics simulations. The simulations
revealed that the MCU-EMRE complex rapidly reached equilibrium and maintained
stability during the simulation in the absence of Berberine. At 100 ns, the Rg value, a
measure of compactness, significantly increased from 3.734 nm to 4.982 nm upon
Berberine addition (Fig. 4C), indicating a more spread-out and less compact complex
structure. This observation was further corroborated by the increased RMSD values
(reflecting average atomic displacement) in the presence of Berberine (Fig. 4D).
Together, these results suggest that Berberine induces a looser and less stable
complex structure (fig. S4B). Furthermore, hydrogen bond analysis revealed a
reduction in the average number of bonds between MCU and EMRE from 10.540 to
6.287 upon Berberine treatment (fig. S4, C and D), signifying a weakening of
interactions between the two proteins. This finding aligns with the significant
decrease in binding energy observed between MCU and EMRE in the presence of
Berberine (from -100.02 ± 17.85 kcal/mol to -29.52 ± 27.07 kcal/mol, fig. S4E).
Collectively, these data strongly support the conclusion that Berberine disrupts the
MCU-EMRE complex by weakening the interactions between MCU and EMRE.

Next, we sought to clarify that whether Berberine's disruption of the MCU-
EMRE binding activity depends on its interaction with the MCU JML domain. We
co-transfected cells with MCU^{WT} or the Berberine (BBR)-binding deficient mutant
MCU^{HAAF} alongside EMRE. Both MCU^{WT} and MCU^{HAAF} were capable of interacting
with EMRE, indicating that the mutation in the JML domain itself does not affect its

assembly with EMRE (Fig. 4E). The addition of Berberine disrupted the assembly
between MCU^{WT} and EMRE; however, when Berberine could not bind to the MCU
JML domain, as with MCU^{HAAF}, Berberine's presence did not influence the MCU-
EMRE assembly. These findings, together with our molecular dynamics simulation
results (Fig. 4, C and D, and fig. S4, B to E), suggest that Berberine's modulation of
mitochondrial Ca^{2+} uptake is contingent upon its interaction with MCU. This
interaction subsequently disrupts the MCU-EMRE complex assembly and ultimately
leads to the inhibition of the MCU channel, thereby preventing Ca^{2+} entry into the
mitochondria (Fig. 4F).

Berberine suppresses mitochondrial Ca^{2+} overload and protects against

myocardial ischemia-reperfusion injury

We next sought to explore the physiological relevance of our findings. It's well
established that excessive Ca^{2+} entry into mitochondria can lead to mitochondrial Ca^{2+}
overload. This overload subsequently triggers the opening of the permeability
transition pore (PTP), causing mitochondrial swelling, and eventually, cell death (*21*).
Cardiac-specific conditional knockout of MCU could protect mitochondria by
preventing the opening of mPTP *in vitro* and *in vivo* (*10, 11*). The mPTP opening can
be monitored by measuring mitochondrial absorbance (*22*). In an *in vitro* context, we
observed that Berberine inhibited Ca^{2+} overload-induced mitochondrial swelling (Fig.
5A). This effect paralleled those of Ru360 and cyclosporin A (CsA) (*23*), a known
inhibitor of mPTP opening, suggesting that Berberine might confer cellular protection
under conditions of mitochondrial Ca^{2+} overload.

To validate this hypothesis, we first established a model of
hypoxia/reoxygenation (H/R) injury in neonatal rat cardiomyocytes and embryonic rat
cardiomyocyte-derived H9c2 cell line. Consistent with previous studies (*24, 25*), H/R
injury led to a significantly elevated mitochondrial Ca^{2+} levels, indicated by
fluorescent intensity of 4mtGCaMP6 or Rhod2 (Fig. 5, B and C, and fig. S5, A and
B). Furthermore, Berberine pretreatment alleviated this mitochondrial Ca^{2+} overload
(Fig. 5, B and C, and fig. S5, A and B). To assess the potential cardioprotective effect
of Berberine *in vivo*, a mouse acute cardiac ischemia/reperfusion (I/R) model was
established (Fig. 5D). Pretreatment of Berberine (24h plus 15 min before reperfusion)
significantly reduced I/R induced myocardial infarct size (Fig. 5, E and F).
Additionally, Berberine ameliorated cardiomyocyte death, evidenced by decreased
cTNl and LDH concentration (Fig. 5, G and H). TUNEL staining experiments further
confirmed the protective effect of Berberine against myocadiac I/R injury (Fig. 5I). In
summary, these data indicate that Berberine exhibits therapeutic effects in myocardial
ischemia/reperfusion injury via counteracting mitochondrial Ca^{2+} overload.

Discussion:

Our virtual screening approach using an FDA-approved drug library identified
Berberine, a natural alkaloid derived from medicinal plants, as a potent MCU
inhibitor. Berberine exerts its inhibitory effect by directly binding to the MCU and
disrupting the assembly of MCU-EMRE complex, consequently hindering Ca^{2+} entry
into mitochondrial matrix. Notably, Berberine significantly reduces mitochondrial
Ca^{2+} overload, thereby protecting cardiomyocytes from ischemia-reperfusion (I/R)-
induced apoptosis *in vitro* and *in vivo*. These results underscore the potential of
Berberine as a promising MCU-inhibiting drug for clinical application.

Berberine, a well-characterized plant alkaloid with a long history of traditional
use, has been employed for treating various conditions, including diabetes,
hyperlipidemia, and cardiovascular diseases (*26*). Its antimicrobial properties are also
well-documented (*27*). Despite its extensive therapeutic application, Berberine's
precise molecular targets remains elusive. Our study provides compelling evidence
that Berberine targets MCU, effectively impeding mitochondrial Ca^{2+} uptake through
direct binding. Importantly, Berberine may overcome limitations observed in existing
MCU inhibitors such as Ru360, DS16570511 (*8*), mitoxantrone (*9*), and MCU-i11
(*28*)—a specific MICU1-targeting inhibitor—in animal experiments. DS16570511,
for instance, exhibits off-target effects on mitochondrial membrane potential, thereby
limiting its application (*29*). Similarly, mitoxantrone, a topoisomerase inhibitor, is
known to have cardiotoxic effects (*30*). Berberine's well-established safety profile *in*
vivo positions it as a strong candidate for further exploration in clinical settings.

Our studies, along with previous studies, consistently demonstrate Berberine's
capacity to provide significant cardioprotection in the context of myocardial I/R
injury (*31*). Clinical studies further support its potential, indicating that Berberine can
modulate inflammatory markers in patients suffering from acute myocardial infarction
(*32*). Berberine's protective effects are multifaceted, encompassing antioxidant, anti-
inflammatory, and anti-apoptotic activities. A key aspect of this protection is
Berberine's ability to maintain mitochondrial integrity and high activity of the
respiratory complexes (*33-35*), as evidenced by its regulation of Bcl-2 and Bax
proteins, which are crucial for mitochondrial membrane stability and the prevention of
apoptotic factor release (*33, 36*). Our findings establish MCU as a key cardiac target
of Berberine, revealing a core mechanism for its protective effects in I/R injury.
Given that mitochondrial Ca^{2+} overload occurring rapidly within the initial 20 minutes
post-ischemia (*25*), early intervention with Berberine is of paramount importance.
Furthermore, Berberine's potential interactions with other targets and its slower-acting
mechanisms warrant further investigation to fully understand its therapeutic potential.

Our findings reveal that Berberine binds directly to the MCU near the
juxtamembrane loop (JML) domain, at residues Y281, Y289, Y291, and A294. The

JML is essential for channel gate formation and its interaction with EMRE is critical
for Ca^{2+} channel activation (18). Mutational analysis of Y281 and Y289 highlights
their significance in MCU-EMRE assembly and the EMRE-mediated conformational
opening of MCU (20). Y281A mutants significantly disrupts EMRE binding (20),
paralleling the loss of Ca^{2+} uptake observed with Berberine treatment. Notably, Y289,
a key hydrophobic residue within the JML, facilitates EMRE-dependent MCU
opening (20). Substituting Y289 with hydrophilic residues impairs channel opening,
underscoring the critical role of hydrophobic interactions at this site for EMRE
binding and MCU activation. Berberine's hydrophobic surface (37) might promote its
integration near the JML, disrupting interactions between critical residues like Y281
and Y289, and fostering a closed MCU conformation unfavorable to EMRE
association. While our molecular dynamics simulations support this hypothesis, the
complex mechanisms of MCU gating warrant further structural studies for full
elucidation.

In conclusion, our study identifies Berberine as a novel MCU inhibitor with
significant potential for clinical applications. Revealing its inhibitory effects on MCU
and providing insights into the complex Berberine-MCU-EMRE dynamics, our
findings lay a solid foundation for the development of targeted drug strategies aimed
at mitochondrial Ca^{2+} regulation. Structural analogs of Berberine could be engineered
for enhanced specificity and safety as MCU inhibitors. Furthermore, the role of MCU
dysfunction is becoming increasingly understood, not only in cardiac
ischemia/reperfusion injury but also for other diseases, such as the recently reported
dependency of venetoclax-resistant leukemic stem cells (LSCs) on mitochondrial
calcium for survival(38). These findings position Berberine as a valuable starting
point for developing novel interventions in diseases associated with dysregulated
mitochondrial Ca^{2+} homeostasis.

Materials and Methods

Experimental animals

All animals used in this study were maintained in the pathogen-free barrier animal facility at the National Center of Biomedical Analysis. Animal care was monitored daily by certified veterinary staff and laboratory personnel. 10-week-old mice in the C57BL/6J background and pregnant Sprague-Dawley rats were purchased from the Beijing Vital River Laboratory Animal Technology Company.

Murine model of myocardial ischemia

The 10 week-male-C57BL/6 mice (>22 g) were randomly assigned to specific groups. Berberine (5 mg/kg) or vehicle was administered intraperitoneally twice (24 hours and 15 minutes prior) to ischemia-reperfusion (I/R) surgery as described (*39*). Briefly, pentobarbital (70 mg/kg i.p.) was used to induce and maintain anesthesia. Mice were then ventilated on a Harvard rodent respirator via a tracheostomy. A midline sternotomy was performed, with a reversible snare occlude placed around the left anterior descending coronary artery. I/R injury was initiated by tightening the snare for 45 minutes, then releasing it. After reperfusion for 24 hours, the mice were sacrificed for further detection and measurement.

For serum LDH concentration and cTnI level, blood samples were centrifuged at 3,000 rpm for 10 min and then detected using an LDH kit (Sigma-Aldrich, Cat# MAK066) and a mouse cardiac troponin I ELISA kit (ThermoFisher, Cat#EEL112) according to the manufacturer's instructions.

For the measurement of the area-at-risk (AAR) and infarct size, the heart was initially excised post-reocclusion of the coronary artery at the previous occlusion site. The heart ascending aorta was cannulated (distal to the sinus of Valsalva) and then perfused retrogradely with 0.05% Alcian blue to delineate the AAR. Following AAR visualization, the heart was frozen at -80 □ for 5 min, cut into ~1 mm slices, and the unstained infarcted (IF) region was lucifagal visualized using 1% 2,3,5-triphenyl-tetrazolium chloride at 37 □ for 15 min. Infarct and left ventricular areas were determined by planimetry with Image J. The infarct size was calculated as infarct area divided by area at risk (IF/AAR).

Molecular Docking Screening

The 3D structures of MCU (PDB: 6XJV) were retrieved from the Protein Data Bank. A set of 2,816 FDA-approved drugs were converted to 3D structures using Open Babel. Preparation and parameterization of the receptor protein and ligands were conducted with AutoDock Tools (ADT3). Docking grid documents were created using AutoGrid of sitemap, and docking simulation were performed with AutoDock Vina (1.2.0) (*40*). The score was ranked according to the value of the binding energy

368   of protein-ligand complex. Finally, the protein-ligand interaction figure was generated
369   by PyMOL.

Cell culture

To establish HeLa cells stably expressing 4mt-GCaMP6s, we constructed a
lentiviral vector (pLVX-puro) containing the 4mt-GCaMP6s insert. This vector was
transfected into HEK293T cells along with the packaging plasmids pSPAX2 and
pVSV-G using a 4:3:1 ratio. Stable transfectants were selected using 1 µg/ml
puromycin. HeLa cells were cultured at 37°C in DMEM supplemented with 10% FBS
and 1% penicillin-streptomycin. A similar approach was employed to express MCU
WT and MCU HAAF in the HeLa cell line. Initially, HeLa cells were infected with
the pLKO.1-shMCU lentivirus and subsequently selected with 1 µg/ml puromycin.
Knockout HeLa cells were then infected with lentivirus carrying pLVX-mCherry-N1
vectors harboring either MCU^{WT} or MCU^{HAAF}. Cells expressing mCherry were
isolated using flow cytometry. Knockout of endogenous MCU and expression of
exogenous MCU-mCherry were confirmed by Western blot analysis. All other
mammalian cell lines used in this study were from the American Type Culture
Collection. All the cell lines were fully authenticated and tested free of mycoplasma.

Mitochondrial Ca^{2+} modulator Screening

Hela cells expressing 4mt-GCaMP6s were permeabilized by a 1 min perfusion
with intracellular buffer (130 mM KCl, 10 mM NaCl, 2 mM K$_2$HPO$_4$, 5 mM Succinic
acid, 5 mM Malic acid, 1 mM Pyruvate, 0.5 mM ATP, 0.1 mM ADP, MgCl$_2$, 20 mM
HEPES, pH 7, 37 □) supplemented with 10 µM digitonin and 50 µM EGTA. Then the
buffer was changed to intracellular buffer supplemented with 10 µM compounds from
primary screen hits of FDA-approved drugs for 10 min. Mitochondrial Ca^{2+} uptake
was evoked by adding 100 µM CaCl$_2$ to the buffer. Ca^{2+} imaging was performed
using a DeltaVision Deconvolution microscope (GE Healthcare) equipped with a ×20
objective. Exposure time was set to 25 ms for FITC. Images were acquired at 1 frame
per 3 seconds with an EM-CCD camera. Images were analysed using Image J
software. All images were background-corrected by subtracting mean pixel values of
a cell-free region of interest.

Molecular dynamics (MD) simulations

Structural data for MCU and EMRE monomers were retrieved from the Protein
Data Bank (PDB) via the entry code 6XJV. The preprocessing phase employed
Gromacs 2018 as the simulation suite, coupled with the Amber99SB-ildn force field
for encompassing protein and small molecule interactions. A 10×10×10 nm^3 water
box, utilizing the TIP3P model, was constructed to solvate the system, ensuring a
minimal 1.2 nm buffer between the protein surfaces and the box boundaries. This was
followed by ion addition for electrostatic equilibrium.

For treating electrostatic interactions, we implemented the Particle-mesh Ewald
(PME) method. Energy minimization was executed through the steepest descent
method over 50,000 steps, setting both the Coulombic and van der Waals interaction
cutoffs at 1 nm. Equilibration phases under NVT (constant number of particles,
volume, and temperature) and NPT (constant number of of particles, pressure, and
temperature) settings preceded a 100 ns MD simulation, conducted under standard
laboratory conditions of temperature and pressure. Non-bonded interactions were
applied with a 10 Å cutoff. Thermal and pressure stability were maintained at 300 K
and 1 bar, respectively, utilizing the V-rescale thermostat and the Berendsen barostat.

This rigorous preparation and execution strategy yielded a robust framework for
accurately simulating the complex's behavior in a physiological context, culminating
in the acquisition of data such as radius of gyration (Rg), root-mean-square deviation
(RMSD), number of hydrogen bonds (H-bonds), and total free energy, to assess
structural stability, conformational dynamics, interaction strength, and energetics
throughout the simulation period.

Ca^{2+} measurements

For imaging Ca^{2+} transients evoked by treatments with an agonist, cells labelled
with Calbryte590 or 4mt-GCaMP6 were incubated at 37 □ with Krebs-Ringer
modified buffer (KRB), which contains 125 mM NaCl, 5 mM KCl, 1 mM Na_3PO_4, 1
mM $MgSO_4$, 5.5 mM glucose and 20 mM HEPES (pH 7.4). After 10 μM histamine
stimulation, a series of images were captured every 3 s using a DeltaVision
Deconvolution microscope (GE Healthcare). The recorded images were analyzed and
quantified using Image J software (NIH).

For image analysis, background correction was first performed frame by frame
by subtracting the intensity of a nearby cell-free region from the signal of the imaged
cell. F_0 is calculated as the initial background-subtracted fluorescence intensity. F
indicates the background-subtracted fluorescence intensity at each time point. $\Delta F/F_0=$
$(F – F_0)/F_0$, indicates the Ca^{2+} concentration change after stimulation.

Mitochondrial membrane potential measurements

For confocal microscopy, adherent HeLa cells were stained with
Tetramethylrhodamine (TMRM) at a final concentration of 100 nM in fresh culture
medium. After a 30-minute incubation period at 37°C to allow for adequate dye
uptake, cells were treated with the compounds of interest for an additional 30 minutes
at the same temperature. Subsequently, cells were washed to remove excess dye and
unbound compounds. Imaging was performed on a DeltaVision confocal microscope
(GE Healthcare). Acquired images were then processed and quantified using ImageJ
software.

For flow cytometric analysis, Hela cells were digested and stained with 100 nM
Tetramethylrhodamine (TMRM) in fresh medium for 30 min at 37□. Compounds
were added for 30 min at 37□. Then cells were washed twice and resuspended in PBS
containing 7AAD. Samples were analyzed using cytoFLEX flow cytometer (Beckman
Coulter).

Western blot analysis.

For western blot analysis, cells were lysed in RIPA buffer (50 mM Tris-HCl (pH
8.0), 150 mM NaCl, 1% (v/v) Nonidet P-40, 0.5% sodium deoxycholate, 0.1% SDS,
protease inhibitor cocktail). Protein samples were resolved by SDS-PAGE and
transferred onto PVDF membranes. Membranes were immunoblotted with the
indicated primary antibodies. Antibodies were used at the following concentrations:
anti-MCU, rabbit polyclonal, Cell Signaling Technology, Cat# 14997 (1:1000); anti-
SNAP, rabbit polyclonal, New England Biolabs, Cat# P9310S (1:1000).

Immunoprecipitation

HEK293T cells were transfected with the indicated plasmids using
Lipofectamine^{TM} 2000 transfection reagent (ThermoFisher, Cat# 11668500). After
transfection for 24 h, cells were treated with 10 μM Berberine for 2 hours before
harvested. Cells were lysed with RIPA buffer (50 mM Tris-HCl (pH 8.0), 150 mM
NaCl, 1% (v/v) Nonidet P-40, 0.5% sodium deoxycholate, protease inhibitor
cocktail).

Cell lysates were immunoprecipitated by anti-Flag M2 affinity gel (Selleck, Cat#
B23102) overnight at 4 □. Beads were washed 5 times in RIPA buffer followed by
eluting bound protein complexes in 2× SDS dye. Whole cell lysates and co-
precipitation samples were analyzed by western blot.

Biotin pulldown assay

For *in vivo* Biotin pulldown assay, HEK293T cells were treated with Biotin or
Biotin-labeled Berberine (10 μM, 2 h). Cells were lysed and harvested with RIPA
buffer, which containing 50 mM Tris-HCl, 150 mM NaCl, 1% NP-40 and 0.5%
sodium deoxycholate. Balanced Streptavidin Agarose beads (ThermoFisher, Cat#
20357) were added and incubated overnight at 4 °C. The beads were washed with
RIPA buffer adding 0.03% Tween 5 minutes for 5 times. The indicated proteins were
detected by Immunoblotting.

For the *in vitro* biotin pulldown assay, biotin or biotin-labeled berberine (2 μg)
was pre-incubated with either recombinant protein (2 μg) or in vitro
transcribed/translated protein (Promega, Cat# L1170) for 8 hours at 4°C. The mixture
was then incubated with streptavidin agarose beads (Thermo Fisher, Cat# 20357)
overnight at 4°C. The beads were washed as previously described and the bound
proteins were analyzed by immunoblotting.

Immunofluorescence and confocal microscopy

For visualizing the intracellular distribution of Berberine, HeLa cells were
grown on coverslips, incubated with 50 nM MitoTracker™ DeepRed (ThermoFisher,
Cat# M22426) in fresh medium for 15 min at 37 °C. 10 μM Berberine was added for
10 min. After treatment, the medium was replaced with phenol red-free DMEM
supplemental with 10% FBS. Then cells were visualized using a DeltaVision
Deconvolution microscope (GE Healthcare). Images of Berberine and MitoTracker™
DeepRed signals, acquired by FITC and Cy5 filters respectively, were merged to
visualize their co-localization in mitochondria.

For visualizing the intracellular distribution of Biotin-labeled Berberine,
HeLa cells were grown on coverslips, incubated with 10 μM Biotin-Berberine for 10
min at 37 °C. After treatment, cells were fixed in 4% paraformaldehyde for 5 min.
Next, the cells were incubated with 0.3% Triton X-100 in PBS for 10 min on ice and
then blocked in 3% BSA in PBS for at least 1 h. The primary antibody was incubated
overnight at 4°C followed by secondary antibody incubation. Finally, the cells were
mounted using 80% glycerol. Antibodies were used at the following concentrations:
anti-COXIV, rabbit polyclonal, Cell Signaling Technology, Cat# 4850 (1:200); Alexa
Fluor 488-conjugated streptavidin antibody, ThermoFisher, Cat# S11223 (1:400);
Alexa Fluor 546-conjugated anti-Rabbit Secondary Antibody, ThermoFisher, Cat#
A11035 (1:400). Images were acquired using a DeltaVision Deconvolution
microscope.

Mitochondrial isolation and swelling assay

For isolating cardiac mitochondria, mouse hearts were minced in mitochondrial
isotonic buffer (225 mM mannitol, 75 mM sucrose, 5 mM MOPS, 0.5 mM EGTA and
2 mM taurine, pH 7.25) supplemented with protease inhibitor cocktail, and
subsequently homogenized using a dounce homogenizer. The homogenate was
initially centrifuged for 10 min at $700 \times g$ and the resulting supernatant was next spun
for 10 min at $12,000 \times g$ to pellet the mitochondria.

Mitochondrial swelling was measured as previously described (*23*). In brief,
isolated cardiac mitochondria (100 μg) were resuspended in swelling buffer (120 mM
KCl, 10 mM Tris-HCl, 5 mM MOPS, 5 mM Na_2HPO_4, 10 mM glutamate, 2 mM
malate and 0.1 mM EGTA) in a total volume of 200 μl. Pore opening was induced
and detected by the addition of 500 μM of total $CaCl_2$ while monitoring absorbance at
540 nm. Where indicated, Berberine (10 μM), cyclosporin A (3 μM) or Ru360 (3 μM)
was added.

Isolation and culture of primary neonatal rate cardiomyocytes (NRCMs)

NRCMs were isolated and cultured as described previously (*41*). Briefly,
cardiomyocytes were isolated from the neonatal hearts of <24h-old Sprague-Dawley
rats using PBS containing 0.03% trypsin and 0.04% collagenase type II. The digestion

was halted by Dulbecco's modified Eagle's medium (DMEM), supplemented with
10% fetal bovine serum (FBS). After differential attachment for 90 min, the cells in
suspension were then plated and cultured in DMEM with 10% FBS at 37°C with 5%
CO_2 for 24 h.

Hypoxia/reoxygenation (H/R) injury model

The hypoxia/reoxygenation (H/R) injury of NRCMs or H9c2 cells were achieved
in incubator (Thermo Fisher Scientific, Madison, WI, USA) filled with 94 % N_2, 5 %
CO_2, and 1% O_2 in DMEM deprived of glucose and sodium pyruvate for 4h, followed
by 8 h-reoxygenation by transferring the cells to a normoxia incubator with normal
supplement of oxygen and glucose.

To examine the protective effect of BBR on H/R-induced Ca^{2+} overload,
NRCMs infected with Ad-4mt-GCaMP6 or H9c2 cells stained with Rhod2 were
treated with BBR at a final concentration 10 μM throughout the H/R period. Then
cells were visualized using a DeltaVision Deconvolution microscope (GE
Healthcare). To assess mitochondrial Ca^{2+} level using Rhod2, cells labeled with
Rhod2 were illuminated at 546 nm, and fluorescence was collected through a TRITC
filter. To assess mitochondrial Ca^{2+} level using 4mt-GCaMP6, cells labeled with 4mt-
GCaMP6 were illuminated at 488 and 390 nm, and fluorescence was collected
through a FITC filter. Images were analysed using Image J software. All images were
background-corrected by subtracting mean pixel values of a cell-free region of
interest.

Statistics and reproducibility.

Statistical comparisons between only two groups were carried out using unpaired
Student's t test or the Mann-Whiney U test when a normal distribution could not be
assumed. For comparisons across multiple groups, a one-way analysis of variance
(ANOVA) was employed. Furthermore, when examining the effects of two
independent variables or the interaction between them on a dependent variable, a two-
way ANOVA was utilized to discern any significant differences. Statistical
calculations were carried out using GraphPad Prism 6.0. We tested data for normality
and variance, and considered a *P* value of less than 0.05 as significant. All
experiments were performed three or more times independently under identical or
similar conditions.

Acknowledgments

The authors thank D.D. Stefani for providing 4mt-GCaMP6 plasmids, and K. Wang and X. Xu for assistance with microscopy assays. This work was supported by the National Natural Science Foundation of China (grant numbers 82025028, 92354304 and 82000107), the National Key R&D Program of China (2021YFA1300203 and 2020YFA0113300), China Postdoctoral Science Foundation (2022M723897).

Author Contributions

Xin Pan conceived the study. Haixin Zhao designed and carried out most of the experiments with the assistance from Siqi Chen, Nian Cao, and Wenjun Wu. Haixin Zhao, Siqi Chen, Guangqin Liu and Haizhen Zhu performed calcium measurements and biochemical experiments. Haixin Zhao, Nian Cao and Jun Gao contributed to mouse studies. Haixin Zhao and Wenjun Wu performed Screening and Data Analysis. Dingyi Lu and Lingmin Zeng performed immunostaining and flow cytometry analysis. Jiayi Chen, Teng Li, Ting Li, Weina Zhang, Qing Xia, Tao Zhou, Ai-Ling Li, Xue-Min Zhang, and Xin Pan contributed to interpreting the results. Xin Pan and Ai-Ling Li supervised the research. Xin Pan and Haixin Zhao wrote the paper.

Declaration of Interests

The authors declare no competing interests.

Reference:

1. Kamer KJ & Mootha VK (2015) The molecular era of the mitochondrial calcium uniporter. *Nat Rev Mol Cell Biol* 16(9):545-553.

2. Giorgi C, Marchi S, & Pinton P (2018) The machineries, regulation and cellular functions of mitochondrial calcium. *Nat Rev Mol Cell Biol* 19(11):713-730.

3. Murphy E & Steenbergen C (2021) Regulation of Mitochondrial Ca(2+) Uptake. *Annu Rev Physiol* 83:107-126.

4. Bernardi P, Rasola A, Forte M, & Lippe G (2015) The Mitochondrial Permeability Transition Pore: Channel Formation by F-ATP Synthase, Integration in Signal Transduction, and Role in Pathophysiology. *Physiol Rev* 95(4):1111-1155.

5. Bauer TM & Murphy E (2020) Role of Mitochondrial Calcium and the Permeability Transition Pore in Regulating Cell Death. *Circ Res* 126(2):280-293.

6. Moore CL (1971) Specific inhibition of mitochondrial Ca++ transport by ruthenium red. *Biochem Biophys Res Commun* 42(2):298-305.

7. Ying WL, Emerson J, Clarke MJ, & Sanadi DR (1991) Inhibition of mitochondrial calcium ion transport by an oxo-bridged dinuclear ruthenium ammine complex. *Biochemistry* 30(20):4949-4952.

8. Kon N, *et al.* (2017) DS16570511 is a small-molecule inhibitor of the mitochondrial calcium uniporter. *Cell Death Discov* 3:17045.

9. Arduino DM, *et al.* (2017) Systematic Identification of MCU Modulators by Orthogonal Interspecies Chemical Screening. *Mol Cell* 67(4):711-723 e717.

10. Luongo TS, *et al.* (2015) The Mitochondrial Calcium Uniporter Matches Energetic Supply with Cardiac Workload during Stress and Modulates Permeability Transition. *Cell Rep* 12(1):23-34.

11. Kwong JQ, *et al.* (2015) The Mitochondrial Calcium Uniporter Selectively Matches Metabolic Output to Acute Contractile Stress in the Heart. *Cell Rep* 12(1):15-22.

12. Lambert JP, *et al.* (2019) MCUB Regulates the Molecular Composition of the Mitochondrial Calcium Uniporter Channel to Limit Mitochondrial Calcium Overload During Stress. *Circulation* 140(21):1720-1733.

13. Zhao H, *et al.* (2019) AMPK-mediated activation of MCU stimulates mitochondrial Ca(2+) entry to promote mitotic progression. *Nat Cell Biol* 21(4):476-486.

14. Baughman JM, *et al.* (2011) Integrative genomics identifies MCU as an essential component of the mitochondrial calcium uniporter. *Nature* 476(7360):341-345.

15. De Stefani D, Raffaello A, Teardo E, Szabo I, & Rizzuto R (2011) A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter. *Nature* 476(7360):336-340.

16. Jin M, *et al.* (2020) Synthesis of a novel fluorescent berberine derivative convenient for its subcellular localization study. *Bioorg Chem* 101:104021.

17. Pereira GC, *et al.* (2007) Mitochondrially targeted effects of berberine [Natural Yellow 18, 5,6-dihydro-9,10-dimethoxybenzo(g)-1,3-benzodioxolo(5,6-a) quinolizinium] on K1735-M2 mouse melanoma cells: comparison with direct effects on isolated mitochondrial fractions. *J Pharmacol Exp Ther* 323(2):636-649.

18. Wang Y, *et al.* (2019) Structural Mechanism of EMRE-Dependent Gating of the Human

615   Mitochondrial Calcium Uniporter. *Cell* 177(5):1252-1261 e1213.

616   19.   Lee SK*, et al.* (2016) Structural Insights into Mitochondrial Calcium Uniporter Regulation by
617   Divalent Cations. *Cell Chem Biol* 23(9):1157-1169.

618   20.   Van Keuren AM*, et al.* (2020) Mechanisms of EMRE-Dependent MCU Opening in the
619   Mitochondrial Calcium Uniporter Complex. *Cell Rep* 33(10):108486.

620   21.   Rasola A & Bernardi P (2011) Mitochondrial permeability transition in Ca(2+)-dependent
621   apoptosis and necrosis. *Cell Calcium* 50(3):222-233.

622   22.   Blattner JR, He L, & Lemasters JJ (2001) Screening assays for the mitochondrial permeability
623   transition using a fluorescence multiwell plate reader. *Anal Biochem* 295(2):220-226.

624   23.   Pan X*, et al.* (2013) The physiological role of mitochondrial calcium revealed by mice lacking
625   the mitochondrial calcium uniporter. *Nat Cell Biol* 15(12):1464-1472.

626   24.   Paillard M*, et al.* (2013) Depressing mitochondria-reticulum interactions protects
627   cardiomyocytes from lethal hypoxia-reoxygenation injury. *Circulation* 128(14):1555-1565.

628   25.   Petersen CE*, et al.* (2023) Increased mitochondrial free Ca(2+) during ischemia is suppressed,
629   but not eliminated by, germline deletion of the mitochondrial Ca(2+) uniporter. *Cell Rep*
630   42(7):112735.

631   26.   Gasmi A*, et al.* (2024) Berberine: Pharmacological Features in Health, Disease and Aging.
632   *Curr Med Chem* 31(10):1214-1234.

633   27.   Liang J, Huang X, & Ma G (2022) Antimicrobial activities and mechanisms of extract and
634   components of herbs in East Asia. *RSC Adv* 12(45):29197-29213.

635   28.   Di Marco G*, et al.* (2020) A High-Throughput Screening Identifies MICU1 Targeting
636   Compounds. *Cell Rep* 30(7):2321-2331 e2326.

637   29.   Payne R, Li C, Fernandez-Garcia E, Vais H, & Foskett K (2019) The MCU Inhibitor
638   Ds16570511 has Off-Target Effects on Mitochondrial Membrane Potential. *Biophysical*
639   *Journal* 116(3).

640   30.   Sawyer DB, Peng X, Chen B, Pentassuglia L, & Lim CC (2010) Mechanisms of anthracycline
641   cardiac injury: can we identify strategies for cardioprotection? *Prog Cardiovasc Dis*
642   53(2):105-113.

643   31.   Liu DQ*, et al.* (2019) Berberine protects against ischemia-reperfusion injury: A review of
644   evidence from animal models and clinical studies. *Pharmacol Res* 148:104385.

645   32.   Qing Y, Dong X, Hongli L, & Yanhui L (2018) Berberine promoted myocardial protection of
646   postoperative patients through regulating myocardial autophagy. *Biomed Pharmacother*
647   105:1050-1053.

648   33.   Wang Y, Liu J, Ma A, & Chen Y (2015) Cardioprotective effect of berberine against
649   myocardial ischemia/reperfusion injury via attenuating mitochondrial dysfunction and
650   apoptosis. *Int J Clin Exp Med* 8(8):14513-14519.

651   34.   Hu F*, et al.* (2024) Berberine inhibits excessive autophagy and protects myocardium against
652   ischemia/reperfusion injury via the RhoE/AMPK pathway. *Int J Mol Med* 53(5).

653   35.   Ding L*, et al.* (2021) Berberine improves dietary-induced cardiac remodeling by upregulating
654   Kruppel-like factor 4-dependent mitochondrial function. *Biol Chem* 402(7):795-803.

655   36.   Zhao GL*, et al.* (2016) Berberine protects rat heart from ischemia/reperfusion injury via
656   activating JAK2/STAT3 signaling and attenuating endoplasmic reticulum stress. *Acta*
657   *Pharmacol Sin* 37(3):354-367.

37.   Chu M*, et al.* (2018) Polypharmacology of Berberine Based on Multi-Target Binding Motifs.

Front Pharmacol 9:801.

38.   Sheth AI*, et al.* (2024) Targeting Acute Myeloid Leukemia Stem Cells Through Perturbation
of Mitochondrial Calcium. *Cancer Discov*.

39.   Zhang J*, et al.* (2022) Novel CaMKII-delta Inhibitor Hesperadin Exerts Dual Functions to
Ameliorate Cardiac Ischemia/Reperfusion Injury and Inhibit Tumor Growth. *Circulation*
145(15):1154-1168.

40.   Trott O & Olson AJ (2010) AutoDock Vina: improving the speed and accuracy of docking
with a new scoring function, efficient optimization, and multithreading. *J Comput Chem*
31(2):455-461.

41.   Lan C*, et al.* (2022) Inhibition of DYRK1A, via histone modification, promotes
cardiomyocyte cell cycle activation and cardiac repair after myocardial infarction.
EBioMedicine 82:104139.

 Figure Legends

Fig. 1. The discovery of Berberine as a novel MCU inhibitor. (A) Schematic of
screening workflow. (B) Snapshot of cytosolic (left, red) and mitochondrial (right,
green) Ca^{2+} transient images from time-lapse movies of representative HeLa cells
stained with Calbryte590 and 4mtGCaMP6 following 10 μM Histamine (His)
stimulation. Cells were pretreated with various concentrations of Berberine. The time
displayed on the images is in seconds. Scale bar, 50 μm. (C) Representative traces of
mitochondrial Ca^{2+} level (indicated by 4mtGCaMP6) in HeLa cells stimulated with 10
μM Histamine and pretreated with different concentrations of Berberine. Data are
shown as mean ± s.e.m. n = 40 cells for each condition. (D) Measurement of the
maximal amplitudes of the mitochondrial Ca^{2+} traces in (C). Data are shown as the
mean ± s.e.m. n = 40 cells for each condition. *P* value was analyzed using one-way
ANOVA. (E) Representative traces of cytosolic Ca^{2+} level (indicated by Calbryte590)
in HeLa cells stimulated with 10 μM Histamine and pretreated with different
concentrations of Berberine. Data are shown as the mean ± s.e.m. n = 40 cells for each
condition. (F) Measurement of the maximal amplitudes of the mitochondrial Ca^{2+}
traces in E. Data are shown as the mean ± s.e.m. n = 40 cells for each condition. *P*
value was analyzed using one-way ANOVA. (G) Traces of mitochondrial Ca^{2+} level
(indicated by fluorescence of 4mt-GCaMP6) in digitonin-permeabilized HeLa cells
upon the addition of Ca^{2+} under varying concentrations of Berberine. (H) Dose-
dependent inhibition of mitochondrial Ca^{2+} uptake activity by Berberine with an IC_{50}
of 2.202 μM.

Fig. 2. Berberine directly binds to MCU. (A) Chemical structure of Biotin-
Berberine. (B) Representative immunofluorescent images of HeLa cells treated with
Biotin-Berberine (green) and stained for mitochondria (COXIV, red). Scale bar, 10
μm. (C) Traces of normalized fluorescence intensity spatial profiles through the white
line shown in (B). (D) Western blot analysis from Biotin-Berberine pulldown assay
using HEK293T cell lysates. Cells were treated with Biotin-Berberine (10 μM, 2 h).
(E) Western Blot analysis from *in vitro* pulldown assay investigating interactions
between GST-MCU and Biotin-Berberine under varying concentrations of Berberine.
(F) Binding ability of Berberine to GST-MCU in MST assays.

Fig. 3. Berberine targets JML domain of MCU to inhibit Ca^{2+} uptake. (A)
Schematic representation of MCU truncation mutants utilized for domain-mapping
studies. Numerical values in parentheses correspond to the amino acid sequences
encompassed by each construct. Key structural domains are labeled: NTD (N-terminal
domain), CC (coiled-coil domain), OJMH (outer juxtamembrane helix), TM
(transmembrane helix), DIME (DIME motif, Ca^{2+} filter), JML (juxtamembrane loop),
and CTD (C-terminal domain). (B) Western blot analysis of Biotin-Berberine

pulldown assays using *in vitro* transcribed and translated SNAP-tagged MCU
truncation mutants as shown in (A). (C) Molecular docking model revealing the
interaction between human MCU and Berberine. (D) Western blot analysis of biotin-
berberine pulldown assays using *in vitro* transcribed and translated SNAP-tagged
MCU mutants (amino acids 275-295, encompassing the JML region). Wild-type
residues Y281, Y289, Y291, and A294 were mutated to AAAF, HAAF, and RAAF,
respectively. (E) Western blot analysis of endogenous MCU and exogenous MCU-
mCherry expression in HeLa stable cell lines with MCU wild-type (WT) and
MCU^{HAAF} mutants. (F) Traces of mitochondrial Ca^{2+} levels, as indicated by the
fluorescence intensity of 4mt-GCaMP6, in digitonin-permeabilized HeLa cells
following Ca^{2+} addition. The cells include control HeLa, and stable cell lines
expressing mcherry-tagged MCU^{WT}and MCU^{HAAF} mutants, under conditions with and
without 10 μM Berberine. (G) Measurement of the maximal amplitudes of the
mitochondrial Ca^{2+} traces in (F). Data are shown as the mean $\pm$ s.e.m. n = 45 cells for
each condition. *P* value was analyzed using two-way ANOVA.

Fig. 4. Berberine inhibits Ca^{2+} uptake by disrupting MCU-EMRE complex
assembly. (A) Western blot analysis depicting the co-immunoprecipitation of Flag-
tagged MCU with SNAP-tagged EMRE, MCU, MICU1, and MICU2 in HEK293T
cells, conducted with or without the presence of 10 μM Berberine for 2 hours. (B)
Western blot analysis of GST pulldown assays performed with *in vitro* transcribed
and translated Flag-tagged EMRE and recombinant GST-MCU, with and without the
addition of 2 μg Berberine. (C) Molecular dynamics simulations showing the radius
of gyration (Rg) of the MCU-EMRE complex in the presence and absence of
Berberine. (D) Molecular dynamics simulations depicting the root-mean-square
deviation (RMSD) of the MCU-EMRE complex with and without Berberine. (E)
Western blot analysis of the co-precipitation of Flag-tagged MCU wild-type (WT) or
Flag-tagged MCU^{HAAF} mutant with SNAP-EMRE in HEK293T cells, treated with 10
μM Berberine for 2 hours or untreated. (F) Proposed model for Berberine-mediated
MCU gating. Left panel, EMRE binds to MCU and stabilizes the luminal gate of
MCU in the open conformation. Right panel, Berberine integrates into the MCU,
inducing a closed conformation that disrupts the assembly of the MCU-EMRE
complex and seals the MCU's luminal gate.

Fig. 5. Berberine protects against myocardial ischemia/reperfusion injury
through inhibiting mitochondrial Ca^{2+} overload. (A) Mitochondrial swelling in
response to Ca^{2+} challenge (500 μM CaCl$_2$). Swelling is inhibited with Berberine (10
μM), Ru360 (3 μM) or CsA (3 μM). (B) Representative images of neonatal rat
cardiomyocytes expressing 4mtGCaMP6. Cells were either treated with 10 μM
Berberine or not, then subjected to hypoxia (3 h) followed by reoxygenation (12 h)
(H/R). Fluorescence at ex488 (green) represents the Ca^{2+}-dependent signal, while
fluorescence at ex390 (blue) indicates the Ca^{2+}-independent signal. The ratio

(pseudocolour) of ex488/ex390 indicates the level of mitochondrial Ca^{2+}. Scale bar,

10 μm. (C) Measurement of mitochondrial Ca^{2+} level indicated by 4mtGCaMP6

fluorescence ratio (ex488/ex405) in (B). Data are shown as the mean ± s.e.m. n = 50

cells for each condition. *P* value was analyzed using one-way ANOVA. (D) The

experimental scheme of the mice with acute I/R injury (45 minutes ischemia followed

by 24 hours reperfusion) with vehicle or Berberine pretreatment (24 hours and 15

minutes before the onset of ischemia). (E) Evans Blue and 2, 3, 5-triphenyltetrazolium

chloride (TTC) staining of myocardium after I/R injury. The blue area indicates non-

ischemic area. The red indicates viable tissue at risk. The white indicates infarct area.

(F) Quantification of infarct size (IF), area at risk (AAR) of mice subjected to acute

I/R injury (as protocols in D). n = 9 or 10 mice. (G-I). Cardiac cell death indexed by

serum cTnI (G, n=12 [sham], 6 [sham+BBR], 11 [I/R], and 12 [I/R+BBR]), serum

LDH concentration (H, n=13 [sham], 6 [sham+BBR], 11 [I/R], and 12 [I/R+BBR]),

and myocardial TUNEL-positive cells (I, n=8 [sham], 6 [sham+BBR], 11 [I/R], and

12 [I/R+BBR]) of the mice subjected to I/R injury (as the protocols in D) with vehicle

or Berberine treatment (5 mg/kg, ip.24 h and 15min before I/R injury). *P* value was

analyzed using one-way ANOVA. Scale bar, 50 μm.

Supplementary Material

Fig. S1. Screening identifies Berberine as an inhibitor of mitochondrial Ca^{2+}
uptake. (A) Histogram plot of the 2,816 FDA-approved drug library based on the
free-energy docking score with MCU. (B) Traces of mitochondrial Ca^{2+} level
(indicated by fluorescence of 4mt-GCaMP6) in digitonin-permeabilized HeLa cells
upon the addition of Ca^{2+} in the presence of DMSO, Ru360, or drugs. (C) Graphical
representation of drug screen results. 'Z-score' ranks drugs based on maximal
amplitudes of the traces in B. (D) The list of the top 7 compounds with mitochondrial
Ca^{2+} uptake inhibitory activity. (E) Measurement of mitochondrial membrane
potential indicated by TMRM fluorescence intensity. Data are shown as the mean $\pm$
s.e.m. n = 50 cells for each condition. *P* value was analyzed using one-way ANOVA.

Fig. S2. Berberine selectively inhibits mitochondrial Ca^{2+} uptake without
impairing cytosolic Ca^{2+} signaling, mitochondrial membrane potential, or cell
viability. (A) Representative traces of cytosolic Ca^{2+} level (indicated by GCaMP6) in
10 μM Histamine-stimulated HeLa cells pretreated with different concentration of
Berberine. Data are shown as the mean $\pm$ s.e.m. n = 50 cells for each condition. (B)
Measurement of the maximal amplitudes of the cytosolic Ca^{2+} traces in A. Data are
shown as the mean $\pm$ s.e.m. n = 50 cells for each condition. *P* value was analyzed
using one-way ANOVA. (C) Representative traces of mitochondrial Ca^{2+} level
(indicated by 4mt-GCaMP8) in 10 μM Histamine-stimulated HeLa cells pretreated
with different concentration of Berberine. Data are shown as the mean $\pm$ s.e.m. n = 50
cells for each condition. (D) Measurement of the maximal amplitudes of the
mitochondrial Ca^{2+} traces in C. Data are shown as the mean $\pm$ s.e.m. n = 50 cells for
each condition. *P* value was analyzed using one-way ANOVA. (E) Flow cytometry
analysis of TMRM staining in HeLa cells pretreated with different concentrations of
Berberine and 50 μM CCCP. Shown is one experiment that is representative of three
similar experiments. (F) Quantitative analysis of relative fluorescence of TMRM
staining after different concentrations of Berberine treatment and 50 μM CCCP
treatment. Data are shown as the mean $\pm$ s.e.m. n = 3 independent experiments. (G)
Cell viability of HeLa cells, U2OS cells and HEK293T cells after different
concentration of Berberine treatment was assessed at indicated time and normalized to
day 0 in each group. Data are shown as the mean $\pm$ s.e.m. n = 3.

Fig. S3. Berberine directly targets MCU. (A) Representative images of Berberine
(left panel) treated HeLa cells co-stained with MitoTracker DeepRed (middle panel).
The merged image is shown in the right panel. Scale bar, 10 μm. (B) Traces of
normalized fluorescence intensity spatial profiles through the white line shown in A.
(C) Traces of mitochondrial Ca^{2+} level (indicated by fluorescence of 4mt-GCaMP6)

in digitonin-permeabilized HeLa cells upon the addition of Ca^{2+} in the presence of
different concentration of Berberine or Biotin-Berberine. Data are shown as the mean
± s.e.m. n = 40 cells for each condition. (D) Measurement of the maximal amplitudes
of the mitochondrial Ca^{2+} traces in C. Data are shown as the mean ± s.e.m. n = 40
cells for each condition. *P* value was analyzed using two-way ANOVA. (E) Western
Blot analysis of Biotin-Berberine pulldown assay in MCU-Flag overexpressed
HEK293T cells. Cells were treated with 2 h 10 μM Biotin or 10 μM Biotin-Berberine.
(F) Two-dimensional diagrams illustrating the binding mode of berberine with MCU.
(G) Sequence alignment of the putative region (amino acids 275-295) surrounding the
JML domain of MCU proteins from 16 diverse species.

Fig. S4. Berberine disrupts MCU-EMRE assembly. (A) Western Blot analysis of
co-precipitation of the Flag-EMRE with SNAP-MCU in HEK293T cells in the
presence of 10 μM Berberine or not for 2 hours. (B) Molecular dynamics simulations
illustrating the assembly conformations of the MCU-EMRE complex in the presence
and absence of Berberine. (C) Molecular dynamics simulations reveal the hydrogen
bonding interactions between MCU and EMRE under conditions with and without
Berberine. (D) Molecular dynamics simulations displaying the number of hydrogen
bonds between MCU and EMRE in the presence and absence of Berberine. (E)
Molecular dynamics simulations demonstrating the binding free energy between
MCU and EMRE with and without Berberine. The components of the binding free
energy are detailed as follows: van der Waals energy (VDWAALS), electrostatic
energy (EEL), polar solvation energy (EGB), non-polar solvation energy (ESURF),
total gas phase free energy (GGAS, calculated as the sum of VDWAALS and EEL),
total solvation free energy (GSOLV, calculated as the sum of EGB and ESURF), and
the total free energy (TOTAL, calculated as the sum of GSOLV and GGAS).

Fig. S5. Berberine attenuates mitochondrial Ca^{2+} overload in cardiomyocytes
under pathological conditions. (A-B) Measurement of mitochondrial Ca^{2+} level
(indicated by Rhod2) in H9C2 cells. Cells were pretreated with 10 μM Berberine or
not, and then were subjected to hypoxia (3 h) followed by reoxygenation (12 h) (H/R).
Relative fluorescence intensity of Rhod2 were measured in B. Data are shown as the
mean ± s.e.m. n = 30 cells (Control and BBR groups), n = 40 cells (H/R and
H/R+BBR groups). *P* value was analyzed using one-way ANOVA. Scale bar, 10 μm.

Fig. 1. The discovery of Berberine as a novel MCU inhibitor.

Fig. 2. Berberine directly binds to MCU.

Fig. 3. Berberine targets JML domain of MCU to inhibit Ca²⁺ uptake.

Fig. 4. Berberine inhibits Ca²⁺ uptake by disrupting MCU-EMRE complex assembly.

Fig. 5. Berberine protects against myocardial ischemia/reperfusion injury through inhibiting mitochondrial Ca²⁺ overload.