1 Chaparral wildfire shifts the functional potential for soil pyrogenic organic
2 matter and nitrogen cycling
3
4
5
6

7 M. Fabiola Pulido-Chavez[1,*], Amelia R. Nelson[2], Michael J Wilkins[2], Sydney I.
8 Glassman[1,*]
9
10
11
[1]Department of Microbiology and Plant Pathology, University of California-
Riverside, 900 University Ave, Riverside, CA, 92521
[2]Department of Soil and Crop Sciences, Colorado State University, 307
University Ave, Fort Collins, CO 80521
[3]Department of Environmental Sciences, University of California-Riverside, 900
University Ave, Riverside, CA, 92521
[41]Department of Ecology and Evolutionary Biology, University of California-
Irvine
20
21
22
23 *Corresponding authors: Sydney I Glassman, sydneyg@ucr.edu, and M. Fabiola
24 Pulido-Chavez, fabspulido@gmail.com
25
26 Keywords: Pyrophilous bacteria, PyOM, Nitrogen,

Abstract

Wildfires alter soil chemistry and microbial communities, opening niche

space for pyrophilous or "fire-loving" microbes to initiate secondary succession.

However, wildfires also increase soil nitrogen (N) and transform woody

vegetation into difficult-to-degrade pyrogenic organic matter (PyOM), a

potential carbon (C) source for pyrophilous microbes. Whether pyrophilous

microbes can degrade these post-fire resources has significant implications for

global C and N cycling, including C sequestration and greenhouse gas emissions.

To determine temporal patterns in the functional potential for PyOM degradation

and N cycling, we performed shotgun metagenomics on 30 burned and unburned

soils collected at 17, 25, 34, 131, and 376 days after a high-severity wildfire in a

Southern California fire-adapted chaparral shrubland. Not only did overall

profiles of C and N cycling genes differ between burned and unburned plots, but

burned plots showed dramatic increases from 17 to 376 days in genes responsible

for degrading PyOM (167%) and cycling N (117%). In contrast, gene

abundances for PyOM degradation and N cycling remained stable over time in

unburned plots. Genes encoding the degradation pathways for two PyOM

degradation intermediate pathways, catechol and protocatechuate, revealed that

the easier-to-degrade ortho-cleavage pathways dominated all time points in the

burned plots. Genes encoding N retention pathways (assimilatory and

dissimilatory nitrate reduction) dominated N cycling pathways in burned plots

for the entire year. Finally, dominant pyrophilous bacterial taxa in the genera

Massilia and *Noviherbaspirillium* encoded different PyOM and N processing

pathways, suggesting that traits enabling adaptation to wildfires, like post-fire

resource acquisition, drive microbial secondary succession.

Introduction

Soil microbial communities support fundamental post-fire soil processes

such as N immobilization thereby facilitating N storage (1) and C cycling

through decomposition, C sequestration (2) and C loss as CO_2 during respiration

(3). However, understanding the complex functional dynamics of the post-fire

soil microbiomes remains challenging. With shifting fire regimes resulting in

increased fire frequency and severity (4), understanding early changes in the

functional potential of post-fire soil microbial communities is necessary to

determine how microbes drive secondary successional dynamics and alter

important ecosystem functions such as those that affect C sequestration and

greenhouse gas (GHG) emissions.

Wildfires often cause reductions in microbial richness, shifting

community composition and enabling pyrophilous or "fire-loving" microbes to

initiate secondary succession (5-8). Bacterial succession following high-severity

wildfires are initiated by the Firmicute and Actinobacteria phyla in various

ecosystems including, Southern California Chaparral (8), and redwood-tanoak

(5), coniferous (9), and boreal forests (6). At the same time, pyrophilous

microbes can trigger a change in resource use that can affect microbial

successional dynamics, including successional directionality and community

metabolism (10, 11). We previously hypothesized that post-fire microbial

succession is initiated by thermotolerators, followed by fast-growing ruderal

taxa, who are later supplanted by taxa that capable of capitalizing on post-fire

resources, for example bacteria in the genera *Noviherbaspirillium* and

*Blastococcu*s (8).This implies that early post-fire taxa might perform metabolic

functions that prime the system for colonization by more diverse and complex

bacterial species with more specialized metabolic functions, potentially resulting

in a new microbial equilibrium. However, whether these simultaneous functional

changes are occurring and subsequently affecting microbial successional

dynamics is unknown.

Fires primarily alter the C cycle through the combustion of vegetation

(12) and producing abundant PyOM, an aromatic and chemically complex C

source known for its resistance to microbial degradation (13, 14). The

aromaticity of PyOM requires microbes to encode genes for cleaving aromatic

rings thereby metabolizing and acquiring C (15). Bacteria in the genera *Massilia*

and *Noviherbaspirillium* are considered putative PyOM degraders (16), and the

abundant pyrophilous fungus *Pyronema domesticum* was found to degrade

PyOM in laboratory studies (17). Therefore, it is likely that other pyrophilous

microbes have the genomic potential to degrade this abundant post-fire C source.

However, wildfires can also increase the abundance of short-chain alkanes (18-

20), such as propane, which are easier to degrade C sources preferentially

selected by microbes for consumption (21-23). However, whether early

successional post-fire microbes have the functional potential to degrade PyOM

over alkanes or whether C source selection changes with succession is currently

unknown.

In addition to altering the C cycle, fires alter the N cycle through N losses

via volatilization, reduced urea hydrolysis (24), and a short-term enrichment of

bioavailable N (25-27). Indirect wildfire effects, such as removal of the

vegetation, can also increase soil NO_3^- and NH_4^+ (27, 28), increase soil

temperature, and decrease soil moisture, resulting in increased aerobic conditions

that reduce nitrification (27, 29) and can further impact the microbial

community. While microbial immobilization—the uptake of NO_3^- and NH_4^+ for

microbial biomass—can decrease denitrification by limiting NO_3^- availability, it

can also lead to NO_2 emissions (30, 31). Despite this, denitrification and urea

hydrolysis still occur, potentially producing N_2O, a potent greenhouse gas (30-

34). Consequently, understanding how pyrophilous microbes treat the increased

post-fire bioavailable N has strong implications for GHG emissions. Moreover,

post-fire microbial communities likely exhibit varying responses to N depending

on the ecosystem type. For example, research in Spanish shrublands three years

post-fire, demonstrated increased relative abundances of genes associated with

NH_3 assimilation, ammonification, and N_2 fixation (35). In contrast, forested

ecosystems in North America showed an increased genomic potential for

nitrification and a decrease in denitrification during the first decade after the fire

(36). Time since fire also alters the processing of post-fire N, highlighting the

complexity of predicting the impact of post-fire microbial succession on N

cycling and trace and GHG emissions over time.

We previously sampled a chaparral wildfire at high temporal resolution

and showed rapid secondary successional dynamics in the first year post-fire (8).

For bacteria, we observed six major compositional turnover events, and we

suggest that underlying community metabolic changes might be driving

secondary succession (8). Specifically, thermotolerators in the Firmicutes

(*Bacillus, Paenibacillus,* and Clostridiales) initially dominated, but were

replaced later in the year by taxa with putative N and PyOM degrading

capabilities, including *Massilia* and *Noviherbaspirillum* (8) and potential PyOM

degrative capability of *Blastococcus* (37). From this dataset, we selected a subset

of burned and unburned soils from 5 time points (17, 25, 34, 131, and 376 days

post-fire) corresponding to high microbial compositional turnover. We

conducted a gene-centric metagenomic analysis to determine how temporal

changes in pyrophilous communities relate to post-fire bacterial functions and

successional dynamics. Specifically, we examined the relative abundance of

genes involved in driving PyOM degradation and N cycling over time. We tested

the hypotheses that the genes encoding potential for H1) degradation of wildfire-

induced C sources like PyOM and alkane and H2) processing of abundant

bioavailable N will increase over time in burned plots. We further hypothesized

H3) that these changes in gene abundances would correspond with bacterial

community shifts and the increased relative abundance of pyrophilous bacteria

with the putative abilities to use high post-fire abundances of PyOM and N.

Methods

Sampling

In August-September 2018, the Holy Fire burned 94 km^2 of Manzanita

(*Arctostaphylos*) dominated Chaparral shrublands in the Cleveland National

Forest in Southern California. Seventeen days post-fire, we established nine plots

(6 burned and 3 unburned), selecting for similarity in pre-fire vegetation,

elevation, and aspect. Each plot contained four 1 m^2 subplots in each cardinal

direction for temporal soil sampling of the top 10 cm of mineral soil beneath the

ash or the duff (Fig. S1). Soil samples were collected at nine time points within

the first-year post-fire (17, 25, 34, 67, 95, 131, 187, 286, and 376 days). DNA

was extracted from 0.25g of 2mm sieved soil and we performed 16S and ITS2

Illumina MiSeq sequencing of all samples (8). From this data, we selected 5 time

points (17, 25, 34, 131, and 376 days post-fire) with extreme bacterial

compositional turnover events, and representative burned and unburned plots

with high enough quantity and quality DNA for metagenomic sequencing.

Metagenomic sequencing

From each of the 5 timepoints, we selected 3 burned and 3 unburned

subplots for a total of 30 metagenomes (15 burned and 15 unburned). Samples

were chosen by selecting the DNA samples with the following criteria, 2.5 ng/µl

and 230/260 above 1.8, from unburned plot 8 (Fig. S1a) and from two closely

situated burned plots with the most similar microbial richness and biomass (plots

2 and 3; Fig. S1a). Metagenomic library preparation and sequencing were

performed at the Department of Energy Joint Genome Institute (JGI) in

Berkeley, California, for 17 samples and at the University of California, Irvine

(UCI) Genomics High-Throughput Facility for 13 samples. Libraries were

prepared from the extracted DNA using the Kapa Biosystems library preparation

kit at JGI and the Illumina Nextera Flex library preparation kit at UCI and were

sequenced on the Illumina NovaSeq 6000 platform on an S4 flow cell (151 bp

paired-end reads and a depth of 25 Gb/sample). See supplementary data 1 for

additional information regarding metagenomic sequencing.

Metagenomics binning

Sequence adapters were removed from the raw reads using BBduk-
v38.89 (sourceforge.net/projects/bbmap/) with the following parameters
(ktrim=r, k=23, mink=11, hdist=1). Raw reads were trimmed with Sickle – v1.33
(38) , and the quality of trimmed reads was assessed with FastQC – v0.11.2 (39).
Trimmed reads were assembled into contiguous sequences (contigs) with the de
novo de Bruijn assembler MEGAHIT – v1.2.9 (40) with the following
parameters (min kmer of 27, max of 127, and increment of 10). Per-contig
coverage per sample was calculated using CoverM contig version v0.6.0 (41)
with the trimmed mean method to retain only mappings >95% identity and
minimum alignment of 75% (42). Assembled contigs (>2,500 bp) were binned
using MetaBAT2 v2.12.1 with default parameters (43) to assemble metagenome-
assembled genomes (MAGs). MAG quality was estimated using checkM v1.1.2
(42) and taxonomy was assigned using GTDB-Tk v2.1.1 (44). To create the
final medium and high-quality MAG database, low-quality MAG were removed
and the dataset was dereplicated using dRep v3.0.0 (45). See supplementary data
1 for additional information regarding metagenomic assemblies.

Contig annotation

To assess the degradation of aromatic hydrocarbons, such as PyOM, we
focused our PyOM contig annotation on catechol and protocatechuate, two
intermediate aerobic degradation pathways (46) and short-chain alkane C as the

easier to degrade C source. To improve retention and annotation of the samples

in a comprehensive manner, all assembled contigs were annotated using three

different databases. First, contigs were annotated using HMMER (47), with

hidden Markov models (HMMs) against Kofamscan HMMs (48) to identify

specific genes related to aromatic compound degradation (i.e., catechol and

protocatechuate and inorganic N cycling) according to previously established

methods (11). Second, we used the Calgary approach to AnnoTating

HYDrocarbon (CANT-HYD) degradation genes, which uses HMMs for

annotating marker genes involved in aerobic and anaerobic hydrocarbon

degradation pathways (49). Third, we used the Distilled and Refined Annotation

of Metabolism, DRAM (50) annotation tool to profile our contigs through

multiple databases (i.e., KEGG (51), UniRef90 (52), MEROPS (53), and

Mmseqs2 (54) with the best hits (based on a bit score of 60) reported for each

database. The contig dataset was normalized by calculating the contig length

corrected trimmed mean (geTMM) of M values (55) using EdgeR (56),

normalizing by library depth and gene length as has been previously used for

metagenomics data (57-59). The results from all three annotation methods were

concatenated into one large file, and genes related to PyOM, alkane degradation,

N, and urea cycling were selected for all downstream analyses. In the case where

contigs contained duplicated annotations, the annotations were manually curated,

and the best functional annotation was determined by selecting the annotation

with the lowest e-value.

We used results from a DESeq analysis (see details below) to identify

genes of interest, then used NCBI BLAST (60) to link the MAG catalog to these

dominant genes. We filtered the BLAST matches to include only hits with 100%

identity and a bit score of at least 100. Therefore, we identified the specific taxa

that encoded key genes, including those related to PyOM and N cycling

pathways. For a comprehensive overview illustrating the bioinformatic

procedures employed, please refer to the Fig. S2.

Statistical analysis

All statistical analyses and figures were done in R version 4.2.3 (61).

First, we determined how fire, time, and their interaction impacted the overall

composition of genes for (H1) PyOM or (H2) N cycling with Permutational

Multivariate Analysis of Variance (PERMANOVA;(62)) as implemented in the

adonis2 function of the vegan package (63) with 9999 permutations with plot as

random effect. Results were visualized using nonmetric multidimensional scaling

(NMDS). Ordinations were based on Bray–Curtis distance matrixes using the

ordinate function in the phyloseq package (64). Since alkanes increase post-fire

and are preferentially selected by microbes as a C source, we separated alkanes

from the PyOM genes for all downstream analyses, allowing us to determine if

this preferential uptake also occurs in post-fire systems. Moreover, since urease

is vital to hydrolyze urea into ammonia (NH_3^+) (65), an essential process that

makes N readily available for plant uptake, we further separated the urea

pathway from the N cycling processes. Separation of urea from N pathway,

allows us to determine the effect of fire on these important ureases and thus

helping us understand if the post-fire microbes possess ureases which can help us

better understand post-fire ecosystem restoration and N cycling.

Second, to determine if bacterial functional changes mirrored the rapid

compositional changes we previously observed, we plotted the top most

abundant bacterial genera in bar plots at 17, 25, 34, 131, and 376 days post-fire

(8). We also calculated the homogeneity of variance and used permutest with

9999 permutations and plot as random effect to determine which time points

differed for burned and unburned pathways independently. We used principal

coordinates analysis (PCoA) to visualize the differences in the homogeneity of

variance of each functional pathway with the vegan package using Bray-Curtis

dissimilarities.

Third, we independently performed generalized negative binomial

regression on burned and unburned plots for each pathway to test how the

number of PyOM, alkane, N, and urea cycling genes changed over time. The

nestedness level was tested using null model and Akaike criteria. Burned PyOM,

alkaline, and urea models included plot, and unburned models included subplot

as random effect. Burned N models required plot, subplot, and time, and

unburned N models required time and subplot as random effects. For all models,

time was scaled and centered. We visualized how the microbial functional

potential changes over time by plotting the sum geTMM of each PyOM, alkane,

N, and urea pathway using phyloseq version 1.42.0 (64) for each time point in

burned and unburned communities independently.

Lastly, to identify the PyOM, alkane, N, and urea genes that were

differentially overrepresented in the burned versus the unburned plots at each

sampling time point, we employed DESeq2 v1.38.3 (66). Five separate models in

DESeq2 were defined, one per timepoint, to assess the response of functional

gene potential and independently identify the difference in response to fire over

time for each pathway. Results are based on the DESeq Wald test (adjusted p-

value < 0.05 and absolute log2 fold-change >0). The total count of differentially

abundant genes was visualized as a bar plot for each independent pathway over

time. Moreover, we used the differentially abundant genes at each time point in

the burned communities (p_{adj}<0.05) to create a metabolic map for PyOM and N

degradation pathways. Furthermore, to link the most abundant genes to their

taxonomy, we first selected the genes of interest (most abundant genes based on

DESeq results at each time point, log2fold change >0 and p < 0.05) and linked

them to their respective taxonomy using our MAG dataset using BLAST

(Bethesda, 2008). Taxonomy and functional annotation of the selected group of

genes were plotted using ggplot2. All statistical scripts (R scripts) are publicly

available on GitHub: https://github.com/pulidofabs/Chaparral-Metagenomes.

Results

We assembled 1.2 Tb of metagenomics sequencing data, resulting in the

annotation of 2,266,623 genes relevant to post-fire microbial metabolisms.

Identified taxa represented across all MAGs belonged to the phyla

Acidobacteriota, Proteobacteria, and Verrucomicrobiota *(8)*. Additional Of

particular interest, we assembled MAGs of pyrophilous bacteria in the genera

Massilia and *Noviherbaspirillium*, two of the most dominant genera in our

chaparral study(8), contig information, MAG quality and taxonomy information,

and contig taxonomy information for the dominant selected genes are available

in Supplementary Data 1C.

Wildfire altered the profile of functional genes for PyOM and N

Fire significantly impacted the profile of functional genes associated with

N and PyOM cycling, leading to significant changes over time (Fig. 1; Table S1).

Wildfire was the strongest driver of differences in functional potential,

explaining 15% of the variation for genes affiliated with the processing of PyOM

(Fig. 1a) and urea (Fig. 1d) and 16% for alkanes (Fig. 1b), and the remaining N

cycling genes (Fig. 1c; Table S1). Time had smaller but significant impacts on

the functional profiles of these genes, explaining 5% of the variance in PyOM,

N, and urea cycling genes and 6% for alkane genes (Table S1). A significant fire-

by-time interaction explained 5% of the variation for PyOM and urea and 6% for

N but was not significant for alkanes (Table S1).

While there was no change in the turnover or community-level alterations

of the profile of functional genes for PyOM (Fig. 2a), alkane degradation (Fig.

2c), N (Fig. 2e) or urea cycling (Fig. 2g) in the unburned plots (betadisper

p>0.05), the burned plots experienced multiple functional profile turnover

periods for PyOM (Fig. 2b), alkane (Fig. 2d), N (Fig. 2f) and urea cycling genes

(Fig. 2h; beta-disper < 0.05; Table S2). Overall, genes associated with urea

processing experienced more turnover events (at 25, 34, and 131 days post-fire)

than those associated with the degradation of PyOM, N cycling (34 and 131

days) or alkane degradation (25 and 131 days; Fig. 2).

Time differentially affected PyOM degradation pathways

There was a significant and positive fire by time interaction such that the

geTMM normalized gene abundances associated with PyOM or alkane

processing increased over time in the burned (Fig. 3b,d), but not unburned plots

(Fig. 3a,c; Table S3; S4). This increase is attributed to the increasing number of

PyOM genes (Fig. S3), which correlates with increased bacterial diversity and

abundance over time (Fig. S4b). Moreover, time differentially affected the

different steps of PyOM (Fig. 3b; Table S5) and alkane cycling pathways (Fig.

3d; Table S6) in burned plots such that we detected temporal increases in

geTMM values associated with catechol and protocatechuate cycling (Fig. S5;

Table S5). However, the β-ketoadipate and all alkane cycling pathways (i.e.,

alkane and propane pathways) remained unchanged over time (Fig. S5; Table

S6).

Time differentially affected N degradation pathways

Similarly, wildfire also had a significant fire by time interaction effect for

N and urea such that the geTMM values increased over time in burned (Fig. 4b,

d) but not unburned plots (Fig. 4a,c), when compared to 17 days post-fire (Table

S3,S4). This increase in geTMM values is due to increased total number of N

cycling genes (Fig. S6), which could be due to increases in bacterial dominance

over time (Fig. S4). Moreover, time differentially affected the various steps of

the N cycling pathways, resulting in increased geTMM values for NO_3^-

assimilation (both assimilatory and dissimilatory) and nitrification in the burned

plots compared to 17 days post-fire ($p < 0.04$; Fig. 4b; Fig. S7, Table S7).

However, denitrification was unaffected by time in the burned plots but did

significantly decrease in unburned plots over time (Fig. S7, Table S7). For the

urea cycle pathways, both the urea and ornithine geTMM values significantly

increased over time in burned plots, whereas there was no effect of time in the

unburned plots (Table S8).

Wildfire selects for ortho-cleavage aromatic degradation pathways

Wildfire resulted in a higher number of PyOM degradation genes that

were significantly differentially abundant within the burned communities than

the unburned communities with particular increases at 131 days post-fire (692

compared to 258; $p_{adj} < 0.05$; Fig. 5a). However, the functional potential for

PyOM degradation genes differed across the burned plots over time ($p_{adj} < 0.05$;

Fig. 5a, 6). Overall, the ortho-cleavage pathway for the protocatechuate and

catechol pathways and the β-ketoadipate pathway were differentially abundant in

the burned plots throughout the year (17-131d; $p_{adj} < 0.05$; Fig. 5a, 6). In contrast,

the meta-cleavage pathways were only differentially abundant in the early post-

fire timepoints, with catechol at 17-25 days and protocatechuate at 17 and 131

days post-fire (p_{adj} <0.05; Fig. 5a, 6). While PyOM degradation genes were

detected in the unburned plots at 17-34 days post-fire, it's important to note that

these genes are specifically related to lignin degradation (e.g., *ligABCIJKM,*

galC; Fig. 5a, 6). Moreover, alkane degradation pathways, propane 2-

monooxygenase, alkane hydroxylase, and alkane monooxygenase were only

overrepresented in the burned plots at 17d post-fire (p_{adj} <0.05; Fig. 5b).

Wildfire increased genes for N retention

371   Wildfires resulted in more differentially abundant N cycling genes within

the burned communities than the unburned communities (143 compared to 55;

Fig. 5c, 7). At all time points, the genes encoding N retention pathways, such as

assimilatory NO_3^- (*nasABC, NRT, narK, nrtP*, and *nirA*) and dissimilatory NO_2^-

(*nirBD, narBlVHY, and nxrB*) reductase and NO_3^- assimilation (*narG, narZ,*

nrxA) pathways were differentially abundant in the burned microbial

communities (p_{adj}<0.05; Fig. 5c, 7). In contrast to the unburned plots, where urea

cycling pathways were only overrepresented at 17 and 25 days, in the burned

plots, all urea degradation pathways were significantly abundant at 25, 34, and

131 days post-fire (p_{adj}<0.05; Fig. 5d,7).

Genomic potential of dominant taxa differed in burned and unburned

communities

384   The genomic potential for catechol and protocatechuate ortho-cleavage

and the β-ketoadipate pathway in the burned plots (*fad* genes) were

overrepresented in MAGs representing the phyla Proteobacteria, Actinobacteria,

and Acidobacteriota (Fig. 8*; Supplementary file 1c). Pathway genes for catechol

ortho-cleavage (*CatABC)* and β-ketoadipate pathway (*pcaDLJ, fadA)* were

differentially encoded by MAG Holy_395.fa, representing the pyrophilous

Proteobacteria *Massilia* (Fig. 8), which dominated the 16S sequences in the

burned plots across all time points (Fig. S4b). The protocatechuate ortho-

cleavage pathway (*pcaBCH*) was encoded by MAGs representing the

Burkholderiales-SG8-41 Solirubrobacteraceae (MAG Holy_206.fa) and the

pyrophilous bacterial genus *Noviherbaspirillium* (Fig. 8, MAG Holy_399.fa),

which dominated at 131 and 376 days post-fire in the burned plots (Fig. S4b).

Additionally, genes related to the catechol meta-cleavage pathway (*dmpBC*,

xylEG) were encoded by the MAGs affiliated with the bacterial families

Xanthobacteraceae-PALSA-894 (MAG Holy_369.fa) and Chthoniobacterales-

AV80 (*CatE*, MAG Holy_436.fa; Fig. 8). Furthermore, protocatechuate meta-

cleavage pathway genes (*ligClK, galC*) were differentially abundant in MAGs

representing the dominant *Noviherbaspirillium* (MAG Holy_399.fa) and

Xanthobacteraceae-PALSA-894 (MAG Holy_368.fa and Holy_369.fa; Fig. 8).

Although catechol, protocatechuate, and β-ketoadipate genomic potential were

differentially abundant at 17 and 25 days post-fire in the unburned plots, the

genes and the genera differed and were driven by the MAGs representing

Bradyrhizobium (MAG Holy_362.fa) and Acidimicrobiales-*AC-14* (MAG

Holy_71.fa; *pcaD, fadAl; praC, xylH, pracH, ligABC*; Fig. 8).

The genes encoding NO_3^- retention pathways, both assimilatory NO_3^-

reductase (*nasABC, nirA*) and dissimilatory NO_3^- reductase (*nirBD*), were

differentially abundant in MAGs representing the family Actinobacteriota

Arthrobacter (*NasABC*; MAG Holy_91.fa)*,* Micromonosporaceae (*NasABC*;

NirBD; MAG Holy_114.fa) and *Pseudonocardia* (*NirBD*; MAG Holy_153.fa;

Fig. 8). The pyrophilous *Massilia* (MAG Holy_395.fa) also encoded genes for N

retention, both dissimilatory NO_3^- reductase (*nirD*) and NO_3^- assimilation (*NRT,*

narK, nrtP, nasA; Fig. 8). Furthermore, genes associated with NO_3^- assimilation

retention pathways (*NRT, narK, nrtP, nasA*) were differentially abundant within

the Actinobacteria, specifically in the genus *Arthrobacter* (MAG Holy_91.fa)

and family Micromonosporaceae (MAG Holy_114.fa; Fig. 8). Like PyOM

genes, NO_3^- assimilation retention pathways , reduction, and denitrification in the

unburned plots at 17 days post-fire were differentially abundant in the non-

pyrophilous bacterial genus *Bradyrhizobium* (MAG Holy_362.fa and

Holy_361.fa; *NRT, narK, nrtP, nasAC, napA*; Fig. 8).

Discussion

We found that high-severity fire dramatically altered the genomic

functional potential of the post-fire soil microbiome, concurrent with large

changes in the bacterial community composition (Pulido-Chavez et al., 2023).

Wildfire increased the geTMM of PyOM degradation genes in burned plots over

time, and our analyses revealed that catechol and protocatechuate ortho-cleavage

(i.e., β-ketoadipate) pathways are the preferred pathway for PyOM degradation.

We also found that wildfire increased the geTMM values for N cycling genes in

burned plots over time, selecting for genes encoding the N retention pathways,

via conversion of NO_3^- to biomass and NO_2 to NH_3^+, with significant

implications for trace and GHG emissions. Finally, we found that pyrophilous

bacterial taxa that dominated our post-fire chaparral communities (i.e.,

Burkholderiales, *Massilia,* and *Noviherbaspirillium*) (Pulido-Chavez et al., 2023)

encoded multiple but different PyOM degradation and N cycling pathways,

suggesting that traits enabling adaptation to wildfires like post-fire resource

acquisition drive microbial succession.

PyOM geTMM values increased over time in burned plots

We previously observed patterns of secondary succession in chaparral

and hypothesized that these changes aligned with the ability of post-fire

microbes to use post-fire resources (8). Here, we found that the normalized

abundance of genes encoding for PyOM degradation increased over time and

were differentially abundant in the burned plots. Given the previously observed

increases in bacterial species richness and diversity over time ((8), Fig. S4), and

the concurrent increase in total number of differentially abundant genes in this

study, it is plausible that the elevated geTMM is associated with the emergence

of additional bacterial species encoding functional potential for PyOM

degradation.

We hypothesized that the succession of post-fire microbes is such that

thermotolerant microbes dominate first, followed by fast-colonizers, which are

then replaced by microbes capable of capitalizing on post-fire resources, such as

PyOM and N (8). Our results support those inferences and indicate that MAGs

representing pyrophilous bacteria Proteobacteria *Noviherbaspirillum* and

Massilia, which dominate in the first year post-fire, encode multiple genes for

PyOM and N degradation particularly at 131 days post-fire (Fig. 6). Indeed,

studies involving PyOM treatments (67-69) and incubation studies (70, 71) have

demonstrated an increase in the relative abundance of *Noviherbaspirillum* and

Massilia, indicating their active degradation of PyOM as a C source. In our study

(5), *Noviherbaspirillium* increased in abundance over time and dominated at 131

and 376 days post-fire, whereas *Massilia* dominated at all post-fire time points

but was particularly dominant at 34 and 131 days post-fire.

Here, we showed that *Massilia* encoded genes for PyOM degradation via

the catechol and β-ketoadipate ortho-cleavage pathway while

Noviherbaspirillium favored the protocatechuate ortho and meta-cleavage

pathways for PyOM degradation. The difference in their degradation pathways

may explain the delayed appearance of *Noviherbaspirillum* during succession, as

microbes using the meta-cleavage pathway often experience a growth lag and

delayed substrate utilization due to regulatory difficulties in inducing

degradation of intermediate PyOM pathways (72). In contrast, the dominance of

Massilia could be attributed to its genomic potential for rapid access to C via the

catechol ortho-cleavage pathway.

While we had previously shown successional patterns for post-fire fungal

communities ((5), Fig. S4), metagenomic analysis for fungi lags behind that of

bacteria, so no fungal-associated MAGs were detected in our data. However,

current research using whole transcriptome sequencing revealed that *Aspergillus*,

a dominant fungus across all of our post-fire timepoints (8), possesses some

genes involved in the β-ketoadipate pathway (73). Thus, it is possible that the

temporal increase in PyOM cycling genes observed in our study is partly due to

an increase in fungal diversity over time. However, further analyses are required

to better develop this hypothesis.

Fire also selects for microbial taxa that encode genes for alkane

degradation. Previous studies have shown that extractable levels of acetate

increase post-fire in subsurface soil (74). In this study, we demonstrate that

bacteria in the early post-fire stages, specifically at 17 and 25 days post-fire,

possess the genomic capacity to use this abundant post-fire resource, as indicated

by the higher abundance of genes such as *ACSS1_2, acs*, and *pta* associated with

the activation of acetate for use in central metabolism. This suggests that early

post-fire bacteria might prefer to degrade labile C sources that support rapid

growth, enabling these populations to dominate the ecosystem. However, later in

the succession process, the potential depletion of labile C sources leads to

community shifts, with microbes capable of using more complex aromatic C

sources rapidly dominating the community. Together, these findings highlight

the intricate interactions that may occur post-fire and suggest that metabolic

handoffs and cooperative resource use may play crucial roles in shaping post-fire

successional and ecosystem dynamics.

Post-fire microbes preferentially retain N and increase nitrification

Soil N cycling, primarily mediated by microbes, is crucial in the post-fire

environment and has strong implications for trace and GHG emissions (25).

Coinciding with significant increases in bioavailable N after the fire, our study

revealed that geTMM values for genes encoding nitrification, assimilatory, and

dissimilatory N reductase increased over time and exhibited differential

abundance in the burned plots. These results are supported by complementary

post-fire studies in coniferous forests (36, 75-78) and chaparral (79) ecosystems

indicating enrichment of nitrification in the immediate aftermath of wildfire. Our

previous research indicated an increase in *Massilia* and *Paenibacillus* over time

following the fire (8), and previous research shows *Paenibacillus* can fix

nitrogen (80, 81) supporting the observed increase in genes related to

nitrification, assimilatory, and dissimilatory N reductase in our study.

Additionally, N immobilization peaked after 131 days, coinciding with increased

abundance of *Arthrobacter* and *Micromonosporaceae*, both of which were found

to encode genes for N immobilization. Thus, our research suggests that during

succession, simultaneous functional changes are occurring and subsequently

driving bacterial successional dynamics, whereby a few bacterial taxa with more

specialized metabolic functions dominate and control bacterial successional

dynamics.

Wildfire-induced short-term N losses via increased microbial retention of

NO_3^- can be detrimental to post-fire ecosystem recovery (79), especially when

microbes, the major players in N retention (79, 82), are also affected by the fire.

Our study revealed an overrepresentation of denitrification genes (*nirK, narG,*

narZ, nxrA) in early post-fire pyrophilous bacteria (17-25 days post-fire), which

may exacerbate N losses in the form of N_2O, potent greenhouse gases. These

findings are consistent with a review of wildfire-affected ecosystems (83).

However, N losses are transient, as post-fire microbes, including those associated

with the genera *Arthrobacter, Massilia, Blastocatellia*, and the family

Micromonosporaceae, have the capacity to immobilize N through the

assimilatory NO_3- reduction pathway, incorporating it into biomass. Moreover,

the genera *Massilia, Pseudonocardia,* and *Micromonosporaceae* can further

convert NO_2^- to NH_3^+ via dissimilatory NO_2^- reductase (*nirBD*), making this

resource readily available for plant and microbial use. These results demonstrate

that post-fire microbes possess the genomic potential to retain N in the

ecosystem. Previous observations of N retention processes, such as microbial N

immobilization, have been documented in various wildfire-affected systems (84),

including chaparral (85), and in field studies involving PyOM additions (86). In

this study, we observed an increase in the abundance of genes related to N

incorporation by MAGs representing pyrophilous genera *Massilia* and

Arthrobacter consistent with prior research highlighting the N-reducing and

assimilating capabilities of both *Massilia* and *Arthrobacter* (87-90). Furthermore,

metagenomic studies have previously shown enrichment of *nirB/D* genes (N

retention genes) four years post-fire (36). These collective results underscore the

role that dominant pyrophilous bacteria in fire-adapted chaparral communities

may play in the assimilation and retention of ecosystem N, playing critical roles

in mitigating N losses and facilitating ecosystem restoration.

Post-fire chaparral microbes preferentially degrade PyOM via the ortho-

cleavage degradation pathway

550   Microbial taxa enriched in the post-fire soils encoded the ortho-cleavage

catechol and protocatechuate pathways for PyOM degradation more so than the

meta-cleavage pathways. However, the bacterial taxa implicated in PyOM

degradation only contained functional genes for part of a specific degradation

pathway, suggesting that post-fire bacterial communities may thrive via

cooperative interactions and metabolic handoffs. Prior studies have shown that

the β-ketoadipate pathway (e.g., ortho-cleavage pathway) is the most common

method for aromatic degradation and the dominant route for aromatic

degradation for lignin-degrading bacteria (91, 92). These results are also

supported by data from post-fire coniferous forest soils, which revealed the

presence of the ortho-cleavage pathway in over 50% of MAGs (11) and

intriguingly suggests that high-severity fires may select for ortho-cleavage

pathways regardless of the biome. Furthermore, we found that the genes for the

full catechol (*catABC*) and protocatechuate (*pcaHGBCDLJ*) ortho-cleavage

degradation pathways were present across all the burned plots for the entire year.

The differentially abundant ortho-cleavage pathway genes were linked to MAGs

representing multiple taxa in the Proteobacteria phylum, including

Noviherbaspirillum (Holy_399.fa and Holy_399.fa), *Massilia* (Holy_395.fa), and

Xanthobacteraceae PALSA-89 (Holy_368.fa and Holy_369.fa), the

Acidobacteriota phylum, Chthoniobacterales AV80 (Holy_436.fa) and the

Actinobacteriota phylum, *Pseudonocardia* (Holy_153.fa), and

Solirubrobacteraceae (Holy_206.fa). Previous research indicated that bacteria

preferentially use the catechol ortho-cleavage pathway while fungi preferentially

use the protocatechuate ortho-cleave pathway (46), suggesting that there is

interspecies competition for the same resource pool between pyrophilous

bacteria and fungi.

The catechol and protocatechuate orthocleavage pathways are

biochemically conserved with high diversity in the level of gene organization

(93, 94), which may indicate that pyrophilous microbes from wildfire-adapted

ecosystems such as chaparral have evolved mechanisms to capitalize on PyOM

as a C source. This is corroborated by the high relative abundance of

differentially abundant PyOM degradation genes in our burned plots. Our

findings indeed revealed differentially abundant PyOM genes, which were linked

to MAGs representing the Actinomycetales *Arthrobacter* (Holy_91.fa),

Pseudonocardia (Holy_153.fa), Micromonosporaceae (Holy_114.fa), as well as

Noviherbaspirillum (Holy_399.fa and Holy_399.fa), and *Massilia* (Holy_395.fa

and Holy_395.fa). Previous research shows that *Arthrobacter* (11) and

Burkholderiales encode genes for aromatic compound catabolism (95),

supporting our results. The ortho-cleavage pathway provides a growth advantage

to microbes (96) by efficiently converting C to cell mass (97-99) and both

pyrophilous *Massilia* and *Noviherbaspirillum* are fast-growing bacteria (100,

101) that dominated this system (8) and contained genes for PyOM degradation.

Together, these results suggest that in a post-fire system, where C is in the form

of commonly considered recalcitrant PyOM, pyrophilous bacteria prefer the

easier-to-degrade PyOM processing pathway (ortho-cleavage pathway), rapidly

converting PyOM to biomass and thus dominating and initiating microbial

succession, as previously suggested (8).

Conclusions

Our high-resolution gene-centric metagenomic analysis allowed us to

detect the rapid enrichment of genomic functional potential for the processing of

PyOM, inorganic N, alkanes, and urea in burned plots during the first post-fire

year, coinciding with the dominance of pyrophilous bacteria in the post-fire soils.

We found that MAGs representing the putative pyrophilous *Massilia* and

Noviherbaspirillium preferentially encode the ortho-cleavage catechol and

protocatechuate degradation pathways while assimilating N to acquire energy,

potentially minimizing GHG emissions. Interestingly, post-fire bacteria

employed distinct pathways for the degradation of post-fire resources such as

PyOM and N, suggesting that metabolic handoffs are common within post-fire

bacterial communities. This indicates that traits which have evolved for survival

after fires, especially those associated with acquiring resources, are the primary

drivers of bacterial secondary successional dynamics. While the capability of

post-fire bacteria to capitalize on PyOM and N has been previously hypothesized

(8, 71), and shown for some taxa (11, 17, 35), our research unveils previously

unknown functional changes over a high resolution successional timeline, filling

a critical gap created by limited post-fire sampling efforts. Thus, our work

significantly contributes to our understanding of post-wildfire bacterial ecology

and provides valuable insights into the intricate dynamics of bacterial

communities in fire-affected ecosystems.

3.7 Acknowledgments
621
We thank the Joint Genome Institute (JGI) Community Sequencing Project
(CSP) New Investigator Award 507364 to SIG, the Bureau of Land Management
(BLM) Joint Fire Science Program (JFSP) Graduate Research Innovation
(GRIN) Award 012641□002 to MFPC and SIG, the United States Department of
Agriculture (USDA)-NIFA Award #2022-67014-36675 to SIG, and the
Department of Energy (DOE) BER Award #DE-SC0023127 to SIG and MJW
for funding this project. We further thank the Cleveland National Forest and the
Trabuco Ranger District, including District Ranger Darrel Vance and Emily
Fudge, Jeffrey Heys, Lauren Quon, Jacob Rodriguez, and Victoria
Stempniewicz, for their help with permitting. We thank JGI Metagenome
Program, including Nicole Shapiro, Natasha Brown, Emiley Eloe-Fadrosh, and
Simon Roux, for their help in processing and submitting our metagenomic
samples. Lastly, we thank Judy A. Chung, Aral C. Greene, Elizah Stephens,
Dylan Enright, and Sameer S. Saroa for fieldwork assistance and James
Randolph for molecular assistance. We thank Jason Stajich and Pete Homyak
and our anonymous reviewers for helpful feedback on the manuscript.
638
639

640   **3.8 Competing Interests**
641
The authors declare no competing or conflicts of interest.

3.9 Data Availability Statement

The metagenomic reads and bacterial MAGs have been deposited to the
National Center for Biotechnology Information Sequence Read Archive
under BioProject PRJNA761539. Amplicon sequence reads for bacterial
16S rRNA and fungal ITS2 have been deposited to the National Center
for Biotechnology Information Sequence Read Archive under BioProject
PRJNA761539. All statistical R scripts/codes are available on GitHub

and include the sample metadata required for analysis
https://github.com/pulidofabs/Chaparral-Metagenomes.

3.10 References

1. Turner MG, Smithwick EAH, Metzger KL, Tinker DB, Romme WH.
    Inorganic nitrogen availability after severe stand-replacing fire in the
    Greater Yellowstone ecosystem. Proceedings of the National Academy of
    Sciences. 2007;104(12):4782–9.
2. Raza T, Qadir MF, Khan KS, Eash NS, Yousuf M, Chatterjee S, et al.
    Unrevealing the potential of microbes in decomposition of organic matter
    and release of carbon in the ecosystem. Journal of Environmental
    Management. 2023;344:118529.
3. Liu W, Zhang Z, Li J, Wen Y, Liu F, Zhang W, et al. Effects of fire on the
    soil microbial metabolic quotient: A global meta-analysis. CATENA.
    2023;224:106957.
4. Riley KL, Loehman RA. Mid-21st-century climate changes increase
    predicted fire occurrence and fire season length, Northern Rocky
    Mountains, United States. Ecosphere. 2016;7(11):e01543.
5. Enright DJ, Frangioso KM, Isobe K, Rizzo DM, Glassman SI. Mega-fire in
    Redwood Tanoak Forest Reduces Bacterial and Fungal Richness and
    Selects for Pyrophilous Taxa that are Phylogenetically Conserved.
    Molecular Ecology. 2022;31:2475–93.
6. Whitman T, Whitman E, Woolet J, Flannigan MD, Thompson DK, Parisien
    M-A. Soil bacterial and fungal response to wildfires in the Canadian boreal
    forest across a burn severity gradient. Soil Biology and Biochemistry.
    2019;138:107571.
7. Fox S, Sikes BA, Brown SP, Cripps CL, Glassman SI, Hughes K, et al. Fire
    as a driver of fungal diversity — A synthesis of current knowledge.
    Mycologia. 2022;114(2):1-27.
8. Pulido-Chavez MF, Randolph JWJ, Zalman C, Larios L, Homyak PM,
    Glassman SI. Rapid bacterial and fungal successional dynamics in first year
    after chaparral wildfire. Molecular Ecology. 2023;32(7):1685-707.
9. Caiafa MV, Nelson AR, Borch T, Roth HK, Fegel TS, Rhoades CC, et al.
    Distinct fungal and bacterial responses to fire severity and soil depth across
    a ten-year wildfire chronosequence in beetle-killed lodgepole pine forests.
    Forest Ecology and Management. 2023;544:121160.
10. Luo Y, Huang Y, Xu R-x, Qian B, Zhou J-w, Xia X-l. Primary and
    Secondary Succession Mediate the Accumulation of Biogenic Amines
    during Industrial Semidry Chinese Rice Wine Fermentation. Applied and
    Environmental Microbiology. 2020;86(17):e01177-20.

11. Nelson AR, Narrowe AB, Rhoades CC, Fegel TS, Daly RA, Roth HK, et
al. Wildfire-dependent changes in soil microbiome diversity and function.
Nat Microbiol. 2022;7(9):1419-30.
12. Nave L, Vance. E, . CS, Curtis. P. Fire effects on temperate forest soil C
and N storage. Ecol Appl. 2011;4:1189-201.
13. Knicker H. Pyrogenic organic matter in soil: Its origin and occurrence, its
chemistry and survival in soil environments. Quaternary International.
2011;243(2):251-63.
14. Bird MI, Wynn JG, Saiz G, Wurster CM, McBeath A. The Pyrogenic
Carbon Cycle. Annual Review of Earth and Planetary Sciences.
2015;43(1):273-98.
15. Haddock JD. Aerobic Degradation of Aromatic Hydrocarbons: Enzyme
Structures and Catalytic Mechanisms. Handbook of Hydrocarbon and
Lipid Microbiology2010. p. 1057-69.
16. Perez-Pantoja D, Donoso R, Agullo L, Cordova M, Seeger M, Pieper DH,
et al. Genomic analysis of the potential for aromatic compounds
biodegradation in Burkholderiales. Environ Microbiol. 2012;14(5):1091-
117.
17. Fischer MS, Stark FG, Berry TD, Zeba N, Whitman T, Traxler MF.
Pyrolyzed Substrates Induce Aromatic Compound Metabolism in the Post-
fire Fungus, Pyronema domesticum. Front Microbiol. 2021;12:729289.
18. Badía D, Martí C, Aguirre AJ, Aznar JM, González-Pérez JA, De la Rosa
JM, et al. Wildfire effects on nutrients and organic carbon of a Rendzic
Phaeozem in NE Spain: Changes at cm-scale topsoil. CATENA.
2014;113:267-75.
19. González-Pérez JA, González-Vila FJ, Almendros G, Knicker H. The
effect of fire on soil organic matter—a review. Environment International.
2004;30(6):855-70.
20. Kuhn TK, Krull ES, Bowater A, Grice K, Gleixner G. The occurrence of
short chain n-alkanes with an even over odd predominance in higher plants
and soils. Organic Geochemistry. 2010;41(2):88-95.
21. Abbasian F, Lockington R, Mallavarapu M, Naidu R. A Comprehensive
Review of Aliphatic Hydrocarbon Biodegradation by Bacteria. Appl
Biochem Biotechnol. 2015;176(3):670-99.
22. Moucawi J, Fustec E, Jambu P, Amblfs A, Jacquesy R. Biooxidation of
added and natural hydrocarbons in soils: Effect of iron. Soil Biology and
Biochemistry. 1981;13(5):335-42.
23. Thomas CL, Jansen B, van Loon EE, Wiesenberg GLB. Transformation of
n-alkanes from plant to soil: a review. SOIL. 2021;7(2):785-809.
24. Goberna M, Garcia C, Insam H, Hernandez MT, Verdu M. Burning fire-
prone Mediterranean shrublands: immediate changes in soil microbial
community structure and ecosystem functions. Microb Ecol.
2012;64(1):242-55.
25. Dannenmann M, Díaz-Pinés E, Kitzler B, Karhu K, Tejedor J, Ambus P, et
al. Postfire nitrogen balance of Mediterranean shrublands: Direct

combustion losses versus gaseous and leaching losses from the postfire soil
mineral nitrogen flush. Global Change Biology. 2018;24(10):4505-20.
26. Johnson DW, Curtis PS. Effects of forest management on soil C and N
storage: meta analysis. Forest Ecology and Management. 2001;140(2-
3):227-38.
27. Neary DG, Klopatek CC, DeBano LF, Ffolliott PF. Fire effects on
belowground sustainability: a review and synthesis. Forest Ecology and
Management. 1999;122(1-2):51-71.
28. DeBano LF, Neary DG, Ffolliott PF. Fire Effects on Ecosystems. 1998.
29. Ayiti OE, Babalola OO. Factors Influencing Soil Nitrification Process and
the Effect on Environment and Health. Frontiers in Sustainable Food
Systems. 2022;6.
30. Chen H, Ma K, Huang Y, Yao Z, Chu C. Stable Soil Microbial Functional
Structure Responding to Biodiversity Loss Based on Metagenomic
Evidences. Frontiers in Microbiology. 2021;12.
31. Hoang HG, Thuy BTP, Lin C, Vo D-VN, Tran HT, Bahari MB, et al. The
nitrogen cycle and mitigation strategies for nitrogen loss during organic
waste composting: A review. Chemosphere. 2022;300:134514.
32. Singh J, Kunhikrishnan A, Bolan NS, Saggar S. Impact of urease inhibitor
on ammonia and nitrous oxide emissions from temperate pasture soil cores
receiving urea fertilizer and cattle urine. The Science of the Total
Environment. 2013;465:56-63.
33. Yoon S, Song B, Phillips RL, Chang J, Song MJ. Ecological and
physiological implications of nitrogen oxide reduction pathways on
greenhouse gas emissions in agroecosystems. FEMS Microbiology
Ecology. 2019;95(6):fiz066.
34. Thomson AJ, Giannopoulos G, Pretty J, Baggs EM, Richardson DJ.
Biological sources and sinks of nitrous oxide and strategies to mitigate
emissions. Philosophical Transactions of the Royal Society B: Biological
Sciences. 2012;367(1593):1157-68.
35. Cobo-Díaz JF, Fernández-González AJ, Villadas PJ, Robles AB, Toro N,
Fernández-López M. Metagenomic Assessment of the Potential Microbial
Nitrogen Pathways in the Rhizosphere of a Mediterranean Forest After a
Wildfire. Microbial Ecology. 2015;69(4):895-904.
36. Dove NC, Tas N, Hart SC. Ecological and genomic responses of soil
microbiomes to high-severity wildfire: linking community assembly to
functional potential. ISME J. 2022;16(7):1853-63.
37. Guo M, Shang X, Ma Y, Zhang K, Zhang L, Zhou Y, et al. Biochars
assisted phytoremediation of polycyclic aromatic hydrocarbons
contaminated agricultural soil: Dynamic responses of functional genes and
microbial community. Environmental Pollution. 2024;345.
38. Joshi NA FJ. Sickle: A sliding-window, adaptive, quality-based trimming
tool for FastQ files 2011;Version 1.33.
786
39. Andrews S. FastQC: A Quality Control Tool for High Throughput
Sequence Data [Online]. 2010.

40.   Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: An ultra-fast
single-node solution for large and complex metagenomics assembly via
succinct de Bruijn graph. arXiv:14097208 [q-bio]. 2014.
41.   Woodcroft BJ. CoverM. Queensland University of Technology Centre for
Microbiome Research: School of Biomedical Sciences; 2007.
42.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW.
CheckM: assessing the quality of microbial genomes recovered from
isolates, single cells, and metagenomes. Genome Res. 2015;25(7):1043-55.
43.   Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for
accurately reconstructing single genomes from complex microbial
communities. PeerJ. 2015;3:e1165.
44.   Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit
to classify genomes with the Genome Taxonomy Database. Bioinformatics.
2019:btz848.
45.   Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and
accurate genomic comparisons that enables improved genome recovery
from metagenomes through de-replication. The ISME Journal.
2017;11(12):2864-8.
46.   Fuchs G, Boll M, Heider J. Microbial degradation of aromatic compounds -
from one strategy to four. Nat Rev Microbiol. 2011;9(11):803-16.
47.   Eddy SR. Accelerated Profile HMM Searches. PLOS Computational
Biology. 2011;7(10):e1002195.
48.   Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et
al. KofamKOALA: KEGG Ortholog assignment based on profile HMM
and adaptive score threshold. Bioinformatics. 2020;36(7):2251-2.
49.   Khot V, Zorz J, Gittins DA, Chakraborty A, Bell E, Bautista MA, et al.
CANT-HYD: A Curated Database of Phylogeny-Derived Hidden Markov
Models for Annotation of Marker Genes Involved in Hydrocarbon
Degradation. Frontiers in Microbiology. 2022;12.
50.   Shaffer M, Borton MA, McGivern BB, Zayed AA, Rosa SLL, Solden LM,
et al. DRAM for distilling microbial metabolism to automate the curation
of microbiome function. bioRxiv. 2020:2020.06.29.177501.
51.   Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes.
Nucleic Acids Research. 2000;28(1):27-30.
52.   Suzek BE, Huang H, McGarvey P, Mazumder R, Wu CH. UniRef:
comprehensive and non-redundant UniProt reference clusters.
Bioinformatics. 2007;23(10):1282-8.
53.   Rawlings ND, Waller M, Barrett AJ, Bateman A. *MEROPS* : the database
of proteolytic enzymes, their substrates and inhibitors. Nucleic Acids
Research. 2014;42(D1):D503-D9.
54.   Steinegger M, Söding J. MMseqs2 enables sensitive protein sequence
searching for the analysis of massive data sets. Nat Biotechnol.
2017;35(11):1026-8.
55.   Smid M, Coebergh van den Braak RRJ, van de Werken HJG, van Riet J,
van Galen A, de Weerd V, et al. Gene length corrected trimmed mean of
M-values (GeTMM) processing of RNA-seq data performs similarly in

intersample analyses while improving intrasample comparisons. BMC
Bioinformatics. 2018;19(1):236.

56. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data.
Bioinformatics (Oxford, England). 2010;26(1):139-40.

57. Capo E, Broman E, Bonaglia S, Bravo AG, Bertilsson S, Soerensen AL, et
al. Oxygen☐deficient water zones in the Baltic Sea promote
uncharacterized Hg methylating microorganisms in underlying sediments.
Limnology and Oceanography. 2021;67(1):135-46.

58. Leleiwi I, Rodriguez-Ramos J, Shaffer M, Sabag-Daigle A, Kokkinias K,
Flynn RM, et al. Exposing new taxonomic variation with inflammation - a
murine model-specific genome database for gut microbiome researchers.
Microbiome. 2023;11(1):114.

59. Santos-Medellin C, Blazewicz SJ, Pett-Ridge J, Firestone MK, Emerson
JB. Viral but not bacterial community successional patterns reflect extreme
turnover shortly after rewetting dry soils. Nat Ecol Evol. 2023;7(11):1809-
22.

60. Bethesda. Building a BLAST database with your (local) sequences:BLAST
Command Line Applications User Manual. National Center for
Biotechnology Information. 2008.

61. RCoreTeam. R: A language and environment for statistical computing. R
Foundation for Statistical Computing. Vienna, Austria.2023.

62. Anderson MJ. Permutational Multivariate Analysis of Variance
(PERMANOVA). Wiley StatsRef: Statistics Reference Online: John Wiley
& Sons, Ltd; 2017. p. 1-15.

63. Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, et al.
vegan: Community Ecology Package. 2022.

64. McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible
Interactive Analysis and Graphics of Microbiome Census Data. . PLOS
ONE. 2013;8(4):e61217.

65. Fernández-García V, Marcos E, Reyes O, Calvo L. Do Fire Regime
Attributes Affect Soil Biochemical Properties in the Same Way under
Different Environmental Conditions? Forests. 2020;11(3):274.

66. Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol.
2014;15(12):550.

67. Chen Z, Tu X, Meng H, Chen C, Chen Y, Elrys AS, et al. Microbial
process-oriented understanding of stimulation of soil N2O emission
following the input of organic materials. Environmental Pollution.
2021;284:117176.

68. Zeba N, Berry TD, Fischer MS, Traxler MF, Whitman T. Soil carbon
mineralization and microbial community dynamics in response to PyOM
addition. preprint. Ecology; 2023 2023/06/22/.

69. Cheng H, Tang G, Wang S, Rinklebe J, Zhu T, Cheng L, et al. Combined
remediation effects of biochar and organic fertilizer on immobilization and
dissipation of neonicotinoids in soils. Environ Int. 2022;169:107500.

70.   Song H-K, Song W, Kim M, Tripathi BM, Kim H, Jablonski P, et al.
Bacterial strategies along nutrient and time gradients, revealed by
metagenomic analysis of laboratory microcosms. FEMS Microbiology
Ecology. 2017;93(fix114).
71.   Woolet J, Whitman T. Pyrogenic organic matter effects on soil bacterial
community composition. Soil Biology and Biochemistry.
2020;141:107678.
72.   Johnson CW, Beckham GT. Aromatic catabolic pathway selection for
optimal production of pyruvate and lactate from lignin. Metabolic
Engineering. 2015;28:240-7.
73.   Sgro M, Chow N, Olyaei F, Arentshorst M, Geoffrion N, Ram AFJ, et al.
Functional analysis of the protocatechuate branch of the β-ketoadipate
pathway in Aspergillus niger. Journal of Biological Chemistry.
2023;299(8):105003.
74.   Blank RR, Allen F, Young JA. Extractable Anions in Soils following
Wildfire in a Sagebrush-Grass Community. Soil Science Society of
America Journal. 1994;58(2):564-70.
75.   Dove NC, Klingeman DM, Carrell AA, Cregger MA, Schadt CW. Fire
alters plant microbiome assembly patterns: integrating the plant and soil
microbial response to disturbance. New Phytologist. 2021;230(6):2433-46.
76.   Kurth VJ, Fransioli N, Perez FZ, Hart SC, Gehring CA. Stand-replacing
wildfires alter the community structure of wood-inhabiting fungi in
southwestern ponderosa pine forests of the USA | Elsevier Enhanced
Reader. 2013.
77.   Dove NC, Safford HD, Bohlman GN, Estes BL, Hart SC. High-severity
wildfire leads to multi-decadal impacts on soil biogeochemistry in mixed-
conifer forests. Ecol Appl. 2020;30(4):e02072.
78.   Kurth VJ, Hart SC, Ross CS, Kaye JP, Fule PZ. Stand-replacing wildfires
increase nitrification for decades in southwestern ponderosa pine forests.
Oecologia. 2014;175(1):395-407.
79.   Hanan EJ, Tague C, Schimel JP. Nitrogen cycling and export in California
chaparral: the role of climate in shaping ecosystem responses to fire.
Ecological Monographs. 2016;87(1):76-90.
80.   Monciardini P, Cavaletti L, Schumann P, Rohde M, Donadio S.
Conexibacter woesei gen. nov., sp. nov., a novel representative of a deep
evolutionary line of descent within the class Actinobacteria. Int J Syst Evol
Microbiol. 2003;53(Pt 2):569-76.
81.   Slepecky RA, Hemphill HE. The Genus Bacillus—Nonmedical. In:
Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E,
editors. The Prokaryotes: Volume 4: Bacteria: Firmicutes, Cyanobacteria.
New York, NY: Springer US; 2006. p. 530-62.
82.   Knicker H. How does fire affect the nature and stability of soil organic
nitrogen and carbon? A review. Biogeochemistry. 2007;85(1):91-118.
83.   Stephens EZ, Homyak PM. Post-fire soil emissions of nitric oxide (NO)
and nitrous oxide (N2O) across global ecosystems: a review.
Biogeochemistry. 2023.

84. Xu W, Elberling B, Ambus PL. Fire increases soil nitrogen retention and
alters nitrogen uptake patterns among dominant shrub species in an Arctic
dry heath tundra. Sci Total Environ. 2022;807(Pt 3):150990.
85. Goodridge BM, Hanan EJ, Aguilera R, Wetherley EB, Chen Y-J,
D'Antonio CM, et al. Retention of Nitrogen Following Wildfire in a
Chaparral Ecosystem. Ecosystems. 2018;21(8):1608-22.
86. Bai SH, Reverchon F, Xu C-Y, Xu Z, Blumfield TJ, Zhao H, et al. Wood
biochar increases nitrogen retention in field settings mainly through abiotic
processes. Soil Biology and Biochemistry. 2015;90:232-40.
87. Bailey AC, Kellom M, Poret-Peterson AT, Noonan K, Hartnett HE,
Raymond J. Draft Genome Sequence of Massilia sp. Strain BSC265,
Isolated from Biological Soil Crust of Moab, Utah. Genome
Announcements. 2014;2(6):e01199-14, 2/6/e-14.
88. <Cacciari et al. - 1986 - Ammonium assimilation in an Arthrobacter sp.
"fluo.pdf>.
89. Cacciari I, Lippi D, Pietrosanti T, Pietrosanti W. Ammonium assimilation
in an Arthrobacter sp. "fluorescens". Archives of Microbiology.
1986;145(2):113-5.
90. Liu Y, Zhang Y, Huang Y, Niu J, Huang J, Peng X, et al. Spatial and
temporal conversion of nitrogen using Arthrobacter sp. 24S4-2, a strain
obtained from Antarctica. Front Microbiol. 2023;14:1040201.
91. Granja-Travez RS, Persinoti GF, Squina FM, Bugg TDH. Functional
genomic analysis of bacterial lignin degraders: diversity in mechanisms of
lignin oxidation and metabolism. Appl Microbiol Biotechnol.
2020;104(8):3305-20.
92. Weiland F, Kohlstedt M, Wittmann C. Guiding stars to the field of dreams:
Metabolically engineered pathways and microbial platforms for a
sustainable lignin-based industry. Metabolic Engineering. 2022;71:13-41.
93. Parke D, D'Argenio DA, Ornston LN. Bacteria Are Not What They Eat:
That Is Why They Are So Diverse. Journal of Bacteriology.
2000;182(2):257-63.
94. Buchan A, Neidle EL, Moran MA. Diverse organization of genes of the
beta-ketoadipate pathway in members of the marine Roseobacter lineage.
Appl Environ Microbiol. 2004;70(3):1658-68.
95. Pérez-Pantoja D, Donoso R, Agulló L, Córdova M, Seeger M, Pieper DH,
et al. Genomic analysis of the potential for aromatic compounds
biodegradation in Burkholderiales. Environmental Microbiology.
2012;14(5):1091-117.
96. Song J, Sung J, Kim YM, Gerben J Z, Kim E. Roles of the meta- and the
ortho-Cleavage Pathways for the Efficient Utilization of Aromatic
Hydrocarbons by Sphingomonas yanoikuyae Bl. Korean Journal of
Microbiology. 2002;38(4):245-.
97. Aghapour AA, Moussavi G, Yaghmaeian K. Biological degradation of
catechol in wastewater using the sequencing continuous-inflow reactor
(SCR). J Environ Health Sci Eng. 2013;11:3.
98. <Basha et al. - Recent advances in the Biodegradation of Phenol A.pdf>.

99.   Basha KM, Rajendran A, Thangavelu V. Recent advances in the
974   Biodegradation of Phenol: A review. 2010;1:219-34.
975   100. Baldani JI, Rouws L, Cruz LM, Olivares FL, Schmid M, Hartmann A. The
976   Family Oxalobacteraceae. In: Rosenberg E, DeLong EF, Lory S,
977   Stackebrandt E, Thompson F, editors. The Prokaryotes. Berlin, Heidelberg:
978   Springer Berlin Heidelberg; 2014. p. 919-74.
979   101. Li X, Gou X, Long D, Ji Z, Hu L, Xu D, et al. Physiological and metabolic
980   analysis of nitrate reduction on poly-gamma-glutamic acid synthesis in
981   Bacillus licheniformis WX-02. Archives of Microbiology.
982   2014;196(11):791-9.
983

Figure 1. Non-metric multidimensional scaling (NMDS) of geTMM normalized a) PyOM b) alkane c) Nitrogen and d) urea cycling genes within the microbial communities of burned and unburned soil samples (color) and time since fire (days; shape). Treatment and time since fire effects based on PERMANOVA (p <0.05). Ellipse represents the 95% confidence level.

Figure 2. Principal component analysis (PCoA) of the mean and standard error Bray-Curtis community dissimilarity at each timepoint for the unburned and burned a,b) PyOM; c,d) Alkane; e, f) nitrogen and g, h) urea functional potentials. Note that timepoints further apart with nonoverlapping standard error bars indicate a community turnover, whereas overlapping standard error bars represent a lack of compositional turnover, with the first sampling time point at 17 days representing the base level. Thus, the first turnover was initiated at 25 days. There were no significant turnover dates for any gene functional potentials in the unburned plots. The days with significant turnovers in the burned plots are listed in the upper right corners. Significance based on betadisper, which measures the dispersion/variability of the data within groups as a means to measure community turnover.

Figure 3. Summed geTMM normalized gene abundance over time for a,b) PyOM and c, d) Alkane degradation in the unburned (left panel) and burned (right panel). Colors are based on the general function for each gene.

Figure 4. Summed geTMM normalized gene abundance over time for a,b) Nitrogen and c, d) Urea cycling genes in the unburned (left panel) and burned (right panel). Colors are based on the general function for each gene.

Figure 5. Total number of genes that were differentially abundant in the burned plots (brown) versus the unburned plots (blue green; x axis) per each metabolic pathway, colored by each sampling time point for a) PyOM, b) Alkane, Nitrogen and d) Urea cycling. Significance based on DESeq2 Wald test (adjusted P value < 0.05 and absolute log2 fold-change >0).

Figure 6. The catechol and protocatechuate degradation pathways with arrows indicating genes that were differentially expressed in the burned treatment at each time point (17, 25, 34, 131 and 376 days post-fire; Wald's test in DESeq2; $p<0.05$). Line type represents catechol (solid) and protocatechuate (dashed) cycling pathways and gray solid lines represent pathways that were not expressed during post-fire year one. Numbers in parentheses represent the mean log2fold change (DESeq2) for each represented gene and are colored by the timepoint in which that gene was significantly abundantly represented.

Figure 7. The nitrogen cycle with arrows indicating genes that were differentially represented in the burned treatment at each time point (17, 25, 34, 131, and 376 days post-fire; Wald's test in DESeq2; $p<0.05$). Line type represents the different nitrogen cycling pathways, colored by timepoint and gray solid lines represent pathways that were not expressed during post-fire year one. Numbers in parentheses represent the mean log2fold change (DESeq2) for each represented gene and are colored by the timepoint in which that gene was significantly abundantly represented.

Figure 8. Taxonomic annotation for the most dominant, differentially abundant PyOM, nitrogen and urea genes based on DESeq analysis (adjusted P value < 0.05 and absolute log2 fold-change >0). Taxonomy based on blasting contigs against the Metagenome Assembled Genomes (MAGs) represented in the x-axis.