1 Chaparral wildfire shifts the functional potential for soil pyrogenic organic matter and
2 nitrogen cycling
3
4

5 M. Fabiola Pulido Barriga[1+], Amelia R. Nelson[2], Peter M. Homyak[3], Michael J. Wilkins[2],
6 Sydney I. Glassman[1,+]
7
8
9
[1]Department of Microbiology and Plant Pathology, University of California- Riverside, 900
University Ave, Riverside, CA, 92521
[2]Department of Soil and Crop Sciences, Colorado State University, 307 University Ave,
Fort Collins, CO, 80521
[3]Department of Environmental Sciences, University of California-Riverside, 900 University
Ave, Riverside, CA, 92521
16
17
18
19 + Corresponding authors: Sydney I Glassman, sydneyg@ucr.edu, and M. Fabiola Pulido
20 Barriga, fabspulido@gmail.com

21
22
23 Keywords: Pyrophilous soil bacteria, PyOM, metagenomics, gene-centric, nitrogen
24 retention, ortho-cleavage pathway, β-ketoadipate
25
26
27 Running head: Wildfire Alters Soil PyOM and N Cycling genes
28
29
30
31

Abstract

Wildfires reshape soil microbiomes and chemistry, enhancing nitrogen availability and leaving behind pyrogenic organic matter (PyOM), a difficult to degrade carbon substrate potentially used by pyrophilous or "fire-loving" microbes. Understanding whether pyrophilous bacteria can metabolize post-fire resources is critical for predicting the fate of both carbon and nitrogen. We explored how secondary succession of pyrophilous bacteria align with changes in functional gene composition, particularly genes related to PyOM degradation and microbial nitrogen metabolism using shotgun metagenomics on 30 burned and unburned soils collected at 17, 25, 34, 131, and 376 days after a high-severity wildfire in a fire-adapted chaparral in Southern California. In burned soils, genes for PyOM degradation increased over time by 167% and for inorganic nitrogen cycling by 117%, while unburned soils showed no significant changes. Genes encoding catechol and protocatechuate, intermediates in the PyOM degradation pathway, indicate that the easier-to-degrade ortho-cleavage pathways consistently dominated the burned plots. These pathways co-occurred with genes for nitrification and nitrogen retention, including assimilatory and dissimilatory nitrate reduction to ammonia (DNRA). We also identified 446 bacterial metagenome-assembled genomes (MAGs) and linked gene profiles to dominant taxa and found that increases in genes for PyOM and nitrogen cycling paralleled the dominance of pyrophilous *Massilia* and *Noviherbaspirillum*, which encoded distinct pathways for PyOM and inorganic nitrogen use over time. Together, these findings reveal previously unrecognized functional shifts in bacterial communities over a high-resolution successional timeline, providing insights into the long-term impact of fire on microbial-mediated ecosystem processes that shape soil carbon and nitrogen dynamics.

1.1 Introduction

Soil microbial communities support fundamental post-fire soil processes, including nitrogen (N) cycling through processes like immobilization, denitrification, and leaching (Turner et al., 2007), and carbon (C) cycling through sequestration and CO_2 release during decomposition (Liu et al., 2023a; Raza et al., 2023). Despite evidence of rapid post-fire microbial succession (Pulido-Chavez et al., 2023), the functional dynamics of post-fire soil microbiomes remain poorly understood, largely due to the limited availability of high-resolution temporal sampling. As fire regimes increase in frequency and severity (Cunningham et al., 2024), understanding early functional changes in post-fire soil microbiomes is crucial for determining how microbes drive secondary succession and influence ecosystem functional recovery.

Wildfires have profound and lasting effects on global C and N cycling. Fires produce pyrogenic organic matter (PyOM)—a persistent C substrate formed during the incomplete combustion of vegetation (Knicker, 2011; Nave et al., 2011; Bird et al., 2015). PyOM's complex chemical structure, with highly condensed aromatic and O-aromatic components (Chatterjee et al., 2012; De la Rosa et al., 2012; Chen et al., 2022), makes it difficult for microbes to degrade, typically breaking down over centuries (Singh et al., 2012), leading to long-term effects on post-fire nutrient dynamics (Haddock, 2010; Knicker, 2011; Bird et al., 2015). However, in-lab research suggests that PyOM breaks down into smaller structures within 10 months (Singh et al., 2014), which, while still containing aromatic components, may be more accessible to microbial enzymes, potentially facilitating microbial PyOM degradation. Additionally, wildfires alter soil chemistry by increasing pH, ammonium (NH_4^+), nitrate (NO_3^-), and urease activity (Ayiti and Babalola, 2022) which are

intensified under high-severity fires that alter soil temperature and moisture (Neary et al.,

1999; Moya et al., 2018), further impacting N cycling processes like nitrification and

denitrification. However, wildfires typically reduce bacterial richness and shift community

composition, favoring pyrophilous ("fire-loving") bacteria, particularly from the Firmicutes

and Actinobacteria phyla (Whitman et al., 2019; Enright et al., 2022; Pulido-Chavez et al.,

2023). Yet, it remains unclear whether these bacteria possess the genes necessary to degrade

PyOM and utilize the available post-fire N.

Pyrophilous bacteria can shape successional dynamics by altering resource use and

driving predictable shifts in species composition (Pulido-Chavez et al., 2023) and

metabolism (Luo et al., 2020; Nelson et al., 2022). They may drive secondary succession

through a trait-based transition from thermotolerance to fast growth and ultimately PyOM

utilization (Pulido-Chavez et al., 2023). In this context, early fast-growers metabolize labile

C substrates, priming or preparing the soil for later-arriving microbes capable of degrading

complex aromatic compounds in PyOM. This process likely relies on functional succession

(Ni et al., 2023), where taxa integrate sequentially based on metabolic traits (Jackrel et al.,

2019), potentially stabilizing microbial communities and enhancing ecosystem recovery.

However, whether these functional changes occur simultaneously or influence microbial

successional dynamics remains unclear.

Bacterial degradation of complex aromatic C substrates, such as lignin or PyOM

occurs through the catechol and protocatechuate pathways, following either the meta- or

ortho-cleavage pathway (Fuchs et al., 2011). The ortho-cleavage (β-ketoadipate) pathway is

favored (Harwood and Parales, 1996; Parke et al., 2000; Buchan et al., 2004), due to its

efficiency, higher energy yield, and enzyme specificity (Harwood and Parales, 1996; Bugg

et al., 2011; Fuchs et al., 2011), which may be particularly relevant in post-fire landscapes

(20). For example, *Massilia*, a well-known pyrophilous genus in the Burkholderiales

(Enright et al., 2022; Pulido-Chavez et al., 2023), may degrade both lignin (Wang L, 2016)

and PyOM (Pérez-Pantoja et al., 2012) via the ortho-cleavage pathway (Pérez-Pantoja et al.,

2012). Beyond aromatic compounds, wildfires also increase short-chain alkanes (González-

Pérez et al., 2004; Badía et al., 2014), such as propane, a relatively more labile C source

preferentially consumed by microbes (Moucawi et al., 1981; Abbasian et al., 2015; Thomas

et al., 2021). However, it remains undocumented how and when pyrophilous bacteria

metabolize these different C sources or how their C substrate preference shifts during post-

fire microbial succession.

Microbial metabolism also plays a crucial role in post-fire N cycling, influencing

nutrient availability and ecosystem recovery. While post-fire greenhouse gas (GHG)

emissions are common (Stephens and Homyak, 2023), microbial communities can mitigate

these emissions by converting abundant NH☐☐ and NO☐☐ into microbial biomass via

assimilatory and dissimilatory nitrate reduction (DNRA) (Neary et al., 1999). This process

may shape microbial succession by favoring taxa with NO☐☐ assimilation genes. Indeed,

rapid post-fire shifts in microbial composition (Pulido-Chavez et al., 2023) likely reflect

changes in microbial metabolism, influencing the recovery of inorganic N, such as NH☐☐

and NO☐☐ which undergo rapid changes in the first post-fire years (Stephens and Homyak,

2023). Notably, pyrophilous bacteria analyzed 3–10 years post-fire retain substantial N-

cycling and PyOM degradation genes (Dove et al., 2022; Nelson et al., 2022; Whitman et

al., 2022), indicating their adaptation to shifting soil nutrient conditions. This suggests that

microbial communities adapt to shifting post-fire conditions and may influence ecosystem

recovery through functional changes that drive microbial succession.

Here, we used shotgun metagenomics to examine how pyrophilous bacteria drive

post-fire microbial functional shifts in C and N cycling, leveraging soils where we

previously characterized microbiomes across nine time points from 2 weeks to 1 year post-

fire (Pulido-Chavez et al., 2023). Focusing on five key time points corresponding to major

bacterial turnover events (Pulido-Chavez et al., 2023), we examined whether bacterial

secondary succession aligns with shifts in functional gene composition, particularly genes

related to PyOM degradation and microbial N metabolism. We used metagenomically

assembled genomes (MAGs) to link functional changes to dominant pyrophilous bacteria

over time. We tested the hypotheses that H1) The relative abundance of PyOM degradation

genes will increase over time as alkane degradation genes decrease, reflecting microbial

adaptation to PyOM as a C source; H2) Genes involved in microbial N metabolism will

increase over time, indicating shifts in bacterial functional potential in response to post-fire

N availability; and H3) Shifts in gene abundances will correspond with bacterial community

shifts, particularly the increasing dominance of pyrophilous bacteria over time.

1.2 Methods

1.2.1 Sampling

In August-September 2018, the Holy Fire burned 94 km² of Manzanita-dominated

chaparral in Southern California. Using BAER soil burn severity maps (USGS), we

established nine plots (six burned, three unburned) 17 days post-fire, selecting sites with

similar pre-fire vegetation, elevation, and aspect. These plots spanned Entisols (Typic

Xerorthents) mapped in the Cieneba series and Mollisols (Lithic Haploxerolls) in the Friant

series (Pulido-Chavez et al., 2023). Each plot contained four 1 m² subplots in each cardinal

direction for temporal soil sampling of the top 10 cm of mineral soil beneath the ash or duff

layer (Fig. S1). To assess soil burn severity at the plot level, we used ash depth as a proxy,

averaging three measurements per subplot. Additional details of the experimental design can

be found in Pulido et al. (Pulido-Chavez et al., 2022). Soil samples were collected at nine

time points ranging from 2 weeks to 1 year post-fire and stored at -80°C within 24h of

collection (Pulido-Chavez et al., 2023). DNA was extracted from 0.25g of 2mm sieved soil

and we performed 16S rRNA and ITS2 Illumina MiSeq sequencing of all samples (Pulido-

Chavez et al., 2023). Building on our previous study of successional dynamics, we selected

the five time points with the largest bacterial turnover events (17, 25, 34, 131, and 376 days

post-fire) and sufficient high-quality DNA for metagenomic sequencing (Pulido-Chavez et

al., 2023).

1.2.2 Metagenomic sequencing

From each of the five timepoints, we selected 3 burned and 3 unburned subplots,

totaling 30 metagenomes. Samples with DNA concentrations of 2.5 ng/µl and 230/260

above 1.8 were selected for metagenomes. Samples were collected from unburned plot 8

and two closely situated burned plots with similar microbial richness and biomass (plots 2

and 3; Fig. S1a). Metagenomic library preparation and sequencing were performed at the

Department of Energy Joint Genome Institute (JGI) for 17 samples and at the University of

California, Irvine (UCI) Genomics High-Throughput Facility for 13 samples. Libraries were

prepared using the Kapa Biosystems library preparation kit at JGI and the Illumina Nextera

Flex library preparation kit at UCI. Sequencing was performed on the Illumina NovaSeq

6000 platform with 151 bp paired-end reads and a depth of 25 Gb/sample.

1.2.3 Metagenomics binning

Sequence adapters were removed from the raw reads using BBduk-v38.89

(sourceforge.net/projects/bbmap/) with the following parameters (ktrim=r, k=23, mink=11,

hdist=1). Raw reads were trimmed with Sickle – v1.33 (Joshi NA, 2011), and the quality of

trimmed reads was assessed with FastQC – v0.11.2 (Andrews, 2010). Trimmed reads were

assembled into contiguous sequences (contigs) with the de novo de Bruijn assembler

MEGAHIT – v1.2.9 (Li et al., 2014) with a min kmer of 27, max of 127, and increment of

10. Per-contig coverage per sample was calculated using CoverM contig v0.6.0

(Woodcroft, 2007) with the trimmed mean method—i.e., the average number of aligned

reads per position after removing the most deeply and shallowly covered regions—while

retaining only mappings with >95% identity and a minimum alignment length of 75%

(Parks et al., 2015). Assembled contigs (>2,500 bp) were binned using MetaBAT2 v2.12.1

with default parameters (Kang et al., 2015) to assemble metagenome-assembled genomes

(MAGs). MAG quality was estimated using checkM v1.1.2 (Parks et al., 2015) and

taxonomy was assigned using GTDB-Tk v2.1.1 (Chaumeil et al., 2019). To create the final

medium and high-quality MAG database, low-quality MAGs (<50% completion, >10%

contamination (Bowers et al., 2017) were removed and the dataset was dereplicated using

dRep v3.0.0 (Olm et al., 2017).

1.2.3 Contig annotation

We assessed microbial preferences for inorganic versus organic N cycling, focusing

on urea, the largest contributor to the organic N cycle through hydrolysis into NH☐☐

(Fernández-García et al., 2020). For C cycling, we focused on short-chain alkanes as a more

readily degradable C source and PyOM genes, focusing on catechol and protocatechuate,

key intermediates of aerobic aromatic degradation (Fuchs et al., 2011), as the more complex

C sources.

To improve retention and to comprehensively annotate metagenomes, we annotated

all assembled contigs using three different databases. First, contigs were annotated using

HMMER (Eddy, 2011), with hidden Markov models (HMMs) against Kofamscan HMMs

(Aramaki et al., 2020) to identify specific genes according to previously established

methods (Nelson et al., 2022). Second, we used the curated Calgary approach to

AnnoTating HYDrocarbon degradation genes (CANT-HYD) database, which uses HMMs

for annotating marker genes involved in aerobic and anaerobic hydrocarbon degradation

pathways (Khot et al., 2022). Third, we used DRAM (Shaffer et al., 2020), an annotation

tool to profile our contigs through multiple databases including KEGG (Kanehisa and Goto,

2000), UniRef90 (Suzek et al., 2007), MEROPS (Rawlings et al., 2014), and Mmseqs2

(Steinegger and Söding, 2017) with the best hits, based on a bit score of 60, reported for

each database. Together, these methods improve annotation accuracy by cross-validating

results, capturing both broad metabolic functions and specialized enzymatic activities, and

improving confidence in gene function assignments. Contigs counts across samples were

normalized using the contig length corrected trimmed mean of M values (geTMM) (Smid et

al., 2018) using EdgeR (Robinson et al., 2010), adjusting for library depth and gene length

(Capo et al., 2021; Leleiwi et al., 2023; Santos-Medellin et al., 2023). For contigs with

duplicate annotations, we manually curated to select the annotation with the lowest e-value

as the best functional annotation.

We identified genes of interest using DESeq analysis (details below) and linked

them to our MAG catalog via NCBI BLAST (Bethesda, 2008), filtering to include only hits

with 100% identity and a bit score ≥ 100, which we used to identify the specific taxa that

encoded key genes related to PyOM, alkane, and organic and inorganic N cycling pathways.

See Supplementary Data 1 for additional information regarding metagenomic sequencing,

assemblies, and annotations, and see Fig. S2 for an overview of bioinformatic procedures

employed.

1.2.4 Statistical analysis

All statistical analyses and figures were done in R version 4.2.3 (RCoreTeam, 2023).

To determine how fire, time, and their interaction impacted the C and N gene composition,

we employed Permutational Multivariate Analysis of Variance (PERMANOVA;(Anderson,

2017)) via adonis2 in the vegan package (Oksanen et al., 2022) with 9999 permutations with

plot as random effect and visualized with nonmetric multidimensional scaling (NMDS).

Ordinations were based on Bray–Curtis distance matrixes using the ordinate function in the

phyloseq package (McMurdie and Holmes, 2013).

To determine if bacterial functional changes mirrored previously observed rapid

community compositional changes (Pulido-Chavez et al., 2023), we plotted the most

abundant bacterial genera identified with 16S amplicons at each post-fire timepoint. We also

calculated the homogeneity of variance using permutest with 9999 permutations and plot as

random effect to determine which time points differed for burned and unburned pathways

independently. Principal coordinates analysis (PCoA) was used to visualize the differences

in the homogeneity of variance of each functional pathway employing Bray-Curtis

dissimilarities in the vegan package.

We performed generalized negative binomial regressions on burned and unburned

plots for each pathway to test how the number of PyOM, alkane, inorganic and organic N

cycling genes changed over time. Nestedness level was assessed using null model and

Akaike criteria. For burned plots, models for PyOM, alkane and organic N included plot as a

random effect, while unburned plots included subplot. In contrast, inorganic N models for

burned plots included plot, subplot and time as random effects, while unburned models for

both organic and inorganic N included time and subplot as random effects. Time was scaled

and centered for all models. We visualized changes in microbial functional potential over

time by plotting the sum geTMM for each pathway of interest at each time point, separately

for burned and unburned plots, using phyloseq version 1.42.0 (McMurdie and Holmes,

2013).

Lastly, to identify the PyOM, alkane, inorganic and organic N genes that were

differentially abundant in burned versus unburned plots at each sampling time point, we

employed DESeq2 v1.38.3 (Love et al., 2014). To evaluate the response of functional gene

potential over time, we created separate DESeq2 models for each pathway at each time

point, determining significance with the Wald test and an adjusted p-value of < 0.05 and an

absolute log2 fold-change > 0. The total count of differentially abundant genes was

visualized as a bar plot for each pathway over time. We used the differentially abundant

genes from the burned communities to create metabolic maps for PyOM and inorganic N

degradation pathways at each time point. Genes identified as abundant by DESeq at each

time point were linked to their respective taxonomy using our MAG dataset via BLAST

(Bethesda, 2008). The taxonomy and functional annotation of the selected genes were

visualized using ggplot2. All R scripts are publicly available on GitHub:

https://github.com/pulidofabs/Chaparral-Metagenomes.

1.3 Results

Metagenomic sequencing of 30 burned and unburned chaparral soils generated 1.2

Tb of data and 2,266,623 annotated genes relevant to post-fire microbial metabolisms. The

significantly abundant genes, as determined by the DESeq results, were linked to 17 of the

446 MAGs identified from this dataset (see Supplementary Data 1C for additional details on

MAG quality and taxonomy). MAGs include the pyrophilous bacterial genera *Massilia* and

Noviherbaspirillium that previously dominated our ASV dataset (Pulido-Chavez et al.,

2023).

1.3.1 Wildfire altered the profile of functional genes for PyOM and inorganic N cycling

Fire and time significantly impacted abundances of functional genes associated with

degradation of PyOM (Fig. 1a), alkanes (Fig. 1b), and cycling of inorganic (Fig. 1c) and

organic N (Fig. 1d). Further, a significant fire-by-time interaction affected degradation of

PyOM and cycling of inorganic and organic N, such that genes changed over time in burned

but not unburned plots (Table S1). Fire was the greatest driver of differences in functional

gene profiles, accounting for 15% to 16% of the variation, with time and fire-by-time

interactions accounting for 5-6% of the variance in C and N gene functional profiles (Table

S1).

292   While unburned plots showed no turnover in functional gene profiles for PyOM

(Fig. 2a), or alkane degradation (Fig. 2c), or inorganic (Fig. 2e) and organic N cycling genes

(Fig. 2g), burned plots exhibited multiple functional turnover events for PyOM (Fig. 2b),

alkane (Fig. 2d), inorganic (Fig. 2f), and organic N genes (Fig. 2h; Table S2). Specifically,

PyOM degradation and inorganic N cycling genes turned over at 34 and 131 days, alkane

degradation genes at 25 and 131 days, and organic N processing genes at 25, 34, and 131

compared to 17 days post-fire (Fig. 2).

1.3.2 Time differentially affected PyOM cycling pathways

301   A significant fire-by-time interaction led to a 167% increase in total geTMM-

normalized gene abundance for PyOM degradation in burned plots from 17 to 376 days

post-fire (Fig. 3b), while alkane cycling genes increased by 91% (Fig. 3d). In contrast,

PyOM and alkane gene abundances remained relatively stable in unburned plots (Fig. 3a;

Table S3). This time-dependent increase was also observed at the individual pathway level,

where PyOM degradation pathways, including catechol, protocatechuate, and β-ketoadipate

degradation (Fig. S3; Table S5), and the propane pathway for alkane degradation increased

over time in burned but not unburned plots (Fig. S4; Table S6). This pattern likely reflects

post-fire bacterial diversity recovery (Fig. S5b), which increases in burned plots over time

as does the total number of PyOM degradation genes (Fig. S6).

1.3.3 Time differentially affected N cycling pathways

A significant fire-by-time interaction led to a 117% increase in total geTMM-

normalized gene abundance for inorganic N cycling in burned plots from 17 to 376 days

post-fire, while organic N cycling genes increased by 108% (Fig. 4; Tables S3, S4). In

contrast, both inorganic and organic N gene abundances remained relatively stable in

unburned plots. This temporal increase was also evident at the pathway level, with

nitrification, DNRA, and NO□□ assimilation increasing in burned plots (Fig. S7; Table S7).

This pattern may also be linked to bacterial diversity recovery (Fig. S4d), which increases in

burned plots over time as does the total number of inorganic N cycling genes (Fig. S8).

Interestingly, denitrification genes (NO□□ → NO□□ → NO; *narGZ, nrxA, nirK*)

significantly decreased over time in unburned plots but remained stable in burned plots (Fig.

S7; Table S7). Organic N cycling pathways did not differ between burned and unburned

plots over time (Fig. S9; Table S8), though two pathways increased in abundance overtime

in burned plots (Fig. S9).

1.3.4 Wildfire selects for ortho-cleavage aromatic degradation pathways

Wildfire resulted in more differentially abundant PyOM degradation genes in burned

compared to unburned soils (692 compared to 258; $p_{adj} < 0.05$; Fig. 5a), with significant

temporal variation in burned plots ($p_{adj} < 0.05$; Figs. 5a, 6). Ortho-cleavage pathways,

including protocatechuate, catechol and β-ketoadipate, were differentially abundant and

dominated burned plots from 17 to 131 days post-fire ($p_{adj} < 0.05$; Fig. 5a, 6). In contrast, the

meta-cleavage pathway, specifically protocatechuate meta-cleavage, was only dominant at

17 days post-fire ($p_{adj} < 0.05$; Fig. 5a, 6). Pathways involving phthalate, toluene, and trans-

cinnamate, which feed into the catechol or protocatechuate pathways, were also

differentially abundant in burned plots between 17 and 131 days post-fire ($p_{adj} < 0.05$; Fig.

5a, 6). Conversely, unburned plots were enriched in lignin degradation genes (e.g.,

ligABCIJKM and *galC*) at 17 and 25 days (Fig. 5a, 6). Additionally, alkane degradation

genes for acetate (*ACSS1_2, acs*, and *pta*) were present at all post-fire time points (Fig. 5b),

while genes for propane 2-monooxygenase, alkane hydroxylase, and alkane monooxygenase

were overrepresented only in burned plots at 17 days post-fire ($p_{adj} < 0.05$; Fig. 5b).

1.3.5 Wildfire increased genes for N retention

Wildfire increased the number of differentially abundant N cycling genes in burned

compared to unburned soils (143 vs. 55; Fig. 5c, 7). At all timepoints, genes involved in

NO_3^- retention pathways, including assimilatory (*nasABC, NRT, narK, nrtP, nirA* and *narB)*

and dissimilatory (DNRA) reductase genes (*nirBD and narIV)* were differentially abundant

in burned plots ($p_{adj} < 0.05$; Fig. 5c, 7). In contrast, denitrification genes (*nirK*, *narGHYZ*)

were differentially abundant in burned plots at early post-fire time points (17 and 25 days;

$p_{adj} < 0.05$; Fig. 5c, 7). However, genes associated with N☐O and N☐ production were not

differentially abundant at any time point throughout the year. Additionally, nitrification

genes (*hao, pmoA-amoA, pmoB-amoB*) were differentially abundant in burned plots at 131

days post-fire ($p_{adj} < 0.05$; Fig. 5c, 7). Lastly, while urea cycling pathways were

overrepresented in unburned plots at 17 and 25 days, they were differentially abundant in

burned plots from 17 to 131 days post-fire ($p_{adj} < 0.05$; Fig. 5d, 7).

1.3.6 PyOM genomic potential of dominant taxa differed in burned versus unburned soils

Genes for catechol and protocatechuate ortho-cleavage and the β-ketoadipate

pathway (*fad* genes) were overrepresented in MAGs from burned plots, particularly within

Proteobacteria, Actinobacteria, and Acidobacteriota (Fig. 8*;* Supplementary file 1c).

Pyrophilous *Massilia*, which dominated the 16S rRNA gene sequences in burned plots

across time (Fig. S5b), was represented by MAGs Holy_71 and Holy_91 (Fig. 8) and

encoded genes for catechol ortho-cleavage (*catABC*) and the β-ketoadipate pathways

(*pcaDLJ, fadA*). The protocatechuate ortho-cleavage pathway (*pcaCH*) was encoded by

Noviherbaspirillium (MAGs Holy_91, Holy_361, Holy_395, and Holy_399), which

dominated the burned plots at 131 and 376 days (Fig. S4b), the Burkholderiales-SG8-41

Solirubrobacteraceae (MAG Holy_399) and Chthoniobacterales-AV80 (MAG Holy_206;

Fig. 8).

Catechol meta-cleavage genes (*dmpBC*, *xylEG*) were encoded by MAGs for

Proteobacteria, Xanthobacteraceae-PALSA-894 (MAG Holy_368) and Verrucomicrobiota,

Chthoniobacterales-AV80 (*catE*, MAG Holy_206; Fig. 8). Lastly, protocatechuate meta-

cleavage pathway genes (*ligClK, galC*) in burned plots were differentially abundant by

Noviherbaspirillium (MAGs Holy_91, 381 and 395) and Xanthobacteraceae-*PALSA-894*

(MAG Holy_368; Fig. 8). In contrast, unburned plots were enriched for lignin degrading

genes (*pcaD, fadAl, praC, xylH, pracH)*, at 17-25 days, encoded by MAGs representing

Bradyrhizobium (MAG Holy_369) and Acidimicrobiales-AC-14 (MAG Holy_399).

1.3.7 Nitrogen genomic potential of dominant taxa differed in burned versus unburned

soils

Genes encoding NO_3^- retention pathways, including assimilatory NO_3^- reductase

(*nasABC, nirA*) and DNRA (*nirBD*), were differentially abundant in MAGs representing

pyrophilous taxa within the Actinobacteriota, such as *Arthrobacter* (*nasABC*; MAG

Holy_114)*,* Micromonosporaceae (*nasABC*; *nirBD*; MAG Holy_395) and *Pseudonocardia*

(*nirBD*; MAG Holy_362; Fig. 8). The Proteobacteria *Massilia* (MAG Holy_71) also

encoded genes for N retention, including DNRA (MAG Holy_71; *nirD*) and NO_3^-

assimilation (*NRT, narK, nrtP, nasA*; Fig. 8). Additionally, *Arthrobacter* (MAG Holy_114)

and the family *Micromonosporaceae* (MAG Holy_395) encoded genes for NO□□

assimilation retention pathways (*NRT, narK, nrtP, nasA*; Fig. 8). In contrast, unburned soils

had significant abundances of NO□□ assimilation, NO□□ dissimilatory reductase, and

denitrification pathways only at 17 days post-fire (Fig. 8), encoded by non-pyrophilous

genera particularly *Bradyrhizobium* (MAG Holy_369, Holy_46; NRT, narK, nrtP, nasAC,

napA; Fig. 8).

1.4 Discussion

Wildfire significantly altered the genomic functional potential of post-fire soil

bacteria across five timepoints, spanning from two weeks to one year post-fire, mirroring

significant shifts in bacterial community composition (Pulido-Chavez et al., 2023). Notably,

the genomic potential for PyOM degradation increased progressively over time in the

burned but not unburned plots, with catechol and protocatechuate ortho-cleavage (i.e., β-

ketoadipate) pathways emerging as the primary bacterial mechanisms for PyOM

degradation. Concurrently, N cycling genes increased over time in burned plots, with a

notable increase in genes for nitrification and a shift towards genes associated with N

retention, such as facilitating the conversion of NO☐☐ to microbial biomass and NO☐☐ to

NH☐. Finally, the increase in functional genes for PyOM degradation and N cycling aligned

with the dominance of pyrophilous bacterial genera, including *Massilia* and

Noviherbaspirillum, each encoding distinct pathways for PyOM and N cycling. These

findings suggest that resource acquisition traits drive bacterial secondary succession in post-

fire soils, underscoring the critical role of pyrophilous microbes in shaping soil C and N

dynamics and enhancing post-fire functional restoration.

1.4.1 Bacterial functional capacities shift from labile C to PyOM across succession

We previously hypothesized that post-fire microbial succession transitions from an

initial dominance of thermotolerant microbes to fast-colonizers, culminating in microbes

capable of exploiting post-fire resources, such as N and PyOM (Pulido-Chavez et al., 2023).

Using metagenomes, we corroborated this hypothesis, revealing increased abundance of

PyOM degradation genes in burned but not unburned plots over time, coinciding with the

recovery of bacterial species richness and diversity. This suggest that successful post-fire

genera require metabolic pathways for PyOM degradation to establish and dominate the

post-fire system.

Post-fire succession involved trade-offs in C use, with early colonizers favoring

labile C and later taxa shifting to more complex, aromatic C sources. During the first 17-25

days, bacteria encoding genes for acetate metabolism, a prevalent post-fire labile C source

(Blank et al., 1994; Certini, 2005; Kutscha and Pflugl, 2020) became dominant (*ACSS1_2,*

acs and *pta)*, indicating that early successional bacteria are well-equipped to degrade labile

C sources to support rapid growth and dominance. As labile C diminished (Fig. 5), bacterial

communities transitioned to taxa capable of degrading complex aromatic C sources like

PyOM (Fig. 8) (Pulido-Chavez et al., 2023). Indeed, by 131 days post-fire, taxa encoding

multiple PyOM degradation genes, such as *Noviherbaspirillum* and *Massilia* became

dominant, supporting findings from studies using PyOM treatments (Chen et al., 2021;

Cheng et al., 2022; Zeba et al., 2023) and incubation experiments (Song et al., 2017; Woolet

and Whitman, 2020) that highlight these genera's role in using PyOM as a C source.

In our study, *Noviherbaspirillium* increased in relative abundance over time,

dominating 16S sequences at 131 and 376 days post-fire, while *Massilia* remained

consistently dominant throughout the year (4). The timing difference likely reflects their

distinct PyOM degradation strategies. *Massilia* uses the simpler catechol and β-ketoadipate

ortho-cleavage pathway, providing immediate access to C resources, which may explain its

early and continuous dominance. In contrast, *Noviherbaspirillum* relies on the more

complex protocatechuate ortho- and meta-cleavage pathway, which is a slower process due

to its regulatory complexities (Johnson and Beckham, 2015). Since the protocatechuate

pathway is also associated with a growth lag and delayed substrate utilization (Johnson and

Beckham, 2015), this could potentially explain the delayed emergence of

Noviherbaspirillum, which might take longer to adapt and begin degrading PyOM

effectively. These findings suggest that the success of microbial taxa post-fire depends on

their ability to use different C sources, with some taxa adapting more quickly to available

resources due to their metabolic pathways. Together, these findings suggest that microbial

succession and dominance in post-fire environments are influenced by the functional

potential for distinct degradation pathways for labile C and PyOM overtime.

Previous research suggest interspecies resource competition between pyrophilous

bacteria and fungi, with bacteria encoding the catechol ortho-cleavage pathway and fungi

encoding the protocatechuate ortho-cleave pathway (Fuchs et al., 2011). Our bacterial-

optimized bioinformatic pipelines yielded no fungal MAGs, but our ASV dataset captured

fungal ASVs throughout the year (Pulido-Chavez et al., 2023), highlighting a

methodological limitation. Indeed, transcriptome data show that *Aspergillus*, a dominant

fungal genus in our ITS2 data (Pulido-Chavez et al., 2023), encodes PyOM degradation

genes via the β-ketoadipate pathway (Sgro et al., 2023). Thus, we hypothesize that fungi

contribute additional PyOM degradation genes, requiring bioinformatic advances to fully

explore this potential.

1.4.2 Post-fire chaparral bacteria preferentially degrade PyOM via the ortho-cleavage

degradation pathway

The greater number of differentially abundant PyOM degradation genes in burned

compared to unburned plots, together with the dominance of genes for the complete catechol

(*catABC*) and protocatechuate (*pcaHGBCDLJ*) ortho-cleavage pathways in burned soils

throughout the year, suggests that post-fire bacteria may preferentially use ortho-cleavage

over meta-cleavage pathways. This aligns with findings from post-fire coniferous forests,

where over 50% of MAGs contained the ortho-cleavage pathway (Nelson et al., 2022), a

pathway also enriched following oil spills, which similarly contain high levels of aromatic

compounds (Brock Melissa et al., 2025). There results suggest that high-severity fires,

which generate substantial PyOM or aromatics (Singh et al., 2012), may favor this pathway

in both chaparral and forest biomes. Indeed, the ortho-cleavage pathway is the primary route

for aerobic aromatic degradation (Granja-Travez et al., 2020; Weiland et al., 2022), known

for its efficiency in processing diverse aromatic compounds (Bugg et al., 2011), converting

C into biomass (Basha et al., 2010; Aghapour et al., 2013), and supporting rapid bacterial

growth (Song et al., 2002). However, while both catechol and protocatechuate ortho-

cleavage pathways are widespread across bacterial species, their gene arrangements vary

(Parke et al., 2000; Buchan et al., 2004), which can influence how effectively bacteria break

down and utilize PyOM as a C source in different substrates. Here, we found that dominant

Proteobacteria MAGs, such as *Noviherbaspirillum* and *Massilia* along with other MAGs

including Xanthobacteraceae PALSA-89, and the Acidobacteriota Chthoniobacterales AV80

and Solirubrobacteraceae, contain genes for catechol (*catABC*) and protocatechuate

(*pcaHGBCDLJ*) ortho-cleavage degradation pathways. Additionally, we found a timing

difference in *Massilia* and *Noviherbaspirillum* emergence (17 days vs 131 days), which

supports the idea that their distinct PyOM degradation strategies may reflect variations in

gene arrangements within these pathways, influencing their ability to utilize PyOM as a C

source. However, further research is needed to explore their specific gene arrangements.

Together, our results suggest that pyrophilous bacteria in wildfire-adapted ecosystems, like

chaparral, may have evolved a preference for the easier-to-degrade ortho-cleavage pathway

to convert PyOM into biomass, thereby facilitating the exploitation of PyOM as a C source

and promoting microbial secondary succession

1.4.3 Post-fire microbiomes express genes associated with inorganic N retention and

nitrification

We found a notable increase in the genomic potential for nitrification and N

retention over time in burned soils. As bioavailable N diminished, bacterial communities

likely transitioned to taxa specialized in N retention, potentially favoring increases in

biomass, which is consistent with the dominance of *Massilia* MAGs*,* which accounted for ~

60% of 16S sequences throughout the year (Pulido-Chavez et al., 2023) and encoded key

genes for N retention (Assimilatory: *nasABC, NRT, narK, nrtP, nirA* and *narB* and DNRA:

nirBD and narIV). By 131 days post-fire, succession was marked by the emergence of

bacterial genera *Arthrobacter*, *Blastocatellia*, and Micromonosporaceae MAGs, which

exhibited high genomic potential for N immobilization via the assimilatory NO_3^-

reduction pathway. These findings support observations of microbial N immobilization in

post-fire environments (Xu et al., 2022), including chaparral (Goodridge et al., 2018) and

studies showing the enrichment of key N retention genes (*nirB/D*) up to four years post-fire

(Dove et al., 2022). Moreover, studies involving PyOM additions have shown that PyOM

facilitates N retention in burned soils (Bai et al., 2015; Yu et al., 2023), suggesting that post-

fire N retention by altering microbiomes may represent a common strategy for stabilizing

nutrient cycling in fire disturbed ecosystems. Additionally, *Massilia, Pseudonocardia*, and

Micromonosporaceae have the capacity to reduce NO_3^- to NH_4^+ via dissimilatory

NO_3^- reductase (*nirB/D*), trading off a mobile and reactive anion for a cation more often

retained on soil exchange sites (Kalisz and Stone, 1980; Homyak et al., 2021). The observed

enrichment of N retention genes over time and the presence of microbial taxa capable of N

immobilization and assimilation, align with the known N reducing and assimilating

functions of *Massilia* and *Arthrobacter* (Cacciari et al., 1986; Bailey et al., 2014; Liu et al.,

2023b).

1.4.4. Denitrification and urea genes increase in early post-fire chaparral

Nitrogen losses from volatilization during fires and leaching afterward often surpass

the ecosystem's capacity to sequester N (Turner et al., 2007; Stephens and Homyak, 2023),

especially if fires disrupt N-cycling microbes (Knicker, 2007; Hanan et al., 2016). Our

findings indicate that the post-fire surge in bioavailable N (Wan et al., 2001) stimulates

bacterial activity, leading to an overrepresentation of denitrification genes (*nirK, narG,*

narZ) during early secondary succession (17-25 days). However, genes for complete nitrate

reduction to N_2 via denitrification ($NO \rightarrow N_2O \rightarrow N_2$) were notably rare, despite studies

reporting production of N_2O and N_2 in post-fire environments (Stephens and Homyak,

2023). While we could not detect the full suite of denitrification genes, complete reduction

to N_2 and partial reduction to N_2O may have still occurred, but may have gone undetected

because we mostly sampled during dry conditions that constrain denitrification (Krichels et

al., 2023). Nevertheless, our measurements suggest that early post-fire bacteria contain a

high number of denitrification genes, potentially creating conditions that accelerate N loss

via gaseous pathways. Additionally, urea cycling pathways were enriched in burned plots

from 17-to-131 days post-fire, but only from 17-to-25 days in unburned plots. This suggest

that post-fire microbes exploit urea as a N source, converting it to NH_4^+ for growth, thus

favoring microbes with urea cycling genes in the N rich environment. Together, our results

show that bacteria with specialized N metabolism dominate and mediate post-fire

succession, transitioning from denitrification to nitrification and ultimately expressing genes

for N retention as bioavailable N declines.

1.5 Conclusions

Our high-resolution gene-centric metagenomic analysis reveals rapid enrichment of

genomic functional potential for PyOM, alkanes, and inorganic N in burned soils over the

first post-fire year. These functional shifts align with the dominance of pyrophilous bacteria,

such as *Massilia* and *Noviherbaspirillium,* which encode pathways for ortho-cleavage

catechol and protocatechuate degradation, while potentially assimilating N into biomass.

This suggest that resource acquisition traits are key drivers of bacterial secondary succession

in fire-affected soils. The incomplete genomic potential for PyOM and N pathways within a

single bacterial genus, combined with distinct degradation routes employed by different

bacteria, suggests metabolic handoffs between groups. Specifically, some taxa degrade

complex compounds, while others utilize byproducts, indicating widespread resource

transfers in wildfire-affected communities. While the ability of post-fire bacteria to

capitalize on PyOM and N has been previously hypothesized (Woolet and Whitman, 2020;

Pulido-Chavez et al., 2023), and shown for some taxa (Cobo-Díaz et al., 2015; Fischer et al.,

2021; Nelson et al., 2022), our research provides the first direct evidence supporting our

conceptual model of trait trade-offs among bacteria (Enright et al., 2022), demonstrating

that specific functional traits drive bacterial succession in fire-impacted soils. These findings

have significant implications for predicting how bacterial communities influence C

sequestration and N cycling in post-fire ecosystems. Moreover, by revealing previously

unrecognized functional shifts in bacterial communities across a high-resolution

successional timeline, we fill a critical gap in post-fire ecological research and provide

essential insights into the long-terms impact of fire on microbial-mediated ecosystem

processes.

563
564
565   **1.6 Acknowledgments**
566
We thank the Joint Genome Institute (JGI) Community Sequencing Project (CSP) New
Investigator Award 507364 to SIG, the Bureau of Land Management (BLM) Joint Fire
Science Program (JFSP) Graduate Research Innovation (GRIN) Award 012641□002 to
MFPC and SIG, the United States Department of Agriculture (USDA)-NIFA Award #2022-
67014-36675 to SIG and PMH, and the Department of Energy (DOE) BER Award #DE-
SC0023127 to SIG, PMH and MJW for funding this project. We further thank the Cleveland
National Forest and the Trabuco Ranger District, including District Ranger Darrel Vance
and Emily Fudge, Jeffrey Heys, Lauren Quon, Jacob Rodriguez, and Victoria Stempniewicz,
for their help with permitting. We thank JGI Metagenome Program, including Nicole
Shapiro, Natasha Brown, Emiley Eloe-Fadrosh, and Simon Roux, for their help in
processing and submitting our metagenomic samples. Lastly, we thank Judy A. Chung, Aral
C. Greene, Elizah Stephens, Dylan Enright, and Sameer S. Saroa for fieldwork assistance
and James Randolph for molecular assistance. We thank Jason Stajich and anonymous
reviewers for helpful feedback on the manuscript.

581
582
583   **1.7 Competing Interests**

584
585   The authors declare no competing or conflicts of interest.

586
587
588
589
590   **1.8 Data Availability Statement**
591

The metagenomic reads and bacterial MAGs have been deposited to the National
Center for Biotechnology Information Sequence Read Archive under BioProject
PRJNA761539. Amplicon sequence reads for bacterial 16S rRNA and fungal ITS2
have been deposited to the National Center for Biotechnology Information Sequence
Read Archive under BioProject PRJNA761539. All statistical R scripts/codes are
available on GitHub and include the sample metadata required for analysis
https://github.com/pulidofabs/Chaparral-Metagenomes.

1.9 References

1. Abbasian, F., Lockington, R., Mallavarapu, M., Naidu, R., 2015. A Comprehensive Review of Aliphatic Hydrocarbon Biodegradation by Bacteria. Appl Biochem Biotechnol 176, 670-699.
2. Aghapour, A.A., Moussavi, G., Yaghmaeian, K., 2013. Biological degradation of catechol in wastewater using the sequencing continuous-inflow reactor (SCR). Journal of Environmental Health Science and Engineering 11, 3.
3. Anderson, M.J., 2017. Permutational Multivariate Analysis of Variance (PERMANOVA), Wiley StatsRef: Statistics Reference Online. John Wiley & Sons, Ltd, pp. 1-15.
4. Andrews, S., 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data [Online].
5. Aramaki, T., Blanc-Mathieu, R., Endo, H., Ohkubo, K., Kanehisa, M., Goto, S., Ogata, H., 2020. KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics 36, 2251-2252.
6. Ayiti, O.E., Babalola, O.O., 2022. Factors Influencing Soil Nitrification Process and the Effect on Environment and Health. Frontiers in Sustainable Food Systems 6.
7. Badía, D., Martí, C., Aguirre, A.J., Aznar, J.M., González-Pérez, J.A., De la Rosa, J.M., León, J., Ibarra, P., Echeverría, T., 2014. Wildfire effects on nutrients and organic carbon of a Rendzic Phaeozem in NE Spain: Changes at cm-scale topsoil. CATENA 113, 267-275.
8. Bai, S.H., Reverchon, F., Xu, C.-Y., Xu, Z., Blumfield, T.J., Zhao, H., Van Zwieten, L., Wallace, H.M., 2015. Wood biochar increases nitrogen retention in field settings mainly through abiotic processes. Soil Biology and Biochemistry 90, 232-240.
9. Bailey, A.C., Kellom, M., Poret-Peterson, A.T., Noonan, K., Hartnett, H.E., Raymond, J., 2014. Draft Genome Sequence of Massilia sp. Strain BSC265, Isolated from Biological Soil Crust of Moab, Utah. Genome Announcements 2, e01199-01114, 01192/01196/e01199-01114.
10. Basha, K.M., Rajendran, A., Thangavelu, V., 2010. Recent advances in the Biodegradation of Phenol: A review. 1, 219-234.
11. Bethesda, 2008. Building a BLAST database with your (local) sequences:BLAST Command Line Applications User Manual. National Center for Biotechnology Information.
12. Bird, M.I., Wynn, J.G., Saiz, G., Wurster, C.M., McBeath, A., 2015. The Pyrogenic Carbon Cycle. Annual Review of Earth and Planetary Sciences 43, 273-298.
13. Blank, R.R., Allen, F., Young, J.A., 1994. Extractable Anions in Soils following Wildfire in a Sagebrush-Grass Community. Soil Science Society of America Journal 58, 564-570.
14. Bowers, R.M., Kyrpides, N.C., Stepanauskas, R., Harmon-Smith, M., Doud, D., Reddy, T.B.K., Schulz, F., Jarett, J., Rivers, A.R., Eloe-Fadrosh, E.A., Tringe, S.G., Ivanova, N.N., Copeland, A., Clum, A., Becraft, E.D., Malmstrom, R.R., Birren, B., Podar, M., Bork, P., Weinstock, G.M., Garrity, G.M., Dodsworth, J.A., Yooseph, S., Sutton, G., Glockner, F.O., Gilbert, J.A., Nelson, W.C., Hallam, S.J., Jungbluth, S.P., Ettema, T.J.G., Tighe, S., Konstantinidis, K.T., Liu, W.T., Baker, B.J., Rattei, T., Eisen, J.A., Hedlund, B., McMahon, K.D., Fierer, N., Knight, R., Finn, R.,

650 Cochrane, G., Karsch-Mizrachi, I., Tyson, G.W., Rinke, C., Genome Standards, C.,
651 Lapidus, A., Meyer, F., Yilmaz, P., Parks, D.H., Eren, A.M., Schriml, L., Banfield,
652 J.F., Hugenholtz, P., Woyke, T., 2017. Minimum information about a single
653 amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of
654 bacteria and archaea. Nature Biotechnology 35, 725-731.

15. Brock Melissa, L., Tavares-Reager Joana, F., Dong, J., Larkin Alyse, A., Lam, T.,
Pineda, N., Olivares Christopher, I., Mackey Katherine, R.M., Martiny Adam, C.,
2025. Bacterial response to the 2021 Orange County, California, oil spill was
episodic but subtle relative to natural fluctuations. Microbiology Spectrum 0,
e02267-02224.

16. Buchan, A., Neidle, E.L., Moran, M.A., 2004. Diverse organization of genes of the
beta-ketoadipate pathway in members of the marine Roseobacter lineage. Applied
and Environmental Microbiology 70, 1658-1668.

17. Bugg, T.D., Ahmad, M., Hardiman, E.M., Rahmanpour, R., 2011. Pathways for
degradation of lignin in bacteria and fungi. Nat Prod Rep 28, 1883-1896.

18. Cacciari, I., Lippi, D., Pietrosanti, T., Pietrosanti, W., 1986. Ammonium assimilation
in an Arthrobacter sp. "fluorescens". Archives of Microbiology 145, 113-115.

19. Capo, E., Broman, E., Bonaglia, S., Bravo, A.G., Bertilsson, S., Soerensen, A.L.,
Pinhassi, J., Lundin, D., Buck, M., Hall, P.O.J., Nascimento, F.J.A., Björn, E., 2021.
Oxygen□deficient water zones in the Baltic Sea promote uncharacterized Hg
methylating microorganisms in underlying sediments. Limnology and Oceanography
67, 135-146.

20. Certini, G., 2005. Effects of fire on properties of forest soils: a review. Oecologia
143, 1-10.

21. Chatterjee, S., Santos, F., Abiven, S., Itin, B., Stark, R.E., Bird, J.A., 2012.
Elucidating the chemical structure of pyrogenic organic matter by combining
magnetic resonance, mid-infrared spectroscopy and mass spectrometry. Organic
Geochemistry 51, 35-44.

22. Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., Parks, D.H., 2019. GTDB-Tk: a
toolkit to classify genomes with the Genome Taxonomy Database. Bioinformatics,
btz848.

23. Chen, H., Wang, J.J., Ku, P.J., Tsui, M.T., Abney, R.B., Berhe, A.A., Zhang, Q.,
Burton, S.D., Dahlgren, R.A., Chow, A.T., 2022. Burn Intensity Drives the
Alteration of Phenolic Lignin to (Poly) Aromatic Hydrocarbons as Revealed by
Pyrolysis Gas Chromatography-Mass Spectrometry (Py-GC/MS). Environ Sci
Technol 56, 12678-12687.

24. Chen, Z., Tu, X., Meng, H., Chen, C., Chen, Y., Elrys, A.S., Cheng, Y., Zhang, J.,
Cai, Z., 2021. Microbial process-oriented understanding of stimulation of soil N2O
emission following the input of organic materials. Environmental Pollution 284,
117176.

25. Cheng, H., Tang, G., Wang, S., Rinklebe, J., Zhu, T., Cheng, L., Feng, S., 2022.
Combined remediation effects of biochar and organic fertilizer on immobilization
and dissipation of neonicotinoids in soils. Environ Int 169, 107500.

26. Cobo-Díaz, J.F., Fernández-González, A.J., Villadas, P.J., Robles, A.B., Toro, N.,
Fernández-López, M., 2015. Metagenomic Assessment of the Potential Microbial

Nitrogen Pathways in the Rhizosphere of a Mediterranean Forest After a Wildfire.
Microbial Ecology 69, 895-904.

27. Cunningham, C.X., Williamson, G.J., Bowman, D., 2024. Increasing frequency and
intensity of the most extreme wildfires on Earth. Nature Ecology & Evolution 8,
1420-1425.

28. De la Rosa, J.M., Faria, S.R., Varela, M.E., Knicker, H., González-Vila, F.J.,
González-Pérez, J.A., Keizer, J., 2012. Characterization of wildfire effects on soil
organic matter using analytical pyrolysis. Geoderma 191, 24-30.

29. Dove, N.C., Tas, N., Hart, S.C., 2022. Ecological and genomic responses of soil
microbiomes to high-severity wildfire: linking community assembly to functional
potential. ISME J 16, 1853-1863.

30. Eddy, S.R., 2011. Accelerated Profile HMM Searches. PLOS Computational
Biology 7, e1002195.

31. Enright, D.J., Frangioso, K.M., Isobe, K., Rizzo, D.M., Glassman, S.I., 2022. Mega-
fire in Redwood Tanoak Forest Reduces Bacterial and Fungal Richness and Selects
for Pyrophilous Taxa that are Phylogenetically Conserved. Molecular Ecology 31,
2475–2493.

32. Fernández-García, V., Marcos, E., Reyes, O., Calvo, L., 2020. Do Fire Regime
Attributes Affect Soil Biochemical Properties in the Same Way under Different
Environmental Conditions? Forests 11, 274.

33. Fischer, M.S., Stark, F.G., Berry, T.D., Zeba, N., Whitman, T., Traxler, M.F., 2021.
Pyrolyzed Substrates Induce Aromatic Compound Metabolism in the Post-fire
Fungus, Pyronema domesticum. Front Microbiol 12, 729289.

34. Fuchs, G., Boll, M., Heider, J., 2011. Microbial degradation of aromatic compounds
- from one strategy to four. Nature Reviews Microbiology 9, 803-816.

35. González-Pérez, J.A., González-Vila, F.J., Almendros, G., Knicker, H., 2004. The
effect of fire on soil organic matter—a review. Environment International 30, 855-
870.

36. Goodridge, B.M., Hanan, E.J., Aguilera, R., Wetherley, E.B., Chen, Y.-J.,
D'Antonio, C.M., Melack, J.M., 2018. Retention of Nitrogen Following Wildfire in
a Chaparral Ecosystem. Ecosystems 21, 1608-1622.

37. Granja-Travez, R.S., Persinoti, G.F., Squina, F.M., Bugg, T.D.H., 2020. Functional
genomic analysis of bacterial lignin degraders: diversity in mechanisms of lignin
oxidation and metabolism. Appl Microbiol Biotechnol 104, 3305-3320.

38. Haddock, J.D., 2010. Aerobic Degradation of Aromatic Hydrocarbons: Enzyme
Structures and Catalytic Mechanisms, Handbook of Hydrocarbon and Lipid
Microbiology, pp. 1057-1069.

39. Hanan, E.J., Tague, C., Schimel, J.P., 2016. Nitrogen cycling and export in
California chaparral: the role of climate in shaping ecosystem responses to fire.
Ecological Monographs 87, 76-90.

40. Harwood, C.S., Parales, R.E., 1996. The B-ketoadipate Pathway and The Biology of
Self-Identity. Annual Review of Microbiology 50, 553-590.

41. Homyak, P.M., Slessarev, E.W., Hagerty, S., Greene, A.C., Marchus, K., Dowdy,
K., Iverson, S., Schimel, J.P., 2021. Amino acids dominate diffusive nitrogen fluxes
across soil depths in acidic tussock tundra. New Phytologist 231, 2162-2173.

42. Jackrel, S.L., Gilbert, J.A., Wootton, J.T., 2019. The Origin, Succession, and
Predicted Metabolism of Bacterial Communities Associated with Leaf
Decomposition. mBio 10.
43. Johnson, C.W., Beckham, G.T., 2015. Aromatic catabolic pathway selection for
optimal production of pyruvate and lactate from lignin. Metabolic Engineering 28,
240-247.
44. Joshi NA, F.J., 2011. Sickle: A sliding-window, adaptive, quality-based trimming
tool for FastQ files
45. . Version 1.33.
46. Kalisz, P.J., Stone, E.L., 1980. Cation Exchange Capacity of Acid Forest Humus
Layers. Soil Science Society of America Journal 44, 407-413.
47. Kanehisa, M., Goto, S., 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes.
Nucleic Acids Research 28, 27-30.
48. Kang, D.D., Froula, J., Egan, R., Wang, Z., 2015. MetaBAT, an efficient tool for
accurately reconstructing single genomes from complex microbial communities.
PeerJ 3, e1165.
49. Khot, V., Zorz, J., Gittins, D.A., Chakraborty, A., Bell, E., Bautista, M.A., Paquette,
A.J., Hawley, A.K., Novotnik, B., Hubert, C.R.J., Strous, M., Bhatnagar, S., 2022.
CANT-HYD: A Curated Database of Phylogeny-Derived Hidden Markov Models
for Annotation of Marker Genes Involved in Hydrocarbon Degradation. Frontiers in
Microbiology 12.
50. Knicker, H., 2007. How does fire affect the nature and stability of soil organic
nitrogen and carbon? A review. Biogeochemistry 85, 91-118.
51. Knicker, H., 2011. Pyrogenic organic matter in soil: Its origin and occurrence, its
chemistry and survival in soil environments. Quaternary International 243, 251-263.
52. Krichels, A.H., Jenerette, G.D., Shulman, H., Piper, S., Greene, A.C., Andrews,
H.M., Botthoff, J., Sickman, J.O., Aronson, E.L., Homyak, P.M., 2023. Bacterial
denitrification drives elevated N(2)O emissions in arid southern California drylands.
Sci Adv 9, eadj1989.
53. Kutscha, R., Pflugl, S., 2020. Microbial Upgrading of Acetate into Value-Added
Products-Examining Microbial Diversity, Bioenergetic Constraints and Metabolic
Engineering Approaches. Int J Mol Sci 21.
54. Leleiwi, I., Rodriguez-Ramos, J., Shaffer, M., Sabag-Daigle, A., Kokkinias, K.,
Flynn, R.M., Daly, R.A., Kop, L.F.M., Solden, L.M., Ahmer, B.M.M., Borton,
M.A., Wrighton, K.C., 2023. Exposing new taxonomic variation with inflammation -
a murine model-specific genome database for gut microbiome researchers.
Microbiome 11, 114.
55. Li, D., Liu, C.-M., Luo, R., Sadakane, K., Lam, T.-W., 2014. MEGAHIT: An ultra-
fast single-node solution for large and complex metagenomics assembly via succinct
de Bruijn graph. arXiv:1409.7208 [q-bio].
56. Liu, W., Zhang, Z., Li, J., Wen, Y., Liu, F., Zhang, W., Liu, H., Ren, C., Han, X.,
2023a. Effects of fire on the soil microbial metabolic quotient: A global meta-
analysis. CATENA 224, 106957.
57. Liu, Y., Zhang, Y., Huang, Y., Niu, J., Huang, J., Peng, X., Peng, F., 2023b. Spatial
and temporal conversion of nitrogen using Arthrobacter sp. 24S4-2, a strain obtained
from Antarctica. Front Microbiol 14, 1040201.

58. Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol 15, 550.
59. Luo, Y., Huang, Y., Xu, R.-x., Qian, B., Zhou, J.-w., Xia, X.-l., 2020. Primary and
Secondary Succession Mediate the Accumulation of Biogenic Amines during
Industrial Semidry Chinese Rice Wine Fermentation. Applied and Environmental
Microbiology 86, e01177-01120.
60. McMurdie, P.J., Holmes, S., 2013. Phyloseq: An R Package for Reproducible
Interactive Analysis and Graphics of Microbiome Census Data. . PLoS One 8,
e61217.
61. Moucawi, J., Fustec, E., Jambu, P., Amblfs, A., Jacquesy, R., 1981. Biooxidation of
added and natural hydrocarbons in soils: Effect of iron. Soil Biology and
Biochemistry 13, 335-342.
62. Moya, D., Gonzalez-De Vega, S., Garcia-Orenes, F., Morugan-Coronado, A.,
Arcenegui, V., Mataix-Solera, J., Lucas-Borja, M.E., De Las Heras, J., 2018.
Temporal characterisation of soil-plant natural recovery related to fire severity in
burned Pinus halepensis Mill. forests. Sci Total Environ 640-641, 42-51.
63. Nave, L., Vance., E., ., C.S., Curtis., P., 2011. Fire effects on temperate forest soil C
and N storage. Ecol Appl 4, 1189-1201.
64. Neary, D.G., Klopatek, C.C., DeBano, L.F., Ffolliott, P.F., 1999. Fire effects on
belowground sustainability: a review and synthesis. Forest Ecology and
Management 122, 51-71.
65. Nelson, A.R., Narrowe, A.B., Rhoades, C.C., Fegel, T.S., Daly, R.A., Roth, H.K.,
Chu, R.K., Amundson, K.K., Young, R.B., Steindorff, A.S., Mondo, S.J., Grigoriev,
I.V., Salamov, A., Borch, T., Wilkins, M.J., 2022. Wildfire-dependent changes in
soil microbiome diversity and function. Nature Microbiology 7, 1419-1430.
66. Ni, G., Lappan, R., Hernández, M., Santini, T., Tomkins, A.G., Greening, C., 2023.
Functional basis of primary succession: Traits of the pioneer microbes. Environ
Microbiol 25, 171-176.
67. Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara,
R., Solymos, P., Stevens, M., Szoecs, P., Wagner, H., Barbour, M., Bedward, M.,
Bolker, B., Borcard, D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S.,
Evangelista, H., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M.,
Lahti, L., McGlinn, D., Ouellette, M., Ribeiro Cunha, E., Smith, T., Stier, A., Ter
Braak, C., Weedon, J., 2022. vegan: Community Ecology Package.
68. Olm, M.R., Brown, C.T., Brooks, B., Banfield, J.F., 2017. dRep: a tool for fast and
accurate genomic comparisons that enables improved genome recovery from
metagenomes through de-replication. The ISME Journal 11, 2864-2868.
69. Parke, D., D'Argenio, D.A., Ornston, L.N., 2000. Bacteria Are Not What They Eat:
That Is Why They Are So Diverse. Journal of Bacteriology 182, 257-263.
70. Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W., 2015.
CheckM: assessing the quality of microbial genomes recovered from isolates, single
cells, and metagenomes. Genome Research 25, 1043-1055.
71. Pérez-Pantoja, D., Donoso, R., Agulló, L., Córdova, M., Seeger, M., Pieper, D.H.,
González, B., 2012. Genomic analysis of the potential for aromatic compounds
biodegradation in Burkholderiales. Environ Microbiol 14, 1091-1117.

72. Pulido-Chavez, M.F., Randolph, J.W.J., Zalman, C., Larios, L., Homyak, P.M.,
Glassman, S.I., 2022. Bacterial and fungal communities experience rapid secondary
succession during the first year following wildfire in California chaparral. bioRxiv,
p. 2021.2012.2007.471678.
73. Pulido-Chavez, M.F., Randolph, J.W.J., Zalman, C., Larios, L., Homyak, P.M.,
Glassman, S.I., 2023. Rapid bacterial and fungal successional dynamics in first year
after chaparral wildfire. Molecular Ecology 32, 1685-1707.
74. Rawlings, N.D., Waller, M., Barrett, A.J., Bateman, A., 2014. *MEROPS* : the
database of proteolytic enzymes, their substrates and inhibitors. Nucleic Acids
Research 42, D503-D509.
75. Raza, T., Qadir, M.F., Khan, K.S., Eash, N.S., Yousuf, M., Chatterjee, S., Manzoor,
R., Rehman, S.u., Oetting, J.N., 2023. Unreveling the potential of microbes in
decomposition of organic matter and release of carbon in the ecosystem. Journal of
Environmental Management 344, 118529.
76. RCoreTeam, 2023. R: A language and environment for statistical computing. R
Foundation for Statistical Computing., Vienna, Austria.
77. Robinson, M.D., McCarthy, D.J., Smyth, G.K., 2010. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics (Oxford, England) 26, 139-140.
78. Santos-Medellin, C., Blazewicz, S.J., Pett-Ridge, J., Firestone, M.K., Emerson, J.B.,
2023. Viral but not bacterial community successional patterns reflect extreme
turnover shortly after rewetting dry soils. Nature Ecology & Evolution 7, 1809-1822.
79. Sgro, M., Chow, N., Olyaei, F., Arentshorst, M., Geoffrion, N., Ram, A.F.J.,
Powlowski, J., Tsang, A., 2023. Functional analysis of the protocatechuate branch of
the β-ketoadipate pathway in Aspergillus niger. Journal of Biological Chemistry 299,
105003.
80. Shaffer, M., Borton, M.A., McGivern, B.B., Zayed, A.A., Rosa, S.L.L., Solden,
L.M., Liu, P., Narrowe, A.B., Rodríguez-Ramos, J., Bolduc, B., Gazitua, M.C.,
Daly, R.A., Smith, G.J., Vik, D.R., Pope, P.B., Sullivan, M.B., Roux, S., Wrighton,
K.C., 2020. DRAM for distilling microbial metabolism to automate the curation of
microbiome function. bioRxiv, 2020.2006.2029.177501.
81. Singh, N., Abiven, S., Maestrini, B., Bird, J.A., Torn, M.S., Schmidt, M.W., 2014.
Transformation and stabilization of pyrogenic organic matter in a temperate forest
field experiment. Glob Chang Biol 20, 1629-1642.
82. Singh, N., Abiven, S., Torn, M.S., Schmidt, M.W.I., 2012. Fire-derived organic
carbon in soil turns over on a centennial scale. Biogeosciences 9, 2847-2857.
83. Smid, M., Coebergh van den Braak, R.R.J., van de Werken, H.J.G., van Riet, J., van
Galen, A., de Weerd, V., van der Vlugt-Daane, M., Bril, S.I., Lalmahomed, Z.S.,
Kloosterman, W.P., Wilting, S.M., Foekens, J.A., Ijzermans, J.N.M., Coene,
P.P.L.O., Dekker, J.W.T., Zimmerman, D.D.E., Tetteroo, G.W.M., Vles, W.J.,
Vrijland, W.W., Torenbeek, R., Kliffen, M., Carel Meijer, J.H., vd Wurff, A.A.,
Martens, J.W.M., Sieuwerts, A.M., on behalf of the, M.s.g., 2018. Gene length
corrected trimmed mean of M-values (GeTMM) processing of RNA-seq data
performs similarly in intersample analyses while improving intrasample
comparisons. BMC Bioinformatics 19, 236.

84. Song, H.-K., Song, W., Kim, M., Tripathi, B.M., Kim, H., Jablonski, P., Adams,
J.M., 2017. Bacterial strategies along nutrient and time gradients, revealed by
metagenomic analysis of laboratory microcosms. FEMS Microbiology Ecology 93.
85. Song, J., Sung, J., Kim, Y.M., Gerben J, Z., Kim, E., 2002. Roles of the meta- and
the ortho-Cleavage Pathways for the Efficient Utilization of Aromatic Hydrocarbons
by Sphingomonas yanoikuyae Bl. Korean Journal of Microbiology 38, 245-245.
86. Steinegger, M., Söding, J., 2017. MMseqs2 enables sensitive protein sequence
searching for the analysis of massive data sets. Nature Biotechnology 35, 1026-1028.
87. Stephens, E.Z., Homyak, P.M., 2023. Post-fire soil emissions of nitric oxide (NO)
and nitrous oxide (N2O) across global ecosystems: a review. Biogeochemistry.
88. Suzek, B.E., Huang, H., McGarvey, P., Mazumder, R., Wu, C.H., 2007. UniRef:
comprehensive and non-redundant UniProt reference clusters. Bioinformatics 23,
1282-1288.
89. Thomas, C.L., Jansen, B., van Loon, E.E., Wiesenberg, G.L.B., 2021.
Transformation of *n*-alkanes from plant to soil: a review. SOIL 7, 785-809.
90. Turner, M.G., Smithwick, E.A.H., Metzger, K.L., Tinker, D.B., Romme, W.H.,
2007. Inorganic nitrogen availability after severe stand-replacing fire in the Greater
Yellowstone ecosystem. Proceedings of the National Academy of Sciences 104,
4782–4789.
91. Wan, S., Hui, D., Luo, Y., 2001. Fire Effects on Nitrogen Pools and Dynamics In
Terretrial Ecosystems: A Meta-Analysis Ecological Applications 11, 1349-1365.
92. Wang L, N.Y., Tang YQ, Song XM, Cao K, Sun LZ, Wang ZJ, Wu XL. , 2016.
Diverse Bacteria with Lignin Degrading Potentials Isolated from Two Ranks of
Coal. . Front Microbiol, 7:1428.
93. Weiland, F., Kohlstedt, M., Wittmann, C., 2022. Guiding stars to the field of dreams:
Metabolically engineered pathways and microbial platforms for a sustainable lignin-
based industry. Metabolic Engineering 71, 13-41.
94. Whitman, T., Whitman, E., Woolet, J., Flannigan, M.D., Thompson, D.K., Parisien,
M.-A., 2019. Soil bacterial and fungal response to wildfires in the Canadian boreal
forest across a burn severity gradient. Soil Biology and Biochemistry 138, 107571.
95. Whitman, T., Woolet, J., Sikora, M., Johnson, D.B., Whitman, E., 2022. Resilience
in soil bacterial communities of the boreal forest from one to five years after wildfire
across a severity gradient. Soil Biology and Biochemistry 172.
96. Woodcroft, B.J., 2007. CoverM. School of Biomedical Sciences, Queensland
University of Technology Centre for Microbiome Research.
97. Woolet, J., Whitman, T., 2020. Pyrogenic organic matter effects on soil bacterial
community composition. Soil Biology and Biochemistry 141, 107678.
98. Xu, W., Elberling, B., Ambus, P.L., 2022. Fire increases soil nitrogen retention and
alters nitrogen uptake patterns among dominant shrub species in an Arctic dry heath
tundra. Sci Total Environ 807, 150990.
99. Yu, L., Homyak, P.M., Li, L., Gu, H., 2023. Succession of bacterial community
structure in response to a one-time application of biochar in barley rhizosphere and
bulk soils. Elementa: Science of the Anthropocene 11.
100. Zeba, N., Berry, T.D., Fischer, M.S., Traxler, M.F., Whitman, T., 2023. Soil carbon
mineralization and microbial community dynamics in response to PyOM addition.
Ecology.

Figure 1. Non-metric multidimensional scaling (NMDS) of geTMM normalized a) PyOM b) alkane c) inorganic nitrogen and d) organic nitrogen cycling genes within the microbial communities of burned and unburned soil samples (color) and time since fire (days; shape). Fire and time since fire effects and stress based on PERMANOVA (p <0.05). Ellipse represents the 95% confidence level.

Figure 2. Principal component analysis (PCoA) of the mean and standard error Bray-Curtis community dissimilarity at each timepoint for the unburned and burned a,b) PyOM; c,d) alkane; e, f) inorganic nitrogen and g, h) organic N (urea) functional potentials. Note that timepoints further apart with nonoverlapping standard error bars indicate a community turnover, whereas overlapping standard error bars represent a lack of compositional turnover, with the first sampling time point at 17 days representing the base level. Thus, the first turnover was initiated at 25 days. There were no significant turnover (TO) dates for any gene functional potentials in the unburned plots. The days with significant turnovers in the burned plots are listed in the upper right corners. Significance based on betadisper, which measures the dispersion/variability of the data within groups to measure community turnover.

Figure 3. Summed geTMM-normalized gene abundance over time for PyOM degradation in (a) unburned and (b) burned plots, and alkane degradation in (c) unburned and (d) burned plots. Colors represent gene function. Generalized Negative Binomial Models indicate a significant increase in relative abundance over time in burned plots, while unburned plots remained stable (Table S1).

Figure 4. Summed geTMM-normalized gene abundance over time for inorganic N cycling genes in (a) unburned and (b) burned plots, and organic N cycling in (c) unburned and (d) burned plots. Colors represent gene function. Generalized Negative Binomial Models indicate a significant increase in relative abundance over time in burned plots, while unburned plots remained stable (Table S1).

Figure 5. Total number of genes that were differentially abundant in the burned plots (brown) versus the unburned plots (blue green; x axis) per each metabolic pathway, colored by each sampling time point for a) PyOM, b) alkane, c) inorganic nitrogen and d) organic N (urea) cycling. Significance based on DESeq2 Wald test (adjusted P value < 0.05 and absolute log2 fold-change >0).

Figure 6. The catechol and protocatechuate degradation pathways with arrows indicating genes that were differentially expressed in the burned treatment at each time point (17, 25, 34, 131 and 376 days post-fire; Wald's test in DESeq2; $p<0.05$). Line type represents catechol (solid) and protocatechuate (dashed) cycling pathways and gray solid lines represent pathways that were not expressed during post-fire year one. Numbers in parentheses represent the mean log2fold change (DESeq2) for each represented gene and are colored by the timepoint in which that gene was significantly abundantly represented.

Figure 7. The nitrogen cycle with arrows indicating genes that were differentially represented in the burned treatment at each time point (17, 25, 34, 131, and 376 days post-fire; Wald's test in DESeq2; $p<0.05$). Line type represents the different nitrogen cycling pathways, colored by timepoint and gray solid lines represent pathways that were not expressed during the first year post-fire. Numbers in parentheses represent the mean log2fold change (DESeq2) for each represented gene and are colored by the timepoint in which that gene was significantly abundantly represented.

Figure 8. Taxonomic annotation for the most dominant, differentially abundant PyOM, inorganic nitrogen and organic N (urea) genes based on DESeq analysis (adjusted P value < 0.05 and absolute log2 fold-change >0). Taxonomy based on blasting contigs against the Metagenome Assembled Genomes (MAGs) represented in the x-axis.