Title:

The effect of extended cold storage and the use of extenders on

motility and swimming kinematics of the sperm of shortnose

sturgeon

Authors:

Kenzie H. Melanson[ab*]

Matthew K. Litvak[a]

[a]Mount Allison University, 62 York Street, Sackville, New

Brunswick E4L 1E2, Canada

[b]Saint Mary's University, 923 Robie St, Halifax, Nova Scotia B3H

3C3, Canada

*Corresponding author

Address: Department of Biology, Gairdner Building, Mount

Allison University, 53 York St, Sackville NB, E4L 1C9, Canada

Email: kmelanson@mta.ca

ABSTRACT

Shortnose sturgeon are listed as a species of special concern in

Canada and as endangered in the US. Increasing knowledge about

this species, particularly in the area of reproductive biology, will

better the management of wild populations and aid in the

development of assisted reproduction protocols. However, access

to wild sperm is limited, so short-term and long-term storage of

sperm from sturgeon is crucial for reproductive studies. Here we

report on testing and development of a short-term storage protocol

for shortnose sturgeon. Milt samples were collected from wild

shortnose sturgeon caught in the Wolastoq River. Subsets of semen

were mixed with different extenders with or without oxygen;

control treatments without extenders were also run. We used

computer-assisted sperm analysis (CASA) to determine sperm

motility and swimming kinematics for the different treatments. All

groups were examined immediately after collection and treatment

application, and then 1, 2, and 7 days after storage in a fridge (4ºC)

for experiment 1, and days 1, 3, 7, 10, 14, 17, 21, and 24 for

experiment 2. The response variables motility, curvilinear velocity

(VCL), linearity (LIN), and wobble (WOB) showed an overall

decrease over time with differences between extender treatments.

While untreated milt maintained some motility up to day 21, the

addition of an extender reduced decline in motility and improved

longevity up to day 24. Milt treated with the Park and Chapman

extender had the slowest motility decline of extenders used, and

milt treated with the modified Tsvetkova extender showed less

potential for contamination.

Keywords: Short-term storage: Shortnose sturgeon: Sperm

extender: Sperm motility

1. Introduction

Sturgeon, valued for their caviar and meat, are the most

endangered vertebrate group in the world, with all 27 species listed

on the International Union for Conservation of Nature (IUCN) Red

List (IUCN, 2023). This is due to habitat fragmentation from dams,

water regulation, pollution, and fisheries persecution, through

either directed fisheries and/or by-catch or poaching (Litvak, 2010;

Bronzi et al., 2011).

Shortnose sturgeon (*Acipenser brevirostrum*) has been

listed as a species of special concern since 1980 and endangered in

the US since 1967 (COSEWIC, 2015). Shortnose sturgeon in

Canada and the US are protected from commercial harvesting,

however they are still vulnerable to threats that affect other

sturgeon species (DFO, 2014; COSEWIC, 2015; Kynard et al.,

2016). Declines in sturgeon have increased interest in aquaculture

development and conservation (Giberson and Litvak, 2003; Litvak,

2010; Bronzi et al., 2011).

In many endangered fish species, including sturgeon, milt

quantity is limited in availability (Gilroy and Litvak, 2019) due to

low numbers and a short breeding season. This is exacerbated by

lack of synchronization of egg and sperm production in both the

lab and the field settings (Kime et al., 2001; Butts et al., 2012).

Development of protocols for both short-term and long-term

storage of sperm are crucial for artificial reproductive efforts in

conservation and aquaculture (Alavi and Cosson, 2005; Barulin

and Shumski, 2022). Short-term storage of refrigerated milt is

inexpensive. Stored milt can be used for assisted reproduction,

sperm quality analysis, and development of cryopreservation

protocols (Contreras et al., 2020). This is particularly important to

defeat timing availability issues of sperm and eggs and for control

of aquaculture and conservation genetic diversity. However, few

short-term sperm storage protocols exist for fishes.

Sturgeon sperm, like that of teleost fish, are immotile in the

testis and in seminal plasma (Ginzburg, 1972; Alavi and Cosson,

2005). Activation is controlled by hypo-osmolality or a decrease of

extracellular K+ ions when sperm are exposed to water during

reproduction (Gallis, 1991; Alavi et al., 2004; Alavi et al., 2012;

Alavi et al., 2019). Several factors can affect the quality and

viability of stored sperm, such as individual male variability and

storage conditions (Aramli et al., 2013). Other factors include

temperature, sperm concentration, and oxidative stress (Barulin

and Shumski, 2022), as well as types of extenders (Park and

Chapman, 2005).

Extender solutions serve as diluents to increase the volume

of the samples, protect the sperm cells, and provide optimal

conditions as a source of energy for fertilization (Park and

Chapman, 2005). Extenders should mimic the seminal fluid

composition, as ions at the right concentration will prevent motility

activation and maintain osmotic equilibrium between the medium

and the sperm cells (Bobe and Labbe, 2008; Alavi et al., 2019).

Park and Chapman (2005) developed an artificial extender to be

used with milt from Gulf of Mexico sturgeon and shortnose

sturgeon. Other extenders for sturgeon semen include Hank's

Balanced Salt Solution (HBSS) (Park and Chapman, 2005), and

modified Tsvetkova (Dorsey et al., 2011), as well as other

combinations of ions at similar osmolalities (Shazada et al., 2024).

Here, we focused on comparing the effect of different

extenders and oxygen on the motility and swimming kinematics of

shortnose sturgeon sperm over several days. We ran two

experiments. Experiment 1 (2022) compared motility and

swimming kinematics of control samples (no extenders) with those

treated with the Park and Chapman extender or HBSS extenders.

Experiment 2 (2023) compared sperm motility and swimming

kinematics of control samples (no extenders) with those treated

with the Park and Chapman extender or the modified Tsvetkova

extender, and all treatments with and without pure oxygen.

2. Materials and methods

2.1.   Animal care protocol

This study was performed under the guidelines of the

Canadian Council of Animal Care: Mount Allison Animal Care

Protocols 102193 and 103376.

2.2.   Fishing

Shortnose sturgeon were caught using gillnets (DFO

Scientific Fishing Permit 330697) with 5" stretch mesh between

May 13th and May 20th, 2022 for experiment 1, and May 19th and

20th 2023 for experiment 2, in the lower reaches of the Wolastoq

River (aka Saint John River) in New Brunswick, 11 km below the

Mactaquac dam. Fishing was attempted three times for experiment

1 (21 fish total), and four times (18 fish total) for experiment 2,

with nets being left in the river between 30 and 60 minutes per

attempt. Individuals were placed in a 400 L circular holding tank

on the boat and transferred two 1035 L insulated tanks (D337

Saeplast) onshore. A pump supplied the onshore tanks with river

water. All fish were tagged using passive integrated transponder

(PIT: Biomark) tags for identification during sperm collection.

Five males were used for experiment 1 and four males were used

in experiment 2.

2.3.   Sperm collection

Fish were taken out of the holding tank and identified with

a PIT tag reader (Biomark HPR Lite). Most fish were already

spermiating during handling, so no other identification was

necessary to determine the sex of the fish. Shop towels were used

to gently dry the urogenital opening to avoid urine contamination.

Collection syringes (either 20 or 60 ml) were equipped with a ~10

cm length of 1/8" inner diameter, 1/4" outer diameter Tygon

tubing to collect sperm. A small cut was made near the end of the

tubing to prevent the tubing from suctioning to the urogenital

canal. If there was not an immediate inflow of milt when the

syringe was drawn back, slight pressure was applied to the

abdomen to stimulate the flow of milt. For all fish, milt was

continuously collected until the syringe could not be drawn back

any further. Once collected, syringes with milt were taken to a

20x8' mobile lab, equipped with power, lights, and an air

conditioner, and immediately put into a fridge (~4°C). Analysis

was conducted immediately after milt collection. Milt was

collected from fish one at a time, to ensure that the application of

treatments and analysis would be done immediately.

2.4.   Treatments

2.4.1. *Experiment 1*

The extenders used for experiment 1 were HBSS (1X 21-020-CM, diluted to ~100mOsm) (Corning, 2013) and the Park and Chapman extender (Park and Chapman, 2005). There were six treatments in total per fish (Figure 1). Tygon tubing was used to attach the collection syringe to a new 20ml syringe, and 1/6 of collected milt was dispersed into the new syringe, Tygon tubing was reused so long as there was only milt from the same fish being transferred. Milt was kept in the fridge until it was needed. Four 50 ml centrifuge tubes were filled with one of two extenders at a 1:2 (milt to extender) ratio to the measured milt in the syringes. Four of six syringes were individually added to one of four tubes, so each tube had 1/6 amount of milt and an extender. These tubes were shaken to mix, and then pulled back into the labeled syringes. The two syringes with no extender had the same amount of milt but were left undiluted. Each syringe contained less than 10 ml of final solution to ensure each syringe was half empty to allow air exchange and circulation. Syringes were stored in Ziploc bags in the fridge, with the plunger of the syringe towards the coldest part of the fridge. Air exchange was done twice a day for each syringe, by pushing out the old air and sucking in new air. This was done in the morning and evening. Samples were only taken out of the

182   fridge when analysis was being conducted, and they were

183   immediately placed back in the fridge once aliquots were taken.

184   *2.4.2. Experiment 2*

The extenders used for experiment 2 were modified

Tsvetkova (Dorsey et al., 2011: 600 ml of a solution made with tris

acid buffer and 40 ml of a solution made with tris base buffer, to

achieve a neutral pH) and the Park and Chapman extender (Park

and Chapman, 2005). There were twelve treatments in total per

fish (Figure 1). Tygon tubing was used to attach the collection

syringe to a new 20 ml syringe, and 1/12 of collected milt was

dispersed into the new syringe, as in Experiment 1, Tygon tubing

was reused so long as there was only milt from the same fish being

transferred. Milt was kept in the fridge until it was needed. Eight

50 ml centrifuge tubes were filled with one of two extenders at a

1:3 (milt to extender) ratio to the measured milt in the syringes.

Eight of twelve syringes were individually added to one of eight

tubes, so each tube had 1/12 amount of milt and an extender. These

tubes were shaken to mix, and then pulled back into the labeled

syringes. The four syringes with no extender had the same amount

of milt but was left undiluted. Six of twelve syringes were treated

with oxygen using Tygon tubes fed into a Ziploc bag full of

oxygen. All twelve syringes contained less than 10 ml of final

solution to ensure each syringe was half empty to allow air

exchange and circulation. All syringes were stored in Ziploc bags

in the fridge, with the plunger towards the coldest part of the

fridge. Air exchange was done after each analysis (every 3-4 days)

by pushing out the old air/oxygen and drawing in new air/oxygen.

Samples were only taken out of the fridge when the analysis was

being conducted, and they were immediately placed back in the

fridge once aliquots were taken for analysis.

2.5.   Analysis of Sperm Activity

Motility and kinematic parameter analysis of sperm was

conducted with an Olympus CX41 microscope and CEROS II

(Hamilton Thorne) CASA system at a magnification of 20x using

20 μm Hamilton Thorne Inc. 2X-CEL slides. Sperm density for

each fish was calculated by adding 2 μL of undiluted sample and

18 μL of 1% bovine serum albumin (BSA), for activation and to

prevent sticking of sperm, to a blank slide. A different dilution was

used for each fish, calculated to reach a target visible cell count of

40-60 sperm per field of view. During analysis slides were kept at

river temperature (8^{o}C) using a Linkam PE95/T95 system

controller. The room temperature and humidity was controlled

with an air conditioner and de-humidifier to prevent condensation

on the cooled slides. 2.0 mL vials and 0.5 mL vials were labelled

with the treatment type and each replicate. These vials were placed

in a microtube holder which was placed on a paper towel and

plastic sheet over ice in the cooler. The plastic bag and paper towel

ensured no direct contact of the holder/vials with ice. A few drops

(1-3) of milt were added to each 0.5 mL labeled vial,

corresponding to the treatment in the syringe, and the syringes

were placed back in the fridge. BSA was then measured into each

2.0 mL vial in correspondence to the values calculated in an Excel

sheet to reach the target dilution (as mentioned previously) based

on sperm density for each fish. Samples for analysis were mixed

one at a time to avoid activation until they could be analyzed. The

calculated amount of sample to reach the appropriate dilution was

taken from the 0.5 mL vial and added to the 2.0 mL vial. The 2.0

mL vial with sample was then shaken slightly to mix. A 20 μL

volume of this mixture was then pipette onto a blank slide and put

under the microscope to be analyzed. Video captures of 60

frames/second, lasting ~2 seconds were taken immediately on the

CASA system after a cover slip was placed on, and a stopwatch

was started. The sample was followed for 1 minute for both

experiments with 3-5 videos captured during this time. These

analyses were repeated for each sample on days 0, 1, 2, and 7 for

experiment 1, and on days 0, 3, 7, 10, 14, 17, 21, and 24 for

experiment 2. A CSV file was created automatically by CASA

system at the end of each minute increment, with data from all

videos for each minute.

2.6. Seminal fluid testing

Seminal fluid testing was performed for both experiments. Milt

was spun in a Thermo Scientific Sorvall Legend Micro 17

Microcentrifuge until a sperm pellet formed (~ 9 minutes on 1000

rpm). Seminal fluid was then separated by pipette into 2 μL tubes.

Osmolality was tested using Model 3320 Advanced Instruments

Inc. osmometer for both experiments, and pH was tested for

experiment 2 only, with a Mettler Toledo pH/ion tester.

2.7. Variables analyzed

In addition to motility, we chose to analyze three related

swimming kinematic factors: curvilinear velocity (VCL), linearity

(LIN), and wobble (WOB). VCL was chosen as it is most

representative of the sperm velocity. LIN was chosen as it is the

straight-line velocity divided by the VCL. WOB was chosen as it is

the average path velocity divided by the VCL. Motility, LIN, and

WOB are represented as percentages, and VCL is in μm/s. Sperm

were classified as motile if they had a VCL>10 µm/s (Xin et al.,

2020; Shazada et al., 2024).

2.8. Statistical analysis

CSV files from CASA were combined using a Python

script, to create one dataset with all data. Only motile sperm were

used to assess kinematics, and motility was calculated as a mean

sum of motile and nonmotile sperm. Figures were created using

full, untransformed data. Data were analyzed using RStudio (R

Core team, 2023), version 4.2.2, and significance was assessed at

$P<0.05$. Linear mixed models were made for all four dependent

variables (VCL, WOB, LIN and motility), using the function *lmer*

in the package *lme4* (Bates et al., 2015). Post-hoc tests to

determine differences between extenders were done using

emmeans (Lenth, 2023), which uses estimated marginal means

(Searle et al., 1980) to interpret differences. An ANOVA was run

on experiment 2 to examine the interaction between day and

extender. Days 14-24 were omitted from experiment 2 linear

mixed models due to the large number of zeros, and increased

contamination after day 10. An ANOVA was run comparing the

last day of motility (VCL<10 µm/s) and post-hoc Tukey tests were

run to determine differences. Normality and homogeneity of

variances were checked using visual analysis of residuals

(histogram, qq-line plot, and plot of points against residuals of the

model), and by examining the skewness of the residuals.

Dependent variables that had residuals with a skewness that was

not symmetric (greater than 0.5 or less than -0.5) or that did not

pass visual homogeneity of variance and normality tests were

transformed using *BoxCoxTrans* in the package *caret* (Kuhn,

2008). For experiment 1, all dependent variables were transformed

except VCL. For experiment 2, WOB, the last day of motility, and

day 7 for each variable as well as all days of motility for the

experiment 2 ANOVA needed to be transformed.

The percent decrease in motility for each treatment was

calculated by dividing the difference in motility from day 0 and

day 7 by the starting motility (day 0), and was an average of all

fish from each experiment.

3.   Results

3.1.   Experiment 1

The number of days sperm were held showed a statistically

significant decrease on all parameters (Table 1, Figure 2a). Post-

hoc comparisons revealed that VCL, LIN, and WOB on days 0, 1,

and 2 were significantly higher than values on day 7. There was a

significant effect of extender on VCL, and post-hoc testing

revealed that the VCL of sperm treated with either extender

decreased at a slower rate than sperm left untreated (Table 1).

When comparing the last day of motility between treatments there

was no statistically significant effect of extender (Table 2). The

percent decrease in motility over the 7-day period was lowest for

extender P (41.21%), and highest for no extender (N) (72.38%).

3.2.   Experiment 2

The amount of days sperm were held showed a statistically

significant decrease on all parameters (Table 1, Figure 2b). There

were statistically significant interactions between day and extender

for all parameters (Table 1) necessitating breaking apart the data

and running individual ANOVAs by day. ANOVAs run on each

day showed that on day 3 the LIN and WOB of sperm treated with

the Park and Chapman extender was significantly higher than that

of sperm treated with the modified Tsvetkova extender, and the

VCL of sperm treated with the Park and Chapman extender was

significantly higher than that of sperm left untreated (Table 3). On

day 10, the VCL, LIN, and WOB of sperm treated with either

extender was significantly higher than that of sperm left untreated

(Table 3).

When comparing the last day of motility between treatments

there was a statistically significant effect of extender, and no

statistically significant effect of oxygen. Post-hoc testing

demonstrates that motility decreased significantly less for sperm

treated with extender P or extender M than for sperm treated with

no extender (N) (Table 2). The percent decrease in motility over

the first 7 days was lowest for extender P with oxygen (42.42%),

and highest for no extender (N) without oxygen (72.29%).

3.2.1. Contamination

Contamination by protozoans was seen on slides as early as

day 10, for both oxygen treatments with extender P. All aliquots

treated with extender P were contaminated by day 17 regardless of

oxygen treatments. Protozoans were seen in one aliquot treated

with extender M and no oxygen on day 17, and on day 21 five

more aliquots treated with extender M had protozoans visible, four

of these were oxygen treated, and by day 24, eight aliquots treated

with extender M were recorded as having protozoans. Only three

aliquots treated with no extender (N) had visible protozoans, two

starting on day 14 and one on day 17. Sperm treated with the Park

and Chapman extender had no motility after 14 days, likely due to

protozoan contamination. The sperm treated with the modified

Tsvetkova extender had minimal protozoan contamination when

compared to the Park and Chapman extender.

3.3.   Osmolality and pH measurements

The modified Tsvetkova extender had a pH of 7.7, and an

osmolality of 68. The Park and Chapman extender had a pH of 7.6

and an osmolality of 108 for experiment 2 (only tested in

experiment 2). HBSS was adjusted to have an osmolality of around

100. The osmolalities and pH for each extender fell within the

range of the osmolalities (40-147 mOsm/kg) and pH (7.95-8.68)

for shortnose sturgeon seminal fluid in this study.

4. Discussion

The goal of our study was to determine a short-term storage

protocol that allows for longevity of sperm motility and swimming

kinematics for shortnose sturgeon sperm.

The percent decrease for motility for both experiments indicate

that the Park and Chapman extender maintained the highest level

of motility and swimming kinematic performance over the first 7

days compared to the other treatments. However, the sperm treated

with the modified Tsvetkova extender had better motility and

performance from day 10 until the end of the experiment. Sperm

treated with either extender performed better than sperm stored

without extenders. The last day of motility analysis shows that

sperm treated with the modified Tsvetkova extender maintains

motility for the longest period of time; it was the only treatment

that lasted until day 24. Sperm treated with the Park and Chapman

dropped in motility earlier than sperm treated with Tsevtkova,

likely due to protozoan contamination. Samples treated with the

modified Tsvetkova extender had lower visible protozoan

contamination than the samples treated with the Park and Chapman

extender or the control non-extender/treated samples. This finding

could be important in future extender research as the Tsevtkova

extender appears to not only prolong the life of sperm, but also

decreases contamination by protozoans.

Even with no treatment, motility did persist up to 24 days,

indicating that even without an extender swimming kinematics can

be maintained. This is important to note as the main goal of the

study was to determine how long sperm can last in short-term

storage so it can be transferred to the lab to be cryopreserved,

without having to take cryopreservation equipment into the field.

When examining sperm kinematics and parameters, it is

important to keep in mind the biological significance of these

sperm kinematics in terms of sperm quality. The classification of

sperm quality varies by researcher. Motility values less than 20%

(Xin et al., 2020) or 30% (Dorsey et al., 2011) have been classified

as poor quality, values between 20-80% (Xin et al., 2020) or 30-

90% (Dorsey et al., 2011) have been classified as moderate quality,

and values above 80% (Xin et al., 2020) or 90% (Dorsey et al.,

2011) have been classified as high quality. The definition of

motility also varies among researchers. In the present study we

have defined motility as sperm with a VCL >10 μm/s (Xin et al.,

2020; Shazada et al., 2024). Other studies have classified sperm as

motile with a VCL >3 μm/s (Dzyuba et al., 2012; Shaliutina et al.,

2013; Dzyuba et al., 2015; Barulin and Shumski, 2022), or as

uncontrollably spinning in place (DiLauro et al., 1994), and some

do not report their motility criteria (Park and Chapman, 2005;

Aramli et al., 2013) making comparisons between studies difficult.

Longevity of sperm motility should also be considered

when establishing motility criteria. For example, gamete contact

time has a large effect on fertilization success (Butts et al., 2009),

as sperm needs to be able to reach an egg. For sturgeon, their

sperm can remain motile for up to several minutes and even hours

in some cases (Ginzburg, 1972; Alavi et al., 2004; Gilroy and

Litvak, 2019). Because of this relatively long period of motility, a

VCL of both 3 μm/s or 10 μm/s may be sufficient to achieve

fertilization. Therefore, defining minimum VCL for motility could

significantly impact interpretation of sperm quality. It is clear that

scientists working in this area of research need to specify motility

criteria for comparisons within and between species.

While sperm in our study were still motile for up to 24

days, sperm after day 10 would be classified as poor quality in

terms of fertilization capacity, as motility was less than 20%

(Dorsey et al., 2011; Xin et al., 2020); after day 10, more than 80%

of sperm had a VCL <10 μm/s (our motility definition).

Extenders tested in this experiment have been used to prolong

sperm motility and kinematics in *Acipenser* sperm in previous

studies. While all extenders used in these previous studies had

similar pH and osmotic pressure there was still a noticeable

difference in motility between extenders used (Table 4). Many of

these studies found that the Park and Chapman extender had the

longest lasting motility. This is likely because of the sugars used,

and the phosphate content (Dorsey et al., 2013). Dorsey et al.

(2013) speculated that sturgeon sperm cells possess glucose pores

such as those found in sperm cells found in other species (Kim and

Moley, 2008; Sancho et al., 2008), which demonstrates that

glucose and phosphate are important for ATP stores and

production, which are crucial for sperm motility. This would

explain the difference in results we saw between HBSS and the

Park and Chapman extender, as HBSS contains dextrose rather

than glucose. This could also explain why sperm treated with the

Park and Chapman extender in experiment 2 appear to decrease

motility at a slower rate (prior to contamination) than the modified

Tsvetkova extender, as the modified Tsvetkova extender contains

sucrose rather than glucose.

In experiment 1 we did not increase oxygen content. For

experiment 2 we added oxygen content as a factor, in addition to

testing a different extender. Our results on the addition of oxygen

were not conclusive, however we did see higher motility and

swimming kinematic with sperm stored with oxygen after day 17

for all treatments (Figure 2b). In other studies, motility of

Acipenser sperm stored in an oxygenated atmosphere was superior

to that of other studies stored in an atmosphere with only air

(DiLauro et al., 1994; Barulin and Shumski, 2022). However, the

addition of oxygen can cause an increase of reactive oxygen

species. The increase in both reactive oxygen species and

heterotrophic protists is one of the most limiting factors of the

storage life of sperm without cryopreservation (Barulin and

Shumski, 2022), as was likely the case with experiment 2. While

these protozoans did not appear to physically latch on or interact

with the motile (or non-motile) sperm, it is likely that they would

be competing for resources from the seminal plasma and extender

mix (primarily the sugars used in the extenders). This would cause

sperm longevity to decrease due to the lack of energy availability,

as we observed in the samples treated with the Park and Chapman

extender. It appears that while glucose is the preferred energy

source for sperm stored in this experiment, it is likely also the

preferred energy for the protozoans found in our experiment.

The most promising results from previous research were found

by Park and Chapman (2005), who were able to maintain a long

motility time for samples with a mixture of Shortnose sturgeon and

Gulf of Mexico sturgeon sperm for 28 days without oxygen. This

is likely because their study appears to be the only study on

Acipenser sperm that used a shaker. Consistently shaking the

stored milt in its tube allows for both oxygen and the fluid itself to

be moved around within the tubes, therefore allowing sperm

greater access to oxygen and resources from seminal fluid or added

extenders. While their study was successful, it is important to

consider that they did not examine sperm longevity separately by

species. It may be that the presence of seminal fluid from Gulf of

Mexico sturgeon enhanced longevity of the shortnose sturgeon

sperm used in their experiment.

Even with the promising results from our study, future research

should include the addition of a shaker to increase oxygen and

particle movement within the stored seminal fluid, as well as

testing antibiotics to lower the likelihood of contamination from

bacteria.

5. Conclusion

Our findings lend insight into which extender is most effective

during short-term storage for shortnose sturgeon sperm, how

oxygen influences sperm longevity, and how these elements can

contribute to developing a protocol for short-term storage of sperm

in the field. This is crucial information to determine, because of the

limited availability of sperm and the extreme risk of extinction for

this vertebrate group. One of the main avenues to save sturgeons is

artificial reproduction, which requires viable sperm that can be

held until females can be stripped. Though cryopreservation is

useful for banking sperm samples, it is impractical for field

settings as the process is intricate and requires expensive and

fragile equipment. Discovering methods to refine short-term

storage protocols will allow sturgeon sperm to remain viable

during transport for various applications such as assisted

reproduction, reproductive research, and development of

cryopreservation protocols. This work documents short-term

storage of shortnose sturgeon sperm directly in the field using a

mobile lab and demonstrates how this protocol can be further

adjusted to be as practical and efficient as possible.

Funding

Natural Sciences and Engineering Research Council of Canada

Discovery Grant (RGPIN-2019-07138); New Brunswick

Innovation Fund Research Assistantship Program; Career

Launchers Canada Internship Program grants to MKL.

Acknowledgments

The authors thank Danny Glassman, Alex Giroux, Naomi

Meed, and Ahmad Hayat for their help with data collection.

CRediT authorship contribution statement

Kenzie H. Melanson and Matthew K.

Litvak: Conceptualization, Methodology, statistical analysis,

writing, and editing.

Declaration of competing interest

Authors declare no conflict of interest involved in this study.

References

Alavi, S., Cosson, J., 2005. Sperm motility in fishes. I. Effects of

temperature and pH: a review. Cell. Biol. Int. 29, 101–110.

https://doi.org/10.1016/j.cellbi.2004.11.021

Alavi, S.M.H., Cosson, J., Karami, M., Mojazi Amiri, B.,

Akhoundzadeh, M.A., 2004. Spermatozoa motility in the

Persian sturgeon, *Acipenser persicus*: effects of pH, dilution

rate, ions and osmolality. Reproduction 128, 819–828.

https://doi.org/10.1530/rep.1.00244

Alavi, S.M.H., Hatef, A., Pšenička, M., Kašpar, V., Boryshpolets,

S., Dzyuba, B., Cosson, J., Bondarenko, V., Rodina, M., Gela,

D., Linhart, O., 2012. Sperm biology and control of

reproduction in sturgeon: (II) sperm morphology, acrosome

reaction, motility and cryopreservation. Rev. Fish Biol. Fish.

22, 861–886. https://doi.org/10.1007/s11160-012-9270-x

Alavi, S.M.H., Cosson, J., Bondarenko, O., Linhart, O., 2019.

Sperm motility in fishes: (III) diversity of regulatory signals

from membrane to the axoneme. Theriogenology 136, 143–

165. https://doi.org/10.1016/j.theriogenology.2019.06.038

Aramli, M.S., Kalbassi, M.R., Nazari, R.M., Aramli, S., 2013.

Effects of short-term storage on the motility, oxidative stress,

and ATP content of Persian sturgeon (*Acipenser persicus*)

sperm. Anim. Reprod. Sci. 143, 112–117.

https://doi.org/10.1016/j.anireprosci.2013.10.010

Barulin, N.V., Shumski, K.L., 2022. Morphological, Biochemical

and Functional Changes in Sperm of the Siberian Sturgeon

Acipenser baerii (Acipenseridae) during Its Short-Term

Storage. J. Ichthyol. 62, 1476–1489.

https://doi.org/10.1134/S0032945222060030

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting

Linear Mixed-Effects Models Using lme4. J. Stat. Soft. 67.

https://doi.org/10.18637/jss.v067.i01

Bobe, J., Labbe, C., 2008. Chilled Storage of Sperm and Eggs, in:

Cabrita, E., Robles, V., Herráez, P. (Eds.), Methods in

Reproductive Aquaculture, Marine Biology. CRC Press, pp.

219–235. https://doi.org/10.1201/9780849380549.sec4

Bronzi, P., Rosenthal, H., Gessner, J., 2011. Global sturgeon

aquaculture production: an overview: Global sturgeon

aquaculture production. J. Appl. Ichthyol. 27, 169–175.

https://doi.org/10.1111/j.1439-0426.2011.01757.x

Butts, I.A.E., Roustaian, P., Litvak, M.K., 2012. Fertilization

strategies for winter flounder: effects of spermatozoa density

and the duration of gamete receptivity. Aquat. Biol. 16, 115–

124. https://doi.org/10.3354/ab00439

Butts, I.A.E., Trippel, E.A., Litvak, M.K., 2009. The effect of

sperm to egg ratio and gamete contact time on fertilization

success in Atlantic cod *Gadus morhua* L. Aquaculture 286,

89–94. https://doi.org/10.1016/j.aquaculture.2008.09.005

Contreras, P., Dumorné, K., Ulloa-Rodríguez, P., Merino, O.,

Figueroa, E., Farías, J.G., Valdebenito, I., Risopatrón, J.,

2020. Effects of short-term storage on sperm function in fish

semen: a review. Rev. Aquac. 12, 1373–1389.

https://doi.org/10.1111/raq.12387

Corning Incorporated, 2013. HBSS (Hank's Balanced Salt

Solution) Formulation.

COSEWIC, 2015. COSEWIC assessment and status report on the

shortnose sturgeon *Acipenser brevirostrum* in Canada.

Committee on the Status of Endangered Wildlife in Canada.

Ottawa. xii + 48 pp. www.registrelep-

sararegistry.gc.ca/default_e.cfm.

DFO, 2014. Distinctiveness and Status of the Saint John River

Population of Shortnose Sturgeon *Acipenser brevirostrum*.

DFO Can. Sci. Advis. Sec. Sci. Resp. 2014/43.

DiLauro, M.N., Krise, W.F., Hendrix, M.A., Baker, S.E., 1994.

Technical Notes: Short-Term Cold Storage of Atlantic

Sturgeon Sperm. Prog. Fish Cult. 56, 143–144.

https://doi.org/10.1577/1548-

8640(1994)056<0143:TNSTCS>2.3.CO;2

Dorsey, K.M., Guthrie, H.D., Welch, G.R., Mohler, J., Theisen,

D.D., Siewerdt, F., Vinyard, B.T., Woods, L.C., 2011. Quality

Assessment of Wild Atlantic Sturgeon Semen under

Conditions of Short-Term Storage. N. Am. J. Aquacult. 73,

418–425. https://doi.org/10.1080/15222055.2011.629945

Dzyuba, B., Boryshpolets, S., Shaliutina, A., Rodina, M.,

Yamaner, G., Gela, D., Dzyuba, V., Linhart, O., 2012.

Spermatozoa motility, cryoresistance, and fertilizing ability in

sterlet *Acipenser ruthenus* during sequential stripping.

Aquaculture 356–357, 272–278.

https://doi.org/10.1016/j.aquaculture.2012.05.006

Dzyuba, V., Cosson, J., Dzyuba, B., Yamaner, G., Rodina, M.,

Linhart, O., 2015. The antioxidant system of seminal fluid

during in vitro storage of sterlet *Acipenser ruthenus* sperm.

Fish. Physiol. Biochem. 42, 563–568.

https://doi.org/10.1007/s10695-015-0159-1

Gallis, J.L., Fedrigo, E., Jatteau, P., Bonpunt, E., Billard, R., 1991.

Siberian sturgeon spermatozoa: effects of dilution, pH,

osmotic pressure, sodium and potassium ions on motility, in:

Williot, P. (Ed.), *Acipenser*: Actes Du Premier Colloque

International Sur l'esturgeon. Centre National du Machisme

Agricole du Genie Rural des Eaux et des Forets, Bordeaux,

pp. 143–151.

Giberson, A.V., Litvak, M.K., 2003. Effect of Feeding Frequency

on Growth, Food Conversion Efficiency, and Meal Size of

Juvenile Atlantic Sturgeon and Shortnose Sturgeon. N. Am. J.

Aquacult. 65, 99–105. https://doi.org/10.1577/1548-

8454(2003)65<99:EOFFOG>2.0.CO;2

Gilroy, C.E., Litvak, M.K., 2019. Swimming kinematics and

temperature effects on spermatozoa from wild and captive

shortnose sturgeon (*Acipenser brevirostrum*). Anim. Reprod.

Sci. 204, 171–182.

https://doi.org/10.1016/j.anireprosci.2019.03.022

Ginzburg, A.S., 1972. Fertilization in fishes and the problem of

polyspermy. Israel Program for Scientific Translation,

Jerusalem.

IUCN, 2023. IUCN Red List of Threatened Species [WWW

Document]. http://www.iucnredlist.org/.

Kim, S.T., Moley, K.H., 2008. Paternal effect on embryo quality in

diabetic mice is related to poor sperm quality and associated

with decreased glucose transporter expression. Reproduction

136, 313–322. https://doi.org/10.1530/REP-08-0167

Kime, D.E., Rurangwa, E., Ollevier, F., 2001. Computer-assisted

sperm analysis CASA as a tool for monitoring sperm quality

in fish. Comp. Biochem. Physiol., Part C: Toxicol.

Pharmacol. 130, 425–433.

Kuhn, M., 2008. Building Predictive Models in R Using the caret

Package. J. Stat. Soft. 28.

https://doi.org/10.18637/jss.v028.i05

Kynard, B., Bolden, S., Kieffer, M., Collins, M., Brundage, H.,

Hilton, E.J., Litvak, M., Kinnison, M.T., King, T., Peterson,

D., 2016. Life history and status of Shortnose Sturgeon

(*Acipenser brevirostrum* LeSueur, 1818). J. Appl. Ichthyol.

32, 208–248. https://doi.org/10.1111/jai.13244

Lenth, R., 2023. _emmeans: Estimated Marginal Means, aka

Least-Squares Means_. R package version 1.8.5.

https://CRAN.R-project.org/package=emmeans.

Litvak, M., 2010. The sturgeons (family: Acipenseridae), in:

François, N.L., Jobling, M., Carter, C., Blier, P. (Eds.),

Finfish Aquaculture Diversification. CABI, UK, pp. 178–199.

https://doi.org/10.1079/9781845934941.0178

Park, C., Chapman, F., 2005. An Extender Solution for the Short-

Term Storage of Sturgeon Semen. N. Am. J. Aquacult. 67,

52–57. https://doi.org/10.1577/FA03-068.1

R Core Team, 2023. R: A language and environment for statistical

computing. R Foundation for Statistical Computing. Vienna,

Austria. https://www.R-project.org/.

Sancho, S., Casas, I., Ekwall, H., Saravia, F., Rodríguez-Martinez,

H., Rodríguez-Gil, J.E., Flores, E., Pinart, E., Briz, M.,

Pruneda, A., Bussalleu, E., Yeste, M., Fàbrega, A., Bonet, S.,

2008. Sperm cryopreservation in Iberian boars: Study of the

sperm quality and distribution of the two hexose transporters

GLUT-3 and GLUT-5. Theriogenology 70, 1399.

https://doi.org/10.1016/j.theriogenology.2008.06.067

Searle, S.R., Speed, F.M., Milliken, G.A., 1980. Population

Marginal Means in the Linear Model: An Alternative to Least

Squares Means. Am. Stat. 34, 216–221.

https://doi.org/10.2307/2684063

Shaliutina, A., Hulak, M., Gazo, I., Linhartova, P., Linhart, O.,

2013. Effect of short-term storage on quality parameters,

DNA integrity, and oxidative stress in Russian (*Acipenser*

gueldenstaedtii) and Siberian (*Acipenser baerii*) sturgeon

sperm. Anim. Reprod. Sci. 139, 127–135.

https://doi.org/10.1016/j.anireprosci.2013.03.006

Shazada, N.E., Zhang, S., Siddique, M.A.M., Cheng, Y., Rodina,

M., Alavi, S.M.H., Linhart, O., 2024. Optimizing extenders

for short-term storage of Sterlet (Acipenser ruthenus) sperm.

Aquaculture Reports 37, 102216.

https://doi.org/10.1016/j.aqrep.2024.102216

Xin, M., Cheng, Y., Rodina, M., Tučková, V., Shelton, W.L.,

Linhart, O., 2020. Improving motility and fertilization

capacity of low-quality sperm of sterlet *Acipenser ruthenus*

during storage. Theriogenology 156, 90–96.

https://doi.org/10.1016/j.theriogenology.2020.07.004

710
711

Figures and tables

Figure 1. Treatments for experiment 1 and experiment 2.

Experiment 1 had six total treatments, experiment 2 had twelve

total treatments.

a)

b)

Figure 2. Mean motility, VCL, LIN, and WOB of sperm over time

from experiment 1 (a), and experiment 2 (b). Day 0 represents day

of sperm collection. Extenders are denoted by colours, H for

HBSS, P for Park and Chapman, M for modified Tsvetkova, and N

for no extender. Oxygen treatments for experiment 2 are denoted

by shapes, O for oxygen treatment and N for no oxygen treatment.

Table 1. Results from experiment 1 and experiment 2 swimming

kinematics statistical analysis using LMER. All parameters

decreased over each day. Post-hoc tests are shown for extender

effects on VCL for 2022 as there was no interaction.

Experiment	Analysis	Factor	VCL	LIN	WOB	MOT
1	Linear mixed model	Extender	0.01261*	0.2509	0.2538	0.3775
		Day	4.216E-06*	0.0003964*	6.324E-05*	0.0001541*
		Extender:Day	0.3698	0.9629	0.8399	0.06711
	Extender effect post-hoc, VCL					
		H-N	0.04430*			
		H-P	0.6251			
		N-P	0.01120*			
Experiment	Analysis	Factor	VCL	LIN	WOB	MOT
2	Linear mixed model	Extender	0.04480*	0.1224	0.08730	0.6774
		Oxygen	0.9242	0.8952	0.9398	0.4411
		Day	1.390E-05*	3.970E-05*	0.0001025*	3.860E-07*
		Extender:Oxygen	0.9838	0.8039	0.7625	0.6568
		Extender:Day	0.0002848*	5.410E-07*	9.070E-09*	0.002208*
		Oxygen:Day	0.3649	0.5362	0.5306	0.8724
		Extender:Oxygen:Day	0.6192	0.4672	0.4573	0.8686
*P<0.05						
H= HBSS						
N= No extender						
P= Park and Chapman extender						

735   **Table 2.** Results from experiment 1 and experiment 2 last day of

736   motility ANOVA's.

Experiment	Analysis	Factor	Last Day Motile
1	ANOVA	Extender	0.9350
2	ANOVA	Extender	6.470E-07*
	Tukey	N-M	1.800E-06*
		P-M	2.270E-05*
		P-N	0.7168
*P<0.05			
H= HBSS			
N= No extender			
P= Park and Chapman extender			
M= modified Tsvetkova extender			

Table 3. Results of Tukey comparisons from the ANOVA for the

effect of extenders on each day for experiment 2.

Day	Comparison	VCL	LIN	WOB	MOT
0	N-M	0.9447	0.07206	0.02567*	0.1814
	P-M	0.1322	0.008045*	0.003116*	0.9869
	P-N	0.06773	0.6506	0.7152	0.1356
3	N-M	0.9385	0.1590692	0.1973831	0.8901148
	P-M	0.09357	0.0026512*	0.0035696*	0.3572449
	P-N	0.04414*	0.2272558	0.2214556	0.1667551
7	N-M	0.3405	0.9587	0.9137	0.9851
	P-M	0.6237	0.9447	0.5914	0.9527
	P-N	0.8780	0.8219	0.8326	0.9905
10	N-M	0.0002980*	0.00005600*	0.00002270*	0.6938
	P-M	0.4979	0.2292	0.1821	0.9878
	P-N	0.008559*	0.008847*	0.005989*	0.7825
*$P<0.05$					
N= No extender					
P= Park and Chapman extender					
M= modified Tsvetkova extender					

Table 4. Results of short-term storage of *Acipenser* sperm from

eight previous studies.

Species	Motility Decrease	Atmosphere	Extender	Days	Motility Definition	Author
A. persicus	100%	NO	N	9	Not mentioned	Aramli et al., 2013
A. baerii	100%	O (2 mg o2/L)	N	5	VCL>3 μm/s	Barulin and Shumski, 2022
A. baerii	100%	O (4 mg o2/L)	N	6	VCL>3 μm/s	Barulin and Shumski, 2022
A. baerii	100%	O (5.5 mg o2/L)	N	8	VCL>3 μm/s	Barulin and Shumski, 2022
A. baerii	100%	O (12 mg o2/L)	N	11	VCL>3 μm/s	Barulin and Shumski, 2022

A. oxyrinchus	60%	O	N	17	Cells uncontrollably spinning in place were classified as nonmotile	DiLauro et al., 1994
A. oxyrinchus	0%	O (96%)	P	7	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	37.90%	NI (99%)	P	7	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	0%	O (96%)	M	7	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	95.20%	NI (99%)	M	7	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	0%	O (96%)	N	7	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	93.30%	NI (99%)	N	7	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	40%*	O (100%)	PL	21	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	50%*	O (100%)	PB	21	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	70%*	O (100%)	N	21	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	40%*	O (21%)	PL	21	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	50%*	O (21%)	PB	21	Not mentioned	Dorsey et al., 2011
A. oxyrinchus	70%*	O (21%)	N	21	Not mentioned	Dorsey et al., 2011
A. ruthenus	30%*	NO	N	1.5	VCL>3 μm/s	Dzyuba et al., 2015
A. oxyrinchus desotoi	30.70%	NO	P	21	Not mentioned	Park and Chapman, 2005
A. brevirostrum	50%	NO	P	28	Not mentioned	Park and Chapman, 2005
A. oxyrinchus desotoi	100%	NO	N	21	Not mentioned	Park and Chapman, 2005
A. brevirostrum	100%	NO	N	21	Not mentioned	Park and Chapman, 2005
A. gueldenstaedtii	100%	NO	N	9	VCL>3 μm/s	Shaliutina et al., 2013
A. baerii	100%	NO	N	9	VCL>3 μm/s	Shaliutina et al., 2013
Acipenser ruthenus	88.52%	NO	N	4	VCL>10 μm/s	Shazada et al., 2019
Acipenser ruthenus	71.98%	NO	E17**	6	VCL>10 μm/s	Shazada et al., 2019
Acipenser ruthenus	52.4%	NO	E18**	6	VCL>10 μm/s	Shazada et al., 2019
Acipenser ruthenus	47.42%	NO	E19**	6	VCL>10 μm/s	Shazada et al., 2019
Acipenser ruthenus	49.01%	NO	E20**	6	VCL>10 μm/s	Shazada et al., 2019
	*estimated from graphs **see publication for extender recipe					
	N= No extender P= Park and Chapman extender M= modified Tsvetkova extender NO= No atmosphere treatment O= Oxygen treatment					

	PL= Park and Chapman diluted after collection in lab
	PB= Park and Chapman diluted immediately after collection

752