Cell size and selection for stress-induced cell fusion in unicellular eukaryotes

Xiaoyuan Liu[1*], Jonathan W. Pitchford[1,2], George W. A. Constable[1],

1 Department of Mathematics, University of York, York, North Yorkshire, United Kingdom **2** Department of Biology, University of York, York, North Yorkshire, United Kingdom

*xiaoyuan.liu@york.ac.uk (XL)

Abstract

In unicellular organisms, sexual reproduction typically begins with the fusion of two cells (plasmogamy) followed by the fusion of their two haploid nuclei (karyogamy) and finally meiosis. Most work on the evolution of sexual reproduction focuses on the benefits of the genetic recombination that takes place during meiosis. However, the selection pressures that may have driven the early evolution of binary cell fusion, which sets the stage for the evolution of karyogamy by bringing nuclei together in the same cell, have seen less attention. In this paper we develop a model for the coevolution of cell size and binary cell fusion rate. The model assumes that larger cells experience a survival advantage from their larger cytoplasmic volume. We find that under favourable environmental conditions, populations can evolve to produce larger cells that undergo obligate binary cell fission. However, under challenging environmental conditions, populations can evolve to subsequently produce smaller cells under binary cell fission that nevertheless retain a survival advantage by fusing with other cells. The model thus parsimoniously recaptures the empirical observation that sexual reproduction is typically triggered by adverse environmental conditions in many unicellular eukaryotes and draws conceptual links to the literature on the evolution of multicellularity.

Author summary

Sexual reproduction is commonly observed, both in eukaryotic microorganisms and in higher multicellular organisms. Sex has evolved despite numerous apparent costs, including investment in finding a partner and the energetic requirements of sexual reproduction. Binary cell fusion is a process that sets the stage for sexual reproduction by bringing nuclei from different cells into contact. Here, we provide a mathematical explanation of the advantage conferred by binary cell fusion due to increased cell mass. We show that when unicellular organisms have the option to invest in either cell fusion or cell mass, they can evolve to fuse together as rapidly as possible in the face of adverse environments, instead of increasing their mass. These results are consistent with the empirical observation that sexual reproduction is often triggered by environmental stress in unicellular eukaryotes. Our results imply an advantage to cell fusion, which helps to shed light on the early evolution of sexual reproduction itself.

Introduction

Although the details of the early evolution of sexual reproduction in the last common eukaryotic common ancestor (LECA) are shrouded in mystery, it is argued that the emergence of eukaryotic sex began with the evolution of cell–cell fusion and meiosis [1] in an archaeal ancestor [2,3]. This step can be further broken down into the evolution of binary cell fusion, the one spindle apparatus, homologous pairing and chiasma, and finally reduction, division and syngamy [4]. The vast majority of theoretical studies investigating the evolution of sexual reproduction have focused on later stages of this evolutionary trajectory, namely the conditions that give rise to a selective pressure for genetic recombination [5–8]. However, comparatively few studies have investigated the selective pressures that may have first given rise to binary cell fusion, which may have facilitated the evolution of a host of other eukaryotic traits [9], including the homologous pairing and meiotic recombination, by bringing nuclei together in the same cell.

Hypotheses for the evolution of binary cell fusion often rely on hybrid fitness advantage. It has been suggested that selection for cell–cell fusions might have initially been driven by "selfish" transposons and plasmids [10–12], or negative epistatic

interactions between mitochondrial mutations [13, 14]. However, once a heterokaryotic cell has been formed (binucleate with nuclei from both parental cells), the advantage of hybrid vigor and the masking of deleterious mutations could lead to the maintenance of cell fusion [4]. Such benefits are required to alleviate costs to cell-fusion, which include selfish extra-genomic elements in the cytoplasm [15] and cytoplasmic conflict [16, 17].

In these previous studies on the evolution of binary cell fusion, the effect of changing environmental conditions is not considered. However, in many extant unicellular organisms, binary cell fusion (and the karyogamy and genetic recombination that follow) occur in response to challenging environmental conditions [18] such as starvation (*Tetrahymena* [19, 20]) and depleted nitrogen levels (*Chlamydomonas reinhardtii* [21] and *Saccharomyces pombe* [22]). Meanwhile in benign conditions with abundant resources these species reproduce asexually using binary cell fission. The mechanisms that drive selection for genetic recombination under challenging environmental conditions are well-studied [23]; recombination can facilitate adaptation to a novel environment [24, 25] and evolving to engage in more sex when fitness is low (fitness-associated sex) can allow an organism to maximise the advantages of sex while minimising the costs [26, 27]. However, this focus on the benefits of recombination leaves space to ask whether binary cell fusion itself could be selected for as a stress response, even in the absence of any genetic advantages.

In this paper, we do not account for the genetic factors discussed above. Instead, we focus only on how the survival advantage associated with increasing cytoplasmic volume might select for binary cell fusion; this relies on the physiological advantages conferred by cell-cell fusion and is independent of the question of the genetic advantages (and disadvantages) of sexual reproduction. This alternative perspective offers useful new insights that can be compared with empirical observation.

That size-based processes could play a role in the early evolution of sexual reproduction has empirical and theoretical support. The "food hypothesis" [28] suggests that metabolic uptake could drive horizontal gene transfer in bacteria and archaea, with DNA molecules providing nutrients for the receiving cell [29, 30]. Indeed, horizontal gene transfer has been shown experimentally to be an important source of carbon and nutrients in bacteria [31, 32]. Binary cell fusion is possible in bacteria (where it has been shown to come with selective benefits from mixed cytoplasm [33]) and archaea [34, 35].

Meanwhile amongst eukaryotes, the benefits of increasing cytoplasmic volume are understood to be strong enough to drive selection for the sexes themselves [36,37]. That early selection for syngamy may have been driven by the survival benefits of larger cytoplasmic volume was argued verbally by Thomas Cavalier-Smith. He suggested that syngamy's initially prime role was "to make zygotes larger and increase their survival rate by being able to store more solid food reserves" [38] and that this selective pressure would have been particularly important under stressful conditions "where life was threatened already by famine and the cell had less to lose and more to gain by fusing with others" [39]. However this hypothesis has until now not been explored mathematically. In suggesting a mechanistic hypothesis for the evolution of binary cell fusion, our work has interesting parallels with [40], where an advantage to cell fusion is identified in terms of shortening the cell-cycle.

Moving to consider potential physiological benefits of binary cell fusion naturally leads to work on the evolution of multicellularity. While multicellularity and binary cell fusion are clearly biologically distinct, from a modelling perspective they share similarities in that they can involve the "coming together" of cells to produce a larger complex [41]. Multicellularity achieved via aggregation allows organisms to rapidly adapt to novel environments that favour increased size [42,43]. It has also been suggested that the genetic nonuniformity of such aggregates may also make them well-suited to resource limited environments [44,45], echoing the hybrid vigor hypotheses for the evolution of early syngamy [4]. Relatively few theoretical studies have investigated the evolution of facultative aggregation in response to changing environments [43]. However in the context of clonal multicellularity ("staying together") such environments have been considered more extensively [46,47]. In this clonal context, the evolutionary dynamics act primarily on fragmentation modes [48,49] (e.g. how a "parental" multicellular complex divides to form new progeny). Interestingly the same quality-quantity trade-off arises here [46] as drives selection for the sexes [36] (anisogamy, gametes of differing sizes); larger daughter cells (or gametes) are more able to withstand unfavourable environmental conditions, while smaller cells can be produced in larger quantities.

In this paper we adapt the classic Parker-Baker-Smith [36] (PBS [37]) model for the evolution of sexes in order to investigate the evolution of binary cell fusion. This builds

on recent work that investigates how the possibility of parthenogenetic reproduction can

drive selection for oogamy in eukaryotes [50]. We assume for simplicity that parental

cells undergo a number of cell-divisions. The size of daughter cells is a compound

evolvable trait determined both by the size of parental cells and the number of cell

divisions. Daughter cells are then introduced to a pool in which they can undergo

binary cell fusion, with larger fused cells experiencing a survival advantage. Unlike in

the classic PBS model, the fusion rate is also a trait subject to evolution. In the

following sections we proceed to outline some insights developed from numerical

simulations before going on to develop analytical results for the model in a fixed

environment. Finally we introduce switching environments and show that under plastic

phenotypic responses, facultative binary cell fusion in response to harsh environmental

conditions can evolve.

Model

Many theories posit that the evolution of sex took place in the context of a

haploid-dominant life cycle, whereby haploid unicellular organisms reproduce asexually

via mitosis and the formation of a diploid cell is a transient state that follows

syngamy [38, 39, 51]. This is consistent with many extant unicellular eukaryotes, with

haploid dominant life cycles prevalent in fungi [52], the norm in chlorophyte and

charophyte algae [53] and accounting for approximately 30% of protist life-cycles [54].

For clarity we will therefore couch our model in terms of a population of free-living

haploid cells and draw examples from these extant organisms to motivate our model.

However we also stress that as our model is primarily concerned with cell mass it is

somewhat agnostic to the genetic details of ploidy. Thus the same model can be used in

the context of theories that suggest diploidy preceded the evolution of syngamy [55],

albeit with different interpretation of the content of the cell nucleus.

Among extant unicellular haploid species that feature facultative sexual

reproduction, the green alga *C. reinhardtii* is a particularly useful model organism to

consider, retaining as it does important features of the LECA [56]. We therefore briefly

describe its life cycle here. *C. reinhardtii* is an aquatic, single-celled haploid

organism [57]. Under benign conditions, cells reproduce asexually; cells grow to increase

their volume more than ten fold, before undergoing a series of n mitoses to produce 2^n [111] daughter cells, or zoospores. Following dissolution of the mother cell wall, these [112] daughter cells are released in order to complete the cycle. [113]

This vegetative mode of reproduction is discontinued under nitrogen-limited [114] conditions [58]. In this second scenario, *C. reinhardtii* cells instead undergo [115] differentiation to form sexually competent gamete cells (gametogenesis). Broadly, these [116] cells are morphologically similar to vegetative cells [59] but show enhanced motility, low [117] photosynthetic activity and mating structures that allow for cell-cell fusion [60, 61]. [118] Fusion between cells is not indiscriminate, but rather restricted to occurring between [119] cells of opposite mating types [62]. These two mating types (denoted plus and minus) [120] are genetically determined at the haploid level and can be understood as ancestral [121] versions of the sexes [63], but with no size dimorphism between the gametes (*C.* [122] *reinhardtii* is isogamous). Should nitrogen levels rise, these gametes can de-differentiate [123] into vegetatively reproducing cells [64]. However, should nitrogen levels remain low, [124] gametes of opposite mating type are chemotactically attracted to each other and fuse to [125] form a binucleate cell [64]. The cell nuclei then fuse to form a diploid zygote, or [126] zygospore, that develops a thick cell wall that is resistent to environmental stress before [127] entering a stage of dormancy. Restoration of benign environmental conditions triggers [128] meiosis in the zygospore and the production of four haploid daughter cells capable of [129] vegetative growth (germination). [130]

As addressed in the introduction, our mathematical model takes inspiration from the [131] classic PBS model for the evolution of anisogamy [36] (the production of sex cells of [132] different sizes). However, whereas such models typically consider the binary cell fusion [133] (fertilization) rate a fixed parameter, we here treat it as a trait subject to evolution. In [134] doing so, our work builds on [50], where the same core model (albeit with a different [135] biological framework) was used to investigate the evolution of anisogamy in species [136] capable of parthenogenesis (sex cell development in the absence of fertilization). Here [137] we begin by reframing the model in the biological context of the early evolution of [138] sexual reproduction, before introducing the possibility of plastic phenotypic responses [139] (which were not considered in [50]) into the model. [140]

Model of cell division, growth, and survival

We begin by considering a population of individual mature cells of total mass E. Each mature cell has a mass M, such that the total number of mature cells is E/M. The cells may then undergo $n \geq 0$ rounds of binary fission. Each resulting daughter cell then has mass $m = M/2^n$, while the total number of daughter cells in the population is $(2^n E)/M$. Following this fission phase, each daughter cell is subjected to an extrinsic mass-dependent mortality, such that larger daughter cells are more likely to survive into the next growth cycle (see upper half of Fig 1). While many choices for such a mass-dependent function are possible, in this paper we focus primarily on the Vance survival function [65], defined as

$$ S(m; \beta) = \exp\left(-\frac{\beta}{m}\right) . \tag{1} $$

Here the parameter β describes the magnitude of the mortality process (i.e. the harshness of the environment); for a given value of m, an increase in β decreases the survival probability of daughter cells. We therefore refer to β as the environmental harshness parameter, with high β corresponding to harsh environments in which survival is difficult, and low β corresponding to more benign environments in which even cells of modest mass have a high probability of surviving. We note that Eq. (1) is both mathematically simple (being defined by only one parameter, β) and has the biologically reasonably property of capturing the principle of a minimum cell size (see also Figure 7) and is thus a common assumption in the literature [66–68]. Surviving cells then go on to grow to mass M and seed the next generation, which will again consist of E/M cells. In this way, E can be understood as setting the carrying capacity of the population in terms of its total mass.

We now wish to incorporate evolutionary dynamics to the simple picture described above. We note that the mass of daughter cells, $m = M/2^n$, is dependent both on the mature cell size M and the number of cell divisions, n. For simplicity we therefore treat m as a genetically determined continuous trait, and explore its evolution using simulations. We note that this treatment is biologically equivalent to the evolving the cell sizer mechanism in *Chlamydomonas*, which operates to ensure that daughter cells have a uniform size [53].

Fig 1. Schematic for model life cycle. Cells (green circles) are characterized by different genotypes (orange and blue nuclei) that control the non-recombining traits m_i and α_i (respectively the daughter cell mass and fusion rate of the i^{th} genotype). Daughter cell mass is a compound parameter that depends both on the mature cell mass M_i and the number of cell divisions n_i ($m_i = M_i/2^{n_i}$) and is treated as a single continuous parameter. Following release of the maternal cell wall, daughter cells are exposed to each other. Should genotypes in the population evolve non-zero fusion rates ($\alpha_i > 0$), binary cell fusion between cells is possible (see Eq. (3)). With a finite time-window for cell fusion, T, only a fraction of cells will be fused at the end of this fusion period. Following this both fused and unfused cells are exposed to a mass-dependent survival function (see Eq. (1)) under which larger fused cells experience a survival benefit as a result of their increased cytoplasmic volume. Surviving fused cells can grow larger due to their increased size at the beginning of the growth period. The mathematical model assumes that the size of daughter cell size of each genotype is conserved (see Eq. (4)). This comes with the implicit assumption the fused cells undergo an additional round of cell division that allows for segregation of nuclei in daughter cells. Costs to cell fusion can arise due to failed cell fusion, hindered growth of the fused cell and failed segregation, as described in the main text. These periods during which these costs can manifest are highlighted in red and captured mathematically by the parameter $1 \geq C \geq 0$.

Suppose that we have S genotypes in the population, such that each genotype produces daughter cells of mass m_i for $i \in S$. We further suppose that initially the frequency of the i^{th} genotype is f_i, such that the number of mature cells of the i^{th} type at the beginning of a growth cycle is $f_i E/M_i$. These mature cells then produce $f_i E/m_i$ daughter cells, of which a fraction $S(m_i; \beta)$ survive to the next growth cycle. Renormalizing by the total number of surviving cells, the frequency of type i cells that

survive, f_i' is

$$f_i' = \frac{f_i \frac{E}{m_i} \exp\left(-\frac{\beta}{m_i}\right)}{\sum_{j=1}^{S} f_j \frac{E}{m_j} \exp\left(-\frac{\beta}{m_j}\right)} . \tag{2}$$

Since each of these surviving cells grows to a size M_i, the next growth cycle begins with
$f_i' E/M_i$ mature cells.

Note that to increase the number of daughter cells they produce (E/m_i) mature
cells can grow to smaller sizes (reduced M_i, which increases the number of mature cells
in the population) or increase their number of cell-divisions (increased n_i) . However by
decreasing M_i and increasing n_i, individuals also produce smaller daughter cells that
are more vulnerable to extrinsic mortality. The size of daughter cells is thus subject to a
quality-quantity trade-off and over multiple iterations of the deterministic cycles
described above, certain genotypes may come to dominate the population. Mutants are
introduced stochastically at the start of a growth cycle with average rate μ (the number
of growth cycles between successive mutations is taken from a geometric distribution
parameterized by rate μ). Mutant genotypes are also chosen stochastically; ancestral
genotypes are chosen with probability f_i, and mutants either increase or decrease the
daughter cell mass trait of their ancestor (i.e. $m_{S+1} = m_i \pm \delta m$), each with an unbiased
50% probability. The parameter δm thus captures the size of mutational steps in trait
space.

Fig 2 summarizes the outcome of such evolutionary dynamics. Fig 2A shows that
the population evolves towards an evolutionarily stable strategy (ESS) in m for a given
environment. Should the environment suddenly become harsher (via an increase in β)
the population evolves towards a new ESS, in which daughter cells are larger (i.e.
daughter cells evolve to become larger to withstand more adverse conditions).

Model additionally incorporating binary cell fusion

We now modify the model to allow for the possibility of binary cell fusion following the
cell fission described above (see Fig 1). Daughter cells may now fuse to form a
binucleated cell (e.g. a dikaryon [69], in which the cytoplasm of the contributing cells
are mixed but their nuclei or nucleoids remain distinct [38]) or remain a mononucleated
cell (with a single nucleus or nucleoid). The rate of cell fusion is given by α. We assume

that this second trait is genetically linked to the daughter cell mass trait, such that each of the S genotypes is now defined by a trait pair (m_i, α_i). Cell fusions occur as a result of mass-action kinetics. For simplicity, we assume that fusions between distinct genotypes occur at their average fusion rate, $(\alpha_i + \alpha_j)/2$. Denoting by N_i the number of unfused cells of genotype i and F_{ij} the number of fused cells resulting from daughter cells of genotype i and j we have

$$
\begin{aligned}
\frac{\mathrm{d}N_i}{\mathrm{d}t} &= -\sum_{j=1}^{S} \left(\frac{\alpha_i + \alpha_j}{2} \right) N_i N_j \,, \quad N_i(0) = f_i \frac{2^{n_i} E}{M_i} = f_i \frac{E}{m_i} \\
\frac{\mathrm{d}F_{ij}}{\mathrm{d}t} &= \left(\frac{\alpha_i + \alpha_j}{2} \right) N_i N_j \,, \qquad F_{ij}(0) = 0 \,,
\end{aligned}
\tag{3}
$$

with $\alpha_i \geq 0 \, \forall \, i$. These fusion dynamics are allowed to run for a fixed period T. When $\alpha = 0$ all cells remain mononucleated. Rates $\alpha > 0$ can be biologically understood as the evolution of any traits the promote cell fusion, such as the loss of the cell wall in the unicellular red algae *Galdieria* [70] or the formation of a mating structure in the unicellular green algae *Chlamydomonas* [71]. For $\alpha > 0$, some proportion of daughter cells will have fused in the period T, leaving some cells unfused; the total number of unfused cells of type i is then $N_i(T)$ and the total number of fused cells with i and j nuclei are $F_{ij}(T)$.

We assume that cells that remain unfused at the end of the fusion period T can still survive and subsequently grow vegetatively. Essentially we are assuming that at early stages of this evolutionary trajectory daughter cells are not overly specialized for strict cell fusion (e.g. full gametic differentiation). This is akin to asexual reproduction in cell wall-less *Galdieria* [70] and gametic de-differentiation in *Chlamydomonas* following exposure nitrogen compounds that can be used for growth [58, 72]. For simplicity, we further assume that any potential dedifferentiation is costless. Unfused cells again survive according to Eq. (1) at a mass-dependent rate. The final number of unfused cells that survive of type i is therefore given by $S(m_i; \beta)N_i(T)$.

Fused cells will receive a survival advantage from their increased mass, $m_i + m_j$. As these fused cells enter the growth period with a larger inital immature cell size, we also assume that fused cells can grow to a larger mature size, $M_i + M_j$. In order for this mature cell to produce daughter cells of the same size as unfused cells (e.g. controlled

Fig 2. Selection for cell fusion as an alternative to increased cell size in response to a harsh environment. Stochastic simulations of evolutionary trajectories when the system is subject to a switch from the benign environment ($\beta_1 = 0.5$, green region) to the harsh environment ($\beta_2 = 2.2$, orange region) at growth cycle 500. Panel A illustrates the case where the fusion rate is held at $\alpha = 0$, representing the scenario where the physiological machinery for fusion has not evolved. Panel B illustrates the case where fusion rate is also subject to evolution. Remaining model and simulation parameters are given in Supporting Information S4 and the initial condition is $(m(0), \alpha(0)) = (1.16, 0)$.

by a sizer mechanism as in *Chlamydomonas* [53]) we implicitly assume an additional round of cell division that allows for the production of mononucleated progeny through vegetative segregation [73] (or alternatively through plasmid segregation machinery [74]), as illustrated in Fig 1.

In practice, the evolution of the machinery required for cell-fusion (and subsequent segregation) is likely to come with costs. These costs include: (i) cell-fusion failure [75]

in the initial fusion phase; (ii) selfish extra-genomic elements in the cytoplasm [15], cytoplasmic conflict [16,17] and maintenance of a binucleated cell [76] hindering growth in the growth phase; (iii) the possibility of binucleated cells failing to form mononucleated progeny (i.e. failed segregation) in the vegetative segregation phase. These phases with potential costs are highlighted in red in Fig 1. We account for these costs with a single parameter $1 \geq C \geq 0$. For (i) costs arising solely from cell fusion failure, $(1 - C)F_{ij}(T)$ fused cells survive to be exposed to the mass-dependent survival phase. For (ii) costs arising solely from inhibited growth, mature cells would reach a size $(1 - C)(M_i + M_j))$. For (iii) costs arising solely from failed segregation, only $(1 - C)F_{ij}(T)S(m_i + m_j)$ surviving mature cells would successfully segregate, with the remainder dying.

Analogously to Eq. (4) we can now write down the change in frequency of type i the frequency of type i cells that survive, f'_i is

$$f'_i = \frac{N_i(T)\exp\left(-\frac{\beta}{m_i}\right) + \sum_{j=1}^{S}(1 - C)F_{ij}(T)\exp\left(-\frac{\beta}{m_i+m_j}\right)}{\sum_{k=1}^{S}\left[N_k(T)\exp\left(-\frac{\beta}{m_k}\right) + \sum_{j=1}^{S}(1 - C)F_{kj}(T)\exp\left(-\frac{\beta}{m_k+m_j}\right)\right]}. \tag{4}$$

We now explore the coevolution of daughter cell mass, m, and fusion rate, α. We assume that mutations in α occur independently of m but at the same fixed rate μ, where μ is measured in units of (number of growth cycles)$^{-1}$ (see also Supporting Information S2.5 and [50]). Analogously to mutations in daughter cell mass, ancestral genotypes are chosen proportional to their frequency, f_i, with mutant genotypes inheriting their parental genotype plus or minus some trait deviation, $\delta\alpha$; $\alpha_{S+1} = \alpha_i \pm \delta\alpha$.

In Fig 2B, we see that in the benign environment, α remains at zero, and the population evolves towards an ESS in m as in Fig 2A. However now when the population is introduced to a harsher environment, the evolutionary dynamics differ from those in Fig 2A (where α was held artificially at zero). Rather than cells evolving to be larger, we see a different response emerging; selection for binary cell fusion ($\alpha > 0$).

The result above is in some sense surprising. Despite the presence of additional survival costs associated with binary cell-fusion, selection for non-zero fusion rates (rather than increased daughter cell size) persists in the harsh environment. We explain

the emergence of this behaviour mathematically in the Mathematical Analysis and
Results section.

Switching environments with phenotypic plasticity

We have seen in Fig 2B that selection for cell fusion can evolve in response to harsh
environmental conditions. We are particularly interested in exploring when such cell
fusion can evolve as a plastic stress response to changing environmental conditions, as
observed in the sexual reproduction of unicellular organisms such as
Tetrahymena [19,20], *C. reinhardtii* [21] and *S. pombe* [22]. We recall that while the
core model in a fixed environment (described above) has been analysed in [50], the
possibility of phenotypic plasticity was not considered.

We model environmental change as switching between two environments β_1 and β_2.
If $\beta_2 > \beta_1$, then β_2 is the harsher environment in which cells have a lower survival
probability (see Eq. (1)). For clarity, we will keep this convention for the remainder of
the paper. We will also allow for phenotypic plasticity such that the population can
evolve different strategies in different environments. The population's evolutionary state
is now described by four traits; the daughter cell mass in environments 1 and 2 (m_1 and
m_2) and fusion rate in these environments (α_1 and α_2). We assume that these traits are
all linked, so that recombination between the traits is not possible. For simplicity we
initially assume that any cost of phenotypic switching or environmental sensing is
negligible and that this plastic switching is instantaneous upon detection of the change
in environmental conditions.

Implementation of Numerical Simulations

The stochastic simulations of the evolutionary trajectories are also implemented using a
Gillespie algorithm [77] where successive mutations and environmental switching events
occur randomly with geometrically distributed waiting times. The rates of mutations μ
and environmental switching λ are measured in units of (number of growth cycles)$^{-1}$.
In the simulations, multiple traits coexist under a mutation-selection balance (see
Supporting Information and [50] and [78] for more detail), which allows us to account
for variations in selection strengths in simulations of our evolutionary trajectories.

As in [50], environmental switching is modeled as a discrete stochastic telegraph
process, with the time spent in each environment distributed geometrically, the discrete
analogue of an exponential distribution since the time spent in each environment is
measured in the number of discrete generations. The population spends an average of
$\tau_1 = 1/\lambda_{1\to2}$ in environment 1 and $\tau_1 = 1/\lambda_{2\to1}$ in environment 2, where $\lambda_{i\to j}$ is the
transition rate from environment i to j.

Mathematical Analysis and Results

In order to analyse the dynamics of the model, we use tools from adaptive
dynamics [79], assuming that traits are continuous and that mutations have small effect
to develop equations for the evolutionary dynamics of traits m and α. We have
previously derived these equations in [50] for the case without phenotypic plasticity.
In [50] we were primarily concerned with the evolution of suppressed fertilization
(gamete fusion) in sexual eukaryotes that had evolved the capacity for asexual
reproduction. Mathematically, this equates to considering initial evolutionary conditions
$\alpha(0) > 0$. In contrast, in the present paper we focus on the initial evolution of cell
fusion, which is equivalent to considering possible evolutionary outcomes when $\alpha(0) = 0$.

For orientation, we begin by describing the possible evolutionary trajectories from an
initial state $(m(0), \alpha(0)) = (m_0, 0)$ in a fixed environment (i.e. when the parameter β,
which measures the harshness of the environment (see Eq. (1)), is constant). We then
move on to exploring the consequences of these dynamics for a population in switching
environments with a plastic phenotypic response.

Fixed Environment

Adopting the classical assumptions of adaptive dynamics [79, 80], we obtain equations
for the selective gradient on traits m and α, which we write as $H_m(m, \alpha; \beta, C)$ and
$H_\alpha(m, \alpha; \beta, C)$ respectively (see Supporting Information S1 and [50]). We find that
along the zero-fusion boundary, the selection gradient $H_\alpha(m, 0; \beta, C)$ can be negative.
While this makes sense mathematically (it is equivalent to the additional *production* of
cells during the fusion window, see Eq. (3)) it makes no sense biologically (the
population's cell mass is not conserved over the cell division and fusion period). We

therefore introduce a biologically realistic boundary at $\alpha = 0$. The equations for the

evolutionary dynamics of m and α are then given by

$$
\frac{\mathrm{d}m}{\mathrm{d}\tau} = H_m(m,\alpha;\beta,C)
$$

$$
\frac{\mathrm{d}\alpha}{\mathrm{d}\tau} =
\begin{cases}
H_\alpha(m,\alpha;\beta,C) & \text{if } \alpha > 0 \text{ or } H_\alpha(m,0;\beta,C) \geq 0 \\
0 & \text{if } \alpha = 0 \text{ and } H_\alpha(m,0;\beta,C) < 0 \,,
\end{cases}
\tag{5}
$$

with

$$
\begin{aligned}
H_m(m,\alpha;\beta,C) &= -\frac{4m(m-\beta) + E\alpha T(1-C)e^{\frac{\beta}{2m}}(4m-\beta)}{4m^2(m + E\alpha T(1-C)e^{\frac{\beta}{2m}})} \\
H_\alpha(m,\alpha;\beta,C) &= -m\frac{(1 - (1-C)e^{\frac{\beta}{2m}})\ln(1 + \frac{E\alpha T}{m})}{2\alpha(E\alpha T(1-C)e^{\frac{\beta}{2m}} + m)} \,.
\end{aligned}
\tag{6}
$$

In total we find three possible attractors for the evolutionary dynamics (see Supporting

Information Section S2), which we outline below.

We begin by considering the dynamics along the $\alpha = 0$ boundary. For $\alpha = 0$, the

equation for the evolutionary dynamics of m simplifies to

$$
\frac{\mathrm{d}m}{\mathrm{d}\tau} = \frac{\beta - m}{m^2}
\tag{7}
$$

while the inequality $H_\alpha(m,0;\beta,C) < 0$ simplifies to

$$
m > -\frac{\beta}{2\ln(1-C)} \,.
\tag{8}
$$

These equations, together with the condition $\alpha \geq 0$, suggest that a fixed point exists on

the $\alpha = 0$ boundary at

$$
(m^*, \alpha^*) = (\beta, 0) \,.
\tag{9}
$$

Meanwhile for large α we find an additional diverging trajectory. In the limit $\alpha \to \infty$,

we find

$$
\left.\frac{\mathrm{d}m}{\mathrm{d}\tau}\right|_{\alpha \to 0} = -\frac{(4m - \beta)}{4m^2} \,, \qquad \left.\frac{\mathrm{d}\alpha}{\mathrm{d}\tau}\right|_{\alpha \to 0} = 0 \,.
\tag{10}
$$

These equations suggest that an attracting manifold $m = \beta/4$ exists in the limit $\alpha \to \infty$

$$
m^*_{\alpha \to \infty} = \frac{\beta}{4} \,.
\tag{11}
$$

The third and final potential evolutionary attractor is found by solving
$H_m(m, \alpha; \beta, C) = 0$ and $H_\alpha(m, \alpha; \beta, C) = 0$ for arbitrary m and α. We obtain

$$(m^*, \alpha^*) = \left(-\frac{\beta}{2\ln(1-C)}, -\frac{\beta}{ET} \frac{1 + 2\ln(1-C)}{2\ln(1-C) + \ln(1-C)^2} \right) . \qquad (12)$$

Unlike the previous evolutionary attractors (see Eqs. (9-11)), this fixed point is a
function of the cost to cell-fusion C. It is therefore useful to focus on the behaviour of
this fixed point as C is increased in order to characterize the behaviour of the
population in different parameter regimes, as illustrated in Fig. 3.

When $C = 0$, the fixed point given in Eq. (12) yields a negative, and therefore
biologically nonphysical, value of α^*. The remaining biologically relevant evolutionary
attractors are an unstable saddle at the fixed point along the $\alpha = 0$ boundary (see
Eq. (9)) and a diverging trajectory towards $\alpha \to \infty$ along the $m^* = \beta/4$ manifold (see
Eq. (11)), as illustrated in Fig. 3(A). As C is increased, α^* at the fixed point given in
Eq. (12) also increases, eventually crossing the $\alpha = 0$ boundary when $C = 1 - e^{-1/2}$ (see
expressions for m^* in Eqs. (9) and (12)). At this point the fixed point given by Eq. (12)
crosses the fixed point along the $\alpha = 0$ boundary and a bifurcation occurs, yielding a
saddle fixed point on the interior and rendering the fixed point at the boundary stable
(see Fig. 3(B)). At this stage the basin of attraction for the stable fixed point on the
$\alpha = 0$ boundary only contains larger values of m for initial conditions with $\alpha(0) = 0$.

As the cost to cell fusion, C, becomes larger still, the basin of attraction for the
stable fixed point on the $\alpha = 0$ boundary grows to encompass the entire $\alpha = 0$
boundary. We denote the critical value of C at which this occurs as C_{Base}. At this
stage, should m_0 be sufficiently small, an initial increase in fusion can be observed but
the population is ultimately attracted to a zero-fusion state at long times, as illustrated
in Fig. 3(C). Finally as C is increased to very large values, the interior fixed point
crosses the diverging trajectory with $\alpha \to \infty$ when $C = 1 - e^{-2}$ (see expressions for m^*
in Eqs.(11) and (12)). This results in another transcritical bifurcation, following which
selection for α along the $m = \beta/4$ manifold becomes negative, and the fixed point given
in Eq. (12) becomes biologically nonphysical (see Fig. 3(D)). This bifurcation then
leaves the zero fusion state as the only stable fixed point.

The scenarios described above for $C > C_{\text{Base}}$ are qualitatively similar, in that they

Fig 3. Fusion costs govern the basin of attraction for the zero-fusion state in a fixed environment. Each panel gives the phase portrait for the co-evolutionary dynamics of (m, α) in a fixed environment (see Eq. (5)). (A) High fusion rates are the only evolutionary outcome when costs to cell fusion are low, with an unstable fixed point at $(m^*, \alpha^*) = (\beta, 0)$ (open red circle) and selection for increasing α on the invariant manifold $m = 1/\beta$ (blue arrow). Purple arrow illustrates the point on the $m = 0$ boundary when selection for α becomes positive (see Eq. (8)). (B) Under low-intermediate fusion costs, a third saddle fixed point emerges (see Eq. (12)), undergoing a bifurcation with $(m^*, \alpha^*) = (\beta, 0)$ and rendering it stable (solid red disk). There are now two evolutionary outcomes, depending on initial conditions; trajectories either increase in α along the invariant manifold, $(m^*, \alpha^*) \to (\beta/4, \infty)$, or are attracted to the $(m^*, \alpha^*) = (\beta, 0)$ fixed point. (C) For intermediate to high fusion costs, the separatrix of the saddle fixed point (red arrows) crosses the origin, rendering the entire $m = 0$ boundary within the basin of attraction of the $(m^*, \alpha^*) = (\beta, 0)$ fixed point. While for low initial cell masses the fusion rate may initially increase, ultimately the population will evolve to a low fusion state. (D) Under high fusion costs the saddle fixed point crosses the invariant manifold $m = 1/\beta$, reversing selection for the fusion rate such that $d\alpha/d\tau$ is now negative along this line. In this scenario all initial conditions evolve towards the $(m^*, \alpha^*) = (\beta, 0)$ fixed point. Coloured dashed lines show the average over multiple stochastic realizations of the simulated model (see also Fig. S1) with initial conditions $m(0) = 0.02$ (blue), $m(0) = 0.5$ (orange), $m(0) = 1.75$ (green) and all in an initial state without fusion ($\alpha(0) = 0$). Each simulated trajectory is run for 1.5×10^6 growth cycles and remaining parameters are given in Supporting Information S4.

Fig 4. Critical cost above which cell fusion becomes impossible to evolve increases with cell density and time available for cell fusion. Blue line shows numerical solutions of Eq. (13). For values of $C > C_{\text{Base}}$, obligate cell fusion remains a potential evolutionary outcome (see Eq. (14)). The value of $C > C_{\text{Base}}$ is bounded above and below by the bifurcation points identified in the main text (black dashed lines, see Eq. (15)).

feature a single evolutionary endpoint for all trajectories with initial conditions $(m(0), \alpha(0)) = (m_0, 0)$ (see in Fig. 3(C-D)). In order to determine C_{Base} analytically, we need to obtain the value of C for which the separatrix, marking the boundary of the basin of attraction for the fixed point $(m^*, \alpha^*) = (\beta, 0)$, intersects the origin point $(m, \alpha) = (0, 0)$. We can obtain a linear approximation for this separatrix using the stable directions of the fixed interior saddle fixed point given in Eq. (12) (see red arrows in Fig. 3(B-C)). In the Supporting Information Section S2.4 we show this is given approximately by the solution to

$$
\begin{aligned}
8(1 + 2\ln(1 - C_{\text{Base}}))^2 \left[5 + \ln(1 - C_{\text{Base}}) \left(5 + 2\ln(1 - C_{\text{Base}})\right)\right] \\
- (ET)^2 \left(2 + \ln(1 - C_{\text{Base}})\right)^4 \ln\left(\frac{6}{2 + \ln(1 - C_{\text{Base}})} - 3\right) = 0 \,.
\end{aligned}
\tag{13}
$$

The root of this equation can be understood as being governed by the compound parameter ET (see Fig. 4). When ET is small (i.e. the density of cells entering the fusion pool is low or the period allowed for fusion short), the root approaches the solution to $1 + 2\ln(1 - C_{\text{Base}}) = 0$. Meanwhile when ET is large the root approaches the solution to $2 + \ln(1 - C_{\text{Base}}) = 0$. These limits to the solution for C_{Base} are precisely the bifurcation points identified previously.

Summarizing the results above, we therefore have three possible evolutionary scenarios starting from initial conditions $(m(0), \alpha(0)) = (m_0, 0)$. Which of these scenarios we observe depends on the cost to cell fusion, C. When C is low, the only evolutionary endpoint is the invariant manifold at high fusion rate. In this scenario,

obligate fusion is the only evolutionary outcome, irrespective of m_0. For intermediate costs to cell fusion, there are two possible evolutionary outcomes, depending on the initial daughter cell mass m_0: if m_0 is small, selection acts to increase fusion rate and obligate fusion is the ESS; if conversely if m_0 is large, the state of no cell fusion is the ESS. Finally when costs to fusion are high selection for decreased fusion rate acts regardless of the initial value of m_0. Mathematically

$$(m^*, \alpha^*) \to \begin{cases} (\beta/4, \infty) & \text{if } 1 - \frac{1}{\sqrt{e}} > C \geq 0 \\ (\beta/4, \infty) \text{ or } (\beta, 0) & \text{if } C_{\text{Base}} > C \geq 1 - \frac{1}{\sqrt{e}} \\ (\beta, 0) & \text{if } C > C_{\text{Base}} \end{cases} \qquad (14)$$

where

$$1 - \frac{1}{e^2} > C_{\text{Base}} \geq 1 - \frac{1}{\sqrt{e}} \, . \qquad (15)$$

We note that it has been previously shown in [50] that the first of these conditions has additional consequences in this model for the evolution of suppressed fertilization in eukaryotic macrogametes. Should the costs to fertilization (cell fusion) exceed $C = 1 - e^{-1/2}$, evolutionary branching can occur at large values of the fusion rate α. However this branching happens at a later evolutionary stage than the focus of this study, the early emergence of binary cell fusion.

We conclude this section by addressing the key biological result that arises from this analysis; cell fusion is uniformly selected for even under moderately high costs (with a fraction of up to $C \approx 0.39$ of fused cells failing to survive) and can even be selected for under higher costs (up to $C \approx 0.86$) given necessary initial conditions and parameters. In the context of the evolution of early binary cell fusion, this provides a surprising nascent advantage to cell fusion. This advantage could even help compensate for other short-term costs arising from the later evolution of sex and recombination. The selective advantage experienced by fusing cells comes from their increased cytoplasmic volume, which leads to increased survival probabilities.

Switching environments with phenotypic plasticity

We now consider the case where evolution acts on the same traits as before, but where the environment is subject to change and the population can evolve a specific response to each environment. We find that under the assumption of costless phenotypic plasticity, the evolutionary dynamics in each environment are decoupled (see Supporting Information Section S3.2);

$$\frac{\mathrm{d}m_1}{\mathrm{d}\tau} = P_1 H_m(m_1, \alpha_1; \beta_1, C), \qquad \frac{\mathrm{d}m_2}{\mathrm{d}\tau} = P_2 H_m(m_2, \alpha_2; \beta_2, C)$$
$$\frac{\mathrm{d}\alpha_1}{\mathrm{d}\tau} = P_1 H_\alpha(m_1, \alpha_1; \beta_1, C), \qquad \frac{\mathrm{d}\alpha_2}{\mathrm{d}\tau} = P_2 H_\alpha(m_2, \alpha_2; \beta_2, C) \tag{16}$$

where $H_m(m, \alpha; \beta, C)$ and $H_\alpha(m, \alpha; \beta, C)$ retain the functional form in Eq. (5) and we again restrict the dynamics to the biologically reasonable $\alpha \geq 0$. Note that the key effect of environmental switching here is to moderate the rate of evolution, with each trait pair (m_i, α_i) evolving at a rate proportional to the time the population is exposed to environment i.

While the *dynamics* of the population in each environment is decoupled, the evolutionary *trajectories* in each environment are coupled by the initial trait values for the population in each environment, which we assume are the same;

$$m_1(0) = m_2(0) = m_0, \quad \text{and} \quad \alpha_1(0) = \alpha_2(0) = \alpha_0 = 0, \tag{17}$$

i.e. the population begins in a phenotypically undifferentiated state. As addressed, we are particularly interested in the case where cell fusion emerges as a plastic response, being selected for in one environment and against in the other. Considering the scenarios summarized in Eq. (14), this restricts our attention to the case of intermediate costs to cell fusion, $C_{\mathrm{Base}} > C \geq 1 - e^{-1/2}$, for which both the zero fusion and the obligate fusion states are potential evolutionary outcomes.

As Eqs. (16) are only coupled through their shared initial conditions, m_0 and α_0, the choice of these initial conditions is an important consideration. Since we are interested in the initial evolution of binary cell fusion, it is natural to assume that the population evolves from a state of zero fusion, $\alpha_0 = 0$. Deciding on a plausible initial daughter cell mass, m_0, takes more thought. We will consider two parsimonious choices.

Fig 5. Selection for cell fusion as an alternative to increased cell size in response to a harsh environment. Illustrative phase portrait for co-evolutionary dynamics of $(m_1, \alpha_1, m_2, \alpha_2)$ in a switching environment with phenotypic switching that exhibits facultative binary cell fusion. In both environment 1 (panel A) and environment 2 (panel B) the cost to cell fusion is $C = 0.6$, purple circles represent the initial condition $(m_1(0), \alpha_1(0)) = (m_2(0), \alpha_2(0)) = (\beta_1, 0)$, and orange circles represent the initial condition $(m_1(0), \alpha_1(0)) = (m_2(0), \alpha_2(0)) = (m^*_{\text{BH}}, 0)$ (see Eq. (20)). The red shaded regions shows the approximate region of the basin of attraction for the stable zero fusion fixed points $(m^*, \alpha^*) = (\beta_1, 0)$ (see red disk, panel (A)) and $(m^*, \alpha^*) = (\beta_2, 0)$ (see red disk, panel (B)). Red arrows indicate the intersection of these basins with the $\alpha = 0$ boundary. Environmental parameters are $\beta_1 = 0.5$ and $\beta_2 = 2.2$ making environment 1 the more benign environment, in which the population typically spends a proportion $P_1 = 0.7$ of its time.

One parsimonious choice would be that the population is already adapted to either environment 1 or environment 2 and that m_0 is given by an evolutionary fixed point in one of these environments (this is the situation illustrated in Fig. 2). Suppose that the population was first exposed to environment 1 (with $\beta = \beta_1$), and that in this environment there was no selective pressure to evolve cell fusion (see Eq. (9)). The population would then initially reside at the zero-fusion fixed point of environment 1,

$$m_1(0) = m_2(0) = m_0 = \beta_1 \,, \tag{18}$$

as illustrated by the purple circles in Fig. 5). In order for selection for cell fusion to act in environment 2, $m_0 = \beta_1$ must lie outside the basin of attraction of the stable zero-fusion fixed point $m^* = \beta_2$ in environment 2. In general, the conditions required for this scenario to hold must be obtained numerically. However when ET is large (e.g.

in high-density environments), the condition simplifies analytically to

$$\beta_2 > 2\ln\left(\frac{1}{1-C}\right)\beta_1\,,\tag{19}$$

i.e. the second environment must be sufficiently harsher than the first environment in
order for obligate cell fusion to evolve as a stress response. This is illustrated in
Fig. 6(A).

An alternative choice for the initial cell mass assumes that the population has
evolved under exposure to both environment 1 and environment 2, but has not yet
evolved phenotypically plastic responses. We have shown in [50] (see also Supporting
Information S3.1) that for $\alpha = 0$, the ESS for daughter cell cell mass in such a scenario
is a bet-hedging strategy;

$$m_{\mathrm{BH}}^* = P_1\beta_1 + (1 - P_1)\beta_2\,.\tag{20}$$

Note that this is intuitively equivalent to the $\alpha = 0$ evolutionary strategy in
environment 1 (weighted by the probability of being in environment 1) plus the
evolutionary strategy in environment 2 (weighted by the probability of being in
environment 2), as illustrated by the orange circles in Fig. 5). We have shown in [50]
that Eq. (20) is an accurate approximation if the timescale of environmental switching
is faster than the evolutionary timescale $(\lambda_{i\to j} > \mu)$.

We must now determine whether the initial condition m_{BH}^* lies in the basins of
attraction of $(m^*, \alpha^*) = (\beta_1, 0)$ in environment 1 and $(m^*, \alpha^*) = (\beta_2, 0)$ in
environment 2. Again this in general requires a numerical approach. However we show
in Supporting Information Section S3.2.2 that analytic progress can be made in the
limit of large ET. We find that in the cost region in which a facultative response is
possible $(1 - e^{-2} > C \geq 1 - e^{-1/2})$, $(m_{\mathrm{BH}}^*, 0)$ is within the basin of attraction of
$(m^*, \alpha^*) = (\beta_1, 0)$ in environment 1 if

$$\beta_2 > \beta_1\left(\frac{P_1}{P_1 - 1} + \frac{1}{2(P_1 - 1)\log(1 - C)}\right)\,,\tag{21}$$

Fig 6. Conditions on cell fusion cost and deterioration in environment required to observe cell fusion as a facultative stress response. Regions are calculated in the limit of large ET and show when fusion is selected for in both environments (blue, $1 - e^{-1/2} > C$) and in neither environment (red, e.g. $C > 1 - e^{-2} > C$) and when fusion is selected for in only the harsher environment (purple and pink). On the interval $1 - e^{-2} > C > 1 - e^{-1/2}$ the evolutionary outcome depends on the ratio of environmental parameters and on the initial evolutionary conditions. In panel (A) the population is initially adapted to environment 1, and the condition for facultative cell fusion in environment 2 is given in Eq. (19). In panels (B-D) the population has initially adapted a bet-hedging strategy to the two environments. Cell fusion remains at zero in environment 1 but evolves in environment 2 if Eq. (21) hold but Eq. (22) is violated (see purple regions). Conversely cell fusion evolves in environment 1 but is held at zero in environment 2 if Eq. (21) is violated but Eq. (22) holds (see pink regions).

and $(m_{\mathrm{BH}}^*, 0)$ is within the basin of attraction of $(m^*, \alpha^*) = (\beta_2, 0)$ in environment 2 if ⁠462

$$\beta_2 < \beta_1 \left(\frac{2P_1 \log(1-C)}{2(P_1 - 1)\log(1-C) - 1} \right). \tag{22}$$

If only one of these conditions holds, then facultative cell fusion can be observed in one ⁠463

environment but not the other. Further, if facultative cell fusion does occur, it will ⁠464

always occur in response to the harsher of the two environments. ⁠465

In Fig. S7(B-D) we illustrate how the possible evolutionary outcomes vary as $P_1 = \tau_1/(\tau_1 + \tau_2)$ (the probability of being in environment 1) is reduced (note that the plot for $P_1 \to 1$ tends to Fig. S7(A), see Eq. (20)). We see that as the time spent in environment 1 is reduced, so too is the parameter regime in which one would expect to see facultative cell fusion in response to a harsher environment 2. However we also see the emergence of a region in which environment 2 is more benign ($\beta_2 < \beta_1$) and in which by symmetry facultative cell fusion can emerge in response to the harsher environment 1 (see pink regions in Fig. S7(B-D)). In Figs. S5-S7 of the Supporting Information, we show that these analytic results are good predictors of the dynamics obtained via numerical simulation.

In summary, for both scenarios described above, we see how facultative binary cell fusion can evolve as a response to harsh environmental conditions that lower the survival probability of daughter cells. Such a response is possible if costs to cell fusion are intermediate and if there is an appreciable increase in environmental harshness, β, between the environments. We have also seen that regular exposure to both environments before the evolution of cell fusion has taken place (such that the population has previously adapted a bet-hedging strategy in daughter cells mass) may reduce the parameter region over which we expect to see cell fusion as a stress response. However we note that in many sexually reproducing unicellular eukaryotes, the environmental conditions that trigger sexual reproduction (and cell fusion during plasmogamy) are very rare; sexual reproduction is estimated to be triggered approximately once in every 2000 generations in *Saccharomyces paradoxus* and once in every 770 generations in *C. reinhardtii* [81,82]. These values would be equivalent to $P_1 \approx 1$ in our model, which is the regime in which facultative binary cell fusion occupies the broadest region of parameter space.

Alternative Survival Functions: The importance of minimum cell size

Throughout the mathematical analysis above, we have considered one specific form of survival function, namely the Vance survival function Eq. (1). In order to illustrate the robustness of our results, we explore the effect of alternative survival functions in

Fig 7. Various choices for the survival function. The blue line illustrates the Vance survival function given in Eq. (1) for $\beta = 1$. We note that this function implicitly captures the feature of a minimum viable cell size as which the survival of cells drops precipitously (see vertical dashed blue line at $m = \beta/4$). The green dashed line shows the survival function given in Eq. (23) with $\gamma = 1$ and $m_{\min} = 0$. Here cell survival decays linearly with mass for small m. The orange dashed line shows Eq. (23) with $\gamma = 0.75$ and $m_{\min} = 0.5$.

Supporting Information Section S5. These functions are illustrated in Fig. 7. We show   496

that an important feature of the Vance survival function is that it implicitly accounts   497

for very small cells having a low survival probability (the growth of Eq. (1) is sub-linear   498

when $m < \beta/2$ as Eq. (1) is convex in this region). Conversely the function   499

$1 - \exp(-\gamma m)$ is linear in m when m is small. Consequently we find that the   500

evolutionary dynamics generated by the survival function $1 - \exp(-\gamma m)$ lead to   501

selection for the production of infinitely many (and infinitely small) daughter cells. In   502

order to correct for this behavior, we can modify the function to account for a minimum   503

viable cell size;   504

$$S(m;\gamma) = \begin{cases} 1 - e^{-\gamma m} & \text{if} \quad m \geq m_{\min} \\ 0 & \text{if} \quad m < m_{\min}. \end{cases} \tag{23}$$

The incorporation of this minimum cell size allows us to identify analogous parameter   505

regimes as identified when using the Vance survival function. In particular we once   506

again find that binary cell fusion can be selected for as a stress response should one   507

environment be sufficiently harsher. The exact parameters at which this occurs are   508

dependent on the choice of $m_{\min}$ and the fusion cost, C. However one notable point of   509

departure is the maximum value for C under which cell fusion can be selections for.   510

Whereas we have shown that the Vance survival function can allow selection for cell   511

fusion up to $C \approx 0.86$, using Eq. 23 leads to a maximum cell fusion cost of $C = 0.5$.   512

Discussion

The evolution of sexual reproduction and its consequences for the subsequent evolutionary trajectory of populations is of general importance in biology [7, 13, 83]. In this paper we have illustrated a reversal of the classic two fold cost of sex that appears in organisms with distinct sexes [84]; in unisexual [85], unicellular organisms, binary cell fusion can be selected for, even in the presence of substantial costs, due to a survival benefit that comes from increased mass. These results allow us to quantitatively assess the verbal hypothesis that syngamy evolved allow cells to store more food reserves and thus increase their survival rate [38]. It is particularly interesting that the benefits conferred to cell fusion through increased cytoplasmic mass are sufficient to withstand remarkably high costs; "obligate sexuality" is the *only* evolutionary outcome with costs equivalent to a loss of $\sim 39\%$ cells that attempt to fuse, and remains a potential outcome with costs of up to $\sim 86\%$ of fused cells dying.

Perhaps most interesting is the case of switching environments with phenotypic plasticity. Here we find under a broad set of biologically reasonable conditions (costs to cell fusion equivalent to $39\% - 86\%$ additional mortality to fused cells and at least moderate changes in environmental quality) that high fusion rates are selected for in harsh environments and zero fusion rates are maintained in benign environments. This behaviour parsimoniously recapitulates the empirically observed reproductive strategies of numerous facultatively sexual species, including *C. reinhardtii* [21] and *S. pombe* [22]. This mechanism, under which cell fusion evolves to increase the survival probability of daughter cells, provides a complementary perspective on the frequent evolution of survival structures (resistant to environmental stress) that form following the formation of a zygote. These include ascospores in fungi [86] and zygote-specific stress-resistant stress wall in *C. reinhardtii* [87]. Note that such correlations between sexual reproduction and the formation of survival structures are not as easily explained under genetic explanations for the evolution of sexual reproduction, where engaging in both behaviour at once constitutes a simultaneous (and therefore potentially costly) change in genotype and temporal dislocation in environment [88, 89].

The results above are particularly interesting in the case of the evolution of early binary cell-fusion as a first step in the evolution of sexual reproduction. While most

studies focus on the genetic benefits of cell-fusion [90] (including a functionally-diploid dikaryotic cell [4]), or the genetic benefits of mixed cytoplasm [13, 14] (which can also come with costs [15–17, 41, 91–93]), the mechanism at play here is purely physiological. Yet, as addressed above, it naturally captures the empirical observation of binary cell-fusion as response to challenging environmental conditions, a feature absent in these earlier models. While the mechanism does not explain the evolution of sexual reproduction and genetic recombination itself, it does provide a nascent advantage to binary cell fusion that sets the stage for the evolution of sex by bringing nuclei from different cells into contact for prolonged periods. The mechanism also shows the potential to counter short-term costs associated with the initial formation of a binucleated cell. In this way the mechanism could facilitate the transition from horizontal gene transfer [94, 95] to meiotic recombination, which is advantageous when genome sizes increase in length [8]. Conceivably, if genetic recombination is beneficial for myriad genetic reasons in the long-term [8, 96], it would seem natural that it would be instigated when the opportunity arises (i.e. when physiological survival mechanisms bring nuclei into close contact). We note that it is obviously possible that the first diploid cells arose by errors in endomitosis [39, 51, 55] (essentially doubling the chromosome number within a single cell) and that meiosis first evolved in this context. Such a sequence of events is still compatible with our very general model, which can alleviate short-term costs of sex such as the energy involved in finding a partner and undergoing fusion [84]. In either scenario sexual reproduction may not be only a direct response to environmental variability [97, 98], but also to the correlated formation of a survival structure.

More generally, it is interesting to note that the conditions for facultative sexuality (e.g. harsh environmental conditions) broadly coincide with those for facultative multicellularity in both bacteria and eukaryotes, with starvation triggering the formation of fruiting bodies in myxobacteria [99, 100] and flocking in yeast [101, 102]. Meanwhile in *C. reinhardtii*, the formation of multicellular palmelloids and aggregates are an alternate stress response to sexual reproduction [103], as are the formation of fruiting bodies in *D. discoideum* [104]. In this multicellular context, the sexual behavior of *D. discoideum* is particularly interesting, as once formed, the zygote attracts hundreds of neighboring cells that are then cannibalized for the provision of a

macrocyst [105]. Conceptually, these various survival strategies are reminiscent of the mechanism in our model of the evolution of binary cell fusion in response to challenging environments. However it would be particularly interesting to extend our model to the multiple-fusion case of *D. discoideum*, where many cells forgo reproductive output to contribute to the macrocyst [106], to see under what conditions the results he have obtained here still qualitatively hold.

One element absent from our model is the fusion of multiple cells, which is likely to be selected for under the assumptions implicit in our model. There would clearly be an upper-limit on the number of fusions selected for, arising from the likely multiplicative effect of the fusion cost C. However in this context, it is interesting to note that one of the hypotheses for the evolution of self-incompatible mating types is as a signal to prevent the formation of polyploid cells [107]. Such a mechanism could also prevent the formation of trikaryotic cells should the cost of multiple fusions be too great. Extending our modeling in this direction may help explain the second stage in models for the evolution of eukaryotic sex, the regulation of cell–cell fusion [1].

The selection pressure for binary cell fusion is dependent on the formation of a binucleated cell with an increased survival benefit arising from its larger size. Although it is reasonable to assume that a single mononuclear cell has lower total resource requirements than a multicellular complex of the same size [108], we have not considered the detailed energetics of the maintenance of two nuclei [9]. In these respects incorporating dynamic energy budget theory into the model would be an important next step [109] as it would provide a clear distinction between the survival benefits of fused cells and unfused bicellular complexes. Within our modelling framework, these two structures are broadly similar [41]. However as we have shown, increased cell-cell attraction can be selected for even in the presence of large costs that one might expect under binary cell fusion but not associate with the formation of a bicellular complex.

We have assumed for simplicity a simple cell division scheme; parental cells undergo n rounds of symmetric division to produce 2^n daughter cells. In the context of multicellularity, switching environments have been shown to promote binary fragmentation [46]. However non-symmetric modes can be selected for [48] reflecting the diverse modes of facultative multicellular life cycles observed in bacteria [110]. It would be interesting to incorporate our results into models of cell division that account

explicitly for growth [111, 112] to determine how these results for multicellular [608] organisms carry over to the unicellular scenario, and further how they may affect those [609] we have shown here. [610]

Finally, we have not explicitly modelled any sources of cytoplasmic or genetic [611] conflict [113], which we have for simplicity included in the fusion cost C. Nevertheless, [612] social conflict does emerge in this model. In a recent paper we have shown how [613] evolutionary branching can arise, with some individuals producing fewer larger cells and [614] others producing more numerous but smaller daughter cells [50]. This branching is [615] driven by the same evolutionary forces that drive selection for anisogamy [114], in which [616] context this can be viewed as sexual conflict [115]. That social conflict should arise in [617] the formation of multicellular aggregates is well understood [41, 100, 116]. However these [618] models typically assume cells of fixed size [47, 117]. Combining the insights derived from [619] the evolution of anisogamy literature with the theory developed in the multicellularity [620] literature represents another promising research direction. [621]

As addressed above, trade-offs between cell fusion rate and mass [118], [622] cell-energetics, inbreeding, and the possibility of multiple cell-fusion events offer [623] interesting avenues to extend this analysis. In addition, we have not accounted for the [624] discrete nature of divisions leading to daughter cells, costs to phenotypic switching, [625] non-local trait mutations, or pre-existing mating types. More generally, extending our [626] mathematical approach leveraging adaptive dynamics to switching environments in [627] other facultatively sexual populations might prove particularly fruitful [62, 119]. [628]

In this paper we have adapted the classic PBS model [36] in two key ways; allowing [629] the fusion rate to evolve and subjecting the population to switching environments. In [630] doing so, we have shown its capacity to parsimoniously capture the evolution of obligate [631] binary cell fusion, obligate binary cell fission, and stressed induced binary cell fusion in [632] unicellular organisms. These results offer particularly interesting implications for the [633] evolution of binary cell-fusion as a precursor to sexual reproduction, as well as [634] suggesting common mechanistic links between the evolution of binary cell fusion and [635] multicellularity. Moreover, our analysis emphasises the importance of exploring the [636] coevolutionary dynamics of a range of evolutionary parameters, and of developing [637] computational and mathematical approaches to elucidate facultative sexual [638] reproduction. [639]

Supporting information

S1 Text. **Supporting information.pdf**

Acknowledgments

This work has made use of Viking high performance computing service at the University of York.

Author Contributions

Conceptualization: Jonathan Pitchford, George Constable

Methodology: Xiaoyuan Liu, George Constable, Jonathan Pitchford

Formal Analysis: Xiaoyuan Liu, George Constable

Software: Xiaoyuan Liu

Investigation: Xiaoyuan Liu

Supervision: George Constable, Jonathan Pitchford

Writing- Original draft preparation: Xiaoyuan Liu, George Constable

Writing- review & editing: Jonathan Pitchford, George Constable

References

1. Goodenough U, Heitman J. Origins of eukaryotic sexual reproduction. Cold Spring Harbor perspectives in biology. 2014;6(3):a016154.

2. Eme L, Spang A, Lombard J, Stairs CW, Ettema TJ. Archaea and the origin of eukaryotes. Nature Reviews Microbiology. 2017;15(12):711–723.

3. Eme L, Tamarit D, Caceres EF, Stairs C, de Anda V, Schon ME, et al. Inference and reconstruction of the heimdallarchaeial ancestry of eukaryotes. BioRxiv. 2023; p. 2023–03.

4. Maynard Smith J. The Evolution of Sex. Cambridge UK: Cambridge University Press; 1978.

5. Muller HJ. The relation of recombination to mutational advance. Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis. 1964;1(1):2–9.

6. Van Valen L. The red queen. The American Naturalist. 1977;111(980):809–810.

7. Hartfield M, Keightley PD. Current Hypotheses for the evolution of sex and recombination. Integr Zool. 2012;7(2):192–209.

8. Colnaghi M, Lane N, Pomiankowski A. Genome expansion in early eukaryotes drove the transition from lateral gene transfer to meiotic sex. Elife. 2020;9:e58873.

9. Lane N. Energetics and genetics across the prokaryote-eukaryote divide. Biology direct. 2011;6:1–31.

10. Rose MR. The contagion mechanism for the origin of sex. Journal of Theoretical Biology. 1983;101(1):137–146.

11. Hickey DA. Selfish DNA: a sexually-transmitted nuclear parasite. Genetics. 1982;101(3-4):519–531.

12. Hickey DA. Molecular symbionts and the evolution of sex. Journal of Heredity. 1993;84(5):410–414.

13. Radzvilavicius AL. Evolutionary dynamics of cytoplasmic segregation and fusion: mitochondrial mixing facilitated the evolution of sex at the origin of eukaryotes. Journal of Theoretical Biology. 2016;404:160–168.

14. Tilquin A, Christie JR, Kokko H. Mitochondrial complementation: a possible neglected factor behind early eukaryotic sex. Journal of Evolutionary Biology. 2018;31(8):1152–1164.

15. Harrison E, MacLean R, Koufopanou V, Burt A. Sex drives intracellular conflict in yeast. Journal of Evolutionary Biology. 2014;27(8):1757–1763.

16. Hurst LD, Hamilton WD. Cytoplasmic fusion and the nature of sexes. Proceedings of the Royal Society of London Series B: Biological Sciences. 1992;247(1320):189–194.

17. Hutson V, Law R. Four steps to two sexes. Proceedings of the Royal Society of London Series B: Biological Sciences. 1993;253(1336):43–51.

18. Speijer D. What can we infer about the origin of sex in early eukaryotes? Philosophical Transactions of the Royal Society B: Biological Sciences. 2016;371(1706):20150530.

19. Phadke S. Sex ratios, mate choice, and diversity in multi-sex systems of ciliates. University of Houston; 2011.

20. Umen JG. Genetics: Swinging ciliates' seven sexes. Current Biology. 2013;23(11):R475–R477.

21. Bishop C. Life cycle control of Chlamydomonas reinhardtii. Genome Biology. 2003;4(1):1–2.

22. Sieber B, Coronas-Serna JM, Martin SG. A focus on yeast mating: From pheromone signaling to cell-cell fusion. In: Seminars in Cell & Developmental Biology. Elsevier; 2022.

23. Ram Y, Hadany L. Condition-dependent sex: who does it, when and why? Philosophical Transactions of the Royal Society B: Biological Sciences. 2016;371(1706):20150539.

24. Fisher RA. The genetical theory of natural selection. Clarendon; 1930.

25. Waxman D, Peck JR. Sex and adaptation in a changing environment. Genetics. 1999;153(2):1041–1053.

26. Hadany L, Otto SP. The evolution of condition-dependent sex in the face of high costs. Genetics. 2007;176(3):1713–1727.

27. Hadany L, Otto SP. Condition-dependent sex and the rate of adaptation. the american naturalist. 2009;174(S1):S71–S78.

28. Vos M, Didelot X. A comparison of homologous recombination rates in bacteria and archaea. The ISME journal. 2009;3(2):199–208.

29. Redfield R. Genes for breakfast: The have-your-cake and-eat-lt-too of bacterial transformation. Journal of Heredity. 1993;84(5):400–404.

30. Redfield RJ. Do bacteria have sex? Nature Reviews Genetics. 2001;2(8):634–639.

31. Palchevskiy V, Finkel SE. Escherichia coli competence gene homologs are essential for competitive fitness and the use of DNA as a nutrient. Journal of bacteriology. 2006;188(11):3902–3910.

32. Narra HP, Ochman H. Of what use is sex to bacteria? Current Biology. 2006;16(17):R705–R710.

33. Shitut S, Shen MJ, Claushuis B, Derks RJ, Giera M, Rozen D, et al. Generating heterokaryotic cells via bacterial cell-cell fusion. Microbiology Spectrum. 2022;10(4):e01693–22.

34. Kuwabara T, Minaba M, Ogi N, Kamekura M. Thermococcus celericrescens sp. nov., a fast-growing and cell-fusing hyperthermophilic archaeon from a deep-sea hydrothermal vent. International journal of systematic and evolutionary microbiology. 2007;57(3):437–443.

35. Naor A, Gophna U. Cell fusion and hybrids in Archaea: prospects for genome shuffling and accelerated strain development for biotechnology. Bioengineered. 2013;4(3):126–129.

36. Parker GA, Baker RR, Smith V. The origin and evolution of gamete dimorphism and the male-female phenomenon. Journal of Theoretical Biology. 1972;36(3):529–553.

37. Lehtonen J. The Legacy of Parker, Baker and Smith 1972: Gamete Competition, the Evolution of Anisogamy and Model Robustness. Cells. 2021;10(3):573.

38. Cavalier-Smith T. Origin of the cell nucleus, mitosis and sex: roles of intracellular coevolution. Biology Direct. 2010;5:1–78.

39. Cavalier-Smith T. Cell cycles, diplokaryosis and the archezoan origin of sex. Archiv für Protistenkunde. 1995;145(3-4):189–207.

40. Mancebo Quintana J, Mancebo Quintana S. A Short-Term Advantage for Syngamy in the Origin of Eukaryotic Sex: Effects of Cell Fusion on Cell Cycle Duration and Other Effects Related to the Duration of the Cell Cycle—Relationship between Cell Growth Curve and the Optimal Size of the Species, and Circadian Cell Cycle in Photosynthetic Unicellular Organisms. International Journal of Evolutionary Biology. 2012;2012.

41. Tarnita CE, Taubes CH, Nowak MA. Evolutionary construction by staying together and coming together. Journal of theoretical biology. 2013;320:10–22.

42. Pentz JT, Márquez-Zacarías P, Bozdag GO, Burnetti A, Yunker PJ, Libby E, et al. Ecological advantages and evolutionary limitations of aggregative multicellular development. Current Biology. 2020;30(21):4155–4164.

43. Staps M, Van Gestel J, Tarnita CE. Emergence of diverse life cycles and life histories at the origin of multicellularity. Nature ecology & evolution. 2019;3(8):1197–1205.

44. Bonner JT. Cellular slime molds. vol. 2127. Princeton University Press; 2015.

45. Hamant O, Bhat R, Nanjundiah V, Newman SA. Does resource availability help determine the evolutionary route to multicellularity? Evolution & Development. 2019;21(3):115–119.

46. Pichugin Y, Park HJ, Traulsen A. Evolution of simple multicellular life cycles in dynamic environments. Journal of the Royal Society Interface. 2019;16(154):20190054.

47. Tang S, Pichugin Y, Hammerschmidt K. An environmentally induced multicellular life cycle of a unicellular cyanobacterium. Current Biology. 2023;33(4):764–769.

48. Pichugin Y, Traulsen A. Evolution of multicellular life cycles under costly fragmentation. PLoS computational biology. 2020;16(11):e1008406.

49. Gao Y, Pichugin Y, Gokhale CS, Traulsen A. Evolution of reproductive strategies in incipient multicellularity. Journal of the Royal Society Interface. 2022;19(188):20210716.

50. Liu X, Pitchford JW, Constable GWA. Adaptive dynamics, switching environments and the origin of the sexes. bioRxiv. 2024; p. 2024–03.

51. Wilkins AS, Holliday R. The evolution of meiosis from mitosis. Genetics. 2009;181(1):3–12.

52. Lee SC, Ni M, Li W, Shertz C, Heitman J. The evolution of sex: a perspective from the fungal kingdom. Microbiology and molecular biology reviews. 2010;74(2):298–340.

53. Umen JG. Green algae and the origins of multicellularity in the plant kingdom. Cold Spring Harbor perspectives in biology. 2014;6(11):a016170.

54. Rizos I, Frada MJ, Bittner L, Not F. Life cycle strategies in free-living unicellular eukaryotes: Diversity, evolution, and current molecular tools to unravel the private life of microorganisms. Journal of Eukaryotic Microbiology. 2024; p. e13052.

55. Hurst LD, Nurse P. A note on the evolution of meiosis. Journal of Theoretical Biology. 1991;150(4):561–563.

56. Cross FR, Umen JG. The Chlamydomonas cell cycle. The Plant Journal. 2015;82(3):370–392.

57. Harris EH. Chlamydomonas sourcebook. vol. 2. Academic Press San Diego; 1989.

58. Sager R, Granick S. Nutritional control of sexuality in Chlamydomonas reinhardi. The Journal of general physiology. 1954;37(6):729.

59. Innami R, Miyamura S, Okoshi M, Nagumo T, Ichihara K, Yamazaki T, et al. Gamete dimorphism of the isogamous green alga (Chlamydomonas reinhardtii), is regulated by the mating type-determining gene, MID. Communications Biology. 2022;5(1):1333.

60. Lv Y, Han F, Liu M, Zhang T, Cui G, Wang J, et al. Characteristics of N 6-methyladenosine modification during sexual reproduction of Chlamydomonas reinhardtii. Genomics, Proteomics & Bioinformatics. 2023;21(4):756–768.

61. Misamore MJ, Gupta S, Snell WJ. The Chlamydomonas Fus1 protein is present on the mating type plus fusion organelle and required for a critical membrane adhesion event during fusion with minus gametes. Molecular biology of the cell. 2003;14(6):2530–2542.

62. Czuppon P, Constable GWA. Invasion and extinction dynamics of mating types under facultative sexual reproduction. Genetics. 2019;213(2):567–580.

63. Constable GWA, Kokko H. The rate of facultative sex governs the number of expected mating types in isogamous species. Nat Ecol Evol. 2018;2(7):1168–1175.

64. Goodenough U, Lin H, Lee JH. Sex determination in Chlamydomonas. In: Seminars in cell & developmental biology. vol. 18. Elsevier; 2007. p. 350–361.

65. Vance RR. On reproductive strategies in marine benthic invertebrates. The American Naturalist. 1973;107(955):339–352.

66. Levitan DR. Optimal egg size in marine invertebrates: theory and phylogenetic analysis of the critical relationship between egg size and development time in echinoids. The American Naturalist. 2000;156(2):175–192.

67. Bulmer MG, Parker GA. The evolution of anisogamy: a game-theoretic approach. Proc R Soc Lond B. 2002;269:2381–2388.

68. Lehtonen J, Kokko H. Two roads to two sexes: unifying gamete competition and gamete limitation in a single model of anisogamy evolution. Behav Ecol Sociobiol. 2011;65:445–459. doi:10.1007/s00265-010-1116-8.

69. Kruzel EK, Hull CM. Establishing an unusual cell type: how to make a dikaryon. Current opinion in microbiology. 2010;13(6):706–711.

70. Hirooka S, Itabashi T, Ichinose TM, Onuma R, Fujiwara T, Yamashita S, et al. Life cycle and functional genomics of the unicellular red alga Galdieria for

elucidating algal and plant evolution and industrial use. Proceedings of the National Academy of Sciences. 2022;119(41):e2210665119.

71. Martin NC, Goodenough UW. Gametic differentiation in Chlamydomonas reinhardtii. I. Production of gametes and their fine structure. The Journal of cell biology. 1975;67(3):587–605.

72. Matsuda Y, Shimada T, Sakamoto Y. Ammonium ions control gametic differentiation and dedifferentiation in Chlamydomonas reinhardtii. Plant and cell physiology. 1992;33(7):909–914.

73. Rodilla V. Origin and evolution of binucleated cells and binucleated cells with micronuclei in cisplatin-treated CHO cultures. Mutation Research/Genetic Toxicology. 1993;300(3-4):281–291.

74. Ebersbach G, Gerdes K. Plasmid segregation mechanisms. Annu Rev Genet. 2005;39(1):453–479.

75. Hall AE, Rose MD. Cell fusion in yeast is negatively regulated by components of the cell wall integrity pathway. Molecular biology of the cell. 2019;30(4):441–452.

76. Warr J. A mutant of Chlamydomonas reinhardii with abnormal cell division. Microbiology. 1968;52(2):243–251.

77. Gillespie DT. Exact stochastic simulation of coupled chemical reactions. The journal of physical chemistry. 1977;81(25):2340–2361.

78. Rafaluk-Mohr C, Ashby B, Dahan DA, King KC. Mutual fitness benefits arise during coevolution in a nematode-defensive microbe model. Evolution Letters. 2018;2(3):246–256.

79. Brännström Å, Johansson J, Von Festenberg N. The hitchhiker's guide to adaptive dynamics. Games. 2013;4(3):304–328.

80. Kisdi É, Stefan A, Geritz H. Adaptive dynamics: a framework to model evolution in the ecological theatre. Journal of mathematical biology. 2010;61(1):165.

81. Tsai IJ, Bensasson D, Burt A, Koufopanou V. Population genomics of the wild yeast Saccharomyces paradoxus: quantifying the life cycle. Proceedings of the National Academy of Sciences. 2008;105(12):4957–4962.

82. Hasan AR, Ness RW. The genomic landscape of recombination rate variation in Chlamydomonas reinhardtii reveals a pronounced effect of linked selection. bioRxiv. 2018; p. 340992.

83. Agrawal AF. Evolution of sex: why do organisms shuffle their genotypes? Current Biology. 2006;16(17):R696–R704.

84. Lehtonen J, Jennions MD, Kokko H. The many costs of sex. Trends in ecology & evolution. 2012;27(3):172–178.

85. Heitman J. Evolution of sexual reproduction: A view from the fungal kingdom supports an evolutionary epoch with sex before sexes. Fungal Biology Reviews. 2015;29(3):108 – 117. doi:https://doi.org/10.1016/j.fbr.2015.08.002.

86. Nieuwenhuis BP, James TY. The frequency of sex in fungi. Philosophical Transactions of the Royal Society B: Biological Sciences. 2016;371(1706):20150540.

87. Waffenschmidt S, Woessner JP, Beer K, Goodenough UW. Isodityrosine cross-linking mediates insolubilization of cell walls in Chlamydomonas. The Plant Cell. 1993;5(7):809–820.

88. Gerber N, Kokko H. Abandoning the ship using sex, dispersal or dormancy: multiple escape routes from challenging conditions. Philosophical Transactions of the Royal Society B: Biological Sciences. 2018;373(1757):20170424.

89. Kokko H. When synchrony makes the best of both worlds even better: How well do we really understand facultative sex? The American Naturalist. 2020;195(2):380–392.

90. Otto SP, Lenormand T. Resolving the paradox of sex and recombination. Nature Reviews Genetics. 2002;3(4):252–261.

91. Hastings. The costs of sex due to deleterious intracellular parasites. Journal of Evolutionary Biology. 1999;12(1):177–183.

92. Togashi T, Cox PA, Bartelt JL. Underwater fertilization dynamics of marine green algae. Mathematical biosciences. 2007;209(1):205–221.

93. Nowak MA. Five rules for the evolution of cooperation. science. 2006;314(5805):1560–1563.

94. Bowler C, Allen AE, Badger JH, Grimwood J, Jabbari K, Kuo A, et al. The Phaeodactylum genome reveals the evolutionary history of diatom genomes. Nature. 2008;456(7219):239–244.

95. Schönknecht G, Chen WH, Ternes CM, Barbier GG, Shrestha RP, Stanke M, et al. Gene transfer from bacteria and archaea facilitated evolution of an extremophilic eukaryote. Science. 2013;339(6124):1207–1210.

96. Neiman M, Lively CM, Meirmans S. Why sex? A pluralist approach revisited. Trends in ecology & evolution. 2017;32(8):589–600.

97. Crow J. An advantage of sexual reproduction in a rapidly changing environment. Journal of Heredity. 1992;83(3):169–173.

98. Bürger R. Evolution of genetic variability and the advantage of sex and recombination in changing environments. Genetics. 1999;153(2):1055–1069.

99. Kaiser D, Manoil C, Dworkin M. Myxobacteria: cell interactions, genetics, and development. Annual Reviews in Microbiology. 1979;33(1):595–639.

100. Bonner JT. The origins of multicellularity. Integrative Biology: Issues, News, and Reviews: Published in Association with The Society for Integrative and Comparative Biology. 1998;1(1):27–36.

101. Verstrepen K, Derdelinckx G, Verachtert H, Delvaux F. Yeast flocculation: what brewers should know. Applied microbiology and biotechnology. 2003;61(3):197–205.

102. Day TC, Márquez-Zacarías P, Bravo P, Pokhrel AR, MacGillivray KA, Ratcliff WC, et al. Varied solutions to multicellularity: The biophysical and evolutionary

consequences of diverse intercellular bonds. Biophysics reviews. 2022;3(2):021305.

103. de Carpentier F, Lemaire SD, Danon A. When unity is strength: the strategies used by Chlamydomonas to survive environmental stresses. Cells. 2019;8(11):1307.

104. Strassmann JE, Queller DC. Evolution of cooperation and control of cheating in a social microbe. Proceedings of the National Academy of Sciences. 2011;108(supplement_2):10855–10862.

105. Douglas T, Queller DC, Strassmann JE. Social amoebae mating types do not invest unequally in sexual offspring. Journal of evolutionary biology. 2017;30(5):926–937.

106. Nickerson AW, Raper KB. Macrocysts in the life cycle of the Dictyosteliaceae. II. Germination of the macrocysts. American Journal of Botany. 1973;60(3):247–254.

107. Perrin N. What uses are mating types? The "developmental switch" model. Evolution: International Journal of Organic Evolution. 2012;66(4):947–956.

108. Raven JA. Allometry and stoichiometry of unicellular, colonial and multicellular phytoplankton. New phytologist. 2008;181:295–309.

109. Kooijman SALM. Dynamic energy budget theory for metabolic organisation. Cambridge university press; 2010.

110. Koyama T, Yamada M, Matsuhashi M. Formation of regular packets of Staphylococcus aureus cells. Journal of Bacteriology. 1977;129(3):1518–1523.

111. Iyer-Biswas S, Wright CS, Henry JT, Lo K, Burov S, Lin Y, et al. Scaling laws governing stochastic growth and division of single bacterial cells. Proceedings of the National Academy of Sciences. 2014;111(45):15912–15917.

112. Marantan A, Amir A. Stochastic modeling of cell growth with symmetric or asymmetric division. Physical Review E. 2016;94(1):012405.

113. Christie JR, Schaerf TM, Beekman M. Selection against heteroplasmy explains the evolution of uniparental inheritance of mitochondria. PLoS genetics. 2015;11(4):e1005112.

114. Maire N, Ackermann1 M, Doebeli M. Evolutionary branching and the evolution of anisogamy. Selection. 2002;2(1-2):119–131.

115. Parker GA. The sexual cascade and the rise of pre-ejaculatory (Darwinian) sexual selection, sex roles, and sexual conflict. Cold Spring Harbor perspectives in biology. 2014;6(10):a017509.

116. Tarnita CE, Washburne A, Martinez-Garcia R, Sgro AE, Levin SA. Fitness tradeoffs between spores and nonaggregating cells can explain the coexistence of diverse genotypes in cellular slime molds. Proceedings of the National Academy of Sciences. 2015;112(9):2776–2781.

117. Pichugin Y, Peña J, Rainey PB, Traulsen A. Fragmentation modes and the evolution of life cycles. PLoS computational biology. 2017;13(11):e1005860.

118. Siljestam M, Martinossi-Allibert I. Anisogamy does not always promote the evolution of mating competition traits in males. The American Naturalist. 2024;203(2):230–253.

119. Krumbeck Y, Constable GWA, Rogers T. Fitness differences suppress the number of mating types in evolving isogamous species. Royal Society Open Science. 2020;7:192126.