CTCF N-terminal Domain Regulates Clustered Protocadherin Gene Expression by Enhancing Cohesin Processivity

Yijun Zhang, Qiang Wu∗

Center for Comparative Biomedicine, Ministry of Education Key Laboratory of Systems Biomedicine, State Key Laboratory of Medical Genomics, Institute of Systems Biomedicine, Shanghai Jiao Tong University, Shanghai 200240, China

∗Corresponding author: qiangwu@sjtu.edu.cn

Abstract

CTCF architects 3D genome folding by anchoring or forestalling cohesin loop extrusion, but the exact mechanism remains obscure. Here, using clustered protocadherins (*cPcdh*) as model genes, we report that CTCF stabilizes or facilitates cohesin loop extrusion by enhancing its processivity. Specifically, we show that, compared with the *Pcdh* α and γ clusters, the *Pcdhβ* cluster is greatly affected upon CTCF$^{Y226A/F228A}$ mutation in the N-terminal domain. Given the long-range distance of the *Pcdhβ* cluster from the distal enhancer, this finding has interesting implications in CTCF regulation of cohesin processivity along the linear chromatin during DNA loop extrusion. In particular, the effect on cohesin processivity upon CTCF$^{Y226A/F228A}$ mutation is conspicuously similar to that of WAPL overexpression, demonstrating that, contrary to the general view of blocking or forestalling cohesin, CTCF actually enhances or facilitates cohesin loop extrusion.

Keywords: Clustered protocadherin; cohesin processivity; CTCF YDF motif; Gene regulation; enhancer-promoter interactions; loop extrusion

Introduction

The *cPcdh* locus comprises three closely-linked clusters of α, β, and γ, spanning a large region of ~1 Mb genomic DNA in mice and encoding 58 distinct PCDH protein isoforms required for specifying an enormous diversity of neural connectivity in the brain (Wu and Maniatis, 1999; Wu et al., 2001). The genomic organization of the *Pcdh* α and γ clusters is similar, each including a repertoire of large variable exons in a tandem array and three shared downstream constant exons (Wu and Maniatis, 1999; Wu et al., 2001). Each variable exon is transcribed independently under the control of its own promoter and then spliced to the respective set of downstream constant exons to encode diverse PCDH isoforms. However, members of the *Pcdhβ* cluster each are only made up of a single variable exon without constant ones and thus do not undergo splicing (Tasic et al., 2002; Wu and Maniatis, 1999) (Figure 1A). In addition, all *cPcdh* except c-type genes exhibit stochastic and monoallelic expression patterns in the developing and adult central nervous system (Canzio et al., 2019; Esumi et al., 2005; Jia et al., 2020; Kiefer et al., 2023; Lv et al., 2022; Wu and Maniatis, 1999). Moreover, the encoded PCDH proteins act as vital cell-surface tags for neural circuit formation, providing individual neurons with a unique molecular 'identifier' required for dendritic self-avoidance and axonal tiling in distinct brain regions (Chen et al., 2017; Dong et al., 2023; Ing-Esteves et al., 2018; Kiefer et al., 2023; Lefebvre et al., 2012; Lv et al., 2022; Mountoufaris et al., 2017; Schreiner and Weiner, 2010; Thu et al., 2014). Dysregulation of the *cPcdh* genes is highly relevant to human mental diseases, including autism spectrum disorders (ASD), Down syndrome, and Alzheimer's disease (AD) (Flaherty and Maniatis, 2020; Jia and Wu, 2020).

The identification of *cis* regulatory elements (CREs) in the *cPcdh* gene clusters has shed significant insights into the mechanism involved in the *cPcdh* promoter choice. Initial DNase I hypersensitivity assay has characterized several CREs, including a cluster of downstream enhancers of *HS7* and *HS5-*

1 (super-enhancer) for the *Pcdhα* cluster (Kehayova et al., 2011; Ribich et al., 2006). Subsequent studies have identified a large cluster of downstream enhancers of *HS7L*, *HS5-1L*, *HS18-20*, *HS21*, and *HS22* (super-enhancer) for the *Pcdh β* and *γ* clusters (Ge et al., 2023; Guo et al., 2012; Guo et al., 2015; Jia et al., 2020; Yokota et al., 2011). Interestingly, members of all three *Pcdh* clusters contain tandem-arrayed CTCF (CCCTC-binding factor) sites in both promoters and enhancers. A striking arrangement of CTCF sites within clustered promoters and enhancers is that the CTCF sites in the promoter regions are in the opposite orientation to those in the enhancer regions, forming convergent CTCF pairs (Figure 1A) (Ge et al., 2023; Guo et al., 2012; Guo et al., 2015; Jia et al., 2020). The CTCF/cohesin-mediated long-distance chromatin interactions between distal enhancers and target promoters are the 3D genomic basis of the *cPcdh* promoter choice (Ge et al., 2023; Guo et al., 2012; Guo et al., 2015; Jia et al., 2020; Zhou et al., 2021). Specifically, the tandem directional CTCF sites within the downstream super-enhancers form spatial contacts with the proximal and distal promoter CBS elements through continuous active cohesin "loop extrusion" to erase the genomic-distance biases in favor of stochastic promoter choice (Ge et al., 2023; Jia et al., 2020; Kiefer et al., 2023). However, the physiological functions of the dynamics of cohesin loop extrusion in the *cPcdh* locus remain poorly understood.

Here, through CRISPR editing of the YDF motif—the key cohesin interacting moiety—of the CTCF N-terminal domain (NTD) in *N2a* cells *in vitro* and in mice *in vivo*, in conjunction with RNA-seq, ChIP-seq, 4C, and Hi-C analyses, we found that members of the *Pcdhβ* cluster are highly susceptible to the perturbed cohesin processivity owing to their distal genomic arrangement relative to the downstream super-enhancer region. Our data broaden the molecular logic that generates Pcdh isoform diversity required for neural circuit assembly via a unique 3D genome organization.

Results

Alteration of *Pcdhβ* expression patterns upon CTCF ADA mutation

Given that CTCF/cohesin-mediated DNA looping between the active promoter and target enhancer is required for the stochastic activation of the *cPcdh* genes (Canzio et al., 2019; Guo et al., 2012), we investigated how the improper positioning of cohesin affects the promoter choices of the *cPcdh* genes. To this end, we mutated the highly conserved amino acid motif YDF to ADA using DNA-fragment editing through CRISPR single-cell screening in Neuro-2a (*N2a*) cells. Specifically, we screened 166 single-cell clones and obtained two independent homozygous single-cell clones of CTCF ADA mutation (Figure S1A, B).

We performed RNA-seq experiments to assess expression alterations of members of the three *Pcdh* clusters upon CTCF ADA mutation (Figure 1B-D). Remarkably, we observed a significant downregulation in the expression levels of a large repertoire of the *Pcdhβ* cluster (Figure 1C). By contrast, it appears that the effects on the *Pcdh* α and γ clusters were less pronounced than those on the *Pcdhβ* cluster (Figure 1B and D). The distinct transcriptional alterations observed within the *cPcdh* locus suggest that members of the *Pcdh* β and $\alpha\gamma$ clusters exhibit varying degrees of sensitivity to CTCF ADA mutation. As a control, we found no significant alteration of *Ctcf* expression levels upon ADA mutation (Figure 1E). Genome-wide expression analyses revealed that ~5,000 genes exhibited a significant alteration of expression levels upon CTCF ADA mutation. Of note, we observed a higher number of genes downregulated compared to those upregulated (Figure S1C, D).

Figure 1

Figure 1. Altered *cPcdh* expression patterns in the CTCF ADA mutant. (**A**) Schematic of the mouse cPcdh locus. The *Pcdhα* cluster is regulated by a downstream super-enhancer composed of *HS7* and *HS5-1* whereas the *Pcdhβγ* clusters are regulated by a downstream super-enhancer composed of a cluster of enhancers of *HS7L-HS22*. The location and orientation of CTCF sites are indicated by arrowheads. Var, variable; Con, constant; SE, super-enhancer. (**B-D**) RNA-seq of wild type (WT) and CTCF ADA mutant (ADA) Neuro-2a (*N2a*) clones (n=3). (**E**) Quantification of the expression levels of the *Ctcf* gene. (**F**) The representative photograph of the WT and heterozygous ADA P0 mice, with their genotypes indicated. (**G-I**) Manhattan plots showing localized enrichments (using 1Mb^{sw}) of downregulated genes in the heterozygous P0 (**G**) and adult (**H**) cortical, as well as olfactory bulb (**I**) tissues, with *cPcdh* as a top-ranking locus. (**J-L**) Quantitative analyses of the *Ctcf* gene in the heterozygous P0 (**J**) and adult (**K**) cortical, as well as olfactory bulb (**L**) tissues. NS, non-significant.

Generation of heterozygous *Ctcf*$^{Y226A/F228A}$ mice

To further elucidate the crucial role of the YDF motif within the CTCF NTD in regulating cohesin processivity on DNA *in vivo*, we generated for the first time the *Ctcf*$^{Y226A/F228A}$ mice by pronuclear microinjection using the CRISPR/Cas9 system (Figure S1). Briefly, we designed single guide RNAs (sgRNAs) programming Cas9 to a region adjacent to the genomic sequences encoding the CTCF Y226 and F228 residues. A single-stranded oligodeoxynucleotide (ssODN) donor was co-injected with each sgRNA and Cas9 mRNA to introduce the targeted mutations by homology-directed repair (HDR) (Figure S1E, F). By genotyping 61 P0 chimeric mice, we obtained one mouse with the precise CTCF ADA mutation (Figure S2A) and 14 mice with random indels. The chimeric *Ctcf*$^{Y226A/F228A}$ mouse was crossed with a wild-type C57BL/6J mouse to generate heterozygous CTCF ADA mutant mice. We obtained two heterozygous *Ctcf*$^{Y226A/F228A}$ mice from a total of 5 litters with only 19 pups, suggesting that this CTCF mutation may affect its litter size or fertility (Figure S2B, C). Although the heterozygous *Ctcf*$^{Y226A/F228A}$ mice appear to be normal compared to the wild-type littermates (Figure 1F; Figure S2D-I), we were unable to obtain homozygous *Ctcf*$^{Y226A/F228A}$ mice after repeated breeding attempts. Consequently, we performed all experiments using heterozygous mice.

Prominent *cPcdh* expression alterations in heterozygous *Ctcf*$^{Y226A/F228A}$

mice

We performed RNA-seq experiments using the heterozygous P0 and adult cortical as well as olfactory bulb tissues to further investigate the functional implications of CTCF ADA mutation on gene expression. Consistent with the 3D genome architectural role of CTCF, we found a broad spectrum of genes that display significant expression alterations. Remarkably, 9.3% of the downregulated genes are members of the *Pcdh* clusters, resulting in a unique, ~231-fold enrichment for downregulated transcripts in the P0 cortex at this locus, as compared to the rest of the genome (P = 6.48×10^{-56} by Poisson test; 1-Mb sliding window (1Mb^{sw}) applied to n = 302 transcripts) (Figure 1G). There is a similar unique enrichment of *cPcdh* genes for downregulated transcripts in both adult cortex and olfactory bulb although the enrichment is less pronounced in the olfactory bulb (Figure 1H and I). As controls, the expression levels of *Ctcf* in the mutant mice are comparable to those in wild-type littermates, suggesting that CTCF ADA mutations do not perturb *Ctcf* expression (Figure 1J-L). Finally, we calculated the proportion of mutated bases of *Ctcf* transcripts and observed that the ratios of mutated bases were comparable to those of wild-type bases (Figure S1G-I).

CTCF YDF motif regulates quantitative *Pcdhβ* expression

Differential gene expression analyses in the heterozygous P0 cortex, adult cortex, and adult olfactory bulb revealed that among all of the downregulated *cPcdh* genes, the vast majority are members of the *Pcdhβ* cluster (Figure 2A-C). We then calculated expression alterations of each *cPcdh* gene. Remarkably, almost all members of the *Pcdhβ* cluster displayed a significant decrease of expression levels in both the neocortical and olfactory bulb tissues upon CTCF ADA mutation (Figure 2D-F). In addition, members of the *Pcdhβ* cluster exhibited the largest decrease of expression levels compared to the other two clusters in the heterozygous *Ctcf*$^{Y226A/F228A}$ mice (Figure 2G-I). This suggests that CTCF ADA mutation has a much stronger effect on members of the *Pcdhβ*

A Fold change of gene expression in P0 cortex

B Fold change of gene expression in adult cortex

C Fold change of gene expression in adult olfactory bulb

D
Pcdhβ cluster

Pcdhγ cluster

Pcdhγ c-type

E
Pcdhβ cluster

Pcdhγ cluster

Pcdhγ c-type

F
Pcdhβ cluster

Pcdhγ cluster

Pcdhγ c-type

G Fold change of cPcdh genes in P0 cortex

H Fold change of cPcdh genes in adult cortex

I Fold change of cPcdh genes in adult olfactory bulb

Figure 2. Prominent *Pcdhβ* expression alterations upon CTCF ADA mutation. (**A-C**) Volcano plots showing all of the downregulated *cPcdh* genes in the heterozygous, compared to their WT littermates, P0 (n=4) (**A**) and adult (n=3) (**B**) cortical, as well as olfactory bulb (n=3) (**C**) tissues. (**D-F**) Shown are expression alterations of repertoires of the *Pcdh β* and *γ* clusters in the heterozygous P0 (n=4) (**D**) and adult (n=3) (**E**) cortical, as well as olfactory bulb (n=3) (**F**) tissues. (**G-I**) Bar plots depicting fold changes of gene expression levels of *cPcdh* genes in the heterozygous P0 (n=4) (**G**) and adult (n=3) (**H**) cortical, as well as olfactory bulb (n=3) (**I**) tissues.

cluster than that on those of the *Pcdh α* and *γ* clusters.

CTCF ADA mutation affects *cPcdh* expression in a genomic distance-biased manner

Increasing evidence has implicated that cohesin-mediated DNA "loop-extrusion" determines *cPcdh* gene choice via spatiotemporal regulation of long-distance chromatin contacts between distal enhancers and target variable promoters (Canzio et al., 2019; Guo et al., 2012; Guo et al., 2015; Jia et al., 2020; Kiefer et al., 2023; Monahan et al., 2012). We asked whether the decreased levels of *cPcdh* gene expression upon CTCF ADA mutation are truly attributed to the DNA looping mediated by cohesin. To this end, we performed ChIP-seq experiments with a specific antibody against RAD21, a cohesin subunit, in both our cellular and mouse models (Figure 3; Figure S3 and S4).

Interestingly, we observed a genomic distance-biased decrease of cohesin enrichments for members of *cPcdh* genes, suggesting defects in cohesin processivity upon CTCF ADA mutation (Figure 3A and B). Specifically, in the context of the *Pcdhα* cluster, we found a significant decrease of RAD21 enrichments at *Pcdh α8*, *α9*, and *αc1* (Figure 3A and B; Figure S3), which have CTCF sites associated with their promoters (Figure 1A). By contrast, as an internal control, the RAD21 enrichments remain largely unchanged at the *HS7* enhancer (Figure S3E), which has no associated CTCF site (Figure 1A) (Kehayova et al., 2011). In the context of the *Pcdhβγ* clusters, ChIP-seq experiments revealed a significant decrease of the deposition of RAD21 at the super-enhancer region as well as at the variable promoter regions of the *Pcdhβγ*

Figure 3

Figure 3. CTCF YDF motif regulates expression levels of *cPcdh* genes in a genomic distance-biased manner. (**A**) ChIP-seq profiles of RAD21 at the *cPcdh* locus from WT and CTCF ADA mutant *N2a* clones. (**B**) Top: Linear genomic distance of *cPcdh* promoters with respect to their enhancers. Bottom: Scatterplots showing fold changes of RAD21 enrichments at members of three *Pcdh* gene clusters in *N2a* cell clones as a function of the relative linear genomic distance of their promoters to the respective enhancers. (**C**) Scatterplots showing fold changes of expression levels of *cPcdh* genes in *N2a* cell clones as a function of relative linear genomic distance between variable promoters and their respective enhancers. (**D**) ChIP-seq profiles of RAD21 at the *cPcdh* locus in WT and heterozygous CTCF ADA mice. (**E**) Top: Linear genomic distance of *cPcdh* promoters with respect to their enhancers. Bottom: Scatterplots showing fold changes of RAD21 enrichments at the *cPcdh* locus in the heterozygous CTCF ADA mice as a function of the linear genomic distance between target variable promoters and their respective enhancers. (**F**) Scatterplots showing fold changes of expression levels of *cPcdh* genes as a function of the linear genomic distance between target variable promoters and their respective enhancers in the heterozygous ADA mice.

clusters (Figure 3A and B; Figure S3F and G). In particular, the decrease of RAD21 enrichment at the *Pcdhβ* cluster is much more prominent than that at the *Pcdh* α and γ clusters (Figure S3A-C). Interestingly, the genomic distance-biased decrease of cohesin enrichments is consistent with decreased expression levels of members of *cPcdh* genes upon CTCF ADA mutation in *N2a* cell clones (Figure 3B and C).

We observed a similar genomic distance-biased decrease of expression levels of chosen *cPcdh* genes in the heterozygous *Ctcf*$^{Y226A/F228A}$ mice *in vivo* (Figure 3D-F; Figure S4). As controls, the enrichment of CTCF itself appears to be unchanged upon CTCF ADA mutation (Figure S5 and S6). Taken together, these data reveal a genomic distance-biased decrease of *cPcdh* gene expression in both cellular and mouse models.

CTCF ADA mutation affects cohesin processivity

To investigate alterations of cohesin-mediated chromatin interactions between distal enhancers and target variable promoters upon CTCF ADA mutation, we performed *in situ* Hi-C experiments and analyzed contact matrices of the *cPcdh* locus. We first evaluated the impact of the ADA mutation on the TAD boundaries

using the directionality index algorithm and observed a significant weakening of the strength of *cPcdh* superTAD and subTAD boundaries (Figure 4A) (Guo et al., 2015; Jiang et al., 2017). In addition, we observed a significant reduction of chromatin contacts within the entire *cPcdh* superTAD upon CTCF ADA mutation (Figure 4B), consistent with the significant decrease of RAD21 enrichments at the *cPcdh* locus (Figure 3D-F). Moreover, there is a significant decrease of long-distance chromatin interactions between the *HS5-1* enhancer and *Pcdhα* variable promoters (Figure 4C). Furthermore, quantitative high-resolution chromosome conformation capture with *HS5-1bL*, *HS18*, or *HS19-20* as an anchor (QHR-4C) revealed a significant decrease of long-range chromatin interactions between the downstream super-enhancer and its target *Pcdh$\beta\gamma$* clusters (Figure 4D). Finally, we observed a genomic distance-biased defect in the chromatin contact probability for the *cPcdh* genes upon CTCF ADA mutation (Figure 4E). In particular, although members of both *Pcdh* β and γ clusters are activated by the same downstream super-enhancer via cohesin-mediated loop extrusion (Ge et al., 2023; Jia et al., 2020; Kiefer et al., 2023), the impact on the *Pcdhβ* cluster is much stronger than that on the *Pcdhγ* cluster owing to the greater distance from *Pcdhβ* to the super-enhancer (Figure 4E). Finally, Hi-C analyses revealed that CTCF ADA mutation mainly affects chromatin loops of large sizes genome-wide (Figure S7). Together, these data strongly suggest a cohesin-processivity defect upon CTCF ADA mutation.

Figure 4

Figure 4. CTCF ADA mutation results in lower cohesin processivity and decreased spatial contact probability between distal promoter-enhancer pairs. (**A**) Hi-C contact maps, directionality index scores, and loop intensities around *cPcdh* locus at the chromosome 18 showing a widespread weakening of chromatin loop strength upon CTCF ADA mutation. The differential contact heatmap is also shown. (**B**) Close-up of contact maps in the *cPcdh* locus at 10-Kb resolution, alongside the differential contact heatmap at the *cPcdh* locus between heterozygous CTCF ADA mutant and WT mice. (**C,D**) Shown are QHR-4C profiles using *HS5-1*, *HS5-1bL*, *HS18*, or *HS19-20* as an anchor. (**E**) Scatterplots depicting fold changes of contact probabilities of *cPcdh* genes as a function of the linear genomic distance between target variable promoters and their respective enhancers in mouse and cellular models. (**F**) A model for CTCF regulation of *cPcdh* gene expression patterns in a genomic distance-biased manner. CTCF ADA mutation interferes the interaction with cohesin, resulting in its short residence time on chromatin fibers and low processivity. This leads to prominent dysregulation of members of the *Pcdhβ* cluster.

Discussion

CTCF/cohesin are central for *cPcdh* gene choice

Given that there are approximately 86 billion neurons in the human brain and individual neurons display highly-branched axons and dendrites that intricately connect with one another in diverse brain regions with minimal overlap, it is enigmatic how neurons generate such sophisticated synaptic connectivity to organize precise neural networks for information processing. The cPCDH proteins—a subgroup of cadherin-like cell adhesion molecules—emerge as central candidates for proper wiring of neural circuits (Chen et al., 2017; Dong et al., 2023; Fan et al., 2018; Kiefer et al., 2023; Lv et al., 2022; Mountoufaris et al., 2018; Mountoufaris et al., 2017; Sanes and Zipursky, 2020; Wu and Jia, 2021; Wu and Maniatis, 1999). The CTCF/cohesin-mediated long-distance chromatin contacts between distal super-enhancers and target variable promoters are central for *cPcdh* gene choice during neural development (Canzio et al., 2019; Ge et al., 2023; Guo et al., 2012; Guo et al., 2015; Jia et al., 2020; Kiefer et al., 2023). Here we have, for the first time, generated a *Ctcf$^{Y226A/F228A}$* mutant mouse model and provided strong genetic evidence that *Ctcf$^{Y226A/F228A}$* mutation severely impacts expression levels of the repertoire of the *cPcdh* genes. In particular, RNA-seq, in conjunction with 4C and Hi-C,

showed that members of the $Pcdh\beta$ cluster are strongly affected in both single-cell *N2a* clones *in vitro* and in heterozygous mice *in vivo* owing to the greatest distance from the super-enhancer. Given the stochastic and monoallelic nature of *cPcdh* gene expression in the brain, it is particularly remarkable that prominent dysregulation of members of the $Pcdh\beta$ cluster could be observed even in the heterozygous CTCF ADA mutants.

Constitutive CTCF deletion in mice leads to early zygotic lethality, indicating an essential role of CTCF during embryo development (Fedoriw et al., 2004; Moore et al., 2012). In addition, conditional CTCF deletion in neurons results in a marked decrease in dendritic arborization and spine density owing to a significant reduction in the expression levels of 53 isoforms of the *cPcdh* genes (Hirayama et al., 2012). We observed that the majority of downregulated *cPcdh* genes are members of the $Pcdh\beta$ cluster upon $Ctcf^{Y226A/F228A}$ mutation in both mouse and cellular models. This observation is reminiscent of the significant reduction of expression levels of mostly members of the $Pcdh\beta$ cluster in the heterozygous *Nipbl* deletion mice (Kawauchi et al., 2009; Remeseiro et al., 2013). Finally, the regulation of *cPcdh* genes, especially the distal ones, in the brain is related to the SA1-localization at the variable promoters (Remeseiro et al., 2012).

Emerging CTCF-related neurodevelopmental disorders resulting from numerous *de novo* heterozygous CTCF mutations frequently manifest as syndromic intellectual disability and neurodevelopmental delay such as microcephaly (Chen et al., 2019; Price et al., 2023; Valverde de Morales et al., 2023). In addition, heterozygous mutations in subunits of cohesin complex (SMC1, SMC3, RAD21) or its chromatin loader (NIPBL) cause Cornelia de Lange syndrome (CdLS) which also frequently manifests as intellectual disability and neurodevelopmental delay (Mannini et al., 2013; Nasmyth and Haering, 2009). The important role of CTCF/cohesin in regulating *cPcdh* gene expression suggests a neuropathogenic explanation for syndromic intellectual

disability from CTCF/cohesin mutations. Indeed, genetic, epigenetic, and 3D genomic mutations and the resulting dysregulation of *cPcdh* gene expression are related to the etiology of numerous neuropsychiatric diseases (Flaherty and Maniatis, 2020; Jia and Wu, 2020). The fact that expression patterns of the *cPcdh* genes are compromised in CTCF ADA mutation, *Nipbl* and *SA1* depletion or mutation, along with the prominent *cPcdh* brain-wiring function, suggest that the dysregulation of *cPcdh* genes resulted from defective cohesin processivity may underlie the pathogenesis of various neurodevelopmental disorders of cohesinopathy.

CTCF N-terminal domain promotes cohesin processivity

The asymmetrical blocking of extruding cohesin complexes by CTCF at opposite CBS elements in enhancers and promoters leads to the formation of long-distance chromatin loops and determines the promoter choice of *cPcdh* genes (Ge et al., 2023; Guo et al., 2012; Guo et al., 2015; Jia et al., 2020; Lu et al., 2019). Specifically, tandem CTCF sites located at *cPcdh* variable promoters function as topological chromatin insulators to erase the proximity-biased cohesin loop extrusion and enhance the activation of distal promoters (Jia et al., 2020). In addition, antisense lncRNA transcription leads to DNA demethylation and CTCF deposition and ensures balanced choice of both distal and proximal promoters (Canzio et al., 2019). Moreover, CDCA7 (cell division cycle associated 7) associated DNA methylation loss translates into aberrant upregulation of *cPcdh* genes (Vukic et al., 2024), especially members of the *Pcdhβ* cluster which are distal to the enhancers. The orientation-specific function of CTCF sites are achieved by a combination of its stringent CTCF anti-parallel recognition (Hashimoto et al., 2017; Yin et al., 2017) and polar interactions between CTCF NTD and cohesin complex (Li et al., 2020; Nora et al., 2020; Pugacheva et al., 2020; Zhang et al., 2023). Given that NTD truncation of CTCF compromises its stalling activity of cohesin loop extrusion and leads to decreased cohesin enrichments at CTCF sites (Li et al., 2020;

Nora et al., 2020; Pugacheva et al., 2020), it is surprising that the ADA mutation does not result in enhanced cohesin processivity. Nevertheless, the compromised cohesin processivity upon CTCF ADA mutation (Figure 4) is consistent with the role of CTCF in promoting distal promoter-enhancer interactions and the topological insulation function of tandem CTCF sites (Jia et al., 2020).

Previous studies systematically dissected the enhancer elements of the *Pcdh* gene clusters. The hypersensitive sites *HS5-1* and *HS7*, situated downstream of the *Pcdhα* cluster, were identified as a composite or super-enhancer for the *Pcdhα* cluster (Canzio et al., 2019; Guo et al., 2012; Guo et al., 2015; Kehayova et al., 2011; Kiefer et al., 2023; Monahan et al., 2012; Ribich et al., 2006; Zhou et al., 2021). Additionally, the super-enhancer of *HS5-1bL*, *HS18*, and *HS19-20* predominantly regulates the expression levels of the *Pcdh* β and γ clusters (Ge et al., 2023; Guo et al., 2012; Guo et al., 2015; Jia et al., 2020; Kiefer et al., 2023; Yokota et al., 2011). The linear genomic distance of the variable promoters relative to the super-enhancers and CTCF/cohesin-mediated specific promoter-enhancer looping interactions are keys for *cPcdh* gene choice. Interestingly, WAPL (wings apart-like protein homolog) dosage in different brain regions functions as a rheostat in regulating *cPcdh* isoform diversity via cohesin processivity (Kiefer et al., 2023). Specifically, high WAPL levels in serotonergic neurons lead to low cohesin processivity and promote expression of enhancer-proximal *αc2*; whereas low WAPL levels in the olfactory bulb result in high cohesin processivity, which leads to promoter choice of the enhancer-distal alternate isoforms (Kiefer et al., 2023).

We report that the CTCF ADA mutation leads to low cohesin processivity and compromised expression of the members of the *Pcdhβ* cluster, which have the greatest distance from the distal super-enhancer. We propose a model in which the CTCF ADA mutation loses its competition with WAPL for binding to cohesin, resulting in low processivity and compromised expression of the

Pcdhβ genes (Figure 4F). The YDF motif of CTCF N-terminal domain interacts with the conserved essential surface (CES) of cohesin (Li et al., 2020; Zhang et al., 2023). It was initially proposed that the CTCF YDF motif competes with the WAPL N-terminal FGF motifs for binding to cohesin CES (Li et al., 2020). However, recent extensive AlphaFold analyses of cohesin subcomplexes suggest that CTCF YDF motif most probably competes with the WAPL C-terminal domain for binding to cohesin CES (Nasmyth et al., 2023). An alternative mechanism of CTCF in enhancing cohesin processivity could be through interference with the recruitment of WAPL by PDS5 (precocious dissociation of sisters 5), thus compromising the cohesin releasing activity (Nasmyth et al., 2023; Nora et al., 2020). Whatever the underlying mechanism, our data suggest CTCF YDF has an anti-cohesin releasing activity by competing with WAPL. The exact mechanism by which CTCF and WAPL regulate cohesin processivity or chromatin association awaits future studies.

Methods

Cell culture

Mouse Neuro-2a (*N2a*) cells were cultured in a modified Eagle's medium (Gibco 11095080) supplemented with 10% fetal bovine serum (FBS) (Sigma-Aldrich F0193), 1% penicillin-streptomycin (Gibco 15140122), and 1% Non-Essential Amino Acids (Gibco 11140050). Typically, *N2a* cells were incubated at 37°C in a humidified incubator with 5% (v/v) CO_2.

For cell passaging, when cells reached about 80% confluency, the culture medium was aspirated from the cell culture dish. The cells were then washed gently once with 1 ml of 1 × PBS (Gibco 70011044) (*Supplementary file 2*) to remove residual medium and cell debris. 1 ml of 0.25% trypsin-EDTA (Gibco 25200056) was added, followed by incubation at 37°C for 3 min to detach cells from the dish surface. Next, 1 ml of cell culture medium was added to neutralize the trypsin. The cell suspension is then transferred to a new 15-ml centrifuge tube and centrifuged at 900 rpm for 5 min at room temperature. The supernatant was discarded, and 1 ml of fresh medium was added to resuspend the cell pellet. Finally, cells are sub-cultured via transferring 250 μl of cell suspension into a new dish containing 5 ml of fresh culture medium.

Plasmid construction

The construction of plasmids (*Supplementary file 2*) for the generation of CTCF ADA mutant (ADA) Neuro-2a (*N2a*) clones was carried out as previously described (Li et al., 2015; Shou et al., 2018). In brief, to generate constructs for sgRNA transcription, 2 μl each of two complementary oligonucleotides (Mut_sgRNA_F and Mut_sgRNA_R) (*Supplementary file 1*) at a concentration of 20 μM and 2 μl of NEBuffer 2 (NEB B7002S) were added into an Eppendorf tube, and the reaction volume was adjusted to 20 μl with nuclease-free water. The reaction mixture was heated in boiling water for 1 min, then allowed to gradually cool to room temperature to form double-stranded DNA (dsDNA) with

5′ overhangs of 'ACCG' and 'AAAC'. In addition, 1 μl of *Bsa*I (NEB R3733S) was added to another Eppendorf tube containing 3 μg of *pGL3* plasmid at a total volume of 50 μl, and the mixture was incubated at 37°C for 2 h to obtain a *Bsa*I-linearized *pGL3* vector. Finally, the annealed dsDNA is then cloned into a *Bsa*I-linearized *pGL3* vector with *U6* promoter to enable sgRNA transcription with the U6 polymerase.

For construction of donor plasmids used in homologous recombination, the left homologous arm of 883-bp was generated by PCR (95°C, 3 min; 95°C, 15 s, 58°C, 15 s, 72°C, 1 min for 35 cycles; and a final extension at 72°C, 5 min) from total mouse genomic DNA with a forward primer (Left_Mut_F) (*Supplementary file 1*) containing vector backbone sequences and a reverse primer (Left_Mut_R) (*Supplementary file 1*) containing the designed CTCF ADA mutation and a synonymous mutation at its 5' end. In addition, the right homologous arm of 885-bp were generated by PCR (95°C, 3 min; 95°C, 15 s, 60°C, 15 s, 72°C, 1 min for 35 cycles; and a final extension at 72°C, 5 min) from total mouse genomic DNA with a forward primer (Right_Mut_F) (*Supplementary file 1*) containing the same mutations and a reverse primer (Right_Mut_R) (*Supplementary file 1*) also containing vector backbone sequences. Moreover, to obtain the vector backbone, 1 μl of *Eco*RI (NEB R3101S) and 1 μl of *Hin*dIII (NEB R3104S) were added to an Eppendorf tube containing 2 μg of *pUC19* plasmid at a total volume of 50 μl, and the mixture was incubated at 37°C for 2 h to obtain a linearized *pUC19* vector. To construct the donor plasmid, the two homologous arms with designed mutations and the linearized vector backbone were joined using the Gibson Assembly (Vazyme C113) according to the protocol recommended by the manufacturer. The oligonucleotides for plasmid construction were listed in *Supplementary file 1*. All plasmids were validated by Sanger sequencing.

CRISPR screening of CTCF ADA single-cell clones

The generation of CTCF ADA single-cell clones was performed using CRISPR/Cas9-mediated DNA fragment editing (Li et al., 2015; Shou et al., 2018). Initially, cells are seeded in a 6-cm culture dish and allowed to reach ~80% confluency. The mixture containing 2 μg of the Cas9 plasmid, 1 μg of the sgRNA plasmid, and 1 μg of the donor plasmid, along with 10 μl of transfection reagent P3000 (Invitrogen L3000015) and 100 μl of MEM (Gibco 11095080) was gently mixed with another mixture consisting of 100 μl of MEM (Gibco 11095080) and 10 μl of Lipo3000 (Invitrogen L3000015). Mixtures were then incubated at room temperature for 15 min. Subsequently, the transfection mixture was added to the cells, and the cells were continuously incubated for additional 2 days. The medium was replaced with fresh MEM each day to remove harmful substances and cell debris. Two days after transfection, puromycin (Solarbio P8230-25mg) was added to the cell growth medium at a concentration of 2 μg/ml to initiate the antibiotic selection for 4 days. Finally, the cells are cultured for an additional week in antibiotic-free medium to allow for recovery, before being used for screening single-cell clones.

To screen single-cell clones, the cultured cell mixture was transferred to a new tube and centrifuged at 2,000 × g to pellet the cells. The supernatant was discarded, and the cell pellets were lysed in 30 μl of Solution A (25 mM NaOH) at 95°C for 20 min and neutralized with 30 μl of Solution B (25 mM Tris-HCl pH 6.8). The genomic DNA of the cultured cell mixture was then screened for the targeted mutations by PCR (95°C, 3 min; 95°C, 15 s, 58°C, 15 s, 72°C, 25 s for 35 cycles; and a final extension at 72°C, 5 min) using specific primers with the 3' end of the forward primer matching the designed mutation and a synonymous mutation (geno_F1_Mut and geno_R1) (*Supplementary file 1*). The cell mixture with the targeted mutations was diluted and seeded into eight 96-well culture plates, with approximately one cell per well. The plates were then cultured for a period of 14 days to let the single cell to grow into a colony. The colonies were picked under a microscope and genotyped by PCR (95°C, 3 min; 95°C, 15 s,

58°C, 15 s, 72°C, 25 s for 35 cycles; and a final extension at 72°C, 5 min) using specific primers (primer pairs: geno_F1_Wt and geno_R1 for WT band detection; primer pairs: geno_F1_Mut and geno_R1 for CTCF ADA band detection) (*Supplementary file 1*).

We screened a total of 166 clones and obtained two homozygous CTCF ADA *N2a* clones. The homozygous CTCF ADA single-cell clones were validated by Sanger sequencing. The primer sequences used for CRISPR DNA-fragment editing were listed in *Supplementary file 1*.

Synthesis of the sgRNA and Cas9 mRNA for microinjection by *in vitro* transcription

The generation of CRISPR components was performed as previously described (Ge et al., 2023; Jia et al., 2020). Briefly, the sgRNA targeting sequence was systematically selected using a computational program (Moreno-Mateos et al., 2015). The template for sgRNA *in vitro* transcription was generated by PCR with a pair of primers. The forward primer (T7_Mut-sgRNA-F1 or T7_Mut-sgRNA-F2) (*Supplementary file 1*) includes a T7 promoter sequence (5'-TAATACGACTCACTATA-3'), the sgRNA-target sequence, and the sgRNA scaffold sequence (5'-GTTTTAGAGCTAGAAATAG-3') (Sangon). The common reverse primer (sgRNA-R) matches the sgRNA backbone (*Supplementary file 1*). The total volume of PCR reaction is 50 μl, containing 1 μl of forward primer, 1 μl of common reverse primer, 25 μl of 2 × Phanta Buffer (Vazyme P505-d1), 1 μl of dNTP (Vazyme P505-d1), 1 μl of Phanta (Vazyme P505-d1), and 10 ng of the *pGL3* plasmid as a template, to amplify (95°C, 3 min; 95°C, 20 s, 63°C, 25 s, 72°C, 10 s for 35 cycles; and a final extension at 72°C, 1 min) a linear DNA template for sgRNA *in vitro* transcription.

The PCR product was purified using the PCR purification kit (Qiagen 28104) and eluted in 30 μl of nuclease-free water. The volume of the purified product was brought to 400 μl with RNase-free water (Invitrogen 10977023), followed

by the addition of 400 μl of phenol-chloroform (ACMEC AC13309) to remove RNase. The sample was then vigorously vortexed to mix thoroughly and centrifuged at 17,000 × g for 15 min at room temperature. DNA was precipitated by transferring 320 μl of supernatant to a new RNase-free microcentrifuge tube and adding 32 μl of 3 M sodium acetate (pH 5.5) (Invitrogen AM9740), 2.5 μl of RNA-grade glycogen (Thermo Scientific R0551), 640 μl of ice-cold ethanol. The sample was vortexed and then stored at -80°C for at least 30 min. The precipitated DNA was pelleted by centrifuging at 12,000 × g for 20 min at 4°C and washed with 1 ml of freshly-made 75% ice-cold ethanol. Finally, the washed sample was then centrifuged at 12,000 × g for 5 min at 4°C, and the DNA pellet was resuspended in 30 μl of RNase-free water (Invitrogen 10977023).

The sgRNA *in vitro* transcription was performed using the MEGAshortscript Kit (Invitrogen AM1354) following the protocol recommended by the manufacturer. In brief, 500 ng of purified product was mixed with 2 μl each of ATP, CTP, GTP, and UTP, along with 2 μl of 10 × Buffer and 2 μl of T7 RNA polymerase (Invitrogen AM1354). The sample was mixed gently and incubated at 37°C for 3 h, followed by the addition of 1 μl of TURBO DNase (Invitrogen AM1354). It was then incubated at 37°C for an additional hour. The transcribed product was brought to a total volume of 100 μl with RNase-free water (Invitrogen 10977023) to purify the RNA. The sgRNA was recovered using the MEGAclear Kit (Invitrogen AM1908) according to the protocol recommended by the manufacturer. In brief, the sample was gently mixed with 350 μl of Binding buffer (Invitrogen AM1908), followed by the addition of 250 μl of ethanol and mixing. The sample was then applied to the filter column (Invitrogen AM1908) and left to stand by 5 min, followed by centrifugation at 14,000 × g for 1 min. After discarding the flow-through, the column was washed twice with 500 μl of Washing solution (Invitrogen AM1908) at 14,000 × g for 1 min. After discarding the Washing solution, the column was centrifuged at 14,000 × g for 1 min to completely remove the Washing solution. Finally, the sgRNA was eluted from

the filter with 30 μl of Elution solution (containing 0.2 mM EDTA) at 14,000 × g for 1 min.

The concentration of the purified RNA was measured using NanoDrop 2000 Spectrophotometer. The A260/A280 ratio of high-quality RNA is approximately 2.0, while the A260/A230 ratio is at least 2.2. The sgRNA was then divided into aliquots of 2.5 μg per tube and stored at -80°C to avoid freeze-thaw cycles. The primer sequences used for the sgRNA generation were listed in *Supplementary file 1*.

To synthesis the Cas9 mRNA, 3 μl of *Xba*I (NEB R0145S) was used to digest 6 μg of the *pcDNA3.1-Cas9* vector, which can express Cas9 under control of the *T7* promoter, in a total volume of 100 μl to obtain a linearized vector. The linearized vector was purified through a gel extraction column (Qiagen 28704). The linearized product was eluted in 30 μl of nuclease-free water and brought to 400 μl with RNase-free water, followed by the addition of 400 μl of phenol-chloroform (ACMEC AC13309) to completely remove RNase. The sample was vigorously shake until the solution turns white and centrifuged at 17,000 × g for 15 min at room temperature. After ethanol precipitation, the DNA pellet was resuspended in 15 μl of RNase-free water.

The Cas9 mRNA was transcribed *in vitro* from the purified linearized product using the mMACHINE T7 ULTRA kit (Invitrogen AM1345) according to the manufacturer's instructions. Briefly, 1 μg of *Xba*I-linearized product was mixed with 10 μl of 2 × NTP/ARCA, 2 μl of 10 × Buffer, and 2 μl of T7 RNA polymerase (Invitrogen AM1345) in a total volume of 20 μl. The sample was then incubated at 37°C for 1 h, followed by the addition of 1 μl of TURBO DNase (Invitrogen AM1345) and incubation at 37°C for 1 h. The transcription reaction was added with a poly(A) tailing mixture containing 36 μl of RNase-free water, 20 μl of 5 × *E*-PAP Buffer, 10 μl of 25 mM $MnCl_2$, and 10 μl of ATP solution, and 4 μl of *E*-PAP enzyme (Invitrogen AM1345) at a total volume of 100 μl. The tailing reaction was incubated at 37°C for 45 min. The Cas9 mRNA was also

recovered using the MEGAclear Kit (Invitrogen AM1908). The concentration of the purified RNA was measured using the spectrophotometer (NanoDrop 2000). Finally, the Cas9 mRNA was divided into aliquots of 5 µg per tube and stored at -80°C to avoid freeze-thaw cycles.

Generation of CTCF ADA mutation mice by homologous recombination using CRISPR/Cas9 system via microinjection

C57/BL6J and Institute of Cancer Research (ICR) mice were maintained at 23°C in an SPF (specific pathogen free) facility following a 12-h (7:00–19:00) light and 12-h (19:00–7:00) dark schedule. All animal experimental procedures were approved by the Institutional Animal Care and Use Committee (IACUC) of Shanghai Jiao Tong University (protocol#: A2016041). Briefly, for mice superovulation, the Pregnant mare serum gonadotropin (PMSG) (Solarbio P9970) or human chorionic gonadotropin (hCG) (MCE HY-107953), supplied as a lyophilized powder, is resuspended at a final concentration of 100 U/ml using 0.9% sterile saline (Sigma-Aldrich S3014) solution and then divided into aliquots of 1 ml per tube. The optimal age of mice prepared for superovulation lies between 3 and 6 weeks. Twelve mice were induced to super-ovulate by intraperitoneal injection of 10 U of PMSG on day 1 at 3 pm. On day 3 at 1 pm, 10 U of hCG was administrated. After the injection of hCG, each female mouse is placed in a cage with a stud male for crossing.

Next morning, the females were checked for a copulation plug, and usually 7-10 out of 12 females will be plugged. The unplugged superovulated females can be used for a second time. Typically, about 25 zygotes can be obtained from a 3- to 6-week-old superovulated C57/BL6J, yielding a total of ~200 zygotes for microinjection. The plugged mice were culled by cervical dislocation, followed by dissection of the oviduct and transference of the oviduct into pre-warm M2 medium containing hyaluronidase (Sigma-Aldrich H4272-30MG) at a final concentration of 0.3 mg/ml. The zygotes were pipetted up and down a few

times using a mouth-controlled glass capillary until the cumulus cells fell off. The zygotes were then transferred to a droplet containing no hyaluronidase (Sigma-Aldrich H4272-30MG) to rinse off the hyaluronidase solution and cell debris. The zygotes can be kept at 37°C in a humidified incubator with 5% (v/v) CO_2 until needed. 15-20 zygotes were placed into an M2 medium droplet in a chamber for microinjection.

To prepare a chamber for microinjection, the M2 medium droplets were set up the day before the zygote collection to allow for gas equilibrium at 37°C in a humidified incubator with 5% (v/v) CO_2. Briefly, 40-μl drops of M2 medium were dispensed on the bottom of a 35-mm culture dish. The droplets were flooded with 3 ml of mineral oil (Sigma-Aldrich M5310). In addition, another 35-mm culture dish containing 3 drops of 25-μl each was used as an injection chamber, which was flooded with 2.5 ml of mineral oil (Sigma-Aldrich M5310) and equilibrized at 37°C in a humidified incubator with 5% (v/v) CO_2.

The injection mixture containing 5 μg of Cas9 mRNA, 2.5 μg of sgRNA, and 10 μg of ssODN was mixed in advance at a total volume of 50 μl and was divided into aliquots of 10 μl per tube. The mixture was loaded into the injection capillary using a mouth-controlled glass capillary and injected into the zygote at a very low pressure, delivering 2 picoliters. Once all of the zygotes in the injection chamber were injected, they should be immediately transferred back into a droplet of M2 medium (Sigma-Aldrich MR-015), which is flooded with mineral oil (Sigma-Aldrich M5310), and incubated at 37°C/5% CO_2 incubator for 30 min to recover. Only zygotes exhibiting normal morphology were implanted into the infundibulum of surrogate pseudo-pregnant ICR females.

For zygotes implantation, the surrogate pseudo-pregnant ICR females were weighed and anesthetized by intraperitoneal injection, and placed on the lid of a 10-cm dish. The back of the mouse was wiped with tissues soaked in 75% ethanol. The fur around the dorsal midline was removed using fine scissors. A single incision was made along the dorsal midline, perpendicular to

the vertebral column, just below where the last rib ended. A smaller incision in the body wall was made to expose the reproductive tract. The fat pads were pulled out, and the ovary, oviduct, and uterus came with the fat pads. The microinjected zygotes were then loaded into the transfer pipette. First, a small amount of M2 medium was taken up into the pipette, followed by an air bubble, then M2 medium, and then a second air bubble. The zygotes were drawn up in a minimal volume of M2 medium, followed by an additional small bubble at the tip of the pipette. The infundibulum of the oviduct was gently cradled by blunt forceps, and the prepared transfer pipette was inserted into the infundibulum. The zygotes were blown into the ampulla, and the reproductive tract was put back into the body cavity. Finally, the implanted ICR females were then placed into a clean cage on a warming plate until they recovered from the anesthetic.

To obtain surrogate pseudo-pregnant ICR mice for zygote implantation, the females in estrus were mated with the sterile males the day before implantation. To prepare sterile ICR males, 8-week-old ICR males were chosen to undergo the vasectomy operation. The males were weighed and anesthetized by intraperitoneal injection. A small 15-mm incision was made along the midline of the abdomen using fine scissors. A similar-sized incision in the body wall was made to expose the tissue. The testicular fat pads were pulled out, and the testis, vas deferens, and epididymis came with the fat pads. The vas deferens was held by fine forceps and cauterized with the red-hot tips of another forceps. Next, the testis was carefully placed back inside the body cavity. The same procedures were repeated to cauterize the other vas deferens. The body wall and skin were sewn up using wound clips. The sterile males were then placed into a clean cage on a warming plate until they recovered from the anesthetic. After 14 days of recovery, a female ICR was mated with the sterile male to verify the male's inability to fertilize. If the female was plugged but did not become pregnant, it suggested that the vasectomy operation was successful.

Mouse Genotyping

The chimeric F0 mice were genotyped by PCR and Sanger sequencing using specific primer pairs. In brief, a small piece of the mouse tail was snipped into an Eppendorf tube. The tail was lysed in 30 µl of Solution A (25 mM NaOH) at 95°C for 20 min and neutralized with 30 µl of Solution B (25 mM Tris-HCl pH 6.8). The 2 µl of the neutralized tail lysis solution was used as a template to screen for the targeted mutations by PCR (95°C, 3 min; 95°C, 15 s, 58°C, 15 s, 72°C, 15 s for 40 cycles; and a final extension at 72°C, 5 min) using specific primers (geno_226228_F and geno_226228_R) (*Supplementary file 1*). All of the PCR products were sent for Sanger sequencing to exam the genotype of the chimeric F0 mice. We obtained one chimeric mouse with the precise CTCF ADA mutation and 14 chimeric mice with random indels out of a total of 61 P0 chimeric mice. The targeted chimeric mouse was then crossed with WT mice to obtain heterozygous F1 mice. From a total of 5 litters, we obtained two heterozygous CTCF ADA F1 mice out of a total of only 19 pups. The primer sequences used for mouse genotyping were listed in *Supplementary file 1*.

RNA-seq

RNA-seq was carried out as previously described with minor modifications (Guo et al., 2015). The P0 pups or adult mice were decapitated (according to the guidelines approved by IACUC of Shanghai Jiao Tong University). The skin and skull of the head were carefully removed, and the brain was placed in the 35-mm dish filled with ice-cold PBS. The cortical or olfactory bulb tissues were then dissected out using fine forceps and immediately transferred to a new microcentrifuge tube containing 500 µl of ice-cold TRIzol Reagent (Invitrogen 15596018). The sample was homogenized using a pestle and then centrifuged at 12,000 × g for 10 min at 4°C. The supernatant was transferred into a new microcentrifuge tube, and 100 µl of chloroform was added. The sample was then incubated at room temperature for 3 min after vigorous shaking for 15 sec.

After centrifuging at 12,000 × g for 15 min at 4°C, the upper aqueous phase was carefully transferred into a new RNase-free microcentrifuge tube, followed by the addition of 250 µl of isopropanol. The mixture was inverted several times and incubated at room temperature for 10 min. After centrifuging at 12,000 × g for 10 min at 4°C, the supernatant was discarded, and the RNA pellet was washed once with 1 ml of freshly-made 75% ethanol, air-dried for 5 min, and dissolved in 30 µl of nuclease-free water. The concentration of RNA was measured using the spectrophotometer (NanoDrop 2000). The high-quality RNA, with an A260/A280 ratio of ~2.0, is subjected to library preparation.

The polyadenylated mRNA was isolated using oligo(dT) coupled to magnetic beads (Vazyme N401) from 1 µg of total RNA and fragmented by heating at 94°C for 8 min. The first cDNA strand was synthesized by reverse transcription with a random primer (N6). After synthesizing the second strand DNA, the Y-shaped adapter-HTS was ligated to both ends (*Supplementary file 1*). The ligated DNA was cleaned by AMPure XP Beads (Beckman A63881). Finally, the RNA-seq library was synthesized through PCR (98°C, 30 s; 98°C, 10 s, 60°C, 30 s, 72°C, 30 s for 10 cycles; and a final extension at 72°C, 5 min) by combining the reagents (2.5 µl of P5 primer (*Supplementary file 1*), 2.5 µl of P7 primer (*Supplementary file 1*), and 25 µl of VAHTS HiFi Amplification Mix (Vazyme NR604)). All RNA-seq libraries were sequenced on an Illumina NovaSeq 6000 platform.

ChIP-seq

Chromatin immunoprecipitation was carried out as previously described with minor modifications (Guo et al., 2015). Briefly, cortical tissues were dissected out using fine forceps and immediately transferred to a dish containing ice-cold NeuroBasal medium (Gibco 10888022). The tissues were then torn into pieces using fine forceps and transferred to an Eppendorf tube containing 700 µl of 0.25% trypsin-EDTA. The sample was incubated at 37°C for 15 min with

continuous shaking at 300 rpm. The trypsin digestion of tissues was quenched by the addition of 700 μl of 10% FBS in the PBS buffer. After filtering the dissociated tissues into a single-cell suspension, the volume was adjusted to 5 ml using 10% FBS in the PBS for crosslinking. The single cells were crosslinked by adding 335 μl of 16% formaldehyde (Thermo Scientific 28906) and incubated at room temperature for 10 min. The reaction was then quenched by the addition of 500 μl of 2 M glycine (Invitrogen 15527013). After centrifuging, the cell pellets were washed twice with 1 ml of ice-cold 1 × PBS (containing 1 × protease inhibitors (Roche 04693116001)). The washed cell pellets were then lysed twice with 1 ml of ice-cold Buffer I (10 mM Tris-HCl pH 7.5, 0.1% sodium deoxycholate (Sigma-Aldrich 30970), 0.15 M NaCl (Sigma-Aldrich S3014), 0.1% SDS (Sigma-Aldrich 71736), 1% Triton X-100 (Sigma-Aldrich T8787), 1 mM EDTA (Sigma-Aldrich E1644), and 1 × protease inhibitors (Roche 04693116001)) at 4°C for 10 min with continuous rotations at 9 rpm. The lysed cells were sonicated with 33% power at a train of 30-sec sonication with 30-sec interval for 30 cycles using the Sonics Vibra-Cell to obtain DNA fragments ranging from 200 to 500 bp. The insoluble debris was discarded by centrifuging at 14,000 × g for 10 min at 4°C. The supernatant cell lysate was initially pre-cleared with 50 μl of protein A/G-agarose beads (Millipore 16-157) for 2 h at 4°C with gentle rotations. The sample was centrifuged at 2,000 × g for 1 min, and the supernatant was then transferred to a new microcentrifuge tube. Finally, an antibody specifically against CTCF (Millipore 07-729) or RAD21 (Abcam ab992) (*Supplementary file 2*) was added to the supernatant for immunoprecipitation at 4°C overnight with slow rotations.

On the following day, the antibody-protein-DNA complexes were purified with 50 μl of protein A/G agarose beads by incubating at 4°C for 3 h with slow rotations. The beads were washed once with 1 ml of ice-cold Buffer I, once with 1 ml of ice-cold Buffer I with 0.4 M NaCl, thrice with ice-cold Buffer I without NaCl, and finally with 1 ml of ice-cold LiCl buffer (0.5 M LiCl (Sigma-Aldrich

L4408), 50 mM Tris-HCl pH 7.5, 1 mM EDTA, 1% NP-40 (Sigma-Aldrich I8896), 0.7% sodium deoxycholate), by incubating at 4°C for 10 min with slow rotations, respectively. The crosslinked complexes were eluted from the beads using 400 μl of elution buffer (50 mM Tris-HCl pH 8.0, 10 mM EDTA, 1% SDS) at 65°C for 1 h while shaking at 1,000 rpm, followed by reverse crosslinking by incubating at 65°C overnight while shaking at 1,000 rpm. After the addition of 400 μl of phenol-chloroform, the DNA was extracted by vigorous vortexing and centrifuging at 17,000 × g for 15 min at room temperature. After ethanol precipitation, the DNA pellet was resuspended in 50 μl of 10 mM Tris-HCl pH 7.5. The concentration of the DNA was measured using Qubit fluorescent dye (Vazyme EQ121-01). The purified DNA was subjected to high-throughput library preparation following the method described above in the RNA-seq section. All ChIP-seq libraries were sequenced on an Illumina NovaSeq 6000 platform.

In situ Hi-C

In situ Hi-C experiments were carried out following the previous protocol with minor modifications (Guo et al., 2015; Zhang et al., 2024). Briefly, after single-cell isolation from the cortical tissues, one million cells were cross-linked with 1% formaldehyde (Thermo Scientific 28906) at room temperature for 10 min with slow rotations. The crosslinked cells were washed twice with 1 × PBS (Gibco 70011044) and lysed twice with 250 μl of ice-cold lysis buffer (10 mM Tris-HCl pH 8.0, 10 mM NaCl, 0.2% NP-40) for 10 min at 4°C to obtain nuclei. Subsequently, the nuclei were permeabilized in 50 μl of 0.05% SDS at 62°C for 10 min, then quenched by the addition of Triton X-100, and digested with 100 U of *Mbo*I (NEB R0147L) in a total volume of 250 μl overnight at 37°C with shaking at 900 rpm. After heat inactivating *Mbo*I at 62°C for 20 min, the digested DNA ends were filled in and labeled by adding a mixture that included DNA Polymerase I, Large (Klenow) Fragment (NEB M0210S), biotin-14-dATP (Thermo 19524016), as well as dCTP, dTTP, and dGTP (NEB N0446S). The

reaction was then incubated at 37°C for 1 h.

After labeling the ends of restriction fragments, the digested nuclei were proximally ligated with 2 μl of T4 DNA ligase (NEB M0202S) at room temperature for 4 h. The ligated chromatin was digested using 4 μl of 20 mg/ml proteinase K (Invitrogen AM2546) at 55°C for 4 h and then reverse crosslinked overnight at 68 °C overnight. The DNA was purified through phenol-chloroform extraction followed by ethanol precipitation as described above. After precipitation, the DNA was dissolved in 130 μl of 10 mM Tris-HCl pH 8.0. Sonication of the DNA was carried out using a Bioruptor system at high intensity levels. The process consisted of 15 cycles, each comprising 30-sec sonication followed by 30-sec interval. DNA fragments of 300–500 bp were selected using AMPure XP beads (Beckman A63881) and subsequently captured using streptavidin beads (Thermo 11206D) to enrich the biotin-labeled DNA fragments. The biotin-DNA on the beads was stringently washed twice with 2-min of vortices at 55°C. The purified DNA was used as a template for library preparation as described above. The library was then subjected to sequencing on the MGI-T7 platform via the PE150 sequencing strategy.

QHR-4C

Quantitative high-resolution chromosome conformation capture copy (QHR-4C) experiments were performed as previously described (Jia et al., 2020) with minor modifications. In brief, five million cells were harvested as described above. The cells were resuspended in 10 ml of MEM medium, and then crosslinked with 2% formaldehyde (Thermo Scientific 28906) at room temperature with slow rotations for 10 min. The reaction was quenched by the addition of 700 μl of 2 M glycine (Invitrogen 15527013). The fixed cells were spun down at 900 × g for 5 min at 4°C and then washed twice with 1 ml of ice-cold PBS (Gibco 70011044). Cells were lysed twice with 1 ml of ice-cold 4C permeabilization buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 5 mM EDTA,

0.2% SDS, 0.5% NP40, 1% Triton X-100, and 1 × protease inhibitors) with slow rotations for 10 min at 4°C. Cells were then resuspended in 73 μl of nuclease-free water, followed by the addition of 10 μl of 10 × *Dpn*II buffer (NEB R0543L) and 2.5 μl of 10% SDS, and incubation at 37°C with constant shaking at 900 rpm. After 1 h, 12.5 μl of 20% Triton X-100 (Sigma-Aldrich T8787) was added to the sample, and the sample was incubated at 37°C for an additional hour with constant shaking at 900 rpm. Next, 8 μl of *Dpn*II (NEB R0543L) was added to the sample, followed by an overnight incubation at 37°C with constant shaking at 900 rpm. The enzyme was heat-inactivated at 65°C for 20 min. The sample was centrifuged at 2,000 × g for 10 min, and the supernatant was carefully removed to not disturb the cell pellets. The cell pellets were then resuspended with 1,600 μl of nuclease-free water. The digested nuclei were ligated with 2 μl of T4 DNA ligase (NEB M0202S) in a total ligation volume of 2 ml at 16°C for 12 h.

The next day, the ligated chromatin was digested with 8 μl of 10 mg/ml Protease K (Invitrogen AM2546) at 65°C for 2 h. The DNA was extracted using phenol-chloroform extraction, followed by ethanol precipitation as described above. The DNA pellet was then dissolved in 50 μl of nuclease-free water and sonicated using a Bioruptor system with a low energy setting. Sonication was performed in cycles of 30-sec on and 30-sec off for 12 cycles to obtain DNA fragments ranging from 200-600 bp. The concentration of the sonicated DNA was measured using the Qubit fluorescent dye (Vazyme EQ121-01). Next, 5 μg of DNA was used as a template for linear amplification with a specific 5' biotin-tagged primer complementary to the viewpoint. The linear amplification was conducted in a total volume of 100 μl, including 50 μl of 2 × Buffer, 0.8 μl of dNTP, 1.6 μl of Phanta (Vazyme P505-d1), and 1 μl of 1 μM 5' biotin-tagged primer (*Supplementary file 1*), using the program (95°C, 2 min; 95°C, 15 s, 58°C, 25 s, 72°C, 1 min 30 s for 80 cycles; and a final extension at 72°C, 5 min). The amplified products were denatured at 95°C for 5 min and then chilled on ice to

obtain biotinylated single-stranded DNA (ssDNA). The biotinylated ssDNA was ligated with 2 μl of 50 μM Adapter-4C (*Supplementary file 1*) adapters in a total volume of 45 μl for 12 h at 16°C and purified by streptavidin magnetic C1 beads (Invitrogen 65001). The QHR-4C libraries were generated via PCR amplification (94°C, 2 min; 94°C, 10 s, 60°C, 15 s, 72°C, 1 min for 19 cycles; and a final extension at 72°C, 5 min) using Phanta high-fidelity DNA polymerase (Vazyme P505-d1). The amplified PCR products were purified using the High-Pure PCR Product Purification kit (Roche 11732676001) and sequenced on an Illumina NovaSeq 6000 platform. The primers for QHR-4C are listed in *Supplementary file 1*.

Bioinformatics analyses

RNA-seq analysis

Raw FASTQ data files were mapped to the mouse reference genome (GRCm38/mm10) using the HISAT2 (v2.2.1) (Kim et al., 2019) with default parameters. Subsequently, Cufflinks (v2.2.1) (Trapnell et al., 2010) was employed to estimate transcript expression levels in FPKM (Fragments Per Kilobase of exon per Million mapped fragments). Gene expression analyses were conducted with a minimum of three replicates, presenting data as mean ± SEM. Statistical significance of gene expression changes was determined using unpaired Student's *t*-test in GraphPad Prism (v8.4.2) and R (v4.1.1), with significance levels denoted as follows: '*' for $P \leq 0.05$, '**' for $P \leq 0.01$, and '***' for $P \leq 0.001$.

For the genome-wide analyses of differentially expressed genes (DEGs), featureCounts (v2.0.1) (Liao et al., 2014) was used to generate the raw count matrix. The count matrix was then assayed using the R package DESeq2 (Love et al., 2014) to detect DEGs, employing parameters of absolute $\log_2 FC > 0.5$ and *padj* < 0.05. Subsequently, a volcano plot depicting DEGs was generated using the R package ggplot2 (v3.5.0) (Ginestet, 2011).

Manhattan plots illustrating the localized enrichments of downregulated genes were produced using the R package CMplot (v4.5.1) (Jiang et al., 2017; Yin et al., 2021). In brief, the mouse reference genome (GRCm38/mm10) was partitioned into non-overlapping 1-Mb bins. Subsequently, the genomic locations of the downregulated genes were intersected with these 1-Mb bins to determine the frequency of hits in each bin. The occurrence frequency of each signal within 1-Mb bins was then modeled using a Poisson distribution. The maximum-likelihood estimator for the λ was determined based on the mean number of occurrences. Subsequently, the Poisson models were employed to compute the probability of observing each signal within every 1-Mb bin. The probabilities were documented within each bin and subjected to CMplot (v4.5.1) (Yin et al., 2021) to generate Manhattan plots.

Scatterplots showing fold changes of expression levels of *cPcdh* genes as a function of the linear genomic distance between promoters and enhancers were generated using ggplot2 (v3.5.0) (Ginestet, 2011; Kiefer et al., 2023). Briefly, the fold change for each *cPcdh* gene was calculated based on the RNA-seq data. The enhancer element associated with the *Pcdh* α, β, and γ clusters are *HS5-1*, *HS18-22*, and *HS5-1bL*, respectively. The 5' untranslated regions (UTRs) of each gene were served as anchor points for calculating the distances between the genes and enhancers. Subsequently, the fold changes for each gene were paired with their respective distances from the enhancers based on gene names. These data were used to create the scatterplots by utilizing ggplot2 (v3.5.0) (Ginestet, 2011). A smooth curve is added to the scatterplot to facilitate the observation of trends in the data using ggplot2 (v3.5.0) (Ginestet, 2011).

Bar plots displaying the proportions of mutated bases in high-throughput RNA-seq reads were generated using ggplot2 (v3.5.0) (Ginestet, 2011). The counts of mutated bases were visualized and summarized using the Integrative Genomics Viewer (IGV) (Thorvaldsdóttir et al., 2013). The ratio was computed

manually and then plotted using ggplot2 (v3.5.0) (Ginestet, 2011) to create the bar graphs.

ChIP-seq analysis

The raw sequencing data were mapped to the mouse reference genome (GRCm38/mm10) using Bowtie2 (v2.2.5) (Langmead and Salzberg, 2012) with default settings. Samtools (v1.15.1) (Li et al., 2009) was utilized to convert the Sequence Alignment/Map (SAM) format into a Binary Alignment/Map (BAM) format and to index the resulting BAM files. Subsequently, the BAM files were normalized to RPKM (Reads Per Kilobase per Million mapped reads) for comparative analysis of occupancy levels across samples, utilizing the bamCoverage module of the deepTools suite (Ramirez et al., 2016). The bedGraph files, generated from the deepTools program (Ramirez et al., 2016), were then uploaded to the UCSC Genome Browser (Kent et al., 2002) for visualization of genomic regions of interest. Peak calling was conducted using MACS2 (v2.2.7.1) (Zhang et al., 2008) with default parameters.

Boxplots illustrating the ChIP-seq profiles of RAD21 in the *cPcdh* locus were generated using ggplot2 (v3.5.0) (Ginestet, 2011). Statistical significance of changes of RAD21 enrichment was computed using the R package rstatix (v0.7.0), with significance levels denoted as following: '*' for $P \leq 0.05$, '**' for $P \leq 0.01$, and '***' for $P \leq 0.001$.

Scatterplots showing fold changes of RAD21 enrichments at the *cPcdh* genes as a function of the linear genomic distance between promoters and enhancers were created using ggplot2 (v3.5.0) (Ginestet, 2011). Briefly, RAD21 enrichments for each *cPcdh* gene were calculated based on the ChIP-seq data by calculating the FPKM of all peaks. The fold changes in RAD21 enrichments in heterozygous mice were then computed by comparing with the WT littermates. Distances between genes and enhancers were calculated using the 5' untranslated regions (UTRs) of each gene as anchor points. The fold

changes in Rad21 enrichments for each gene were paired with their corresponding distances from the enhancers based on gene names.

Hi-C analysis

Raw FASTQ reads were mapped to the mouse reference genome assembly (GRCm38/mm10) using the HiC-Pro pipeline (Servant et al., 2015) (*Supplementary file 3*). The resulting allValidPairs files were converted to the hic format files using the hicpro2juicebox script from the HiC-Pro toolkit, enabling analysis with Juicer tools (Durand et al., 2016). Subsequently, the hic format was converted to cooler format using hic2cool (v0.8.3). To ensure accuracy, the contact matrix was normalized to a contact depth of 100 million based on *cis*-interactions using the Knight-Ruiz (KR) method.

The Hi-C contact maps for targeted genomic regions were generated using the fancplot module in FAN-C toolkits (Kruse et al., 2020), and the differential contact heatmaps were constructed using Seaborn, a Python library. At a 100-Kb resolution, the Hi-C *cis*-Eigenvector 1 values and Pearson's correlation matrix were calculated utilizing the Juicer software (Durand et al., 2016). Subsequently, the resulting *cis*-Eigenvector 1 values were converted into the bedGraph format for visualization in the UCSC genome browser (Kent et al., 2002). Scatterplots with a heat density were depicted using the R package ggpointdensity (v0.1.0).

TADs were called using the Hidden Markov Models (HMM) approach, as reported in a previous study (Dixon et al., 2012). The dense matrix was subjected to DI_from_matrix script to calculate the Directionality Index (DI) score. The DI score was then utilized to predict the hidden states of all bins via HMM models. Subsequently, bins with consecutive identical states were merged into regions. Regions containing less than 3 bins or with a median posterior bin probability below 0.99 were excluded from the analysis using a Perl script. Ultimately, TADs were defined as regions extending downstream

from an upstream boundary and upstream from a downstream boundary. The DI score was transformed into the bedGraph format to visualize in the UCSC genome browser.

To identify loops in the contact matrix, Chromosight (v1.6.3) (Matthey-Doret et al., 2020) was employed to detect chromatin loops with a genomic distance greater than 4 Kb. Subsequently, specific loops were analyzed using the Aggregate Peak Analysis (APA) with cooltools (v0.6.1) (Open2C et al., 2022) and coolpup.py (v1.1.0) (Flyamer et al., 2020).

QHR-4C analysis

Raw FASTQ files were mapped to the mouse reference genome (GRCm38/mm10) using Bowtie2 (v2.2.5) (Langmead and Salzberg, 2012) with default parameters. The reads per million (RPM) values were calculated using r3Cseq (Thongjuea et al., 2013) program (v1.20). Bedgraph files generated by r3Cseq (v1.20) were visualized in the UCSC genome browser.

Scatterplots depicting fold changes of contact probabilities of *cPcdh* genes as a function of the linear genomic distance between target variable promoters and their respective enhancers were created using ggplot2 (v3.5.0). Interaction values were assigned to the variable exons of each member of the *cPcdh* genes, and the fold changes in contact probabilities upon CTCF ADA mutation were quantified. Gene-enhancer distances were computed utilizing the 5' untranslated regions (UTRs) of each gene as reference points. The fold change data for each gene were matched with distances from the enhancers based on gene names. These datasets were employed in ggplot2 (v3.5.0) to produce scatterplots, with a smoothing curve included to facilitate the visualization of data trends.

Acknowledgements

We would like to express our gratitude to all members of our group for their insightful discussions. This work was supported by grants to Q.W. from the National Natural Science Foundation of China (32330016) and the National Key R&D Program of China (2022YFC3400200).

Author Contributions

Q.W. conceived and supervised the project. Y.Z. performed experiments and analyzed data. Y.Z. and Q.W. wrote the manuscript.

References

Canzio, D., Nwakeze, C. L., Horta, A., Rajkumar, S. M., Coffey, E. L., Duffy, E. E., Duffié, R., Monahan, K., O'Keeffe, S., Simon, M. D., et al. 2019. Antisense lncRNA Transcription Mediates DNA Demethylation to Drive Stochastic Protocadherin α Promoter Choice. **Cell** 177:639-653. DOI: https://doi.org/10.1016/j.cell.2019.03.008, PMID: 30955885

Chen, F., Yuan, H. M., Wu, W. Y., Chen, S. K., Yang, Q., Wang, J., Zhang, Q., Gui, B. H., Fan, X., Chen, R. M., et al. 2019. Three additional de novo mutations in Chinese patients help to define an emerging neurodevelopmental disorder. **American Journal of Medical Genetics Part C-Seminars in Medical Genetics** 181:218-225. DOI: https://doi.org/10.1002/ajmg.c.31698, PMID: 30893510

Chen, W. V., Nwakeze, C. L., Denny, C. A., O'Keeffe, S., Rieger, M. A., Mountoufaris, G., Kirner, A., Dougherty, J. D., Hen, R., Wu, Q., et al. 2017. Pcdhαc2 is required for axonal tiling and assembly of serotonergic circuitries in mice. **Science** 356:406-411. DOI: https://doi.org/10.1126/science.aal3231, PMID: 28450636

Dixon, J. R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J. S., Ren, B. 2012. Topological domains in mammalian genomes identified by analysis of chromatin interactions. **Nature** 485:376-380. DOI: https://doi.org/10.1038/nature11082, PMID: 22495300

Dong, H., Li, J., Wu, Q., Jin, Y. 2023. Confluence and convergence of Dscam and Pcdh cell-recognition codes. **Trends in Biochemical Sciences** 48:1044-1057. DOI: https://doi.org/10.1016/j.tibs.2023.09.001, PMID: 37839971

Durand, N. C., Shamim, M. S., Machol, I., Rao, S. S., Huntley, M. H., Lander, E. S., Aiden, E. L. 2016. Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. **Cell Systems** 3:95-98. DOI: https://doi.org/10.1016/j.cels.2016.07.002, PMID: 27467249

Esumi, S., Kakazu, N., Taguchi, Y., Hirayama, T., Sasaki, A., Hirabayashi, T., Koide, T., Kitsukawa, T., Hamada, S., Yagi, T. 2005. Monoallelic yet combinatorial expression of variable exons of the protocadherin-α gene cluster in single neurons. **Nature Genetics** 37:171-176. DOI: https://doi.org/10.1038/ng1500, PMID: 15640798

Fan, L., Lu, Y., Shen, X., Shao, H., Suo, L., Wu, Q. 2018. Alpha protocadherins and Pyk2 kinase regulate cortical neuron migration and cytoskeletal dynamics via Rac1 GTPase and WAVE complex in mice. **eLife** 7:e35242. DOI: https://doi.org/10.7554/eLife.35242, PMID: 29911975

Fedoriw, A. M., Stein, P., Svoboda, P., Schultz, R. M., Bartolomei, M. S. 2004. Transgenic RNAi reveals essential function for CTCF in H19 gene imprinting. **Science** 303:238-240. DOI: https://doi.org/10.1126/science.1090934, PMID: 14716017

Flaherty, E., Maniatis, T. 2020. The role of clustered protocadherins in neurodevelopment and neuropsychiatric diseases. **Current Opinion in Genetics & Development** 65:144-150. DOI: https://doi.org/10.1016/j.gde.2020.05.041, PMID: 32679536

Flyamer, I. M., Illingworth, R. S., Bickmore, W. A. 2020. Coolpup.py: versatile pile-up analysis of Hi-C data. **Bioinformatics** 36:2980-2985. DOI: https://doi.org/10.1093/bioinformatics/btaa073, PMID: 32003791

Ge, X., Huang, H. Y., Han, K. Q., Xu, W. J., Wang, Z. X., Wu, Q. 2023. Outward-oriented sites within clustered CTCF boundaries are key for intra-TAD chromatin interactions and gene regulation. **Nature Communications** 14:8101. DOI: https://doi.org/10.1038/s41467-023-43849-0, PMID: 38062010

Ginestet, C. 2011. ggplot2: Elegant Graphics for Data Analysis. **Journal of the Royal Statistical Society Series a-Statistics in Society** 174:245-245. DOI: https://doi.org/10.1111/j.1467-985X.2010.00676_9.x, PMID:

Guo, Y., Monahan, K., Wu, H. Y., Gertz, J., Varley, K. E., Li, W., Myers, R. M., Maniatis, T., Wu, Q. 2012. CTCF/cohesin-mediated DNA looping is required for protocadherin α promoter choice. **PNAS** 109:21081-21086. DOI: https://doi.org/10.1073/pnas.1219280110, PMID: 23204437

Guo, Y., Xu, Q., Canzio, D., Shou, J., Li, J. H., Gorkin, D. U., Jung, I., Wu, H. Y., Zhai, Y. N., Tang, Y. X., et al. 2015. CRISPR Inversion of CTCF Sites Alters Genome Topology and Enhancer/Promoter Function. **Cell** 162:900-910. DOI: https://doi.org/10.1016/j.cell.2015.07.038, PMID: 26276636

Hashimoto, H., Wang, D., Horton, J. R., Zhang, X., Corces, V. G., Cheng, X. 2017. Structural basis for the versatile and methylation-dependent binding of CTCF to DNA. **Molecular Cell** 66:711-720. DOI: https://doi.org/10.1016/j.molcel.2017.05.004, PMID: 28529057

Hirayama, T., Tarusawa, E., Yoshimura, Y., Galjart, N., Yagi, T. 2012. CTCF is required for neural development and stochastic expression of clustered Pcdh genes in neurons. **Cell Reports** 2:345-357. DOI: https://doi.org/10.1016/j.celrep.2012.06.014, PMID: 22854024

Ing-Esteves, S., Kostadinov, D., Marocha, J., Sing, A. D., Joseph, K. S., Laboulaye, M. A., Sanes, J. R., Lefebvre, J. L. 2018. Combinatorial effects of alpha- and gamma-protocadherins on neuronal survival and dendritic self-avoidance. **The Journal of Neuroscience** 38:2713-2729. DOI: https://doi.org/10.1523/JNEUROSCI.3035-17.2018, PMID: 29439167

Jia, Z. L., Li, J. W., Ge, X., Wu, Y. H., Guo, Y., Wu, Q. 2020. Tandem CTCF sites function as insulators to balance spatial chromatin contacts and topological enhancer-promoter selection. **Genome Biology** 21:75. DOI: https://doi.org/10.1186/s13059-020-01984-7, PMID: 32293525

Jia, Z. L., Wu, Q. 2020. Clustered Protocadherins Emerge as Novel Susceptibility Loci for Mental Disorders. **Frontiers in Neuroscience** 14:587819. DOI:

https://doi.org/10.3389/fnins.2020.587819, PMID: 33262685

Jiang, Y., Loh, Y. E., Rajarajan, P., Hirayama, T., Liao, W., Kassim, B. S., Javidfar, B., Hartley, B. J., Kleofas, L., Park, R. B., et al. 2017. The methyltransferase SETDB1 regulates a large neuron-specific topological chromatin domain. **Nature Genetics** 49:1239-1250. DOI: https://doi.org/10.1038/ng.3906, PMID: 28671686

Kawauchi, S., Calof, A. L., Santos, R., Lopez-Burks, M. E., Young, C. M., Hoang, M. P., Chua, A., Lao, T. T., Lechner, M. S., Daniel, J. A., et al. 2009. Multiple Organ System Defects and Transcriptional Dysregulation in the Mouse, a Model of Cornelia de Lange Syndrome. **PLoS Genetics** 5:e1000650. DOI: https://doi.org/10.1371/journal.pgen.1000650, PMID: 19763162

Kehayova, P., Monahan, K., Chen, W. S., Maniatis, T. 2011. Regulatory elements required for the activation and repression of the protocadherin-α gene cluster. **PNAS** 108:17195-17200. DOI: https://doi.org/10.1073/pnas.1114357108, PMID: 21949399

Kent, W. J., Sugnet, C. W., Furey, T. S., Roskin, K. M., Pringle, T. H., Zahler, A. M., Haussler, D. 2002. The human genome browser at UCSC. **Genome Research** 12:996-1006. DOI: https://doi.org/10.1101/gr.229102, PMID: 12045153

Kiefer, L., Chiosso, A., Langen, J., Buckley, A., Gaudin, S., Rajkumar, S. M., Servito, G. I. F., Cha, E. S., Vijay, A., Yeung, A., et al. 2023. WAPL functions as a rheostat of Protocadherin isoform diversity that controls neural wiring. **Science** 380:eadf8440. DOI: https://doi.org/10.1126/science.adf8440, PMID: 37347873

Kim, D., Paggi, J. M., Park, C., Bennett, C., Salzberg, S. L. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. **Nature Biotechnology** 37:907-915. DOI: https://doi.org/10.1038/s41587-019-0201-4, PMID: 31375807

Kruse, K., Hug, C. B., Vaquerizas, J. M. 2020. FAN-C: a feature-rich framework for the analysis and visualisation of chromosome conformation capture data. **Genome Biology** 21:303. DOI: https://doi.org/10.1186/s13059-020-02215-9, PMID: 33334380

Langmead, B., Salzberg, S. L. 2012. Fast gapped-read alignment with Bowtie 2. **Nature Methods** 9:357-359. DOI: https://doi.org/10.1038/nmeth.1923, PMID: 22388286

Lefebvre, J. L., Kostadinov, D., Chen, W. V., Maniatis, T., Sanes, J. R. 2012. Protocadherins mediate dendritic self-avoidance in the mammalian nervous system. **Nature** 488:517-521. DOI: https://doi.org/10.1038/nature11305, PMID: 22842903

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., Genome Project Data Processing, S. 2009. The Sequence Alignment/Map format and SAMtools. **Bioinformatics** 25:2078-2079. DOI: https://doi.org/10.1093/bioinformatics/btp352, PMID: 19505943

Li, J. H., Shou, J., Guo, Y., Tang, Y. X., Wu, Y. H., Jia, Z. L., Zhai, Y. N., Chen, Z. F., Xu, Q., Wu, Q. 2015. Efficient inversions and duplications of mammalian regulatory DNA elements and gene clusters by CRISPR/Cas9. **Journal of Molecular Cell Biology** 7:284-298. DOI: https://doi.org/10.1093/jmcb/mjv016, PMID: 25757625

Li, Y., Haarhuis, J. H. I., Cacciatore, A. S., Oldenkamp, R., van Ruiten, M. S., Willems, L., Teunissen, H., Muir, K. W., de Wit, E., Rowland, B. D., et al. 2020. The structural basis for cohesin-CTCF-anchored loops. **Nature** 578:472-476. DOI: https://doi.org/10.1038/s41586-019-1910-z, PMID: 31905366

Liao, Y., Smyth, G. K., Shi, W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. **Bioinformatics** 30:923-930. DOI:

https://doi.org/10.1093/bioinformatics/btt656, PMID: 24227677

Love, M. I., Huber, W., Anders, S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. **Genome Biology** 15:550. DOI: https://doi.org/10.1186/s13059-014-0550-8, PMID: 25516281

Lu, Y. J., Shou, J., Jia, Z. L., Wu, Y. H., Li, J. H., Guo, Y., Wu, Q. 2019. Genetic evidence for asymmetric blocking of higher-order chromatin structure by CTCF/cohesin. **Protein & Cell** 10:914-920. DOI: https://doi.org/10.1007/s13238-019-00656-y, PMID: 31559572

Lv, X. H., Li, S., Li, J. W., Yu, X. Y., Ge, X., Li, B., Hu, S. H., Lin, Y., Zhang, S. B., Yang, J. J., et al. 2022. Patterned cPCDH expression regulates the fine organization of the neocortex. **Nature** 612:503-511. DOI: https://doi.org/10.1038/s41586-022-05495-2, PMID: 36477535

Mannini, L., Cucco, F., Quarantotti, V., Krantz, I. D., Musio, A. 2013. Mutation spectrum and genotype-phenotype correlation in Cornelia de Lange syndrome. **Human Mutation** 34:1589-1596. DOI: https://doi.org/10.1002/humu.22430, PMID: 24038889

Matthey-Doret, C., Baudry, L., Breuer, A., Montagne, R., Guiglielmoni, N., Scolari, V., Jean, E., Campeas, A., Chanut, P. H., Oriol, E., et al. 2020. Computer vision for pattern detection in chromosome contact maps. **Nature Communications** 11:5795. DOI: https://doi.org/10.1038/s41467-020-19562-7, PMID: 33199682

Monahan, K., Rudnick, N. D., Kehayova, P. D., Pauli, F., Newberry, K. M., Myers, R. M., Maniatis, T. 2012. Role of CCCTC binding factor (CTCF) and cohesin in the generation of single-cell diversity of protocadherin-alpha gene expression. **PNAS** 109:9125-9130. DOI: https://doi.org/10.1073/pnas.1205074109, PMID: 22550178

Moore, J. M., Rabaia, N. A., Smith, L. E., Fagerlie, S., Gurley, K., Loukinov, D., Disteche, C. M., Collins, S. J., Kemp, C. J., Lobanenkov, V. V., et al. 2012. Loss of maternal CTCF is associated with peri-implantation lethality of Ctcf null embryos. **PLoS One** 7:e34915. DOI: https://doi.org/10.1371/journal.pone.0034915, PMID: 22532833

Moreno-Mateos, M. A., Vejnar, C. E., Beaudoin, J.-D., Fernandez, J. P., Mis, E. K., Khokha, M. K., Giraldez, A. J. 2015. CRISPRscan: designing highly efficient sgRNAs for CRISPR-Cas9 targeting in vivo. **Nature Methods** 12:982-988. DOI: https://doi.org/10.1038/nmeth.3543, PMID: 26322839

Mountoufaris, G., Canzio, D., Nwakeze, C. L., Chen, W. V., Maniatis, T. 2018. Writing, Reading, and Translating the Clustered Protocadherin Cell Surface Recognition Code for Neural Circuit Assembly. **Annual Review of Cell and Developmental Biology** 34:471-493. DOI: https://doi.org/10.1146/annurev-cellbio-100616-060701, PMID: 30296392

Mountoufaris, G., Chen, W. S. V., Hirabayashi, Y., O'Keeffe, S., Chevee, M., Nwakeze, C. L., Polleux, F., Maniatis, T. 2017. Multicluster Pcdh diversity is required for mouse olfactory neural circuit assembly. **Science** 356:411-413. DOI: https://doi.org/10.1126/science.aai8801, PMID: 28450637

Nasmyth, K., Haering, C. H. 2009. Cohesin: its roles and mechanisms. **Annual Review of Genetics** 43:525-558. DOI: https://doi.org/10.1146/annurev-genet-102108-134233, PMID: 19886810

Nasmyth, K. A., Lee, B. G., Roig, M. B., Löwe, J. 2023. What AlphaFold tells us about cohesin's retention on and release from chromosomes. **Elife** 12:RP88656. DOI: https://doi.org/10.7554/eLife.88656, PMID: 37975572

Nora, E. P., Caccianini, L., Fudenberg, G., So, K., Kameswaran, V., Nagle, A., Uebersohn, A.,

Hajj, B., Le Saux, A., Coulon, A., et al. 2020. Molecular basis of CTCF binding polarity in genome folding. **Nature Communications** 11:5612. DOI: https://doi.org/10.1038/s41467-020-19283-x, PMID: 33154377

Open2C, Abdennur, N., Abraham, S., Fudenberg, G., Flyamer, I. M., Galitsyna, A. A., Goloborodko, A., Imakaev, M., Oksuz, B. A., Venev, S. V. 2022. Cooltools: enabling high-resolution Hi-C analysis in Python. **bioRxiv**. DOI: https://doi.org/10.1101/2022.10.31.514564, PMID:

Price, E., Fedida, L. M., Pugacheva, E. M., Ji, Y. J., Loukinov, D., Lobanenkov, V. V. 2023. An updated catalog of CTCF variants associated with neurodevelopmental disorder phenotypes. **Frontiers in Molecular Neuroscience** 16:1185796. DOI: https://doi.org/10.3389/fnmol.2023.1185796, PMID: 37324587

Pugacheva, E. M., Kubo, N., Loukinov, D., Tajmul, M., Kang, S., Kovalchuk, A. L., Strunnikov, A. V., Zentner, G. E., Ren, B., Lobanenkov, V. V. 2020. CTCF mediates chromatin looping via N-terminal domain-dependent cohesin retention. **PNAS** 117:2020-2031. DOI: https://doi.org/10.1073/pnas.1911708117, PMID: 31937660

Ramirez, F., Ryan, D. P., Gruning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., Heyne, S., Dundar, F., Manke, T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis. **Nucleic Acids Research** 44:W160-W165. DOI: https://doi.org/10.1093/nar/gkw257, PMID: 27079975

Remeseiro, S., Cuadrado, A., Gomez-Lopez, G., Pisano, D. G., Losada, A. 2012. A unique role of cohesin-SA1 in gene regulation and development. **The EMBO Journal** 31:2090-2102. DOI: https://doi.org/10.1038/emboj.2012.60, PMID: 22415368

Remeseiro, S., Cuadrado, A., Kawauchi, S., Calof, A. L., Lander, A. D., Losada, A. 2013. Reduction of Nipbl impairs cohesin loading locally and affects transcription but not cohesion-dependent functions in a mouse model of Cornelia de Lange Syndrome. **Biochimica et Biophysica Acta** 1832:2097-2102. DOI: https://doi.org/10.1016/j.bbadis.2013.07.020, PMID: 23920377

Ribich, S., Tasic, B., Maniatis, T. 2006. Identification of long-range regulatory elements in the protocadherin-α gene cluster. **PNAS** 103:19719-19724. DOI: https://doi.org/10.1073/pnas.0609445104, PMID: 17172445

Sanes, J. R., Zipursky, S. L. 2020. Synaptic Specificity, Recognition Molecules, and Assembly of Neural Circuits. **Cell** 181:536-556. DOI: https://doi.org/10.1016/j.cell.2020.04.008, PMID: 32359437

Schreiner, D., Weiner, J. A. 2010. Combinatorial homophilic interaction between gamma-protocadherin multimers greatly expands the molecular diversity of cell adhesion. **PNAS** 107:14893-14898. DOI: https://doi.org/10.1073/pnas.1004526107, PMID: 20679223

Servant, N., Varoquaux, N., Lajoie, B. R., Viara, E., Chen, C. J., Vert, J. P., Heard, E., Dekker, J., Barillot, E. 2015. HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. **Genome Biology** 16:259. DOI: https://doi.org/10.1186/s13059-015-0831-x, PMID: 26619908

Shou, J., Li, J. H., Liu, Y. B., Wu, Q. 2018. Precise and Predictable CRISPR Chromosomal Rearrangements Reveal Principles of Cas9-Mediated Nucleotide Insertion. **Molecular Cell** 71:498-509. DOI: https://doi.org/10.1016/j.molcel.2018.06.021, PMID: 30033371

Tasic, B., Nabholz, C. E., Baldwin, K. K., Kim, Y., Rueckert, E. H., Ribich, S. A., Cramer, P., Wu,

Q., Axel, R., Maniatis, T. 2002. Promoter choice determines splice site selection in protocadherin α and -γ pre-mRNA splicing. **Molecular Cell** 10:21-33. DOI: https://doi.org/10.1016/S1097-2765(02)00578-6, PMID: 12150904

Thongjuea, S., Stadhouders, R., Grosveld, F. G., Soler, E., Lenhard, B. 2013. r3Cseq: an R/Bioconductor package for the discovery of long-range genomic interactions from chromosome conformation capture and next-generation sequencing data. **Nucleic Acids Research** 41:e132. DOI: https://doi.org/10.1093/nar/gkt373, PMID: 23671339

Thorvaldsdóttir, H., Robinson, J. T., Mesirov, J. P. 2013. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. **Briefings in Bioinformatics** 14:178-192. DOI: https://doi.org/10.1093/bib/bbs017, PMID: 22517427

Thu, C. A., Chen, W. V., Rubinstein, R., Chevee, M., Wolcott, H. N., Felsovalyi, K. O., Tapia, J. C., Shapiro, L., Honig, B., Maniatis, T. 2014. Single-cell identity generated by combinatorial homophilic interactions between alpha, beta, and gamma protocadherins. **Cell** 158:1045-1059. DOI: https://doi.org/10.1016/j.cell.2014.07.012, PMID: 25171406

Trapnell, C., Williams, B. A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M. J., Salzberg, S. L., Wold, B. J., Pachter, L. 2010. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. **Nature Biotechnology** 28:511-515. DOI: https://doi.org/10.1038/nbt.1621, PMID: 20436464

Valverde de Morales, H. G., Wang, H. V., Garber, K., Cheng, X., Corces, V. G., Li, H. 2023. Expansion of the genotypic and phenotypic spectrum of CTCF-related disorder guides clinical management: 43 new subjects and a comprehensive literature review. **American Journal of Medical Genetics Part A** 191:718-729. DOI: https://doi.org/10.1002/ajmg.a.63065, PMID: 36454652

Vukic, M., Chouaref, J., Della Chiara, V., Dogan, S., Ratner, F., Hogenboom, J. Z. M., Epp, T. A., Chawengsaksophak, K., Vonk, K. K. D., Breukel, C. 2024. CDCA7-associated global aberrant DNA hypomethylation translates to localized, tissue-specific transcriptional responses. **Science Advances** 10:eadk3384. DOI: https://doi.org/10.1126/sciadv.adk3384, PMID: 38335290

Wu, Q., Jia, Z. L. 2021. Wiring the Brain by Clustered Protocadherin Neural Codes. **Neuroscience Bulletin** 37:117-131. DOI: https://doi.org/10.1007/s12264-020-00578-4, PMID: 32939695

Wu, Q., Maniatis, T. 1999. A striking organization of a large family of human neural cadherin-like cell adhesion genes. **Cell** 97:779-790. DOI: https://doi.org/10.1016/S0092-8674(00)80789-8, PMID: 10380929

Wu, Q., Zhang, T., Cheng, J. F., Kim, Y., Grimwood, J., Schmutz, J., Dickson, M., Noonan, J. P., Zhang, M. Q., Myers, R. M., et al. 2001. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. **Genome Research** 11:389-404. DOI: https://doi.org/10.1101/gr.167301, PMID: 11230163

Yin, L., Zhang, H., Tang, Z., Xu, J., Yin, D., Zhang, Z., Yuan, X., Zhu, M., Zhao, S., Li, X., et al. 2021. rMVP: A Memory-Efficient, Visualization-Enhanced, and Parallel-Accelerated Tool for Genome-Wide Association Study. **Genomics, Proteomics & Bioinformatics** 19:619-628. DOI: https://doi.org/10.1016/j.gpb.2020.10.007, PMID: 33662620

Yin, M., Wang, J., Wang, M., Li, X., Zhang, M., Wu, Q., Wang, Y. 2017. Molecular mechanism of directional CTCF recognition of a diverse range of genomic sites. **Cell Research** 27:1365-

1377. DOI: https://doi.org/10.1038/cr.2017.131, PMID: 29076501

Yokota, S., Hirayama, T., Hirano, K., Kaneko, R., Toyoda, S., Kawamura, Y., Hirabayashi, M., Hirabayashi, T., Yagi, T. 2011. Identification of the Cluster Control Region for the Protocadherin-β Genes Located beyond the Protocadherin-γ Cluster. **Journal of Biological Chemistry** 286:31885-31895. DOI: https://doi.org/10.1074/jbc.M111.245605, PMID: 21771796

Zhang, H. S., Shi, Z. B., Banigan, E. J., Kim, Y., Yu, H. T., Bai, X. C., Finkelstein, I. J. 2023. CTCF and R-loops are boundaries of cohesin-mediated DNA looping. **Molecular Cell** 83:2856-2871. DOI: https://doi.org/10.1016/j.molcel.2023.07.006, PMID: 37536339

Zhang, M., Huang, H., Li, J., Wu, Q. 2024. ZNF143 deletion alters enhancer/promoter looping and CTCF/cohesin geometry. **Cell Reports** 43:113663. DOI: https://doi.org/10.1016/j.celrep.2023.113663, PMID: 38206813

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nusbaum, C., Myers, R. M., Brown, M., Li, W., et al. 2008. Model-based analysis of ChIP-Seq (MACS). **Genome Biology** 9:R137. DOI: https://doi.org/10.1186/gb-2008-9-9-r137, PMID: 18798982

Zhou, Y., Xu, S., Zhang, M., Wu, Q. 2021. Systematic functional characterization of antisense eRNA of protocadherin alpha composite enhancer. **Genes & Development** 35:1383-1394. DOI: https://doi.org/10.1101/gad.348621.121, PMID: 34531317

Figure S1. Generation of models of CTCF ADA mutation. (**A**,**B**) Genotyping of two independent homozygous single-cell clones of CTCF ADA mutation. (**C**,**D**) Volcano plots of differentially expressed genes in each of the two independent *N2a* cell clones. The downregulated *cPcdh* genes are marked with yellow dots. (**E**) Schematic of CRISPR-mediated DNA editing with a single stranded oligodeoxynucleotide (ssODN) donor for

CTCF ADA mutation in mice. (**F**) Outline of the procedure for generating CRISPR-mediated site-directed mutagenesis in the N-terminal domain of CTCF in mice. (**G-I**) Bar plots showing ratios of mutated bases in high-throughput RNA-seq reads from the heterozygous P0 (**G**) and adult (**H**) cortical, as well as olfactory bulb (**I**) tissues.

Figure S2. Genotyping of mouse models of CTCF ADA mutation. (**A**) Genotyping of the chimeric CTCF ADA mouse by TA cloning and Sanger sequencing. (**B**,**C**) Genotyping of the heterozygous CTCF ADA mice by Sanger sequencing. (**D-I**) Genotyping of a litter of three P0 wild-type mice and three P0 heterozygous CTCF ADA mice by Sanger sequencing.

Figure S3. Decreased enrichments of RAD21 at the promoter and enhancer regions of the *cPcdh* genes in *N2a* cells. (**A-C**) Quantification of alterations of RAD21 enrichments at the *Pcdh* α (**A**) , β (**B**), and γ (**C**) clusters in the cellular model. (**D**) RAD21 ChIP-seq profiles at the *cPcdh* locus in CTCF ADA *N2a* clones compared to the WT clone. (**E-G**) Close-up of RAD21 profiles at the *Pcdh* α (**E**) , β (**F**), and γ (**G**) clusters showing decreased enrichments of RAD21 at *cPcdh* variable exons and enhancers. Note that as internal controls with no

CTCF site, there do not appear decreased RAD21 enrichments at $\alpha c2$ and *HS7*.

Figure S4. Decreased enrichments of RAD21 at the promoter and enhancer regions of the *cPcdh* locus in the mouse brain. (**A-C**) Quantification of alterations of RAD21 enrichments at the *Pcdh* α (**A**), β (**B**), and γ (**C**) clusters in the mouse model. (**D**) RAD21 ChIP-seq profiles at the *cPcdh* locus in heterozygous CTCF ADA mice compared to WT littermates. (**E-G**) Close-up of RAD21 profiles at the *Pcdh* α (**E**), β (**F**), and γ (**G**) clusters showing

decreased enrichments of RAD21 at *cPcdh* variable exons and enhancers.

Figure S5

Figure S5. Enrichments of CTCF at the promoter and enhancer regions of the *cPcdh* genes in *N2a* cells. (**A**) CTCF ChIP-seq profiles at the *cPcdh* locus in CTCF ADA *N2a* clones compared to the WT clone. (**B-D**) Close-up of CTCF profiles at the *Pcdh* α (**B**) , β (**C**), and γ (**D**) clusters showing no decrease of CTCF enrichments at *cPcdh* variable exons and enhancers.

Figure S6

Figure S6. Enrichments of CTCF at the promoter and enhancer regions of the c*Pcdh* locus in the mouse brain. (**A**) ChIP-seq profiles of CTCF at the *cPcdh* locus in WT and heterozygous CTCF ADA mice. (**B-D**) Close-up of CTCF profiles at the *Pcdh* α (**B**), β (**C**), and γ (**D**) clusters.

Figure S7. CTCF YDF motif enables cohesin to have a long residence time on chromatin. (**A**) Violin plot showing the distribution of loop sizes in gained, lost, and unchanged regions, showing that unchanged loops display the smallest size, whereas lost loops typically tend to have the largest size, and gained loops are in an intermediate range. (**B**) Violin plot

depicting the distribution of lengths of merged, split, and unchanged TADs, showing that the length of split TADs tends to be significant longer than that of the merged or unchanged TADs. (**C-E**) Aggregate peak analysis (APA) of signals of the gained (**C**), lost (**D**), or unchanged (**E**) loops. The APA analysis revealed a significant decrease in loop strength of lost loops, alongside a notable increase in looping strength based on the gained loops upon CTCF ADA mutation. The unchanged loops also exhibited a significant decrease in looping strength, indicating that CTCF ADA mutation destabilizes the extruding cohesin complex on chromatin. In each panel, the APA value is indicated in the upper-right corner. (**F**) Bar plot showing numbers of compartment in wild-type and heterozygous CTCF ADA mice. (**G**) Scatterplot displaying the Pearson's correlation of *cis*-eigenvector 1 values between wild-type and heterozygous CTCF ADA mice. (**H**) Heatmaps of Pearson's correlation at chr18: 26,252,991-47,772,017 in wild-type and heterozygous CTCF ADA mice, showing an unchanged compartmentalization upon CTCF ADA mutation. (**I**) Numbers of TADs in wild-type and heterozygous CTCF ADA mice. (**J**) Aggregate TAD analysis (ATA) in wild-type and heterozygous CTCF ADA mice, showing a significant decrease of intra-TAD contacts. (**K**) Pileup of the directionality index scores in 300-Kb regions centered on all TAD boundaries in wild-type and heterozygous CTCF ADA mice, showing the weakened insulation at TAD boundaries. (**L**) Averaged insulation scores in 300-Kb regions centered on all TAD boundaries in wild-type and heterozygous CTCF ADA mice. (**M**) Hi-C loop number in wild-type and heterozygous CTCF ADA mice. (**N**) Aggregate peak analysis (APA) depicting the superimposed signal for genome-wide loops estimated from Hi-C data of wild-type and heterozygous CTCF ADA mice, showing global weakening of chromatin loops upon CTCF ADA mutation. In each panel, the APA value is indicated in the upper-right corner.

Supplementary file 1. Oligonucleotides used in this study

Oligonucleotides for the construction of plasmids for transfection

Mut_sgRNA_F	ACCGTGATTTTGAGGAAGAACAGC
Mut_sgRNA_R	AAACGCTGTTCTTCCTCAAAATCA
Left_Mut_F	AAAACGACGGCCAGTGAATTCGGTTCTTAGAGATAGGGTTTCTCTGT
Left_Mut_R	TTCAGCATCAGCCACAGACACATCCACGTCTTTG
Right_Mut_F	TGTCTGTGGCTGATGCTGAAGAAGAACAGCAGGAAGGACTGC
Right_Mut_R	GACCATGATTACGCCAAGCTTTTGCTGGCTTAACTAGGTGAG

Oligonucleotides for the sgRNA generation for microinjection

T7_Mut-sgRNA-F1	taatacgactcactataGGTGTCTGTGTATGATTTTGgttttagagctagaaatag
T7_Mut-sgRNA-F2	taatacgactcactataggTGATTTTGAGGAAGAACAGCgttttagagctagaaatag
sgRNA-R	AAAAGCACCGACTCGGTGCC
ssODN_Mut	CCAAAAAGAGCAAACTTCGTTACACAGAAGAGGGCAAAGACGTGGATGTGTCTGTGGCTGATGCTGAAGAAGAACAGCAGGAAGGACTGCTGTCTGAGGTTAATGCAGAGAAAGTAGTTG

Oligonucleotides for the cell clone genotyping

geno_F1_Wt	GGATGTGTCTGTGTATGATTTTGAG
geno_F1_Mut	TGTCTGTGGCTGATGCTGAA
geno_R1	CCAGGGCTATGTAGAAGACTCTG

Oligonucleotides for amplifying targeted sites for the mouse genotyping via Sanger sequencing

geno_226228_F	GGAGAAGTGGAGACACTAGAGCAG
geno_226228_R	TAACTTGCATAGAGGCTGTTGTATC

Oligonucleotides for final construction of library for the high-throughput sequencing

Adapter-HTS_F	5'-ACACTCTTTCCCTACACGACGCTCTTCCGATC*T-3'
Adapter-HTS_R	5'-p-GATCGGAAGAGCACACGTCTGAACTCCAGTC-3'
P5 index-1	AATGATACGGCGACCACCGAGATCTACACTATAGCCTACACTCTTTCCCTACACGACGCTCTTCCGATCT
P5 index-2	AATGATACGGCGACCACCGAGATCTACACATAGAGGCACACTCTTTCCCTACACGACGCTCTTCCGATCT
P5 index-3	AATGATACGGCGACCACCGAGATCTACACCCTATCCTACACTCTTTCCCTACACGACGCTCTTCCGATCT
P5 index-4	AATGATACGGCGACCACCGAGATCTACACGGCTCTGAACACTCTTTCCCTACACGACGCTCTTCCGATCT
P5 index-5	AATGATACGGCGACCACCGAGATCTACACAGGCGAAGACACTCTTTCCCTACACGACGCTCTTCCGATCT
P5 index-6	AATGATACGGCGACCACCGAGATCTACACTAATCTTAACACTCTTTCCCTACACGACGCTCTTCCGATCT
P5 index-7	AATGATACGGCGACCACCGAGATCTACACCAGGACGTACACTCTTT

	CCCTACACGACGCTCTTCCGATCT
P5 index-8	AATGATACGGCGACCACCGAGATCTACACGTACTGACACACTCTTT CCCTACACGACGCTCTTCCGATCT
P5 index-9	AATGATACGGCGACCACCGAGATCTACACGACCTGTAACACTCTTT CCCTACACGACGCTCTTCCGATCT
P5 index-10	AATGATACGGCGACCACCGAGATCTACACATGTAACTACACTCTTT CCCTACACGACGCTCTTCCGATCT
P5 index-11	AATGATACGGCGACCACCGAGATCTACACGTTTCAGAACACTCTTT CCCTACACGACGCTCTTCCGATCT
P5 index-12	AATGATACGGCGACCACCGAGATCTACACCACAGGATACACTCTTT CCCTACACGACGCTCTTCCGATCT
P7-index-1	CAAGCAGAAGACGGCATACGAGATCGAGTAATGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-2	CAAGCAGAAGACGGCATACGAGATTCTCCGGAGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-3	CAAGCAGAAGACGGCATACGAGATAATGAGCGGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-4	CAAGCAGAAGACGGCATACGAGATGGAATCTCGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-5	CAAGCAGAAGACGGCATACGAGATTTCTGAATGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-6	CAAGCAGAAGACGGCATACGAGATACGAATTCGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-7	CAAGCAGAAGACGGCATACGAGATAGCTTCAGGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-8	CAAGCAGAAGACGGCATACGAGATGCGCATTAGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-9	CAAGCAGAAGACGGCATACGAGATCATAGCCGGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-10	CAAGCAGAAGACGGCATACGAGATTTCGCGGAGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-11	CAAGCAGAAGACGGCATACGAGATGCGCGAGAGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT
P7-index-12	CAAGCAGAAGACGGCATACGAGATCTATCGCTGTGACTGGAGTTCA GACGTGTGCTCTTCCGATCT

Oligonucleotides for QHR-4C experiments

Adapter-4C_F	5'-GACGTGTGCTCTTCCGATCTGNNNNNNN-3'NH2 C6
Adapter-4C_R	5'-p-CAGATCGGAAGAGCACACGTC-3'NH2 C6
biotin_mHS5-1	5'-biotin–GTGGCTTTGTTACTCTAGGAACAG
biotin_mSEbc	5'-biotin–AACTATGCTCAGGGCCTCTGG
biotin_mSEd	5'-biotin–TCCCTGCCAGCTGAGCTAAG
biotin_mSEe	5'-biotin–TTAGAGGACACGCTGCCGTC
qhr-mhs51-F	AATGATACGGCGACCACCGAGATCTACACTATAGCCTACACTCTTT CCCTACACGACGCTCTTCCGATCTGGAGGTTAAAGCAAAGACTAAG

	ATC
qhr-m-se-bc-F	AATGATACGGCGACCACCGAGATCTACACATAGAGGCACA
	CTCTTTCCCTACACGACGCTCTTCCGATCTGAGGCTTATG
	GGCCTTACAGATC
qhr-m-se-d-F	AATGATACGGCGACCACCGAGATCTACACCCTATCCTACACTCTTT
	CCCTACACGACGCTCTTCCGATCTACAGCAGGCAGCAGCTGATC
qhr-m-se-e-F	AATGATACGGCGACCACCGAGATCTACACGGCTCTGAACACTCTTT
	CCCTACACGACGCTCTTCCGATCTCCAGATTCTGAACCCGACAGAT
	C

Supplementary file 2. Reagents and plasmids used in this study

Antibodies

Anti-CTCF Antibody	Millipore	Cat# 07-729
Anti-Rad21 Antibody	Abcam	Cat# ab992

Chemicals

MEM	Gibco	Cat# 11095080
MEM-NEAA	Gibco	Cat# 11140050
Penicillin-streptomycin	Gibco	Cat# 15140122
Fetal Bovine Serum	Sigma-Aldrich	Cat# F0193
PBS	Gibco	Cat# 70011044
0.25% trypsin-EDTA	Gibco	Cat# 25200056
*Bsa*I	NEB	Cat# R3733S
*Eco*RI	NEB	Cat# R3101S
*Hin*dIII	NEB	Cat# R3104S
*Xba*I	NEB	Cat# R0145S
*Mbo*I	NEB	Cat# R0147L
*Dpn*II	NEB	Cat# R0543L
Lipofectamine 3000 reagents	Invitrogen	Cat# L3000015
Puromycin dihydrochloride	Solarbio	Cat# P8230-25mg
Phanta Max Super-Fidelity DNA Polymerase	Vazyme	Cat# P505-d1
RNase-free water	Invitrogen	Cat# 10977023
Phenol-chloroform	ACMEC	Cat# AC13309
MEGAshortscript Kit	Invitrogen	Cat# AM1354
mMACHINE T7 ULTRA kit	Invitrogen	Cat# AM1345
MEGAclear Kit	Invitrogen	Cat# AM1908
Sodium acetate	Invitrogen	Cat# AM9740
Glycogen	Thermo Scientific	Cat# R0551
PMSG	Solarbio	Cat# P9970
hCG	MCE	Cat# HY-107953
Sodium Chloride	Sigma-Aldrich	Cat# S3014
Hyaluronidase	Sigma-Aldrich	Cat# H4272-30MG
M2 medium	Sigma-Aldrich	Cat# MR-015
Mineral oil	Sigma-Aldrich	Cat# M5310
TRIzol Reagent	Invitrogen	Cat# 15596018

AMPure XP beads	Beckman	Cat# A63881
16% Formaldehyde	Thermo Scientific	Cat# 28906
Glycine	Invitrogen	Cat# 15527013
Sodium deoxycholate	Sigma-Aldrich	Cat# 30970
SDS	Sigma-Aldrich	Cat# 71736
Triton X-100	Sigma-Aldrich	Cat# T8787
EDTA	Sigma-Aldrich	Cat# E1644
LiCl	Sigma-Aldrich	Cat# L4408
NP40	Sigma-Aldrich	Cat# I8896
NeuroBasal medium	Gibco	Cat# 10888022
EDTA-free protease inhibitor	Roche	Cat# 04693116001
Protein A-agarose beads	Millipore	Cat# 16-157
RNase A	Thermo Scientific	Cat# EN0531
Proteinase K	Invitrogen	Cat# AM2546
Equalbit 1 × dsDNA HS Assay Kit	Vazyme	Cat# EQ121-01
DNA Polymerase I, Large (Klenow) Fragment	NEB	Cat# M0210S
Biotin-14-dATP	Thermo Scientific	Cat# 19524016
Deoxynucleotide (dNTP) Solution Set	NEB	Cat# N0446S
T4 DNA ligase	NEB	Cat# M0202S
Streptavidin magnetic C1 beads	Invitrogen	Cat# 65001
Dynabeads M-280	Thermo Scientific	Cat# 11206D

Commercial assays

ClonExpress MultiS One Step Cloning Kit	Vazyme	Cat# C113
VAHTS Universal RNA-seq Library Prep Kit for Illumina	Vazyme	Cat# NR604
VAHTS Universal DNA Library Prep Kit for Illumina	Vazyme	Cat# ND607
QIAquick PCR Purification Kit	Qiagen	Cat# 28104
MinElute Gel Extraction Kit	Qiagen	Cat# 28604
High-Pure PCR Product Purification kit	Roche	Cat# 11732676001

Plasmids

pGL3-U6-sgRNA-vector	This study	N/A
pcDNA3.1-Cas9-vector	This study	N/A
pUC19_CtcfMut-vector	This study	N/A

Supplementary file 3. *In situ* Hi-C statistics

	WT_rep1	WT_rep2	ADA_rep1	ADA_rep2
Valid_interaction_pairs	1,169,250,659	1,282,103,567	1,056,315,746	1,164,695,035
Valid_interaction_pairs_FF	289,994,816	318,174,399	263,315,179	289,472,201
Valid_interaction_pairs_RR	289,921,136	318,100,330	263,312,165	289,512,479
Valid_interaction_pairs_RF	288,683,416	316,540,127	262,692,179	288,544,162
Valid_interaction_pairs_FR	300,651,291	329,288,711	266,996,223	297,166,193
Dangling_end_pairs	66,522,829	75,925,553	46,541,563	76,450,325
Religation_pairs	33,099,080	36,340,974	16,334,596	27,451,754
Self_Cycle_pairs	3,555,453	3,745,368	2,968,630	2,581,732
Single-end_pairs	0	0	0	0
Filtered_pairs	0	0	0	0
Dumped_pairs	467,118	397,900	413,642	261,100
Valid_interaction	1,169,250,659	1,282,103,567	1,056,315,746	1,164,695,035
Valid_interaction_rmdup	1,075,932,537	1,176,201,570	869,554,809	1,019,127,664
Trans_interaction	373,187,735	395,027,651	402,791,776	490,664,581
Cis_interaction	702,744,802	781,173,919	466,763,033	528,463,083
Cis_shortRange	153,890,242	171,454,035	82,565,133	99,359,414
Cis_longRange	548,854,560	609,719,884	384,197,900	429,103,669