1 **Bacteriophage infection drives loss of β-lactam resistance in methicillin-resistant**
2 ***Staphylococcus aureus***
3

4 My Tran*, Angel J. Hernandez Viera*, Patricia Q. Tran, Charlie Y. Mo
5 Department of Bacteriology, University of Wisconsin-Madison, USA
6 *Denotes equal contribution
7 Correspondence: cymo@wisc.edu

**Abstract**

Bacteriophage (phage) therapy has been proposed as a means to combat drug-resistant bacterial pathogens. Infection by phage can select for mutations in bacterial populations that confer resistance against phage infection. However, resistance against phage can yield evolutionary trade-offs of biomedical use. Here we report the discovery of staphylococcal phages that cause different strains of methicillin-resistant *Staphylococcus aureus* (MRSA) to become sensitized to $\beta$-lactams, a class of antibiotics against which MRSA is typically highly resistant. MRSA cells that survive infection by these phages display significant reductions in minimal inhibitory concentration against different $\beta$-lactams compared to uninfected bacteria. Phage-treated MRSA further exhibited attenuated virulence phenotypes in the form of reduced hemolysis and clumping. Sequencing analysis revealed that the different MRSA strains evolved unique genetic profiles during infection. These results suggest complex evolutionary trajectories in MRSA during phage predation and open up new possibilities to reduce drug resistance and virulence in MRSA infections.

*Staphylococcus aureus* is a Gram-positive bacterium found on the skin and
mucosa of humans and animals. It causes a wide range of human diseases, from
endocarditis to pneumonia.[1] Methicillin-resistant *S. aureus* (MRSA) is a form of *S. aureus*
that displays high levels of drug resistance towards β-lactam antibiotics, such as
penicillins and cephalosporins. β-lactams act by inhibiting the activity of transpeptidase
enzymes during peptidoglycan synthesis in bacterial cell walls.[2] β-lactams are one of the
most commonly prescribed drug classes, and many are designated as "Critically
Important" antimicrobials by the World Health Organization.[3]
A chief mediator of β-lactams resistance in MRSA is the SCCmec cassette, a
mobile genetic element that carries the resistance gene *mecA.* MecA encodes for the
penicillin-binding protein 2A (PBP2a), a transpeptidase that has a low affinity for β-
lactams.[4] This lower affinity permits PBP2a to participate in peptidoglycan synthesis even
in the presence of β-lactams, ultimately resulting in cell survival. In addition to *mecA*,
numerous MRSA strains carry β-lactamases, such as BlaZ, that degrade β-lactams, thus
further contributing to drug resistance. MRSA infections pose a considerable public health
risk as they are notoriously difficult to treat and are widespread in communities and
hospital settings. In 2019, MRSA alone accounted for more than 100,000 deaths
attributable to drug-resistant infections worldwide.[5] Developing novel solutions to combat
MRSA is a major focus in academia and industry.
Bacteriophages (phages) are viruses that infect and kill bacteria, posing one of the
greatest existential threats to bacterial communities. Some estimates suggest that 40%
of all bacterial mortality worldwide is caused by phage predation.[6] Due to this lethality,
phage therapy has been proposed as an attractive alternative for combating drug-
resistant bacterial infections. In phage therapy, lytic phages are administered to kill the
bacterial pathogen(s) causing an infection. Phages offer certain advantages over
traditional antibiotics: they are highly specific for their hosts by reducing off-target killing;
they self-amplify and evolve, enabling the rapid generation of new phage variants with
improved activities; and they are generally regarded as safe, based on extremely rare
cases of toxicity in animals and patients.[7] Indeed, against staphylococcal infections, over
a dozen promising case studies and clinical trials have been reported in the past decade.[8]
Despite these advances, routine use of phage therapy is still met with challenges.
Chief among these challenges is the inevitable rise of phage resistance as phage
predation exerts a strong selective pressure on bacterial populations. According to one
analysis that focused on phage therapy outcomes, resistance against phage evolved in
75 percent of human clinical cases in which the evolution of resistance was monitored.[9]
Mutations represent a chief pathway by which bacteria evolve resistance against phage.
To date, the best characterized phage resistance mutations strains involve alterations of
the cell surface that affect phage attachment. For most lytic phages, the first step of their
infection involves attachment of the phage particle to a receptor molecule (or multiple
molecules) on the cell surface; these receptors are often proteins or sugar moieties, which
are recognized by phage proteins.[8,10,11] Mutations in these receptor molecules can
prevent phage binding, leading to resistance. For example, in *Escherichia coli*, mutations
in the cell-wall protein LamB confer resistance against lambda phage infection, while in
*S. aureus*, mutations that modify the glycopolymers in the cell wall, such as wall teichoic
acid (WTA), have been shown to limit the infection by phages.[11,12] Complicating the
picture, studies have revealed that a plethora of additional host mechanisms, including

dedicated anti-phage defense systems, can impact the evolution of resistance against
phage.[13,14] These problems highlight the importance of developing phage treatment
strategies that minimize or capitalize on the evolution of phage resistance.[15]
A unique aspect of phage therapy is the possibility to exploit evolutionary trade-
offs to combat resistant pathogens. A genetic trade-off is defined as an evolved trait that
confers a fitness advantage against a particular selective pressure at the expense of
reduced fitness against an unselected pressure. Across many different species of bacteria,
such fitness trade-offs are known to occur between phage resistance and antibiotic
resistance (Figure 1A). For example, selection by a lytic phage PYO[Sa] on the methicillin-
sensitive *S. aureus* Newman selected for mutations in the *femA* aminoacyltransferase,
leading to increased resistance against phage infection but reduced resistance towards
antibiotics.[16] These studies raise the question of whether such phage/antibiotic resistance
trade-offs can be induced in actual MRSA strains, which would have significant
therapeutic implications. Employing phages to drive the evolution of such beneficial trade-
offs could serve as a means to reduce drug resistance. However, very little is known about
how frequently these trade-offs occur in different bacterial species, or through what
molecular mechanisms they work.
Here we investigate how infection by staphylococcal phages causes three different
MRSA strains to evolve sensitivity to different types of β-lactam antibiotics. Interestingly,
despite being treated by the same phage, the loss of resistance phenotype is mediated
by mutations distinct to each MRSA strain. In addition to resistance, phage infection also
causes a reduction in hemolysis and clumping, which are key virulence phenotypes in
MRSA. Finally, we isolated a mutant version of the phage that displays enhanced killing,
improved resistance reduction, and reduced virulence phenotypes across all three MRSA
strains.
94
**Results**
**Identification of ΦStaph1N with activity against multiple MRSA strains**
We subjected three MRSA strains – MRSA252 (USA200), MW2 (USA400), and
LAC (USA300) – to a panel of staphylococcal phages using plaquing assays (Table S1,
Figure S1). All three MRSA strains are pathogenic isolates that have been implicated in
human infection and disease and have been used as representative examples for
studying MRSA.[17–19] Of the phages tested, one phage, ΦStaph1N, was able to form
distinct plaques on all three MRSA strains (Figure 1B). ΦStaph1N belongs to the myovirus
family of lytic staphylococcal phages, with a genome size of ~138 kB.[20,21]
Despite its ability to infect all three MRSA strains, ΦStaph1N infection was often
unable to effectively lyse MRSA cultures. For example, both MW2 and LAC displayed
incomplete lysis in liquid culture at multiplicities of infection (MOIs) of 0.1 or lower (Figure
S2). At an MOI of 0.1, we observed that all three MRSA strains lysed within 2-3 hours
following phage infection but recovered high cell density after passaging one percent of
the culture into fresh media. We therefore hypothesized that predation by ΦStaph1N
selected for mutations conferred resistance against ΦStaph1N infection, allowing the
culture to regrow. Indeed, ΦStaph1N was unable to form plaques on recovered MRSA
cultures that survived in the initial ΦStaph1N infection (Figure 1B).
Next, we asked whether phage resistance against ΦStaph1N infection could cause
a trade-off in drug resistance in MRSA. Since both phage and β-lactams interface with

the bacterial cell wall, we hypothesized that phage resistance mutations could impact β-
lactam activity. Parental strains of MRSA252, MW2, and LAC displayed higher minimal
inhibitory concentrations (MICs) of ≥48 µg/mL against the β-lactams oxacillin (OXA),
cefazolin (CEF), amoxicillin (AMX), and amoxicillin & clavulanic acid (AMX+CA), visually
indicated by their ability to form lawns surrounding antibiotic strips (Figure 1C, Table S2).
The MRSA strains were sensitive to vancomycin (VANC; MICs = 1.5 µg/mL), which
inhibits cell wall synthesis through a different mechanism than β-lactams.
By contrast, all strains that survived ΦStaph1N infection displayed a reduction in
resistance against OXA, CEF, and AMX+CA, with fold reductions in MIC between 10 and
1000-fold (Figure 1D). No changes in MIC were observed with VANC. These observations
suggest that the loss of resistance might be specific to the mechanism of action of β-
lactams. Furthermore, phage-treated strains remained resistant to AMX alone, likely
caused by the activity of β-lactamases present in the three MRSA strains. Together, these
results suggest that predation by ΦStaph1N causes three different MRSA strains to lose
their resistance towards β-lactams.
Next, we asked how infection at lower MOIs with ΦStaph1N affected β-lactam
resistance in MRSA. We infected the three MRSA strains with ΦStaph1N at MOIs ranging
from 0.01 to $10^{-5}$, isolated the surviving MRSA cells, and tested the survivors for their MIC
against oxacillin; each condition was tested in three independent replicates (Table S2).
For MRSA252, we observed a ~3-order of magnitude fold reduction of MIC in the
surviving cells at an MOI of $10^{-5}$. With MW2, the reduction of MIC markedly decreased
with lower phage levels, showing on average no reduction in MIC at MOIs of $10^{-3}$ or lower
(Table S2). For LAC, two of the replicates displayed an order of magnitude reduction of
MIC at an MOI of $10^{-4}$, while one replicate did show a strong decrease in MIC at the same
MOI. These results suggest that for MRSA252, MOIs as low as $10^{-5}$ of ΦStaph1N infection
can still drive the loss of resistance, while for MW2 and LAC, higher MOIs of phage are
needed to ensure the same outcome of reduced β-lactam resistance.
We then asked whether the reduction in β-lactam can be achieved with a phage
from a different taxonomical group. The MRSA LAC strain is sensitive to infection by
ΦNM1γ6, a lytic version of the temperate phage ΦNM1, displaying plaquing comparable
to ΦStaph1N (Table S1, Figure S1). ΦNM1γ6 belongs to the siphovirus family and was
derived from the *S. aureus* Newman strain.[22–24] We therefore examined whether infection
with ΦNM1γ6 could also reduce β-lactam resistance in LAC. We infected LAC with
ΦNM1γ6 at an MOI of 0.1 and measured the MIC against oxacillin of the surviving cells.
Here, recovered LAC cultures did not show a reduction in MICs (Figure S3). This suggests
that ΦNM1γ6 is unable to reduce resistance in susceptible MRSA hosts, raising the
possibility that different phages vary in their ability to change β-lactam resistance in
MRSA.
153
**Discovery of mutant in ΦStaph1N with enhanced activity against MRSA**
A key advantage of phage therapy over conventional pharmaceutical antibiotics is
that phages themselves can evolve. For ΦStaph1N, we noticed that while the phage was
able to plaque all three MRSA strains, its plaque-forming efficiency was reduced on the
MW2 and LAC strains (Figure 1B, Figure S1). ΦStaph1N plaques on MW2 and LAC
strains were hazy and the overall efficiency of plaquing was approximately two orders of
magnitude less than that with MRSA252. Unexpectedly, we consistently observed

smaller, clear plaques arising in the larger, hazy plaques of LAC (Figure 2A). This led us
to hypothesize that these clear plaques are caused by a mutant ΦStaph1N that plaques
more efficiently on LAC. We therefore isolated phage clones from these single plaques
and tested for their activity against MRSA. This mutant phage, which we called Evo2, is
able to plaque on LAC and MW2 strains with higher efficiency, displaying comparable
plaquing to MRSA252 (Figure 2A, Figure S1). In growth curve experiments, we further
observed that Evo2 lyses MRSA cultures at lower MOIs compared to ΦStaph1N (Figure
S2). Notably, Evo2 exhibits lytic activity against MW2 and LAC even at an MOI of 0.0001,
a concentration at which ΦStaph1N does not show any detectable activity against the two
strains.
We sequenced the genome of Evo2 to determine the genetic mechanism driving
this enhanced activity. We observed a single point mutation in ORF141 that induces a
premature stop codon (Figure S4). Sequence analysis with HHpred predicts ORF141 to
be a putative DNA binding protein with an HTH motif (PDB: 2LVS, E-value: 2.5e-9). We
speculate that this protein is a transcriptional regulator that when inactivated by a
nonsense mutation, increases ΦStaph1N infectivity. Future studies will center on
determining the mechanism of this mutation and why Evo2 only evolved in the LAC strain.
Given Evo2's enhanced activity against MRSA, we then asked how predation by
Evo2 impacted β-lactam resistance in our three MRSA strains. We infected MRSA252,
MW2, and LAC with Evo2 at an MOI of 0.1 and measured the MICs against β-lactams
after 48 hours of passaging. Broadly, similar to the ΦStaph1N infection, infection by Evo2
reduced the MICs of the three MRSA strains against OXA, CEF, and AMX+CA, while
MICs against AMX alone and VAN did not change significantly (Figure 2B).
We then tested how different MOIs of Evo2 impacted β-lactam resistance in MRSA
(Table S2). As with ΦStaph1N, we infected the three MRSA strains with Evo2 at varying
MOIs from $10^{-2}$ to $10^{-5}$ and measured the MIC against oxacillin in the evolved MRSA;
each condition was tested in three independent replicates. For most of the MRSA252
cultures, we did not see a significant difference in the fold reduction of MIC between
ΦStaph1N and Evo2. Across all three strains, we observed that replicate cultures across
different MOIs were unable to recover growth following Evo2 infection (Table S2).
However, cultures of MW2 and LAC that did regrow displayed a loss of oxacillin
resistance. Thus, overall Evo2 displays a higher infectivity against the three MRSA strains
and a greater potency in reducing β-lactam resistance.
194
**Genomic mutations in MRSA strains following phage infection**
Following these MIC tests, we hypothesized that the phage-treated MRSA stains
evolved mutations that increased their sensitivity to β-lactams. We sequenced the
genomes of three clonal isolates (A-C) from each MRSA strain that underwent ΦStaph1N,
Evo2, or a mock infection. We observed that each MRSA strain evolved distinct mutation
profiles (Figure 3A). Irrespective of the strain and phage treatment, most mutations were
predicted to result in substitutions in the protein product, followed by truncations (Figure
S5). Cluster of Orthologous Genes (COG) variants associated with transcription, cell
wall/membrane/envelope biogenesis, coenzyme transport and metabolism, and defense
mechanisms, were the most commonly found categories. Mutations in annotated genes
that appeared at least twice across the clonal replicates are summarized in Table 1.
Information on all detected genetic variants is listed in Supplementary Table 1.

**Table 1. Mutated genes in MRSA following infection with phages ΦStaph1N or Evo2**

| Gene | Description | Strain | Phage infection | Reference |
|------|-------------|--------|-----------------|-----------|
| *sarA* | Transcriptional regulator of antibiotic resistance and virulence | MW2 | Evo2 | [25,26] |
| *mgrA* | Transcriptional regulator of antibiotic resistance and virulence | MW2 | ΦStaph1N, Evo2 | [27,28] |
| *rpoB* | Transcriptional regulator of antibiotic resistance | LAC | ΦStaph1N, Evo2 | [29] |
| *arlR* | Transcriptional regulator of antibiotic resistance and virulence | LAC | ΦStaph1N, Evo2 | [28,30,31] |
| *spoVG* | Transcriptional regulator of antibiotic resistance and virulence | LAC | ΦStaph1N, Evo2 | [32,33] |
| *cysE* | Cysteine and methionine synthesis, serine O-acetyltransferase | MW2 | Evo2 | [34] |
| *metK* | Cysteine and methionine synthesis, S-adenosylmethionine (SAM) synthetase | MW2 | ΦStaph1N, Evo2 | [35] |
| *trpF* | Phenylalanine, tyrosine and tryptophan synthesis, phosphoribosylanthranilate isomerase | LAC | ΦStaph1N, Evo2 | [36] |
| *femA* | Peptidoglycan synthesis, pentaglycine synthesis | MRSA252 | ΦStaph1N, Evo2 | [37,38] |
| *murE* | Peptidoglycan synthesis, UDP-MurNAc tripeptide synthesis | MW2 | ΦStaph1N, Evo2 | [39] |
| *trpS* | Aminoacyl-tRNA synthesis, tryptophanyl-tRNA synthesis | MW2 | ΦStaph1N, Evo2 | |
| *ytqA* | tRNA modifications, mnm$^5$s$^2$U synthesis | MW2 | ΦStaph1N, Evo2 | [40] |
| *yvcD* | Unknown | MW2 | Evo2 | |
| *natA* | ABC transporter | MW2 | ΦStaph1N, Evo2 | [41] |
| *tcaB* | Predicted multidrug efflux pump | MW2 | ΦStaph1N, Evo2 | [42] |

In all MRSA strains, we found mutations that could account for the loss in β-lactam resistance. In MRSA252, both ΦStaph1N and Evo2 infection were selected for frameshift or nonsense mutations in the *femA* gene that would inactive the protein product. FemA is required for the synthesis of the pentaglycine branch on *S. aureus* Lipid II, the peptidoglycan precursor (Table 1). Deletions of *femA* have been shown to increase susceptibility to β-lactams even when PBP2a (encoded by *mecA*) is expressed.[37,38] These results thus provide a potential mechanism for how some MRSA252 cells lose β-lactam resistance after phage selection. At the same time, we noted the presence of other uncharacterized mutations in phage-treated MRSA252. For example, clone A of ΦStaph1N-treated cells carried 2 mutations: a frameshift in *femA* and a substitution mutation in an uncharacterized protein; meanwhile, clone B displayed a substitution mutation in *pfkA*, a predicted ATP-dependent 6-phosphofructokinase and mutation in an intergenic region; clone C showed a substitution mutation in a putative transport protein, called *yueF* (Figure 3B, Supplementary Table 1). The role of these mutations in mediating phage resistance or β-lactam sensitivity, if any, remains unknown.

In MW2, we found mutations in two transcriptional regulators, *mgrA* and *sarA* (Figure 3A, Table 1). Both *mgrA* and *sarA* belong to the family of MarR (multiple antibiotic resistance regulator)/SarA (staphylococcal accessory regulator A) proteins, which regulate drug resistance and virulence in *S. aureus*.[25–27] In ΦStaph1N-treated MW2, only *mgrA* was mutated, while in Evo2-treated MW2, clones also showed nonsense mutations in *sarA*. We also found mutations in *metK* and *ytqA* which are both predicted to be associated with S-adenosylmethione: *metK* synthesizes SAM, while *ytqA* belongs to the radical SAM enzyme family and is predicted to be involved in tRNA modification.[35,40] Phage-treated MW2 also displayed mutations in other genes, including *tcaB and murE*.

Notably, each clonal replicate had multiple mutations in the genome, while by contrast
untreated MW2 cells only displayed a deletion in an intergenic region that is not present
in any of the phage-treated samples.
In the LAC strain, we observed a third, distinct mutational pattern (Figure 3A). Of
note, we found nonsense mutations in *arlR,* which is part of the *arlRS* two-component
signaling system (Table 1). The activity of ArlRS has been implicated in *S. aureus*
virulence, pathogenicity, and oxacillin resistance.[30,31] Further, we observed substitution
mutations in *spoVG*, which is a transcription factor regulating the expression genes
involved in a variety of functions, including cell wall metabolism (Table 1).[32] Indeed,
SpoVG activates the expression of *femA*.[33] Studies have shown that SpoVG modulates
β-lactam antibiotic resistance by modulating cell wall synthesis.[33] Similar to the other two
MRSA strains, phage-treated clones of LAC showed multiple mutations in their genomes,
although their association with β-lactam sensitivity remains unclear. Finally, we observed
mutations in *prsA* and *bioA* that appeared in both the mock and phage treatment
conditions, suggesting that these mutations do not arise due to phage selection.
248
**Phage-treated MRSA strains display reduced virulence phenotypes**
*S. aureus* strains encode a wide variety of virulence factors that facilitate bacterial
infection, and previous studies have used MW2 and LAC strains as models of MRSA
virulence.[27,28] In MW2 and LAC, we observed the evolution of nonsense mutations in
known virulence regulators, such as *mgrA*, *arlR*, and *sarA*. We therefore hypothesized
that phage-treated MRSA cells could also display altered virulence phenotypes in addition
to reduced β-lactam resistance. We first tested the ability of MRSA strains that survived
phage predation for their ability to form biofilms in a Crystal Violet assay. We found that
Evo2 infection of MRSA252 resulted in a significant reduction in Crystal Violet absorption
compared to the parental strain. However, we show no significant difference in Crystal
Violet absorption between parental, mock- and phage-treated MW2 and LAC strains
(Figure S6).
Next, we tested whether phage infection could affect the hemolysis of rabbit blood
cells. Hemolysis is mediated by the secretion of toxins, such as alpha toxins, and plays
an important role in MRSA infection.[43] Expression of these toxins is regulated by virulence
pathways that comprise numerous transcription factors, including *mgrA*, *arlR*, and *sarA.*
Parental MW2 and LAC colonies lysed rabbit blood cells on blood agar plates, producing
distinct halos of clearance around the bacterial cells. As expected, for untreated LAC, the
total area of hemolysis was on average 3-fold larger than that of untreated MW2 (~210
mm$^2$ vs ~70 mm$^2$, respectively); with MRSA252, by contrast, lysis was not detected
(Figure 4A). Following treatment with ΦStaph1N, we observed that MW2 and LAC
displayed a reduced area of hemolysis by 4 to 5-fold. With Evo2-treated cells, we found
that in MW2 the fold reduction was comparable to that of ΦStaph1N-treated cells.
However, for Evo2-treated LAC, loss of hemolysis was even more pronounced, with two
of the replicates showing no detectable hemolysis.
*S. aureus* is also able to bind to fibrinogen, resulting in the agglutination, or
clumping, of bacteria. Clumping is thought to have several functions in the context of
staphylococcal infections. Clumps are likely to be more resistant to clearance by the
immune system, partly because they may be too large to be phagocytosed by
neutrophils.[28] Similar to the expression of hemolysins, previous work has shown that

clumping in MW2 and LAC is regulated by a network of virulence regulators. For MW2
and LAC, we observed that phage-treated cells displayed less clumping than the mock-
treated or parental strain. For MW2, ΦStaph1N infection resulted in a modest reduction,
while Evo2 infection resulted in a reduction of approximately 3-fold (Figure 4B). In LAC,
we found that both ΦStaph1N and Evo2 treatment resulted in comparable reductions of
clumping in surviving cells.
285
**Discussion**
Here we report the discovery that infection by the myovirus ΦStaph1N and its
derivative Evo2 can drive MRSA populations to evolve favorable genetic trade-offs
between phage and β-lactam resistance. MRSA strains that survived phage infection
were resistant to phage; however, they displayed up to a 1000-fold decrease in their MICs
against β-lactam antibiotics. Sequencing revealed that each of the three MRSA strains
tested evolved distinct mutational profiles following phage infection. Of these mutations,
a significant portion of the affected genes are involved in transcriptional regulation,
antibiotic resistance, and cell maintenance. Concurring with the sequencing data, phage-
treated MRSA strains displayed attenuated virulence phenotypes in the form of reduced
hemolysis and cellular agglutination.
Our system highlights the multitude of evolutionary paths that different bacterial
strains can undertake to evolve resistance against phage. Most staphylococcal phages
are thought to adsorb to *S. aureus* by recognizing wall teichoic acid (WTA). Prior work
has revealed that phage resistance in *S. aureus* can arise from the disruption of genes
involved in the synthesis and modification of WTA, such as *tagO or tarM*.[44] Some MRSA
strains also alter cell wall glycosylation through dedicated genes encoded on
prophages.[11] Additionally, the presence of cell wall-bound Protein A has been shown to
decrease phage absorption, likely by masking WTA.[45] However, our results suggest that
the evolution of phage resistance can potentially impact networks of genes in the different
MRSA strains, including metabolic genes and transcription factors. For example, both
*mgrA* and *arlRS* regulate the expression of proteins associated with the cell wall, such as
the Giant Staphylococcal Surface Protein (GSSP).[28] We hypothesize that mutations in
these regulators might perturb the composition of proteins in the cell wall, which in turn
can disrupt phage binding or prevent disrupt injection of the phage nucleic acid into the
cytoplasm. Further studies are needed to dissect the effect of these gene networks on
phage resistance and β-lactam resistance.
Our study also shows some of the challenges associated with determining
phage/antibiotic resistance trade-offs. Not only do different bacterial strains evolve distinct
mutations, but individual clones can accumulate multiple mutations in their genome. Such
complex dynamics have been observed in other coevolving bacteria-phage systems
before and can serve as useful models to study evolution. However, they make it difficult
to predict broader phenotypic outcomes in the bacteria following phage infection. For
example, phage resistance has also been shown to confer so-called "trade-ups," where
a non-selected trait is enhanced (e.g. cross-resistance between phage and antibiotic). In
our study, we cannot rule out that our phage-treated MRSA has evolved (or will evolve)
trade-ups. Although further research is required to test this possibility, the potential clinical
benefit of reduced resistance and virulence in MRSA may outweigh the risks associated
with trade-ups.

Despite evolving distinct mutational profiles, the three MRSA strains nonetheless
displayed a convergence of phenotypes in the form of enhanced phage resistance,
reduced β-lactam resistance, and attenuated virulence. We posit that this convergence is
caused by the involvement of the cell wall in all three phenotypic outcomes. Phages must
interface with the cell wall to adsorb, β-lactams target proteins associated with the
maintenance of the cell wall, and many *S. aureus* virulence factors are embedded within
the cell wall or need to get secreted through it. The integrity and composition of the cell
wall can be determined by numerous genes, ranging from single proteins involved in cell
wall synthesis, such as *femA*, to transcriptional regulators that control the expression of
cell wall-associated proteins, such as *mgrA*. Thus, the mutational patterns observed in
each MRSA strain might reflect unique genetic solutions that enable the bacteria to adapt
to the phage predation, while also maximizing the fitness for that particular strain.
From a biomedical perspective, exploiting phage/antibiotic resistance trade-offs
have been proposed as a means to combat resistance in bacterial pathogens.[15] Not only
could phage treatments reduce the bacterial load of an infection, but also potentially
resensitize the bacterial population to antibiotics that it was previously resistant against.
For example, in our study, we observed that the MIC of phage-treated MRSA against
oxacillin dropped below 2 µg/mL, which is designated by the FDA as the oxacillin MIC
susceptible breakpoint for *S. aureus*. We speculate that infection by ΦStaph1N, Evo2, or
an equivalent phage can facilitate the re-deployment of β-lactams in the treatment against
MRSA infections. Additionally, phage-treated MRSA might also lose virulence and
become more susceptible to immune clearance, which can further improve treatment
outcomes. Future studies will need to examine this hypothesis in animal infection models.
348

**Acknowledgments**
MT is supported by the SciMed GRS program at UW-Madison; CYM is supported by start-
up funds from the Department of Bacteriology at UW-Madison. We are grateful to Dr.
Wilmara Salgado-Pabón for providing us with MRSA strains and helpful suggestions. We
thank the Marraffini and Hatoum-Aslan laboratories for providing us with bacteriophage
samples. We also thank all members of the Mo lab for their scientific input.

**Online Methods**
**Strains and culture conditions**
The bacterial strains used in this study are listed in Table S1. Unless otherwise indicated,
all MRSA strains (LAC, MRSA252, MW2) were grown in Brain Heart Infusion (BHI) media
at 37 ºC with shaking (235 RPM).
361

**Plate-based plaque assay**
Bacterial lawns were prepared by mixing 100 µL of an overnight culture with 5 mL of
melted BHI agarose (top agar). The bacteria and top agar mixture were poured onto a
solid BHI plate. The plate was dried for 10 minutes. 10-fold serial dilutions ($10^0$ -$10^{-7}$
unless otherwise noted) of phage were then spotted on the bacterial lawn. Plates were
then incubated at 37 ºC for 16 hours. Phage titer in plaque-forming units per µL (pfu/µL)
was then calculated.
369

**Phage infection assay**

MRSA strains were plated onto BHI agar plates and grown overnight. Individual colonies
from the parental strains (also referred to as P0) were picked. Single colonies were
inoculated in a round bottle tube containing 5 mL BHI broth. The cultures were incubated
at 37 ºC, 235 RPM for 24 hours. The grown P0 cultures were then diluted 1:100 into fresh
5 mL BHI broth. At the early log phase (OD ~0.3), the bacterial cultures were treated with
phage at an MOI of 0.1, unless indicated otherwise. The treated bacterial cultures were
incubated at 37 ºC with shaking (235 RPM) for 24 hours. The cultures were then
passaged 1:100 into fresh 5 mL BHI broth. This passage was then grown at 37 ºC, 235
RPM, for another 24 hours. Surviving cultures were then used for both phenotypic assays
and sequencing experiments. As a negative control, MRSA strains were passaged using
the steps described above without phage treatment (mock).
382
**MIC assay**
Bacterial lawns were normalized to contain $1x10^8$ CFU/mL bacteria mixed with top agar
for a total volume of 5 mL. The bacteria and top agar mixture is poured onto a solid BHI
plate. The plate was dried for 10 minutes. MIC with increasing concentrations of
antibiotics were placed on the semi-dried bacterial lawn and allowed to dry for 10 minutes.
The plates are then incubated at 37 ºC overnight. For analysis, the plates were imaged,
and the MIC of the bacteria was determined. The MIC is determined at the edge of the
inhibition ellipse intersects the side of the strip.
391
392 **Rabbit blood hemolysis**
393 Phage-treated or mock-treated cultures were diluted to an OD600 of 0.1, 5 µL of this
394 dilution was spotted on rabbit blood TSB agar plates and incubated at 37 ºC for 24 hours.
395 The area of clearance was determined by the following formula:
396 $[\pi (\text{diameter of clearance}/2)^2] - [\pi(\text{diameter of bacterial spot}/2)^2]$
397
**Clumping assay**
Clumping assays were performed as described previously.[46] In short, overnight cultures
were diluted 1:100 and incubated at 37 ℃ until the cultures reached an $OD_{600}$ of 1.5. At
this point, 1.5 mL of culture was washed two times and resuspended with PBS.
Lyophilized human plasma was added for a final concentration of 1.25%. Resuspended
cells were left to sit statically at room temperature. 100uL were taken from the top of the
cell suspensions in 30-minute intervals and the $OD_{600}$ was measured.
405
**Biofilm assay**
Biofilm assay was performed using the crystal violet method as outlined.[47] In brief,
overnight cultures grown in BHI at 37 ºC were back diluted 1:10 into a 96-well bottomed
microwell plate. The plates were incubated at 37 ºC overnight. The contents in the plate
were discarded and washed with PBS. Biofilm fixation was done with the addition of
sodium acetate (2%). Crystal violet (0.1%) was used for staining followed by a final wash
with PBS. Absorbance at 600 nm was read using a spectrophotometer.
413
**DNA sequencing and genome assembly**
Following published protocols, genomic DNA from bacteria and phage was
isolated using phenol-chloroform extraction. Purified DNA was sent to Plasmidsaurus and

SeqCenter for Nanopore and Illumina sequencing, respectively. Reference genomes for
bacterial strains were assembled using Flye v2.9.3 with default settings for long-reads.
This resulted in 1 singular contig assemblies for 252 (2902592bp, 125x) and MW2
(2820460, 600x), and 3 contigs for LAC (2907712, 645x). Phage assemblies for Evo2
and ΦStaph1N were done with the SPADES assembler v3.15.5.
Open-reading frames (ORFs) were called on the assembled bacterial genomes
using Prodigal v.2.6.3,[48] resulting in a gene-feature file (GFF), and translated genes as
.faa and .fna formats. We used BLASTp (Accessed June 4th, 2024), against the protein
BLAST database swissprot_2023-06-28, with an expectation value cutoff of 0.001. The
top hit for each ORF was used as the final functional annotation.
427
**Mutation identification**
A total of 27 genomic samples were collected. DNA was extracted and sent for
long-read sequencing using Oxford Nanopore Technology (Plasmidsaurus, San
Francisco, USA). Reads were filtered using filtlong v0.2.1 using default settings
(https://github.com/rrwick/Filtlong) and with the -p flag 95 (keeping 95% of the best reads).
Quality-filtered long reads were mapped against the respective genomes using minimap2
2.22-r1101[49] resulting in 1 alignment file output per sample (.sam file). The read mapping
software Minimap2[49] was selected because of its suitability to map long reads. Samtools
v1.20[50] was used to convert the .sam files into .bam files, sort the bam file, index the bam
files, and generate a coverage table for each position along the alignment.
Reference files (fasta and GFF files) and the 53 "sorted.bam" alignments files were
imported into Geneious. Variant calling was performed using Geneious Prime Geneious
Prime ® 2024.0.2 (www.geneious.com), using the custom settings: 10% coverage,
minimum 95% variant frequency threshold, and the option for "Analyze effect of variants
on translation" checked.  Variant results and genome annotations table files were
exported as a tab-separated-table, and visualized using R v4.4.0, mostly with the
tidyverse package. Sequencing data processing, quality filtering, and mapping were
performed at the Center for High-Throughput Computing (https://chtc.cs.wisc.edu/).
BLAStp was performed on usegalaxy.eu (Accessed June 4th, 2024).

# Figures

448
449
**Figure 1. Infection by bacteriophage ΦStaph1N drives the loss of β-lactam resistance in MRSA. A.** Schematic of the experimental setup. Drug-resistant (Abx$^R$), phage-sensitive (Phage$^S$) bacterial cultures are infected with phage. The population of infected cells is passaged and allowed to recover. The surviving cell population is resistant to phage infection (Phage$^R$) but has evolved sensitivity to antibiotics (Abx$^S$). **B.** ΦStaph1N infects MRSA strains MRSA252, MW2, and LAC (left panel). Following infection with ΦStaph1N, evolved cultures of the three MRSA strains are resistant to ΦStaph1N (right panel). **C.** ΦStaph1N-treated, evolved MRSA strains show significant loss of resistance against oxacillin, compared to the parental strains. Loss of resistance is indicated by the area of bacterial clearance surrounding the antibiotic resistance strip. **D.** ΦStaph1N treatment causes loss resistance against different β-lactams. Plotted are the fold reductions of minimal inhibitory concentration (MIC) between treated and mock-treated cells. OXI = oxacillin; CEF = cefazolin; AMX = amoxicillin; AMX+CA = amoxicillin & clavulanic acid; VANC = vancomycin.

**Figure 2. Evo2 is a variant of ΦStaph1N with higher activity against MRSA**. **A.** Single Evo2 plaques appear inside hazy ΦStaph1N plaques on LAC (Left-most panel). Compared to ΦStaph1N, Evo2 shows comparable infectivity towards MRSA252 but improved infectivity towards MW2 and LAC (Right-most panels). **B.** Similar to ΦStaph1N, Evo2 infection reduces β-lactam resistance in MRSA.

**Figure 3. Phage infection of MRSA strains produces distinct mutational profiles**. **A.** Coding sequences (CDS) with mutations from the three MRSA strains following phage treatment or mock treatment. For each strain, three isolates were sequenced and their mutations identified. Mutations are color-coded based on the number of occurrences among the three replicates. **B.** Categories of genes with mutations that arose in each MRSA strain and treatment condition.

**Figure 2. Phage treatment of MRSA results in attenuated virulence phenotypes. A.**
MW2 and LAC strains display hemolytic activity on rabbit blood agar plates, while
MRSA252 does not. **B**. Phage-treated MW2 and LAC strains display reduced hemolysis
compared to uninfected cells. **C**. Surviving cultures of MW2 and LAC treated with either
ΦStaph1N (blue) or Evo2 (red) show reduced clumping rates compared to mock
untreated cells (teal).

## Supplemental data

**Figure S1. Phage sensitivity of MRSA strains.** Efficiencies of plaquing of phages on MRSA252, MW2, and LAC. Phages were 10-fold serially diluted and spotted onto top agar overlays of each strain. Phage Evo2 shows the highest plaquing efficiency on all three strains.

**Figure S2. Growth curves of MRSA strains under varying levels of phage infection.**
MRSA252, MW2, and LAC cultures were infected with either ΦStaph1N or Evo2 at the
indicated multiplicity of infection (MOI). The optical density ($OD_{600}$) of the cultures was

495   monitored on an automated plate reader. Each condition was tested in 3 independent
496   replicates.

| Oxacillin | | Cefazolin | | Amoxacillin | | Amoxacillin +clavulanic acid | | Vancomycin | |
|---|---|---|---|---|---|---|---|---|---|
| Mock | ΦStaph1N | Mock | ΦStaph1N | Mock | ΦStaph1N | Mock | ΦStaph1N | Mock | ΦStaph1N |

497
498

**Figure S3. Phage ΦNM1γ6 infection LAC does not drive the loss of β-lactam resistance.** LAC cultures were infected with phage at an MOI 0.1 and allowed to recover. The surviving cells were then assayed for their MICs against different β-lactams and vancomycin using antibiotic strips.

**Figure S4. Sequencing analysis of Evo2.** Evo2 is a mutant form of ΦStaph1N with a nonsense mutation in ORF141. The A to C mutation (marked by the arrow) in Evo2 converts Serine 77 of ORF141 into a stop codon.

507
508 **Figure S5. Types of polymorphisms in MRSA strains following infection by phage**
509 **or a mock treatment.** Plotted are the polymorphisms that were found in a gene with an
510 assigned COG category.

**Figure S6. Effect of phage infection on biofilm formation in MRSA strains.** Cultures
were infected or mock-infected with either ΦStaph1N or Evo2. RP62a is a strain of *S.*
*epidermidis* with known biofilm forming capability, while LM1680 is a derivative of RP62a
that has lost biofilm forming ability.[51,52] Biofilm biomass was assessed by staining with
Crystal Violet. Solubilized crystal violet was quantified by measuring absorbance at 600
nm. Values represent averages and standard deviations of three replicates. Statistical
significance was determined with a two-tailed t-test.

519 **Table S1. Bacterial strains and bacteriophages used in this study**

| Strain Name | Comments | Source or Reference |
|---|---|---|
| MRSA MW2 (USA400) | | Baba *et al.*, 2002 |
| MRSA LAC (USA300) | | Voyich *et al.*, 2005 |
| MRSA252 (USA200) | | Holden *et al.*, 2004 |
| *S. epidermidis* RP62a | methicillin-resistant biofilm-producing *S. epidermidis* | Christensen *et. al.*, 1982 |
| *S. epidermidis* LAM1680 | Derived from *S. epidermidis* RP62a; carries genomic deletion that inactivates biofilm production | Jiang *et al.*, 2016 |
| ΦStaph1N | | Łobocka *et al.*, 2012 |
| Evo2 | Derived from ΦStaph1N | This study |
| ΦNM1$\gamma$6 | Lytic version of ΦNM1 | Marraffini laboratory |
| ΦNM4$\gamma$4 | Lytic version of ΦNM4 | Marraffini laboratory |
| Φ12 | | Marraffini laboratory |
| Andhra | Infects *S. epidermidis* | Hatoum-Aslan laboratory |

521  **Table S2. MIC against oxacillin of MRSA strains treated with different MOIs of phage**

| | MRSA252 | | | | | |
|---|---|---|---|---|---|---|
| | Staph1N | | | Evo2 | | |
| MOI | Rep 1 | Rep 2 | Rep 3 | Rep 1 | Rep 2 | Rep 3 |
| 10^-2 | 0.25 | 0.125 | 0.38 | NG | 2 | 0.5 |
| 10^-3 | NG | 0.94 | 0.19 | 1 | 0.75 | 1 |
| 10^-4 | 0.5 | 0.25 | 0.19 | 0.75 | 1 | 0.5 |
| 10^-5 | 0.25 | 0.38 | NG | 0.38 | NG | NG |
| Mock | >256 | >256 | >256 | >256 | >256 | >256 |
| | MW2 | | | | | |
| | Staph1N | | | Evo2 | | |
| MOI | Rep 1 | Rep 2 | Rep 3 | Rep 1 | Rep 2 | Rep 3 |
| 10^-2 | 3 | 24 | 24 | 4 | NG | NG |
| 10^-3 | 32 | 24 | 48 | 4 | NG | NG |
| 10^-4 | 48 | 96 | 32 | 3 | NG | NG |
| 10^-5 | 96 | 64 | 24 | 2 | NG | NG |
| Mock | 96 | 48 | 32 | 96 | 48 | 32 |
| | LAC | | | | | |
| | Staph1N | | | Evo2 | | |
| MOI | Rep 1 | Rep 2 | Rep 3 | Rep 1 | Rep 2 | Rep 3 |
| 10^-2 | NG | NG | 2 | 0.064 | NG | NG |
| 10^-3 | NG | 3 | 1.5 | 0.032 | NG | NG |
| 10^-4 | 32 | 1.5 | 1 | NG | NG | NG |
| 10^-5 | 32 | 16 | 0.38 | NG | NG | NG |
| Mock | 32 | 48 | 48 | 32 | 48 | 48 |

522  NG: no growth detected

## References

1.  Gorwitz, R. J. *et al.* Changes in the Prevalence of Nasal Colonization with *Staphylococcus aureus* in the United States, 2001–2004. *J INFECT DIS* **197**, 1226–1234 (2008).
2.  Tomasz, A. The Mechanism of the Irreversible Antimicrobial Effects of Penicillins: How the Beta-Lactam Antibiotics Kill and Lyse Bacteria. *Annu. Rev. Microbiol.* **33**, 113–137 (1979).
3.  World Health Organization. *Critically Important Antimicrobials for Human Medicine*. (World Health Organization, Geneva, 2019).
4.  Zapun, A., Contreras-Martel, C. & Vernet, T. Penicillin-binding proteins and β-lactam resistance. *FEMS Microbiol Rev* **32**, 361–385 (2008).
5.  Murray, C. J. L. *et al.* Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet* **399**, 629–655 (2022).
6.  Suttle, S. W. Viruses and Nutrient Cycles in the Sea. **49**, (1999).
7.  Kortright, K. E., Chan, B. K., Koff, J. L. & Turner, P. E. Phage Therapy: A Renewed Approach to Combat Antibiotic-Resistant Bacteria. *Cell Host & Microbe* **25**, 219–232 (2019).
8.  Hatoum-Aslan, A. The phages of staphylococci: critical catalysts in health and disease. *Trends in Microbiology* **29**, 1117–1129 (2021).
9.  Oechslin, F. Resistance Development to Bacteriophages Occurring during Bacteriophage Therapy. *Viruses* **10**, 351 (2018).
10. Azam, A. H. & Tanji, Y. Peculiarities of Staphylococcus aureus phages and their possible application in phage therapy. *Appl Microbiol Biotechnol* **103**, 4279–4289 (2019).
11. Gerlach, D. *et al.* Methicillin-resistant Staphylococcus aureus alters cell wall glycosylation to evade immunity. *Nature* **563**, 705–709 (2018).
12. Meyer, J. R. *et al.* Repeatability and Contingency in the Evolution of a Key Innovation in Phage Lambda. *Science* **335**, 428–432 (2012).
13. Bernheim, A. & Sorek, R. The pan-immune system of bacteria: antiviral defence as a community resource. *Nat Rev Microbiol* **18**, 113–119 (2020).
14. Maynard, N. D. *et al.* A Forward-Genetic Screen and Dynamic Analysis of Lambda Phage Host-Dependencies Reveals an Extensive Interaction Network and a New Anti-Viral Strategy. *PLoS Genet* **6**, e1001017 (2010).
15. Oromí-Bosch, A., Antani, J. D. & Turner, P. E. Developing Phage Therapy That Overcomes the Evolution of Bacterial Resistance. *Annu. Rev. Virol.* **10**, 503–524 (2023).
16. Berryhill, B. A., Huseby, D. L., McCall, I. C., Hughes, D. & Levin, B. R. Evaluating the potential efficacy and limitations of a phage for joint antibiotic and phage therapy of *Staphylococcus aureus* infections. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2008007118 (2021).
17. Voyich, J. M. *et al.* Insights into Mechanisms Used by *Staphylococcus aureus* to Avoid Destruction by Human Neutrophils. *The Journal of Immunology* **175**, 3907–3919 (2005).

18. Holden, M. T. G. *et al.* Complete genomes of two clinical *Staphylococcus aureus* strains: Evidence for the rapid evolution of virulence and drug resistance. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 9786–9791 (2004).

19. Kobayashi, S. D. *et al.* Comparative analysis of USA300 virulence determinants in a rabbit model of skin and soft tissue infection. *J Infect Dis* **204**, 937–941 (2011).

20. Xia, G. & Wolz, C. Phages of Staphylococcus aureus and their impact on host evolution. *Infection, Genetics and Evolution* **21**, 593–601 (2014).

21. Deghorain, M. & Van Melderen, L. The Staphylococci Phages Family: An Overview. *Viruses* **4**, 3316–3335 (2012).

22. Mo, C. Y. *et al.* Type III-A CRISPR immunity promotes mutagenesis of staphylococci. *Nature* **592**, 611–615 (2021).

23. Goldberg, G. W., Jiang, W., Bikard, D. & Marraffini, L. A. Conditional tolerance of temperate phages via transcription-dependent CRISPR-Cas targeting. *Nature* **514**, 633–637 (2014).

24. Baba, T., Bae, T., Schneewind, O., Takeuchi, F. & Hiramatsu, K. Genome Sequence of *Staphylococcus aureus* Strain Newman and Comparative Analysis of Staphylococcal Genomes: Polymorphism and Evolution of Two Major Pathogenicity Islands. *J Bacteriol* **190**, 300–310 (2008).

25. Li, L. *et al.* The Global Regulon *sarA* Regulates β-Lactam Antibiotic Resistance in Methicillin-Resistant *Staphylococcus aureus* In Vitro and in Endovascular Infections. *J Infect Dis.* **214**, 1421–1429 (2016).

26. Zielinska, A. K. *et al.* SARA -mediated repression of protease production plays a key role in the pathogenesis of *S taphylococcus aureus* USA 300 isolates. *Molecular Microbiology* **86**, 1183–1196 (2012).

27. Crosby, H. A. *et al.* The Staphylococcus aureus Global Regulator MgrA Modulates Clumping and Virulence by Controlling Surface Protein Expression. *PLoS Pathog* **12**, e1005604 (2016).

28. Kwiecinski, J. M. *et al.* Staphylococcus aureus uses the ArlRS and MgrA cascade to regulate immune evasion during skin infection. *Cell Reports* **36**, 109462 (2021).

29. Panchal, V. V. *et al.* Evolving MRSA: High-level β-lactam resistance in Staphylococcus aureus is associated with RNA Polymerase alterations and fine tuning of gene expression. *PLoS Pathog* **16**, e1008672 (2020).

30. Bai, J. *et al.* The role of ArlRS in regulating oxacillin susceptibility in methicillin-resistant *Staphylococcus aureus* indicates it is a potential target for antimicrobial resistance breakers. *Emerging Microbes & Infections* **8**, 503–515 (2019).

31. Walker, J. N. *et al.* The Staphylococcus aureus ArlRS Two-Component System Is a Novel Regulator of Agglutination and Pathogenesis. *PLoS Pathog* **9**, e1003819 (2013).

32. Schulthess, B. *et al.* The σ $^B$ -Dependent *yabJ-spoVG* Operon Is Involved in the Regulation of Extracellular Nuclease, Lipase, and Protease Expression in Staphylococcus aureus. *J Bacteriol* **193**, 4954–4962 (2011).

33. Liu, X., Zhang, S. & Sun, B. SpoVG Regulates Cell Wall Metabolism and Oxacillin Resistance in Methicillin-Resistant Staphylococcus aureus Strain N315. *Antimicrob Agents Chemother* **60**, 3455–3461 (2016).

34. Chen, C. *et al.* Characterization of serine acetyltransferase (CysE) from methicillin-resistant Staphylococcus aureus and inhibitory effect of two natural products on CysE. *Microbial Pathogenesis* **131**, 218–226 (2019).

35. Markham, G. D., DeParasis, J. & Gatmaitan, J. The sequence of metK, the structural gene for S-adenosylmethionine synthetase in Escherichia coli. *Journal of Biological Chemistry* **259**, 14505–14507 (1984).

36. Proctor, A. R. & Kloos, W. E. Tryptophan Biosynthetic Enzymes of *Staphylococcus aureus*. *J Bacteriol* **114**, 169–177 (1973).

37. Maidhof, H., Reinicke, B., Blümel, P., Berger-Bächi, B. & Labischinski, H. femA, which encodes a factor essential for expression of methicillin resistance, affects glycine content of peptidoglycan in methicillin-resistant and methicillin-susceptible Staphylococcus aureus strains. *J Bacteriol* **173**, 3507–3513 (1991).

38. Srisuknimit, V., Qiao, Y., Schaefer, K., Kahne, D. & Walker, S. Peptidoglycan Cross-Linking Preferences of Staphylococcus aureus Penicillin-Binding Proteins Have Implications for Treating MRSA Infections. *J. Am. Chem. Soc.* **139**, 9791–9794 (2017).

39. Gardete, S. *et al.* Role of *murE* in the Expression of β-Lactam Antibiotic Resistance in *Staphylococcus aureus*. *J Bacteriol* **186**, 1705–1713 (2004).

40. Jaroch, M. *et al.* Alternate routes to mnm$^5$s$^2$U synthesis in Gram-positive bacteria. *J Bacteriol* **206**, e00452-23 (2024).

41. Kobayashi, K. *et al.* Comprehensive DNA Microarray Analysis of *Bacillus subtilis* Two-Component Regulatory Systems. *J Bacteriol* **183**, 7365–7370 (2001).

42. Maki, H., McCallum, N., Bischoff, M., Wada, A. & Berger-Bächi, B. *tcaA* Inactivation Increases Glycopeptide Resistance in *Staphylococcus aureus*. *Antimicrob Agents Chemother* **48**, 1953–1959 (2004).

43. Otto, M. Staphylococcus aureus toxins. *Current Opinion in Microbiology* **17**, 32–37 (2014).

44. Jurado, A., Fernández, L., Rodríguez, A. & García, P. Understanding the Mechanisms That Drive Phage Resistance in Staphylococci to Prevent Phage Therapy Failure. *Viruses* **14**, 1061 (2022).

45. Moller, A. G. *et al.* Genes Influencing Phage Host Range in Staphylococcus aureus on a Species-Wide Scale. **6**, (2021).

46. Crosby, H. A., Kwiecinski, J. M. & Horswill, A. R. In Vitro Assay for Quantifying Clumping of Staphylococcus aureus. in *Staphylococcus aureus* (ed. Rice, K. C.) vol. 2341 31–36 (Springer US, New York, NY, 2021).

47. Feoktistova, M., Geserick, P. & Leverkus, M. Crystal Violet Assay for Determining Viability of Cultured Cells. *Cold Spring Harb Protoc* **2016**, pdb.prot087379 (2016).

48. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**, 119 (2010).

49. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100 (2018).

50. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

51. Christensen, G. D., Simpson, W. A., Bisno, A. L. & Beachey, E. H. Adherence of slime-producing strains of Staphylococcus epidermidis to smooth surfaces. *Infect Immun* **37**, 318–326 (1982).

657  52. Jiang, W. *et al.* Dealing with the Evolutionary Downside of CRISPR Immunity:
658     Bacteria and Beneficial Plasmids. *PLoS Genet* **9**, e1003844 (2013).