Title: Navigating "tip fog": Embracing uncertainty in tip measurements

Authors: Jeremy M. Beaulieu[1] and Brian C. O'Meara[2]

[1]Department of Biological Sciences, University of Arkansas; Fayetteville, Arkansas, 72701

USA. Email: jmbeauli@uark.edu (Corresponding Author)

[2]Department of Ecology and Evolutionary Biology, University of Tennessee; Knoxville,

Tennessee, 37996-1610 USA.

Data availability. All scripts and substantial outputs are available at

https://github.com/thej022214/MeasurementError

Author contributions: J.M.B. and B.C.O. contributed equally.

Funding: This study was funded by grants from the National Science Foundation (grant

DEB−1916558 awarded to J.M.B. and grant DEB−1916539 awarded to B.C.O.).

Conflict of Interest Statement: The authors declare that they have no conflict of interests.

Acknowledgments

Conversations with Andy Alverson, James Boyko, James Pease, and Stephen Smith were of help

in generating ideas and providing constructive feedback.

Abstract – Nature is full of messy variation, which serves as the raw material for evolution. However, in comparative biology this variation is smoothed into averages. Overlooking this variation not only weakens our analyses but also risks selecting inaccurate models, generating false precision in parameter estimates, and creating artificial patterns. Furthermore, the complexity of uncertainty extends beyond traditional "measurement error," encompassing various sources of intraspecific variance. To address this, we propose the term "tip fog" to describe the variance between the true species mean and what is recorded, without implying a specific mechanism. We show why accounting for tip fog remains critical by showing its impact on continuous comparative models and discrete comparative and diversification models. We rederive methods to estimate this variance and use simulations to assess its feasibility and importance in a comparative context. Our findings reveal that ignoring tip fog substantially affects the accuracy of rate estimates, with higher tip fog levels showing greater biases from the true rates, as well as affecting which models are chosen. The findings underscore the importance of model selection and the potential consequences of neglecting tip fog, providing insights for improving the accuracy of comparative methods in evolutionary biology.

Keywords: evolutionary rates, measurement error, intraspecific variation, tip fog, macroevolution, hidden Markov model

Introduction

True, one portrait may hit the mark much nearer than another, but none can hit it with any very considerable degree of exactness. So there is no earthly way of finding out precisely what the whale really looks like. And the only mode in which you can derive even a tolerable idea of his living contour, is by going a whaling yourself; but by so doing, you run no small risk of being eternally stove and sunk by him. —Herman Melville, Moby Dick

Messy variation is ubiquitous in nature, as Melville keenly observed. This variation provides the raw material for evolution, while smoothed averages of it are crucial for phylogenetic comparative methods. Typically, multiple individuals — or even one — are used to calculate a mean, which is then used as the representative value for a species in trait evolution, correlation, or diversification studies. While we acknowledge some scatter around this idealized value, we often dismiss it as minor, believing it only slightly reduces the power of our methods, if at all. However, ignoring this variation introduces more than just noise; it can fundamentally undermine analyses. That is, it can change the models we choose as best fits to the data (e.g., Sylvestro et al., 2015), produce misleadingly precise parameter estimates (e.g., Ives et al., 2007), and reveal seemingly compelling patterns that are entirely artificial (e.g., O'Meara & Beaulieu, 2024). Ignoring variance at the tips is not just muddying the waters, it is creating spurious new islands towards which we set sail.

The sources of this variation are numerous and additive. First, every measurement carries uncertainty, a concept ingrained in us from the early days of learning science with quizzes about significant digits and finite resolution of tick marks on our meter sticks. Then there is biological uncertainty, such as the exact length of a squid's tentacle or the precise point where a leaf begins. Sampling uncertainty arises when a subset is used to represent the entire group. Finally, true intraspecific variability exists due to differing optimal conditions, genetic drift within

populations, and phenotypic plasticity. This means that even the most precise measurement from

one population might not reflect the overall species mean.

In comparative biology, we often overlook the complexity of uncertainty. We might

consider "measurement error" as the standard error of the observed measurements for species,

but the common default is to assume this error is zero. Moreover, the array of factors

contributing to this uncertainty extends far beyond what we traditionally categorize as

measurement error. This intraspecific variance has been described by various terms, including

"specific variances" (Cheverud et al., 1985), "residual variation" (Lynch, 1991), "phenotypic

variation" (Felsenstein, 2008), and "measurement error" (Harmon & Losos, 2005; O'Meara et

al., 2006; Silvestro et al., 2015). Some of these terms suggest specific mechanisms. Other terms

are more descriptive but may have meanings outside our field that can cause confusion. To avoid

ambiguity, we propose a new term: "tip fog." This term captures the variance that occurs at the

present between the true species mean derived from the evolutionary process and what an

experimenter records as a value, without being tied to any particular mechanism, and it applies to

characters that are discrete or continuous.

Here, we show why accounting for tip fog is critical by showing its impact on both

continuous and discrete comparative models. We rederive methods to estimate this variance and

assess its feasibility and importance in a comparative context. By highlighting the consequences

of ignoring tip fog, we urge the community to adopt these methods as default standards into all

comparative analyses to avoid the pitfalls of misleading model selection, inferring biased

parameter estimates, and interpreting artificial patterns.

Estimating tip fog for continuous trait evolution

There has been extensive research on the importance of tip fog (under various names) in continuous models, including approaches like independent contrasts (Felsenstein, 2008) and univariate models where model parameters or the comparison of model fit are relevant (Harmon & Losos 2005; Ives et al., 2007; Revell & Reynolds, 2012; Silvestro et al., 2015). All these studies concluded that tip fog can significantly affect the results and recommend its inclusion in analyses, particularly Silvestro et al. (2015). However, its use remains rare. Popular software packages, such as *ouch* (Butler & King, 2004), *surface* (Ingram & Mahler, 2013), and *revBayes* (Hohna et al., 2016), do not allow for accounting for tip fog in their Ornstein-Uhlenbeck models. Packages like *OUwie* (Beaulieu et al., 2012) and *bayou* (Uyeda & Harmon, 2014) allow specifying tip fog as a model parameter but default it to zero; the function fitContinuous in *geiger* (Pennell et al., 2014) also allows it to be inferred, though it also assumes a default value of zero.

The reluctance to incorporate tip fog may stem from the difficulty of obtaining empirical values of sample variance (which is a subset of tip fog) — it is challenging to acquire species means for all traits, let alone the variance in that value. Additionally, any observed value is likely to be an underestimate, capturing only some sources of variation. Estimating tip fog introduces another free parameter into the model, so despite arguments for its potential utility, researchers might opt for simpler models.

The way we parameterize tip fog in *OUwie* is rather straightforward and follows from O'Meara et al. (2006) and Ives et al. (2007). For a standard Brownian motion model, the evolution of a trait over time is modeled as a random walk, with the rate of evolution described

95   by the rate parameter σ^2. The expected variance-covariance matrix, $\mathbf{V}$, reflects the variances and

96   covariances of trait values among different species, based on their shared evolutionary history.

97   To incorporate tip fog, ζ_c, the within-species variance is added to the diagonal elements

98   of $\mathbf{V}$, representing the additional variance in trait values due to factors specific to each species.

99   The modified variance-covariance matrix, $\mathbf{V}^*$, is expressed as:

$$\mathbf{V}^* = \mathbf{V} + \zeta_c \mathbf{I}$$

where $\mathbf{I}$ is the identity matrix. Adding ζ_c to the diagonal of $\mathbf{V}$ allows the model to account for

both the phylogenetic covariance among species as well as estimate the additional variance

within each species. It is possible to have a different ζ_c for each individual species: the variance

in log length for a squishy squid is likely larger than the variance in log length for a crunchy

crustacean species, for example. One might imagine that a ζ_c as a percentage of each species'

mean could also work. In our implementation here, we assumed all species had the same value

for tip fog when estimating it (but not necessarily when simulating).

For more complex models, such as those allowing the rate parameter σ^2 to vary and/or

traits to evolve towards specific trait "optima" (i.e., like Ornstein-Uhlenbeck models) based on a

discrete regime, the addition of ζ_c follows the same formulation described above. The only

difference lies in how $\mathbf{V}$ is constructed. However, the impact of not accounting for tip fog on

evolution rates in such models is quite dramatic.

To illustrate, we conducted a set of simulations where the generating models were a

multiple-rate Brownian motion model (BMS) and a multiple-optima, multiple-rate Ornstein-

Uhlenbeck model (OUMV). We first created an identifiable two-state regime mapping on a

randomly generated 200-tip phylogeny (birth rate set to 0.4 events Myr^{-1}, and death rate of 0.2

events Myr^{-1}) in TreeSim (Stadler 2011) with the root to tip length scaled to one. Under both

generating models, we set the rate for the first regime to $\sigma_1^2 = 1$ units Myr^{-1} and scaled the rate

for the second regime to take on either $\sigma_2^2 = 1.25$, 3, or 5 units Myr^{-1}. For the OUMV model, we

set a global α parameter, which controls the rate of the "pull" towards individual trait means

(here we set $\theta_1 = 1$, and $\theta_2 = 3$), ensuring that the half-life corresponds to 50% the height of the

tree (i.e., $\alpha = 1.39$ units Myr^{-1}).

Tip fog was simulated by resampling each tip value from a normal distribution centered

at the individual species mean and with a standard deviation that was a percentage of the mean

for each species. We generated 100 data sets each, where the percentage varied 10%, 20%, 30%,

40%, and 50% of the mean. Each data set was then evaluated under BM1, BMS, OU1, OUM,

and OUMV models. Estimates for evolutionary rates were summarized as weighted harmonic

averages using Akaike weights (equivalent to weighted arithmetic means of the wait times);

estimates of trait means were based on a weighted arithmetic average. All simulations were

performed in the R package *OUwie* (Beaulieu et al., 2012).

The results of these simulations are presented in Figure 1 and Table S1. In general, not

accounting for tip fog substantially affects the estimates of evolutionary rates in complex models

with multiple rate regimes. As the level of fog increases, the evolutionary rates are increasingly

biased upward, regardless of the regime. For example, with just 30% fog under a BMS

generating model the average rate in regime 1 is 10- to 30-fold higher than the generating model.

In contrast, when ζ_c is estimated as part of the model ($+\zeta_c$), the evolutionary rates generally align

more closely with their true values (for the same models as above, 0.9 to 1.1 times the true

value). However, in the OUMV model, at higher fog levels, the rates exhibit a downward bias in

regime 1 and an upward bias in regime 2, with the upward bias being substantially more

pronounced. Nevertheless, incorporating tip fog into the model substantially improves the rate

estimates compared to not accounting for it.

For the OUMV models, tip fog had a pronounced effect on estimates of the trait mean

(θ_2) for regime 2 only, regardless of the underlying rate or whether fog was included in the

model (Fig. S2). To better understand this, we conducted an additional simulation, assuming an

OUM model where the evolutionary rate was set equal across regimes $\sigma^2 = 2$ and varied θ_2 to be

1.25x, 3x, and 5x on a pectinate tree with an identifiable two-state regime mapping. Results from

these simulations exhibited a similar pattern: in the presence of fog, the estimates of theta for the

second regime are increasingly underestimated as the amount of tip fog increases (Fig. 2).

Examining the confidence regions surrounding estimates of θ_2, α, and σ^2 using *dentist* (Boyko

& O'Meara, 2024) reveals that tip fog introduces greater uncertainty in estimates of α and σ^2,

which incidentally impacts estimates of θ_2, even when ζ_c is included in the model. This

uncertainty is also reflected in model support; the presence of fog leads to some support for

models that include additional complexity, such as BMS and OUMV (Table S1). Overall, these

results suggest that any model fit should be interpreted with consideration of the parameter

estimates and the underlying uncertainty.

Impact of tip fog on discrete trait evolutionary models

The impact of character misassignments on the accuracy and reliability of continuous-time

Markov models remains largely unexplored outside of tree inference contexts (e.g., Felsenstein,

2004; Ho et al., 2007; Rambaut et al., 2009; Kuhner & McGill, 2014; Davis & Navin, 2016).

Misassignments, whether due to data collection errors or legitimate polymorphisms, introduce

uncertainty about the true state of a species and can lead to erroneous estimates of transition rates

163 between states and exaggerated biological patterns (O'Meara & Beaulieu, 2024). Here we apply

164 a framework, first proposed by Felsenstein (2004), for continuous-time Markov models that

165 allows for the simultaneous estimation of character misassignments and state transition rates.

166 Conceptually, this approach is best understood as a type of hidden Markov model (HMM). In

167 such models, the true state is not directly observed, making it "hidden," while the observed data

168 represent a noisy or misclassified version of these states. The goal of the model is to infer the

169 underlying true states by estimating the likelihood of observing the data given the true states. It is

170 rather straightforward in that the general model remains unchanged, except we alter the observed

171 probabilities at the tips.

172  Suppose that i indexes n tips in a tree, and that S_i represents the true underlying state of

173 tip i and that O_i corresponds to the observed state for tip i. When these states are known exactly,

174 as is assumed by any standard continuous-time Markov model, we assume the state of each tip is

175 $P(O_i = o \mid S_i = o) = 1$. In the binary case, o might represent the presence or absence of a particular

176 character state (e.g., woody versus herbaceous plants, feathers versus no feathers). However,

177 when these observations are subject to uncertainty, as is often the case, we assume then that the

178 observed states, O_i, are generated conditionally on the true states, S_i. The probability of the

179 observed state o given that the true state is s can be expressed by $P(O_i = o \mid S_i = s) = 1 - \zeta_{o,s}$ where

180 $\zeta_{o,s}$ defines what we refer to as the "tip fog probability". Thus, the probability at a given tip then

181 becomes $1 - \zeta_{o,s}$ for the observed state and $\zeta_{o,s}$ for the alternative state. For an arbitrary number of

182 states when state i is observed, the probability of each alternative state (i.e., $j \neq i$) is $\zeta_{i,j}$ with the

183 observed state being $1 - \sum_{j \neq i} \zeta_{i,j}$, where $\zeta_{i,j}$ represents the probability that the observed state is i

184 when the true state is j.

The likelihood of the model is obtained by maximizing the standard likelihood formula,

$L = P(D \mid \mathbf{Q}, \mathrm{T}, \zeta_d)$, for observing character states, D, given the continuous-time Markov model,

$\mathbf{Q}$, a fixed topology with a set of branch lengths (denoted by T). Note that we have added an

additional free parameter in the formula, ζ_d, that denotes a single tip fog probability for all

observed states. When $\zeta_d = 0$, the likelihood simply reduces to the likelihood of a standard

continuous-time Markov model. We implemented these "tip fog" models as part of the *corHMM*

package (Beaulieu et al., 2013; Boyko & Beaulieu, 2021), allowing for the specification of as

many tip fog probabilities, ζ_i, as there are unique observed states in the data set.

We conducted a simulation study to assess the impact of ignoring tip fog, as well as the

overall behavior when estimating it. A 200-tip phylogeny was generated (birth rate set to 0.4

events Myr^{-1}, and death rate of 0.2 events Myr^{-1}) in TreeSim (Stadler 2011), which was used to

simulate 100 datasets assuming equal transitions rates of $q_{01} = q_{10} = 0.025$ transitions Myr-1

between binary states. We varied the root age of the phylogeny to take on four ages: 5, 10, 15,

and 20 Myr. In a recent publication (O'Meara & Beaulieu, 2024) we showed that inaccuracies in

character state assignments significantly affect younger trees more due to shorter overall tree

lengths, and so varying clade age was meant to mimic this effect. To simulate tip fog, we

randomly altered the observed state of 1%, 5%, 10%, 15%, or 20% of taxa to be the reverse of its

true state. For each simulation replicate, we fit two classes of models: 1) equal rates (ER, single

rate for all transitions) and all rates different (ARD, two independent rates) without estimating

tip fog probabilities (i.e., referred to as "Default"), and 2) ER and ARD with each estimating a

single ζ_d parameter. Rate estimates within each of the model classes were then summarized by

calculating a weighted harmonic mean of each transition parameter using the Akaike weights

(w_i).

As with continuous trait data, failing to account for tip fog dramatically inflates transition

rates. We found that the magnitude of upward bias increases as the degree of tip fog also

increases (Fig. 3). Even a tip fog probability of 1%, which is expected to result in just two taxa

out of 200 being misassigned to the wrong state, substantially biased transition rates upward. As

expected, the magnitude of the effect did depend on clade age. For example, with 1% tip fog

probability in a 5 Myr tree, rates more than double their true value ($q_{01} = 0.049$ and $q_{10} = 0.069$

transitions Myr^{-1}), while for a clade age of 20 Myr, rates are only slightly upward biased ($q_{01} =$

0.030 and $q_{10} = 0.033$ transitions Myr^{-1}). Nevertheless, when tip fog was 5% and higher, the

model-averaged rates were orders of magnitude higher than the true values regardless of clade

age (Fig. 3). When tip fog probability was estimated as part of the model ($+\zeta_d$), the transition

rates remain relatively stable across different fog levels (Fig. 3). Although younger clades still

show a slight upward bias, it does not systematically increase with increasing tip fog as it does in

the default model fits. Notably, estimates of ζ_d are consistently centered on their true values,

indicating the model effectively infers the degree of tip fog (Fig. 3C).

We also found that failing to account for tip fog substantially impacts model weights.

With just 5% tip fog, the weight shifts almost entirely (>90%) to the ARD model, a pattern that

was consistent across all clade ages (Table S2). When the tip fog probability is estimated under

an ER generating model, the Akaike weights favor the generating ER model with the distribution

of the weights converging towards the null weight expected based solely on the penalty term

(Fig. 4). This indicates that uncorrected tip fog not only inflates rate estimates but can also

erroneously make the data choose too complex models.

To illustrate this further, we also included an HMM model that contained two distinct

rate classes ($q_{01A} = q_{10A}$ and $q_{01B} = q_{10B}$) and an additional transition rate governing the transition

among them (i.e., three transition rates total). When this model was included as part of the

default model set, the ARD and HMM models competed for the highest Akaike weight, with the

ER model showing very little support generally for tip fog values >1% (Fig. S3; Table S3).

Interestingly, the HMM consistently had the highest support in trees with clade ages >10 Myr

(Table S3). When the tip fog probability was estimated as part of the model, support for the

HMM consistently converged toward the null Akaike weight, which was substantially lower than

either the ER or ARD models (Fig. S3).

We were also curious about the performance of the $+\zeta_d$ models when included as part of

a broader set that also included models that did not estimate ζ_d. It may be that even though the

ER + ζ_d fits well when compared to other $+\zeta_d$ models, the added complexity of the additional

parameter reduces the power to detect tip fog when it is present. However, when we pooled the

default and tip fog model sets together and recalculated the Akaike weights, the ER + ζ_d model

emerged as the best model across all fog values except $\zeta_d = 1\%$ (Fig. 4C; Table S2). In that case,

these results indicate that such low fog values are often difficult to distinguish from no fog,

which is expected.

Extending to state-speciation and extinction models

The approach for estimating tip fog from discrete traits extends naturally to state-speciation and

extinction models (SSE; Maddison et al., 2007; FitzJohn et al., 2009; Beaulieu & O'Meara,

2016). We parameterize the initial conditions for $D_{N,i}(t)$ – the probability that a lineage observed

in state i at time t – to be 1 - $\zeta_{i,j}$ for the observed state and $\zeta_{i,j}$ for the alternative state. The initial

conditions for $E_i(0)$, the probability that a lineage in state i at time t would go completely extinct

by the present, remains unaltered. The likelihood calculation then proceeds down the tree as

usual.

255   When the input tree represents a sample of all extant species within a focal clade, the

initial conditions for $D_{N,i}(0)$ must also account for the sampling fraction, f_i, which specifies the

probability that a species with true state i is sampled and included in the tree. For example, if a

tip is observed in state 0 at the present, the initial conditions for $D_{N,0}(0) = f_0 (1 - \zeta_d)$ and

$D_{N,1}(0) = f_1 \zeta_d$ for the alternative state. The initial probability for $E_0(0)$ remains the probability of

a lineage not being present in the phylogeny, either by going extinct or not being sampled, and is

therefore set as $E_i(0) = 1 - f_0$. These modifications have been implemented in the hidden state-

speciation and extinction model in the R package *hisse* (Beaulieu and O'Meara 2016).

263   Unincorporated tip fog should have two effects on the estimation and interpretation of

parameters in an SSE model. First, tip fog should erroneously inflate transition rates, though not

to the degree that they are in continuous-time Markov models. This is because SSE models

jointly estimate speciation, extinction, and state transition processes, making it less likely for the

model to attribute too much to transition rates alone. Instead, character misassignments are more

likely to be absorbed as part of the variation in speciation and/or extinction rates. However, this

benefit comes with a trade-off: the model may homogenize diversification rates among observed

states, which can lead to increased support for models that assume some form of character-

independence. In other words, some of the tips observed in state 0 are actually in state 1, and

vice versa, making the states seem more similar evolutionarily than they should be.

273   To investigate how SSE models behave with tip fog, we simulated scenarios where the

turnover rate $(\lambda + \mu)$ for state 1 was nearly double that of state 0. Specifically, we simulated 100

trees with $\lambda_0 = 0.22$ events Myr^{-1} and $\lambda_1 = 0.42$ events Myr^{-1}, with extinction rates set to 75% of

the speciation rates in both cases, and equal transition rates of 0.025 transitions Myr^{-1}. Each tree

started in state 0 and evolved for 50 Myr, resulting in an average of about 200 taxa per tree. To

avoid patterns from simulation time bias or inflating the effect of tip fog by stopping at a fixed

number of taxa, which can result in a clade with zero length branches, we terminated simulations

at a pre-specified time. Tip fog was introduced by randomly altering the true state of 1%, 5%,

10%, 15%, or 20% of taxa to the incorrect state. For each simulation replicate, we fit two sets of

six models, including both character-independent (e.g., CID-2) and character-dependent models

(e.g., BiSSE), with either equal or asymmetric transition rates (see Table S3). Each model set

either ignored tip fog (referred to as "default") or estimated its probability ($+\zeta_d$). We calculated a

weighted harmonic average of the transition rates across model fits using Akaike weights.

Estimates of turnover rates were summarized as a weighted harmonic mean of the rates

represented at the tips of the tree. That is, for each model, the marginal probability of each state

(and rate class for CID-2) was computed for every tip, and then computed as the weighted

harmonic mean across all models using the Akaike weights.

As expected, our simulations revealed that transition rates became increasingly inflated

with higher degrees of tip fog when not accounting for tip fog in the model (Fig. 5A). In

addition, as expected the magnitude of the rate inflation is muted compared to the continuous-

time Markov models for the same tip fog (Fig. 3). In contrast, when tip fog was estimated as part

of the model, the individual transition rates generally remained close to their true values as did

the estimates of the tip fog probability (Fig. 5C).

Unexpectedly, turnover rates tended to converge as the degree of tip fog increased,

regardless of whether tip fog was estimated (Fig. 5B). We suspect this pattern arises for two

distinct reasons. First, in the default model set that does not include ζ_d, the convergence appears

to be driven by an increasing model weight toward character-independent models as the level of

tip fog increases (Fig. 6A, Table S4). There is increased support for models that do not

differentiate based on the observed character states, leading to more uniform estimates of

turnover rates across different states. In contrast, for the $+\zeta_d$ models, support for character-

dependent models remained stable across different levels of simulated tip fog (Fig. 6B).

The convergence of turnover rates despite the stability in model support for character-

dependence within the $+\zeta_d$ models is likely due to how tip rates are summarized when tip fog is

present. Consider a BiSSE model where the turnover rate is 0.35 events Myr^{-1} for state 0 and

0.70 events Myr^{-1} for state 1. In the absence of tip fog, the rate for a given tip would simply be

the observed state since there is no uncertainty as to the true state. However, with the tip fog

probability being estimated a tip might be observed in say state 1, but there is some uncertainty

as to whether it is state 0 instead. Thus, we must account for this when summarizing the rates

within a given model when tip fog probability is included, and this tends to homogenize

diversification rates as tip fog increases. For instance, if a model is estimated to have a 5% tip

fog probability, the tip rate for a taxon observed in state 1 for that model is calculated as a

weighted average such that $(0.35 \times 0.05) + (0.70 \times 0.95) = 0.68$ events Myr^{-1}. Now, if the tip fog

probability increases to say 20%, as we did in our simulations, the tip rate would adjust

downwards to $(0.35 \times 0.2) + (0.70 \times 0.8) = 0.63$ events Myr^{-1}. This adjustment would also cause

the turnover rate for tips observed in state 0 to gradually increase, contributing to the overall

homogenization of turnover rates, even with clear support for the character-dependent models

included in the model set.

Discussion

While the importance of tip fog in continuous traits has long been acknowledged, we were surprised by how accurately it can be estimated from both continuous and discrete data directly. This is significant because tip fog is not just simply adding another parameter — it represents the extent to which tip data distorts or misrepresents the underlying reality. Given the feasibility of estimating it and the detrimental effects of setting it to zero, we have updated all our software to estimate tip fog by default, and we recommend others do the same [*NB: only once this paper is in press, as a peer review may find an issue we have missed so we do not want to enable this by default until then*]. Anyone converting biological variation into discrete data knows this inevitably leads to fuzzy cases, and this fuzziness is even more pronounced with continuous variation. Ignoring tip fog can result in confidently incorrect conclusions rather than mere uncertainty. Even if practitioners are hesitant to increase model complexity by estimating an additional parameter, selecting an arbitrary value of say, 10%, is likely more accurate than the current practice of assuming tip fog is 0%.

Our implementation of tip fog is straightforward. For continuous traits, we use a constant value across all tips, while for discrete traits, we follow a similar approach in our simulations, though we also allow for varying error rates depending on observed states. For example, in traits like parental care, it is more likely that species with this trait might be missed rather than incorrectly reported as having it. However, there are opportunities to increase complexity. Tip fog for continuous traits could be modeled as a proportion of the observed state instead of a fixed standard deviation. Moreover, different amounts of fog could be estimated for species categorized by factors such as observations from herbaria versus field studies, species with extensive records versus those with fewer, and observations made by undergraduates versus

faculty. Incorporating the number of observations per species could also refine estimates of

continuous tip fog. Exploring different regimes of tip fog across the phylogenetic tree offers

additional avenues for improvement (*sensu* Ives et al., 2007).

We hasten to acknowledge that tip fog is not the only source of uncertainty in

macroevolutionary studies. Uncertainty in topology and branch lengths, as well as

unincorporated heterogeneity in the evolutionary process, also likely play significant roles.

Properly incorporating tip fog does not negate the need to consider these other factors. However,

just as incorporating tree uncertainty by conducting analyses across a set of trees is essential, so

is incorporating tip fog.

The concept of tip fog presents an opportunity to revisit the basic principles of model

selection. Models with or without tip fog estimation can be compared using metrics such as AIC,

AICc, or BIC, all of which account for model complexity. The model with the lowest score on

these metrics offers the best balance between complexity and fit. For instance, if a model that

includes tip fog estimation has a ΔAIC of 0, and a model that forces tip fog to zero has a ΔAIC of

1.4, the model estimating tip fog is superior. However, it is not uncommon to encounter model

choice being based on requiring that a more complex model outperform a simpler one by a

certain arbitrary margin before considering it (i.e., ΔAIC > 2). However, such an approach is

neither necessary nor appropriate (Burnham & Anderson, 2004).

We also note that tip fog is distinct from approaches that account for polymorphism in tip

data, although they are related. For instance, consider a character with states yellow, white, and

red, where one species exhibits polymorphism with some flowers being yellow and others white,

while most species in the clade are uniformly one of the three colors. In such a case, the

likelihood calculation would start with $P(O_i = \text{yellow}) = P(O_i = \text{white}) = 1$ for that species [rather

than 0.5 for each, as noted by Felsenstein (2004)]. This differs from errors such as a researcher

misassigning a yellow specimen as white due to poor lighting. Despite this distinction, tip fog

can still be applied alongside polymorphism to improve model accuracy. In fact, this might be

particularly useful for models of biogeography where polymorphic scoring is a central feature

(Ree & Smith 2008; Bätscher & de Vos, 2024).

Our study focuses on tip fog within traditional macroevolutionary models, which

typically analyze one or a few characters. However, tip fog can significantly impact models that

handle multiple characters, such as those used for inferring phylogenetic trees and networks

(e.g., Kuhner & McGill). One major component of tip fog is sequencing error, which is likely

more substantial than is typically acknowledged. Incorporating tip fog as a default option in tree

inference is particularly sensible given the large volumes of data often available. Yet, popular

tree inference programs like RAxML-NG (Kozlov et al., 2019) and IQ-TREE (Minh et al., 2020)

currently lack this capability. The omission of tip fog is especially critical for branch length

estimates, as unaccounted-for tip fog tends to lengthen terminal branches (which make up over

half of a tree's edges), thereby inflating overall substitution rates. This can lead to an inaccurate

estimation of tree age if a rate calibration is used. When using multiple fossils or other

calibrations, the effect on overall tree age is less predictable but generally results in an increased

ratio of terminal to internal branch lengths. In any event, while some models for cancer tumor

phylogenies incorporate error expectations (Davis & Navin 2016), sequencing error remains a

significant concern in traditional phylogenetic studies (see also Ho et al., 2005).

We note several important caveats. Our analysis has focused on predictable errors, such

as a species mean being off by 10% or a 20% chance of misclassifying a species' state as woody

instead of herbaceous. However, we have not addressed more extraordinary sources of error,

such as entering a unitless mass in milligrams for one species while using grams for others,

confusing range data between a plant and an insect due to homonyms, omitting the sign for

longitude, or recording a missing value as -99, which is then incorrectly treated as a valid state.

Tip fog, as we incorporate it, is unlikely to address these types of errors effectively. While tip fog

can account for certain uncertainties, it relies on the data being fundamentally accurate.

Additionally, tip fog does not correct errors in tree or network topology or branch lengths – these

also remain important to incorporate. While our work uses likelihood and AIC, we expect similar

results using Bayesian methods or using model selection criteria beyond AIC. Even with

Bayesian methods, which deal quite well in uncertainty, existing common approaches essentially

put full prior weight on the data being completely right, not allowing any exploration about the

possibility of nonzero tip fog (but see Revell & Reynolds, 2012). Either adding one or more tip

fog parameters or allowing a looser coupling between observed and actual states, would allow

Bayesian methods to incorporate this important factor.

Finally, several questions remain unanswered. For instance, empirical estimates of

variability at tips, such as the standard deviation of samples, might underrepresent tip fog

because they do not account for factors affecting all modern samples, such as environmental

plasticity. This issue has not been explored in our study; it remains uncertain whether estimating

tip fog directly is more effective than relying on empirical estimates of tip variability or if adding

an "additional" fog parameter to the model would be beneficial. While we have incorporated tip

fog into SSE models for discrete traits, we have not applied it to SSE models of continuous data

(e.g., QuaSSE; FitzJohn, 2010). We also have not explored multivariate models, as discussed by

Felsenstein (2008). Additionally, some simulation results were unexpected. For example, we

anticipated that tip fog would significantly improve estimates of θ in an Ornstein-Uhlenbeck

model, but the effect appears less pronounced than expected. The presence and impact of tip fog

in empirical studies are still unknown; analyzing existing datasets to estimate tip fog and assess

the consequences of ignoring it would be a valuable next step.

Conclusions

Melville warned that those who seek to understand whales must risk their boats being crushed.

Similarly, comparative analyses are at risk of failing due to unrecognized variation from a

myriad of sources — what we term "tip fog." Yet, this risk can be mitigated. By incorporating tip

fog into our standard models, we can improve the accuracy of our inferences and avoid the

pitfalls of confidently incorrect conclusions. As we navigate the complexities of biological data,

making tip fog a standard consideration in our comparative models will better anchor our

analyses.

References

Bätscher, L., & de Vos, J. M. (2024). Avoiding impacts of phylogenetic tip-state-errors on

dispersal and extirpation rates in alpine plant biogeography. *Journal of Biogeography, 51(6)*,

1104-1116. doi: https://doi.org/10.1111/jbi.14811

Beaulieu, J. M. & O'Meara, B. C. (2016). Detecting hidden diversification shifts in models of

trait-dependent speciation and extinction. *Systematic Biology, 65(4)*, 583-601. doi:

https://doi.org/10.1093/sysbio/syw022

Beaulieu, J. M., Jhwueng, D.-C., Boettiger, C., & O'Meara, B. C. (2012). Modeling stabilizing

selection: expanding the Ornstein-Uhlenbeck model of adaptive evolution. *Evolution, 66(8)*,

2369-2383. doi: https://doi.org/10.1111/j.1558-5646.2012.01619.x

Beaulieu, J. M., O'Meara, B. C., & Donoghue, M. J. (2013). Identifying hidden rate changes in the evolution of a binary morphological character: the evolution of plant habit in campanulid angiosperms. *Systematic Biology, 62(5)*, 725-737. doi: https://doi.org/10.1093/sysbio/syt034

Boyko, J. D., & Beaulieu, J. M. (2021). Generalized hidden Markov models for comparative datasets. *Methods in Ecology and Evolution, 12(3)*, 468-478. doi: https://doi.org/10.1111/2041-210X.13534

Boyki, J. D., & O'Meara, B. C. (2024). dentist: quantifying uncertainty by sampling points around maximum likelihood estimates. *Methods in Ecology and Evolution, 15(4)*, 628-638. doi: https://doi.org/10.1111/2041-210X.14297

Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodal inference*. Springer Press.

Butler, M. A., & King, A. A. (2004). Phylogenetic comparative analysis: a modeling approach for adaptive evolution. *The American Naturalist, 164(6)*, 683-695. doi: https://doi.org/10.1086/426002

Cheverud, J. M., Dow, M. M., & Leutenegger, W. (1985) The quantitative assessment of phylogenetic constraints in comparative analyses: sexual dimorphism in body weight among primates. *Evolution, 39(6)*, 1335–1351. doi: https://doi.org/10.1111/j.1558-5646.1985.tb05699.x

Davis, A. & Navin, N. E. (2016). Computing tumor trees from single cells. *Genome Biology, 17*, 113. doi: https://doi.org/10.1186/s13059-016-0987-z

Felsenstein, J. (2004). *Inferring phylogenies*. Sinauer Associates.

Felsenstein, J. (2008). Comparative methods with sampling error and within-species variation:

contrasts revisited and revised. *The American Naturalist, 171(6)*, 713-725. doi:

https://doi.org/10.1086/587525

FitzJohn, R. G. (2010). Quantitative traits and diversification. *Systematic Biology, 59(6)*, 619-

633. doi: https://doi.org/10.1093/sysbio/syq053/

FitzJohn, R. G., Maddison, W. P., & Otto, S. P. (2009). Estimating trait-dependent speciation

and extinction rates from incompletely resolved phylogenies. *Systematic Biology, 58(6)*, 595-

611. doi: https://doi.org/10.1093/sysbio/syp067

Harmon, L. J. & Losos, J. B. (2005). The effect of intraspecific sample size on type I and type II

error rates in comparative studies. *Evolution, 59(12)*, 2705-2710. doi:

https://doi.org/10.1111/j.0014-3820.2005.tb00981.x

Ho, S. Y. W., Phillips, M. J., Cooper, A., & Drummond, A. J. (2005). Time dependency of

molecular rate estimates and systematic overestimation of recent divergence times.

Molecular Biology and Evolution, 22(7), 1561-1568. doi:

https://doi.org/10.1093/molbev/msi145

Ho, S.Y.W., Lancer, R., Phillips, M. J., Barnes, I., Thomas, J. A., Kolokotronis, S.-O., &

Shapiro, B. (2007). *Systematic Biology, 60(3)*, 366-375. doi:

https://doi.org/10.1093/sysbio/syq099

Höhna, S., Landis, M. J., Heath, T. A., Boussau, B., Lartillot, N., Moore, B. R., Huelsenbeck, J.

P., & Ronquist, F. (2016). RevBayes: Bayesian phylogenetic inference using graphical

models and an interactive model-specification language. *Systematic Biology, 65(4)*, 726-736.

doi: https://doi.org/10.1093/sysbio/syw021

Ingram, T., & Mahler, D. L. (2013). SURFACE: detecting convergent evolution from

comparative data by fitting Ornstein-Uhlenbeck models with stepwise Akaike Information

Criterion. *Methods in Ecology and Evolution, 4(5)*, 416-425. doi:

https://doi.org/10.1111/2041-210X.12034

Ives, A. R., Midford, P. E., & Garland, T., Jr. (2007). Within-species variation and measurement

error in phylogenetic comparative methods. *Systematic Biology, 56(2)*, 252-270. doi:

https://doi.org/10.1080/10635150701313830

Kozlov, A. M., Darriba, D., Flouri, T., Morel, B., & Stamatakis, A. (2019). RAxML-NG: a fast,

scalable and user-friendly tool for maximum likelihood of phylogenetic inference.

Bioinformatics, 35(21), 4453-4455. doi: https://doi.org/10.1093/bioinformatics/btz305

Kuhner, M. K., & McGill, J. (2014). Correcting for sequencing error in maximum likelihood

phylogeny inference. *G3 (Bethesda), 4(12)*, 2545-2552. doi:

https://doi.org/10.1534/g3.114.014365

Lynch, M. (1991). Methods for the analysis of comparative data in evolutionary biology.

Evolution, 45(5), 1065-1080. doi: https://doi.org/10.1111/j.1558-5646.1991.tb04375.x

Maddison, W. P., Midford, P. E., & Otto, S. P. (2007). Estimating a binary character's effect on

speciation and extinction. *Systematic Biology, 56(5)*, 701-710. doi:

https://doi.org/10.1080/10635150701607033

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler,

A., & Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic

inference in the genomic era. *Molecular Biology and Evolution, 37(5)*, 1530-1534. doi:

https://doi.org/10.1093/molbev/msaa015

O'Meara, B. C., & Beaulieu, J. M. (2024). Noise leads to the perceived increase in evolutionary

rates over short time scales. *PLoS Computational Biology*, Accepted.

O'Meara, B. C., Ané, C., Sanderson, M. J., & Wainwright, P. C. (2006). Testing for different

rates of continuous trait evolution using likelihood. *Evolution, 60(5)*, 922-933. doi:

https://doi.org/10.1111/j.0014-3820.2006.tb01171.x

Pennell, M. W., Eastman, J. M., Slater, G. J., Brown, J. W., Uyeda, J. C., FitzJohn, R. G., Alfaro,

M. E., & Harmon, L. J. (2014). geiger v2.0: an expanded suite of methods for fitting

macroevolutionary models to phylogenetic trees. *Bioinformatics, 30(15)*, 2216-2218. doi:

https://doi.org/10.1093/bioinformatics/btu181

Rambaut, A., Ho, S. Y. W., Drummond, A. J., & Shapiro, B. (2008). Accommodating the effect

of ancient DNA damage on inferences of demographic histories. *Molecular Biology and*

Evolution, 26(2), 245-248. doi: https://doi.org/10.1093/molbev/msn256

Ree, R. H., & Smith, S. A. (2008). Maximum likelihood inference of geographic range evolution

by dispersal, local extinction, and cladogenesis. *Systematic Biology, 57(1)*, 4-14. doi:

https://doi.org/10.1080/10635150701883881

Revell, L. J., & Reynolds, G. (2012). A new Bayesian method for fitting evolutionary models to

comparative data with intraspecific variation. *Evolution, 66(9)*, 2697-2707. doi:

https://doi.org/10.5061/dryad.7fv08k72

Stadler, T. (2011). Simulating trees with a fixed number of extant species. *Systematic Biology,*

60(5), 676-684. doi: https://doi.org/10.1093/sysbio/syr029

Sylvestro, D., Kostikva, A., Litsios, G., Pearman, P. B., & Salamin, N. (2015). Measurement

errors should always be incorporated in phylogenetic comparative analysis. *Methods in*

Evolution and Evolution, 6(3), 340-346. doi: https://doi.org/10.1111/2041-210X.12337

Uyeda, J. C., & Harmon, L. J. (2014). A novel Bayesian method for inferring and interpreting the

dynamics of adaptive landscapes from phylogenetic comparative data. *Systematic Biology,*

63(6), 902-918. doi: https://doi.org/10.1093/sysbio/syu057

Figure 1. Uncertainty in estimating the evolutionary rate (σ^2) as a function of tip fog, with the generating model assuming various (**A**, **B**) multiple-rate Brownian motion (BMS) and (**C**, **D**) multiple mean, multiple-rate Ornstein-Uhlenbeck (OUMV) models. Panels (**A**) and (**C**) depict cases where the rate for regime 2 is 1.25 times that of regime 1, while (**B**) and (**D**) show a rate 3 times that of regime 1. Tip fog was simulated by resampling each tip value from a normal distribution centered at the individual species mean and with a standard deviation that was a percentage of the mean for each species. Data sets were then evaluated under BM1, BMS, OU1, OUM, and OUMV models, with rates summarized using a weighted harmonic mean based on Akaike weights (see text). Darker boxes indicate rate summarized across models excluding tip fog (Default), which show an upward bias in evolutionary rates as fog levels increase, regardless of the regime. The less saturated boxes represent rates summarized across models that estimate tip fog ($+\zeta_c$), where evolutionary rates generally align more closely with true values. Dashed blue and orange lines indicate the generating values for regimes 1 and 2, respectively.

Figure 2. Uncertainty in estimating the trait means (θ_i) when the generating model is a multiple-mean Ornstein-Uhlenbeck model (OUM), and the simulated data sets contained differing levels of tip fog. (**A**) depicts a scenario where the trait mean for regime 2 is 3 times that of regime 1, whereas (**B**) depicts a scenario where regime 2 is 5 times that of regime 1. Tip fog was simulated by resampling each tip value from a normal distribution centered at the individual species mean and with a standard deviation that was a percentage of the mean for each species. Data sets were evaluated under BM1, BMS, OU1, OUM, and OUMV models, with rates summarized using a weighted mean based on Akaike weights (see text). Darker boxes indicate θ_i summarized across models excluding tip fog (Default); the less saturated boxes represent θ_i summarized across models that estimated tip fog ($+\zeta_c$). Dashed blue and orange lines indicate the generating values for regimes 1 and 2, respectively. In both scenarios, as the amount of tip fog increases, estimates for θ_2 are increasingly underestimated, irrespective of whether tip fog was estimated.

Figure 3. Uncertainty in estimating transition rates using an equal rates (ER) continuous-time
Markov model ($q_{01} = q_{10} = 0.025$ transitions Myr^{-1}) with increasing levels of tip fog. Panels (**A**)
and (**B**) show rate uncertainty for clades aged 5 Myr and 10 Myr, respectively. Younger clades
show greater uncertainty due to shorter tree lengths (see main text). To simulate tip fog, we
randomly altered the observed state of 1%, 5%, 10%, 15%, or 20% of taxa to be the reverse of its
true state. Data sets were evaluated under an equal-rates model (ER, single rate for all
transitions) and an all-rates different model (ARD, two independent rates). Rate estimates within
each of the model classes were then summarized by calculating a weighted harmonic mean of
each transition parameter using the Akaike weights. Darker boxes indicate transition rates
summarized across models excluding tip fog (Default); the less saturated boxes represent
transition rates summarized across models that estimated tip fog ($+\zeta_d$). Dashed blue and orange
lines indicate the generating values for state *0* and state *1*, respectively. (**C**) depicts uncertainty in
ζ_d estimates across all clade ages, with the dashed red line indicating the true ζ_d.

Figure 4. Summaries of model support based on Akaike weight (*w*) for equal rates (ER) and all rates different (ARD) continuous-time Markov models fit (**A**) without tip fog and (**B**) fit including tip fog as a parameter, or (**C**) pooled together as part of an inclusive model set. The generating model for these simulations was an equal rates (ER) continuous-time Markov model ($q_{01} = q_{10} = 0.025$ transitions Myr^{-1}) with increasing levels of tip fog. To simulate tip fog, we randomly altered the observed state of 1%, 5%, 10%, 15%, or 20% of taxa to be the reverse of its true state. Dashed lines in (**A**) and (**B**) represent the null expectation of the Akaike weight as the average Akaike weight if we assume an equal likelihood across all models.

Figure 5. Uncertainty in estimating (**A**) transition rates and (**B**) turnover rates under a state-speciation and extinction model with increasing levels of tip fog. The generating model was a character-dependent model (CD), where state *1* to have turnover rate ($\lambda_1 + \mu_1 = 0.735$ events Myr^{-1}) that was nearly 2x the rate of state *0* ($\lambda_0 + \mu_0 = 0.385$ events Myr^{-1}), with state transitions between *0* and *1* set at 0.025 transitions Myr^{-1}; extinction fraction was set at 0.75 for both regimes. To simulate tip fog, we randomly altered the observed state of 1%, 5%, 10%, 15%, or 20% of taxa to be the reverse of its true state. Data sets were evaluated two sets of six models, including both character-independent (e.g., CID-2) and character-dependent models (e.g., BiSSE), with either equal or asymmetric transition rates (see Table S3) Each model set either ignored tip fog (Default) or estimated its probability ($+\zeta_d$). Rate estimates within each of the model classes were then summarized by calculating a weighted harmonic mean of each rate parameter using the Akaike weights. Darker boxes indicate transition rates summarized across models excluding tip fog (Default); the less saturated boxes represent transition rates summarized across models that estimated tip fog ($+\zeta_d$). Dashed blue and orange lines indicate the generating values for state *0* and state *1*, respectively. (**C**) depicts uncertainty in ζ_d estimates, with the dashed red line indicating the true ζ_d.

Figure 6. Summary of the cumulative support for character-dependence (CD; turnover rate depends on the state) and character-independence (CID; turnover rates do not depend on the state) when the generating model is a character-dependent model that contains increasing levels of tip fog. Panel (**A**) shows how support for character-independence increases with increasing tip fog when tip fog is ignored (Default), whereas (**B**) when tip fog is estimated as part of the model $(+\zeta_d)$ support for character-dependent models remains stable across different levels of simulated tip fog.