Blocking Toxin Function and Modulating the Gut Microbiota: Caffeic Acid and its Derivatives as Potential Treatments for *Clostridioides difficile* Infection

Yan Guo[1#], Yong Zhang[2#], Guizhen Wang[3], Hongtao Liu[1], Jianfeng Wang[1],

Xuming Deng[1], Liuqing He[4*], Jiazhang Qiu[1*]

[1] State Key Laboratory for Diagnosis and Treatment of Severe Zoonotic Infectious Diseases, Key Laboratory for Zoonosis Research of the Ministry of Education, College of Veterinary Medicine, Jilin University, Changchun, China.

[2] Center for Pathogen Biology and Infectious Diseases, State Key Laboratory for Zoonotic Diseases, The First Hospital of Jilin University, Changchun 130021, China.

[3] Measurement Biotechnique Research Center, College of Biological and Food Engineering, Jilin Engineering Normal University, Changchun 130052, China.

[4] Hunan Provincial Key Laboratory of Animal Intestinal Function and Regulation, Hunan International Joint Laboratory of Animal Intestinal Ecology and Health, Laboratory of Animal Nutrition and Human Health, College of Life Sciences, Hunan Normal University, Changsha, China.

#YG and YZ contributed equally to this work.

*Correspondence: Jiazhang Qiu (qiujz@jlu.edu.cn); Liuqin He (heliuqin@hunnu.edu.cn)

Abstract

Clostridioides difficile infection (CDI) is the leading cause of hospital-acquired diarrhea that seriously threatens public health. The disruption of normal gut microbiota by the use of broad-spectrum antimicrobial agents enables *C. difficile* to proliferate in the colon. The emergence and prevalence of hypervirulent *C. difficile* strains result in increased morbidity, mortality, and recurrence rates of CDI, thus creating a pressing need for novel therapeutics. The multi-domain toxins TcdA and TcdB are the primary determinants of CDI pathogenesis, rendering them ideal drug targets in the anti-virulence paradigm. In this study, we identified caffeic acid and its derivatives as active inhibitors of TcdB via a cell-based high-throughput phenotypic screening. Further mechanistic investigations revealed that caffeic acid phenethyl ester (CAPE) could directly bind to TcdB, thus suppressing InsP6-induced autoproteolysis and inhibiting glucosyltransferase activity. CAPE treatment remarkably reduces the pathology of CDI in a murine infection model in terms of alleviated diarrhea symptoms, decreased bacterial colonization, and relieved histopathological lesions. Moreover, CAPE treatment of C. difficile-challenged mice induces a remarkable increase in the diversity and composition of the gut microbiota (e.g., *Bacteroides*) and alterations of gut metabolites (e.g., adenosine, D-proline, and melatonin), which might partially contribute to the therapeutic outcomes of CAPE against CDI. Our results reveal the potential of CAPE as a therapeutic for the management of CDI, or CAPE might serve as a lead compound for the

45   development of antivirulence drugs targeting TcdB.

47   **Keywords:** *Clostridioides difficile*; toxin B; caffeic acid phenethyl ester; gut

48   microbiota; gut metabolites

Introduction

Clostridioides difficile (*C. difficile*) is an obligate anaerobic, spore-forming, gram-positive bacterial pathogen that is the most common cause of hospital-acquired and antibiotic-associated gastrointestinal diseases, including mild diarrhea and life-threatening pseudomembranous colitis [1, 2, 3]. In the United States, *C. difficile* is responsible for almost half a million infections and approximately 29,000 deaths per year, resulting in an estimated annual healthcare-associated cost of up to $4.8 billion [4, 5]. Importantly, the incidence, severity, mortality and healthcare costs of *C. difficile* infection (CDI) have dramatically increased in the 21st century, at least partially owing to the frequent use of antimicrobial agents and the emergence of hypervirulent strains (e.g., PCR ribotypes 027 and 078) [6, 7, 8]. Limited options are available for the treatment of CDI, while antibiotics including metronidazole, fidaxomicin and vancomycin, are the mainstays of clinical practice [9, 10]. However, antibiotic treatment is ineffective in more than 35% of CDI patients and can lead infection recurrence [11, 12]. Moreover, the rapid evolution of antibiotic resistance in *C. difficile* has further compromised the therapeutic efficacy of current treatment options [13]. Therefore, the high morbidity and mortality, high recurrence rates, and emergence of multiple antibiotic-resistant isolates have made this pathogen a major threat to public health; these factors highlight the urgent need for novel therapies.

C. difficile widely exists in the environment and in the intestinal tract of

humans and animals and is mainly transmitted via the fecal-oral route [14]. The

pathogenesis of CDI is correlated with the use of antibiotics, which impair the

structure and composition of the normal gut microbiota, thus allowing *C. difficile*

to proliferate in the colon [14]. Therefore, previous antibiotic therapy is the major

risk factor for the development of CDI. In the colon, the dormant spores of *C.*

difficile germinate into oxygen-sensitive vegetative cells that produce toxins and

cause symptoms of disease [15]. *C. difficile* secretes two large clostridial

exotoxins (TcdA and TcdB), which are directly responsible for the clinical

pathology of CDI [16, 17]. Epidemiological studies suggest that most disease-

causing *C. difficile* strains have genes that encode TcdB [18, 19]. TcdA and TcdB

have identical functional domains: a N-terminal glucosyltransferase catalytic

domain (GTD), a cysteine protease active domain (CPD), a transmembrane

domain (DRBD), and a C-terminal receptor binding domain (CROPs) [20]. When

secreted into the colon, both toxins are internalized by host cells through

receptor-mediated, clathrin-dependent endocytosis. Following a decrease in

endosomal pH, the toxins undergo conformational changes in the DRBD,

resulting in insertion of the toxin into the endosome membrane, pore formation,

and the release of the GTD and CPD domains into the cytosol. Then, CPD

binding with cytosolic inositol hexakisphosphate ($InsP_6$) activates its protease

activity, leading to the cleavage and release of the GTD. Eventually, the GTD

utilizes UDP-glucose to glucosylate Rho and/or Ras family GTPases, thus

disrupting GTPase signaling and resulting in deleterious effects, including

cytopathic "rounding", dysregulation of the actin cytoskeleton, apoptosis, and

pyroptosis [1]. Although both toxins are cytotoxic to host cells, TcdB shows

greater toxicity than TcdA [21].

Disruption of the gut microbiota by antimicrobial agents results in high

recurrence rates, which limits the application of antimicrobial therapy for CDI [14].

Consequently, there is a critical need to develop nonantibiotic treatments for

CDI, either as standalone or augmented therapies. Importantly, strategies

targeting bacterial virulence factors, including toxins, are termed anti-virulence

strategies and have attracted interest in the 21st century [22]. Owing to its

essential role in *C. difficile* pathogenesis, TcdB has become an ideal target for

the development of anti-virulence drugs [23]. In this study, caffeic acid and a

series of its derivatives were found to inhibit TcdB-mediated cell rounding via

cell-based screening. Among these potential inhibitors, caffeic acid phenethyl

ester (CAPE) had the greatest inhibitory effect. Mechanistically, CAPE bound

to both full-length TcdB and the GTD fragment, leading to alterations in the

secondary structure. This interaction suppressed the $InsP_6$-mediated self-

cleavage of TcdB, as well as the enzymatic activity of GTD, thereby decreasing

the level of glucosylated Rac1 in TcdB-treated host cells and further inhibiting

the cytopathology and cytotoxicity of TcdB. Oral administration of CAPE

significantly protected against CDI in models of animal infection. In addition to

direct toxin inhibition, the therapeutic effect may also be attributed to CAPE-

induced alteration of the gut microbiota (e.g., *Bacteroides*) as well as gut

metabolites (e.g., adenosine, D-proline, and melatonin). Overall, our study

revealed that caffeic acid and its derivatives, particularly CAPE, may serve as

promising lead compounds for the development of anti-virulence drugs to treat

CDI.

Results

CAPE inhibits cell rounding induced by TcdB

TcdB can inactivate Rho GTPases via glucosylation, leading to the loss of

cytoskeletal structure and cell-rounding. Consequently, we utilized this

phenotype to identify potential TcdB inhibitors from 2076 chemicals, including

1515 FDA-approved drugs and 561 natural compounds. The minimal

concentration of recombinant His_6-TcdB (0.2 ng/mL) that induced 100% cell

rounding within 1 h was used for drug screening. This concentration of TcdB

was preincubated with 8 μg/mL of the individual compounds for 1 h before the

compounds were added to Vero cell cultures. At 1 h post-TcdB treatment, cell

morphology was visualized under a light microscope (**Figure 1A**). These

analyses allowed us to identify 22 candidate compounds with potent inhibitory

effects (**Figure 1B**). Interestingly, six out of the 22 compounds were caffeic acid

derivatives. Caffeic acid itself only slightly inhibited TcdB-induced cell rounding,

with a 50% inhibitory concentration (IC_{50}) of 527.5 μg/mL (**Table S1**). Notably,

except for rosmarinic acid, the IC_{50} values for the other screened caffeic acid

derivatives were less than 6 μg/mL, suggesting that these compounds potently

inhibited TcdB-mediated toxicity (**Table S1**). CAPE is the phenethyl alcohol

ester of caffeic acid and a main active component of propolis with broad

pharmacological properties, such as anticancer, anti-inflammatory, antiviral,

antibacterial, antifungal, antioxidant, and immunomodulatory activities [24]. As

CAPE has the lowest IC_{50} (3.0 µg/mL) among the active caffeic acid derivatives,

it was selected as the representative compound for subsequent investigations

(**Table S1**). After preincubation with increasing concentrations of CAPE, dose-

dependent inhibition of cell rounding was observed in both Vero and HeLa cells

(**Figure 1C**). In addition, we also determined whether inhibition of TcdB by

CAPE could lead to loss of toxin function *in vivo* using a murine systemic

intoxication model. A lethal dose of recombinant His_6-TcdB (30 ng) was

preincubated with either DMSO or CAPE (2 or 4 µg/mL) for 1 h at 37 °C,

followed by intraperitoneal injection into mice. Notably, all mice in the DMSO

group (n=10) died within 36 h after toxin injection. However, 100% of mice

pretreated with CAPE 4 (µg/mL) toxin survived at the monitoring endpoint (72

h) (n=10) (**Figure 1D**). Taken together, these data indicate that CAPE is a

potent toxin inhibitor of TcdB.

CAPE decreases $InsP_6$-induced autoproteolysis of TcdB and inhibits the

enzymatic activity of GTD

TcdB is a multi-domain containing toxin that induces toxicity in host cells

via an intricate and multistep mechanism. This mechanism involves (i) receptor-

mediated endocytosis, (ii) pore formation and subsequent membrane

translocation, (iii) autoproteolysis and release of GTD, and (iv) the glucosylation

of small GTPases (**Figure 2A**). To determine the mechanism by which CAPE

suppresses the cytotoxicity of TcdB, we first labeled full-length TcdB with Alexa

Fluor 488; then, the concentration of the endocytosed toxin was determined via

flow cytometry. Apparently, incubation of CAPE with TcdB did not significantly

affect its entry into host cells (**Figure 2B**). Next, we stained CAPE-treated cells

with LysoTracker, a fluorescent dye that accumulates in the acidic organelles of

live cells. In contrast to the vacuolar-type H+-ATPase (V-ATPase) inhibitor

bafilomycin A1 (Baf-A1), CAPE treatment did not alter the intensity of

LysoTracker fluorescence staining in either THP-1 or Caco-2 cells, suggesting

that inactivation of TcdB by CAPE is not due to changes in the pH of endosomal

compartments (**Figure 2C-D** and **Figure S1A-B**).

These observations indicate that CAPE might exert its effects after

membrane translocation. Indeed, preincubation of full-length TcdB with CAPE

blocked the $InsP_6$-induced autoproteolysis of TcdB, as revealed by the gradual

decrease in the production of released GTP, followed by the addition of

increasing amounts of CAPE *in vitro* (**Figure 2E-F**). Furthermore, we

determined the glucosyltransferase activity using recombinant GTD in the

absence or presence of different doses of CAPE. Strikingly, incubation with 2

µg/mL CAPE resulted in a significant reduction in GTD activity, whereas 32

µg/mL CAPE compromised 87.5% of the enzymatic activity (**Figure 2G**).

Importantly, in response to suppression of $InsP_6$-induced autoproteolysis and

GTD function, treatment of TcdB-challenged cells with CAPE markedly

decreased the level of glucosylated Rac1 in a dose-dependent manner (**Figure**

2H-I). Taken together, these results suggest that CAPE disrupts InsP$_6$-induced

autoproteolysis of TcdB and inhibits the enzymatic activity of GTD, thereby

decreasing the glucosylation of Rac1 and alleviating cell rounding.

CAPE alters the secondary structure of full-length TcdB and GTD

To further clarify the mechanisms used by CAPE to inactivate TcdB, we

determined the secondary structures of full-length TcdB and GTD in the

absence or presence of CAPE by circular dichroism spectroscopy. Apparently,

incubation of CAPE with full-length TcdB resulted in a significant increase in the

α-helix content (15.4% to 21.0%) and a concomitant decrease in the β-pleated

sheet content (34.5% to 20.9%) (**Figure 3A-B**). Similarly, the secondary

structure of GTD was markedly altered upon incubation with CAPE, with the

proportion of α-helices decreasing from 37.9% to 0.0% and the proportion of β-

pleated sheets decreasing from 37.2% to 10.9% (**Figure 3C-D**). Taken together,

these data suggest that CAPE may interact with full-length TcdB and GTD to

induce conformational changes, thus impairing their functions.

CAPE directly interacts with full-length TcdB and GTD

To investigate the potential interaction between CAPE and TcdB or GTD,

the binding constants were determined by surface plasmon resonance (SPR)

analysis. The results showed that CAPE has strong binding affinity with full-

length TcdB and GTD, and the KD values were 1.269×10^{-5} M and 1.007×10^{-3}

M, respectively (**Figure 4A-B**). These data demonstrate the direct interaction between CAPE and full-length TcdB/GTD.

Next, we performed molecular docking and molecular simulation dynamics analyses to investigate the interaction between CAPE and GTD (**Figure 5A**). The results revealed that Leu265, Asp286, Thr465, Ile466, Leu519, and Trp520 in GTD were potential binding sites for CAPE (**Figure 5A**). The root mean square deviation (RMSD) of the GTD/CAPE complex was stable over a 100 ns period of simulation (**Figure 5B**). Further energy decomposition analysis showed that the side chains of Leu265, Thr465, Leu519, and Trp520 of GTD contributed to binding with CAPE via van der Waals interactions, among which Trp520 had the highest binding energy (**Figure 5C**). Moreover, we mutated each of the residues that were involved in binding and subsequently determined their glucosyltransferase activities in the presence or absence of CAPE. The enzymatic activities of these mutants were similar to that of wild-type GTD (**Figure 5D**). Mutations in GTD$_{L265A}$, GTD$_{T465A}$, GTD$_{I466A}$, and GTD$_{L519A}$ did not affect the inhibitory effect of CAPE (**Figure 5E-I**). However, the inhibition of GTD$_{W520A}$ and GTD$_{D286A}$ by CAPE was notably compromised (**Figure 5J-K**).

CAPE treatment reduces the pathology of CDI in a murine model

The abovementioned observation that CAPE suppresses the function of TcdB prompted us to further evaluate the potential therapeutic effects of this compound in CDI. To this end, we employed a well-established murine model in which infected mice can develop symptoms resembling those observed in

human CDIs (25). First, cytotoxicity measurements revealed that the viability of

both THP-1 and Vero cells was not affected by CAPE treatment (**Figure S2A-**

B), thus indicating low toxicity. Mice were preconditioned with drinking water

containing 0.5 g/L cefoperazone for 5 days, followed by normal drinking water

for 2 days. After challenge with *C. difficile* BAA-1870 via gavage, mice were

then treated daily with either vehicle or 30 mg/kg CAPE by gavage throughout

the disease course (day 0 to day 5) (**Figure 6A**). Severe weight loss in *C.*

difficile-infected mice was observed on days 2 and 3 post infection and was

accompanied by diarrhea (**Figure 6B**). Notably, compared to the infection

control group, CAPE treatment significantly increased the dry/wet weight ratio

of the mouse feces, suggesting the alleviation of diarrhea symptoms (**Figure**

6C). Correspondingly, the weight loss of mice was relieved upon receiving

CAPE (**Figure 6B**). Moreover, fecal pellets collected on day 3 were used to

determine the bacterial burden. The results revealed that CAPE treatment

significantly decreased the number of *C. difficile* CFUs (**Figure 6D**). On day 3

post-*C. difficile* infection, the colon was collected for histopathological analysis.

Remarkably, *C. difficile*-infected mice exhibited conspicuous colon edema,

epithelial cell injury and inflammatory factor infiltration (**Figure 6E**). In contrast,

these histopathological lesions were significantly less common in mice

administered CAPE (**Figure 6F**). Taken together, these data demonstrate the

efficacy of CAPE in the treatment of CDI.

CAPE induces distinct microbiota changes in CAPE mice when compared

 with model mice

Antibiotic-induced dysbiosis of the gut microbiota promotes the

development of CDI [25]. Therefore, we next determined whether CAPE has any

impact on gut microbial communities that may contribute to its therapeutic

efficacy. To this end, on the 3rd day after *C. difficile* challenge, fecal samples

from the model group and CAPE treatment group were collected and subjected

to microbiota analysis via 16S rRNA gene sequencing. Remarkably, as

reflected by the Venn diagram, CAPE treatment resulted in significant changes

in the microbial composition compared to those in the model group, and only

16.42% of the operational taxonomic units (OTUs) were shared (**Figure 7A**). In

addition, both the α- and β-diversity of the gut microbiota in CAPE-treated mice

were significantly greater than those in model mice. Specifically, the increase

in the Chao1 richness estimator of mice in the CAPE treatment group

suggested a greater number of microbial species and increased evenness in

the fecal samples (**Figure 7B-C**); the increase in the Shannon–Wiener diversity

index and the Simpson diversity index reflected a significant increase in

microbiota diversity (**Figure 7D-E**). Principal coordinate analysis (PCoA)

revealed distinct confidence ellipses between the bacterial communities of the

CAPE-treated group and the model group, thereby revealing substantial

differences in their microbiota profiles (**Figure 7F**).

A comparative analysis was performed to examine changes at the phylum

level following CAPE treatment. *Bacteroidota*, *Firmicutes,* and *Proteobacteria*

were the major constituents in both the CAPE treatment and model groups, suggesting that the administration of CAPE did not significantly alter the mouse microbiota at the phylum level (**Figure 7G**). Next, we compared the 10 most abundant species at the genus level. Notably, CAPE treatment markedly elevated the abundance of *Bacteroides* and decreased the abundance of *Muribaculaceae* and *Parabacteroides* (**Figure 7H**). Leveraging linear discriminant analysis of effect sizes (LEfSe), the microbiota was examined at the genus level to identify bacteria exhibiting substantial differences in abundance between the two cohorts (LDA score > 3.5). After CAPE treatment, the relative abundance of *Bacteroides*, *Ruminococcus_gnavus_group*, *Muribaculum* and *Faecalibaculum* OTUs notably increased, while that of *Hungatella*, *Parasutterella* and *Muribaculaceae* OTUs significantly decreased (**Figure 7I**). *Bacteroides*, *Muribaculum* and *Faecalibaculum* improve colonization resistance to *C. difficile* [26, 27, 28]. Importantly, the abundance of these bacterial species increased following CAPE treatment (**Figure 7J-L**). In contrast, the abundance of *Parasutterella*, a bacterium that facilitates *C. difficile* colonization [29], was reduced in mice receiving CAPE (**Figure 7M**). Taken together, these findings suggest that CAPE treatment accelerates microbiota recovery during CDI.

CAPE alters the levels of gut metabolites in *C. difficile*-infected mice

In addition to the crucial role of the gut microbiota in conferring colonization resistance against *C. difficile*, gut microbiota-derived metabolites significantly

contribute to the disease process [30]. Metabolomic analysis was performed to

evaluate the effects of CAPE treatment on gut metabolites in mice. The PLS-

DA model exhibited significant separation of clusters between the model and

CAPE treatment groups (**Figure 8A**). Among the 1599 metabolites, 109

showed substantial differences in levels between the two groups. Specifically,

CAPE-treated mice exhibited 19 metabolites with upregulated expression and

50 metabolites with downregulated expression compared to those in the model

group (fold change > 1.5) (**Figure 8B**). Furthermore, KEGG enrichment

analysis revealed that the differentially abundant metabolites were significantly

enriched in the purine metabolism and bile secretion pathways (**Figure 8C**).

Adenosine is an endogenous purine nucleoside that binds to specific cell

surface receptors to exert anti-inflammatory effects at high concentrations [31].

Importantly, CAPE treatment of *C. difficile*-infected mice can increase the

adenosine concentration in the gut (**Figure 8D**), which might be attributed to its

inhibition of xanthine oxidase [32]. Similarly, the levels of uric acid, the final

product of purine metabolism, were markedly decreased in mice after receiving

CAPE (**Figure 8D**). An earlier study reported that increasing the gut adenosine

concentration with an adenosine deaminase inhibitor prevents *C. difficile* toxin

A-induced enteritis in mice [33]. Therefore, the increase in gut adenosine induced

by CAPE might play an important role in its therapeutic efficacy in CDI.

C. difficile predominantly utilizes sugars and amino acids for energy

production through various metabolic pathways, such as fermentation [34].

Notably, Stickland fermentation, a Clostridia-specific type of amino acid fermentation, is associated with *C. difficile* pathogenesis [34]. D-proline is one of the preferred nutrients used by *C. difficile* in the Stickland reaction and is critical for its growth [35, 36]. Interestingly, treatment of *C. difficile*-infected mice with CAPE significantly reduced gut D-proline levels (**Figure 8D**). In addition, the metabolomics data also revealed a dramatic increase in the level of melatonin (**Figure 8D**), which is a ubiquitous indolamine that was previously shown to decrease the rate of recurrent CDI [37]. Consistently, quantification of the melatonin content in mouse fecal supernatants using a melatonin detection kit revealed elevated levels of melatonin in CAPE-treated mice (**Figure S3**). Moreover, correlation analysis revealed a positive association between *Faecalibaculum* abundance and melatonin levels (**Figure S4**). Taken together, these results indicate that alterations in gut metabolites caused by CAPE treatment might contribute to its therapeutic efficacy in CDI.

Oral administration of melatonin alleviates the symptoms of CDI

Next, we investigated whether any of the upregulated metabolites had a direct influence on TcdB. To this end, we preincubated recombinant TcdB with each of the selected metabolites prior to addition to the cell cultures. Interestingly, treatment with TcdB and 100 μg/mL melatonin significantly alleviated the cell death induced by this toxin (**Figure 9A**). This observation prompted us to investigate the direct therapeutic potential of melatonin in CDI. Therefore, *C. difficile*-infected mice were orally administered 200 mg/kg

melatonin at 24 h intervals. The weight changes and the dry/wet ratio of the stool samples were monitored daily for three days. The results showed that melatonin-treated mice displayed remarkable alleviation of diarrhea, as manifested by notable improvements in weight recovery and a significant increase in the dry/wet weight ratio on the 2nd and 3rd days after *C. difficile* challenge (**Figure 9B-C**). Consistent with these observations, histopathological assessment of colonic tissue sections also revealed attenuated colon damage upon melatonin treatment (**Figure 9D-E**). Taken together, these data indicate that a high dose of exogenous melatonin could protect mice from CDI and that the CAPE-induced increase in endogenous melatonin might improve the therapeutic outcome of patients with CAPE.

Discussion

C. difficile has become the prevalent agent of nosocomial diarrhea and substantially threatens public health. In particular, the increase in morbidity and mortality caused by the prevalence of hypervirulent and drug-resistant strains has become a global concern [7]. In 2019, *C. difficile* was listed by the US Centers for Disease Control and Prevention as an "urgent threat" that surpassed methicillin-resistant *Staphylococcus aureus* (MRSA) and ESBL-producing *Enterobacterales* [38]. *C. difficile* is a commensal bacterium that colonizes the lower gastrointestinal tract [39]. CDI typically occurs in susceptible individuals whose normal gut microbiota is disrupted by the administration of broad-

spectrum antibiotics [39]. Although CDI is considered an antibiotic-associated

disease, the current treatment for CDI still relies on antimicrobial drugs,

including metronidazole, fidaxomicin and vancomycin [9, 10]. Despite antibiotic

therapy, more than 35% of CDI patients are susceptible to recurrent infection,

which is primarily attributable to the intrinsic effects of antimicrobial agents,

which disrupt the microbiota [12]. In addition, long-term administration of

antibiotics promotes the selection of antibiotic-resistant subpopulations in the

gut, thus increasing the risk of infections caused by antibiotic-resistant and

opportunistic bacteria [40]. As a result, the development of nonantibiotic

therapeutics is needed for the treatment of CDI.

Indeed, numerous novel treatments for CDI are being investigated, among

which the anti-virulence strategy represents one of the most promising and

successful therapeutic approaches [22, 41]. Unlike antibiotics, which target

bacterial components that are critical for bacterial growth, anti-virulence drugs

aim to disarm pathogenic bacteria by neutralizing their virulence factors [42, 43].

Therefore, this strategy imposes less selective pressure that directs bacterial

evolution of antimicrobial resistance [43]. The disease progression of CDI is

strictly dependent on large clostridial toxins (TcdA and TcdB), as strains lacking

both toxins (i.e., A⁻B⁻) were completely avirulent and failed to induce any

discernable disease symptoms in a hamster model of infection [44]. Importantly,

epidemic *C. difficile* strains with increased morbidity and mortality rates produce

greater amounts of toxins [45]. Based on these premises, TcdA and TcdB are

considered ideal drug targets for CDI in the anti-virulence paradigm [46]. Indeed,

bezlotoxumab, a fully humanized monoclonal antibody that neutralizes TcdB,

was recently approved by the FDA and used in CDI patients to reduce the rate

of recurrence [47]. This achievement has encouraged researchers to develop

next-generation anti-virulence treatments. In particular, small molecule

inhibitors of toxin production and/or activity have attracted increasing interest

owing to their advantages in cost and the convenience of oral administration [41].

The multidomain architecture and mechanisms of action of clostridial toxins

have provided distinct options for blocking their toxicity. Indeed, previous efforts

have identified diverse inhibitors that act on distinct steps of intoxication, such

as the inhibition of toxin binding to cell receptors, the suppression of endosomal

maturation, and the inactivation of GTD or CPD activities [23, 48, 49, 50, 51]. Although

which approach is the best remains unknown, we believe that an inhibitor that

targets multiple steps of intoxication would be superior to those that act on a

single step. In this study, we identified a series of caffeic acid derivatives that

were potent inhibitors of TcdB via a phenotypic screen. Further mechanistic

study revealed that CAPE, the phenethyl alcohol ester of caffeic acid, could

block $InSP_6$-induced autoproteolysis of TcdB and inactivate its GTD function

(**Figure 10**). Moreover, CAPE treatment significantly protected against CDI in a

mouse model of infection. Our results suggested that caffeic acid derivatives,

particularly CAPE, are promising lead compounds for the development of anti-

virulence drugs that target TcdB. CAPE is a major component of propolis, a

natural bee-produced substance that has been used as a remedy for thousands

of years in Europe and Asia [52]. As a result, further investigations on the

therapeutic potential of propolis against CDI, either in combination with

standard therapy or as a single therapy, would be of great interest.

Antibiotic-induced gut microbiota dysbiosis is a major risk factor for CDI [53].

Thus, reconstituting the gut microbiota is a promising strategy for CDI therapy.

Fecal microbiota transplantation (FMT) is considered an effective CDI

therapeutic strategy and is widely used in clinical practice [54]. Despite the high

cure rate and increasing popularity of FMT, its safety remains controversial.

Another prominent advantage of anti-virulence drugs is that they allow

maintenance of the normal microbiota, which is of critical importance for the

treatment of CDI. Importantly, CAPE treatment of *C. difficile*-infected mice

restored the structure and composition of the microbiota (**Figure 10**). In

particular, the abundance of *Bacteroides*, a critical gut microbiota component

that is required for *C. difficile* resistance [26], was significantly elevated upon

CAPE treatment. Interestingly, *Bacteroides* were also predominant in CDI

patients administered RBX2660 (Rebyota), an FDA-approved fecal microbiota

product [55, 56]. The increased diversity of the gut microbiota might contribute to

the therapeutic effects of CAPE in CDI.

In addition to the microbiota, gut metabolites also play crucial roles in the

pathogenesis of CDI, either by promoting or suppressing disease progression

[30]. For example, bile acids and amino acids promote spore germination and

pathogenesis, while secondary bile acid and short-chain fatty acids aid in

colonization resistance and suppress spore germination [30]. In addition to

restoration of the gut microbiota, metabolomics analysis of mouse fecal

samples revealed dramatic alterations in gut metabolites upon CAPE treatment

(**Figure 10**). The differentially abundant metabolites were significantly enriched

in the purine metabolism and bile metabolism pathways. Remarkably, the level

of adenosine, an endogenous purine nucleoside that is released from cells

upon injury or inflammation, was dramatically increased in CAPE-treated mice.

In contrast, the amount of uric acid, the final product of purine metabolism, was

reduced. Previous studies showed that the increase in adenosine

concentrations caused by the inhibition of adenosine deaminase via EHNA

prevents *C. difficile* TcdA-mediated damage and inflammation [33]. This

protection might be a result of the anti-inflammatory effects of adenosine, which

rely on the activation of its receptors [57]. In addition, we observed a reduction in

the amount of D-proline, which is one of the preferred amino acids utilized by

C. difficile to generate energy through Stickland reactions [35]. Melatonin is a

ubiquitous indolamine that has broad biological activities [58]. Notably, we

detected a significantly increase in the amount of melatonin in the feces of mice

that were administered CAPE. Importantly, an earlier study demonstrated a

decreased risk of recurrent CDI in patients exposed to melatonin, possibly

owing to its antibacterial and anti-inflammatory activities [37]. In this study, we

found that oral administration of exogenous melatonin alleviated the symptoms

of CDI in a mouse model of infection. Although determining the exact mechanism will require further investigation, we propose that the melatonin-mediated inhibition of TcdB intoxication observed in this study might contribute to its therapeutic effect. Taken together, these results indicate that the alterations in gut metabolites caused by CAPE might also contribute to treatment outcomes for CDI patients.

Methods

Cell lines

All cell lines (HeLa, Caco-2, Vero, HEK293T and THP-1) were obtained from ATCC (Manassas, VA). HeLa, Caco-2, Vero and HEK293T cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen, Carlsbad, CA) supplemented with 10% fetal bovine serum (FBS) (Invitrogen) and 1% (w/v) penicillin–streptomycin. THP-1 cells were cultured in Roswell Park Memorial Institute (RPMI) 1640 medium (Invitrogen) supplemented with 10% FBS and 1% (w/v) penicillin–streptomycin. All cell lines were cultured at 37 °C in a 5% CO_2 atmosphere.

Bacterial strains and reagents

C. difficile BAA-1870 (kindly provided by Aiwu Wu, Guangzhou Medical University) was cultured in brain-heart infusion broth (BHI). The compounds were identified from two main sources: FDA-approved drugs obtained from ApexBio Technology (Houston, TX, USA) and small molecule natural

compounds obtained from Chengdu Ruifensi Biotechnology Co., Ltd. (Chengdu,

China). LysoTracker Red DND-99, kanamycin, and cefoperazone sodium were

purchased from Dalian Meilun Biotechnology Co., Ltd. (Dalian, China).

Mice

Male C57BL/6 mice aged 6-8 weeks were purchased from Liaoning

Changsheng Biotechnology Co., Ltd., and housed in a specific pathogen-free

(SPF) facility. The mice lived in a controlled environment with a temperature of

23±2 °C and a humidity of 55±10 °C. Prior to infection, the mice were

acclimated to the environment for a minimum of one week. All mouse

experiments were approved by the Jilin University Institutional Animal Care

Committee, Jilin University and strictly conducted according to the guidelines

of this committee (SY202309036).

Protein expression and purification

The expression and purification of full-length TcdB were conducted

following the method described by *Yang* et al. [59]. In brief, *Bacillus megaterium*

harboring pHis1522-TcdB was inoculated into LB media supplemented with a

final concentration of 10 μg/mL tetracycline. Once the $OD600_{nm}$ reached

approximately 0.3, xylose (0.5% w/v) was added to induce protein expression

at 37 °C for 16 h. Then, 100 mL of the bacterial cells was harvested by

centrifugation, resuspended in 5 mL of lysis buffer containing 20 mM imidazole,

300 mM NaCl, 20 mM NaH_2PO_4, 500 μM EDTA, and protease inhibitor cocktail

(adjusted to pH 8.0), and subjected to sonication. The cell lysate was then

centrifuged at 13,400 × g for 20 min to remove cell debris. The TcdB protein

was purified using nickel column affinity chromatography. The purity of the

protein was confirmed by SDS–PAGE, and the protein was subsequently

dialyzed against buffer containing 10% glycerol, 20 mM Tris (pH 7.5), and 150

mM NaCl. After the protein concentration was determined, the purified TcdB

protein was aliquoted and stored at -80 °C for further use.

For the purification of GTD, amplified DNA products were inserted into

pET28a (Novagen), and the *Escherichia coli* (*E. coli*) strain BL21 (DE3) was

used for the expression and purification of GTD. Transformed *E. coli* pET28a-

GTD was cultured in Luria–Bertani (LB) medium supplemented with 10 µg/mL

kanamycin. Once the OD600 reached 0.6, the expression of proteins was

induced with 0.2 mM isopropyl-β-d-thiogalactoside (IPTG) overnight at 18 °C.

The bacterial cells were then harvested by centrifugation and subjected to

sonication. The cell lysate was then centrifuged at 13,400 × g for 20 min to

remove cell debris. GTD protein was purified using nickel affinity

chromatography and dialyzed in buffer containing 10% glycerol, 20 mM Tris (pH

7.5), and 150 mM NaCl. The GTD mutants were introduced into pET28a-GTD

using the Quick Change Site-Directed Mutagenesis Kit (Stratagene, San Diego,

CA, USA) with the primers listed in **Table S2.** The subsequent purification was

performed using the same procedure as described for wild-type GTD.

Cell rounding assay

Vero or HeLa cells were seeded in 96-well sterile plates using serum-free

DMEM overnight, and the compounds were preincubated with TcdB for 1 h.

After incubation, the TcdB-compound complex was added to cells for 1 h at

37 °C. The cell morphology was visualized under a light microscope.

LysoTracker assay

The pH of endosomes was determined by LysoTracker Red DND-99.

Caco-2 or THP-1 cells in complete DMEM were seeded in 96-well sterile plates

overnight. The media was changed to serum-free DMEM for 1 h, and CAPE

was preincubated with TcdB for 1 h, followed by addition to the cells and

incubation for 2 h. Baf-A1 served as a control. After incubation, LysoTracker

and Hoechst dyes were added and the cells were incubated for 1 h. Excess dye

was removed by changing the media, and alterations in lysosomal red intensity

were documented using a fluorescence microscope.

***In vitro* TcdB autoprocessing assay**

The self-cleavage ability of TcdB primarily depends on its cysteine protease

activity. CAPE (1-16 µg/mL) was preincubated with TcdB for 1 h, followed by

the addition of InsP$_6$ and incubation at 37 °C for 6 h. Subsequently, loading

buffer was added for SDS–PAGE analysis, and the results were visualized

using Coomassie Brilliant Blue staining.

Rac1 glucosylation assay

HEK293T cells were seeded into a 6-well plate. The next day, TcdB and

different concentrations of CAPE (4-32 µg/mL) were added to cells. After 2 h,

the cells were harvested, lysed using lysis buffer, and subjected to

centrifugation at 13,400 × g for 10 min to collect the supernatant. The collected

supernatant was then mixed with loading buffer and subjected to SDS–PAGE

followed by western blot analysis. The glycosylation level of Rac1 was

evaluated by determining the levels of non-glycosylated Rac1 (BD Biosciences,

Mississauga, ON) and total Rac1 (Abcam).

UDP-Glo™ UDP-glucose hydrolase assay

GTD activity was assessed according to the manufacturer's instructions. A

total volume of 16 µL, consisting of 100 nM GTD, various concentrations of

CAPE and glucosylation buffer, was prepared. Four microliters of UDP-glucose

at a final concentration of 25 µM was added to the mixture, which was then

allowed to react at room temperature for 15 min. To stop the reaction, 10 µL of

the reaction mixture was removed and placed in a white polystyrene 96-well

plate containing 10 µL of UDP detection reagent. The reaction took place at

room temperature for 1 h. The luminescence value was measured at a 750 ms

integral, and the GTD activity was determined based on these measurements.

Circular dichroism (CD) spectroscopy

CD spectra of TcdB, TcdB-CAPE, GTD, and GTD-CAPE were obtained

using a CD spectrophotometer (MOS-500; Bio-Logic, France). TcdB or GTD at

a concentration of 0.2 mg/mL was incubated with 32 µg/mL CAPE. Changes in

the secondary structure of TcdB or GTD were assessed at room temperature

using a quartz cuvette with an optical path length of 1 mm. The scan wavelength

range was set from 190 to 250 nm with a resolution of 0.2 nm and a bandwidth

of 1 nm. The secondary structures of TcdB, TcdB-CAPE, GTD, and GTD-CAPE

were analyzed using the BeStSel web server.

Molecular docking

CAPE was obtained as a ligand from PubChem, and its topology was

generated using AmeberTools. The crystal structure of GTD was retrieved from

the Protein Data Bank (PDB) and processed using AutoDockTools. Docking

calculations were performed using AutodockVina, and molecular dynamics

simulations were carried out using Gromacs 2020.6. The RMSD of the receptor

and ligand was monitored to assess conformational changes during the binding

process. The interaction energy was analyzed using the Poisson-Boltzmann

surface area (MM-PBSA) method, which is a molecular mechanics approach.

Flow cytometry

Flow cytometry was used to quantify the cellular uptake of TcdB via

endocytosis. To facilitate detection, full-length TcdB was labeled with Alexa

Fluor®488 NHS Ester (Yeasen Biotechnology Co., Ltd., Shanghai, China). Prior

to cell treatment, Alexa Fluor 488-TcdB was preincubated with CAPE at a

concentration of 32 µg/mL for 30 min. The treated solution was then added to

Caco-2 cells and incubated for 1 h at 37 °C. Following three washes with

phosphate-buffered saline (PBS), the Caco-2 cells were collected, and the

fluorescence signal was measured using flow cytometry.

Cytotoxicity assay

577 Vero or THP-1 cells were plated into 96-well plates. The next day, different

concentrations of CAPE were prepared by diluting them in DMEM and adding

them to the cells. After incubation at 37 °C for 3 h, the viability of cells was

assessed using a Cell Counting Kit-8 (CCK-8) (Beyotime Biotechnology Co.,

Shanghai, China) according to the manufacturer's instructions.

Biacore analysis

The affinity analysis of TcdB or GTD with CAPE was conducted using a

Biacore T100 device (Biacore Inc., Uppsala, Sweden) according to the

manufacturer's protocol. A CM5 chip (General Electric Company, GE) was

utilized to immobilize the TcdB or GTD proteins. Various concentrations of

CAPE were injected, and all steps were performed in running buffer consisting

of 5% DMSO in PBS-P20 (GE). Evaluation version 3.1 software provided by

Biacore and the 1:1 Langmuir combined model were employed for data analysis.

Mouse infection model

For the acute toxicity model, after pretreatment of TcdB with CAPE (2 or 4

μg/mL) for 1 h, the mice were intraperitoneally injected with a complex of TcdB-

solvent or TcdB-CAPE (30 ng of TcdB per mouse). The survival of mice was

observed for 72 h.

For the CDI model, mice were given cefoperazone (0.5 g/L) in their drinking

water for 5 days (from day -7 to day -2), followed by switching to sterilized water

for 2 days. On day 0, the mice were challenged with 1×10^8 colony-forming

units (CFUs) of *C. difficile* BAA-1870 (Ribotype 027) by oral gavage. For the

CAPE or melatonin treatment group, mice were treated with CAPE (30 mg/kg

by oral gavage) or melatonin (200 mg/kg by oral gavage) every day after *C.*

difficile challenge. The weight loss and fecal state of the mice were monitored

daily. Fecal samples were collected on days 0-3 to evaluate the stool water

content as a marker of diarrhea. Additionally, fecal samples on day 3 were

collected to assess *C. difficile* colonization. Colonic tissues were collected on

day 3 post infection, stained with hematoxylin and eosin (H&E) and scored as

previously described [60].

Fecal samples were collected on day 3 post infection, frozen in liquid

nitrogen and transferred to the laboratory with dry ice for 16S rRNA sequencing

and untargeted metabolomics analysis (Novogene Bioinformatics Technology

Co., Ltd., Tianjin, China). The melatonin level in mouse stool supernatant was

determined using the Mouse MT (Melatonin) ELISA Kit (Sangon Biotechnology

Co., Ltd., Shanghai, China).

Statistical analysis

Western blot densitometry and LysoTracker fluorescence intensity were

measured using ImageJ software (NIH, Bethesda, MD, USA). Statistical

analysis was performed using GraphPad Prism 9.0 (GraphPad Software, USA).

Unpaired two-tailed Student's t tests were used for comparing two groups, and

one-way analysis of variance (ANOVA) was used for analyzing data from more

than three groups. The statistical significance was determined using the log-

rank (Mantel–Cox) test (survival rates) and Mann–Whitney U test (tissue

bacterial load) *in vivo*. All the data are expressed as the mean ± standard

deviation (SD). * indicates $p < 0.05$ and ** indicates $p < 0.01$.

Acknowledgments

The authors thank Professor Aiwu Wu (Guangzhou Medical University) for

the *C. difficile* BAA-1870 strain and Professor Hanping Feng (Tufts University)

for the pHis1522-TcdB plasmid. This work was supported by The National Key

Research & Development Program of China (2021YFD1801000), the Natural

Science Foundation of Jilin Province (20230101142JC and 20240101282JC),

and The Fundamental Research Funds for the Central Universities.

Author contributions

Y Guo., LQ He and JZ Qiu designed the study and wrote the manuscript.

Y Guo, Y Zhang, GZ Wang and HT Liu conducted experiments. JF Wang, XM

Deng, LQ He and JZ Qiu supervised the study and revised the manuscript. All

authors read and approved the final manuscript. JZ Qiu and LQ He are

responsible for the overall content as the guarantor.

Disclosure statement

No potential conflict of interest was reported by the author (s).

Data availability

All 16S rRNA gene sequencing data generated and analyzed during the

current study were deposited in the Sequence Read Archive (SRA) under the

BioProject accession number PRJNA1100298. Untargeted metabolomics data

are available via Metabolights with the identifier MTBLS9938.

References

1. Smits WK, Lyras D, Lacy DB, Wilcox MH, Kuijper EJ. Clostridium difficile infection. *Nat Rev Dis Primers* **2**, 16020 (2016).

2. Miller BA, Chen LF, Sexton DJ, Anderson DJ. Comparison of the burdens of hospital-onset, healthcare facility-associated Clostridium difficile Infection and of healthcare-associated infection due to methicillin-resistant Staphylococcus aureus in community hospitals. *Infect Control Hosp Epidemiol* **32**, 387-390 (2011).

3. Kelly CP, LaMont JT. Clostridium difficile--more difficult than ever. *N Engl J Med* **359**, 1932-1940 (2008).

4. Dubberke E. Clostridium difficile infection: the scope of the problem. *J Hosp Med* **7 Suppl 3**, S1-4 (2012).

5. Lessa FC*, et al.* Burden of Clostridium difficile infection in the United States. *N Engl J Med* **372**, 825-834 (2015).

6. Goorhuis A*, et al.* Emergence of Clostridium difficile infection due to a new hypervirulent strain, polymerase chain reaction ribotype 078. *Clin Infect Dis* **47**, 1162-1170 (2008).

7. He M*, et al.* Emergence and global spread of epidemic healthcare-associated Clostridium difficile. *Nat Genet* **45**, 109-113 (2013).

8. Clements AC, Magalhaes RJ, Tatem AJ, Paterson DL, Riley TV. Clostridium difficile PCR ribotype 027: assessing the risks of further worldwide spread. *Lancet Infect Dis* **10**, 395-404 (2010).

9. Cornely OA*, et al.* Fidaxomicin versus vancomycin for infection with Clostridium difficile in Europe, Canada, and the USA: a double-blind, non-inferiority, randomised controlled trial. *Lancet Infect Dis* **12**, 281-289 (2012).

10. Bingley PJ, Harding GM. Clostridium difficile colitis following treatment with metronidazole and vancomycin. *Postgrad Med J* **63**, 993-994 (1987).

11. Feuerstadt P, Theriault N, Tillotson G. The burden of CDI in the United States: a multifactorial challenge. *BMC Infect Dis* **23**, 132 (2023).

12. Hopkins RJ, Wilson RB. Treatment of recurrent Clostridium difficile colitis: a narrative review. *Gastroenterol Rep (Oxf)* **6**, 21-28 (2018).

13. Spigaglia P, Mastrantonio P, Barbanti F. Antibiotic Resistances of Clostridium difficile. *Adv Exp Med Biol* **1050**, 137-159 (2018).

14. Seekatz AM, Young VB. Clostridium difficile and the microbiota. *J Clin Invest* **124**, 4182-4189 (2014).

15. Paredes-Sabja D, Shen A, Sorg JA. Clostridium difficile spore biology: sporulation, germination, and spore structural proteins. *Trends Microbiol* **22**, 406-416 (2014).

16. Lyras D*, et al.* Toxin B is essential for virulence of Clostridium difficile. *Nature* **458**, 1176-1179 (2009).

17. Kuehne SA, Cartman ST, Heap JT, Kelly ML, Cockayne A, Minton NP. The role of toxin A and toxin B in Clostridium difficile infection. *Nature* **467**, 711-713 (2010).

18. Carter GP*, et al.* Defining the Roles of TcdA and TcdB in Localized Gastrointestinal Disease, Systemic Organ Damage, and the Host Response during Clostridium difficile Infections. *mBio* **6**, e00551 (2015).

19. Drudy D, Fanning S, Kyne L. Toxin A-negative, toxin B-positive Clostridium difficile. *Int J Infect Dis* **11**, 5-10 (2007).

20. Chen P*, et al.* Structure of the full-length Clostridium difficile toxin B. *Nat Struct Mol Biol* **26**, 712-719 (2019).

21. Orrell KE, Melnyk RA. Large Clostridial Toxins: Mechanisms and Roles in Disease. *Microbiol Mol Biol Rev* **85**, e0006421 (2021).

22. Rasko DA, Sperandio V. Anti-virulence strategies to combat bacteria-mediated disease. *Nat Rev Drug Discov* **9**, 117-128 (2010).

23. Tam J, Beilhartz GL, Auger A, Gupta P, Therien AG, Melnyk RA. Small molecule inhibitors of Clostridium difficile toxin B-induced cellular damage. *Chem Biol* **22**, 175-185 (2015).

24. Balaha M, De Filippis B, Cataldi A, di Giacomo V. CAPE and Neuroprotection: A Review. *Biomolecules* **11**, (2021).

25. Garland M*, et al.* The Clinical Drug Ebselen Attenuates Inflammation and Promotes Microbiome Recovery in Mice after Antibiotic Treatment for CDI. *Cell Rep Med* **1**, (2020).

26. Li X, *et al.* A strain of Bacteroides thetaiotaomicron attenuates colonization of Clostridioides difficile and affects intestinal microbiota and bile acids profile in a mouse model. *Biomed Pharmacother* **137**, 111290 (2021).

27. Kienesberger B, *et al.* (S)-Reutericyclin: Susceptibility Testing and In Vivo Effect on Murine Fecal Microbiome and Volatile Organic Compounds. *Int J Mol Sci* **22**, (2021).

28. Khanna S, *et al.* Gut microbiome predictors of treatment response and recurrence in primary Clostridium difficile infection. *Aliment Pharmacol Ther* **44**, 715-727 (2016).

29. Dong D, *et al.* Effects of intestinal colonization by Clostridium difficile and Staphylococcus aureus on microbiota diversity in healthy individuals in China. *BMC Infect Dis* **18**, 207 (2018).

30. Gurung B, Stricklin M, Wang S. Gut Microbiota-Gut Metabolites and Clostridioides difficile Infection: Approaching Sustainable Solutions for Therapy. *Metabolites* **14**, (2024).

31. Cronstein BN. Adenosine, an endogenous anti-inflammatory agent. *J Appl Physiol (1985)* **76**, 5-13 (1994).

32. Yong T, *et al.* Caffeic acid phenethyl ester alleviated hypouricemia in hyperuricemic mice through inhibiting XOD and up-regulating OAT3. *Phytomedicine* **103**, 154256 (2022).

33. de Araujo Junqueira AF, *et al.* Adenosine deaminase inhibition prevents Clostridium difficile toxin A-induced enteritis in mice. *Infect Immun* **79**, 653-662 (2011).

34. Neumann-Schaal M, Jahn D, Schmidt-Hohagen K. Metabolism the Difficile Way: The Key to the Success of the Pathogen Clostridioides difficile. *Front Microbiol* **10**, 219 (2019).

35. Bouillaut L, Self WT, Sonenshein AL. Proline-dependent regulation of Clostridium difficile Stickland metabolism. *J Bacteriol* **195**, 844-854 (2013).

36. Jackson S, Calos M, Myers A, Self WT. Analysis of proline reduction in the nosocomial pathogen Clostridium difficile. *J Bacteriol* **188**, 8487-8495 (2006).

37.   Sutton SS, Magagnoli J, Cummings TH, Hardin JW. Melatonin as an Antimicrobial Adjuvant and Anti-Inflammatory for the Management of Recurrent Clostridioides difficile Infection. *Antibiotics (Basel)* **11**, (2022).

38.   2019 AR Threats Report. (2019). [Available from: https://www.cdc.gov/drugresistance/biggest-threats.html].

39.   Abbas A, Zackular JP. Microbe-microbe interactions during Clostridioides difficile infection. *Curr Opin Microbiol* **53**, 19-25 (2020).

40.   Raplee I*, et al.* Emergence of nosocomial associated opportunistic pathogens in the gut microbiome after antibiotic treatment. *Antimicrob Resist Infect Control* **10**, 36 (2021).

41.   Beilhartz GL, Tam J, Melnyk RA. Small Molecules Take A Big Step Against Clostridium difficile. *Trends Microbiol* **23**, 746-748 (2015).

42.   Ranftler C*, et al.* Binding and neutralization of C. difficile toxins A and B by purified clinoptilolite-tuff. *PLoS One* **16**, e0252211 (2021).

43.   Zucca M, Scutera S, Savoia D. Novel avenues for Clostridium difficile infection drug discovery. *Expert Opin Drug Discov* **8**, 459-477 (2013).

44.   Kuehne SA, Collery MM, Kelly ML, Cartman ST, Cockayne A, Minton NP. Importance of toxin A, toxin B, and CDT in virulence of an epidemic Clostridium difficile strain. *J Infect Dis* **209**, 83-86 (2014).

45.   Warny M*, et al.* Toxin production by an emerging strain of Clostridium difficile associated with outbreaks of severe disease in North America and Europe. *Lancet* **366**, 1079-1084 (2005).

46.   Stewart D, Anwar F, Vedantam G. Anti-virulence strategies for Clostridioides difficile infection: advances and roadblocks. *Gut Microbes* **12**, 1802865 (2020).

47.   Deeks ED. Bezlotoxumab: A Review in Preventing Clostridium difficile Infection Recurrence. *Drugs* **77**, 1657-1663 (2017).

48.   Bender KO*, et al.* A small-molecule antivirulence agent for treating Clostridium difficile infection. *Sci Transl Med* **7**, 306ra148 (2015).

49.   Chan H*, et al.* Vitamin D(3) and carbamazepine protect against

Clostridioides difficile infection in mice by restoring macrophage lysosome acidification. *Autophagy* **18**, 2050-2067 (2022).

50. Tam J*, et al.* Host-targeted niclosamide inhibits C. difficile virulence and prevents disease in mice without disrupting the gut microbiota. *Nat Commun* **9**, 5233 (2018).

51. Tam J*, et al.* Intestinal bile acids directly modulate the structure and function of C. difficile TcdB toxin. *Proc Natl Acad Sci U S A* **117**, 6792-6800 (2020).

52. Tek NA, Senture SA, Ersoy N. Is Propolis a Potential Anti-Obesogenic Agent for Obesity? *Curr Nutr Rep*, (2024).

53. Schubert AM, Sinani H, Schloss PD. Antibiotic-Induced Alterations of the Murine Gut Microbiota and Subsequent Effects on Colonization Resistance against Clostridium difficile. *mBio* **6**, e00974 (2015).

54. van Nood E*, et al.* Duodenal infusion of donor feces for recurrent Clostridium difficile. *N Engl J Med* **368**, 407-415 (2013).

55. Lee C*, et al.* Safety of fecal microbiota, live-jslm (REBYOTA()) in individuals with recurrent Clostridioides difficile infection: data from five prospective clinical trials. *Therap Adv Gastroenterol* **16**, 17562848231174277 (2023).

56. Gonzales-Luna AJ, Carlson TJ, Garey KW. Review Article: Safety of Live Biotherapeutic Products Used for the Prevention of Clostridioides difficile Infection Recurrence. *Clin Infect Dis* **77**, S487-S496 (2023).

57. Cavalcante IC*, et al.* Effect of novel A2A adenosine receptor agonist ATL 313 on Clostridium difficile toxin A-induced murine ileal enteritis. *Infect Immun* **74**, 2606-2612 (2006).

58. Ma N, Zhang J, Reiter RJ, Ma X. Melatonin mediates mucosal immune cells, microbial metabolism, and rhythm crosstalk: A therapeutic target to reduce intestinal inflammation. *Med Res Rev* **40**, 606-632 (2020).

59. Yang G*, et al.* Expression of recombinant Clostridium difficile toxin A and B in Bacillus megaterium. *BMC Microbiol* **8**, 192 (2008).

60. Reeves AE, Theriot CM, Bergin IL, Huffnagle GB, Schloss PD, Young VB. The interplay between microbiome dynamics and pathogen dynamics in a murine model of Clostridium difficile Infection. *Gut*

868   *Microbes* **2**, 145-158 (2011).
869

Figures

Figure 1. CAPE suppresses TcdB-mediated cell rounding.

(A) Schematic of the process used for the identification of small molecules that

inhibit the cell rounding phenotype induced by TcdB. TcdB (0.2 ng/mL) was

preincubated with 8 µg/mL of individual compounds prior to addition to the cell

cultures. **(B)** Active compounds were identified from the chemical library

consisting of 2076 chemicals. The red dots represent caffeic acid and its

derivatives. **(C)** Dose-dependent CAPE inhibition of TcdB-induced rounding of

Vero (upper) and HeLa (lower) cells. Scale bar, 100 µm. The dose titration

curves are presented on the right. **(D)** The survival rates of mice that were

intraperitoneally injected with CAPE-pretreated or native TcdB (30 ng). * $p < 0.05$

and ** $p < 0.01$ by log-rank (Mantel–Cox) test.

Figure 2. Mechanism by which CAPE suppresses TcdB intoxication.

(A) Schematic illustration of the multistep mechanism of TcdB intoxication of host cells. **(B)** Determination of internalized TcdB levels. Alexa Fluor 488-labeled TcdB was incubated with 32 µg/mL CAPE for 30 min before it was added to the cell culture. The cellular uptake of TcdB was measured by flow cytometry analysis. **(C)** Lysosomal activity in host cells treated with CAPE was measured via LysoTracker staining. The nuclei were stained with DAPI. Bar, 100 µm. **(D)** The intensities of LysoTracker staining were quantified using ImageJ. * p <0.05; ** p <0.01 using one-way ANOVA. **(E-F)** CAPE inhibits $InsP_6$-induced autoprocessing of full-length TcdB. Two micrograms of TcdB was preincubated with increasing concentrations of CAPE for 1 h. Self-cleavage of TcdB was initiated by the addition of 3 µg of $InsP_6$ and the cells were incubated for 6 h at 37 °C. Reactions were terminated with 5x SDS loading buffer, and the release of GTD was visualized by CBB staining of the SDS–PAGE gel **(E)**. The level of released GTD in panel E was quantified by ImageJ **(F)** * p <0.05; ** p

<0.01 using one-way ANOVA. **(G)** Dose-dependent inhibition of the GTD

activity of TcdB by CAPE. Recombinant GTD of TcdB was mixed with different

concentrations of CAPE (0-32 µg/mL) and incubated for 15 min at 25 ℃. GTD

activity was measured by the UDP-Glo™ glycosyltransferase assay. **(H-I)**

Intracellular Rac1 glucosylation in cells treated with TcdB. Native or CAPE-

pretreated TcdB was added to the cell culture and cells were incubated for 2 h.

The cells were lysed, and the glucosylation of Rac1 was analyzed by western

blotting using antibodies recognizing total Rac1 and unglycosylated Rac1 **(H)**.

The level of Rac1 glucosylation shown in H was quantified by ImageJ **(I)**. * *p*

<0.05; ** *p* <0.01 using one-way ANOVA.

Figure 3. CAPE alters the secondary structure of TcdB and GTD.

(A-B) The secondary structure of TcdB **(A)** or GTD **(B)** in the presence or absence of 32 μg/mL CAPE was determined by CD spectroscopy. **(C-D)** Calculated CD spectra of CAPE-treated TcdB **(C)** and GTD **(D)**. The wavelength for CD spectroscopy was set at 190-250 nm.

Figure 4. Direct interaction of CAPE with TcdB and GTD.

(A-B) Full-length TcdB or GTD was immobilized on a CM5 chip, followed by the

injection of various concentrations of CAPE. The response unit (RU) value and

affinities of CAPE interactions with TcdB **(A)** and GTD **(B)** were determined by

a Biacore T100.

Figure 5. Asp286 and Trp520 in the GTD are critical for CAPE-mediated

inhibition of GTD.

(A) Three-dimensional structure determination of GTD with the CAPE complex

was performed by molecular modeling. **(B)** The RMSD values of the GTD-

CAPE complex. **(C)** Decomposition of the binding energy on a per-residue basis

in the binding sites of the GTD-CAPE complex. **(D)** Glucosyltransferase activity

of wild-type GTD and the GTD mutants. **(E-K)** Residual activity of wild-type

GTD **(E)** and its mutants GTD$_{L265A}$ **(F)**, GTD$_{T465A}$ **(G)**, GTD$_{I466A}$ **(H)**, GTD$_{L519A}$ **(I)**,

GTD$_{W520A}$ **(J)** and GTD$_{D286A}$ **(K)** in the presence of increasing concentrations of

CAPE (0-4 µg/mL). * p <0.05; ** p <0.01 using one-way ANOVA.

Figure 6. CAPE treatment blocks the pathology of CDI in mice.

(A) Schematic of the process used for the establishment of the CDI mouse

model. **(B)** Weight changes in CDI mice after CAPE or vehicle treatment (n=10).

(C) The fecal dry/wet weight ratio in CDI mice treated with CAPE or vehicle

(n=6). * p <0.05; ** p <0.01 using one-way ANOVA. **(D)** Bacterial colonization

in the feces of CDI mice treated with CAPE or vehicle (n=15). * p <0.05; ** p

<0.01 using the Mann–Whitney U test. **(E)** Pathological analysis of colon

sections from CDI mice treated with CAPE or vehicle. The black boxes are the

regions that were magnified and are shown underneath. **(F)** Histopathological

scores of vehicle-treated and CAPE-treated mice in terms of cell infiltration

(orange), epithelial damage (purple), and edema (green). * p <0.05; ** p <0.01

949   using one-way ANOVA.

Figure 7. The influence of CAPE treatment on the diversity and composition of the gut microbiota in CDI mice.

(A) Venn diagram illustrating the numbers of differentially abundant bacteria identified using a single model and their cooperation. **(B-E)** Quantification of the α diversity of the gut microbiota according to the observed species richness **(B-C),** Shannon index **(D)**, and Simpson index **(E)**. **(F)** PCoA showing β diversity based on the weighted UniFrac distance. **(G-H)** Relative abundance of the top 10 predominant bacteria classified at the phylum **(G)** and genus **(H)** levels. **(I)** Significant changes in flora among the model and CAPE groups, as measured by LEfSe analysis (LDA score [log 10] > 3.5). **(J-M)** Relative abundances of *Bacteroides* **(J)**, *Muribaculum* **(K)**, *Faecalibaculum* **(L)**, and *Parasutterella* **(M)** in mice in the model and CAPE treatment groups. * *p* <0.05; ** *p* <0.01 using an unpaired Student's t test.

Figure 8. Alterations in gut metabolites in CAPE-treated mice with CDI.

(A) The score plots show orthogonal partial least square discriminant analysis

(PLS-DA) results. **(B)** Volcano plot representing the upregulated or

downregulated gut metabolites in the CAPE treatment group compared to those

in the model group. **(C)** Enrichment bubble map of the top 20 metabolic

pathways in the CAPE treatment and model groups. **(D)** The 26 potential

biomarkers are displayed in a heatmap, along with hierarchical clustering

analysis.

Figure 9. Oral administration of melatonin alleviates the pathology of CDI.

(A) The impact of upregulated metabolites on the cytotoxicity of TcdB. TcdB (0.2

ng/mL) was incubated with each of the metabolites (100 µg/mL) for 1 h at 37

°C before being added to the Vero cell culture. At 3 h after TcdB treatment, cell

death was evaluated with a CCK-8 kit. **(B)** Body weight changes in melatonin-

treated CDI mice (n=5). **(C)** The fecal dry/wet weight ratio in CDI mice after

melatonin treatment (n=5). **(D)** Histopathological analysis of colon tissue from

982   CDI mice treated with melatonin. **(E)** Histopathological scores of melatonin-

983   treated CDI mice. * p <0.05; ** p <0.01 using one-way ANOVA.

985

Figure 10. A model for CAPE-mediated suppression of TcdB cytotoxicity

and protection of mice from CDI.

The intoxication of cells by TcdB involves multistep mechanisms. Direct

interaction of CAPE with full-length TcdB blocks $InsP_6$-induced autoproteolysis,

resulting in decreased production of active GTD. The binding between GTD and

CAPE inhibits its glucosyltransferase activity, thus decreasing the glucosylation

level of Rac1. CAPE treatment of CDI mice markedly restored the diversity and

composition of the gut microbiota and induced changes in gut metabolites,

which might have contributed to the therapeutic outcomes of CDI following

CAPE treatment.

A TcdB

N — GTD — CPD — DRBD — CROPs — C

1. Binding
2. Endocytosis
3. Endosome
4. Membrane insertion
5. Autoprocessing
6. Rho inactivation
7. Pathogenic effects

Extracellular

Target cell

B

Count

Alexa-flour488-TcdB

Ctrl
TcdB
TcdB-CAPE

C

Lysotracker | DAPI | Merge

Blank
TcdB
Baf-A1
CAPE (8 μg/mL)
CAPE (32 μg/mL)

D

Intensity of LysoTracker

Ctrl   BafA1   TcdB   CAPE (8 μg/mL)   CAPE (32 μg/mL)

E

TcdB	+	+	+	+	+	+	+
InsP$_6$	–	+	+	+	+	+	+
CAPE	–	–	1	2	4	8	16 (μg/mL)

TcdB full-length
TcdB processed
"Released GTD"

Coomassie Blue

F

Released GTD

Ctrl   0   1   2   4   8   16

CAPE (μg/mL)

G

GTD activity (%)

0   2   4   8   16   32

CAPE (μg/mL)

H

| TcdB | – | + | + | + | + | + |
| CAPE | – | – | 4 | 8 | 16 | 32 (μg/mL) |

40
35
25
15

Unglycosylated Rac1

IB: Unglycosylated Rac1

| TcdB | – | + | + | + | + | + |
| CAPE | – | – | 4 | 8 | 16 | 32 (μg/mL) |

40
35
25
15

Total Rac1

IB: Total Rac1

I

Unglycosylated Rac1/ Total Rac1

Ctrl   0   4   8   16   32

CAPE (μg/mL)

A *C. difficile* challenge and treatment of CAPE (30 mg/kg; intragastrical administration; every day)

C57BL/6 6-8 weeks

Cefoperazone in drinking water (0.5 g/L)

Day -7

Day -2 Change to sterile water

Day 0

Day 3 Sampling

Day 5 Endpoint monitoring

B Weight (percent of day 0)

Ctrl · Model · CAPE

n = 10

Days

C % of water in stool

Ctrl · Model · CAPE

Days

D CFUs/g (log10)

Model CAPE

E

Ctrl   Model   CAPE

* Cellular infiltration   * Epithelial damage   * Edema

F Histopathological scores

Cellular infiltration
Epithelial damage
Edema

Ctrl   Model   CAPE

A THP-1

Cell viability (%) vs CAPE (μg/mL)

B Vero

Cell viability (%) vs CAPE (μg/mL)