Blocking toxin function and modulating the gut microbiota: caffeic acid phenethyl ester as a potential treatment for *Clostridioides difficile* infection

Yan Guo[1#], Yong Zhang[2#], Guizhen Wang[3], Hongtao Liu[1], Jianfeng Wang[1], Xuming Deng[1], Liuqing He[4*], Jiazhang Qiu[1*]

[1] State Key Laboratory for Diagnosis and Treatment of Severe Zoonotic Infectious Diseases, Key Laboratory for Zoonosis Research of the Ministry of Education, College of Veterinary Medicine, Jilin University, Center for Pathogen Biology and Infectious Diseases, The First Hospital of Jilin University, Changchun, China.

[2] Center for Pathogen Biology and Infectious Diseases, State Key Laboratory for Zoonotic Diseases, The First Hospital of Jilin University, Changchun 130021, China.

[3] Measurement Biotechnique Research Center, College of Biological and Food Engineering, Jilin Engineering Normal University, Changchun 130052, China.

[4] Hunan Provincial Key Laboratory of Animal Intestinal Function and Regulation, Hunan International Joint Laboratory of Animal Intestinal Ecology and Health, Laboratory of Animal Nutrition and Human Health, College of Life Sciences, Hunan Normal University, Changsha, China.

#YG and YZ contributed equally to this work.

23 [*]Correspondence: Jiazhang   Qiu   (qiujz@jlu.edu.cn);   Liuqin   He

24 (heliuqin@hunnu.edu.cn)

Abstract

Clostridioides difficile infection (CDI) is the leading cause of hospital-acquired diarrhea that seriously threatens public health. The disruption of normal gut microbiota by the use of broad-spectrum antimicrobial agents enables C. difficile to proliferate in the colon. The emergence and prevalence of hypervirulent C. difficile strains result in increased morbidity, mortality, and high recurrence rates of CDI, thus creating a pressing need for novel therapeutics. The multi-domain toxins TcdA and TcdB are the primary determinants of CDI pathogenesis, rendering them ideal drug targets in the anti-virulence paradigm. In this study, we identified caffeic acid and its derivatives from natural compounds library as active inhibitors of TcdB via a cell-based high-throughput phenotypic screening. Further mechanistic investigations revealed that caffeic acid phenethyl ester (CAPE) could directly bind to TcdB, thus suppressing InsP$_6$-induced autoproteolysis and inhibiting glucosyltransferase activity. CAPE treatment remarkably reduces the pathology of CDI in a murine infection model in terms of alleviated diarrhea symptoms, decreased bacterial colonization, and relieved histopathological lesions. Moreover, CAPE treatment of *C. difficile*-challenged mice induces a remarkable increase in the diversity and composition of the gut microbiota (e.g., *Bacteroides* spp.) and alterations of gut metabolites (e.g., adenosine, D-proline, and melatonin), which might partially contribute to the therapeutic outcomes of CAPE against CDI. Our results reveal the potential of CAPE as a therapeutic

48   for the management of CDI, or CAPE might serve as a lead compound for the

49   development of antivirulence drugs targeting TcdB.

51   **Keywords:** *Clostridioides difficile*; toxin B; caffeic acid phenethyl ester; gut

52   microbiota; gut metabolites

Introduction

Clostridioides difficile (*C. difficile*) is an obligate anaerobic, spore-forming,

gram-positive bacterial pathogen that is the most common cause of

hospital-acquired and antibiotic-associated gastrointestinal diseases, including

mild diarrhea and life-threatening pseudomembranous colitis [1, 2, 3]. In the

United States, *C. difficile* is responsible for almost half a million infections and

approximately 29,000 deaths per year, resulting in an estimated annual

healthcare-associated cost of up to $4.8 billion [4, 5]. Importantly, the incidence,

severity, mortality and healthcare costs of *C. difficile* infection (CDI) have

dramatically increased in the 21st century, at least partially owing to the

frequent use of antimicrobial agents and the emergence of hypervirulent

strains (e.g., PCR ribotypes 027 and 078) [6, 7, 8]. Limited options are available

for the treatment of CDI, while antibiotics including metronidazole, fidaxomicin

and vancomycin, are the mainstays of clinical practice [9, 10]. However, antibiotic

treatment is ineffective in more than 35% of CDI patients and can lead

infection recurrence [11, 12]. Moreover, the rapid evolution of antibiotic resistance

in *C. difficile* has further compromised the therapeutic efficacy of current

treatment options [13]. Therefore, the high morbidity and mortality, high

recurrence rates, and emergence of multiple antibiotic-resistant isolates have

made this pathogen a major threat to public health; these factors highlight the

urgent need for novel therapies.

C. difficile widely exists in the environment and in the intestinal tract of

humans and animals and is mainly transmitted via the fecal-oral route [14]. The

pathogenesis of CDI is correlated with the use of antibiotics, which impair the

structure and composition of the normal gut microbiota, thus allowing *C.*

difficile to proliferate in the colon [14]. Therefore, previous antibiotic therapy is

the major risk factor for the development of CDI. In the colon, the dormant

spores of *C. difficile* germinate into oxygen-sensitive vegetative cells that

produce toxins and cause symptoms of disease [15]. *C. difficile* secretes two

large clostridial exotoxins (TcdA and TcdB), which are directly responsible for

the clinical pathology of CDI [16, 17]. Epidemiological studies suggest that most

disease-causing *C. difficile* strains have genes that encode TcdB [18, 19]. TcdA

and TcdB have identical functional domains: a N-terminal glucosyltransferase

catalytic domain (GTD), a cysteine protease active domain (CPD), a

transmembrane domain (DRBD), and a C-terminal receptor binding domain

(CROPs) [20]. When secreted into the colon, both toxins are internalized by host

cells through receptor-mediated, clathrin-dependent endocytosis. Following a

decrease in endosomal pH, the toxins undergo conformational changes in the

DRBD, resulting in insertion of the toxin into the endosome membrane, pore

formation, and the release of the GTD and CPD domains into the cytosol. Then,

CPD binding with cytosolic inositol hexakisphosphate ($InsP_6$) activates its

protease activity, leading to the cleavage and release of the GTD. Eventually,

the GTD utilizes UDP-glucose to glucosylate Rho and/or Ras family GTPases,

thus disrupting GTPase signaling and resulting in deleterious effects, including

cytopathic "rounding", dysregulation of the actin cytoskeleton, apoptosis, and

pyroptosis [1]. Although both toxins are cytotoxic to host cells, TcdB shows

greater toxicity than TcdA [21].

Disruption of the gut microbiota by antimicrobial agents results in high

recurrence rates, which limits the application of antimicrobial therapy for CDI [14].

Consequently, there is a critical need to develop nonantibiotic treatments for

CDI, either as standalone or augmented therapies. Importantly, strategies

targeting bacterial virulence factors, including toxins, are termed anti-virulence

strategies and have attracted interest in the 21st century [22]. Owing to its

essential role in *C. difficile* pathogenesis, TcdB has become an ideal target for

the development of anti-virulence drugs [23]. In this study, caffeic acid and a

series of its derivatives were found to inhibit TcdB-mediated cell rounding via

cell-based screening. Among these potential inhibitors, caffeic acid phenethyl

ester (CAPE) had the greatest inhibitory effect. Mechanistically, CAPE bound

to both full-length TcdB and the GTD fragment, leading to alterations in the

secondary structure. This interaction suppressed the $InsP_6$-mediated

self-cleavage of TcdB, as well as the enzymatic activity of GTD, thereby

decreasing the level of glucosylated Rac1 in TcdB-treated host cells and

further inhibiting the cytopathology and cytotoxicity of TcdB. Oral

administration of CAPE significantly protected against CDI in models of animal

infection. In addition to direct toxin inhibition, the therapeutic effect may also be

attributed to CAPE-induced alteration of the gut microbiota (e.g., *Bacteroides*

spp.) as well as gut metabolites (e.g., adenosine, D-proline, and melatonin).

Overall, our study revealed that caffeic acid and its derivatives, particularly

CAPE, may serve as promising lead compounds for the development of

anti-virulence drugs to treat CDI.

Results

CAPE inhibits cell rounding induced by TcdB

TcdB can inactivate Rho GTPases via glucosylation, leading to the loss of

cytoskeletal structure and cell-rounding. Consequently, we utilized this

phenotype to identify potential TcdB inhibitors from 2076 chemicals, including

1515 FDA-approved drugs and 561 natural compounds. The minimal

concentration of recombinant His_6-TcdB (0.2 ng/mL) that induced 100% cell

rounding within 1 h was used for drug screening. This concentration of TcdB

was preincubated with 8 µg/mL of the individual compounds for 1 h, the

compounds that were not bound to TcdB were removed and then added to

added to Vero cell cultures. At 1 h post-TcdB treatment, cell morphology was

visualized under a light microscope (**Figure 1A**). These analyses allowed us to

identify 22 candidate compounds with potent inhibitory effects (**Figure 1B**).

Interestingly, six out of the 22 compounds were caffeic acid derivatives. The

natural compound library contained eight caffeic acid derivatives, of which

methyl caffeic acid and ferulic acid displayed no efficacy. Caffeic acid itself only

slightly inhibited TcdB-induced cell rounding, with a 50% inhibitory

concentration (IC_{50}) of 527.5 µg/mL (1.86 mM) (**Table S1**). Notably, except for

rosmarinic acid, the IC_{50} values for the other screened caffeic acid derivatives

were less than 6 µg/mL, suggesting that these compounds potently inhibited

TcdB-mediated toxicity (**Table S1**). CAPE is the phenethyl alcohol ester of

caffeic acid and a main active component of propolis with broad

pharmacological properties, such as anticancer, anti-inflammatory, antiviral,

antibacterial, antifungal, antioxidant, and immunomodulatory activities [24]. As

CAPE has the lowest IC_{50} (3.0 µg/mL) among the active caffeic acid

derivatives, it was selected as the representative compound for subsequent

investigations (**Table S1**). After preincubation with increasing concentrations of

CAPE, dose-dependent inhibition of cell rounding was observed in both Vero

and HeLa cells (**Figure 1C**). The protective influence of CAPE against

TcdB-induced Vero cell death remained evident even when the treatment

duration was prolonged to 12 h (**Figure S1A-B**).

In addition, we also determined whether inhibition of TcdB by CAPE could

lead to loss of toxin function *in vivo* using a murine systemic intoxication model.

A lethal dose of recombinant His_6-TcdB (30 ng) was preincubated with either

DMSO or CAPE (2 or 4 µg/mL) for 1 h at 37 ^{o}C, followed by intraperitoneal

injection into mice. Notably, all mice in the DMSO group (n=10) died within 36

h after toxin injection. However, 100% of mice pretreated with CAPE 4 (µg/mL)

toxin survived at the monitoring endpoint (72 h) (n=10) (**Figure 1D**).

Furthermore, we evaluated the efficacy of CAPE following an intraperitoneal

injection by administering it 2 h prior to the injection of TcdB, in order to gauge

the survival rate of the mice. The outcomes revealed that a 2 h pretreatment

with CAPE at doses of 30 and 50 mg/kg significantly enhanced the mice's

protection rate by 57.1% (**Figure S2)**. Simultaneously, we assessed the impact

of CAPE on *C.difficile* itself. CAPE markedly repressed *tcdB* gene expression

at a concentration of 16 μg/mL (**Figure S3A**) and inhibited the growth and

spore formation of *C. difficile* BAA-1870 at a concentration of 32 μg/mL

(**Figure S3B-C**). Taken together, these data indicate that CAPE is a potent

toxin inhibitor of TcdB and may serve as a promising antimicrobial agent.

CAPE decreases InsP$_6$-induced autoproteolysis of TcdB and inhibits the

enzymatic activity of GTD

TcdB is a multi-domain containing toxin that induces toxicity in host cells

via an intricate and multistep mechanism. This mechanism involves (i)

receptor-mediated endocytosis, (ii) pore formation and subsequent membrane

translocation, (iii) autoproteolysis and release of GTD, and (iv) the

glucosylation of small GTPases (**Figure 2A**). To determine the mechanism by

which CAPE suppresses the cytotoxicity of TcdB, we first labeled full-length

TcdB with Alexa Fluor 488; then, the concentration of the endocytosed toxin

was determined via flow cytometry. Apparently, incubation of CAPE with TcdB

did not significantly affect its entry into host cells (**Figure 2B**). Next, we stained

CAPE-treated cells with LysoTracker, a fluorescent dye that accumulates in

the acidic organelles of live cells. In contrast to the vacuolar-type H+-ATPase

(V-ATPase) inhibitor bafilomycin A1 (Baf-A1), CAPE treatment did not alter the

intensity of LysoTracker fluorescence staining in either THP-1 or Caco-2 cells,

suggesting that inactivation of TcdB by CAPE is not due to changes in the pH

of endosomal compartments (**Figure 2C-D** and **Figure S4A-B**).

These observations indicate that CAPE might exert its effects after

membrane translocation. Indeed, preincubation of full-length TcdB with CAPE

blocked the $InsP_6$-induced autoproteolysis of TcdB, as revealed by the gradual

decrease in the production of released GTP, followed by the addition of

increasing amounts of CAPE *in vitro* (**Figure 2E-F**). Furthermore, we

determined the glucosyltransferase activity using recombinant GTD in the

absence or presence of different doses of CAPE. Strikingly, incubation with 2

µg/mL CAPE resulted in a significant reduction in GTD activity, whereas 32

µg/mL CAPE compromised 87.5% of the enzymatic activity (**Figure 2G**).

Importantly, in response to suppression of $InsP_6$-induced autoproteolysis and

GTD function, treatment of TcdB-challenged cells with CAPE markedly

decreased the level of glucosylated Rac1 in a dose-dependent manner

(**Figure 2H-I**). Taken together, these results suggest that CAPE disrupts

$InsP_6$-induced autoproteolysis of TcdB and inhibits the enzymatic activity of

GTD, thereby decreasing the glucosylation of Rac1 and alleviating cell

rounding.

CAPE alters the secondary structure of full-length TcdB and GTD

To further clarify the mechanisms used by CAPE to inactivate TcdB, we

determined the secondary structures of full-length TcdB and GTD in the

absence or presence of CAPE by circular dichroism spectroscopy. Apparently,

incubation of CAPE with full-length TcdB resulted in a significant increase in

the α-helix content (15.4% to 21.0%) and a concomitant decrease in the

β-pleated sheet content (34.5% to 20.9%) (**Figure 3A-B**). Similarly, the

secondary structure of GTD was markedly altered upon incubation with CAPE,

with the proportion of α-helices decreasing from 37.9% to 0.0% and the

proportion of β-pleated sheets decreasing from 37.2% to 10.9% (**Figure 3C-D**).

Taken together, these data suggest that CAPE may interact with full-length

TcdB and GTD to induce conformational changes, thus impairing their

functions.

CAPE directly interacts with full-length TcdB and GTD

To investigate the potential interaction between CAPE and TcdB or GTD,

the binding constants were determined by surface plasmon resonance (SPR)

analysis. The results showed that CAPE has strong binding affinity with

full-length TcdB and GTD, and the KD values were 1.269×10^{-5} M and

1.007×10^{-3} M, respectively (**Figure 4A-B**). The interaction between CAPE and

TcdB exhibits a slow approach to equilibrium, which may primarily be attributed

to the rapid occupation of the protein's primary binding sites by the small

molecule during the initial phase. Following this, CAPE proceeds to bind to

sites with secondary or reduced affinity, a process that prolongs the time

required to achieve equilibrium. Furthermore, the possibility that CAPE binds

to multiple sites on TcdB necessitates a duration for the exploration and

occupation of these various sites before equilibrium is established. These data

demonstrate the direct interaction between CAPE and full-length TcdB/GTD.

Next, we performed molecular docking and molecular simulation

dynamics analyses to investigate the interaction between CAPE and GTD

(**Figure 5A**). The results revealed that Leu265, Asp286, Thr465, Ile466,

Leu519, and Trp520 in GTD were potential binding sites for CAPE (**Figure 5A**).

The root mean square deviation (RMSD) of the GTD/CAPE complex was

stable over a 100 ns period of simulation (**Figure 5B**). Further energy

decomposition analysis showed that the side chains of Leu265, Thr465,

Leu519, and Trp520 of GTD contributed to binding with CAPE via van der

Waals interactions, among which Trp520 had the highest binding energy

(**Figure 5C**). Moreover, we mutated each of the residues that were involved in

binding and subsequently determined their glucosyltransferase activities in the

presence or absence of CAPE. The enzymatic activities of these mutants were

similar to that of wild-type GTD (**Figure 5D**). Mutations in GTD_{L265A}, GTD_{T465A},

GTD_{I466A}, and GTD_{L519A} did not affect the inhibitory effect of CAPE (**Figure**

5E-I). However, the inhibition of GTD_{W520A} and GTD_{D286A} by CAPE was notably

compromised (**Figure 5J-K**).

CAPE treatment reduces the pathology of CDI in a murine model

The abovementioned observation that CAPE suppresses the function of

TcdB prompted us to further evaluate the potential therapeutic effects of this

compound in CDI. To this end, we employed a well-established murine model

in which infected mice can develop symptoms resembling those observed in

human CDIs (25). First, cytotoxicity measurements revealed that the viability of

both THP-1 and Vero cells was not affected by CAPE treatment (**Figure**

S5A-B), thus indicating low toxicity. Mice were preconditioned with drinking

water containing 0.5 g/L cefoperazone for 5 days, followed by normal drinking

water for 2 days. After challenge with *C. difficile* BAA-1870 via gavage, mice

were then treated daily with either vehicle or 30 mg/kg CAPE by gavage

throughout the disease course (day 0 to day 5) (**Figure 6A**). Severe weight

loss in *C. difficile*-infected mice was observed on days 2 and 3 post infection

and was accompanied by diarrhea (**Figure 6B**). Notably, compared to the

infection control group, CAPE treatment significantly increased the dry/wet

weight ratio of the mouse feces, suggesting the alleviation of diarrhea

symptoms (**Figure 6C**). Correspondingly, the weight loss of mice was relieved

upon receiving CAPE (**Figure 6B**). Moreover, fecal pellets collected on day 3

were used to determine the bacterial burden. The results revealed that CAPE

treatment significantly decreased the number of *C. difficile* CFUs (**Figure 6D**).

On day 3 post-*C. difficile* infection, the colon was collected for

histopathological analysis. Remarkably, *C. difficile*-infected mice exhibited

conspicuous colon edema, epithelial cell injury and inflammatory factor

infiltration (**Figure 6E**). In contrast, these histopathological lesions were

significantly less common in mice administered CAPE (**Figure 6F**). Taken

together, these data demonstrate the efficacy of CAPE in the treatment of CDI.

CAPE induces distinct microbiota changes in CAPE-treated mice when

compared with model mice

Antibiotic-induced dysbiosis of the gut microbiota promotes the

development of CDI [25]. Therefore, we next determined whether CAPE has any

impact on gut microbial communities that may contribute to its therapeutic

efficacy. To this end, on the 3^{rd} day after *C. difficile* challenge, fecal samples

from the model group and CAPE treatment group were collected and

subjected to microbiota analysis via 16S rRNA gene sequencing. Remarkably,

as reflected by the Venn diagram, CAPE treatment resulted in significant

changes in the microbial composition compared to those in the model group,

and only 16.42% of the operational taxonomic units (OTUs) were shared

(**Figure 7A**). In addition, both the α- and β-diversity of the gut microbiota in

CAPE-treated mice were significantly greater than those in model mice.

Specifically, the increase in the Chao1 richness estimator of mice in the CAPE

treatment group suggested a greater number of microbial species and

increased evenness in the fecal samples (**Figure 7B-C**); the increase in the

Shannon□Wiener diversity index and the Simpson diversity index reflected a

significant increase in microbiota diversity (**Figure 7D-E**). Principal coordinate

analysis (PCoA) revealed distinct confidence ellipses between the bacterial

communities of the CAPE-treated group and the model group, thereby

revealing substantial differences in their microbiota profiles (**Figure 7F**).

A comparative analysis was performed to examine changes at the phylum

level following CAPE treatment. *Bacteroidota*, *Firmicutes,* and *Proteobacteria*

were the major constituents in both the CAPE treatment and model groups,

suggesting that the administration of CAPE did not significantly alter the

mouse microbiota at the phylum level (**Figure 7G**). Next, we compared the 10

most abundant TAXA at the genus level. Notably, CAPE treatment markedly

elevated the abundance of *Bacteroides* and decreased the abundance of

Muribaculaceae and *Parabacteroides* (**Figure 7H**). Leveraging linear

discriminant analysis of effect sizes (LEfSe), the microbiota was examined at

the genus level to identify bacteria exhibiting substantial differences in

abundance between the two cohorts (LDA score > 3.5, $p < 0.05$). After CAPE

treatment,   the   relative   abundance   of   *Bacteroides*,

Ruminococcus_gnavus_group, *Muribaculum* and *Faecalibaculum* OTUs

notably increased, while that of *Hungatella*, *Parasutterella* and *Muribaculaceae*

OTUs significantly decreased (**Figure 7I**). *Bacteroides*, *Muribaculum* and

Faecalibaculum improve colonization resistance to *C. difficile* [26, 27, 28].

Importantly, the abundance of these bacterial species increased following

CAPE treatment (**Figure 7J-L**). In contrast, the abundance of *Parasutterella*, a

bacterium that facilitates *C. difficile* colonization [29], was reduced in mice

receiving CAPE (**Figure 7M**). Taken together, these findings suggest that

CAPE treatment accelerates microbiota recovery during CDI.

CAPE alters the levels of gut metabolites in *C. difficile*-infected mice

In addition to the crucial role of the gut microbiota in conferring

colonization resistance against *C. difficile*, gut microbiota-derived metabolites

significantly contribute to the disease process [30]. Metabolomic analysis was

performed to evaluate the effects of CAPE treatment on gut metabolites in

mice. On the 3[rd] day after infection, the feces of mice in each group were

frozen by liquid nitrogen for metabolomics analysis. The PLS-DA model

exhibited significant separation of clusters between the model and CAPE

treatment groups (**Figure 8A**). Among the 1599 metabolites, 109 showed

substantial differences in levels between the two groups. Specifically,

CAPE-treated mice exhibited 19 metabolites with upregulated expression and

50 metabolites with downregulated expression compared to those in the model

group (fold change > 1.5, $p < 0.05$) (**Figure 8B**). Furthermore, KEGG

enrichment analysis revealed that the differentially abundant metabolites were

significantly enriched in the purine metabolism and bile secretion pathways

(**Figure 8C**). Adenosine is an endogenous purine nucleoside that binds to

specific cell surface receptors to exert anti-inflammatory effects at high

concentrations [31]. Importantly, CAPE treatment of *C. difficile*-infected mice can

increase the adenosine concentration in the gut (**Figure 8D**), which might be

attributed to its inhibition of xanthine oxidase [32]. Similarly, the levels of uric acid,

the final product of purine metabolism, were markedly degraded in mice after

receiving CAPE (**Figure 8D**). An earlier study reported that increasing the gut

adenosine concentration with an adenosine deaminase inhibitor prevents *C.*

difficile toxin A-induced enteritis in mice [33]. Therefore, the increase in gut

adenosine induced by CAPE might play an important role in its therapeutic

efficacy in CDI.

C. difficile predominantly utilizes sugars and amino acids for energy

production through various metabolic pathways, such as fermentation [34].

Notably, Stickland fermentation, a Clostridia-specific type of amino acid

fermentation, is associated with *C. difficile* pathogenesis [34]. D-proline is one of

the preferred nutrients used by *C. difficile* in the Stickland reaction and is

critical for its growth [35, 36]. Interestingly, treatment of *C. difficile*-infected mice

with CAPE significantly reduced gut D-proline levels (**Figure 8D**). In addition,

the metabolomics data also revealed a dramatic increase in the level of

melatonin (**Figure 8D**), which is a ubiquitous indolamine that was previously

shown to decrease the rate of recurrent CDI [37]. Consistently, quantification of

the melatonin content in mouse fecal supernatants using a melatonin detection

kit revealed elevated levels of melatonin in CAPE-treated mice (**Figure S6**).

Moreover, correlation analysis revealed a positive association between

Faecalibaculum abundance and melatonin levels (**Figure S7**). Taken together,

these results indicate that alterations in gut metabolites caused by CAPE

treatment might contribute to its therapeutic efficacy in CDI.

Oral administration of melatonin alleviates the symptoms of CDI

Next, we investigated whether any of the upregulated metabolites had a direct influence on TcdB. To this end, we preincubated recombinant TcdB with each of the selected metabolites prior to addition to the cell cultures. Interestingly, treatment with TcdB and 100 µg/mL melatonin significantly alleviated the cell death induced by this toxin (**Figure 9A**). This observation prompted us to investigate the direct therapeutic potential of melatonin in CDI. Therefore, *C. difficile*-infected mice were orally administered 200 mg/kg melatonin at 24 h intervals. The weight changes and the dry/wet ratio of the stool samples were monitored daily for three days. The results showed that melatonin-treated mice displayed remarkable alleviation of diarrhea, as manifested by notable improvements in weight recovery and a significant increase in the dry/wet weight ratio on the 2^{nd} and 3^{rd} days after *C. difficile* challenge (**Figure 9B-C**). Consistent with these observations, histopathological assessment of colonic tissue sections also revealed attenuated colon damage upon melatonin treatment (**Figure 9D-E**). Taken together, these data indicate that a high dose of exogenous melatonin could protect mice from CDI and that the CAPE-induced increase in endogenous melatonin might improve the therapeutic outcome of patients with CAPE.

Discussion

C. difficile has become the prevalent agent of nosocomial diarrhea and

substantially threatens public health. In particular, the increase in morbidity

and mortality caused by the prevalence of hypervirulent and drug-resistant

strains has become a global concern [7]. In 2019, *C. difficile* was listed by the

US Centers for Disease Control and Prevention as an "urgent threat" that

surpassed methicillin-resistant *Staphylococcus aureus* (MRSA) and

ESBL-producing *Enterobacterales* [38]. *C. difficile* is a commensal bacterium

that colonizes the lower gastrointestinal tract [39]. CDI typically occurs in

susceptible individuals whose normal gut microbiota is disrupted by the

administration of broad-spectrum antibiotics [39]. Although CDI is considered an

antibiotic-associated disease, the current treatment for CDI still relies on

antimicrobial drugs, including metronidazole, fidaxomicin and vancomycin [9, 10].

Despite antibiotic therapy, more than 35% of CDI patients are susceptible to

recurrent infection, which is primarily attributable to the intrinsic effects of

antimicrobial agents, which disrupt the microbiota [12]. In addition, long-term

administration of antibiotics promotes the selection of antibiotic-resistant

subpopulations in the gut, thus increasing the risk of infections caused by

antibiotic-resistant and opportunistic bacteria [40]. As a result, the development

of nonantibiotic therapeutics is needed for the treatment of CDI.

Indeed, numerous novel treatments for CDI are being investigated, among

which the anti-virulence strategy represents one of the most promising and

successful therapeutic approaches [22, 41]. Unlike antibiotics, which target

bacterial components that are critical for bacterial growth, anti-virulence drugs

aim to disarm pathogenic bacteria by neutralizing their virulence factors [42, 43].

Therefore, this strategy imposes less selective pressure that directs bacterial

evolution of antimicrobial resistance [43]. The disease progression of CDI is

strictly dependent on large clostridial toxins (TcdA and TcdB), as strains lacking

both toxins (i.e., A^-B^-) were completely avirulent and failed to induce any

discernable disease symptoms in a hamster model of infection [44]. Importantly,

epidemic *C. difficile* strains with increased morbidity and mortality rates

produce greater amounts of toxins [45]. Based on these premises, TcdA and

TcdB are considered ideal drug targets for CDI in the anti-virulence paradigm

[46]. Indeed, bezlotoxumab, a fully humanized monoclonal antibody that

neutralizes TcdB, was approved by the FDA and used in CDI patients to

reduce the rate of recurrence [47]. This achievement has encouraged

researchers to develop next-generation anti-virulence treatments. In particular,

small molecule inhibitors of toxin production and/or activity have attracted

increasing interest owing to their advantages in cost and the convenience of

oral administration [41]. The multidomain architecture and mechanisms of action

of clostridial toxins have provided distinct options for blocking their toxicity.

Indeed, previous efforts have identified diverse inhibitors that act on distinct

steps of intoxication, such as the inhibition of toxin binding to cell receptors, the

suppression of endosomal maturation, and the inactivation of GTD or CPD

activities [23, 48, 49, 50, 51]. Although which approach is the best remains unknown,

we believe that an inhibitor that targets multiple steps of intoxication would be

superior to those that act on a single step. In this study, we identified a series

of caffeic acid derivatives that were potent inhibitors of TcdB via a phenotypic

screen. Further mechanistic study revealed that CAPE, the phenethyl alcohol

ester of caffeic acid, could block InsP$_6$-induced autoproteolysis of TcdB and

inactivate its GTD function (**Figure 10**). Moreover, CAPE treatment

significantly protected against CDI in a mouse model of infection. Our results

suggested that caffeic acid derivatives, particularly CAPE, are promising lead

compounds for the development of anti-virulence drugs that target TcdB.

CAPE is a major component of propolis, a natural bee-produced substance

that has been used as a remedy for thousands of years in Europe and Asia [52].

As a result, further investigations on the therapeutic potential of propolis

against CDI, either in combination with standard therapy or as a single therapy,

would be of great interest.

Antibiotic-induced gut microbiota dysbiosis is a major risk factor for CDI [53].

Thus, reconstituting the gut microbiota is a promising strategy for CDI therapy.

Fecal microbiota transplantation (FMT) is considered an effective CDI

therapeutic strategy and is widely used in clinical practice [54]. Despite the high

cure rate and increasing popularity of FMT, patients remains limited

acceptance. Another prominent advantage of anti-virulence drugs is that they

allow maintenance of the normal microbiota, which is of critical importance for

the treatment of CDI. Importantly, CAPE treatment of *C. difficile*-infected mice

restored the structure and composition of the microbiota (**Figure 10**). In

particular, the abundance of *Bacteroides*, a critical gut microbiota component

that is required for resistance to *C. difficile* colonization [26, 55, 56, 57], was

significantly elevated upon CAPE treatment. Interestingly, *Bacteroides* were

also predominant in CDI patients administered RBX2660 (Rebyota), an

FDA-approved fecal microbiota product [58, 59]. The increased diversity of the gut

microbiota might contribute to the therapeutic effects of CAPE in CDI.

In addition to the microbiota, gut metabolites also play crucial roles in the

pathogenesis of CDI, either by promoting or suppressing disease progression

[30]. For example, bile acids and amino acids promote spore germination and

pathogenesis, while secondary bile acid and short-chain fatty acids aid in

colonization resistance and suppress spore germination [30]. In addition to

restoration of the gut microbiota, metabolomics analysis of mouse fecal

samples revealed dramatic alterations in gut metabolites upon CAPE

treatment (**Figure 10**). The differentially abundant metabolites were

significantly enriched in the purine metabolism and bile metabolism pathways.

Remarkably, the level of adenosine, an endogenous purine nucleoside that

can regulate immune responses and inflammatory processes by binding to its

receptors, such as its ability to inhibit the release of certain inflammatory

mediators, was dramatically increased in CAPE-treated mice [60, 61]. In contrast,

the amount of uric acid, the final product of purine metabolism, was reduced.

Previous studies showed that the increase in adenosine concentrations

caused by the inhibition of adenosine deaminase via EHNA prevents *C. difficile*

TcdA-mediated damage and inflammation [33]. This protection might be a result

of the anti-inflammatory effects of adenosine, which rely on the activation of its

receptors [62]. In addition, we observed a reduction in the amount of D-proline,

which is one of the preferred amino acids utilized by *C. difficile* to generate

energy through Stickland reactions [35]. Melatonin is a ubiquitous indolamine

that has broad biological activities [63]. Notably, we detected a significantly

increase in the amount of melatonin in the feces of mice that were

administered CAPE. Importantly, an earlier study demonstrated a decreased

risk of recurrent CDI in patients exposed to melatonin, possibly owing to its

antibacterial and anti-inflammatory activities [37]. In this study, we found that oral

administration of exogenous melatonin alleviated the symptoms of CDI in a

mouse model of infection. Although determining the exact mechanism will

require further investigation, we propose that the melatonin-mediated inhibition

of TcdB intoxication observed in this study might contribute to its therapeutic

effect. Taken together, these results indicate that the alterations in gut

metabolites caused by CAPE might also contribute to treatment outcomes for

CDI patients.

Methods

Cell lines

All cell lines (HeLa, Caco-2, Vero, HEK293T and THP-1) were obtained

from ATCC (Manassas, VA). HeLa, Caco-2, Vero and HEK293T cells were

cultured in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen, Carlsbad,

CA) supplemented with 10% fetal bovine serum (FBS) (Invitrogen) and 1%

(w/v) penicillin□streptomycin. THP-1 cells were cultured in Roswell Park

Memorial Institute (RPMI) 1640 medium (Invitrogen) supplemented with 10%

FBS and 1% (w/v) penicillin□streptomycin. All cell lines were cultured at 37 °C

in a 5% CO_2 atmosphere.

Bacterial strains and reagents

C. difficile BAA-1870 (RT027/ST1) was kindly provided by Aiwu Wu

(Guangzhou Medical University) and cultured in brain-heart infusion broth (BHI)

supplemented with 5 g/L yeast extract and 0.1% (w/v) L-cysteine (BHIS). The

compounds were identified from two main sources: FDA-approved drugs

obtained from ApexBio Technology (Houston, TX, USA) and small molecule

natural compounds obtained from Chengdu Ruifensi Biotechnology Co., Ltd.

(Chengdu, China). LysoTracker Red DND-99, kanamycin, and cefoperazone

sodium were purchased from Dalian Meilun Biotechnology Co., Ltd. (Dalian,

China).

Mice

Male C57BL/6 mice aged 6-8 weeks were purchased from Liaoning

Changsheng Biotechnology Co., Ltd., and housed in a specific pathogen-free

(SPF) facility. The mice lived in a controlled environment with a temperature of

23±2 °C and a humidity of 55±10 °C. Prior to infection, the mice were

acclimated to the environment for a minimum of one week. All mouse

experiments were approved by the Jilin University Institutional Animal Care

Committee, Jilin University and strictly conducted according to the guidelines

of this committee (SY202402301).

Protein expression and purification

The expression and purification of full-length TcdB were conducted

following the method described by *Yang* et al. [64]. In brief, *Bacillus megaterium*

harboring pHis1522-TcdB was inoculated into LB media supplemented with a

final concentration of 10 μg/mL tetracycline. Once the $OD600_{nm}$ reached

approximately 0.3, xylose (0.5% w/v) was added to induce protein expression

at 37 °C for 16 h. Then, 100 mL of the bacterial cells was harvested by

centrifugation, resuspended in 5 mL of lysis buffer containing 20 mM imidazole,

300 mM NaCl, 20 mM NaH_2PO_4, 500 μM EDTA, and protease inhibitor cocktail

(adjusted to pH 8.0), and subjected to sonication. The cell lysate was then

centrifuged at 13,400 × g for 20 min to remove cell debris. The TcdB protein

was purified using nickel column affinity chromatography. The purity of the

protein was confirmed by SDS□PAGE, and the protein was subsequently

dialyzed against buffer containing 10% glycerol, 20 mM Tris (pH 7.5), and 150

mM NaCl. After the protein concentration was determined, the purified TcdB

protein was aliquoted and stored at -80 °C for further use.

For the purification of GTD, amplified DNA products were inserted into

pET28a (Novagen), and the *Escherichia coli* (*E. coli*) strain BL21 (DE3) was

used for the expression and purification of GTD. Transformed *E. coli*

pET28a-GTD was cultured in Luria–Bertani (LB) medium supplemented with

10 μg/mL kanamycin. Once the OD600 reached 0.6, the expression of proteins

was induced with 0.2 mM isopropyl-β-d-thiogalactoside (IPTG) overnight at

18 °C. The bacterial cells were then harvested by centrifugation and subjected

to sonication. The cell lysate was then centrifuged at 13,400 × g for 20 min to

remove cell debris. GTD protein was purified using nickel affinity

chromatography and dialyzed in buffer containing 10% glycerol, 20 mM Tris

(pH 7.5), and 150 mM NaCl. The GTD mutants were introduced into

pET28a-GTD using the Quick Change Site-Directed Mutagenesis Kit

(Stratagene, San Diego, CA, USA) with the primers listed in **Table S2.** The

subsequent purification was performed using the same procedure as

described for wild-type GTD.

Reverse transcription-quantitative PCR

Total RNA was extracted by TRIzol and then reverse-transcribed into

complementary DNA by a NovoScript® Plus All-in-one 1st Strand cDNA

Synthesis SuperMix (E047; Novoprotein, China). mRNA expression of *tcdB*

was measured by quantitative PCR with NovoScript®SYBR qPCR SuperMix

Plus (E096; Novoprotein, China) using the primers shown in **Table S3**. 16S

rRNA was used as internal control. The relative mRNA expression levels were

calculated by the delta-delta CT method.

Growth curve assays

Growth curve assay was performed to estimate the influence of CAPE on

the growth of *C. difficile* BAA-1870. The overnight culture of BAA-1870 was

transferred to fresh BHIS medium and incubated at 37°C in an anaerobic

atmosphere. The culture was supplemented with different concentrations of

CAPE ranging from 0 to 64 μg/mL and the optical density at 600 nm was

monitored at 4 h intervals to assess growth of BAA-1870.

Inhibition of spore outgrowth

The impact of CAPE on the sporulation of *C. difficile* BAA-1870 was

assessed using the protocol described by Joshua *et al* [65]. In brief, BAA-1870

cultures were treated with 8 or 32 μg/mL of CAPE for 48 h. The treated

samples were then harvested, subjected to a double wash with PBS, and

incubated in a 60°C water bath for 30 min. Subsequently, 20 μL of each spore

suspension was plated onto BHIS agar supplemented with 0.1% taurine to

facilitate germination. These plates were incubated anaerobically at 37°C for 5

d to allow for spore germination. Following this period, the number of resulting

colonies on the plates was enumerated.

Cell rounding assay

Cell rounding assay was performed according to Kristina *et al*, with

appropriate adjustments [48]. Vero or HeLa cells were seeded in 96-well sterile

plates using serum-free DMEM overnight, and the compounds were

preincubated with TcdB for 1 h. After incubation, the TcdB-compound complex

was added to cells for 1 h at 37 °C. The cell morphology was visualized under

a light microscope.

LysoTracker assay

The pH of endosomes was determined by LysoTracker Red DND-99 [49]. Caco-2 or THP-1 cells in complete DMEM were seeded in 96-well sterile plates overnight. The media was changed to serum-free DMEM for 1 h, and CAPE was preincubated with TcdB for 1 h, followed by addition to the cells and incubation for 2 h. Baf-A1 served as a control. After incubation, LysoTracker and Hoechst dyes were added and the cells were incubated for 1 h. Excess dye was removed by changing the media, and alterations in lysosomal red intensity were documented using a fluorescence microscope.

In vitro **TcdB autoprocessing assay**

The self-cleavage ability of TcdB primarily depends on its cysteine protease activity [48]. CAPE (1-16 μg/mL) was preincubated with TcdB for 1 h, followed by the addition of $InsP_6$ and incubation at 37 °C for 6 h. Subsequently, loading buffer was added for SDS□PAGE analysis, and the results were visualized using Coomassie Brilliant Blue staining.

Rac1 glucosylation assay

Rac1 glucosylation assay was performed following the method described by John *et al* [50]. HEK293T cells were seeded into a 6-well plate. The next day, TcdB and different concentrations of CAPE (4-32 μg/mL) were added to cells. After 2 h, the cells were harvested, lysed using lysis buffer, and subjected to centrifugation at 13,400 × g for 10 min to collect the supernatant. The collected supernatant was then mixed with loading buffer and subjected to SDS□PAGE

followed by western blot analysis. The glycosylation level of Rac1 was

evaluated by determining the levels of non-glycosylated Rac1 (BD Biosciences,

Mississauga, ON) and total Rac1 (Abcam).

UDP-Glo^{TM} UDP-glucose hydrolase assay

GTD activity was assessed according to the manufacturer's instructions [50].

A total volume of 16 µL, consisting of 100 nM GTD, various concentrations of

CAPE and glucosylation buffer, was prepared. Four microliters of UDP-glucose

at a final concentration of 25 µM was added to the mixture, which was then

allowed to react at room temperature for 15 min. To stop the reaction, 10 µL of

the reaction mixture was removed and placed in a white polystyrene 96-well

plate containing 10 µL of UDP detection reagent. The reaction took place at

room temperature for 1 h. The luminescence value was measured at a 750 ms

integral, and the GTD activity was determined based on these measurements.

Circular dichroism (CD) spectroscopy

CD spectra of TcdB, TcdB-CAPE, GTD, and GTD-CAPE were obtained

using a CD spectrophotometer (MOS-500; Bio-Logic, France). TcdB or GTD at

a concentration of 0.2 mg/mL was incubated with 32 µg/mL CAPE. Changes in

the secondary structure of TcdB or GTD were assessed at room temperature

using a quartz cuvette with an optical path length of 1 mm. The scan

wavelength range was set from 190 to 250 nm with a resolution of 0.2 nm and

a bandwidth of 1 nm. The secondary structures of TcdB, TcdB-CAPE, GTD,

and GTD-CAPE were analyzed using the BeStSel web server.

Molecular docking

CAPE was obtained as a ligand from PubChem, and its topology was

generated using AmeberTools. The crystal structure of GTD was retrieved from

the Protein Data Bank (PDB) and processed using AutoDockTools. Docking

calculations were performed using AutodockVina, and molecular dynamics

simulations were carried out using Gromacs 2020.6. The RMSD of the

receptor and ligand was monitored to assess conformational changes during

the binding process. The interaction energy was analyzed using the

Poisson-Boltzmann surface area (MM-PBSA) method, which is a molecular

mechanics approach.

Flow cytometry

Flow cytometry was used to quantify the cellular uptake of TcdB via

endocytosis. To facilitate detection, full-length TcdB was labeled with Alexa

Fluor®488 NHS Ester (Yeasen Biotechnology Co., Ltd., Shanghai, China).

Prior to cell treatment, Alexa Fluor 488-TcdB was preincubated with CAPE at a

concentration of 32 µg/mL for 30 min. The treated solution was then added to

Caco-2 cells and incubated for 1 h at 37 °C. Following three washes with

phosphate-buffered saline (PBS), the Caco-2 cells were collected, and the

fluorescence signal was measured using flow cytometry.

Cytotoxicity assay

Vero or THP-1 cells were plated into 96-well plates. The next day, different

concentrations of CAPE were prepared by diluting them in DMEM and adding

them to the cells. After incubation at 37 °C for 3 h, the viability of cells was

assessed using a Cell Counting Kit-8 (CCK-8) (Beyotime Biotechnology Co.,

Shanghai, China) according to the manufacturer's instructions.

Biacore analysis

The affinity analysis of TcdB or GTD with CAPE was conducted using a

Biacore T100 device (Biacore Inc., Uppsala, Sweden) according to the

manufacturer's protocol. A CM5 chip (General Electric Company, GE) was

utilized to immobilize the TcdB or GTD proteins. Various concentrations of

CAPE were injected, and all steps were performed in running buffer consisting

of 5% DMSO in PBS-P20 (GE). Evaluation version 3.1 software provided by

Biacore and the 1:1 Langmuir combined model were employed for data

analysis.

Mouse infection model

For the acute toxicity model, after pretreatment of TcdB with CAPE (2 or 4

µg/mL) for 1 h, the mice were intraperitoneally injected with a complex of

TcdB-solvent or TcdB-CAPE (30 ng of TcdB per mouse). The survival of mice

was observed for 72 h.

For the CDI model, mice were given cefoperazone (0.5 g/L) in their

drinking water for 5 days (from day -7 to day -2), followed by switching to

sterilized water for 2 days. On day 0, the mice were challenged with 1×10^8

colony-forming units (CFUs) of *C. difficile* BAA-1870 (Ribotype 027) by oral

gavage. For the CAPE or melatonin treatment group, mice were treated with

CAPE (30 mg/kg by oral gavage) or melatonin (200 mg/kg by oral gavage)

every day after *C. difficile* challenge. The weight loss and fecal state of the

mice were monitored daily. Fecal samples were collected on days 0-3 to

evaluate the stool water content as a marker of diarrhea. Additionally, fecal

samples on day 3 were collected to assess *C. difficile* colonization. Colonic

tissues were collected on day 3 post infection, stained with hematoxylin and

eosin (H&E) and scored as previously described [66].

Fecal samples were collected on day 3 post infection, frozen in liquid

nitrogen and transferred to the laboratory with dry ice for 16S rRNA

sequencing and untargeted metabolomics analysis (Novogene Bioinformatics

Technology Co., Ltd., Tianjin, China). The melatonin level in mouse stool

supernatant was determined using the Mouse MT (Melatonin) ELISA Kit

(Sangon Biotechnology Co., Ltd., Shanghai, China).

Statistical analysis

Western blot densitometry and LysoTracker fluorescence intensity were

measured using ImageJ software (NIH, Bethesda, MD, USA). Statistical

analysis was performed using GraphPad Prism 9.0 (GraphPad Software, USA).

Unpaired two-tailed Student's t tests were used for comparing two groups, and

one-way analysis of variance (ANOVA) was used for analyzing data from more

than three groups. The statistical significance was determined using the

log-rank (Mantel–Cox) test (survival rates) and Mann–Whitney U test (tissue

bacterial load) *in vivo*. All the data are expressed as the mean ± standard

deviation (SD). * indicates p <0.05 and ** indicates p <0.01.

Acknowledgments

The authors thank Professor Aiwu Wu (Guangzhou Medical University) for

the *C. difficile* BAA-1870 strain and Professor Hanping Feng (Tufts University)

for the pHis1522-TcdB plasmid. This work was supported by the National Key

Research & Development Program of China (2021YFD1801000), National

Natural Science Foundation of China (32200141), the Natural Science

Foundation of Jilin Province (20230101142JC and 20240101282JC), and The

Fundamental Research Funds for the Central Universities.

Author contributions

702   Y.G., L.Q.H and J.Z.Q designed the study and wrote the manuscript. Y.G.,

703 Y.Z., G.Z.W. and H.T.L. conducted experiments. J.F.W., X.M.D., L.Q.H. and

704 J.Z.Q. supervised the study and revised the manuscript. All authors read and

705 approved the final manuscript. J.Z.Q. and L.Q.H. are responsible for the

706 overall content as the guarantor.

Competing interests

709   The authors declare that they have no competing interests.

Data availability

712   All 16S rRNA gene sequencing data generated and analyzed during the

713 current study were deposited in the Sequence Read Archive (SRA) under the

714 BioProject accession number PRJNA1100298. Untargeted metabolomics data

715 are available via Metabolights with the identifier MTBLS9938.

References

1. Smits WK, Lyras D, Lacy DB, Wilcox MH, Kuijper EJ. Clostridium difficile infection. *Nat Rev Dis Primers* **2**, 16020 (2016).

2. Miller BA, Chen LF, Sexton DJ, Anderson DJ. Comparison of the burdens of hospital-onset, healthcare facility-associated Clostridium difficile Infection and of healthcare-associated infection due to methicillin-resistant Staphylococcus aureus in community hospitals. *Infect Control Hosp Epidemiol* **32**, 387-390 (2011).

3. Kelly CP, LaMont JT. Clostridium difficile--more difficult than ever. *N Engl J Med* **359**, 1932-1940 (2008).

4. Dubberke E. Clostridium difficile infection: the scope of the problem. *J Hosp Med* **7 Suppl 3**, S1-4 (2012).

5. Lessa FC*, et al.* Burden of Clostridium difficile infection in the United States. *N Engl J Med* **372**, 825-834 (2015).

6. Goorhuis A*, et al.* Emergence of Clostridium difficile infection due to a new hypervirulent strain, polymerase chain reaction ribotype 078. *Clin Infect Dis* **47**, 1162-1170 (2008).

7. He M*, et al.* Emergence and global spread of epidemic healthcare-associated Clostridium difficile. *Nat Genet* **45**, 109-113 (2013).

8. Clements AC, Magalhaes RJ, Tatem AJ, Paterson DL, Riley TV. Clostridium difficile PCR ribotype 027: assessing the risks of further worldwide spread. *Lancet Infect Dis* **10**, 395-404 (2010).

9. Cornely OA*, et al.* Fidaxomicin versus vancomycin for infection with Clostridium difficile in Europe, Canada, and the USA: a double-blind, non-inferiority, randomised controlled trial. *Lancet Infect Dis* **12**, 281-289 (2012).

10. Bingley PJ, Harding GM. Clostridium difficile colitis following treatment with metronidazole and vancomycin. *Postgrad Med J* **63**, 993-994 (1987).

11. Feuerstadt P, Theriault N, Tillotson G. The burden of CDI in the United States: a multifactorial challenge. *BMC Infect Dis* **23**, 132 (2023).

12.   Hopkins RJ, Wilson RB. Treatment of recurrent Clostridium difficile
colitis: a narrative review. *Gastroenterol Rep (Oxf)* **6**, 21-28 (2018).

13.   Spigaglia P, Mastrantonio P, Barbanti F. Antibiotic Resistances of
Clostridium difficile. *Adv Exp Med Biol* **1050**, 137-159 (2018).

14.   Seekatz AM, Young VB. Clostridium difficile and the microbiota. *J Clin*
Invest **124**, 4182-4189 (2014).

15.   Paredes-Sabja D, Shen A, Sorg JA. Clostridium difficile spore biology:
sporulation, germination, and spore structural proteins. *Trends Microbiol*
22, 406-416 (2014).

16.   Lyras D*, et al.* Toxin B is essential for virulence of Clostridium difficile.
Nature **458**, 1176-1179 (2009).

17.   Kuehne SA, Cartman ST, Heap JT, Kelly ML, Cockayne A, Minton NP.
The role of toxin A and toxin B in Clostridium difficile infection. *Nature*
467, 711-713 (2010).

18.   Carter GP*, et al.* Defining the Roles of TcdA and TcdB in Localized
Gastrointestinal Disease, Systemic Organ Damage, and the Host
Response during Clostridium difficile Infections. *mBio* **6**, e00551 (2015).

19.   Drudy D, Fanning S, Kyne L. Toxin A-negative, toxin B-positive
Clostridium difficile. *Int J Infect Dis* **11**, 5-10 (2007).

20.   Chen P*, et al.* Structure of the full-length Clostridium difficile toxin B. *Nat*
Struct Mol Biol **26**, 712-719 (2019).

21.   Orrell KE, Melnyk RA. Large Clostridial Toxins: Mechanisms and Roles
in Disease. *Microbiol Mol Biol Rev* **85**, e0006421 (2021).

22.   Rasko DA, Sperandio V. Anti-virulence strategies to combat
bacteria-mediated disease. *Nat Rev Drug Discov* **9**, 117-128 (2010).

23.   Tam J, Beilhartz GL, Auger A, Gupta P, Therien AG, Melnyk RA. Small
molecule inhibitors of Clostridium difficile toxin B-induced cellular
damage. *Chem Biol* **22**, 175-185 (2015).

24.   Balaha M, De Filippis B, Cataldi A, di Giacomo V. CAPE and
Neuroprotection: A Review. *Biomolecules* **11**, (2021).

25.   Garland M*, et al.* The Clinical Drug Ebselen Attenuates Inflammation

and Promotes Microbiome Recovery in Mice after Antibiotic Treatment
for CDI. *Cell Rep Med* **1**, (2020).
807

808 26. Li X*, et al.* A strain of Bacteroides thetaiotaomicron attenuates
809 colonization of Clostridioides difficile and affects intestinal microbiota
810 and bile acids profile in a mouse model. *Biomed Pharmacother* **137**,
811 111290 (2021).
812

813 27. Kienesberger B*, et al.* (S)-Reutericyclin: Susceptibility Testing and In
814 Vivo Effect on Murine Fecal Microbiome and Volatile Organic
815 Compounds. *Int J Mol Sci* **22**, (2021).
816

817 28. Khanna S*, et al.* Gut microbiome predictors of treatment response and
818 recurrence in primary Clostridium difficile infection. *Aliment Pharmacol*
819 *Ther* **44**, 715-727 (2016).
820

821 29. Dong D*, et al.* Effects of intestinal colonization by Clostridium difficile
822 and Staphylococcus aureus on microbiota diversity in healthy
823 individuals in China. *BMC Infect Dis* **18**, 207 (2018).
824

825 30. Gurung B, Stricklin M, Wang S. Gut Microbiota-Gut Metabolites and
826 Clostridioides difficile Infection: Approaching Sustainable Solutions for
827 Therapy. *Metabolites* **14**, (2024).
828

829 31. Cronstein BN. Adenosine, an endogenous anti-inflammatory agent. *J*
830 *Appl Physiol (1985)* **76**, 5-13 (1994).
831

832 32. Yong T*, et al.* Caffeic acid phenethyl ester alleviated hypouricemia in
833 hyperuricemic mice through inhibiting XOD and up-regulating OAT3.
834 *Phytomedicine* **103**, 154256 (2022).
835

836 33. de Araujo Junqueira AF*, et al.* Adenosine deaminase inhibition prevents
837 Clostridium difficile toxin A-induced enteritis in mice. *Infect Immun* **79**,
838 653-662 (2011).
839

840 34. Neumann-Schaal M, Jahn D, Schmidt-Hohagen K. Metabolism the
841 Difficile Way: The Key to the Success of the Pathogen Clostridioides
842 difficile. *Front Microbiol* **10**, 219 (2019).
843

844 35. Bouillaut L, Self WT, Sonenshein AL. Proline-dependent regulation of
845 Clostridium difficile Stickland metabolism. *J Bacteriol* **195**, 844-854
846 (2013).
847

36. Jackson S, Calos M, Myers A, Self WT. Analysis of proline reduction in

the nosocomial pathogen Clostridium difficile. *J Bacteriol* **188**,
8487-8495 (2006).

37. Sutton SS, Magagnoli J, Cummings TH, Hardin JW. Melatonin as an
Antimicrobial Adjuvant and Anti-Inflammatory for the Management of
Recurrent Clostridioides difficile Infection. *Antibiotics (Basel)* **11**,
(2022).

38. 2019   AR   Threats   Report.,   2019.   [Available   from:
https://www.cdc.gov/drugresistance/biggest-threats.html].

39. Abbas   A,   Zackular   JP.   Microbe-microbe   interactions   during
Clostridioides difficile infection. *Curr Opin Microbiol* **53**, 19-25 (2020).

40. Raplee I*, et al.* Emergence of nosocomial associated opportunistic
pathogens in the gut microbiome after antibiotic treatment. *Antimicrob
Resist Infect Control* **10**, 36 (2021).

41. Beilhartz GL, Tam J, Melnyk RA. Small Molecules Take A Big Step
Against Clostridium difficile. *Trends Microbiol* **23**, 746-748 (2015).

42. Ranftler C*, et al.* Binding and neutralization of C. difficile toxins A and B
by purified clinoptilolite-tuff. *PLoS One* **16**, e0252211 (2021).

43. Zucca M, Scutera S, Savoia D. Novel avenues for Clostridium difficile
infection drug discovery. *Expert Opin Drug Discov* **8**, 459-477 (2013).

44. Kuehne SA, Collery MM, Kelly ML, Cartman ST, Cockayne A, Minton NP.
Importance of toxin A, toxin B, and CDT in virulence of an epidemic
Clostridium difficile strain. *J Infect Dis* **209**, 83-86 (2014).

45. Warny M*, et al.* Toxin production by an emerging strain of Clostridium
difficile associated with outbreaks of severe disease in North America
and Europe. *Lancet* **366**, 1079-1084 (2005).

46. Stewart D, Anwar F, Vedantam G. Anti-virulence strategies for
Clostridioides difficile infection: advances and roadblocks. *Gut Microbes*
12, 1802865 (2020).

47. Deeks ED. Bezlotoxumab: A Review in Preventing Clostridium difficile
Infection Recurrence. *Drugs* **77**, 1657-1663 (2017).

48. Bender KO*, et al.* A small-molecule antivirulence agent for treating
Clostridium difficile infection. *Sci Transl Med* **7**, 306ra148 (2015).

893
894 49. Chan H, *et al.* Vitamin D(3) and carbamazepine protect against
895   Clostridioides difficile infection in mice by restoring macrophage
896   lysosome acidification. *Autophagy* **18**, 2050-2067 (2022).
897
898 50. Tam J, *et al.* Host-targeted niclosamide inhibits C. difficile virulence and
899   prevents disease in mice without disrupting the gut microbiota. *Nat*
900   *Commun* **9**, 5233 (2018).
901
902 51. Tam J, *et al.* Intestinal bile acids directly modulate the structure and
903   function of C. difficile TcdB toxin. *Proc Natl Acad Sci U S A* **117**,
904   6792-6800 (2020).
905
906 52. Tek NA, Senture SA, Ersoy N. Is Propolis a Potential Anti-Obesogenic
907   Agent for Obesity? *Curr Nutr Rep*,   (2024).
908
909 53. Schubert AM, Sinani H, Schloss PD. Antibiotic-Induced Alterations of
910   the Murine Gut Microbiota and Subsequent Effects on Colonization
911   Resistance against Clostridium difficile. *mBio* **6**, e00974 (2015).
912
913 54. van Nood E, *et al.* Duodenal infusion of donor feces for recurrent
914   Clostridium difficile. *N Engl J Med* **368**, 407-415 (2013).
915
916 55. Bornet E, Westermann AJ. The ambivalent role of Bacteroides in enteric
917   infections. *Trends Microbiol* **30**, 104-108 (2022).
918
919 56. Pike CM, Theriot CM. Mechanisms of Colonization Resistance Against
920   Clostridioides difficile. *J Infect Dis* **223**, S194-S200 (2021).
921
922 57. Deng H, *et al.* Corrigendum: Bacteroides fragilis Prevents Clostridium
923   difficile Infection in a Mouse Model by Restoring Gut Barrier and
924   Microbiome Regulation. *Front Microbiol* **10**, 601 (2019).
925
926 58. Lee C, *et al.* Safety of fecal microbiota, live-jslm (REBYOTA()) in
927   individuals with recurrent Clostridioides difficile infection: data from five
928   prospective clinical trials. *Therap Adv Gastroenterol* **16**,
929   17562848231174277 (2023).
930
931 59. Gonzales-Luna AJ, Carlson TJ, Garey KW. Review Article: Safety of
932   Live Biotherapeutic Products Used for the Prevention of Clostridioides
933   difficile Infection Recurrence. *Clin Infect Dis* **77**, S487-S496 (2023).
934
60. Hasko G, Linden J, Cronstein B, Pacher P. Adenosine receptors:
therapeutic aspects for inflammatory and immune diseases. Nat Rev

Drug Discov 7, 759-770 (2008).

61.   Shakya AK, Naik RR, Almasri IM, Kaur A. Role and Function of
Adenosine and its Receptors in Inflammation, Neuroinflammation, IBS,
Autoimmune Inflammatory Disorders, Rheumatoid Arthritis and
Psoriasis. Curr Pharm Des 25, 2875-2891 (2019).

62.   Cavalcante IC*, et al.* Effect of novel A2A adenosine receptor agonist
ATL 313 on Clostridium difficile toxin A-induced murine ileal enteritis.
Infect Immun **74**, 2606-2612 (2006).

63.   Ma N, Zhang J, Reiter RJ, Ma X. Melatonin mediates mucosal immune
cells, microbial metabolism, and rhythm crosstalk: A therapeutic target
to reduce intestinal inflammation. *Med Res Rev* **40**, 606-632 (2020).

64.   Yang G*, et al.* Expression of recombinant Clostridium difficile toxin A
and B in Bacillus megaterium. *BMC Microbiol* **8**, 192 (2008).

65.   Fletcher JR*, et al.* Clostridioides difficile exploits toxin-mediated
inflammation to alter the host nutritional landscape and exclude
competitors from the gut microbiota. *Nat Commun* **12**, 462 (2021).

66.   Reeves AE, Theriot CM, Bergin IL, Huffnagle GB, Schloss PD, Young
VB. The interplay between microbiome dynamics and pathogen
dynamics in a murine model of Clostridium difficile Infection. *Gut*
Microbes **2**, 145-158 (2011).

Figures

Figure 1. CAPE suppresses TcdB-mediated cell rounding.

(A) Schematic of the process used for the identification of small molecules that

inhibit the cell rounding phenotype induced by TcdB. TcdB (0.2 ng/mL) was

preincubated with 8 µg/mL of individual compounds prior to addition to the cell

cultures. **(B)** Active compounds were identified from the chemical library

consisting of 2076 chemicals. The red dots represent caffeic acid and its

derivatives. **(C)** Dose-dependent CAPE inhibition of TcdB-induced rounding of

Vero (upper) and HeLa (lower) cells. Scale bar, 100 µm. The dose titration

curves are presented on the right. **(D)** The survival rates of mice that were

intraperitoneally injected with CAPE-pretreated or native TcdB (30 ng). * *p*

<0.05 and ** *p* <0.01 by log-rank (Mantel□Cox) test.

Figure 2. Mechanism by which CAPE suppresses TcdB intoxication.

(A) Schematic illustration of the multistep mechanism of TcdB intoxication of host cells. **(B)** Determination of internalized TcdB levels. Alexa Fluor 488-labeled TcdB was incubated with 32 µg/mL CAPE for 30 min before it was added to the cell culture. The cellular uptake of TcdB was measured by flow cytometry analysis. **(C)** Lysosomal activity in host cells treated with CAPE was measured via LysoTracker staining. The nuclei were stained with DAPI. Bar, 100 µm. **(D)** The intensities of LysoTracker staining were quantified using ImageJ. * $p < 0.05$; ** $p < 0.01$ using one-way ANOVA. **(E-F)** CAPE inhibits InsP$_6$-induced autoprocessing of full-length TcdB. Two micrograms of TcdB was preincubated with increasing concentrations of CAPE for 1 h. Self-cleavage of TcdB was initiated by the addition of 3 µg of InsP$_6$ and the cells were incubated for 6 h at 37 $^{\circ}$C. Reactions were terminated with 5x SDS loading buffer, and the release of GTD was visualized by CBB staining of the

SDS□PAGE gel **(E)**. The level of released GTD in panel E was quantified by

ImageJ **(F)** * p <0.05; ** p <0.01 using one-way ANOVA. **(G)** Dose-dependent

inhibition of the GTD activity of TcdB by CAPE. Recombinant GTD of TcdB was

mixed with different concentrations of CAPE (0-32 µg/mL) and incubated for 15

min at 25 ^{o}C. GTD activity was measured by the UDP-Glo™

glycosyltransferase assay. **(H-I)** Intracellular Rac1 glucosylation in cells treated

with TcdB. Native or CAPE-pretreated TcdB was added to the cell culture and

cells were incubated for 2 h. The cells were lysed, and the glucosylation of

Rac1 was analyzed by western blotting using antibodies recognizing total

Rac1 and unglycosylated Rac1 **(H)**. The level of Rac1 glucosylation shown in

H was quantified by ImageJ **(I)**. * p <0.05; ** p <0.01 using one-way ANOVA.

1005

1006

Figure 3. CAPE alters the secondary structure of TcdB and GTD.

(A-B) The secondary structure of TcdB **(A)** or GTD **(B)** in the presence or absence of 32 μg/mL CAPE was determined by CD spectroscopy. **(C-D)** Calculated CD spectra of CAPE-treated TcdB **(C)** and GTD **(D)**. The wavelength for CD spectroscopy was set at 190-250 nm.

1007

1008

1009

1010

1011

1012

Figure 4. Direct interaction of CAPE with TcdB and GTD.

(A-B) Full-length TcdB or GTD was immobilized on a CM5 chip, followed by

the injection of various concentrations of CAPE. The response unit (RU) value

and affinities of CAPE interactions with TcdB **(A)** and GTD **(B)** were

determined by a Biacore T100.

Figure 5. Asp286 and Trp520 in the GTD are critical for CAPE-mediated inhibition of GTD.

(A) Three-dimensional structure determination of GTD with the CAPE complex was performed by molecular modeling. **(B)** The RMSD values of the GTD-CAPE complex. **(C)** Decomposition of the binding energy on a per-residue basis in the binding sites of the GTD-CAPE complex. **(D)** Glucosyltransferase activity of wild-type GTD and the GTD mutants. **(E-K)** Residual activity of wild-type GTD **(E)** and its mutants GTD$_{L265A}$ **(F)**, GTD$_{T465A}$ **(G)**, GTD$_{I466A}$ **(H)**, GTD$_{L519A}$ **(I)**, GTD$_{W520A}$ **(J)** and GTD$_{D286A}$ **(K)** in the presence of increasing concentrations of CAPE (0-4 µg/mL). * p <0.05; ** p <0.01 using one-way ANOVA.

1032

Figure 6. CAPE treatment blocks the pathology of CDI in mice.

(A) Schematic of the process used for the establishment of the CDI mouse

model. **(B)** Weight changes in CDI mice after CAPE or vehicle treatment

(n=10). **(C)** The fecal dry/wet weight ratio in CDI mice treated with CAPE or

vehicle (n=6). * p <0.05; ** p <0.01 using one-way ANOVA. **(D)** Bacterial

colonization in the feces of CDI mice treated with CAPE or vehicle (n=15). * p

<0.05; ** p <0.01 using the Mann□Whitney U test. **(E)** Pathological analysis of

colon sections from CDI mice treated with CAPE or vehicle. The black boxes

are the regions that were magnified and are shown underneath. **(F)**

Histopathological scores of vehicle-treated and CAPE-treated mice in terms of

cell infiltration (orange), epithelial damage (purple), and edema (green). * p

1044   <0.05; ** p <0.01 using one-way ANOVA.

1045

1046

Figure 7. The influence of CAPE treatment on the diversity and composition of the gut microbiota in CDI mice.

(A) Venn diagram illustrating the numbers of differentially abundant bacteria identified using a single model and their cooperation. **(B-E)** Quantification of the α diversity of the gut microbiota according to the observed species richness **(B-C),** Shannon index **(D)**, and Simpson index **(E)**. **(F)** PCoA showing β diversity based on the weighted UniFrac distance. **(G-H)** Relative abundance of the top 10 predominant bacteria classified at the phylum **(G)** and genus **(H)** levels. **(I)** Significant changes in flora among the model and CAPE groups, as measured by LEfSe analysis (LDA score [log 10] > 3.5). **(J-M)** Relative abundances of *Bacteroides* **(J)**, *Muribaculum* **(K)**, *Faecalibaculum* **(L)**, and *Parasutterella* **(M)** in mice in the model and CAPE treatment groups. * p <0.05; ** p <0.01 using an unpaired Student's t test.

1060

Figure 8. Alterations in gut metabolites in CAPE-treated mice with CDI.

(A) The score plots show orthogonal partial least square discriminant analysis

(PLS-DA) results. **(B)** Volcano plot representing the upregulated or

downregulated gut metabolites in the CAPE treatment group compared to

those in the model group. **(C)** Enrichment bubble map of the top 20 metabolic

pathways in the CAPE treatment and model groups. **(D)** The 26 potential

biomarkers are displayed in a heatmap, along with hierarchical clustering

analysis.

1069

Figure 9. Oral administration of melatonin alleviates the pathology of CDI.

(A) The impact of upregulated metabolites on the cytotoxicity of TcdB. TcdB

(0.2 ng/mL) was incubated with each of the metabolites (100 μg/mL) for 1 h at

37 ^{o}C before being added to the Vero cell culture. At 3 h after TcdB treatment,

cell death was evaluated with a CCK-8 kit. **(B)** Body weight changes in

melatonin-treated CDI mice (n=5). **(C)** The fecal dry/wet weight ratio in CDI

mice after melatonin treatment (n=5). **(D)** Histopathological analysis of colon

1077   tissue from CDI mice treated with melatonin. **(E)** Histopathological scores of

1078   melatonin-treated CDI mice. * p <0.05; ** p <0.01 using one-way ANOVA.

1079

1080

Figure 10. A model for CAPE-mediated suppression of TcdB cytotoxicity and protection of mice from CDI.

The intoxication of cells by TcdB involves multistep mechanisms. Direct interaction of CAPE with full-length TcdB blocks $InsP_6$-induced autoproteolysis, resulting in decreased production of active GTD. The binding between GTD and CAPE inhibits its glucosyltransferase activity, thus decreasing the glucosylation level of Rac1. CAPE treatment of CDI mice markedly restored the diversity and composition of the gut microbiota and induced changes in gut metabolites, which might have contributed to the therapeutic outcomes of CDI following CAPE treatment.

A *C. difficile* challenge and treatment of CAPE (30 mg/kg; intragastrical administration; every day)

C57BL/6
6-8 weeks

Cefoperazone in drinking water (0.5 g/L)

Day -7

Day -2
Change to sterile water

Day 0

Day 3
Sampling

Day 5
Endpoint monitoring

B
Weight (percent of day 0)
Days
Ctrl   Model   CAPE
n = 10

C
% of water in stool
Days
Ctrl   Model   CAPE

D
CFUs/g (log10)
Model   CAPE

E
Ctrl   Model   CAPE

* Cellular infiltration   * Epithelial damage   * Edema

F
Histopathological scores
Cellular infiltration
Epithelial damage
Edema
Ctrl   Model   CAPE

A 1 h

B 12 h

A THP-1

B Vero

1　**Table S1. IC$_{50}$ of the caffeic acid and its derivate on TcdB-mediated**

2　**cell-rounding.**

Compound	Structure	IC$_{50}$ (µg/mL)
Caffeic acid		527.5
Ethyl caffeate		5.9
CAPE		3.0
Echinacoside		5.1
Rosmarinic acid		198.1
Salvianolic acid		3.4
Verbascoside		5.4

5 **Table S2. Primers used in this study.**

Primers	Sequences	Notes
YG101	ctgGGATCCATGAGTTTAGTTAATAG	*gtd* BamHI-F
YG102	ctgGTCGACCTAAAGAGAACCTTCAA	*gtd* SalI-R
YG103	CAGAAGCAGCAGCTGCATTCCACCTTTCTACC AACTCTTGTTC	gtd_{L265A} -F
YG104	GAACAAGAGTTGGTAGAAAGGTGGAATGCAG CTGCTGCTTCTG	gtd_{L265A} -R
YG105	GTATTCCTGGTAACATATCAACAGCTAAATACAT ACCACCAATTTCT	gtd_{D286A} -F
YG106	AGAAATTGGTGGTATGTATTTAGCTGTTGATAT GTTACCAGGAATAC	gtd_{D286A} -R
YG107	CAGGGCCACTTAAGTTAATAGCAGTTTTAACAT CTGGGAAGAA	gtd_{T465A}-F
YG108	TTCTTCCCAGATGTTAAAACTGCTATTAACTTA AGTGGCCCTG	gtd_{T465A}-R
YG109	TGCTTCAGGGCCACTTAAGTTAGCAGTAGTTT TAACATCTGGGAAG	gtd_{I466A}-F
YG110	CTTCCCAGATGTTAAAACTACTGCTAACTTAAG TGGCCCTGAAGCA	gtd_{I466A}-R
YG111	CTTGCATCGTCAAATGACCATGCGCTAGCCAT TTCTTGTTCAGT	gtd_{L519} -F
YG112	ACTGAACAAGAAATGGCTAGCGCATGGTCATT TGACGATGCAAG	gtd_{L519A} -R
YG113	CTCTTGCATCGTCAAATGACGCTAAGCTAGCC ATTTCTTGTTC	gtd_{W520A} -F
YG114	GAACAAGAAATGGCTAGCTTAGCGTCATTTGA CGATGCAAGAG	gtd_{W520A}-R

6
7

8 **Table S3. Sequence of primers used for Reverse transcriptase PCR**

9 **(RT-PCR) analysis.**

Gene	Primer	Sequence (5'-3')
tcdB	Forward	TAGCAAGAAGTTCAGAGAGAGGAT
	Reverse	CTTGTATTTCACCATCTCCAGGA
16S	Forward	GCGTAGGCGGTCTTTCAAG
	Reverse	TCGCAACTGGTGTTCCTCC

11