# Bacterial cross-feeding can promote gene retention by lowering gene expression costs

Short title: Cross-feeding can promote gene retention

Ying-Chih Chuang[1,2,†], Megan G. Behringer[3], Gillian Patton[3],
Jordan T. Bird[4], Crystal E. Love[1], Ankur Dalia[1], and James B. McKinlay[1]*

[1]Department of Biology, Indiana University, Bloomington, IN, USA
[2]Biochemistry Program, Indiana University, Bloomington, IN, USA
[3]Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA
[4]Battelle Memorial Institute, Columbus, OH 43201

*Corresponding author: James B. McKinlay
**Email:** jmckinla@iu.edu

[†] Current address: Department of Biological Sciences, University of Southern California,
Los Angeles, CA, USA

**Competing interests statement:** The authors declare no conflicts of interest.

**Keywords:** cross-feeding; purine; adenine; siderophore, microbial interactions;
Rhodopseudomonas; palustris; Black Queen Hypothesis; mutualism; excretion;
microbial physiology, gene loss

Study Funding information:

This work was supported in part by US Army Research Office grants W911NF-14-1-
0411 and W911NF-17-1-0159, the National Science Foundation CAREER award MCB-
1749489 to JBM, and National Institutes of Health grants R35GM128674 to ABD and
R35GM150625 to MB. Supercomputing resources were supported in part by Lilly
Endowment, Inc., through its support for the IU Pervasive Technology Institute.

**ABSTRACT**

Gene loss is expected in microbial communities when the benefit of obtaining a biosynthetic precursor from a neighbor via cross-feeding outweighs the cost of retaining a biosynthetic gene. However, gene cost primarily comes from expression, and many biosynthetic genes are only expressed when needed. Thus, one can conversely expect cross-feeding to repress biosynthetic gene expression and promote gene retention by lowering gene cost. Here we examined long-term bacterial cocultures pairing *Escherichia coli* and *Rhodopseudomonas palustris* for evidence of gene loss or retention in response to cross-feeding of non-essential adenine. Although *R. palustris* continued to externalize adenine in long-term cultures, *E. coli* did not accumulate mutations in purine synthesis genes, even after 700 generations. *E. coli* purine synthesis gene expression was low in coculture, suggesting that gene repression removed selective pressure for gene loss. In support of this explanation, *R. palustris* also had low transcript levels for iron-scavenging siderophore genes in coculture, likely because *E. coli* facilitated iron acquisition by *R. palustris*. *R. palustris* siderophore gene mutations were correspondingly rare in long-term cocultures but were prevalent in monocultures where transcript levels were high. Our data suggests that cross-feeding does not always drive gene loss, but can instead promote gene retention by repressing costly expression.

**Graphical abstract**

## INTRODUCTION

Changes to the genetic inventory of a given microbe can be influenced by living in a community. An example of this influence is described in the Black Queen Hypothesis (BQH). Based on the card game Hearts, where players avoid holding the costly Queen of Spades, the BQH posits that gene loss (herein we do not distinguish loss-of-function mutations from gene deletions for convenience) will occur when acquiring a resource from neighbor is less costly than producing that resource (1). The BQH involves a producer that generates a public resource, and a beneficiary that uses the resource and is thus subject to beneficial gene loss. The BQH is analogous to the emergence of cheaters that lose genes and benefit by exploiting a public resource. However, unlike a cheater, a beneficiary is less likely to be the same species as the producer and is less likely to harm the producer (1, 2). The BQH, or analogous processes, are now commonly used to explain the prevalence of auxotrophs, which cannot synthesize one or more essential nutrients (3, 4). Indeed, several groups have observed that auxotrophs carrying biosynthetic gene deletions have a fitness advantage over wild-type cells when corresponding biosynthetic precursors are provided (5-9).

A confounding aspect of BQH prediction of gene loss is that nutrient availability can also repress gene expression and thus lower the cost of a gene (3, 10, 11); most of a gene's cost comes from its expression (12-16), and thus repressing gene expression could have similar cost-savings as a loss-of-function mutation. Thus, one can arrive at an opposite prediction wherein cross-feeding could promote gene retention by repressing costly gene expression. This prediction is well-documented from studies of cheaters. For example, cheaters that had lost genes for iron-scavenging siderophores emerged when iron was scare and siderophore genes were expressed, but gene loss was not observed when iron was plentiful and siderophore genes were repressed (11, 17, 18). Thus, nutrient availability alone might not be a good predictor for gene loss and the emergence of beneficiaries.

Previously, we developed cocultures pairing fermentative *Escherichia coli* and phototrophic *Rhodopseudomonas palustris*, wherein each species is dependent on the other for essential nutrients; *E. coli* ferments glucose into organic acids that provide essential carbon to *R. palustris* while *R. palustris* converts $N_2$ gas into $NH_4^+$, providing essential nitrogen to *E. coli* (19). More recently, we discovered that *R. palustris* also externalizes adenine at levels that can support an *E. coli* purine auxotroph (20, 21). There is also suggestive evidence that *E. coli* facilitates *R. palustris* iron acquisition, perhaps through cross-feeding of *E. coli* siderophores (22). Based on the BQH alone, one might predict that the public availability of adenine and siderophores would enrich for *E. coli* adenine auxotrophs and *R. palustris* siderophore mutants over time.

Here we examined long-term monocultures and cocultures for evidence of gene loss due to the availability of non-essential adenine and possible siderophores from a partner. In each case gene retention was correlated with low expression, suggesting that repression of gene expression can decrease the frequency of gene loss.

## RESULTS

**Gene loss is not beneficial when adenine is provided to *E. coli* monocultures.** In anaerobic cocultures where *R. palustris* provided essential nitrogen ($NH_4^+$) to *E. coli* and *E. coli* provided essential carbon (organic acids) to *R. palustris* (19), we learned that *R. palustris* also externalizes enough adenine to support an *E. coli ΔpurH* purine auxotroph (20, 21). Herein, we recognized adenine availability as an opportunity to look for evidence of the BQH; de novo purine synthesis costs more ATP than the purine salvage pathway, and thus could lead to purine synthesis gene loss.

Fitness benefits have been observed for auxotrophic mutants, when the corresponding nutrient is available either as a supplement or via cross-feeding (5-8). Thus, we first tested whether loss of purine biosynthesis would be advantageous in the presence of adenine by comparing the growth of *E. coli ΔpurH* versus wild-type (WT) PFM2 parent cultures. We focused on *E. coli* PFM2 instead of the MG1655 (used previously by our lab) because PFM2 does not carry the *rph* mutation that leads to slower growth in the presence of adenine (23, 24), which can be alleviated by a *ΔpurH* mutant (Chuang and McKinlay in prep). Despite the availability of adenine, the *ΔpurH* mutant did not grow faster than the PFM2 parent in the presence of adenine (Fig 1A, B).

**Figure 1. Adenine availability does not provide a fitness advantage to an engineered *E. coli* PFM2 purine auxotroph. A.** Growth of WT *E. coli* PFM2 and its *ΔpurH* mutant in monoculture with and without 35 µM adenine (ade). **B.** Monoculture growth rates for WT PFM2 vs a corresponding *ΔpurH* mutant with and without 35 µM adenine. 'a,' indicates statistically similar comparisons ($p > 0.01$) as determined using a one-way ANOVA with Tukey's multiple comparisons test. **C.** Invasion-from-rare assay competing a *ΔpurH* mutant against its PFM2 parent in coculture with *R. palustris* CGA676 under $N_2$-fixing conditions where *R. palustris* excretes $NH_4^+$ and adenine. The orange line is the best fit from a linear regression analysis with 95% CI shaded in purple. Corresponding equations and x-intercepts (upper, lower bounds for 95% CI) are shown. Change in frequency = (*E. coli ΔpurH* / (*E. coli* WT + *E. coli ΔpurH*))$_{final}$ − (*E. coli ΔpurH* / (*E. coli* WT + *E. coli ΔpurH*))$_{initial}$. **A-C.** Each data point represents a measurement from a single biological replicate.

We also assessed the fitness impact of a *ΔpurH* mutation by competing the mutant
against the PFM2 parent in coculture with $NH_4^+$- and adenine-excreting *R. palustris*
NifA* (CGA676). We used a range of initial frequencies so that we could simultaneously
test for coexistence by mutual invasibility, where an equilibrium frequency can be
extrapolated from the x-intercept by linear regression analysis (25-27). However, there
was poor linear correlation in the invasion-from-rare assay (Fig 1C). The same was true
when a *ΔpurH*::km$^R$ mutant was used (Fig S1). However, the *ΔpurH* mutant tended to
decrease in frequency relative to parent in cocultures, where adenine is available (Fig
1C and S1). Taken together, our results suggest that a *ΔpurH* mutant does not have a
competitive advantage over PFM2 in the presence of adenine. Thus, at least in this
engineered context, we do not expect that adenine availability to benefit *E. coli* PFM2
purine auxotrophs.

150
**Fig S1. Invasion-from-rare assay competing *E. coli***
***ΔpurH*::km$^R$ against its PFM2 parent in coculture with *R.***
***palustris* CGA676.** The orange line is the best fit from a
linear regression analysis with 95% CI shaded in purple.
Corresponding equations and x-intercepts (upper, lower
bounds for 95% CI) are shown. Change in frequency = (*E.*
*coli ΔpurH* / (*E. coli* WT + *E. coli ΔpurH*))$_{final}$ − (*E. coli ΔpurH*
/ (*E. coli* WT + *E. coli ΔpurH*))$_{initial}$. Each data point
represents a measurement from a single biological replicate.

160
161
162
**Adenine auxotrophs are rare in long-term cocultures.** Although the engineered
*ΔpurH* mutant did not have an obvious fitness advantage, it is possible that (i) there was
a subtle difference in growth rate that could be enriched over many generations and (ii)
other kinds of purine auxotrophy mutations could lead to a fitness advantage. To
account for these possibilities, we looked for the evidence of purine auxotrophs in long-
term monocultures and cocultures of *E. coli* PFM2 and $NH_4^+$- and adenine-excreting *R.*
*palustris* (CGA676; NifA*) that we had maintained for other reasons (Fig 2A; Table S1-
5). Adenine is amply available even in evolved cocultures (Chuang and McKinlay in
prep). Focusing on the latest common timepoint between monoculture vs coculture
treatments (generation 650), we did not observe obvious mutations in purine synthesis
genes in PFM2 in cocultures or monocultures (Fig 2B; Fig S2). When considering other
time points, only one purine biosynthesis gene mutation across was observed in one
monoculture line at generation 204 (*purT* at 54.3%, line 15; Table S2).

176
To account for non-obvious mutations that might lead to purine auxotrophy. We also
screened four *E. coli* isolates from each monoculture and coculture line (40 monoculture
isolates and 40 coculture isolates) for adenine auxotrophy in liquid cultures in 96-well
plates (Fig 2C). Although several monoculture isolates could not grow in minimal media,
suggesting that dependencies might have developed between *E. coli* subpopulations,
only two monoculture isolates were possibly adenine auxotrophs (line 20; Fig 2C). We
suspect that the other auxotrophs are due to mutations in *rpoC*, which were

conditionally enriched in monoculture (Fig S2A). Certain mutations in *E. coli rpoC* are
known to generate polyauxotrophies (28, 29).
186

187
188
189 **Fig S2. Mutations enriched in *E. coli* PFM2 (A) and *R. palustris* CGA676 (B) in**
190 **long-term monocultures and cocultures.** Each data point represents a mutation
191 observed at a frequency >0.1 in an experimental line. Generation: gen.
192
193 No adenine auxotrophs were isolated from cocultures and all coculture isolates grew in
194 defined media, suggesting that our coculture conditions might not favor emergence of
195 other dependencies between subpopulations. Overall, we conclude that *E. coli* adenine
196 auxotrophs are not a major subpopulation in any long term coculture despite the
197 availability of adenine.

198
199
**Fig 2. *E. coli* adenine auxotrophs are not prevalent in long-term cocultures. A.**
Long-term monoculture and coculture conditions. OAcs, organic acids. **B.** Enriched *E.*
*coli* mutations in genes for nucleobase metabolism and regulation of nitrogen
metabolism for long-term monocultures and cocultures. Each point represents a
mutation frequency in a given gene for a given evolutionary line. **C.** Random evolved *E.*
*coli* isolates (generation 650; n=4) were screened for adenine auxotrophy in liquid
monoculture conditions with and without 50 μM adenine. Ancestral strain (Anc) and
*ΔpurH* (*Δ*H) mutant values are also included (n=3). Each point represents a single
measurement for a single isolate (replicate measurements were not made for any
isolate). Each line connects measurements for the same isolate grown with and without
adenine.

**Cross-fed adenine represses purine synthesis gene expression.** One possible
explanation for why *E. coli* purine auxotrophs did not emerge in coculture is because
adenine availability repressed gene expression, thus lowering the purine biosynthesis
gene costs. In support of this notion, we previously saw that *E. coli* MG1655 down-
regulated several purine synthesis genes in coculture versus monoculture (22). To
determine if the same is true for PFM2, we quantified *purH* transcripts by RT-qPCR.
Coculture PFM2 *purH* levels were 20% of those in monoculture (Fig 3A). Adding
adenine to monocultures resulted in a *purH* transcript level that was 12% of that

220 observed without adenine, suggesting that the low expression in coculture was due to
221 adenine availability (Fig 3A).
222
*R. palustris* CGA676 also showed less *purH* transcript in coculture, 50% of that
observed in monoculture (Fig 3B). It is unclear how *E. coli* would influence repression of
*R. palustris purH* expression. However, the lower expression supports previous findings
that *E. coli* does not stimulate *R. palustris* adenine production (20). Adenine availability
and expression of purine biosynthesis genes curiously did not lead to purine
biosynthesis gene mutations in *R. palustris* monocultures. It is possible that *R. palustris*
lacks effective adenine uptake mechanisms. In a separate study, we found that adenine
can be toxic to *R. palustris*, and purine externalizing strains, including the parent of
CGA676 are more resistant to exogenous adenine (Chuang and McKinlay, in prep).
Thus, CGA676 might be pre-dispositioned for adenine externalization rather than
uptake.
234
**Fig 3. Purine biosynthesis genes are down-**
**regulated in coculture.** RT-qPCR
quantification of *purH* transcript levels in *E. coli*
PFM2 (**A**; relative to *hcaT*) and *R. palustris*
CGA676 (**B**; relative to *fixJ*). Adenine (+ ade)
was added to a final concentration of 0.1 mM to
ensure that it was not used up during culture
growth. Values are the mean ± SD, n = 3-4.
Statistically significant differences from the
monoculture condition for each strain were
determined using an unpaired two-tail t-test;
***, $p < 0.001$; ****, $p < 0.0001$.

247
***R. palustris* siderophore gene loss is correlated with expression.** We looked for
other correlations of gene retention and low gene expression in our long-term cultures.
A standout was *R. palustris* siderophore genes which accumulated mutations in
monoculture but not coculture (Fig 4A, B; Fig S2). We previously saw that *R. palustris*
down-regulated siderophore synthesis genes in coculture with *E. coli* MG1655, relative
to monocultures (22). We verified that this trend was also true for *R. palustris* in
coculture with PFM2; RT-qPCR analysis showed that the transcript levels for the
siderophore synthesis gene RPA2390 was 20% in coculture compared to monoculture
(Fig 4C). Adding soluble iron (ammonium ferric citrate) to monocultures decreased
RPA2390 expression to 3% of that observed without added iron, suggesting that the low
expression in coculture was a response to enhanced iron availability in coculture (Fig
4C).
260
*R. palustris* might down-regulate siderophore production in coculture if *E. coli* facilitates
iron acquisition. Others have speculated that *R. palustris* can use siderophores from
other bacteria, because its genome encodes multiple transporters but only one
siderophore synthesis cluster for petrobactin-like siderophores (30, 31); *E. coli* produces
enterobactin siderophores (32). In agreement with this speculation, *E. coli* siderophore

gene expression seemed to respond to iron loss to *R. palustris*; the *entF* transcript level
was 3.3-fold higher in coculture than in monoculture (Fig. 4D). These observations
support a notion that *R. palustris* can use *E. coli* siderophores, however the actual
molecular nature of this relationship is pending a thorough investigation.
270
Despite the increase in *E. coli entF* expression in coculture, *E. coli* siderophore mutants
were rare and inconsistent across coculture lines and were not observed in long-term
monocultures (Table S2, S4). It is possible that this expression range is not enough to
pose a fitness cost, perhaps due to a low iron requirement under fermentative
conditions (e.g., little use for iron-containing cytochromes). In agreement with this
notion, transposon insertions in *E. coli* MG1655 siderophore genes had mainly neutral
effects on fitness in monocultures and cocultures (21).

278
279
**Fig 4. Siderophore gene loss is prevalent in *R. palustris* monocultures where**
**gene expression is high. A, B.** *R. palustris* CGA676 mutation frequencies in
siderophore synthesis genes for each monoculture (**A**) or coculture line (**B**) where
mutations were observed (no mutations were observed in lines 21, 25, and 27). Repeat
colors in a given graph indicate different mutations in the same gene. **C, D.** RT-qPCR
quantification of siderophore synthesis genes in *R. palustris* (**C**; RPA2390 relative to
*fixJ*) and in *E. coli* (**D**; *entF* relative to *hcaT*). Values are the mean ± SD, n = 3.
Statistically significant differences from the monoculture (mono) condition for each strain
were determined using an unpaired two-tail t-test; *, $p < 0.05$; **, $p < 0.01$; co, coculture,
Fe; ammonium ferric citrate.
290
291 **Discussion**
292
In long-term cocultures, the availability of cross-fed adenine, and possibly siderophores,
did not result in corresponding gene loss in a recipient. Our data suggest that the
externalized resource repressed recipient gene expression, thereby lowering gene cost
and selective pressure for gene loss. Conversely, siderophore loss-of-function
mutations accumulated in *R. palustris* monocultures where gene expression was high.
Although there are likely situations where cross-feeding promotes gene loss, there are

also likely situations where cross-feeding instead promotes gene retention through
repressing gene expression.
301
Our results do not counter the BQH, which posits that gene loss will occur if the benefit
of acquiring resources from a neighbor outweighs the cost of gene retention (1). If the
cost of gene retention is low, then gene loss would only have neutral fitness effects,
rather than beneficial effects. Others have suggested that low gene expression is not
enough to promote gene retention because the maintenance and activity of sensor and
regulatory proteins can still drive of gene loss (3). However, the fitness benefits of
mutants lacking biosynthetic genes suggests that benefits can be realized while
regulatory circuits remain intact (5-9). The association between gene cost and gene
expression is well-documented (10-16). Our data, and data from others, suggest that
there are cases where resource availability can lower gene cost simply by repressing
gene expression. For example, repression of an antibiotic resistance gene in *E. coli*
eliminated the fitness cost of that gene (10). In another example closer to the conditions
of the current study, iron-limitation that prompted high siderophore production by *P.*
*aeruginosa* led to a higher frequency of siderophore-deficient mutants compared to iron-
rich conditions where siderophore production was low (11). The *P. aeruginosa* results
were presented in the context of cheating, where gene loss is associated with
exploitation that harms the producer, unlike BQH beneficiaries that should not harm the
producer (1). Yet, the conditions leading to the emergence of cheaters and BQH
beneficiaries are analogous (1). As such, many lessons from literature on cheaters can
apply to BQH scenarios.
322
Insights into BQH gene-loss can also come from resource-rich monocultures, with the
caveat that nutrient access might be well above that from a cross-feeding partner. For
example, repression of gene expression in long-term nutrient-rich *E. coli* monocultures
could explain why the loss of some catabolic genes was not observed in parallel lines
and occurred without fitness effects (33). Repression of gene expression will shift the
cost of a gene towards neutral. Gene loss with neutral fitness effects can still occur, but
enrichment of those mutations would require hitchhiking with a beneficial mutation or
passage through a severe population bottleneck. Cost-neutral gene loss, as opposed to
beneficial gene loss, might also require time, and possibly spatial structure, that were
not part of our experimental design. Should cost-neutral auxotrophic mutants emerge in
nature, the chances of rescue via cross-feeding from a neighbor could be high (34, 35).
334
The extent to which gene repression can promote gene retention will also depend on
the level of expression strength of repression; as noted above, relatively low expression
might explain the lack of *E. coli* siderophore gene mutations. Gene expression can be
noisy, and that noise can be costly and subject to selective optimization in a stable
nutrient regime (36, 37). Noisy gene expression might explain why some engineered
auxotrophs have greater fitness over a prototrophic parent (5-9). In long-term *E. coli*
monocultures, loss-of-function mutations for maltose catabolism were arguably
associated with weak gene repression in the presence of glucose (38). During growth
on glucose, ancestral expression of the maltoporin was half that compared to during
growth on maltose. Several evolved isolates carried mutations that prevented growth on

maltose. However, in an evolved isolate that retained the ability to grow on maltose,
maltoporin expression with glucose was 5% of that with maltose (38), suggesting that
stronger repression contributed to gene retention.

The level of nutrient availability also likely affects the level of gene repression. In other
words, there might be a resource concentration range over which BQH gene loss would
occur. For example, *Acinetobacter baylyi* histidine autotrophs had a fitness advantage
over a prototrophic parent when there was $\leq$ 50 µM histidine, but the advantage was
reversed as histidine levels reached 100-200 µM (6), possibly due to stronger gene
repression in the parent, though this is speculation on our part. One might also consider
that the responses of a prototrophic strain versus an otherwise clonal auxotroph to an
extracellular nutrient can vary widely; whereas an auxotroph would benefit from the
nutrient, the prototroph might experience beneficial, neutral or even detrimental fitness
effects (6). The extent to which loss of a biosynthetic gene can have compounding
fitness effects by affecting expression of other genes deserves investigation.

Our work presents a caveat to the popular expectation of cross-feeding as a driver of
gene loss. Cross-feeding can instead promote gene retention when resource availability
is high enough to repress gene expression and thus lower the cost of a gene. The
likelihood of gene loss then becomes the same as any other gene with a neutral cost.

**MATERIALS AND METHODS**

**Bacterial strains.** *R. palustris* CGA676 is derived from CGA0092 (39) and carries a
*nifA\** mutation that causes $NH_4^+$ excretion under $N_2$-fixing conditions (19, 40). *E. coli*
PFM2 (41, 42) $\Delta purH$::km$^R$ was made via lambda Red recombination (43) using
constructs amplified from KEIO mutants (44) using primers YCC29 (GCG CAA ACG
TTT TCG TTA CAA TGC) and YCC30 (TGC ATT ACC CGG AGC AAC). FLP-mediated
excision was used to remove the kanamycin resistance cassette (km$^R$) to generate the
$\Delta purH$ strain (43).

**Growth conditions.** Anoxic media in test tubes were prepared by bubbling $N_2$ through
10 ml of media in 27-ml anaerobic test tubes, then sealing with rubber stoppers and
aluminum crimps prior to autoclaving. Monocultures and cocultures were grown
horizontally at 30°C with light from a 45 W halogen bulb (430 lumens) and shaking at
150 rpm in minimal M9-derived coculture medium (MDC) (19) with either (i) *E. coli*
monocultures: 25 mM glucose, 10 mM $NH_4Cl$, and cation solution (100X stock: 100 mM
$MgSO_4$ and 10 mM $CaCl_2$); *R. palustris* monocultures: 20 mM sodium acetate and 10
mM $NH_4Cl$; or cocultures: 25 mM glucose and cation solution. Cultures with plasmid-
carrying strains were also supplemented with 100 µg/ml gentamycin or 25 µg/ml
chloramphenicol as appropriate. Starter cultures were inoculated with single colonies.
*R. palustris* starter cultures were grown in MDC with 20 mM acetate and 10 mM $NH_4Cl$.
*E. coli* starter cultures were grown aerobically in lysogeny broth, with 30 µg/ml
kanamycin (km) when appropriate. *E. coli* starter cultures were washed twice in 1 ml
MDC prior to inoculating test cultures or bioassays. Cocultures were inoculated with 0.1
ml each of *R. palustris* and *E. coli* to an initial optical density ($OD_{660}$) of ~0.003 each.

Cultures in 96-well plates were treated similarly except that oxic stock solutions were used to prepare 0.2 ml volumes in each well. Anoxic conditions were achieved by sealing plates inside a BD GasPak EZ large incubation container with 2 anaerobic sachets.

**Experimental evolution.** Founder monocultures of *E. coli* PFM2 and *R. palustris* NifA* CGA676 were grown from a single colony in anoxic MDC with either 25 mM glucose, cation solution, and 3 mM NH₄Cl for PFM2 or 20 mM sodium acetate for CGA676. A single founder monoculture was then used to inoculate 10 monocultures and 10 cocultures. All cultures were grown horizontally without shaking at 30°C with light in MDC. PFM2 monocultures were supplemented with 10 mM glucose, cation solution, 25 mM NaCl, and 2.3 mM NH₄Cl. CGA676 monocultures were supplemented with 25 mM glucose, cation solution, 8 mM disodium succinate, 7.3 mM sodium acetate, 0.25 mM sodium formate, 1.4 mM sodium lactate, and 6.3 mM ethanol. Cocultures were supplemented with 25 mM glucose, cation solution, and 25 mM NaCl. Every 7 days, cultures were vortexed and 0.25 ml was transferred to fresh medium. About every 5 transfers, stocks were frozen in 25% glycerol at -80°C and separate cell pellets from 1 ml samples were frozen for gDNA extraction.

**Analytical procedures.** Cell densities were measured via turbidity ($OD_{660}$) using a Genesys 20 visible spectrophotometer (Thermo-Fisher).

**Competition assays.** Competition assays were conducted in an invasion-from-rare format to consider coexistence by the mutual invasion criterion, where each population can increase when rare (25, 27). Cocultures were started from various initial frequencies (targeting 0.01 – 0.99) of each *E. coli* strain (WT vs Δ*purH* or WT vs Δ*purH*::km$^R$) for a total initial cell density of ~$10^6$ colony forming units (CFU) / ml. *R. palustris* CGA676 was inoculated to an initial density of ~$10^6$ CFU / ml. Frequencies were determined upon inoculation and after 5 days. WT and Δ*purH* mutants were distinguished by counting CFUs on M9 agar with cations and 25 mM glucose and on LB agar and then determining Δ*purH* populations from the difference. When Δ*purH*::km$^R$ mutants were used LB agar included km to allow for direct determination of population size. Change in frequency = (*E. coli* Δ*purH* / (*E. coli* WT + *E. coli* Δ*purH*))$_{final}$ – (*E. coli* Δ*purH* / (*E. coli* WT + *E. coli* Δ*purH*))$_{initial}$ (26).

**Genome sequencing and mutation analysis.** gDNA was purified from cells using a Qiagen DNeasy Blood and Tissue kit following the manufacturer's instructions. Lysis was facilitated after resuspension by adding proteinase K (50 μg/ml final), and incubating at 56°C for 10 min. RNaseA (4 μl, Promega) was then added and the lysate was incubated for 2 min before proceeding.

DNA fragment libraries were made using a NextFlex Bioo Rapid DNA kit and libraries were sequenced using Illumina NextSeq 500 150x150 paired-end runs by the IU Center for Genomics and Bioinformatics. Paired-end reads were pre-processed for quality with cutadapt 3.4 (45) with the following options: -a AGATCGGAAGAGC -A AGATCGGAAGAGC ; -q 15,10; -u 6. Mutations were called using breseq v. 0.32.0 on

polymorphism mode (46). *E. coli* monoculture population sequences were mapped to the MG1655 genome (accession NC_000913); *R. palustris* monoculture population sequences were mapped to a concatenated reference genome consisting of the CGA009 chromosome (accession BX571963), and its plasmid pRPA (accession BX571964). Co-culture sequences were mapped to a concatenation of the *E. coli* and *R. palustris* reference genomes. Polymorphisms that co-occurred in both the monoculture and co-culture datasets were filtered and maintained as a subset to enrich for the most informative variants representing treatment differences (Table S1-4). This filtering step also removed sequence differences between the reference sequences and those of the experimental strains used. Variants were prioritized as mutations of interest if they were detectable at the final two sequencing timepoints and co-occurred across multiple populations in the same locus. All mutations can be found in Tables S1-4. Locus tag conversions can be found in Table S5.

**Reverse transcription quantitative real-time PCR (RT-qPCR).** Cultures received 100 µM adenine or ammonium ferric [iron(III)] citrate as indicated. Cultures were harvested in exponential phase at 0.6-0.8 $OD_{660}$ except for *E. coli* monocultures (+/- adenine experiment), which were harvested at 0.3-0.4 $OD_{660}$ to avoid adenine depletion. Cultures were chilled on ice and pelleted by centrifugation. Lysis, RNA purification, and cDNA generation were performed exactly as described (20). Standard curves were generated using gDNA. Transcripts were quantified as described (20) using the appropriate primers (Table S6) with a Mastercycler ep *realplex* real-time PCR system (Eppendorf). Data was analyzed by *realplex* software using Noiseband. Specificities were validated by melting curves and by the presence of a single band on an agarose gel.

**Statistical analyses.** Statistical analyses of growth and RT-qPCR data were performed using Graphpad Prism (v10).

## ACKNOWLEDGEMENTS

This work was supported in part by US Army Research Office grants W911NF-14-1-0411 and W911NF-17-1-0159, the National Science Foundation CAREER award MCB-1749489 to JBM, and National Institutes of Health grants R35GM128674 to ABD and R35GM150625 to MB. Supercomputing resources were supported in part by Lilly Endowment, Inc., through its support for the IU Pervasive Technology Institute.

We are grateful to J. Gliessman, A. Cairo, S. Kamaran, J. Mazny, and N. Ramli, for maintaining long-term cultures. We thank P. Foster for PFM2 and helpful advice. We also thank A. Drummond, J. Drummond, J.P. Gerdt, J. Lennon, M. Lynch, T. Romeo, and the McKinlay lab for helpful advice.

## References

1.    J. J. Morris, R. E. Lenski, E. R. Zinser, The Black Queen Hypothesis: Evolution of dependencies through adaptive gene loss. *mBio* **3**, e00036-00012 (2012).

2.    M. Ghoul, A. S. Griffin, S. A. West, Toward an evolutionary definition of cheating.
*Evolution* **68**, 318-331 (2014).

3.    G. D'Souza *et al.*, Ecology and evolution of metabolic cross-feeding interactions in
bacteria. *Nat Prod Rep* **35**, 455-488 (2018).

4.    S. Pande, C. Kost, Bacterial unculturability and the formation of intercellular metabolic
networks. *Trends Microbiol.* **25**, 349-361 (2017).

5.    S. Zamenhof, H. H. Eichhorn, Study of microbial evolution through loss of biosynthetic
functions: establishment of "defective" mutants. *Nature* **216**, 456-458 (1967).

6.    G. D'Souza *et al.*, Less is More: Selective advantages can explain the prevalent loss of
biosynthetic genes in bacteria. *Evolution* **68**, 2559-2570 (2014).

7.    D. Dykhuizen, Selection for tryptophan auxotrophs of *Escherichia coli* in glucose-limited
chemostats as a test of the energy conservation hypothesis of evolution. *Evolution* **32**,
125-150 (1978).

8.    S. Pande *et al.*, Fitness and stability of obligate cross-feeding interactions that emerge
upon gene loss in bacteria. *ISME J* **8**, 953-962 (2013).

9.    G. D'Souza, C. Kost, Experimental evolution of metabolic dependency in bacteria. *PLoS*
*Genet.* **12**, e1006364 (2016).

10.    T. N. Nguyen, Q. G. Phan, L. P. Duong, K. P. Bertrand, R. E. Lenski, Effects of carriage
and expression of the Tn10 tetracycline-resistance operon on the fitness of *Escherichia*
*coli* K12. *Mol. Biol. Evol.* **6**, 213-225 (1989).

11.    Z. Dumas, R. Kümmerli, Cost of cooperation rules selection for cheats in bacterial
metapopulations. *J. Evol. Biol.* **25**, 473-484 (2012).

12.    F. C. Neidhardt, J. L. Ingraham, M. Schaechter, *Physiology of the bacterial cell: a*
*molecular approach* (Sinauer Associates Inc, Sunderland, MA, 1990).

13.    M. Lynch, G. K. Marinov, The bioenergetic costs of a gene. *Proc Natl Acad Sci U S A*
**112**, 15690-15695 (2015).

14.    D. M. Stoebel, A. M. Dean, D. E. Dykhuizen, The cost of expression of Escherichia coli
lac operon proteins is in the process, not in the products. *Genetics* **178**, 1653-1660
(2008).

15.    E. Dekel, U. Alon, Optimality and evolutionary tuning of the expression level of a
protein. *Nature* **436**, 588-592 (2005).

16.    Z. Wu *et al.*, Expression level is a major modifier of the fitness landscape of a protein
coding gene. *Nat Ecol Evol* **6**, 103-115 (2022).

17.    F. Harrison, J. Paul, R. C. Massey, A. Buckling, Interspecific competition and
siderophore-mediated cooperation in *Pseudomonas aeruginosa*. *ISME J* **2**, 49-55 (2008).

18.    A. R. T. Figueiredo, A. Wagner, R. Kümmerli, Ecology drives the evolution of diverse
social strategies in *Pseudomonas aeruginosa*. *Mol. Ecol.* **30**, 5214-5228 (2021).

19.    B. LaSarre, A. L. McCully, J. T. Lennon, J. B. McKinlay, Microbial mutualism dynamics
governed by dose-dependent toxicity of cross-fed nutrients. *ISME J* **11**, 337-348 (2017).

20.    Y.-C. Chuang *et al.*, Bacterial adenine cross-feeding stems from a purine salvage
bottleneck *ISME J* **18**, wrae034 (2024).

21.    B. LaSarre, A. M. Deutschbauer, C. E. Love, J. B. McKinlay, Covert cross-feeding
revealed by genome-wide analysis of fitness determinants in a synthetic bacterial
mutualism. *Appl Environ Microbiol* **86**, e00543-00520 (2020).

22. A. L. McCully *et al.*, An *Escherichia coli* nitrogen starvation response is important for
mutualistic coexistence with *Rhodopseudomonas palustris*. *Appl Environ Microbiol* **84**,
e00404-00418 (2018).

23. K. F. Jensen, The *Escherichia coli* K-12 "wild types" W3110 and MG1655 have an rph
frameshift mutation that leads to pyrimidine starvation due to low pyrE expression levels.
*J. Bacteriol.* **175**, 3401-3407 (1993).

24. A. S. Bagnara, L. R. Finch, The effects of bases and nucleosides on the intracellular
contents of nucleotides and 5-phosphoribosyl 1-pyrophosphate in *Escherichia coli*. *Eur.*
*J. Biochem.* **41**, 421-430 (1974).

25. T. N. Grainger, J. M. Levine, B. Gilbert, The invasion criterion: a common currency for
ecological research. *Trends Ecol. Evol.* **34**, 925-935 (2019).

26. S. P. Hammarlund, J. M. Chacón, W. R. Harcombe, A shared limiting resource leads to
competitive exclusion in a cross-feeding system. *Environ. Microbiol.* **21**, 759-771 (2019).

27. P. Chesson, Mechanisms of maintenance of species diversity. *Annu Rev Ecol Syst* **31**,
343-366 (2000).

28. D. Satory, J. A. Halliday, P. Sivaramakrishnan, R. C. Lua, C. Herman, Characterization
of a novel RNA polymerase mutant that alters DksA activity. *J. Bacteriol.* **195**, 4187-
4194 (2013).

29. L. Brown, D. Gentry, T. Elliott, M. Cashel, DksA affects ppGpp induction of RpoS at a
translational level. *J. Bacteriol.* **184**, 4455-4465 (2002).

30. F. W. Larimer *et al.*, Complete genome sequence of the metabolically versatile
photosynthetic bacterium *Rhodopseudomonas palustris*. *Nat. Biotechnol.* **22**, 55-61
(2004).

31. O. Baars, F. M. M. Morel, X. Zhang, The purple non-sulfur bacterium
*Rhodopseudomonas palustris* produces novel petrobactin-related siderophores under
aerobic and anaerobic conditions. *Environ. Microbiol.* **20**, 1667-1676 (2018).

32. L. Cavas, I. Kirkiz, Characterization of siderophores from E*scherichia coli* strains
through genome mining tools: an antiSMASH study. *AMB Express* **12**, 74 (2022).

33. N. Leiby, C. J. Marx, Metabolic erosion primarily through mutation accumulation, and
not tradeoffs, drives limited evolution of substrate specificity in *Escherichia coli*. *PLoS*
*Biol.* **12**, e1001789 (2014).

34. S. Giri, G. Yousif, S. Shitut, L. Oña, C. Kost, Prevalent emergence of reciprocity among
cross-feeding bacteria. *ISME Commun* **2**, 1-7 (2022).

35. S. Giri *et al.*, Metabolic dissimilarity determines the establishment of cross-feeding
interactions in bacteria. *Curr. Biol.* **31**, 5547-5557 e5546 (2021).

36. J. B. McKinlay, G. M. Cook, K. Hards, Microbial energy management-A product of three
broad tradeoffs. *Adv. Microb. Physiol.* **77**, 139-185 (2020).

37. E. J. O'Brien, J. Utrilla, B. O. Palsson, Quantification and classification of *E. coli*
proteome utilization and unused protein costs across environments. *PLoS Comput Biol*
**12**, e1004998 (2016).

38. L. Pelosi *et al.*, Parallel changes in global protein profiles during long-term experimental
evolution in *Escherichia coli*. *Genetics* **173**, 1851-1869 (2006).

39. B. E. Mazny, O. F. Sheff, B. LaSarre, A. McKinlay, J. B. McKinlay, Complete genome
sequence of *Rhodopseudomonas palustris* CGA0092 and corrections to the *R. palustris*
CGA009 genome sequence. *Microbiol Resour Announc* **12**, e0128522 (2023).

40. J. B. McKinlay, C. S. Harwood, Carbon dioxide fixation as a central redox cofactor
recycling mechanism in bacteria. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 11669-11675 (2010).
41. P. L. Foster, H. Lee, E. Popodi, J. P. Townes, H. Tang, Determinants of spontaneous
mutation in the bacterium *Escherichia coli* as revealed by whole-genome sequencing.
*Proc Natl Acad Sci U S A* **112**, E5990-5999 (2015).
42. H. Lee, E. Popodi, H. Tang, P. L. Foster, Rate and molecular spectrum of spontaneous
mutations in the bacterium *Escherichia coli* as determined by whole-genome sequencing.
*Proc Natl Acad Sci U S A* **109**, E2774-2783 (2012).
43. K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in
*Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. USA* **97**, 6640-6645
(2000).
44. T. Baba *et al.*, Construction of *Escherichia coli* K-12 in-frame, single-gene knockout
mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).
45. M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads.
*EMBnet j* **17** (2011).
46. D. E. Deatherage, J. E. Barrick, "Identification of Mutations in Laboratory-Evolved
Microbes from Next-Generation Sequencing Data Using breseq" in Engineering and
Analyzing Multicellular Systems: Methods and Protocols*,* L. Sun, W. Shou, Eds.
(Springer New York, New York, NY, 2014), 10.1007/978-1-4939-0554-6_12, pp. 165-
188.
592

 **Table S6. Primers used in this study**

| Primer | Sequence (5'-3') | Description |
|---|---|---|
| YCC29 | GCG CAA ACG TTT TCG TTA CAA TGC G | 5' of Δ*purH*::*Km* in JW3970 |
| YCC30 | TGC ATT ACC CGG AGC AAC | 3' of Δ*purH*::*Km* in JW3970 |
| YCC80 | TAA GGG AAC CGT GCA TGT G | Forward qPCR primer for *Rp fixJ* |
| YCC81 | GGA TTC GTA CAG CTT GAC CTC | Reverse qPCR primer for *Rp fixJ* |
| YCC97 | ACG TCG TCG TGG TTC TTG | Forward qPCR primer for *Rp purH* |
| YCC98 | CGA AGC CAC CGT CGA TAA A | Reverse qPCR primer for *Rp purH* |
| YCC99 | GCA ACA CGT TCT GCT GAT G | Forward qPCR primer for *Rp RPA2390* |
| YCC100 | CAT TGG TTC TCG GCC TAT CT | Reverse qPCR primer for *Rp RPA2390* |
| YCC91 | AAC CGC ATG GCC CTT ATT | Forward qPCR primer for *Ec entF* |
| YCC92 | GTA TCC AGC AAG CCA AGA AAT G | Reverse qPCR primer for *Ec entF* |
| YCC93 | GTG TTC GAA GGC TTT GAT GG | Forward qPCR primer for *Ec purH* |
| YCC94 | GTG AAG AGC AGC GAC TAT GA | Reverse qPCR primer for *Ec purH* |
| YCC95 | GTC GCA CTT GCC GT AAT C | Forward qPCR primer for *Ec hcaT* |
| YCC96 | GCT GAT GCT GGT GAT GAT TG | Reverse qPCR primer for *Ec hcaT* |