1

Temperature fluctuations influence predictions of landscape-scale patterns of spruce budworm

defoliation

Emily N. Black[a], Deepa S. Pureswaran[b], Katie E. Marshall[a]

[a]Department of Zoology, University of British Columbia, Vancouver, British Columbia, V6T

1Z4, Canada,

[b]Natural Resources Canada, Canadian Forest Service, Atlantic Forestry Centre, 1350, Regent

Street, Fredericton, NB E3C 2G6, Canada

Abstract

The geographic range of insects is heavily influenced by their tolerance for stressful

abiotic conditions, including temperature. While many studies on insect thermal tolerance

consider temperature exposure, the frequency of temperature exposures is emerging as an

important and generally overlooked driver of insect fitness. The (eastern) spruce budworm

(*Choristoneura fumiferana*) is a lepidopteran defoliating pest of coniferous forests across Canada

whose outbreaks lead to large-scale tree mortality. Studies have shown the frequency of

temperature fluctuations affects spruce budworm overwintering survival rates in the laboratory;

however, the influence of temperature fluctuations on spruce budworm defoliation at the

landscape level has not been investigated. We used a species distribution model approach to

evaluate the influence of temperature fluctuations on the distribution and severity of spruce

budworm defoliation. We combined publicly available maps of spruce budworm outbreaks

between 2006-2016 in Quebec with climate, temperature fluctuation, and forest composition

predictors to train a species distribution model. Our model evaluated how predictors influence

spruce budworm defoliation, and compared these results to a model trained without temperature

fluctuations. Additionally, we predicted future spruce budworm defoliation under 2041-2070

climate change conditions using the models trained with and without temperature fluctuation

predictors and compared the results to determine the effect of temperature fluctuations on future

defoliation predictions. We found that model performance improved with the inclusion of

temperature fluctuation predictors, and these predictors ranked highly, relative to predictors in

other categories. The model trained with temperature fluctuation predictors also predicted vastly

different defoliation distribution and severity across Quebec and Ontario than the model trained

without them under climate change conditions. These results reveal the previously overlooked

importance of temperature fluctuations on landscape-scale spruce budworm defoliation and

support their inclusion in insect species distribution models.

Introduction

Environmental conditions determine the spatial distribution, range limits, and fitness of

insects on a regional and landscape scale (Gaston, 2003; Godsoe et al., 2017; Holt, 2009).

According to Hutchinson's ecological niche concept, animals inhabit geographic regions that

meet their abiotic requirements (temperature, pH, radiation), and populations living in conditions

outside their niche face fitness costs that cause local extinction (Holt, 2009; Hortal et al., 2010;

Hutchinson, 1957). In particular, temperature directly affects insect survival, growth,

metabolism, and fecundity, and is therefore a major factor determining species' spatial

distributions and their success within those ranges (Régnière, 2009; Régnière, Powell, et al.,

2012).

Insects living in temperate regions face thermal challenges in their environment which

affect their physiology, and insects must be adapted to these challenges in order persist and

thrive. In regions where temperatures seasonally drop below freezing, insects cannot forage

during the winter, may experience loss of neuromuscular function (Findsen et al., 2014), and risk

tissue damage from freezing and osmotic stress (Storey & Storey, 1989). To survive these

conditions, insects may overwinter in diapause: a long-term state of developmental arrest with a

reduced metabolic rate to conserve energetic resources accumulated during the summer

(Tougeron, 2019). Pre- and post- diapause fitness is highly sensitive to fall, winter, and spring

temperatures (Sinclair, 2015) as unfavourable temperatures can deplete overwintering energy

stores, reduce survival, and induce non-lethal costs such as limited fecundity (Marshall &

Sinclair, 2015; Roe et al., 2024). Insects also have mechanisms to protect against freezing

damage; however, these often lead to trade-offs between conserving resources and protection

against cold damage (Marshall & Sinclair, 2012; Toxopeus & Sinclair, 2018). High seasonal

temperatures also have detrimental effects on insects through induced elevated metabolic rates

and protein denaturation (González-Tokman et al., 2020). This can be overcome with

biochemical and physiological responses (Storey & Storey, 2004); however, as with cold

temperatures these responses incur survival and fitness costs (González-Tokman et al., 2020).

The relationship between stress and temperature is not linear and dependent on many factors;

thus, it is important to study how physiological adaptations to temperature carry over to

environmental responses.

While many physiological and ecological studies use mean temperatures to investigate an

organism's response to temperature across their range, this does not reflect conditions in the wild

where they will experience fluctuations in temperature on multiple time scales (seconds, hours,

days, months, years; Dillon et al., 2016). Temperature fluctuations at different frequencies,

temperatures, seasons, and life stages can have varying and overlapping impacts on insect

physiology and fitness (Marshall & Sinclair, 2012). Frequent temperature fluctuations within an

insect's optimal thermal range can positively impact insect fitness and fecundity by decreasing

energy requirements for growth (Carrington et al., 2013). Previous exposure to high temperatures

in the pea leafminer (*Liriomyza huidobrensis*) leads to higher expression of heat shock proteins

and improved thermotolerance in subsequent heat exposures (Huang et al., 2007). At below-

freezing temperatures, temperature fluctuations can positively or negatively affect insect fitness

depending on the species and effect being studied (Marshall & Sinclair, 2012). Marshall and

Sinclair (2015) investigated the effect of temperature fluctuations on the (eastern) spruce

budworm (*Choristoneura fumiferana*), Lepidoptera: Tortricidae, and found that the frequency of

cold temperature exposures was more detrimental than cold temperature duration or intensity to

overwintering survival. Fluctuations at cold temperatures may also allow for repair of cold-

damaged tissues during winter (Colinet et al., 2006), lead to de-acclimation of insects to winter

conditions that increases the chance of chilling injury (Sobek-Swant et al., 2012), and positively

or negatively affect survival (Marshall & Sinclair, 2010, 2012, 2015). Overall, temperature

fluctuations have a wide range of impacts on insect physiology, survival, and fitness (Marshall &

Sinclair, 2012). However, these findings are rarely extrapolated to studies investigating the

thermal drivers of insect species distributions, including those in temperate regions which will be

impacted by above and below freezing temperature fluctuations (Marshall & Sinclair, 2012,

2015).

104   The spruce budworm is a major North American insect pest found in boreal forests from

Alaska to Newfoundland (Gray, 2008; Marshall & Roe, 2021; Natural Resources Canada, 2021).

Eggs are laid in late summer. First instar larvae hatch and develop into second instars that spin

hibernacula and overwinter in trees until early May. After overwintering, larvae feed voraciously

on coniferous trees before developing into adults. Outbreaks of this pest occur every 30-40 years

and cause landscape scale defoliation: the larvae will eat expanding new buds of host trees

(balsam fir, white spruce, black spruce; Mattson et al., 1991) in significant quantities. Several

years of consecutive defoliation eventually weakens and kills the trees (Gray, 2008). A recent

outbreak in Quebec resulted in 10.5 million hectares of forest damage via defoliation since 2006

(Ministère des Ressources naturelles et des Forêts, 2023). Outbreaks are preventable if

"hotspots" where populations are increasing are identified in the early stages (Johns et al., 2019;

MacLean et al., 2019), which requires significant effort to understand drivers of outbreaks and

defoliation severity, extensive forest monitoring, and prediction of future defoliation hotspots

(Johns et al., 2019).

Climate, including temperature, can modulate the relationship between insect defoliators

such as spruce budworm and host trees, which may drive trends in defoliation across the

landscape. Pureswaran et al. (2019) found that warming temperatures are expected to improve

synchrony between host tree budburst and budworm larval emergence, causing defoliation earlier

in the season and increasing suitability of black spruce as a host (which has generally escaped

defoliation via delayed budburst; Bellemin-Noël et al., 2021). Drought may also increase

defoliation by spruce budworm, as defoliators are expected to perform better during droughts due

to higher soluble nitrogen concentrations in tree foliage (Jactel et al., 2012). In the closely-

related western spruce budworm (*Choristoneura occidentalis* Freeman), drought is associated

with increased risk of defoliation (Flower et al., 2014; Xu et al., 2019); although, this may not be

the case in *Choristoneura fumiferana* (Moise et al., 2019).

Extensive aerial sampling by provincial governmental agencies records the location and

severity of spruce budworm defoliation across Canada, and this scale of sampling lends itself

well to the generation of spruce budworm species and defoliation distribution models (Ministère

des Ressources naturelles et des Forêts, 2023; Natural Resources Canada, 2021). Predictive

distribution models of spruce budworm outbreaks have evaluated the importance of different

environmental factors, including temperature, on outbreak likelihood and defoliation risk on a

landscape scale (Gray, 2008; Li et al., 2020), and have predicted the effects of climate change on

outbreak distribution and severity (Candau & Fleming, 2005; Johns et al., 2019; Régnière, St-

Amant, et al., 2012). In particular, Gray (2008) found that patterns in outbreak distribution were

partially explained by climate variables and forest composition, and that outbreaks were

expected to be longer and more severe under climate change. Candau & Fleming (2005) also

found that the distribution of spruce budworm outbreaks are strongly correlated with

temperature: in particular, winter maximum and minimum temperature.

While climate change will affect overall climatic conditions in temperate North America,

temperature fluctuations are also expected to shift due to climate change and may affect future

spruce budworm defoliation. Wang & Dillon (2014) find that global temperature variability has

increased under climate change, and that this effect is particularly pronounced in temperate and

polar regions. However, this varies by temperature threshold, season, and biome, as the

frequency of extreme cold temperatures (below -20°C) and freezing temperatures (below 0°C)

are expected to decrease in the boreal forest in 2081-2100 climate conditions (Contosta et al.,

2024). The disappearance of extreme cold and freezing temperatures and changes in overall

temperature variability will have far-reaching consequences across the boreal forest biome,

including boreal forest pests such as the spruce budworm (Contosta et al., 2024; Pureswaran et

al., 2019). However, temperature fluctuations have not yet been incorporated into methods such

as species distribution models which can identify the temperature thresholds with the greatest

influence on pest outbreaks and evaluate their role in driving observed distributions (Colinet et

al., 2006; Marshall & Sinclair, 2015). Thus, it is not yet understood how temperature fluctuations

may influence the distribution of spruce budworm defoliation now and in the future, nor their

importance relative to other factors such as host tree distribution and other climatic conditions.

Here, we use a species distribution model framework to investigate how temperature

fluctuations influence historical patterns of spruce budworm defoliation across Quebec from

2006-2016 and predictions of defoliation using a 2041-2070 climate change scenario. Based on

previous work by Marshall and Sinclair (2015) investigating how the number and frequency of

temperature fluctuations affects spruce budworm fitness, we hypothesize that temperature

fluctuations influence the landscape-scale distribution of spruce budworm defoliation. Using

historical records of defoliation collected in Quebec, we trained a spatial random forest

regression model to predict the severity of historical defoliation using three selected general

predictor categories (climate, forest, temperature fluctuations). Using spatial cross-validation we

found that temperature fluctuations are informative to the model, and that predictors in the

temperature fluctuation category are similar in importance as climate predictors and are often

more important than forest predictors. When predicting defoliation using a 2041-2070 climate

change scenario, we find that the model is sensitive to temperature fluctuation predictors, as their

inclusion drastically changes the overall distribution and severity of defoliation compared to a

model trained without temperature fluctuation predictors. Overall, these results support our

hypothesis, and indicate that temperature fluctuations influence the landscape-scale distribution

of spruce budworm defoliation.

Methods
i)   Collecting and aggregating spruce budworm defoliation data

The Ministère des Ressources Naturelles et des Forêts in Quebec conducts annual aerial

flyovers of forests to record damage due to major insect pests, the effects of disturbances such as

forest fires, and the effectiveness of current pest control measures (Ministère des Ressources

naturelles et des Forêts, 2023) . Survey locations are based on the locations of insect damage

from the previous year, and predictive modelling of future outbreaks (Ministère des Ressources

naturelles et des Forêts, 2023). The location and severity of spruce budworm *(Choristoneura*

fumiferana) defoliation patches are recorded during this survey (Ministère des Ressources

naturelles et des Forêts, 2023). Spruce budworm defoliation patches collected from 1992-2006

($n_{polygon}$=4,958), 2007-2013 ($n_{polygon}$=54,867), and 2014-2021 ($n_{polygon}$=26,2826) are stored as

polygons in a publicly available file (see supp. table 1 for links to data sources). The polygons

have the following attributes: year of defoliation, defoliation severity level and score, hectares of

defoliation, Quebec administrative region, and the geometric shape of the defoliation polygon.

Defoliation severity scores in Quebec were 1 – mild canopy damage, 2 – moderate canopy

damage, and 3 – severe canopy damage. Maps of Ontario defoliation were also collected, but

were not used in subsequent analyses due to difficulties standardizing with the Quebec data

(supp. methods, Figure S1).

We aggregated defoliation polygons for defoliation in Quebec on a raster grid using the

terra package (Hijmans et al., 2021; Figure 1). We mapped the presence and absence of

defoliation, and the cumulative defoliation severity of each grid cell (i.e., how susceptible it was

to spruce budworm damage) by summing the annual defoliation severity score of all polygons

covering the centroid of a grid cell and assigning that value to the cell. The raster of spruce

budworm defoliation in Quebec covered an area from -79.57°W to -61.55°W, and 45.42°N to

51.73°N. Grids had an initial resolution of approximately 412m × 412m; however, grids used in

analysis were aggregated to a resolution of 300 arcsec (approximately 6.5km × 6.5km) to match

predictor resolution.

Government records of spruce budworm defoliation indicate that the most recent

outbreak in Eastern Canada began in 2006 (Natural Resources Canada, 2021); thus, we used

defoliation patches from this year onwards in our analysis. To confirm that defoliation polygons

aligned with known outbreak years, we plotted the cumulative amount of defoliation in each grid

cell in which defoliation was observed between 1997 and 2021. Defoliation increased

exponentially across Quebec 2006 onwards (Figure S2). While the outbreak beginning in 2006 is

ongoing (Natural Resources Canada, 2021), high-resolution daily climate records were only

available until 2016. Thus, we only included defoliation from 2006-2016 in analyses.

ii) Collecting and aggregating predictor data

We chose to evaluate the effect of four general predictor categories on defoliation

presence and severity: forest, climate, degree days, and temperature fluctuations. For a list of

data sources, see supp. table 1; for a list of all predictors and their descriptions and assigned

predictor categories, see supp. table 2.

For forest predictors, we obtained raster grids of the forest composition of Canada,

including the distribution of spruce budworm host trees (white spruce, black spruce, balsam fir)

and percent tree cover from the National Forest Inventory (NFI), a collaborative effort by

federal, provincial, and territorial government agencies to generate continuous records of forest

composition across Canada (*National Forest Inventory*, 2021). All grids had an initial resolution

of 250m × 250m, and were upscaled to approximately 6.5km × 6.5km.

For climate, temperature fluctuation, and degree day predictors, we obtained minimum

and maximum daily temperatures and daily precipitation from 2006-2016 across eastern Canada

from the CHELSA (Climatologies at High Resolution for the Earth's Land Surface Areas)

dataset. CHELSA calculated global daily fine-resolution temperature and precipitation values up

to 2016 using downscaling of atmospheric temperatures and climatologies (Karger et al., 2017,

2021). Although many high-resolution climate products exist for our region of interest (e.g. T.

Wang et al., 2016), we selected CHELSA primarily due to its generation of global daily

temperatures which allowed us to generate accurate daily temperature fluctuation counts across a

broad geographic region, as well as its generation of global monthly climate variables under

several climate change scenarios. Additionally, CHELSA's global distribution allows our

methods to be more easily extrapolated to other species and regions of interest across the globe.

From the daily CHELSA temperature values, we calculated the annual mean temperature, annual

mean minimum and maximum temperature, and the absolute annual minimum and maximum

temperature. We calculated these values for each month of the year, averaged across 2006-2016.

We also obtained daily precipitation values from the CHELSA dataset to calculate the mean

yearly and monthly precipitation. For the degree day predictors, using the CHELSA data we

summed the difference between mean daily temperatures and 0°C annually and in each month,

averaged across 2006-2016. Maps of historical climate and degree day data covered an area from

-83.57°W to -57.55°W, and 41.42°N to 55.73°N, and had an initial resolution of 300 arcsec

(approximately 6.5km × 6.5km).

To calculate temperature fluctuation predictors across the landscape, we calculated

repeated temperature exposures, or the number of times the temperature in each grid cell crossed

a particular temperature threshold daily each month (for example, the number of times the

temperature crossed 0°C daily in January) based on laboratory experiments performed by

Marshall and Sinclair (2015). To calculate this, we interpolated the daily minimum and

maximum temperatures from CHELSA to simulate change in daily temperature across an entire

year at each grid cell. Then, we summed the number of times the temperature at each point

crossed above a particular threshold (e.g., 0°C, -10°C) to calculate the frequency of cold

exposure. We then averaged these values across 2006-2016 and transformed the points into a

raster map with a resolution of 300 arcsec (approximately 6.5km × 6.5km). Overall, we

generated a total of 169 climate, forest, and temperature fluctuation predictors (see supp. table 2

for description of all predictors).

To predict defoliation presence and severity in the future, we re-generated model

predictors under simulated climate change conditions from 2041-2070. Since tree species in the

eastern Canadian boreal forest show limited migration ability under climate change conditions

(Boisvert-Marsh et al., 2022), we used the same tree distributions as the historical models

(*National Forest Inventory*, 2021).

To generate climate, degree day, and temperature fluctuation variables under climate

change conditions, we used the monthly CHELSA CMIP5-predicted climatologies for 2041-

2070 (Karger et al., 2017). These represent monthly temperatures in an example year under

climate change scenarios, and were generated using the Geographic Fluid Dynamics Laboratory

Earth System 4 (GFDL-ESM4; 10). We selected climatologies from the SSP370 climate

scenario, which predicts a moderate to high amount of future global emissions (*The SSP*

Scenarios, n.d.). To simulate daily temperatures under this climate change scenario, we used the

chillR package (Luedeling et al., 2023) to generate daily stochasticity in the monthly CHELSA

climate change projections. The *chillR* package uses the CHELSA monthly projections to guide

future climatic trends in the stochastic data, and historical daily temperatures from 2010-2016

CHELSA data to generate stochastic daily minimum and maximum temperatures that follow

annual temperature trends (Figure S3). From the *chillR* daily stochastic weather output, we

generated annual and monthly climatic predictors, degree day predictors, and temperature

fluctuation predictors under SSP370 climate change conditions. As we were predicting into the

future and the range of the eastern sprue budworm is expected to shift (Gray, 2008; Régnière, St-

Amant, et al., 2012), we increased the spatial extent of the predictors to include -96.76°W to -

59.51°W and 43.42°N to 53.75649°N. All predictors had a spatial resolution of approximately

6.5km × 6.5km.

iii) Building and evaluating distribution models of defoliation

To include areas with no defoliation in the models, we marked cells up to 35km away

from the nearest point with defoliation as regions where defoliation was absent, creating a

"buffer region" around the outbreak. We also ensured grid cells used for training the models were

within Quebec provincial boundaries. We used points within this buffer as true absences of

defoliation in the model.

Exploratory modelling showed that random forest regression had the highest r-squared

value (best preliminary performance) amongst several machine learning algorithms tested; thus,

it was selected for further modelling analysis. Furthermore, to generate honest and spatially

unbiased metrics of model performance, spatial cross-validation was used reduce the effect of

spatial autocorrelation on model performance and evaluation. To do this, we used the *spatialRF*

package (Wright & Ziegler, 2017), which can generate spatial random forest models, evaluate the

contributions of predictors locally and to overall model performance, and evaluate model

performance using spatial cross-validation. We chose to evaluate the influence of temperature

fluctuation predictors on defoliation using a random forest regression model predicting

cumulative defoliation severity (i.e. summed severity scores).

Climate, degree day, forest, and temperature fluctuation predictors were highly correlated

with one another, and therefore all predictors initially showed a high degree of overfitting and

resulted in poor overall performance. Thus, we ran an initial model with all 169 predictors and

default *spatialRF* hyperparameters (Wright & Ziegler, 2017) and obtained the ranked relative

importance of each predictor. Then, we kept predictors with a high relative importance in the

initial model, and eliminated less important predictors which have a correlation coefficient over

0.95 and subsequently a variance inflation factor (VIF) higher than 7.5 as compared to higher-

ranking predictors (i.e. if two predictors are highly correlated with one another, the predictor

with a higher importance to the initial model is kept while the other is removed; Zhang et al.,

2023). These values are more conservative than those used in similar models (Dormann et al.,

2013); however, due to the large number of variables which were highly correlated with one

another, we decided more restrictive predictor selection was needed. This significantly reduced

the number of predictors in the final model, as they were reduced from 169 to 12 predictors.

Degree day predictors were not selected for the model; thus, they were absent from subsequent

analysis. Some host tree predictors were not selected during predictor pruning, so they were

added back in to ensure predictions of defoliation under climate change were limited by the

distribution of suitable food and habitat (which are not expected to change significantly under

climate change scenarios; Boisvert-Marsh et al., 2022). We then re-tuned model hyperparameters

using the *spatialRF* function "rf_tuning".

To evaluate the model and determine the importance of selected predictors, we performed

spatial cross-validation with a 4:1 training:testing data split to measure the root-mean-square

error (RMSE) of the regression model, and the permuted increase in model error when individual

predictors were removed (Wright & Ziegler, 2017). For evaluation and climate change

projections, the model predicted the cumulative defoliation severity score. To determine the

effect of temperature fluctuation predictors on model performance, we regenerated the model

without temperature fluctuation predictors and compared its performance to the model with all

predictors. Predictor pruning was re-performed and model hyperparameters were re-tuned for the

model without temperature fluctuation predictors to ensure differences in performance were not

due to initial predictor selection or poorly adapted model hyperparameters (Wright & Ziegler,

2017).

To determine where the model performed more poorly, we investigated whether model

error (residuals) varied predictably across the region. We also generated partial dependence plots

to visualize the relationships between predictors and defoliation severity. Partial dependence

plots show the relationship between the value of a predictor and the value of the predicted

defoliation propensity generated by the model. This can be used to infer the effect of the

predictor on spruce budworm defoliation.

iv)   Predicting defoliation under 2040-2071 climate change conditions

We used the random forest regression to predict regions across Quebec and Ontario (from -

96.76°W to -59.51°W and 43.42°N to 53.75649°N) which may be at risk of defoliation in a

simulated 2040-2071 climate scenario. We provided the model with the CHELSA CMIP5

climatology predictors and existing host tree distributions and had the model predict defoliation

across the region. Then, to determine the effect of temperature fluctuations on predictions of

defoliation under climate change, we re-predicted defoliation under climate change conditions

using the model trained without temperature fluctuation predictors. To measure the effect of

temperature fluctuations, we report deviations in cell-by-cell predictions between the full model

and the model with temperature fluctuation predictors removed and investigated differences in

the distribution of predicted defoliation.

All analyses were performed using R version 4.3.0 (R Core Team, 2021).

341
342   **Results**
343
344   *i)   Overall model performance*
345
Overall, the regression model predicted cumulative defoliation severity with a root-mean-

squared error (RMSE) of 2.64 ± 0.32 (SD) during evaluation (indicating predictions tended to

deviate from actual defoliation severity by a value of 2.64). The model trained without

temperature fluctuation variables had a RMSE of 3.01 ± 0.45 (SD), indicating model

performance slightly decreased when trained without temperature fluctuation predictors.

To understand where the regression model performed more poorly, we investigated the

residuals of the regression model along a gradient of defoliation severity. We tested both a

polynomial and linear fit and found that model error increased along a gradient of defoliation

severity when considering grid cells where defoliation was present (linear regression fit: $R^2 =$

0.37, $F_{1,1479} = 866.3$, $p < 0.001$; Figure 2). While the relationship between residuals and

defoliation severity was fit similarly by a polynomial and linear regression ($\Delta AIC = 0.55$), we

selected the simpler linear model. Grid cells where no defoliation was observed showed a wide

range of residuals, but generally defoliation absence was predicted accurately (y-axis histogram,

Figure 2B).

ii)   Variable contributions

Climate and temperature fluctuation predictors had the highest overall importance in the

random forest regression model, with a mean increase in error (RMSE) when permuted of 1.95 ±

0.21 (SD) and 1.83 ± 0.35 (SD), respectively (Figure 3). Forest predictors had lower overall

importance, with a mean increase in error (RMSE) when permuted of 1.48 ± 0.20 (SD). The

distribution of local (cell-by-cell) importance of the top four most important variables also vary

among predictors, with some showing higher importance localized to areas where defoliation is

present or absent. For example, August mean precipitation generally showed higher importance

in areas where defoliation was present, while January -5°C fluctuations was generally more

important in areas where defoliation was absent (Figure S4). Temperature fluctuation predictors

make up six of 15 predictors in the model and rank relatively highly among other predictors.

Five out of six temperature fluctuation predictors rank higher than all tree distribution predictors.

The relationships between cumulative defoliation severity and model predictors in the

regression model are complex and depend on the quantiles of other predictors (Figure 4).

Generally, defoliation is more severe with lower August mean precipitation and higher April

mean precipitation. Defoliation severity also increases at lower elevations and higher tree cover

(although relationships with specific tree species are variable). There is generally no consistent

relationship between defoliation severity and fluctuation predictors, as these vary across months,

temperatures, and quantiles of other predictors (Figure S5 shows relationship between values of

the four most important predictors and observed and predicted defoliation).

iii)   Climate change predictions

In general, the regression models trained with and without temperature fluctuation were

able to differentiate areas susceptible to defoliation when predicting using CMIP5 predicted

climatologies for 2041-2070 (Figure 5A,B). The model trained with temperature fluctuation

variables had an average predicted cumulative defoliation severity of 2.44 ± 0.68 (SD) and

predicted the highest defoliation severity as 4.87. The model trained without temperature

fluctuation variables predicted higher defoliation on average, with a mean predicted cumulative

defoliation severity of 3.03 ± 0.64 (SD). This model also predicted a higher maximum

cumulative defoliation severity, with a maximum predicted value of 6.70.

The regression model trained with temperature fluctuations predicted severe defoliation

at a lower frequency than the model trained without temperature fluctuations, but in more

northerly regions (Figure 5). The frequency of cells with a predicted cumulative defoliation

severity >4 was much higher in the model trained without temperature fluctuations (AUC_{with}

$_{fluctuations} = 101$; $AUC_{without\ fluctuations} = 2,421$). The average latitude of cells with a defoliation

severity >4 was 51.1 ± 1.54 (SD) in the model trained with temperature fluctuations, and $49.4 \pm$

1.66 (SD) in the model without. However, given higher defoliation severity is observed near to

the highest latitudes predicted on by the model it is possible that the latitudinal extent of the

study area is artificially truncating defoliation predictions.

Discussion

Here, we show that temperature fluctuations are an important predictor of spruce

budworm defoliation from 2006-2016 in Quebec, as including temperature fluctuations improved

the performance of the random forest regression model when predicting cumulative defoliation

severity. Temperature fluctuation predictors ranked highly amongst model predictors and had a

similar mean importance to climate predictors and higher mean importance than forest

predictors. When predicting future defoliation under climate change conditions, the inclusion of

temperature fluctuation predictors drastically reduced predicted defoliation severity across the

landscape. These results suggest that temperature fluctuations play a role in influencing the

landscape-scale defoliation of spruce budworm and highlight the importance of incorporating

physiological measures into species distribution models to improve performance.

The importance of temperature fluctuations to the defoliation regression model was

expected given the physiological importance of temperature fluctuations to spruce budworm.

Marshall and Sinclair (2015) demonstrate that the number and frequency of low temperature

stress events experienced affected spruce budworm survival in the laboratory, which provides

experimental evidence that supports our findings that temperature fluctuations may be important

on a landscape scale. Their results also indicate that the frequency of temperature fluctuations

may be more important than duration and severity of temperature exposure, which was not the

case in our model given the importance of other climatic variables. However, climate may matter

more at large spatial scales than in laboratory experiments due to climate extremes in the

northern and southern spruce budworm distribution margins that define range limits (Marshall &

Roe, 2021). Additionally, while Marshall and Sinclair investigated temperature fluctuations at

below-freezing temperatures, our model suggests that fluctuations at mild temperatures during

the fall and winter may influence defoliation severity over the landscape. How the frequency of

mild temperature exposures may affect spruce budworm and other insects during the fall and

winter has not been directly studied. However, short exposures to warm temperature events are

known to affect the fitness of spruce budworm and other insects, particularly when they occur

prior to or during overwintering in the fall and winter. Roe et al. (2024) find that spruce

budworm are highly sensitive to fall length and temperature, and that a delicate balance is struck

between survival and energetic costs during this period. Prolonged exposure to higher

temperatures prior to overwintering also led to significant delayed post-overwintering mortality

in the green-veined white (*Pieris napi*; Nielsen et al., 2022), and Zhu et al. (2019) find that

increased frequency of extreme heat events during generally mild summer conditions led to a

significant decrease in fitness of an aphid species (*Sitobian avenae*). Thus, while there is no

direct correlation between spruce budworm fitness and mild temperature fluctuations in fall and

winter, it is plausible that a relationship may exist, and we suggest further study to identify

mechanistic explanations.

Shifts in the frequency of temperature fluctuations under climate change may influence

future spruce budworm defoliation distribution and severity, but challenges still exist when

modelling future daily temperatures and how biota will respond to climate change. While the

accuracy of species distribution model predictions generally cannot be evaluated, predicted

defoliation under climate change by the model trained with temperature fluctuations was

dramatically different than the model trained without and suggests that defoliation models are

highly sensitive to these predictors. Overall temperature variability is expected to increase under

climate change (Dillon et al., 2016; G. Wang & Dillon, 2014); however, the frequency of

variation at specific temperature thresholds is strongly driven by warming fall, winter, and spring

temperatures and the non-uniformity of climate change effects across biomes and latitudes

(Contosta et al., 2024; Marshall et al., 2020). This was reflected in our results, as changes in the

frequency of temperature fluctuation predictors under climate change varied widely across the

landscape and among our predictors, as we saw a general decrease in the frequency of mild fall

temperatures, and a general increase in the frequency of mild winter temperatures (unpublished

data). This also highlights uncertainty in species distribution model predictions, as while our

methods could generate stochastic daily temperatures informed by both daily historical

temperatures and future climate trends, this may not accurately forecast trends in temperature

fluctuations (and, by extension, spruce budworm defoliation) in the future. More reliable

methods to predict future daily maximum and minimum temperatures are being developed

(Araya-Osses et al., 2020; Dillon et al., 2016; Requena et al., 2021), but there are still few

resources on a fine spatial and temporal scale useful for large-scale biogeographical forecasting.

The sensitivity of our species distribution model to temperature fluctuations and the lack of fine-

scale future daily weather forecasting reveals uncertainty when predicting the response of biota

to climate shifts. To combat this, we suggest further refinement of our methods and the

development of more fine-scale future climate modelling to clarify the response of spruce

budworm and other insects to daily temperature fluctuations under climate change.

In general, the importance of climate predictors to the random forest regression model

was consistent with findings from previous spruce budworm species distribution models (Candau

& Fleming, 2005; Gray, 2008; Zhang et al., 2023) and environmental factors known to affect

spruce budworm physiology and defoliation. Winter (Dec. and Feb.) maximum temperatures and

spring and summer (Apr. and Aug.) precipitation were the most important climate predictors to

our model. The importance of winter maximum temperatures is consistent with the spruce

budworm distribution model created by Candau and Fleming (2005), who identify this as the

most important variable for predicting defoliation frequency during a 1967-1998 outbreak in

Ontario. Given the effect of winter temperatures on insect cold stress and overwintering survival

and fitness (Marshall & Roe, 2021; Sinclair, 2015), it is logical that winter temperatures may

also drive spruce budworm fitness at the landscape scale. Precipitation was also important for

predicting defoliation; in particular, August precipitation was the most important individual

predictor of defoliation in our regression model. Studies on *C. fumiferana* show that periods of

drought can promote defoliation (Wellington et al., 1950), and that larval development is faster in

summers with dry and sunny weather (Greenbank, 1956); however, spruce budworm have

generally eclosed by August so climatic conditions in this month likely do not have a direct

effect on larval defoliation (Marshall & Roe, 2021). Peak spruce budworm moth exodus typically

occurs in August (Royama, 1984), so lower tropospheric weather conditions in this month

influence dispersal (Boulanger et al., 2017) and may partially explain our results. Our results

match the spruce budworm defoliation model created by Zhang et al. (2023), as they also find

that spring and summer precipitation are strong predictors of defoliation during an outbreak in

Newfoundland. Ultimately, our results align with previous findings on the climatic drivers of

defoliation and underscore the influence of climate on spruce budworm outbreak dynamics.

While our model attempts to capture how environmental factors influence spruce

budworm defoliation, biotic factors may explain the higher model error in areas of severe

defoliation. The dynamics of spruce budworm outbreaks are complex and are likely driven by a

combination of top-down and bottom-up factors (Pureswaran et al., 2016). Thus, higher model

error in areas with high defoliation severity may be explained by spruce budworm population

dynamics and defoliator-host tree relationships. De Grandpré et al. (2022) find that spruce

budworm herbivory significantly increases tree needle nitrogen content, which may lead to a

positive feedback loop of improved host foliage quality promoting further spruce budworm

feeding and development. Additionally, spruce budworm population dynamics may account for

the error distribution of our model. Spatial and temporal dynamics of spruce budworm outbreaks

are likely driven by density-dependent control by natural enemies and spatial synchrony of

outbreaks due to the Moran effect (Pureswaran et al., 2016). Dispersal of gravid moths can also

synchronize outbreaks across the landscape (Régnière & Nealis, 2019), and population spread,

growth rates, and outbreak risk are known to be density-dependent in the spruce budworm

(Régnière et al., 2019). Thus, while models investigating environmental factors influencing

defoliation are useful, additional drivers contributing to outbreak dynamics that are not as easily

modelled such as biotic interactions must also be considered.

Our model provides insights into the factors influencing the distribution and severity of

spruce budworm defoliation; however, these insights are correlative and should be further

supported by physiological mechanisms. Correlative models use climate and environmental

factors to predict the response of an organism to changes in those factors, and have been used

effectively to investigate the drivers of landscape-scale outbreaks (Candau & Fleming, 2005;

Gray, 2008; Zhang et al., 2023). However, they assume that trends in that response can be

extrapolated to conditions which were not explicitly studied (e.g., extrapolating the effect of

current temperature conditions on insect range to conditions under climate change), and do not

consider organism physiology or sub-processes, which can modify the response of the organism

to different conditions. Thus, robust predictions of insect distributions in changing conditions

require an understanding of physiological mechanisms driving their response in addition to

landscape-level studies of abundance and fitness using modelling techniques (Rodríguez et al.,

2019). Thus, findings of our model should be cautiously applied to novel populations and

environmental conditions, given they do not consider mechanistic explanations of spruce

budworm responses. Incorporating the mechanistic response of spruce budworm in distribution

modelling is possible and may be explored in future models using methods such as *NicheMapR*

(Kearney & Porter, 2020; Lembrechts et al., 2019) or hybrid modelling techniques (Tourinho &

Vale, 2023).

Conclusion

In this study, we investigated the hypothesis that temperature fluctuations influence the

landscape-scale distribution of spruce budworm defoliation. Results from our species distribution

model of defoliation suggest temperature fluctuations influence historic spruce budworm

defoliation in Quebec and are as important to the model as climatic variables. Additionally,

inclusion of temperature fluctuation variables drastically decreases predicted defoliation severity

across Ontario and Quebec under climate change conditions. Further study is needed to clarify

the role of biotic drivers of outbreaks on defoliation patterns, and to generate reliable and large-

scale future daily temperatures to more accurately evaluate the response of insects to climate

shifts in the future. Ultimately, elucidating the effects of temperature fluctuations and other

climatic factors on spruce budworm defoliation reveals a link between physiological responses

and landscape-scale fitness. Given the highly destructive nature of spruce budworm, this will

also contribute to our understanding of the drivers of forest pest outbreaks to enable better

prediction and management of regions susceptible to large-scale outbreaks.

Funding statement

ENB was supported by an NSERC Undergraduate Student Research Award and NSERC

Canadian Graduate Scholarship – Master's Award. DSP was supported by the Canadian Forest

Service. KEM was supported by an NSERC Discovery Grant.

Data availability statement

All data used in the study are publicly available and can be found in table S1. Code used in

analyses, full-sized figures, supplementary material, and table S1 containing links to data are

available at osf.io/anxc2/.

Acknowledgements

We would like to thank Dr. Amanda Roe, Ian DeMerchant, and Adam Hogg for their assistance

in accessing and providing spruce budworm defoliation data, as well as the *Ministère des*

Ressources Naturelles et des Forêts and the *National Forest Inventory* for data collection and

curation. We would like to additionally thank Dr. Amanda Roe for her invaluable knowledge on

spruce budworm and project guidance. We thank Dr. Amy Angert and Dr. Bill Milsom for their

comments which significantly improved the preliminary manuscript.

Conflict of Interest

The authors have no conflicts of interest to report.

554
555
Citations:
557

Alvich, J. (n.d.). *Earth System (ESM4)* (World). Retrieved October 5, 2023, from

https://www.gfdl.noaa.gov/earth-system-esm4/

Araya-Osses, D., Casanueva, A., Román-Figueroa, C., Uribe, J. M., & Paneque, M. (2020).

Climate change projections of temperature and precipitation in Chile based on statistical

downscaling. *Climate Dynamics*, *54*(9), 4309–4330. https://doi.org/10.1007/s00382-020-

05231-4

Bellemin-Noël, B., Bourassa, S., Despland, E., De Grandpré, L., & Pureswaran, D. S. (2021).

Improved performance of the eastern spruce budworm on black spruce as warming

temperatures disrupt phenological defences. *Global Change Biology*, *27*(14), 3358–3366.

https://doi.org/10.1111/GCB.15643

Boisvert-Marsh, L., Pedlar, J. H., de Blois, S., Le Squin, A., Lawrence, K., McKenney, D. W.,

Williams, C., & Aubin, I. (2022). Migration-based simulations for Canadian trees show

limited tracking of suitable climate under climate change. *Diversity and Distributions*,

28(11), 2330–2348. https://doi.org/10.1111/ddi.13630

Boulanger, Y., Fabry, F., Kilambi, A., Pureswaran, D. S., Sturtevant, B. R., & Saint-Amant, R.

(2017). The use of weather surveillance radar and high-resolution three dimensional

weather data to monitor a spruce budworm mass exodus flight. *Agricultural and Forest*

Meteorology, *234–235*, 127–135. https://doi.org/10.1016/j.agrformet.2016.12.018

Candau, J. N., & Fleming, R. A. (2005). Landscape-scale spatial distribution of spruce budworm

defoliation in relation to bioclimatic conditions. *Canadian Journal of Forest Research*,

35(9), 2218–2232. https://doi.org/10.1139/x05-078

Carrington, L. B., Armijos, M. V., Lambrechts, L., Barker, C. M., & Scott, T. W. (2013). Effects

of fluctuating daily temperatures at critical thermal extremes on *Aedes aegypti* life-

history traits. *PLOS ONE*, *8*(3). https://doi.org/10.1371/JOURNAL.PONE.0058824

Colinet, H., Renault, D., Hance, T., & Vernon, P. (2006). The impact of fluctuating thermal

regimes on the survival of a cold-exposed parasitic wasp, *Aphidius colemani*.

Physiological Entomology, *31*(3), 234–240. https://doi.org/10.1111/J.1365-

3032.2006.00511.X

Contosta, A. R., Arndt, K. A., Baulch, H. M., Casson, N. J., Harpold, A., Morelli, T. L., Sirén, A.

P. K., & Templer, P. H. (2024). Threshold Changes in Winter Temperature and

Precipitation Drive Threshold Responses Across Nine Global Climate Zones and

Associated Biomes. *Annual Review of Ecology, Evolution, and Systematics*, *55*, 271–300.

https://doi.org/10.1146/annurev-ecolsys-110421-102101

De Grandpré, L., Marchand, M., Kneeshaw, D. D., Paré, D., Boucher, D., Bourassa, S., Gervais,

D., Simard, M., Griffin, J. M., & Pureswaran, D. S. (2022). Defoliation-induced changes

in foliage quality may trigger broad-scale insect outbreaks. *Communications Biology*,

5(1), 1–10. https://doi.org/10.1038/s42003-022-03407-8

Dillon, M. E., Woods, H. A., Wang, G., Fey, S. B., Vasseur, D. A., Telemeco, R. S., Marshall, K.,

& Pincebourde, S. (2016). Life in the Frequency Domain: The Biological Impacts of

Changes in Climate Variability at Multiple Time Scales. *Integrative and Comparative*

Biology, *56*(1), 14–30. https://doi.org/10.1093/ICB/ICW024

Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G.,

Gruber, B., Lafourcade, B., Leitão, P. J., Münkemüller, T., McClean, C., Osborne, P. E.,

Reineking, B., Schröder, B., Skidmore, A. K., Zurell, D., & Lautenbach, S. (2013).

602   Collinearity: A review of methods to deal with it and a simulation study evaluating their

603   performance. *Ecography*, *36*(1), 27–46. https://doi.org/10.1111/j.1600-

604   0587.2012.07348.x

605 Findsen, A., Pedersen, T. H., Petersen, A. G., Nielsen, O. B., & Overgaard, J. (2014). Why do

606   insects enter and recover from chill coma? Low temperature and high extracellular

607   potassium compromise muscle function in *Locusta migratoria*. *Journal of Experimental*

608   *Biology*, *217*(8), 1297–1306. https://doi.org/10.1242/jeb.098442

609 Flower, A., Gavin, D. G., Heyerdahl, E. K., Parsons, R. A., & Cohn, G. M. (2014). Drought-

610   triggered western spruce budworm outbreaks in the interior Pacific Northwest: A multi-

611   century dendrochronological record. *Forest Ecology and Management*, *324*, 16–27.

612   https://doi.org/10.1016/j.foreco.2014.03.042

613 Gaston, K. J. (2003). *The Stucture and Dynamics of Geographic Ranges*. Oxford University

614   Press.

615   https://books.google.com/books/about/The_Structure_and_Dynamics_of_Geographic.ht

616   ml?id=zBqpX1ajLFMC

617 Godsoe, W., Jankowski, J., Holt, R. D., & Gravel, D. (2017). Integrating Biogeography with

618   Contemporary Niche Theory. *Trends in Ecology & Evolution*, *32*(7), 488–499.

619   https://doi.org/10.1016/j.tree.2017.03.008

620 González-Tokman, D., Córdoba-Aguilar, A., Dáttilo, W., Lira-Noriega, A., Sánchez-Guillén, R.

621   A., & Villalobos, F. (2020). Insect responses to heat: Physiological mechanisms,

622   evolution and ecological implications in a warming world. *Biological Reviews*, *95*(3),

623   802–821. https://doi.org/10.1111/BRV.12588

Gray, D. R. (2008). The relationship between climate and outbreak characteristics of the spruce budworm in eastern Canada. *Climatic Change*, *87*(3–4), 361–383. https://doi.org/10.1007/s10584-007-9317-5

Greenbank, D. O. (1956). The role of climate and dispersal in the initiation of outbreaks of the spruce budworm in New Brunswick: I. the role of climate. *Canadian Journal of Zoology*, *34*(5), 453–476. https://doi.org/10.1139/z56-048

Hijmans, R. J., Bivand, R., Forner, K., Ooms, J., & Pebesma, E. (2021). *terra: Spatial Data Analysis* [Computer software]. Comprehensive R Archive Network (CRAN). https://CRAN.R-project.org/package=terra

Holt, R. D. (2009). Bringing the Hutchinsonian niche into the 21st century: Ecological and evolutionary perspectives. *Proceedings of the National Academy of Sciences*, *106*(Supplement 2), 19659–19665. https://doi.org/10.1073/PNAS.0905137106

Hortal, J., Roura-Pascual, N., Sanders, N. J., & Rahbek, C. (2010). Understanding (insect) species distributions across spatial scales. *Ecography*, *33*(1), 51–53. https://doi.org/10.1111/J.1600-0587.2009.06428.X

Huang, L.-H., Chen, B., & Kang, L. (2007). Impact of mild temperature hardening on thermotolerance, fecundity, and Hsp gene expression in *Liriomyza huidobrensis*. *Journal of Insect Physiology*, *53*(12), 1199–1205. https://doi.org/10.1016/j.jinsphys.2007.06.011

Hutchinson, G. E. (1957). Concluding Remarks. *Cold Spring Harbor Symposia on Quantitative Biology*, *22*(0), 415–427. https://doi.org/10.1101/sqb.1957.022.01.039

Jactel, H., Petit, J., Desprez-Loustau, M.-L., Delzon, S., Piou, D., Battisti, A., & Koricheva, J. (2012). Drought effects on damage by forest insects and pathogens: A meta-analysis.

Global Change Biology, *18*(1), 267–276. https://doi.org/10.1111/j.1365-

2486.2011.02512.x

Johns, R. C., Bowden, J. J., Carleton, D. R., Cooke, B. J., Edwards, S., Emilson, E. J. S., James,

P. M. A., Kneeshaw, D., MacLean, D. A., Martel, V., Moise, E. R. D., Mott, G. D.,

Norfolk, C. J., Owens, E., Pureswaran, D. S., Quiring, D. T., Régnière, J., Richard, B., &

Stastny, M. (2019). A Conceptual Framework for the Spruce Budworm Early Intervention

Strategy: Can Outbreaks be Stopped? *Forests*, *10*(10), 910.

https://doi.org/10.3390/F10100910

Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann,

N. E., Linder, H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's

land surface areas. *Scientific Data*, *4*(1), Article 1. https://doi.org/10.1038/sdata.2017.122

Karger, D. N., Wilson, A. M., Mahony, C., Zimmermann, N. E., & Jetz, W. (2021). Global daily

1 km land surface precipitation based on cloud cover-informed downscaling. *Scientific*

Data, *8*(1), 307. https://doi.org/10.1038/s41597-021-01084-6

Li, M., MacLean, D. A., Hennigar, C. R., & Ogilvie, J. (2020). Previous year outbreak conditions

and spring climate predict spruce budworm population changes in the following year.

Forest Ecology and Management, *458*, 117737.

https://doi.org/10.1016/J.FORECO.2019.117737

Luedeling, E., Caspersen, L., & Fernandez, E. (2023). *chillR: Statistical Methods for Phenology*

Analysis in Temperate Fruit Trees (Version 0.75) [Computer software]. https://cran.r-

project.org/web/packages/chillR/index.html

MacLean, D. A., Amirault, P., Amos-Binks, L., Carleton, D., Hennigar, C., Johns, R., &

Régnière, J. (2019). Positive Results of an Early Intervention Strategy to Suppress a

Spruce Budworm Outbreak after Five Years of Trials. *Forests*, *10*(5), Article 5.

https://doi.org/10.3390/f10050448

Marshall, K. E., Gotthard, K., & Williams, C. M. (2020). Evolutionary impacts of winter climate

change on insects. *Current Opinion in Insect Science*, *41*, 54–62.

https://doi.org/10.1016/J.COIS.2020.06.003

Marshall, K. E., & Roe, A. D. (2021). Surviving in a Frozen Forest: The Physiology of Eastern

Spruce Budworm Overwintering. *Physiology*, *36*(3), 174–182.

https://doi.org/10.1152/PHYSIOL.00037.2020

Marshall, K. E., & Sinclair, B. J. (2010). Repeated stress exposure results in a survival-

reproduction trade-off in *Drosophila melanogaster*. *Proceedings. Biological Sciences*,

277(1683), 963–969. https://doi.org/10.1098/rspb.2009.1807

Marshall, K. E., & Sinclair, B. J. (2012). The impacts of repeated cold exposure on insects.

Journal of Experimental Biology, *215*(10), 1607–1613.

https://doi.org/10.1242/JEB.059956

Marshall, K. E., & Sinclair, B. J. (2015). The relative importance of number, duration and

intensity of cold stress events in determining survival and energetics of an overwintering

insect. *Functional Ecology*, *29*(3), 357–366. https://doi.org/10.1111/1365-2435.12328

Mattson, W. J., Haack, R. A., Lawrence, R. K., & Slocum, S. S. (1991). Considering the

nutritional ecology of the spruce budworm in its management. *Forest Ecology and

Management*, *39*, 183–210. https://doi.org/10.1016/0378-1127(91)90176-V

Ministère des Ressources naturelles et des Forêts. (2023). *Aires infestées par la tordeuse des

bourgeons de l'épinette au Québec en 2023* (p. 29).

Moise, E. R. D., Lavigne, M. B., & Johns, R. C. (2019). Density has more influence than drought

on spruce budworm (*Choristoneura fumiferana*) performance under outbreak conditions.

Forest Ecology and Management, *433*, 170–175.

https://doi.org/10.1016/j.foreco.2018.10.031

National Forest Inventory. (2021). https://nfi.nfis.org/en

Natural Resources Canada. (2021). *Spruce budworm*. https://www.nrcan.gc.ca/our-natural-

resources/forests/wildland-fires-insects-disturbances/top-forest-insects-and-diseases-

canada/spruce-budworm/13383

Nielsen, M. E., Lehmann, P., & Gotthard, K. (2022). Longer and warmer prewinter periods

reduce post-winter fitness in a diapausing insect. *Functional Ecology*, *36*, 1151–1162.

https://doi.org/10.1111/1365-2435.14037

Pureswaran, D. S., Johns, R., Heard, S. B., & Quiring, D. (2016). Population Ecology Paradigms

in Eastern Spruce Budworm (Lepidoptera: Tortricidae) Population Ecology: A Century of

Debate Environmental Entomology Advance Access Downloaded from. *Environmental*

Entomology, 1–10. https://doi.org/10.1093/ee/nvw103

Pureswaran, D. S., Neau, M., Marchand, M., De Grandpré, L., & Kneeshaw, D. (2019).

Phenological synchrony between eastern spruce budworm and its host trees increases

with warmer temperatures in the boreal forest. *Ecology and Evolution*, *9*(1), 576–586.

https://doi.org/10.1002/ece3.4779

R Core Team. (2021). *R: A language and environment for statistical computing* [Computer

software]. R Foundation for Statistical Computing.

Régnière, J. (2009). Predicting insect continental distributions from species physiology.

Unasylva, *60*, 37–42.

Régnière, J., Cooke, B. J., Béchard, A., Dupont, A., & Therrien, P. (2019). Dynamics and

Management of Rising Outbreak Spruce Budworm Populations. *Forests*, *10*(9), Article 9.

https://doi.org/10.3390/f10090748

Régnière, J., & Nealis, V. G. (2019). Density Dependence of Egg Recruitment and Moth

Dispersal in Spruce Budworms. *Forests*, *10*(8), Article 8.

https://doi.org/10.3390/f10080706

Régnière, J., Powell, J., Bentz, B., & Nealis, V. (2012). Effects of temperature on development,

survival and reproduction of insects: Experimental design, data analysis and modeling.

Journal of Insect Physiology, *58*(5), 634–647.

https://doi.org/10.1016/J.JINSPHYS.2012.01.010

Régnière, J., St-Amant, R., & Duval, P. (2012). Predicting insect distributions under climate

change from physiological responses: Spruce budworm as an example. *Biological*

Invasions, *14*(8), 1571–1586. https://doi.org/10.1007/S10530-010-9918-1/FIGURES/10

Requena, A. I., Nguyen, T.-H., Burn, D. H., Coulibaly, P., & Nguyen, V.-T.-V. (2021). A temporal

downscaling approach for sub-daily gridded extreme rainfall intensity estimation under

climate change. *Journal of Hydrology: Regional Studies*, *35*, 100811.

https://doi.org/10.1016/j.ejrh.2021.100811

Rodríguez, L., García, J. J., Carreño, F., & Martínez, B. (2019). Integration of physiological

knowledge into hybrid species distribution modelling to improve forecast of

distributional shifts of tropical corals. *Diversity and Distributions*, *25*(5), 715–728.

https://doi.org/10.1111/DDI.12883

Roe, A. D., Wardlaw, A. A., Butterson, S., & Marshall, K. E. (2024). Diapause survival requires a

temperature-sensitive preparatory period. *Current Research in Insect Science*, *5*, 100073.

https://doi.org/10.1016/j.cris.2024.100073

Royama, T. (1984). Population Dynamics of the Spruce Budworm *Choristoneura fumiferana*.

Ecological Monographs, *54*(4), 429–462. https://doi.org/10.2307/1942595

Sinclair, B. J. (2015). Linking energetics and overwintering in temperate insects. *Biology*

Publications, *72*. https://ir.lib.uwo.ca/biologypubhttps://ir.lib.uwo.ca/biologypub/72

Sobek-Swant, S., Crosthwaite, J. C., Lyons, D. B., & Sinclair, B. J. (2012). Could phenotypic

plasticity limit an invasive species? Incomplete reversibility of mid-winter deacclimation

in emerald ash borer. *Biological Invasions*, *14*(1), 115–125.

https://doi.org/10.1007/S10530-011-9988-8/FIGURES/3

Storey, K. B., & Storey, J. M. (1989). Freeze Tolerance and Freeze Avoidance in Ectotherms. In

L. C. H. Wang (Ed.), *Animal Adaptation to Cold* (pp. 51–82). Springer.

https://doi.org/10.1007/978-3-642-74078-7_2

Storey, K. B., & Storey, J. M. (2004). Metabolic rate depression in animals: Transcriptional and

translational controls. *Biological Reviews*, *79*(1), 207–233.

https://doi.org/10.1017/S1464793103006195

The SSP Scenarios. (n.d.). [Page]. DKRZ. Retrieved July 25, 2023, from

https://www.dkrz.de/en/communication/climate-simulations/cmip6-en/the-ssp-scenarios

Tougeron, K. (2019). Diapause research in insects: Historical review and recent work

perspectives. *Entomologia Experimentalis et Applicata*, *167*(1), 27–36.

https://doi.org/10.1111/EEA.12753

Tourinho, L., & Vale, M. M. (2023). Choosing among correlative, mechanistic, and hybrid

models of species' niche and distribution. *Integrative Zoology*, *18*(1), 93–109.

https://doi.org/10.1111/1749-4877.12618

Toxopeus, J., & Sinclair, B. J. (2018). Mechanisms underlying insect freeze tolerance. *Biological*

Reviews, *93*(4), 1891–1914. https://doi.org/10.1111/brv.12425

Wang, G., & Dillon, M. E. (2014). Recent geographic convergence in diurnal and annual

temperature cycling flattens global thermal profiles. *Nature Climate Change*, *4*(11),

Article 11. https://doi.org/10.1038/nclimate2378

Wang, T., Hamann, A., Spittlehouse, D., & Carroll, C. (2016). Locally Downscaled and Spatially

Customizable Climate Data for Historical and Future Periods for North America. *PLOS*

ONE, *11*(6), e0156720. https://doi.org/10.1371/journal.pone.0156720

Wellington, W. G., Fettes, J. J., Belyea, R. M., & Turner, K. B. (1950). Physical and biological

indicators of the development of outbreaks of the spruce budworm, *Choristoneura*

fumiferana (clem.) (Lepidoptera: Tortricidae). *Canadian Journal of Research*, *28d*(6),

308–331. https://doi.org/10.1139/cjr50d-021

Wright, M. N., & Ziegler, A. (2017). ranger: A Fast Implementation of Random Forests for High

Dimensional Data in C++ and R. *JSS Journal of Statistical Software*, *77*.

https://doi.org/10.18637/jss.v077.i01

Xu, B., Hicke, J. A., & Abatzoglou, J. T. (2019). Drought and Moisture Availability and Recent

Western Spruce Budworm Outbreaks in the Western United States. *Forests*, *10*(4), Article

4. https://doi.org/10.3390/f10040354

Zhang, B., Leroux, S. J., Bowden, J. J., Hargan, K. E., Hurford, A., & Moise, E. R. D. (2023).

Species distribution model identifies influence of climatic constraints on severe

defoliation at the leading edge of a native insect outbreak. *Forest Ecology and*

Management, *544*, 121166. https://doi.org/10.1016/j.foreco.2023.121166

Zhu, L., Wang, L., & Ma, C.-S. (2019). Sporadic short temperature events cannot be neglected in

predicting impacts of climate change on small insects. *Journal of Insect Physiology*, *112*,

48–56. https://doi.org/10.1016/j.jinsphys.2018.12.003

785
786   **Figure Captions**
787
Figure 1. Distribution of spruce budworm cumulative defoliation severity across Quebec from
2006-2016, collected from aerial surveys conducted by *Ministère des Ressources Naturelles et*
des Forêts in Quebec. Cumulative defoliation severity calculated by summing annual defoliation
severity levels in each year from 2006-2016. An additional 35km buffer with no defoliation was
added to the mapped defoliation to serve as additional absences. n = 4,146 grid cells (total area =
233,269km^2). Resolution: 6.5km × 6.5km.
794
Figure 2. Regression model error tends to increase in areas with higher observed defoliation
severity. **A)** Distribution of residuals of random forest regression model. Red cells indicate
model underestimated defoliation severity, blue cells indicate model overestimated defoliation
severity. **B)** Relationship between model residuals and actual values of cumulative defoliation
severity. Histograms in margins show density of data across x- and y-axis values. Linear
regression line only fit to cumulative defoliation severity (x-axis) values higher than zero (where
defoliation was present). $n_{present}$ = 1,481 grid cells.
802
803   Figure 3. Climate and temperature fluctuation predictors are generally more important than forest
804   variables in the random forest regression model. Bars represent increase in overall model error
805   when predictor was removed from training model.
806
Figure 4. Relationship between random forest regression model predictors and predicted
cumulative defoliation severity across fixed quantile values of other model predictors. X-axes
indicates value of labelled predictor across outbreak range: temperatures = °C, fluctuations =
frequency per month, tree abundances = % cover, elevation = m, and precipitation = kg m^{-2} s^{-1}
(mean rate of precipitation scaled up by a factor of 100,000 to ensure compatibility with
predictor selection).
813
Figure 5. Predicted defoliation across Quebec, Ontario, and Labrador by the random forest
regression model using CMIP5 predicted climatologies for 2041-2070. Areas covered by lakes
and rivers removed from prediction area. **A)** Distribution of predicted cumulative defoliation
severity of model trained using temperature fluctuations. **B)** Distribution of predicted cumulative
defoliation severity of model trained without temperature fluctuations. **C)** ΔPredicted defoliation
severity between models trained with and without temperature fluctuations. **D)** The model

trained using temperature fluctuations predicted severe (>4 cumulative severity score) defoliation

less and with a more northerly distribution

Value of predictor