Epigenetic control of tetrapyrrole biosynthesis by $^m4$C DNA methylation in a cyanobacterium

Nils Schmidt[1], Nils Stappert[2], Kaori Nimura-Matsune[3], Satoru Watanabe[3], Roman Sobotka[4,5], Martin Hagemann[1]*, Wolfgang R. Hess[2]*

[1]Institute of Biosciences, Department of Plant Physiology, University of Rostock, D-18059 Rostock, Germany

[2]University of Freiburg, Faculty of Biology, Genetics and Experimental Bioinformatics, Schänzlestr. 1, D-79104 Freiburg, Germany

[3]Department of Bioscience, Tokyo University of Agriculture, Tokyo, Japan.

[4]Institute of Microbiology of the Czech Academy of Sciences, Centre Algatech, Třeboň, 379 01, Czech Republic

[5]Faculty of Science, University of South Bohemia, České Budějovice, 370 05, Czech Republic

*Corresponding authors: Martin Hagemann, University of Rostock, Institute of Biosciences, Department of Plant Physiology, A.-Einstein-Str. 3, D-18059 Rostock, Germany; Tel: +49(0)3814986110; Email: martin.hagemann@uni-rostock.de

Wolfgang R. Hess, University of Freiburg, Faculty of Biology, Genetics and Experimental Bioinformatics, Schänzlestr. 1, D-79104 Freiburg, Germany; Tel: +49(0)7612032796; Email: wolfgang.hess@biologie.uni-freiburg.de

Running title: DNA modification and HemJ expression

Abstract

Epigenetic DNA modifications are pivotal in eukaryotic gene expression, but their regulatory significance in bacteria is less understood. In *Synechocystis* 6803, the DNA methyltransferase M.Ssp6803II modifies the first cytosine in the GGCC motif, forming N4-methylcytosine (GG^{m4}CC). Deleting the *sll0729* gene (Δ*sll0729*) caused a bluish phenotype due to reduced chlorophyll levels, which was reversed by suppressor mutations. Re-sequencing of seven suppressor clones revealed a common GGCC to GGTC mutation in the *slr1790* promoter's discriminator sequence, encoding protoporphyrinogen IX oxidase, HemJ, crucial for tetrapyrrole biosynthesis. Transcriptomic and qPCR analyses indicated aberrant *slr1790* expression in Δ*sll0729* mutants. This aberration led to the accumulation of coproporphyrin III and protoporphyrin IX, indicative of impaired HemJ activity. To confirm the importance of DNA methylation in *hemJ* expression, native and mutated *hemJ* promoter variants were introduced into the wild type, followed by *sll0729* deletion. The *sll0729* deletion segregated in strains with the GGTC motif in the *hemJ* promoter, resulting in wild-type-like pigmentation, whereas freshly prepared Δ*sll0729* mutants with the native *hemJ* promoter exhibited the bluish phenotype. These findings demonstrate that *hemJ* is tightly regulated in *Synechocystis* and that N4-methylcytosine is essential for proper *hemJ* expression.

Key words: Cyanobacteria / epigenetic modifications / HemJ / DNA methyltransferase / tetrapyrrole biosynthesis

1. Introduction

DNA base methylation occurs in all kingdoms of life. It affects DNA replication, cell cycle control, DNA mismatch repair, DNA-protein interactions, phenotypic heterogeneity among populations, gene expression, and recognition of intruder DNA.[1] The most common types of methylated bases comprise the endocyclic 5-methylcytosine (^{m5}C), the exocyclic N-6-methyladenine (^{m6}A), and N4-methylcytosine (^{m4}C). Newly methylated bases are established by DNA methyltransferases (MTases) that recognize specific DNA motifs.[2] These modifications do not impair with Watson-Crick base pairing but protrude to the major groove of DNA exposing the methyl group to DNA binding proteins or the transcriptional machinery.[2] Bacterial MTases are commonly associated with restriction endonucleases forming restriction-modification (R-M) systems.[3,4]

However, MTases not associated with a cognate restriction enzyme, named 'orphan' MTases, are also present. Global analyses of the DNA methylome of 230 different bacteria and archaea revealed such orphan MTases in 48% of the examined organisms.[5] The most prominent examples of 'orphan' MTases in prokaryotes are the ^{m5}C MTase Dcm and the ^{m6}A MTases Dam and Ccrm.[6–8] In eukaryotes and prokaryotes, ^{m5}C[9–11] as well as ^{m6}A modifications can alter gene expression if associated with promoter sequences.[12,13] CpG dinucleotides carrying a ^{m5}C are most frequently observed in eukaryotic promoter regions,[10] whereas e.g. the *pap*-operon of *E. coli* is regulated by ON and OFF states caused by the change in methylation of adenines in the promoter region.[14] The effects of ^{m4}C methylation in prokaryotes, however, has remained largely unclear. Changes in transcription and a change in pathogenicity were shown for *Leptospira interrogans*.[15] But, how the occurrence of ^{m4}C in promoter sequences can affect gene expression has still to be elucidated.

Cyanobacteria are the only prokaryotes conducting oxygenic photosynthesis making them ecologically relevant. Their unique physiology and redox chemistry have generated interest for producing various chemicals based on photosynthesis, using CO_2 and solar energy.[16–18] In the model cyanobacterium *Synechocystis* sp. PCC 6803 (*Synechocystis* 6803) five MTases have been identified. We previously showed that: i) the CGATCG sequence motifs in *Synechocystis* 6803 are double-methylated (^{5m}CG^{6m}ATCG) by the

enzymes M.Ssp6803I and M.Ssp6803III; ii) these motifs are abundant within the repeat-
spacer arrays of a subtype I-D CRISPR-Cas system; iii) conjugation efficiency decreased
by 50% in cells lacking ^{5m}C methylation due to deletion of gene *slr0214* encoding
M.Ssp6803I.[19] These results pointed at an important role of DNA methylation in
Synechocystis 6803.

M.Ssp6803II, the DNA methyltransferase encoded by gene *sll0729* is responsible for the
addition of an exocyclic methyl group to the internal cytosine in the motif GGCC, which
occurs 38,512 times in the *Synechocystis* 6803 genome.[20] Mutant strains with a deletion
of gene *sll0729* (Δ*sll0729*) developed a pronounced phenotype, including slower growth,
smaller cell diameter, lowered DNA contents, and a bluish pigmentation caused by a
lowered chlorophyll/phycocyanin ratio,[21] whereas complementation by *sll0729* restored
the phenotype of the wild type (WT). We demonstrated that altered transcript levels of two
genes, *sll0470* and *sll1526*, in Δ*sll0729* were caused by the lacking ^{m4}C methylation in
GGCC elements located in the respective -35 promoter elements.[21] The results suggested
the finding that ^{m4}C plays some essential role in *Synechocystis* 6803. After longer
cultivation times, suppressor mutants with a pigmentation resembling WT appeared
frequently. These suppressor mutants were still ^{m4}C methylation negative, but the nature
of these mutations remained enigmatic.[21]

In this work we have addressed the molecular basis of the suppressor mutants. By
resequencing analysis we found that all lines shared a single point mutation in one GGCC
site recognized by M.Ssp6803II for ^{m4}C methylation, which was changed in all sequenced
suppressors to GGTC. However, this point mutation is not located in a coding region, but
upstream of gene *slr1790* encoding protoporphyrinogen IX oxidase (HemJ), a key enzyme
of tetrapyrrole biosynthesis.[22,23] Transcriptomic analyses revealed a lowered mRNA level
for *slr1790* in Δ*sll0729*, while levels similar to WT were reached in the suppressor mutants.
In a series of reconstitution experiments we demonstrate that the non-balanced *hemJ*
expression due to the non-methylated promoter sequence led to a dramatic over-
accumulation of coproporphyrin III (CoPP) and protoporphyrin IX (PPIX), which is typical
for the inhibited HemJ enzymatic activity (see below). In contrast, the accumulation of
these metabolites became WT-like in the presence of the GGTC promoter variant. Overlay
of the resequencing data with high-resolution data from the genome-wide mapping of

transcriptional start sites suggested that the identified C-->T mutation is located within the
discriminator area of the *slr1790* promoter. Thus, we show that N4 methylation of the first
cytosine in the motif GGCC leading to GG^{m4}CC is essential for the quantitative correct
transcription initiation of the RNA polymerase and how a single nucleotide exchange of
this cytosine in the promoter of *slr1790* can alter gene expression and affect the
physiology of a photosynthetic bacterium. Our results underline the relevance of
epigenetic effects linked to ^{m4}C methylation.

2. Material and Methods

2.1. Bacterial strains and cultivation

Synechocystis sp. PCC 6803 substrain PCC-M[24] was used in all experiments and served
as wild type (WT). Axenic strains were maintained on agar plates with BG11 mineral
medium at 30°C under constant illumination (30 µmol photons m^{-2} s^{-1}; to slow down
formation of suppressor clones). Transformants were selected and segregated on media
containing 50 µg ml^{-1} kanamycin (Km; Sigma), 50 µg ml^{-1} spectinomycin (Spec; Sigma) or
both. For physiological characterization axenic cultures of the different strains were grown
photoautotrophically in BG11 medium, either under slight shaking in Erlenmeyer flasks at
30 µmol photons m^{-2} s^{-1}, or aerated with atmospheric air in a photobioreactor (Multi-
Cultivator MC 1000-OD-WW warm light, Photon Systems Instruments) at 30°C under
continuous illumination of 40 µmol photons m^{-2} s^{-1}. The appearance of suppressor clones
was detected by measuring the absorption spectrum in between 400 and 750 nm (CARY
50 Bio, Varian). Contamination by heterotrophic bacteria was evaluated by spreading 0.2
ml culture aliquots on LB plates. The *E. coli* strain DH5α was used for routine DNA
manipulations. *E. coli* was cultured in LB medium at 37°C. *Synechocystis* 6803 growth
was recorded by measurements of optical density at 720 nm (OD$_{720}$) either manually
(shaking flask) or automatically (Multicultivator).

2.2. Resequencing analysis of suppressor lines

Five-hundred ng of genomic DNA was fragmented to an average length of 500 bp using a Covaris S2 sonication system (Covaris, Inc., MA, USA). Sequencing libraries were prepared using the NEBNext Ultra DNA Library Prep Kit for Illumina (New England Biolabs). Paired-end sequencing was carried out for 150 cycles using the Nextseq500 system (Illumina Inc., CA, USA) according to the manufacturer's specifications. The sequencing reads were trimmed using the CLC Genomics Workbench ver. 9.5.4 (Qiagen) with the following parameters: Phred quality score > 30; removing the 5′ terminal 10 nt; and removing truncated reads <20 nt. Trimmed reads were mapped to the reference genome sequence and plasmids of *Synechocystis* 6803 (accession numbers: CP003265-CP003272) using CLC Genomics Workbench ver. 9.5.4 (Qiagen) with the following parameters: length fraction: 0.8, and similarity fraction: 0.9. To identify SNVs and indels, we used the filter settings as follows: minimum read depth for SNV/indel calling, 20; minimum read depth for the SNV calling, 10. Sequence reads were deposited to the DRA/SRA database with the accession numbers DRR585585–DRR585594, BioProject PRJDB18568.

2.3. Genetic engineering

The isolation of total DNA from *Synechocystis* 6803 was performed as previously described.[20] The GG^{m4}CC methylation was verified by restriction analyses using chromosomal DNA from *Synechocystis* 6803, the restriction endonucleases were used in a 10-fold excess and were incubated for at least 16 h at 37°C to ensure complete digestion. Synthetic primers were deduced from the genome sequence of *Synechocystis* 6803[25] for the specific amplification of the promoter sequence of the *slr1790* gene, coding for the protoporphyrinogen IX oxidase HemJ. Mutagenesis PCR was used to mutate the GG^{m4}CC methylation site of the promoter sequence to GGTC (see **Supplementary Table S4** for primer sequences). For this purpose, a DNA fragment containing the promoter region of *slr1790* was amplified by PCR and cloned into pJET1.2 (Thermo Scientific). The *aadA* gene, conferring Spec resistance from pUC4S (Pharmacia), was introduced into selected restriction sites. Verified constructs were transferred into *Synechocystis* 6803.[26]

For generation of M.Ssp6803II-deficient strains, the *sll0729* gene was deleted as
described.[20] A list of all *Synechocystis* 6803 strains used in the study is given in
Supplementary Table S2.

To check the *slr1790* promoter activity, different promoter variants with single nucleotide
substitutions of the internal cytosine in GGCC were tested. The nucleotide sequence of
the used *slr1790* promoter ranging from -143 to +38 (TSS at +1) was fused to *luxAB*
reporter genes and integrated into the genome. Promoter variants carrying the WT motif
GGCC, GGTC, GGAC and GGGC were generated. For the integration into the genome
the pILA vector[27] was used.

2.4. RNA preparation and qRT-PCR

Cultures were grown under standard light conditions (50 µmol photons m^{-2} s^{-1}) and 30°C
and harvested via centrifugation (5,000 g, 20°C, 10 min) during exponential phase (OD
0.8). The pellet was resuspended in 1 ml PGTX,[28] immediately frozen in liquid nitrogen
and processed further as previously described.[29] Relative amounts of RNA were
quantified by qRT-PCR. Therefore, cDNA was prepared according to the manufacturer's
instructions using 600 ng of each RNA sample with the QuantiTect Reverse Transcription
Kit (QIAGEN). qRT-PCR was performed using the FastGene® qFYR Real-Time PCR
System (NIPPON Genetics). PCR cycles for each run are described in **Supplementary**
Dataset S1 and **S2**. qPCRBIO SyGreen Blue Mix Separate-ROX was used and PCR
reactions were performed according to the protocol (PCR biosystems). Primers
(**Supplementary Table S4**) were synthesized by IDT. All reactions were measured as
technical triplicates and *rnpA* was used as an endogenous control. The data was analyzed
using the qFYR Analyzer Studio Software (Nippon Genetics) and relative quantification of
RNA was calculated using the comparative Ct Method (ΔΔCt). The WT sample was used
as the calibrator. The average of each technical triplicate measurement was displayed as
a bar chart.

2.5. Pigment quantifications

The strains were precultured at 30 µmol photons m^{-2} s^{-1} constant light and 30°C. After seven days the BG11 medium was refreshed and the cultures were grown for another two days (OD_{720} 0.15 – 0.3). All following steps were performed at dark conditions. 2 ml of the culture was collected by centrifugation (4,000 g, 20°C, 5 min). Supernatant was completely removed and the pelleted cells were resuspended in 200 µl A. dest. (HPLC grade, Merck) followed by centrifugation (4,000 g, 20°C, 2 min). The supernatant was discarded and the pellet was resuspended in 100 µl 75% methanol (HPLC grade, Merck) and incubated for 15 min at 20°C. The sample was then centrifuged (4,000 g, 20°C, 2 min) and the supernatant collected. Remaining pellet was resuspended in 80% methanol, incubated for 15 min at 20°C and centrifuged again (4,000 g, 20°C, 2 min.) Supernatant was collected, combined with the supernatant from the first extraction and the resulting solution centrifuged (4,000 g, 20°C, 5 min). The extracted pigments were finally transferred into micro inserts for high performance liquid chromatography (HPLC) analysis, performed as described.[30]

2.6. Microscopy and determination of cell size

Strains were cultivated at 30 µmol photons m^{-2} s^{-1} constant light and 30°C for 10 days. Ten µl of these cultures were transferred on microscope slide, fixed by a cover slip and directly used for microscopy (Olympus BX51). The cell sizes were determined in ImageJ.

3. Results

3.1. The suppressor mutation in ∆*sll0729* maps to a single nucleotide in the *slr1790* promoter

The absence of M.Ssp6803II resulted in the strongest phenotypic differences among the mutants lacking specific DNA methyltransferase activities, including bluish pigmentation due to decreased chlorophyll levels.[20] However, the phenotype was not stable, as a relatively high number of suppressor mutants with WT-like pigmentation appeared after prolonged cultivation of mutant ∆*sll0729*.[21] To identify the mutation(s) leading to reversion

of the phenotype, seven independently isolated suppressor clones were sequenced and
compared to the parental Δ*sll0729* strain. The sequence data revealed several single
nucleotide variations (SNV) and short deletions in the different suppressor mutants
(**Supplementary Table S1**). However, one SNV was found in the DNA of all suppressor
clones. This SNV affected a GGCC motif for M.Ssp6803II-specific DNA methylation
changing it to GGTC within the promoter of *slr1790* (**Fig. 1**).

Figure 1. One SNV upstream of *hemJ* is conserved in all suppressor clones. Upper
panel: The gene *slr1790* encoding HemJ is transcribed from a single promoter with a
transcription start site (arrow, +1) previously mapped to position 256,609 (NCBI reference
sequence accession NC_000911.1) of the forward strand and associated with a 5' UTR
of 89 nt. Shown is the accumulation of sequencing reads in the genome-wide mapping of
transcription start sites by differential RNA-seq.[31] The sharp decline in the sequence
coverage results from the obtained average sequencing read length of 100 nt.[31] Lower
panel: Resequencing of the genomes of seven suppressor clones revealed one
conserved single nucleotide variation (SNV) at position 256,605 of the forward strand
(position -4 with regard to the transcription start) changing C to T. This transition changes
the GGCC methylation motif to GGTC (red T) and is located between the promoter -10
element and the transcription start site, that is, within the promoter discriminator region
(highlighted by the ochre background).

The gene *slr1790* encodes the protoporphyrinogen IX oxidase HemJ.[22,23] HemJ is

essential for the enzymatic protoporphyrin IX (PPIX) synthesis, which is the last common

precursor for the biosynthesis of heme and chlorophyll. Heme is not just an essential

cofactor of cytochromes and other electron carriers and enzymes, but also a precursor for

the synthesis of bilins — the chromophores of phycocyanin and allophycocyanin light

harvesting proteins, and of photoreceptors such as cyanobacteriochromes and

phytochromes.[32] Therefore, a possible defect in the expression of a gene necessary for

tetrapyrrole biosynthesis could be related to the observed pigmentation phenotype of the

Δ*sll0729* mutant.

The conserved SNV in the suppressor mutants at position 256,605 affects specifically the

GGCC motif for M.Ssp6803II-specific DNA methylation in the core promoter of *slr1790*.

The motif is situated immediately downstream of the -10 element and just upstream of the

transcriptional start site (position -4; **Fig. 1**) of this gene. This region is crucial for correct

transcription initiation, because DNA strand separation by RNA polymerase in the

formation of the open complex is controlled by an arranged separation of specific

nucleotides starting from -11 and passing the discriminator sequence.[33]

Overall, there are 38,512 instances of the GGCC motif on one chromosome in the multi-

copy *Synechocystis* 6803 genome.[20] Hence, it is possible that not all of them are fully

methylated. Bisulfite sequencing analysis permits the direct and highly sensitive detection

of ^{m5}C but it can also be used to map ^{m4}C, although ^{m4}C is partially resistant to bisulfite-

mediated deamination. When the assay was used for a global methylation analysis, we

found that about 90% of GGCC methylation sites were methylated in the *Synechocystis*

6803 genome. Among them, bisulfite sequencing showed a complete methylation of the

GGCC at position 256,605 in the DNA isolated from WT cells (**Supplementary Fig. S1**),

which will be absent in mutant Δ*sll0729* with abolished M.Ssp6803II and in suppressor

clones bearing the mutated *slr1790* promoter with GGTC.

3.2. Lack of GG^{m4}CC methylation in *Synechocystis* 6803 impacts *hemJ* expression

We have previously reported that the lacking methylation of GGCC motifs by M.Ssp6803II impacted the expression of genes including *sll0470* and *sll1526*, with $\log_2$FCs of +1.3 and -1.05, respectively [21]. Therefore, it appeared possible that the lack of GG^{m4}CC methylation had also consequences for the expression of *hemJ*. We reanalyzed our existing transcriptome data sets comparing the gene expression in mutant Δ*sll0729* with WT.[21] Indeed, a similarly decreased expression of *hemJ* in Δ*sll0729* compared to WT was observed under different light conditions in an experiment from 2013 (**Supplementary Fig. S2A**). In the transcriptome data from 2014, the expression of *hemJ* was diminished in mutant Δ*sll0729* relative to WT ($\log_2$FC -0.855, below our significance threshold of -1), while in the *sll0729* complementation strain it returned to WT-levels (**Supplementary Fig. S2B**). Finally, in a third dataset from 2017, clearly diminished *hemJ* transcript levels were observed for two freshly generated Δ*sll0729* independent deletion mutants (**Supplementary Fig. S2C**). Thus, these transcriptomic data showed that the expression of *hemJ* was lowered in Δ*sll0729*, in which the methylase M.Ssp6803II specific for cytosine N4-methylation leading to GG^{m4}CC was missing. The quantitative difference was in all cases less than twofold compared to WT, but observed reproducibly in three independent experiments performed over several years.

To verify this observation, the expression of *hemJ* was quantified by qRT-PCR in cells of freshly generated Δ*sll0729* mutants and three independently recovered suppressor clones. Again, the absence of M.Ssp6803II resulted in lowered *hemJ* expression in mutant cells, while the expression returned to levels slightly higher than WT in the suppressor clone (**Fig. 2A**). Furthermore, the impact of different point mutations in the discriminator region of the promoter was tested. For this experiment, we fused the reporter genes *luxAB* to the native *hemJ* promoter with intact GGCC motif or variants with a GGAC, GGTC, or GGGC motif (listed in **Supplementary Table S2**). The exchange of GGCC to GGTC (as found in the suppressor clones) resulted in slightly higher transcript levels than the native GGCC motif in the WT background, whereas the GGAC and GGGC variants showed the highest promoter activities (**Fig. 2B**).

Figure 2. Abundance of transcripts driven by *hemJ* promoter variants in different *Synechocystis* 6803 strains. (A) Expression of *hemJ* in wild type (WT), Δ*sll0729*, or one selected suppressor mutant clone (Suppr. Δ*sll0729*) quantified by qRT-PCR. The bars show averages of technical triplicates of biological triplicates, individual data points are given as well. WT was used as the calibrator. (**B)** qRT-PCR quantification of *luxAB* mRNA levels that were under control of *hemJ* promoter variants, in which the first C in the GGCC motif was unchanged or exchanged to A, T or G. The bars indicate averages of technical triplicates of biological triplicates, individual data points are given as dots. The WT motif GGCC was used as the calibrator for Relative Quantification (RQ).

Significance was calculated with an unpaired Two-Samples Wilcoxon Test using RStudio; *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$) between the strains at corresponding time points (details of analysis and statistical test in **Dataset EV1 and EV2**).

From these experiments we conclude that single nucleotide substitutions in the discriminator region indeed affected the levels of mRNAs produced by the *hemJ* promoter, regardless of whether the native or with the *luxAB* mRNA another transcript was produced. The GGTC motif yielded the quantitatively most similar mRNA level compared to the (methylated) GGCC in the WT background.

3.3. Reconstitution of *Synechocystis* 6803 strains to verify the importance of GGCC methylation for *hemJ* expression

The above experiments clearly showed that the identity of the nucleotide at the -4 position in the *hemJ* promoter determined the level of the transcribed mRNA. The measured quantitative differences were relatively minor, but the high frequency of suppressor mutants at this position for the Δ*sll0729* phenotype suggested that these differences were physiologically meaningful. Therefore, we tested whether the GGCC motif and proper expression of *hemJ* were directly related to the phenotypic stability of Δ*sll0729.* Several strains were generated to address this possibility (**Supplementary Table S2**). Attempts to completely segregate the *slr1790* deletion were unsuccessful, consistent with an earlier report.[22] However, we were able to generate two strains, one with the native *slr1790* promoter (P*slr1790*) and one with the mutated (GGCC substituted by GGTC) *slr1790* promoter (MP*slr1790*) by gene replacement (both strains harbored upstream a spectinomycin resistance gene) (**Supplementary Table S2**). Genotyping and sequence analysis revealed that both strains were completely segregated and stably maintained the respective promoter variants (**Fig. 3A, B**).

Subsequently, these two strains and the corresponding WT served as recipients for a new round of *sll0729* deletions, yielding strains Δ*sll0729,* P*slr1790*/Δ*sll0729,* and MP*slr1790*/Δ*sll0729* (**Supplementary Table S2**). The number of kanamycin-resistant clones obtained differed between strains containing either the native or mutated *hemJ* promoter. In the latter case, more clones appeared and these were almost all completely segregated, whereas transformation of the *sll0729* deletion cassette into the WT or the strain harboring the native promoter variant resulted in lower ratios of clones with completely segregated *sll0729* mutation (**Supplementary Fig. S3**).

Figure 3. Genotyping and phenotyping of *Synechocystis* 6803 strains to verify the importance of GGCC methylation for *hemJ* expression. (**A**) Genotyping via PCR. Upper panel, the gene encoding M.Ssp6803II is not present in strains with Δ*sll0729* background as indicated by the lack of the 827 bp amplicon. Lower panel, presence of the wild-type (WT) or the mutated *hemJ* promoter in the generated strains as indicated by the 639 bp amplicon in WT and Δ*sll0729*, or the 2687 bp fragment in the manipulated strains. The latter fragment is larger due to the *aadA* antibiotic resistance cassette inserted upstream of the promoter. Three independent clones were analyzed per strain. For details of the respective strains, see **Supplementary Table S2**. (**B**) Sequence analysis to verify the intact GGCC motif in the native promoter and its change to GGTC in the clones with mutated promoter sequence. (**C**) Absorption spectra of different strains grown under standard conditions. The spectra were normalized to the absorption at 750 nm.

This finding is in line with our expectations, because the GGTC promoter variant found in

the suppressor clones makes the deletion of M.Ssp6803II less harmful for *Synechocystis*

6803. Interestingly, the original bluish phenotype and decreased growth of mutant

Δ*sll0729* were observed in freshly segregated clones from transformations of the WT and

the strain with the native *hemJ* promoter, while the strain with the mutated *hemJ* promoter

and segregated *sll0729* showed WT-like pigmentation, as observed for newly selected

suppressor clones (**Fig. 3C, Supplementary Fig. S4**).

Subsequently, selected clones were tested in liquid and solid media under different growth

conditions. Many features reported by Gärtner et al.[21] characterizing the initial Δ*sll0729*

mutation were reproduced, such as the smaller cell size, changed pigmentation, and

growth (**Supplementary Fig. S4, Supplementary Table S3**). WT cells displayed an

average area of 4.80 μm^2, whereas all cells of strains with deleted M.Ssp6803II were

smaller (Δ*sll0729*: 3.91 μm^2; P*slr1790*/Δ*sll0729*: 3.52 μm^2; MP*slr1790*/Δ*sll0729*: 3.02

μm^2). Control strains had slightly larger cell sizes (P*slr1790*: 6.00 μm^2; MP*slr1790*: 5.51

μm^2). One of the most obvious changes was related to pigmentation. As previously

reported,[21] fresh mutants lacking M.Ssp6803II had reduced chlorophyll levels if *hemJ* was

driven by its native promoter. In contrast, the GGCC-->GGTC mutation of the *hemJ*

promoter prior to deleting *sll0729* yielded WT-like pigmentation, which was also observed

in suppressor strains. Moreover, segregated Δ*sll0729* mutants in strains with the native

hemJ promoter were less frequent, less viable, and showed slower growth than WT or

Δ*sll0729* mutants in the strain with the GGTC-changed *hemJ* promoter (**Supplementary**

Fig. S5).

Most importantly, all clones with a completely deleted *sll0729* gene were not further able

to methylate GGCC sites, regardless of the strain from which the DNA was isolated. In

contrast, DNA from all strains with an intact *sll0729* gene was resistant to restriction

enzymes that cut only DNA with non-methylated GGCC motifs (**Supplementary Fig. S6**).

Finally, in the suspension of segregated fresh Δ*sll0729* and P*slr1790*/Δ*sll0729* mutant

clones with bluish phenotype, clones with WT-like pigmentation spontaneously appeared

at a high frequency after one to four weeks of cultivation, as observed previously. DNA

was isolated from two of these putative suppressor clones and the *hemJ* promoter region
was PCR-amplified and sequenced. As observed in the genomic analysis of suppressor
lines, the GGCC motif upstream of the *hemJ* transcription start site was again changed at
position -4 to GGTC.

3.4. Effects of epigenetic manipulation on pigmentation

The gene *slr1790* encodes protoporphyrinogen IX oxidase PPO (HemJ),[22,23] the enzyme
converting the colorless protoporphyrinogen IX (PPIX) into red PPIX (**Fig. 4**). This is the
last common step in the biosynthesis of heme and chlorophyll cofactors and of all further
tetrapyrrole derivatives [32]. Therefore, the relatively small differences in *hemJ* transcription
caused by the nature of the nucleotide at the -4 position of the promoter might directly
affect the formation of pigment biosynthesis intermediates.

Figure 4. HemJ catalyzes the enzymatic conversion of protoporphyrinogen IX to PPIX. Scheme of last common steps of the chlorophyll and heme biosynthesis. HemJ is one of three different paralogs of the protoporphyrinogen IX oxidase (PPO) known in nature and is present in most cyanobacteria such as *Synechocystis* 6803. This enzyme oxidizes protoporphyrinogen IX into PPIX, the universal precursor of chlorophylls, hemes and bilins [22,23]. The enzyme names and accession numbers of the corresponding genes in *Synechocystis* 6803 are given.

To obtain a better picture of chlorophyll biosynthesis, intermediates in porphyrin
biosynthesis leading to chlorophyll were quantified in different strains grown under
identical conditions (**Table 1**). The most remarkable differences were found in the
amounts of CoPP and PPIX. CoPP is the non-enzymatically oxidized form of
coproporphyrinogen III, which is the substrate of coproporphyrinogen oxidase, a

preceding enzymatic step of HemJ. These two enzymes are enzymatically coupled, and
the impaired activity of HemJ also affects the decarboxylation of coproporphyrinogen III.[23]
Although PPIX is the product of the HemJ catalytic activity, it can also accumulate in the
cell by spontaneous oxidation of protoporphyrinogen IX after blocking HemJ activity (**Fig.**
4,[23]). Unfortunately, the cellular content of protoporphyrinogen IX cannot be measured
directly by HPLC with fluorescence detection, as this compound is colorless, and its non-
enzymatic oxidation to PPIX is very fast.

Extremely high amounts of PPIX, together with high amounts of CoPP were specifically
observed in strains showing the bluish phenotype (**Supplementary Fig. S4**), that is,
strains with segregated deletion of *sll0729* and native *hemJ* promoter, whereas all other
strains contained low amounts of PPIX and CoPP, similar to WT (**Table 1,**
Supplementary Tables S5 and **S6**).

Table 1. Quantification of chlorophyll synthesis intermediates in different strains of
Synechocystis **6803 with mutated** *sll0729* **gene and/or mutated** *slr1790* **promoter.**
The first committed intermediate of chlorophyll pathway is Mg-protoporphyrin IX (MgP),
which is consequently methylated (MgPME). The MgPME is converted into divinyl
protochlorophyllide (DV-Pchlide) and then, in two steps, into monovinyl chlorophyllide
(MV-Chlide); chlorophyll is finally made by attachment of phytol to the MV-Chlide. Relative
values (peak areas) from six biological replicates analyzed in two runs are shown.
Statistically significant differences (p value ≤ 0.05) are printed in bold. Statistical details
are in **Supplementary Tables S5** and **S6**.

	CoPP	PPIX	MgP	MgPME	DV-Pchlide	MV-Chlide
WT	65.1	49.2	100.3	443.2	200.2	2328.6
Δ*sll0729*	**269.7**	**3921.2**	88.2	340.7	161.1	2359.5
P*slr1790*	51.5	32.4	53.6	254.6	194.6	2336.5
MP*slr1790*	56.6	26.5	46.8	242.6	182.3	1809.9
P*slr1790*/Δ*sll0729*	**430.8**	**3058.4**	82.2	556.2	161.4	3377.3
MP*slr1790*/Δ*sll0729*	34.8	15.3	43.7	147.4	103.5	2752.1

These results indicate that the lowered *hemJ* expression dramatically impacted the
tetrapyrrole biosynthetic pathway leading to the accumulation of the phototoxic

intermediates CoPP and PPIX. We also monitored the levels of later intermediates of the

chlorophyll biosynthetic pathway, but detected no clear differences, as in the case of

CoPP or PPIX. The lower chlorophyll level observed in the *sll0729* mutant (**Fig. 3C**) is

thus most likely caused by extremely high PPIX content, causing strong oxidative stress,

rather than restricted chlorophyll biosynthesis.

4. Discussion

The primary aim of our study was to identify the genetic changes in frequently occurring

suppressor clones that exhibited wild-type-like pigmentation in the presence of abolished

GG^{m4}CC methylation. A similar frequent appearance of suppressor mutations has recently

been reported for other *Synechocystis* 6803 mutants affected in the biosynthesis of

tetrapyrroles,[34,35], which reflects the essential role of this process in the highly pigmented

cyanobacterial cell. Our genome sequencing of suppressor clones for Δ*sll0729* revealed

one common SNV in the GGCC motif immediately downstream of the -10 sequence in the

slr1790 promoter towards GGTC. The absence of GGCC methylation resulted in slightly

diminished *slr1790* expression encoding HemJ, which is involved in porphyrin synthesis,

while the complementation of M.Ssp6803II activity due to ectopic expression of *sll0729*

reversed *slr1790* expression to WT levels.

Overall, our data indicate that the improper expression of *slr1790* due to missing ^{m4}C-

methylation in the *hemJ* promoter resulted in unbalanced porphyrin biosynthesis, leading

to massive accumulation of CoPP and PPIX. These compounds have great potential to

produce reactive oxygen species (ROS), which likely led to the observed strongly reduced

growth and pigmentation phenotype reported here and previously for *Synechocystis*

6803,[21] as well as in corresponding algal and plant mutants.[36,37] We present two main

findings, the physiological relevance of a particular GG^{m4}CC methylation and the dramatic

phenotypic effect of seemingly only slightly disturbed expression of HemJ.

Our data unambiguously show the relevance of GG^{m4}CC methylation at a critical promoter

position for the transcription of *hemJ* and the proper biosynthesis of essential tetrapyrroles

in *Synechocystis* 6803. If this setting evolved as the most physiological promoter

configuration, it should also be expected for related bacteria. There are not many

cyanobacteria with mapped transcription initiations sites, but for the related *Synechocystis*

sp. PCC 6714 (*Synechocystis* 6714) such a dataset exists.[38] The two strains share 2838

protein-coding genes, but also have 845 (*Synechocystis* 6803) and 895 unique genes

(*Synechocystis* 6714)[39] indicating substantial genetic differences. Interestingly, a GGCC

motif is situated at a comparable position between the -10 element and the transcription

start site of *hemJ* in *Synechocystis* 6714, as in *Synechocystis* 6803 (**Supplementary Fig.**

S7). Moreover, *Synechocystis* 6714 possesses the gene D082_01520 (GenBank

accession AIE72681), which encodes a likely ortholog of the DNA methyltransferase

M.Ssp6803II (59% identical and 73% similar amino acids). Hence, the potential impact of

GGCC methylation on *hemJ* expression likely is conserved beyond the here investigated

model strain *Synechocystis* 6803.

The change from C to T is often observed in eukaryotes, where deamination of the C5

methylated cytosine results in thymine, which escapes the DNA repair system. Such a

mechanism is unlikely in the case here because, first, the cytosine in the GGCC motif is

N4-methylated; second, the suppressor mutation occurred in cells in which the

methylation of this motif is abolished due to deletion of *sll0729*. Hence, we assume that

the exchange of C with T was related to the fact that thymine carries a methyl group, which

might mimic the previous GGCC motif methylation and thereby restores the proper

expression of *hemJ*. Both the methyl groups of ^{m4}C and thymidine protrude into the major

groove, thereby making it accessible for DNA-binding factors (**Fig. 5**).[2,40] It has been

shown that an interaction between the methyl group and certain amino acids via CH⋯π

hydrogen bonds[41] of the sigma factor could possibly lead to an stabilized open promoter

complex or facilitate recognition of the promoter sequence by the sigma factor.[42,43]

Additionally, exchanging C with T maintains a pyrimidine base at this position. Our

expression analysis in WT strains bearing a GGCC or GGTC motif containing *hemJ*

promoter in front of the *luxAB* reporter genes verified similar promoter activities of a

methylated WT promoter and the mutated promoter sequence (**Fig. 2B**).

Figure 5. DNA double helix structure of GGC/TC and chemical structures of thymidine and N4-methylcytosine. (A) The methyl groups of thymine (purple) and N4-methylcytosine (orange) protrude into the major groove. DNA double helix structure was generated with ChimeraX.[44] **(B)** Chemical structures of thymidine and N4-methylcytidine. The methyl groups are circled in red. Chemical structures were generated with ChemDraw.

Whether the observed differences in transcription were caused by a direct effect on RNA polymerase kinetics or by a change in transcription factor binding remains unclear. Nevertheless, the GGCC motif methylated by M.Ssp6803II is located in the discriminator sequence of the *hemJ* promoter, which has been shown to have a distinct impact on transcription initiation.[42] The promoter sequence directs the rate of transcription, starting with RNA polymerase binding, formation of the open promoter complex, and promoter escape, and is crucial for transcription initiation.[45] Apparent mild changes in this sequence can lead to major changes in transcription initiation.[46–48] Thus, it is not surprising that the absence of a normally present methyl group at this site can alter gene expression. In

another study, hemi-methylation of an adenine located at position -13 in the promoter

region of the IS10 transposase gene in the GATC nucleotide sequence led to the

activation of transcription.[12] Methylated bases such as ^{m4}C, ^{m5}C, and ^{m6}A affect the melting

properties of the DNA double helix.[49–52] During open complex formation of the promoter

sequence, the melting properties of the DNA double helix lead to altered promoter escape

by the RNA polymerase.[33] In addition, the sequence itself and DNA curvature (bending)

should not be neglected, as they could also be affected by the presence or absence of a

methyl group.[53]

The second main observation of this study was the strong effect of slightly disturbed *hemJ*

expression on the accumulation of certain tetrapyrrole biosynthesis intermediates. This

finding is best explained in the context of a multi-enzyme complex for tetrapyrrole

biosynthesis that supports the channeling of synthesis intermediates towards the metal

chelatases to avoid ROS production, as reported for various organisms.[23,54–56] The

decreased expression of *hemJ* in the absence of GGCC methylation likely led to disturbed

complex formation.

Our observation that the level of the first specific chlorophyll precursor, Mg-protoporphyrin

IX, was almost unaffected in the Δ*sll0729* strain (**Table 1**) indicates that mutant cells still

have sufficient PPIX as a substrate for Mg chelatase. However, protoporphyrinogen IX,

released from the complex, probably immediately oxidizes to PPIX (see **Fig. 4**) and

initiates ROS production, which then results in further metabolic disturbances and

bleaching. Ectopic expression of *hemJ* paralogs in the *hemJ* mutant background of

Synechocystis 6803 was not successful, likely because of disturbed porphyrin complex

formation.[57] This is another indication that the enzyme-bound conversion of these

intermediates in a porphyrin-synthesis complex is necessary for the subsequent correct

channeling of PPIX to the insertion of Fe^{2+} or Mg^{2+} by the respective chelatases. A similar

'bleaching' phenotype as in *sll0729* deletion strain has been described for a *Synechocystis*

ferrochelatase mutant accumulating around 40-times more PPIX than WT. This strain

progressively loses chlorophyll despite upregulation of the chlorophyll pathway.[58]

The massive accumulation of the intermediate PPIX in the *sll0729* mutant appears indeed

a bit counterintuitive; theoretically, the decreased expression of a key enzyme should

result in higher accumulation of its substrate but lower amounts of its products. However,

photodynamic herbicides such as acifluorfen, blocking the activity of protoporphyrinogen

IX oxidase in plants, have essentially the same effect as the *sll0729* deletion in

Synechocystis - a dramatic accumulation of PPIX causing severe oxidative damage.[59] In

the Δ*sll0729* strain, this extreme stress situation then initiates high pressure on the cell to

solve the problem via improved *hemJ* expression by the mutated GGTC site inside the

hemJ promoter.

Collectively, our study revealed an important role of ^{m4}C-methylation for proper gene

expression in *Synechocystis* 6803 and likely other cyanobacteria. Under laboratory

conditions, this regulatory feature seems to be particularly important for the regulation of

porphyrin biosynthesis, which needs to be tightly regulated to ensure substrate channeling

towards metal chelatases to avoid oxidative stress.

^{m4}C-DNA methylation is likely to have many more important functions, because highly

conserved orthologs of the enzyme M.Ssp6803II are encoded in the genomes of many

other cyanobacteria, and many more GGCC motifs exist in the genome sequences of

cyanobacteria.[20] In eukaryotes, epigenetic modifications relevant for gene expression

predominantly involve C5-methylcytosine and rarely N6-methyladenine, whereas the here

investigated N4-methylcytosine modification is frequent in bacteria. However, recently,

the relevance of N4-methylcytosine was discovered as an epigenetic mark in certain

invertebrates which use an enzyme acquired from bacteria by horizontal gene transfer.[60]

These findings and the results described here indicate that more attention should be paid

to the possible epigenetic effects of ^{m4}C-DNA-methylation, both in eukaryotic as well as

in prokaryotic organisms.

Acknowledgments

We thank Klaudia Michl, University of Rostock, for excellent technical assistance and
Ingeborg Scholz, University of Freiburg, for the *luxAB* strains. The help of Prof. Klaus
Herburger, University of Rostock, during the microscopic analyses of cell sizes is highly
appreciated. This study was funded by the German Research Foundation (DFG) (grant
HE 2544/12-2 to WRH, grant HA 2002/17-2 to MH) and by project
CZ.02.01.01/00/22_008/0004624 of the Czech Ministry of Education, Youth and Sports to
RS.

Data availability

The resequencing data produced in this study are available in the DRA/SRA database
with the accession numbers DRR585585–DRR585594, BioProject PRJDB18568.
Previously generated bisulfite raw data are available at
https://www.ncbi.nlm.nih.gov/biosample/8378604 (BioProject ID: PRJNA430784,
BioSample: SAMN08378604, Run: SRX3574087) and previously generated microarray
data in the GEO database at
https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE126282, accession number
GSE126282 (BioProject PRJNA521475).

Author contributions

597 WRH and MH designed the study. NSc and NSt performed most of the experiments and
598 evaluated data. SW and KNM performed resequencing of suppressor clones and data
599 analyses. NSc and RS performed pigment analysis and data analysis. WRH and MH
600 contributed to data analyses. WRH and MH drafted the manuscript. All authors read and
601 approved the final manuscript.

Supplementary Data

604 Supplementary data are available at DNARES online.

References

1. Sánchez-Romero, M. A., Cota, I., and Casadesús, J. 2015, DNA methylation in bacteria: from the methyl group to the methylome. *Curr. Opin. Microbiol.*, **25**, 9–16.
2. Jeltsch, A. 2002, Beyond Watson and Crick: DNA methylation and molecular enzymology of DNA methyltransferases. *Chembiochem Eur. J. Chem. Biol.*, **3**, 274–93.
3. Loenen, W. A. M., and Raleigh, E. A. 2014, The other face of restriction: modification-dependent enzymes. *Nucleic Acids Res.*, **42**, 56–69.
4. Loenen, W. A. M., Dryden, D. T. F., Raleigh, E. A., Wilson, G. G., and Murray, N. E. 2014, Highlights of the DNA cutters: a short history of the restriction enzymes. *Nucleic Acids Res.*, **42**, 3–19.
5. Blow, M. J., Clark, T. A., Daum, C. G., et al. 2016, The epigenomic landscape of prokaryotes. *PLOS Genet.*, **12**, e1005854.
6. Marinus, M. G., and Morris, N. R. 1973, Isolation of Deoxyribonucleic Acid methylase mutants of *Escherichia coli* K-12. *J. Bacteriol.*, **114**, 1143–50.
7. Palmer, B. R., and Marinus, M. G. 1994, The *dam* and *dcm* strains of *Escherichia coli*--a review. *Gene*, **143**, 1–12.
8. Collier, J. 2009, Epigenetic regulation of the bacterial cell cycle. *Curr. Opin. Microbiol.*, **12**, 722–9.
9. Kahramanoglou, C., Prieto, A. I., Khedkar, S., et al. 2012, Genomics of DNA cytosine methylation in *Escherichia coli* reveals its role in stationary phase transcription. *Nat. Commun.*, **3**, 886.
10. Moore, L. D., Le, T., and Fan, G. 2013, DNA methylation and its basic function. *Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol.*, **38**, 23–38.
11. Militello, K. T., Mandarano, A. H., Varechtchouk, O., and Simon, R. D. 2014, Cytosine DNA methylation influences drug resistance in *Escherichia coli* through increased *sugE* expression. *FEMS Microbiol. Lett.*, **350**, 100–6.
12. Roberts, D. 1985, IS10 transposition is regulated by DNA adenine methylation. *Cell*, **43**, 117–30.
13. Camacho, E. M., and Casadesús, J. 2005, Regulation of *traJ* transcription in the Salmonella virulence plasmid by strand-specific DNA adenine hemimethylation. *Mol. Microbiol.*, **57**, 1700–18.
14. van der Woude, M., Braaten, B., and Low, D. 1996, Epigenetic phase variation of the *pap* operon in *Escherichia coli*. *Trends Microbiol.*, **4**, 5–9.
15. Gaultney, R. A., Vincent, A. T., Lorioux, C., et al. 2020, 4-Methylcytosine DNA modification is critical for global epigenetic regulation and virulence in the human pathogen *Leptospira interrogans*. *Nucleic Acids Res.*, **48**, 12102–15.
16. Bolay, P., Dodge, N., Janssen, K., Jensen, P. E., and Lindberg, P. 2024, Tailoring regulatory components for metabolic engineering in cyanobacteria. *Physiol. Plant.*, **176**, e14316.
17. Hagemann, M., and Hess, W. R. 2018, Systems and synthetic biology for the biotechnological application of cyanobacteria. *Curr Opin Biotechnol*, **49**, 94–9.
18. Klähn, S., Opel, F., and Hess, W. R. 2024, Customized molecular tools to strengthen metabolic engineering of cyanobacteria. *Green Carbon*, **2**, 149–63.

19. Scholz, I., Lott, S. C., Behler, J., Gärtner, K., Hagemann, M., and Hess, W. R. 2019,
Divergent methylation of CRISPR repeats and *cas* genes in a subtype I-D CRISPR-
Cas-system. *BMC Microbiol.*, **19**, 147.1-147.11.
20. Hagemann, M., Gärtner, K., Scharnagl, M., et al. 2018, Identification of the DNA
methyltransferases establishing the methylome of the cyanobacterium
Synechocystis sp. PCC 6803. *DNA Res.*, **25**, 343–52.
21. Gärtner, K., Klähn, S., Watanabe, S., et al. 2019, Cytosine N4-methylation via
M.Ssp6803II is involved in the regulation of transcription, fine-tuning of DNA
replication and DNA repair in the cyanobacterium *Synechocystis* sp. PCC 6803.
Front. Microbiol., **10**, 1–14.
22. Kato, K., Tanaka, R., Sano, S., Tanaka, A., and Hosaka, H. 2010, Identification of a
gene essential for protoporphyrinogen IX oxidase activity in the cyanobacterium
Synechocystis sp. PCC6803. *Proc. Natl. Acad. Sci. U. S. A.*, **107**, 16649–54.
23. Skotnicová, P., Sobotka, R., Shepherd, M., Hájek, J., Hrouzek, P., and Tichý, M.
2018, The cyanobacterial protoporphyrinogen oxidase HemJ is a new b-type heme
protein functionally coupled with coproporphyrinogen III oxidase. *J. Biol. Chem.*, **293**,
12394–404.
24. Trautmann, D., Voß, B., Wilde, A., Al-Babili, S., and Hess, W. R. 2012, Microevolution
in cyanobacteria: re-sequencing a motile substrain of *Synechocystis* sp. PCC 6803.
DNA Res., **19**, 435–48.
25. Kaneko, T., Sato, S., Kotani, H., et al. 1996, Sequence analysis of the genome of the
unicellular cyanobacterium *Synechocystis* sp. strain PCC6803. II. Sequence
determination of the entire genome and assignment of potential protein-coding
regions. *DNA Res.*, **3**, 109–36.
26. Scharnagl, M., Richter, S., and Hagemann, M. 1998, The cyanobacterium
Synechocystis sp. strain PCC 6803 expresses a DNA methyltransferase specific for
the recognition sequence of the restriction endonuclease PvuI. *J. Bacteriol.*, **180**,
4116–22.
27. Kunert, A., Hagemann, M., and Erdmann, N. 2000, Construction of promoter probe
vectors for *Synechocystis* sp. PCC 6803 using the light-emitting reporter systems Gfp
and LuxAB. *J. Microbiol. Methods*, **41**, 185–94.
28. Pinto, F. L., Thapper, A., Sontheim, W., and Lindblad, P. 2009, Analysis of current
and alternative phenol based RNA extraction methodologies for cyanobacteria. *BMC
Mol. Biol.*, **10**, 1–8.
29. Kraus, A., Spät, P., Timm, S., et al. 2024, Protein NirP1 regulates nitrite reductase
and nitrite excretion in cyanobacteria. *Nat. Commun.*, **15**, 1911.
30. Pilný, J., Kopečná, J., Noda, J., and Sobotka, R. 2015, Detection and quantification
of heme and chlorophyll precursors using a high performance liquid chromatography
(HPLC) system equipped with two fluorescence detectors. *Bio-Protoc.*, **5**, e1390–
e1390.
31. Kopf, M., Klähn, S., Scholz, I., Matthiessen, J. K. F., Hess, W. R., and Voß, B. 2014,
Comparative analysis of the primary transcriptome of *Synechocystis* sp. PCC 6803.
DNA Res., **21**, 527–39.
32. Czarnecki, O., and Grimm, B. 2012, Post-translational control of tetrapyrrole
biosynthesis in plants, algae, and cyanobacteria. *J. Exp. Bot.*, **63**, 1675–87.

33. Chen, J., Darst, S. A., and Thirumalai, D. 2010, Promoter melting triggered by
bacterial RNA polymerase occurs in three steps. *Proc. Natl. Acad. Sci. U. S. A.*, **107**,
12523–8.
34. Chen, G. E., Hitchcock, A., Mareš, J., et al. 2021, Evolution of Ycf54-independent
chlorophyll biosynthesis in cyanobacteria. *Proc. Natl. Acad. Sci. U. S. A.*, **118**,
e2024633118.
35. Skotnicová, P., Srivastava, A., Aggarwal, D., et al. 2024, A thylakoid biogenesis BtpA
protein is required for the initial step of tetrapyrrole biosynthesis in cyanobacteria.
New Phytol., **241**, 1236–49.
36. Issawi, M., Sol, V., and Riou, C. 2018, Plant photodynamic stress: What's new?
Front. Plant Sci., **9**, 681.
37. Zhang, W., Willows, R. D., Deng, R., et al. 2021, Bilin-dependent regulation of
chlorophyll biosynthesis by GUN4. *Proc. Natl. Acad. Sci. U. S. A.*, **118**, e2104443118.
38. Kopf, M., Klähn, S., Scholz, I., Hess, W. R., and Voß, B. 2015, Variations in the non-
coding transcriptome as a driver of inter-strain divergence and physiological
adaptation in bacteria. *Sci. Rep.*, **5**, 9560.
39. Kopf, M., Klähn, S., Pade, N., et al. 2014, Comparative genome analysis of the
closely related *Synechocystis* strains PCC 6714 and PCC 6803. *DNA Res.*, **21**, 255–
66.
40. Cervi, A. R., Guy, A., Leonard, G. A., Téoule, R., and Hunter, W. N. 1993, The crystal
structure of N4-methylcytosine.guanosine base-pairs in the synthetic hexanucleotide
d(CGCGm4CG). *Nucleic Acids Res.*, **21**, 5623–9.
41. Wang, J., and Yao, L. 2019, Dissecting C-H⋯π and N-H⋯π interactions in two
proteins using a combined experimental and computational approach. *Sci. Rep.*, **9**,
20149.
42. Feklistov, A., Barinova, N., Sevostyanova, A., et al. 2006, A basal promoter element
recognized by free RNA polymerase sigma subunit determines promoter recognition
by RNA polymerase holoenzyme. *Mol. Cell*, **23**, 97–107.
43. Bae, B., Feklistov, A., Lass-Napiorkowska, A., Landick, R., and Darst, S. A. 2015,
Structure of a bacterial RNA polymerase holoenzyme open promoter complex. *eLife*,
4, e08504.
44. Pettersen, E. F., Goddard, T. D., Huang, C. C., et al. 2021, UCSF CHIMERAX:
Structure visualization for researchers, educators, and developers. *Protein Sci.*, **30**,
70–82.
45. Hook-Barnard, I. G., and Hinton, D. M. 2007, Transcription initiation by mix and match
elements: flexibility for polymerase binding to bacterial promoters. *Gene Regul. Syst.*
Biol., **1**, 275–93.
46. Minchin, S., and Busby, S. 1993, Location of close contacts between *Escherichia coli*
RNA polymerase and guanine residues at promoters either with or without consensus
-35 region sequences. *Biochem. J.*, **289**, 771–5.
47. Hook-Barnard, I. G., and Hinton, D. M. 2009, The promoter spacer influences
transcription initiation via σ[70] region 1.1 of *Escherichia coli* RNA polymerase. *Proc.*
Natl. Acad. Sci. U. S. A., **106**, 737–42.
48. Zenkin, N., Kulbachinskiy, A., Yuzenkova, Y., et al. 2007, Region 1.2 of the RNA
polymerase σ subunit controls recognition of the −10 promoter element. *EMBO J.*,
26, 955–64.

49. Engel, J. D., and Von Hippel, P. H. 1978, Effects of methylation on the stability of
nucleic acid conformations. Studies at the polymer level. *J. Biol. Chem.*, **253**, 927–
34.
50. Butkus, V., Klimasauskas, S., Petrauskiene, L., et al. 1987, Synthesis and physical
characterization of DNA fragments containing N4-methylcytosine and 5-
methylcytosine. *Nucleic Acids Res.*, **15**, 8467–78.
51. Buitrago, D., Labrador, M., Arcon, J. P., et al. 2021, Impact of DNA methylation on
3D genome structure. *Nat. Commun.*, **12**, 3243.
52. Rausch, C., Zhang, P., Casas-Delucchi, C. S., et al. 2021, Cytosine base
modifications regulate DNA duplex stability and metabolism. *Nucleic Acids Res.*, **49**,
12870–94.
53. Bae, S.-H., Cheong, H.-K., Cheong, C., Kang, S., Hwang, D. S., and Choi, B.-S.
2003, Structure and dynamics of hemimethylated GATC sites: implications for DNA-
SeqA recognition. *J. Biol. Chem.*, **278**, 45987–93.
54. Ferreira, G. C., Andrew, T. L., Karr, S. W., and Dailey, H. A. 1988, Organization of
the terminal two enzymes of the heme biosynthetic pathway. Orientation of
protoporphyrinogen oxidase and evidence for a membrane complex. *J. Biol. Chem.*,
263, 3835–9.
55. Masoumi, A., Heinemann, I. U., Rohde, M., Koch, M., Jahn, M., and Jahn, D. 2008,
Complex formation between protoporphyrinogen IX oxidase and ferrochelatase
during haem biosynthesis in *Thermosynechococcus elongatus*. *Microbiol. Read.*
Engl., **154**, 3707–14.
56. Medlock, A. E., Shiferaw, M. T., Marcero, J. R., et al. 2015, Identification of the
mitochondrial heme metabolism complex. *PloS One*, **10**, e0135896.
57. Kohata, R., Lim, H., Kanamoto, Y., et al. 2023, Heterologous complementation
systems verify the mosaic distribution of three distinct protoporphyrinogen IX oxidase
in the cyanobacterial phylum. *J. Plant Res.*, **136**, 107–15.
58. Sobotka, R., McLean, S., Zuberova, M., Hunter, C. N., and Tichy, M. 2008, The C-
terminal extension of ferrochelatase is critical for enzyme activity and for functioning
of the tetrapyrrole pathway in *Synechocystis* strain PCC 6803. *J. Bacteriol.*, **190**,
2086–95.
59. Lermontova, I., and Grimm, B. 2000, Overexpression of plastidic protoporphyrinogen
IX oxidase leads to resistance to the diphenyl-ether herbicide acifluorfen. *Plant*
Physiol., **122**, 75–84.
60. Rodriguez, F., Yushenova, I. A., DiCorpo, D., and Arkhipova, I. R. 2022, Bacterial
N4-methylcytosine as an epigenetic mark in eukaryotic DNA. *Nat. Commun.*, **13**,
1072.
779