Transcriptomic profiling of gill biopsies to define predictive markers for seawater survival in farmed Atlantic salmon

Authors:

Lars Grønvold[1], Mattis J. van Dalum[2], Anja Striberny[3], Domniki Manousi[1], Trine Ytrestøyl[4], Turid Mørkøre[1], Solomon Boison[5], Bjarne Gjerde[6], Even Jørgensen[2], Simen R. Sandve[1] & David G. Hazlerigg[2]

[1]Department of Animal and Aquaculture Sciences, Faculty of Bioscience, Norwegian University of Life Sciences, Ås, Norway

[2]Arctic Chronobiology and Physiology, Arctic and Marine Biology, UiT – the Arctic University of Norway, Tromsø, Norway

[3]Nofima AS, The Norwegian Institute of Food, Fisheries, and Aquaculture Research, Tromsø, Norway

[4]Nofima AS, The Norwegian Institute of Food, Fisheries, and Aquaculture Research, Sunndelsøra, Norway

[5]MOWI Genetics AS, Bergen, Norway

[6]Nofima AS, The Norwegian Institute of Food, Fisheries, and Aquaculture Research, Ås, Norway

Correspondence:

David Hazlerigg - david.hazlerigg@uit.no

Simen Sandve - simen.sandve@nmbu.no

Keywords:
Atlantic salmon, smolt, photoperiod, sea water survival, random forest model, gene expression,

Abstract

Wild Atlantic salmon migrate to sea following completion of a developmental process known as parr - smolt transformation (PST), which establishes a seawater (SW) tolerant phenotype. Effective imitation of this aspect of anadromous life-history is a crucial aspect of commercial salmon production, with current industry practice being marred by significant losses during transition from the freshwater (FW) to SW phase of production. The natural photoperiodic control of PST can be mimicked by exposing farmed juvenile fish to a reduced duration photoperiod for at least 6 weeks before increasing the photoperiod in the last 1 - 2 months before SW transfer. While it is known that variations in this general protocol affect subsequent SW performance, there is no uniformly accepted industry standard; moreover, reliable prediction of SW performance from fish attributes in the FW phase remains a major challenge. Here we describe an experiment in which we took gill biopsies 1 week prior to SW transfer from 3000 individually tagged fish raised on 3 different photoperiod regimes during the FW phase. Biopsies were subjected to RNA profiling by Illumina sequencing, while individual fish growth and survival was monitored over 300 days in a SW cage environment, run as a common garden experiment. Using a random forest machine learning algorithm, we developed gene expression-based predictive models for initial survival and stunted growth in SW. Stunted growth phenotypes could not be predicted based on gill transcriptomes, but survival the first 40 days in SW could be predicted with moderate accuracy. While several previously identified marker genes contribute to this model, a surprisingly low weighting is ascribed to sodium potassium ATPase subunit genes, contradicting advocacy for their use as SW readiness markers. However, genes with photoperiod-history sensitive regulation were highly enriched among the genes with highest importance in the prediction model. This work opens new avenues for understanding and exploiting developmental changes in gill physiology during smolt development.

Introduction

In anadromous salmonids, the transformation of freshwater resident juvenile fish (parr) into a migratory form (smolt) which goes to sea is known as smoltification, smolting or parr-smolt transformation (hereafter, PST). PST entails a complex combination of physiological and behavioural changes, amongst which the acquisition of the ability to efficiently maintain water and ionic balance is critical (McCormick 2013).

PST typically occurs after several years of slow growth in freshwater (FW) streams where spawning took place and requires parr to have exceeded a certain minimum threshold size at the end of the growing season preceding PST (Thorpe 1977; Handeland et al. 2013; Sigholt et al. 1995; Strand, Hazlerigg, and Jørgensen 2018). Exposure to a declining autumn photoperiod followed by increasing photoperiod the following spring triggers the process of PST, apparently through a temporally coordinated sequence of endocrine change (McCormick 2013). The photoperiod-dependent seasonal gating of PST ensures a synchronous springtime migration to sea, which is thought to reduce predation losses (Furey et al. 2016; Handeland et al. 1996).

Aquacultural production of Atlantic salmon (*Salmo salar*) depends on the ability to mimic natural PST to produce large numbers of seawater (SW) tolerant juvenile fish for transfer to sea cages where they can grow rapidly. There is no single universally accepted protocol for this commercial process. A widely used strategy for artificial smolt production is to rear hatchlings on a fast growth regime (i.e. continuous light (LL) and a fast growth diet for 6 - 12 months) to rapidly achieve threshold size, and then to expose them to short photoperiod (SP) for several weeks and before finally returning them to LL for the last 1 - 2 months of the FW phase. Based on observations of SW performance, it has been shown that the duration of exposure to SP should be at least six weeks long for LL to induce smelting (Duncan and Bromage 1998). The mechanisms through which PST runs as a photoperiodic history-dependent process remain unknown, and untangling the role of SP exposure in smolt

development is of considerable practical interest, as SP exposure reduces growth rates and

slows aquaculture production (Sigholt, Åsgård, and Staurnes 1998; Imsland, Handeland, and

Stefansson 2014).

To overcome this delaying effect of SP exposure on production schedules, alternative

strategies have emerged in which fish are maintained on constant light throughout the FW

phase, and SW tolerance is achieved through dietary manipulation with salt / tryptophan

enriched diets, water salinity manipulation, or by increasing size of fish (Striberny et al. 2021)

in the run up to SW transfer. At present there is no unifying consensus on the relative merits

of these different approaches, and difficulties in determining SW-readiness under different

protocols probably contribute to industry losses during the SW-transfer phase of salmon

production.

Because of its pivotal role in the regulation of water and ionic fluxes, undergoing a dramatic

switch from a salt conserving (FW) to salt excreting (SW) function, the gill has become the

focus of efforts to optimise commercial smolt production (Takvam et al. 2024). Within the gill,

mitochondria-rich cells (MRCs) are considered the primary drivers of ionic regulation (Evans

2008), and PST includes a pronounced shift in the location and phenotypic attributes of

MRCs in the gill (McCormick 2013; West et al. 2021). During PST, the gill complement of

MRCs shifts from an ion-absorbing FW type to an ion-secreting SW type, and the distribution

of MRCs shifts from the lamellae to the gill filament itself (Pisam et al. 1988; Madsen,

Kiilerich, and Tipsmark 2009). One of the most studied molecular changes in a the gill MRC

during PST is the shift in expression of the genes encoding the ion regulatory Na+, K+-

ATPase (NKA) pump and concurrent increase in NKA protein activity (Reviewed in Takvam et

al. 2024). Associated with this, relative levels of expression of two isoforms of the NKA α-

subunit shifts during PST, with α1a showing higher expression in parr and α1b showing

higher expression in smolts (McCormick, Regish, and Christensen 2009).

From an applied perspective, characterisation of gill transcriptomic changes during PST

provides a potential route to designing marker-based predictive strategies for optimising

smolt production. Current approaches focus on relative expression of NKA alpha subunit

isoforms (Takvam et al. 2024), but there is clearly potential to exploit other aspects of

transcriptomic change to optimise predictive power. We have previously hypothesised that

genes with photoperiod-history dependent regulation in the developing smolt gill would be

interesting candidates for novel smolt status markers (Iversen et al. 2020). In this study we

aim to test this, using a large-scale experiment in which gene expression prior to entering

SW is compared with individual fish performance in the SW phase, including survival after

SW transfer and stunted growth (i.e. loser fish (Noble et al. 2018)). We raised 3000 tagged

fish on 3 different light regimes, produced RNAseq from gill biopsies one week prior to SW

transfer, and subsequently recorded individual survival and growth in SW. We then applied a

machine learning approach to this unique dataset, attempting to link SW performance to FW

gill gene expression profiles. Our results demonstrate the potential for the development of

novel and improved markers for smolt status in salmon aquaculture.

Materials and Methods

Fish rearing experiment

For detailed description of the fish rearing see Gjerde et al. (Gjerde et al. 2024). In brief,

three groups of about 1000 juvenile Atlantic salmon (*Salmo salar*) were exposed to three

different photoperiods; LL (24 hours light), 12:12 (12 hours light / 24 hours) and 8:16 (8 hours

light / 24 hours). The LL group was kept from hatching until sea water (SW) transfer on

continuous light. The 8:16 group was reared on LL from hatching until the fish reached a

body mass of ~50g after which they were exposed to a 6-week period of short photoperiod

exposure (8L:16D - 8 hours day light and 16 hours darkness). After these 6 weeks followed

another 8 weeks on LL before sea water transfer when the fish reached a body mass of

~100g. The 12:12 group was, exposed to 12h day light and 12h darkness for 6 weeks,

following 8 weeks in LL before SW transfer. One week prior to sea water transfer, gill

biopsies were sampled and smolt index (i.e. skin colouration), weight and length were

measured. Approximately one week after gill biopsies were taken, the fish were moved to

one common seawater cage where they were kept for 10 months and daily recording of

mortality were carried out.

Ethical statement

The experiment was performed according to EU regulations concerning the protection of

experimental animals (Directive 2010/63/EU). Appropriate measures were taken to minimize

pain and discomfort. The experiment was approved by the Norwegian Food and Safety

Authority (FOTS id. number 25658).

RNAseq data generation

Gill biopsies were flash frozen on dry ice and sent to QIAGEN (Germany) for RNA-isolation

using the RNeasy Fibrous Tissue Mini kit. Isolated RNA samples were then shipped to

Novogen (UK) for RNA RNA-sequencing library construction which were sequenced using

2*150 bp pair end illumina sequencing.

RNAseq read mapping

Prior to transcript quantification, adaptor sequences were trimmed off using fastp version

0.23.2 (Chen et al. 2018). Read mapping and transcript quantification was done using

Salmon version 1.1.0 (Patro et al. 2017) with the Atlantic salmon transcriptome annotation

from ENSEMBL from the Ssal_v3.1 genome assembly (Ssal_v3.1, GCA_905237065.2) in

the 'selective mapping-mode'. A transcriptome index file was used as decoy sequence to

avoid the incorrect mapping of reads between annotated and unannotated yet highly similar

genomic regions (Srivastava et al. 2020). The options –keepDuplicates and –-*gcBias* was

used transcript quantification to account for the high sequence similarity between some of

the duplicated genes from the salmonid genome duplication (Lien et al. 2016) and correct for

fragment-level GC biases of aligned reads, respectively. To calculate mRNA expression at

the gene-level from output files from *Salmon* the sum of the raw read counts and normalized

gene expression values (transcripts per million) were done with the R package tximport (Soneson, Love, and Robinson 2015).

Differential gene expression analyses between samples with different photoperiod history

We carried out three different analyses of differential gene expression using gill RNAseq data to define a set of core genes whose expression during PST in the gill consistently depends upon photoperiodic history in the FW phase. Two of these analyses used previously published datasets (Iversen et al. 2020, 2021) (experiments 1 and 2) and the third analysis used data from this study (experiment 3) (Figure S1). In experiment 1 (Figure S1A) we contrasted 7 months old salmon (reared in fresh water) either raised on continuous light (LL group, n=6) or exposed to 60 days of short days (8L:16D) followed by 50 days in LL (SPLL group, n=6). In experiment 2 (Figure S1B) we contrasted 11 months old salmon (reared in fresh water) exposed to 2 weeks of short photoperiod (2w group, 8L:16D, n=6) with fish exposed to 8 weeks of short photoperiod (8w group, n=6). Both groups experienced an 8-week period on continuous light after the short photoperiod. In experiment 3 (Figure S1C) from this study we contrasted smolt raised on continuous light (LL, n=981) with smolt exposed to 6 weeks of short photoperiod treatments (8L:16D, n=927).

Differential expression in experiment 1 and 2 were analysed with EdgeR v3.42.4. Genes with low expression were filtered (min number of reads summed across all fish per group was 10) and normalized using trimmed means of M-values. A quasi-likelihood negative binomial generalized log-linear model with 0 as intercept was fitted to the data. Differentially expressed genes were identified in pairwise contrasts using empirical Bayes F-tests. P-values were adjusted for multiple testing using the BH method. Since large sample numbers have been shown to exaggerate false positive rates when applying standard differential gene expression methods (Li et al. 2022), DEGs from experiment 3 were identified using Wilcoxon rank-sum test. False discovery rate thresholds were defined as the square root of the minimum sample size per group, resulting in adjusted p-value cutoffs of <= 0.05 for

experiments 1-2 and <=0.002 for experiment 3. Finally, upset plots of shared significant

DEGs were made with the R package upsetR (Conway, Lex, and Gehlenborg 2017) to define

a core set of photoperiodic history dependent smolt development genes identified as DEGs

in all three analyses.

Differential gene expression between fish with different SW performance

DEG analyses was also used to identify gene expression differences in gill samples from FW

between smolts with different growth and mortality outcomes in SW. First, we contrasted

gene expression in fish from the LL group which died within 40 days post seawater transfer

versus fish that were still alive in November, approximately five months after sea transfer (i.e.

excluding fish that died of disease or was otherwise lost). Second, we contrasted gene

expression differences between loser fish with extremely low growth and 'winners' (i.e. fish

with normal growth r). Classification of "losers" was based on the growth rate (i.e. gained

weight divided by initial weight) where the threshold of 0.5 was decided based on the

minimum between the two peaks in the growth rate distribution (see Figure 2A). Read counts

were normalized across samples using EdgeR's TMM method and then converted to Counts

Per Million (CPM). These values were subsequently log-transformed to log2(CPM+1). Genes

with zero reads in more than 90% of the samples were excluded, resulting in the removal of

approximately 9,000 out of 47,000 genes. The final dataset of log-transformed expression

values was used for Wilcoxon rank-sum tests. Fold change values were calculated as the

difference of the mean log transformed expression values of the contrasted samples.

Random forest prediction

To build prediction models for SW survival and growth using FW gill gene expression data

from this study (experiment 3) we used the RandomForestClassifier from the sklearn python

module   (Pedregosa et al. 2011)   with   parameters   n_estimators=1000   and

class_weight='balanced' (otherwise defaults). We included the log transformed gene

expression as well as the commonly used smolt marker NKA ratio as features. NKA ratio was

calculated by subtracting the log transformed gene expression of NKA α1a

(ENSSSAG00000088099) from NKA α1b (ENSSSAG00000041746). This is equivalent to

dividing α1b/α1a in the non-transformed expression values. Samples were randomly split into

80% training samples and 20% test samples. This was repeated 100 times, i.e. random split

and model training, recording the importance scores of the genes and the prediction

probabilities of each sample. The mean of the importance scores and the prediction

probabilities of the test samples are reported in the analyses. We performed RF analyses for

both 'early death' and 'loser fish' outcomes. Both early death and loser fish was defined as

outlined for the DEG analyses.

Identification of Gill Cell-Type Specific Genes Using Single Nuclei RNAseq

To identify genes specific to different cell types in gill tissues, we utilized the single nuclei

RNA sequencing (snRNAseq) data provided by West et al. (2021). This dataset comprises

snRNAseq analyses performed on gill tissues sampled at various stages of smoltification

under different photoperiod regimes. Nuclei were assigned to 20 clusters based on their

expression patterns using Seurat (Hao et al. 2024). Each cluster was subsequently

annotated with a cell type using known marker genes and gene ontology (GO) analysis.

The expression levels of 2,000 variable genes were averaged across the nuclei in each cell-

type cluster and each genes was assigned to the assigned cell type of the cluster with the

highest average expression level. Since the original gene annotations from West et al. (2021)

were based on NCBI identifiers, we converted these to Ensembl gene IDs using the

g:Profiler's g:Convert utility (Kolberg et al. 2023). This conversion resulted in 1,354 cell-type

specific Ensembl genes mapped from 1,271 NCBI genes (i.e. not all genes could be mapped

and some had multiple matches).

Cell type enrichment analysis

Fisher's exact test using the *fisher.test()* in R (R Core Team 2021) was used to assess

whether the differentially expressed genes (DEGs) in early death are associated with certain

cell types. DEGs were identified based on their importance scores, with those exceeding 5e-

5 (top ~10%) being classified as DEGs. These DEGs were further divided into up-regulated

or down-regulated based on the fold change. For each cell type, we evaluated the

independence between a gene's classification as a DEG and its association with that

particular cell type. An odds ratio greater than one from this test suggests an enrichment of

DEGs that are specifically expressed in that cell type. Specifically, for up-regulated DEGs, it

suggests a higher abundance of the corresponding cell type in the gill samples from fish that

died early, and conversely, for down-regulated DEGs, it indicates a greater abundance in the

surviving fish. The p-values were adjusted for multiple comparisons using the *p.adjust()*

function in R (R Core Team 2021) with the Benjamini & Hochberg.

Results

Defining a core set of photoperiodic history dependent smolt gill development genes

Our reanalyses of the datasets from Iversen et al. (2020, 2021) (experiments 1 and 2) identified 781 and 6,923 genes with significant different expression levels between gill samples from fish with different photoperiodic history during the PST (Figure S1D). DEG analyses between gills from fish with different photoperiodic histories in this study (i.e. LL *vs* 8L:16D and LL *vs* 12L:12D) identified a total of 14,209 genes which were DEGs in one or both contrasts (Figure S1D). Finally, we intersected these three analyses to define a core set of 219 genes whose expression during PST gill consistently depends upon photoperiodic history in the FW phase (Table S1).

Random forest prediction models for mortality and growth stagnation

The main aim of this study was to test if smolt gill gene expression profiles in fresh water could be used to predict seawater performance at the individual fish level, and secondarily to determine whether candidate markers defined by their sensitivity to photoperiodic history constituted useful SW performance predictors. We made RF prediction models for two seawater outcomes; early mortality (<40 days after seawater transfer) and stagnation of growth, referred to as loser fish.

Early mortality was substantial (>8%, Figure 1A) in the LL group, while extremely low in both groups of fish exposed to reduced photoperiods during the FW phase (12L:12D, n=8 fish = 0.8% mortality) and (8L:16D, n=6 fish = 0.6% mortality). Hence, we only used the LL group to develop the RF prediction model for early mortality.

We tested the RF model on random subsets of LL fish not used in model training. This demonstrated good prediction capability (Figure 1B) with an average area under the receiver operating characteristic (auROC) of 0.82. In other words, in 82% of the predictions the model will correctly assign a higher absolute risk of mortality to a randomly selected fish that will die compared to a randomly selected fish which will not die.

Although an auROC > 0.80 is considered to be clinical useful in medicine (Çorbacıoğlu and

Aksel 2023), this prediction model will have relatively high false positive rate. Nevertheless, a

ranked list of the importance scores of the RF model features (i.e. gene expression levels)

can also be used to gain insights into biological processes which contribute to SW

performance. We therefore performed an in-depth analysis of the gene importance scores.

Interestingly, the most used molecular marker for smolt status, the two individual genes

forming the NKA-subunits (NKAα1a and NKAα1b) had low importance scores in the

prediction model (Figure 1C). We also included a proxy for the commercially available

SmoltVision test (Pharmaq Analytic, Norway) which is the ratio of mRNA levels for the two

NKA-subunits (NKAα1b:NKAα1a). Surprisingly, this NKA-ratio feature was only the 35[th] most

important feature in the RF model (Figure 1C, Table S2). However, another well-known

smolt-gill gene, the cystic fibrosis transmembrane regulator (CFTR) (Hiroi and McCormick

2012), had the highest importance in our RF model (Figure 1C). In addition, the top

predictive genes (importance > 0.001, 37 genes, Table S2) also contained other genes linked

to ion-transport such as the sodium/potassium-transporting ATPase subunit alpha-3 and the

patj gene, which is involved in tight junction formation in mammals (Shin, Straight, and

Margolis 2005).

We then explored the predictive importance of genes with photoperiod-history dependent

regulation (intersection between DEGs in experiment 1, 2, and this study, see Figure S1) for

initial SW survival. Among the 219 photoperiodic history-dependent genes, 48% were in the

top 10% most important genes in the RF model (yellow coloured points in Figure 1C). This

was a significantly higher percentage than expected by chance (Fisher test p-value=9.53 x

10^{-47}).

The development of the smolt gill involves a pronounced remodelling of its cellular

composition, notably in terms of its complement of chloride cells and of immune cell types

(McCormick 2013; West et al. 2021). Leveraging recently published data on gill cell type-

specific markers (West et al. 2021), we linked gene expression differences between surviving

fish and those which died early after seawater transfer to the cell types these differentially

expressed genes were associated with. Genes associated with mitochondria-rich cells and

one subtype of epithelial cells (1) were significantly more likely (FDR<0.05) to be expressed

more strongly in survivors compared to fish dying within 40 days of SW transfer (Figure 1D).

Genes associated with seven cell types, including three immune-system related cells, were

significantly more likely (FDR<0.05) to be expressed at lower levels in survivors compared to

fish that died within 40 days of SW transfer (Figure 1D).

Finally, we also used a similar RF approach to predict loser fish status, which is a common

challenge in aquaculture. Short photoperiod exposure protocols were not associated with

loser fish status; hence we combined all loser fish in one group (Figure 2A). Our results

clearly demonstrate that gill gene expression in the FW phase is not possible to use to train

an RF model to predict loser fish status in SW (Figure 2B).

Discussion

In the present study we aimed at identifying and validating gene expression markers for

smolt status by using gill gene expression at the end the smolt rearing in FW to predict

survival and performance in the SW phase. We could not predict loser fish outcomes based

on FW gill transcriptomes, indicating that this phenomenon is not mechanistically linked to

smolt gill development and ion regulation ability. Conversely, the prediction model for SW

survival had reasonable discriminatory ability (auROC=0.84), and results from the RF model

(Figure 2C) provide novel biological insights into the smolt gill and development of new

biomarkers for smolt status.

The most common molecular biomarker for smolt status is the NKA-subunit gene expression

ratio (NKAα1b: NKAα1a) (Reviewed in Takvam et al. 2024). Surprisingly, in this study (Figure

1C) the NKA-ratio was only ranked as the 35[th] most important feature in the prediction model

for early mortality in SW. This result is in line with a recent experiment by Kahw et al. (2021),

which found large variation in NKA-ratio related to fish genetics and smolt production

protocols (0+ vs 1+) and concluded that NKA-ratio was not a robust biomarker for smolt

status. Hence, Kahw et al. (2021) and our findings in this study questions the reliability of the

NKA-ratio as a good predictor for seawater survival and performance in aquaculture smolt

production.

Our large-scale transcriptomic approach using gene expression profiles from individual fish

therefore represents an important step towards identifying new and improved biomarkers for

smolt status. One interesting aspect of our list of putative smolt biomarkers (Table S1 and

S2) is the highly significant overrepresentation of core photoperiod-history sensitive genes

(Fisher test, p-value = 9.529355e-47, Figure 1C). The most predictive gene CFTR (Figure

1C) was not in the core photoperiod-history sensitive gene set, however CFTR was deemed

photoperiodic-history sensitive in two out of three experiments (Table S1). Among the top 36

features with highest prediction model importance (> 0.001), 12 were in the core photoperiod

sensitive gene list, including calpain 2, egln2, and TEFa. Interestingly, calpain2 was ranked

as the gene with the strongest photoperiod-history sensitive regulation during PST in Iversen

et al. (Iversen et al. 2020). Egln2 is known to interact with hypoxia inducible factors (HIFs) in

human studies according to the Molecular Interaction Database

(https://www.ebi.ac.uk/intact/) and in zebrafish larvae there is a bidirectional crosstalk

between HIFs and glucocorticoid signalling (Marchi et al. 2020). Furthermore, the D-element

transcription factor TEFa, which is implicated in light - circadian interactions in zebrafish

(Vatine et al. 2009), is a core part of photoperiod transduction in mammals (Dardente et al.

2010) and is directly responsive to cortisol in salmon (West et al. 2020). The presence of

egln2 and TEFa among the most predictive genes thus suggests a glucocorticoid- signalling

link to PST gill development and initial SW survival. In conclusion, our results points to a

mechanistic link between poor smolt development in the LL group and dysregulation of

molecular developmental processes that are naturally regulated by photoperiod signals.

Focusing on these "winter signal" affected genes is therefore an exciting avenue in future

research and validation of novel biomarkers for smolt physiology.

One caveat related to our prediction model results is that the data we used to build the model

was limited to data from fish raised on LL. This was because initial survival was >99% in

smolts raised using exposure to short photoperiods. Even though an LL protocol can

compromise the development of hypo-osmoregulatory ability as well as other smolt

development characteristics, our results are still of high relevance to the industry due to

extensive use of LL in smolt production.

359 Author contributions:

Mattis J. van Dalum: investigation, formal analysis, visualization, writing

Bjarne Gjerde: conceptualization, experimental design, manuscript editing

Solomon Boison: conceptualization, experimental design, manuscript editing

Lars Grønvold: investigation, formal analysis, writing, manuscript editing

David G. Hazlerigg: conceptualization, funding acquisition, writing and editing, supervision

Even Jørgensen: fish rearing and sampling, manuscript editing

Turid Mørkøre: fish phenotype recording, manuscript editing

Simen R. Sandve: conceptualization, funding acquisition, writing and editing, supervision

Anja Striberny: fish rearing and sampling, writing

Domniki Manousi: bioinformatic data analyses, manuscript editing

Trine Ytrestøyl: fish phenotype recording, manuscript editing

371 Acknowledgements

372   Thanks to MOWI for providing fish for this experiment and Sofie Robertson for exploring the

373   use of RF prediction models using gene expression data.

Funding acknowledgements

375   This study was funded by the Norwegian Seafood Research Fund (project no. 901589).

Data availability

377   All transcriptome data from the new experiments presented in this paper is deposited at ENA

378   under the project number $\texttt{ERP131717}$.

Figure and table legends

Figure 1: Random forest predictions of early mortality in seawater. A) Early mortality cases in LL group. B) Random forest model predictions for individual fish probability for mortality in an early phase (<40 days) following seawater transfer. Predictions are based on gill tissue gene expression profiles from smolts in fresh water prior to seawater transfer. C) Scatterplot of random forest model importance scores *vs* differential gene expression adjusted p-values from a contrast between fish dying early after seawater transfer and surviving fish. Yellow coloured dots represents genes with photoperiod history dependent regulation. Red dots represents classical smolt physiology states associated genes. D) Fisher test odds-ratios (i.e. enrichment) for cell type biassed/specific genes which are expressed higher and lower in surviving fish compared to fish dying early (<40 days) after seawater transfer.

Figure 2. Random forest predictions of loser fish status in seawater. A) Classification of loser fish and winner fish based on growth rates in seawater. B) Random forest model predictions for individual fish classification of loser fish (low or no growth in seawater) and winner fish (normal growth rates) following seawater transfer. F) Scatterplot of random forest model importance scores *vs* differential gene expression adjusted p-values from a contrast between fish classified as loser fish (low or no growth in seawater) and winner fish (normal growth rates) following seawater transfer.

Figure S1. Defining a core set of photoperiod history-sensitive genes
A) Experiment 1: a ´short day´ experiment comparing the effect of FW rearing in continuous light (LL) throughout with exposure to 8-h light / 24-h for a 6 week period. B) Experiment 2: a ´winter length´ experiment in which the duration of exposure to 8-h light / 24-h was either 2- or 8 weeks. C) Experiment 3: data from this study in which the effect of rearing on LL throughout were compared with exposure to either 12-h or 8-h light / 24-h for 6 weeks. D) Upset plot showing differentially expressed genes (DEGs) in each of the experiments. DEGs were filtered down to a common set of genes that had photoperiod sensitive regulation in all

three experiments (FDR adjusted p-value <= 0.002 for experiment 3 and <=0.05 for

experiments 1, and 2).

Table S1. Results from all DEG analyses and RF model

Table S2. Top predictive genes for mortality in the initial SW phase

References

Chen, Shifu, Yanqing Zhou, Yaru Chen, and Jia Gu. 2018. "Fastp: An Ultra-Fast All-in-One
FASTQ Preprocessor." *Bioinformatics* 34 (17): i884–90.
https://doi.org/10.1093/bioinformatics/bty560.

Conway, Jake R, Alexander Lex, and Nils Gehlenborg. 2017. "UpSetR: An R Package for the
Visualization of Intersecting Sets and Their Properties." *Bioinformatics* 33 (18): 2938–40.
https://doi.org/10.1093/bioinformatics/btx364.

Çorbacıoğlu, Şeref Kerem, and Gökhan Aksel. 2023. "Receiver Operating Characteristic
Curve Analysis in Diagnostic Accuracy Studies: A Guide to Interpreting the Area under the
Curve Value." *Turkish Journal of Emergency Medicine* 23 (4): 195–98.
https://doi.org/10.4103/tjem.tjem_182_23.

Dardente, Hugues, Cathy A Wyse, Mike J Birnie, Sandrine M Dupré, Andrew S I Loudon,
Gerald A Lincoln, and David G Hazlerigg. 2010. "A Molecular Switch for Photoperiod
Responsiveness in Mammals." *Current Biology* 20 (24): 2193–98.
https://doi.org/10.1016/j.cub.2010.10.048.

Duncan, Neil J, and Niall Bromage. 1998. "The Effect of Different Periods of Constant Short
Days on Smoltification in Juvenile Atlantic Salmon (Salmo Salar)." *Aquaculture* 168 (1–4):
369–86. https://doi.org/10.1016/S0044-8486(98)00363-9.

Evans, David H. 2008. "Teleost Fish Osmoregulation: What Have We Learned since August
Krogh, Homer Smith, and Ancel Keys." *American Journal of Physiology. Regulatory,*
Integrative and Comparative Physiology 295 (2): R704-13.
https://doi.org/10.1152/ajpregu.90337.2008.

Gjerde, Bjarne, Solomon Antwi Boison, David Hazlerigg, Trine Ytrestoyl, Turid Morkore,
Even Jorgensen, Anj Striberny, and Simen Rod Sandve. 2024. "Impact of Three Light
Smoltification Regimes on Performance and Genetic Parameters of Traits in Atlantic
Salmon." *BioRxiv*, April. https://doi.org/10.1101/2024.04.02.587721.

Handeland, S O, A K Imsland, B Th Björnsson, and S O Stefansson. 2013. "Long-Term
Effects of Photoperiod, Temperature and Their Interaction on Growth, Gill Na^+, K^+-ATPase
Activity, Seawater Tolerance and Plasma Growth-Hormone Levels in Atlantic Salmon
Salmo Salar." *Journal of Fish Biology* 83 (5): 1197–1209. https://doi.org/10.1111/jfb.12215.

Hao, Yuhan, Tim Stuart, Madeline H Kowalski, Saket Choudhary, Paul Hoffman, Austin
Hartman, Avi Srivastava, et al. 2024. "Dictionary Learning for Integrative, Multimodal and
Scalable Single-Cell Analysis." *Nature Biotechnology* 42 (2): 293–304.
https://doi.org/10.1038/s41587-023-01767-y.

Hiroi, Junya, and Stephen D McCormick. 2012. "New Insights into Gill Ionocyte and Ion
Transporter Function in Euryhaline and Diadromous Fish." *Respiratory Physiology &*
Neurobiology 184 (3): 257–68. https://doi.org/10.1016/j.resp.2012.07.019.

Houde, Aimee Lee S, Oliver P Günther, Jeffrey Strohm, Tobi J Ming, Shaorong Li, Karia H
Kaukinen, David A Patterson, Anthony P Farrell, Scott G Hinch, and Kristina M Miller.
2019. "Discovery and Validation of Candidate Smoltification Gene Expression Biomarkers
across Multiple Species and Ecotypes of Pacific Salmonids." *Conservation Physiology* 7
(1): coz051. https://doi.org/10.1093/conphys/coz051.

Iversen, Marianne, Teshome Mulugeta, Børge Gellein Blikeng, Alexander Christopher West,
Even Hjalmar Jørgensen, Simen Rød Sandven, and David Hazlerigg. 2020. "RNA Profiling
Identifies Novel, Photoperiod-History Dependent Markers Associated with Enhanced
Saltwater Performance in Juvenile Atlantic Salmon." *Plos One* 15 (4): e0227496.
https://doi.org/10.1371/journal.pone.0227496.

Iversen, Marianne, Teshome Mulugeta, Alexander West, Even H Jørgensen, Samuel A M
Martin, Simen Rød Sandve, and David Hazlerigg. 2021. "Photoperiod-Dependent
Developmental Reprogramming of the Transcriptional Response to Seawater Entry in
Atlantic Salmon (Salmo Salar)." *G3 (Bethesda, Md.)*, March.
https://doi.org/10.1093/g3journal/jkab072.

Khaw, Hooi Ling, Bjarne Gjerde, Solomon A Boison, Elise Hjelle, and Gareth F Difford. 2021.
"Quantitative Genetics of Smoltification Status at the Time of Seawater Transfer in Atlantic
Salmon (Salmo Salar)." *Frontiers in Genetics* 12 (November): 696893.
https://doi.org/10.3389/fgene.2021.696893.

Kolberg, Liis, Uku Raudvere, Ivan Kuzmin, Priit Adler, Jaak Vilo, and Hedi Peterson. 2023.
"G:Profiler-Interoperable Web Service for Functional Enrichment Analysis and Gene
Identifier Mapping (2023 Update)." *Nucleic Acids Research* 51 (W1): W207–12.
https://doi.org/10.1093/nar/gkad347.

Li, Yumei, Xinzhou Ge, Fanglue Peng, Wei Li, and Jingyi Jessica Li. 2022. "Exaggerated
False Positives by Popular Differential Expression Methods When Analyzing Human
Population Samples." *Genome Biology* 23 (1): 79. https://doi.org/10.1186/s13059-022-
02648-4.

Lien, Sigbjørn, Ben F Koop, Simen R Sandve, Jason R Miller, Matthew P Kent, Torfinn
Nome, Torgeir R Hvidsten, et al. 2016. "The Atlantic Salmon Genome Provides Insights
into Rediploidization." *Nature* 533 (7602): 200–205. https://doi.org/10.1038/nature17164.

Madsen, Steffen S, Pia Kiilerich, and Christian K Tipsmark. 2009. "Multiplicity of Expression
of Na+,K+-ATPase {alpha}-Subunit Isoforms in the Gill of Atlantic Salmon (Salmo Salar):
Cellular Localisation and Absolute Quantification in Response to Salinity Change." *Journal*
of Experimental Biology 212 (Pt 1): 78–88. https://doi.org/10.1242/jeb.024612.

Marchi, Davide, Kirankumar Santhakumar, Eleanor Markham, Nan Li, Karl-Heinz Storbeck,
Nils Krone, Vincent T Cunliffe, and Fredericus J M van Eeden. 2020. "Bidirectional
Crosstalk between Hypoxia-Inducible Factor and Glucocorticoid Signalling in Zebrafish
Larvae." *PLoS Genetics* 16 (5): e1008757. https://doi.org/10.1371/journal.pgen.1008757.

McCormick, S D, A M Regish, and A K Christensen. 2009. "Distinct Freshwater and
Seawater Isoforms of Na+/K+-ATPase in Gill Chloride Cells of Atlantic Salmon." *Journal of*
Experimental Biology 212 (Pt 24): 3994–4001. https://doi.org/10.1242/jeb.037275.

McCormick, S D. 2013. "Smolt Physiology and Endochrinology." In *Euryhaline Fishes; Fish*
Physiology. Vol. 32.

Patro, Rob, Geet Duggal, Michael I Love, Rafael A Irizarry, and Carl Kingsford. 2017.
"Salmon Provides Fast and Bias-Aware Quantification of Transcript Expression." *Nature*
Methods 14 (4): 417–19. https://doi.org/10.1038/nmeth.4197.

Pedregosa, Fabian, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion,
Olivier Grisel, Mathieu Blondel, et al. 2011. "Scikit-Learn: Machine Learning in Python."
Journal of Machine Learning Research.

Pisam, M, P Prunet, G Boeuf, and A Rambourg. 1988. "Ultrastructural Features of Chloride
Cells in the Gill Epithelium of the Atlantic Salmon, Salmo Salar, and Their Modifications
during Smoltification." *The American Journal of Anatomy* 183 (3): 235–44.
https://doi.org/10.1002/aja.1001830306.

R Core Team. 2021. *R: A Language and Environment for Statistical Computing* (version
v4.3). Computer software. Vienna, Austria: R Foundation for Statistical Computing.

Shin, Kunyoo, Sam Straight, and Ben Margolis. 2005. "PATJ Regulates Tight Junction
Formation and Polarity in Mammalian Epithelial Cells." *The Journal of Cell Biology* 168 (5):
705–11. https://doi.org/10.1083/jcb.200408064.

Sigholt, Trygve, Magne Staurnes, Hans J. Jakobsen, and Torbjørn Åsgård. 1995. "Effects of
Continuous Light and Short-Day Photoperiod on Smolting, Seawater Survival and Growth
in Atlantic Salmon (Salmo Salar)." *Aquaculture* 130 (4): 373–88.
https://doi.org/10.1016/0044-8486(94)00349-S.

Soneson, Charlotte, Michael I Love, and Mark D Robinson. 2015. "Differential Analyses for
RNA-Seq: Transcript-Level Estimates Improve Gene-Level Inferences. [Version 2; Peer
Review: 2 Approved]." *F1000Research* 4 (January): 1521.
https://doi.org/10.12688/f1000research.7563.2.

Srivastava, Avi, Laraib Malik, Hirak Sarkar, Mohsen Zakeri, Fatemeh Almodaresi, Charlotte
Soneson, Michael I Love, Carl Kingsford, and Rob Patro. 2020. "Alignment and Mapping
Methodology Influence Transcript Abundance Estimation." *Genome Biology* 21 (1): 239.
https://doi.org/10.1186/s13059-020-02151-8.

Strand, Jo E T, David Hazlerigg, and Even H Jørgensen. 2018. "Photoperiod Revisited: Is
There a Critical Day Length for Triggering a Complete Parr-Smolt Transformation in
Atlantic Salmon Salmo Salar?" *Journal of Fish Biology* 93 (3): 440–48.
https://doi.org/10.1111/jfb.13760.

Striberny, Anja, Daniel E. Lauritzen, Juan Fuentes, Marco A. Campinho, Pasqualina
Gaetano, Vilma Duarte, David G. Hazlerigg, and Even H. Jørgensen. 2021. "More than
One Way to Smoltify a Salmon? Effects of Dietary and Light Treatment on Smolt
Development and Seawater Growth Performance in Atlantic Salmon." *Aquaculture* 532
(February): 736044. https://doi.org/10.1016/j.aquaculture.2020.736044.

Takvam, Marius, Kristina Sundell, Henrik Sundh, Naouel Gharbi, Harald Kryvi, and Tom Ole
Nilsen. 2024. "New Wine in Old Bottles: Modification of the Na^+/K^+ ATPase Enzyme
Activity Assay and Its Application in Salmonid Aquaculture." *Reviews in Aquaculture* 16
(3): 1087–98. https://doi.org/10.1111/raq.12887.

Thorpe, J E. 1977. "Bimodal Distribution of Length of Juvenile Atlantic Salmon (*Salmo Salar*
L.) under Artificial Rearing Conditions." *Journal of Fish Biology* 11 (2): 175–84.
https://doi.org/10.1111/j.1095-8649.1977.tb04111.x.

Vatine, Gad, Daniela Vallone, Lior Appelbaum, Philipp Mracek, Zohar Ben-Moshe, Kajori
Lahiri, Yoav Gothilf, and Nicholas S Foulkes. 2009. "Light Directs Zebrafish Period2

Expression via Conserved D and E Boxes." *PLoS Biology* 7 (10): e1000223.
https://doi.org/10.1371/journal.pbio.1000223.

West, Alexander C, Marianne Iversen, Even H Jørgensen, Simen R Sandve, David G
Hazlerigg, and Shona H Wood. 2020. "Diversified Regulation of Circadian Clock Gene
Expression Following Whole Genome Duplication." *PLoS Genetics* 16 (10): e1009097.
https://doi.org/10.1371/journal.pgen.1009097.

West, Alexander C, Yasutaka Mizoro, Shona H Wood, Louise M Ince, Marianne Iversen,
Even H Jørgensen, Torfinn Nome, et al. 2021. "Immunologic Profiling of the Atlantic Salmon
Gill by Single Nuclei Transcriptomics." *Frontiers in Immunology* 12 (May): 669889.
https://doi.org/10.3389/fimmu.2021.669889.

Furey, Nathan B, Scott G Hinch, Arthur L Bass, Collin T Middleton, Vanessa Minke-Martin,
and Andrew G Lotto. 2016. "Predator Swamping Reduces Predation Risk during Nocturnal
Migration of Juvenile Salmon in a High-Mortality Landscape." *The Journal of Animal
Ecology* 85 (4): 948–59. https://doi.org/10.1111/1365-2656.12528.

Handeland, S O, T Järvi, A Fernö, and S O Stefansson. 1996. "Osmotic Stress,
Antipredatory Behaviour, and Mortality of Atlantic Salmon (*Salmo Salar*) Smolts."
Canadian Journal of Fisheries and Aquatic Sciences 53 (12): 2673–80.
https://doi.org/10.1139/f96-227.

Imsland, Albert Kjartansson, Sigurd Olav Handeland, and Sigurd Olav Stefansson. 2014.
"Photoperiod and Temperature Effects on Growth and Maturation of Pre- and Post-Smolt
Atlantic Salmon." *Aquaculture International* 22 (4): 1331–45.
https://doi.org/10.1007/s10499-014-9750-1.

Noble, C, K Gismervik, M H Iversen, and J Kolarevic. 2018. "Welfare Indicators for Farmed
Atlantic Salmon: Tools for Assessing Fish Welfare."

Sigholt, Trygve, Torbjørn Åsgård, and Magne Staurnes. 1998. "Timing of Parr-Smolt
Transformation in Atlantic Salmon (Salmo Salar): Effects of Changes in Temperature and
Photoperiod." *Aquaculture* 160 (1–2): 129–44. https://doi.org/10.1016/S0044-
8486(97)00220-2.