Tree nursery environments and their effect on early trait variation

Annika Perry[1*], Joan K. Beaton[2], Jenni A. Stockan[2], Glenn R. Iason[2], Joan E.
Cottrell[3] and Stephen Cavers[1]

[1]*UK Centre for Ecology & Hydrology (UKCEH), Bush Estate, Midlothian, EH26 0QB,*
UK

[2]*Ecological Sciences, James Hutton Institute, Craigiebuckler, Aberdeen, AB15 8QH,*
UK

[3]*Northern Research Station, Forest Research, Roslin, EH25 9SY, UK*

Corresponding author: annt@ceh.ac.uk

Abstract

Despite the major role of nurseries in raising young plants and trees prior to
transplantation, not enough is known about how the nursery climate impacts the
growth and development of plants from germination through to maturity. It is
important for forestry practitioners to understand the effect that different nursery
environments may have on early stage growth as these may exceed differences due
to genetic variation and can confound the use of early stage traits for selection. Here,
a replicated progeny-provenance experiment of the economically and ecologically
important species Scots pine (*Pinus sylvestris* L.) was established in three
environmentally distinct nurseries in Scotland and traits including survival, growth,
form and phenology were measured. Temperature variation and photoperiod were
the only uncontrolled environmental variables during this period, and their effect on
measured traits was found to be significant among nurseries from the first growing
season onwards. Trait interactions were not consistent between nurseries, indicating
that the effectiveness of using proxy traits to select for desirable characteristics may
depend on the environment in which the trees are grown. This study is the first in a
series that will examine trait variation in Scots pine from seedlings to mature trees
and highlights the importance of carefully considering and accounting for the nursery
environment when growing trees for subsequent transplantation.

Introduction

Globally, forests function as the major terrestrial carbon sink (Pan *et al.* 2011) and
increasing the area of forest cover is vital to achieving net zero targets (Westaway *et*
al. 2023). In addition, increased forest cover would help tackle biodiversity loss,
increase filtering of pollutants, provide natural management of flood risk and
contribute to soil stabilisation and water purification (Holl and Brancalion 2020).
Ongoing afforestation efforts create an increased need for planting stock, which is
most commonly supplied as nursery-grown saplings (Brancalion and Holl 2020) and
this places the forest nursery in a pivotal and crucially important position. In Great

Britain alone, forest nurseries produced around 173 million saplings for the year
2022-2023 (Forestry Commission 2023).

In their role as the primary location for raising young trees, choices made by
nurseries have been shown to profoundly influence size and development of young
trees, effects that may be observable long after transplantation. For example, the
choice of fertiliser (Gruffman *et al.* 2012), growing medium (Heiskanen and Rikala
2000), container type, root management, watering regime (Villar-Salvador *et al.*
1999) and timing of operational activities (such as transplantation, Luoranen 2018)
during early life stages can affect root growth, seedling morphology, and
susceptibility to pests and pathogens (Selander and Immonen 1992). Whilst soil,
water and nutrients can be controlled quite precisely by nursery managers, the
geographic location of a nursery and any protection measures used, such as
glasshouse cover, determine other critical components of the nursery environment.
As variables like temperature and photoperiod can strongly affect the growth,
development and biochemistry of seedlings (Downs and Borthwick 1956;
Vapaavuori, Rikala and Ryyppö 1992; Nerg *et al.* 1994; Domisch, Finér and Lehto
2001) with effects lasting well beyond transplantation (Dhar *et al.* 2015), the nursery
environment can be hugely influential.

The use of locally sourced seed is an explicit requirement when planting native
species in the UK (Herbert, Samuel and Patterson 1999), but the location of the
nursery in which the resulting plants are raised is rarely specified. For example, the
contemporary guiding document for UK forestry – The UK Forestry Standard (Forest
Research 2023) - does not comment on nursery management, nor does it connect
nursery practice to its recommendations on seed sourcing. Previous publications,
such as the Forestry Nursery Practice (Aldhous and Mason 1994), do consider site
selection factors, but largely from the point of view of maximising plant performance,
minimising risk and logistical considerations. In practice, choice of nursery is
generally governed by what the market offers and by economics. Consequently, the
nursery site may be environmentally distant from both the site of seed origin and
planting site of raised seedlings. If the effect of the nursery environment on plant
phenotype is large, then the effect of local seed sourcing may be minimal in
comparison. A quantitative evaluation of the impact of the nursery environment on
trait variance is therefore potentially very important to seed sourcing policy, nursery
practice and to the long-term success of planting initiatives.

The impact of the early growing environment is also a critical factor to consider in
tree improvement and breeding programmes, particularly with regards to its effect on
survival and on variation in desirable traits. Traditional breeding practice, which
makes selections at around half rotation length (Zobel and Talbert 1984), is highly
inefficient for trees with long rotation times and there is much interest in methods for
screening trait variation at the nursery stage (Wu *et al.* 1997). However, trait values
measured prior to transplantation (ca. two years) rarely predict values in later years

(Burdon, Bannister and Low 1992; Hong, Fries and Wu 2015; Dong *et al.* 2018)

(apart from species with very short rotation lengths, e.g. *Eucalyptus grandis* in

Colombia, (Osorio, White and Huber 2003)). The use of 'proxy' traits to enable early

selection relies on the stability of trait-trait relationships among environments, which

may (Fukatsu *et al.* 2011) or may not be (Marron, Dillen and Ceulemans 2007)

identifiable. Significant variation in age-age and trait-trait correlations among

environments (Williams 1988; Osorio, White and Huber 2003) means that nursery

effects may confound selection based on early evaluation of trait variation.

Identifying such nursery effects is particularly relevant for widely planted,

economically relevant species. *Pinus sylvestris* L. (Scots pine) is a globally important

tree species with a natural range that extends from southern Spain to northern

Finland, and from western Scotland to far-eastern Russia. As a foundation species

with high conservation value, it has been shown to exhibit high adaptive trait

variation at both regional and international scales (Salmela *et al.* 2011; Perry *et al.*

2016; Donnelly *et al.* 2018; Benavides *et al.* 2021; Ramírez-Valiente *et al.* 2021).

Pinus sylvestris is also an economically valuable timber production species and is

widely planted both within and beyond its natural range: it is estimated that it

occupies more than 20 percent of the productive forest area in the EU (Mason and

Alía 2000) and, within Great Britain, it is the second most abundantly grown species

in tree nurseries after Sitka spruce (*Picea sitchensis*, Forestry Commission 2023).

Globally, *Pinus sylvestris* grows (both naturally and planted) across a vast range. Its

distribution is extremely environmentally heterogeneous and the species has

adapted to survive across a massive spectrum of temperature profiles, from hugely

variable to relatively stable. *Pinus sylvestris*' ability to grow across a huge

environmental gradient is also associated with a strong phenotypic variation.

Common garden trials are widely used in forestry to quantify the genetic component

of phenotypic variation. Replicated common gardens in different environments can

be used to examine the environmental component of phenotypic variation. Nurseries,

in their efforts to minimise environmental variation within a site and to standardise

conditions for each plant, are functionally common gardens. By making use of a

controlled, replicated common garden experimental design in nursery settings, we

can identify the contribution of the nursery environment to phenotypic variation. This

allows evaluation of the importance of the nursery environment to longer term

development of the plant and establishes a baseline for interpreting field trial results

in years to come. In addition, these results will enable age-age correlations to be

explored as the trees mature in field trials.

For this study, we compared early life stage traits of *P. sylvestris* seedlings grown in

three nursery environments (subsequently planted in multisite common garden field

trials in 2012, for full description see Beaton et al., 2022; Perry et al., 2024b). The

plants in each nursery trial had a common genetic background (seed collected

systematically from across the natural range in Scotland and arranged in a fully

replicated design), were grown under standard soil, nutrient and watering regimes
and were managed and measured by the same team of people. As the first paper in
a series exploring adaptive trait variation in these trials we focus here exclusively on
the effect of nursery environment on overall trait variation (although the experimental
design allows for genetic effects on early trait variation to be examined, we do not
deal with these here). We quantified the environmental effect on early growth and
development and survival and examined relationships within and among traits.

Methods

Source of material

Seedling source and nursery environments are detailed in Beaton *et al.* (2022).
Briefly, seed was collected from 10 mother trees from each of 21 provenances of *P.*
sylvestris from across the native range in Scotland in March 2007. Following
stratification, seeds were germinated at the James Hutton Institute, Aberdeen
(latitude 57.133214, longitude -2.158764) in June 2007 and transplanted into
individual pots. The full final collection consisted of 210 families (10 families from
each of 21 provenances) each consisting of 24 half-sibling progeny (total 5,040
individuals).

Nursery environment

After transfer into individual pots, and with the intention of eventually using these
plants to establish field based provenance trials, 8 seedlings per family were moved
to one of three nurseries (total 1,680 seedlings per nursery) in July 2007: an
unheated nursery glasshouse (NG) at the James Hutton Institute, Aberdeen; an
outdoor nursery in the west of Scotland (NW) at Inverewe Gardens, (latitude
57.775714, longitude -5.597181); an outdoor nursery in the east of Scotland (NE) at
the James Hutton Institute (location as above). In each nursery, trees were arranged
in 40 randomised complete blocks, where each block contained two trees per family
(total 42 trees). Watering was automatic in NG, and manually, as required, for NE
and NW. The watering regime ensured that water was never limiting and
waterlogging of seedlings was avoided. No artificial light was provided in any of the
nurseries. In May 2010, the pots containing the seedlings from NG were moved
outdoors to Glensaugh in Aberdeenshire (latitude 56.89, longitude -2.54).

Abiotic variation among nurseries was controlled as much as possible by growing the
seedlings under standard soil, nutrient and watering regimes. The major sources of
abiotic variation which could not be/were not controlled were temperature and wind,
of which temperature was measured throughout their growth in the nursery
environments. Biotic interactions were not controlled or measured during this period.

Hourly temperature was recorded at each of the nursery sites using data loggers
from July 2007 until December 2010. Temperature data loggers were suspended 50

cm above the ground under an aluminium foil-covered funnel. Daily minimum,
maximum and average temperatures (between 0000-2300 each day) and the daily
temperature variances were calculated for each nursery over this period (Perry et al.,
2024a). Daily variance was also partitioned into separate values for the day and
night period, where day variance was estimated between the hours of sunrise and
sunset between 0000-2300 and night variance was estimated between the hours of
sunset and sunrise between 0000-2300. Sunset and sunrise for each day at each
nursery were obtained from www.timeanddate.com.

On the few occasions on which more than one hourly record per 24 hours was
missing, the daily temperatures were replaced with average daily temperature data
derived from the nearest weather stations to the nurseries (information provided by
the National Meteorological Library and Archive, Met Office, UK) for the period July
2007 to December 2010). Weather stations were: Poolewe (0.7 miles from NW);
Craibstone (4.3 miles from NE and NG) and Glensaugh No 2 (0.1 miles from NG
when moved outdoors to Glensaugh). Where weather station data were missing
(only relevant for Poolewe over this period) they were replaced with measurements
from the next closest weather station: Aultbea No 2 (4.2 miles from NW). Due to the
differences in the location between the weather stations and the nurseries and the
different methods of recording temperature, data from weather stations were
adjusted prior to inclusion in the 'nursery temperatures' dataset by comparing daily
temperatures across the whole period. Missing logger temperatures were directly
replaced, where possible, by checking the weather station temperature at the
missing time point and calculating the mean logger temperature recorded at all other
instances that the same weather station temperature was recorded. In the few
instances where there were no direct replacements, replacement logger values were
obtained by taking the mean value for the three logger records either side of the
missing logger data point in a dataset which was ordered by the weather station
temperatures. For missing data in NG (prior to May 2010), replacement
temperatures derived from NE were used instead of weather station data. Where
there were days with missing hourly data for one of more of the nurseries,
temperature variance could not be estimated and so these days were excluded from
analyses for all nurseries to avoid skewing results.

The growing season length in each year at each nursery was estimated as the
period bounded by a daily mean temperature >5 °C for >5 consecutive days and
daily mean temperature <5 °C for >5 consecutive days (after 1st July). During the
defined growing season period trees were assumed to be 'growing' whereas outwith
the defined growing season trees were assumed to be 'not growing'. The previously
described temperature variances were compared between these two different
periods. The growing degrees were estimated for each day in the period July 2007 to
December 2010 as the number of degrees above 5 °C. The cumulative growing
degree days (GDD) were estimated by summing the growing degrees for each
period leading up to phenology assessments at each nursery from 1 January in

2008. Chilling days were defined as the days in which the average maximum
temperature did not exceed 5 °C.

Phenotype assessments

203   Seedling phenotypes were assessed for all individuals for a range of traits reflecting
204   different life history characteristics: growth, form, phenology and survival.

i. Growth traits

Growth traits were assessed annually from 2007-2010. Annual increments in height
(HI, mm) and basal stem diameter (DI, mm) were measured as the increase from the
end of one growing season to the end of the next (for comparisons of height
increment across multiple years, height in the first year of growth in 2007 is
considered as the height increment for this year), whilst absolute height (HA, mm)
and absolute basal stem diameter (measured at the root collar: DA, mm) were
measured at the end of each growing season. Relative growth rate (HR) was
estimated as the percentage increase in HI relative to HA in the previous year and
was recorded for 2008 to 2010.

ii. Form traits

216   Form was measured annually from 2007-2009. Needle length (NL, mm) was
217   measured for three randomly selected needles per tree and a mean value was
218   obtained. The total number of buds (Bu) on each seedling was counted in 2008 and
219   2009. Slenderness (HD, the ratio of HA to DA) was recorded in 2010 as a measure
220   of the impact of multiple years of growth on tree form.

iii. Phenology traits

Phenology was assessed weekly during the spring and autumn of 2008. Budburst
timing (BB) was defined as the number of days from 31 March 2008 to the time when
newly emerged green needles were observed. The estimated proportion of trees
which had burst bud in each nursery was calculated at each assessment. To
compare budburst progression across the three sites, the point at which an
estimated 50 % of trees had undergone budburst (BB_{50}) was calculated as follows:

$$BB_{50} = (50 - B_{i<50}) - \frac{B_{i>50} - B_{i<50}}{A_{i>50} - A_{i<50}} + A_{i<50}$$

where $i<50$ is the ith assessment performed at each nursery where < 50 % of trees
had undergone budburst ($i<50$: NE = 2; NG = 3; NW = 2); $i>50$ is the ith assessment
where > 50 % of trees had undergone budburst ($i>50$: NE = 3; NG = 4; NW = 3); A is
the variable associated with each assessment performed at each nursery (A: number
of days after 31 March; or cumulative growing degree days since 1 January); and B

is the percentage of trees observed with budburst at each nursery at each
assessment.

Growth cessation (GC) was defined as the number of days from 10 September 2008
to the date when no further height growth was observed by eye. To compare the
growing season length at each nursery with the duration of growth observed in the
seedlings, 'growth duration' (GD) was calculated for each seedling as the number of
days between budburst and growth cessation.

iv. Survival traits

241 Mortality (number of trees which died by the end of a given growing season as a
242 percentage of the total number of trees alive at the end of the previous growing
243 season) was recorded each year from 2007 to 2010. Survival was recorded as the
244 percentage of surviving trees in each nursery by 2010).

Statistical analysis

Paired t tests were performed using R (R Core Team 2024) to compare nursery
means for annual climatic variables (mean daily temperature, daily variance, growing
degree days) and climatic variables measured within and outside the growing
season (day temperature variance and night temperature variance). Where there
were days with missing hourly data for one of more of the nurseries, temperature
variance could not be estimated and so these days were excluded from analyses for
all nurseries to avoid skewing results.

Coefficient of variation, calculated as standard deviation divided by the mean, was
estimated for each trait in each nursery (Figure S1). Where traits were measured in
multiple years only the most recent year was used.

To evaluate relationships among traits, Pearson's correlation coefficients and
significance values were estimated for trees growing in each nursery using the
'Hmisc' package (Harrell Jr 2024) in R (R Core Team 2024). Absolute height,
absolute basal stem diameter and slenderness were only included for the final
assessment year (2010).

Nested analyses of variance (ANOVA) were performed in Minitab version 21 (Minitab
2024) for all trees with nursery as a fixed effect and block nested within nursery as a
random effect. Nested ANOVA was also used to analyse the distribution of variance
among these traits. In order to account for variance due to relatedness and/or
phenotypic plasticity which would otherwise fall within the residual variance,
additional terms (provenance; family nested within provenance; a nursery and
provenance interaction term) were included but were not considered separately,
given that the focus of this study is on the impact of environment on trait variation.
Tukey's post-hoc tests were performed to determine whether mean values among
nurseries were significantly different from one another for all traits and for mean

temperature, temperature variance (daily, day and night) and growing degree days
for each year. ANOVAs were repeated with all terms included as random effects to
identify their contribution to the total variance.

Results

Climatic variation among nurseries

Trees growing at each of the three nurseries experienced broadly similar mean daily
temperatures for each year between 2007 and 2010 which were not significantly
different among nurseries (Figure 1) with the exception of NE and NG in both 2008
and 2009, where mean daily temperature at NG was significantly warmer than at NE
in both years and warmer in NG than NW in 2009. Trees at NG also experienced the
fewest chilling days (total 119 between 2007 and 2010, compared with NE = 157 and
NW = 123) and had higher daily GDD throughout the majority of the year (Figure 1)
compared to the other two nurseries. In contrast, the daily temperature variance that
trees were exposed to was consistently and significantly lower for trees growing at
NW than for trees growing at either NG or NE (Figure 1). The daily temperature
variance that trees experienced in NG and NE was also significantly different,
although the magnitude of the difference was less than when comparing NW with
either NE or NG (Figure 1).

Figure 1. Climatic variation at the three nurseries for the period July 2007 to
December 2010 when seedlings were growing in individual pots at the nursery sites.
Lines indicate smoothed monthly means for each year. Boxplots indicate values for
the entire year: solid grey lines indicate the median value; the bottom and top of the
boxes indicate the first and third quartile; and the upper and lower whiskers extend to
the highest and lowest values within 1.5 times for interquartile range (no outliers are
shown for ease of visualisation. No data are included for daily temperature variance
in 2007 due to the extent of missing hourly data for this year. Significant differences
among nurseries within each year are indicated with asterisks (*: p = 0.01-0.05; **: p
= 0.001-0.01; ***: p = 0.0001-0.001; ****: p < 0.0001; ns: not significant. Units: mean
temperature, °C; daily temperature variance, °C², growing degree days, number of
degrees above 5 °C for each day.

Differences in temperature variance were most pronounced among nurseries and
consistent among years during the daytime within the growing season period (Figure
S2). During this period, daytime variance was significantly higher at NG than at NE
or NW, and higher at NE than at NW in all years. In contrast, variance during the
night-time was lowest at NG compared to the outdoor nurseries in all years outside

the growing season (mean night temperature variance 2008-2010: NE = 2.40 °C^2;
NG = 1.35 °C^2; NW = 2.17 °C^2).

Trait variation among nurseries

Trees growing in NG had higher mean values for traits relating to growth than those
growing outdoors in NE and NW by 2010 (Table 1): they had thicker stems and grew
taller and faster, on average. They also had larger increments for both height and
basal stem diameter in 2010. Trees grown in NE were the smallest and thinnest with
the slowest growth rate and lowest increments on average in 2010. Mean values for
traits relating to form and survival were lower, and budburst timing was earlier, for
trees growing in NG than those growing in one or both outdoor nurseries (Table 1)
with the exception of needle length.

Table 1. Traits measured in individual seedlings, grouped into the following
categories: 'Growth', 'Form', 'Phenology' and 'Survival'. Overall means and
associated standard errors are provided for each nursery (NE, NG and NW)
separately. Tukey pairwise comparisons are reported as a subscript letter (A, B, C)
after the mean value: nurseries that do not share a letter (within traits for each year)
have significantly different means. Number of observations for each nursery are in
parentheses.

Trait/year code*	Unit	NE	NG	NW
Growth				
DA08	Mm	3.58[A] ± 0.02 (1465)	2.87[C] ± 0.02 (1348)	2.96[B] ± 0.03 (1121)
DA09	Mm	5.74[A] ± 0.02 (1438)	4.80[B] ± 0.02 (1248)	4.69[C] ± 0.04 (1050)
DA10	Mm	6.77[C] ± 0.04 (1419)	8.81[A] ± 0.04 (1247)	7.19[B] ± 0.07 (1029)
DI09	Mm	2.14[A] ± 0.02 (1438)	1.90[B] ± 0.02 (1247)	1.71[C] ± 0.03 (1048)
DI10	Mm	1.08[C] ± 0.02 (1416)	4.01[A] ± 0.04 (1246)	2.48[B] ± 0.04 (1029)
HA08	Mm	81.12[C] ± 0.5 (1192)	100.36[A] ± 0.83 (1319)	88.32[B] ± 0.82 (1014)
HA09	Mm	237.76[A] ± 1.33 (1440)	241.14[A] ± 1.9 (1246)	216.9[B] ± 2.51 (1036)
HA10	Mm	322.80[C] ± 1.69 (1416)	382.73[A] ± 2.03 (1245)	334.87[B] ± 4.13 (775)
HI07	Mm	59.98[A] ± 0.3 (1611)	57.30[B] ± 0.32 (1498)	47.92[C] ± 0.29 (1492)
HI08	Mm	21.59[C] ± 0.46 (1192)	42.93[A] ± 0.73 (1319)	40.14[B] ± 0.80 (1014)
HI09	Mm	164.83[A] ± 1.23 (1180)	142.54[B] ± 1.46 (1218)	138.72[B] ± 2.03 (952)
HI10	Mm	84.20[C] ± 0.75 (1415)	141.48[A] ± 1.24 (1238)	125.84[B] ± 1.85 (767)
HR08	%	40.11[C] ± 1.08 (1192)	77.85[B] ± 1.50 (1319)	90.59[A] ± 2.15 (1014)
HR09	%	212.06[A] ± 2.05 (1180)	150.43[C] ± 1.87 (1218)	162.66[B] ± 2.45 (952)
HR10	%	36.55[B] ± 0.41 (1415)	65.81[A] ± 1.07 (1237)	65.33[A] ± 1.11 (767)
Form				
Bu08	Count	7.14[A] ± 0.07 (1462)	5.79[B] ± 0.09 (1339)	4.55[C] ± 0.09 (1118)
HD08	Ratio	22.56[C] ± 0.16 (1192)	35.46[A] ± 0.27 (1317)	29.74[B] ± 0.28 (1014)
HD09	Ratio	41.67[C] ± 0.21 (1438)	50.24[A] ± 0.35 (1242)	45.56[B] ± 0.35 (1036)
HD10	Ratio	48.50[A] ± 0.25 (1414)	44.21[B] ± 0.26 (1245)	47.76[A] ± 0.36 (774)
NL07	Mm	25.55[B] ± 0.12 (1611)	26.22[A] ± 0.12 (1495)	22.84[C] ± 0.13 (1492)
NL08	Mm	100.42[C] ± 0.36 (1453)	115.96[A] ± 0.56 (1294)	104.92[B] ± 0.62 (1113)
NL09	Mm	31.21[C] ± 0.22 (1440)	54.67[A] ± 0.39 (1248)	49.56[B] ± 0.37 (1050)
Phenology				
BB08	Days since 31/03/08	41.99[B] ± 0.11 (1491)	19.14[C] ± 0.19 (1454)	42.58[A] ± 0.16 (1145)
GC08	Days since 10/09/08	44.61[B] ± 0.53 (1450)	44.75[B] ± 0.56 (1393)	57.61[A] ± 0.35 (1067)
GD08	Days	165.71[C] ± 0.53 (1440)	188.76[A] ± 0.57 (1390)	178.32[B] ± 0.37 (1055)
Survival				
Su10	%	84.88	74.23	61.31
M07	%	4.11	10.83	10.83
M08	%	9.06	9.88	25.03
M09	%	1.64	6.44	6.41
M10	%	1.04	1.27	2.00

325 * *Trait/year code: Letter(s) denote trait, number denotes year of measurement: traits - DA, absolute*
326 *basal stem diameter; DI, basal stem diameter increment; HA, absolute height; HI, height increment;*
327 *HR, relative growth rate; Bu, number of buds; HD, slenderness; NL, needle length; BB, budburst*
328 *timing; GC, growth cessation; GD, growth duration; Su, survival; M, mortality; years - 07, 2007; 08,*
329 *2008; 09, 2009; 10, 2010.*

Trait means were significantly different among the three nurseries for each trait and
each year with the exception of height increment in 2009 and relative growth rate in

2010, both among NG and NW, and growth cessation among NE and NG (Table 1).
The length of needles was considerably longer in all nurseries in 2008 than in either
2007 or 2009 (Figure S3, Table 1). Needles which grew in 2008 were, on average,
3.93 (NE) to 4.59 (NW) times longer than those which grew in 2007, and 2.11 (NW)
to 3.21 (NE) times longer than needles which grew in 2009. Overall survival was
lower for trees in NW compared to the other two nurseries (Table 1) although this
was due to a spike in mortality in 2008 which was thought to be largely due to biotic
factors such as sand fleas, snails and fungal infection (although cause of mortality in
each case was not recorded). Mortality was highest/joint-highest at NW in three of
the four assessment years and was also high in the other year. Mortality at the other
outdoor site (NE) was the lowest of all nurseries across all four years. By 2010, there
were significant differences among nurseries in seedling survival. Mortality was
highest in 2008 at NW (25.03 % annual mortality) which accounted for over half of
the seedling deaths across the period for this nursery. By 2010, mortality was low for
all nurseries.

Trees that died were those which tended, in the previous year, to be shorter and
thinner, formed fewer buds and shorter needles, burst bud and stopped growing later
than trees which remained alive (Figure 2). Exceptions to this were found in only six
out of 57 comparisons (Figure 2): growth cessation in 2008 for trees growing in NG,
some growth traits (absolute and increment height and relative growth rate) for trees
growing in NW in 2008 and needle length in 2009 for trees growing in NW and NE.

Figure 2. Mean trait values for trees that were either alive or dead by the end of the
year after the trait was measured, relative to one another. Where trait means were
identical for each group, relative means are each 0.5 (indicated by dashed line).
Where means are higher for trees which were alive in the following year, the
proportion of the bar that is green exceeds 0.5. Numbers of trees in each group are
indicated within the respective coloured section of each bar. Trait codes are given in
Table 1.

The relative difference in trait values between trees that were alive and those that
died increased as trees got older (Figure 2), although differences over more than
one year cannot be compared for traits relating to phenology). The numbers of trees
in the 'dead' group also reduced over this period, from an average of 221 trees in
2007 to 64 trees in 2008 and 10 trees in 2009 compared to those in the 'alive' group
(average numbers of trees counted for each trait: 1,313 in 2007; 1,198 in 2008;
1,191 in 2009).

Associations between nursery climate and trait variation

Timing of phenological traits measured in 2008 was significantly different among
nurseries (Table 1): budburst was, on average, 23 days earlier at NG than at either
NE or NW; BB_{50} was 38.7 and 38.8 days in NE and NW, respectively, but only 16
days in NG (Figure 3). Over the same period, the difference in GDD among
nurseries was much lower (Figure 3, NE: 207.3; NG: 237.7; NW: 238.1). Trees at NE
required fewer cumulative GDD, and those at NW and NG required more cumulative
GDD to complete budburst throughout the assessment period but only by around 30
degree days. Trees at NW were exposed to two chilling days during the budburst
period (5th and 6th April 2008) but the temperature environments were otherwise
similar at the two outdoor nurseries (NE and NW, Figure 1).

Figure 3. Cumulative percentage of trees reaching budburst, expressed as days
since 31 March 2008 (left) and cumulative GDD since 1 January 2008 (right). Circles
indicate assessment days. The dotted line indicates the day when 50% of trees had
burst bud (BB_{50}).

Despite significantly earlier initiation of budburst in NG, trees ceased growing at
around the same time as those in NE (Table 1) which was on average about 13 days

earlier than in NW. The duration of growth (period between budburst and growth
cessation) therefore varied on average between 166 and 189 days (NE and NG,
respectively) with trees in NW actively growing for an average of 178 days each
year. Mean growth duration was highly significantly positively associated with
growing season length (Figure 4a) although the amount of variation in this trait at
each site was substantial.

392
Figure 4. Linear regressions between growing season length (the period in days
bounded by a daily mean temperature >5 °C for >5 consecutive days and daily mean
temperature <5 °C for >5 consecutive days, after 1st July) and a) growth duration in
2008; and b) relative growth rate 2008 to 2010. Regressions were performed on all
data points. Coloured points are trait values for individual seedlings (jittered for
clarity). Black points indicate trait mean for each nursery. Error bars are one
standard deviation either side of the mean. Mean growth duration: number of days
between budburst and growth cessation. Mean relative growth rate: mm per year as
a proportion of its size at the end of the previous growing season.

There was also a significant positive relationship between mean relative growth rate
and the mean growing season length at each site between 2008 and 2010 (Figure
4b). In-season performance in individual years may be strongly associated with the
length of growing season in both the current and the previous year, due to
provisioning of the bud during the growing season, but these comparisons will be
more powerful using field rather than nursery data, due to the comparatively small
number of datapoints available for the latter.

Relationships within and among traits

Age-age phenotypic correlations were performed for measurements of a given trait
obtained in pairs of years for all trees in each nursery site separately (Figure 5). The
majority of age-age phenotypic correlations were significant across each pair of
years in each of the nurseries, with the exception of needle length (between years
2007 and 2008 in NG; between years 2008 and 2009 in NE), basal stem diameter

increment (between years 2009 and 2010 in NG), and height increment (between
years 2007 and 2009, and 2007 and 2010 both in NG). Traits were consistently
either positively or negatively correlated among years, with the exception of height
increment and needle length. Phenotypic correlations among height increments in
different years for individuals in NG were negative when 2010 is included (in contrast
to the same years in other nurseries for this trait) – this is the year during which the
pots were moved outdoors from the glasshouse. The patterns of pairwise
comparisons among the sites were more similar for the outdoor sites (NE and NW)
than NG. Correlations generally increased in significance as the age of the trees
increased and as the number of years between comparisons decreased (Figure 5).

Figure 5. Correlation coefficient (r) and associated significance values (p) for a given
trait among years at each nursery (NE, NG, NW). On x-axis, Year is shortened,
whereby a pairwise comparison of a trait between years 2007 and 2010 is presented
as '07-10'. Trait codes are detailed in Table 1. Height measured in the first year
(HI07) has been included in both absolute height (HA) and height increment (HI)
comparisons

For simplicity, the number of traits measured across multiple years that were
consistently significantly associated (in the same direction) among years were
reduced prior to subsequent analyses, either to a single (most recent) assessment
year (for traits for which successive measurements were not independent: DA, HA)
or to a mean value among years (HR, DI). Traits which were not consistent among
years (HI, NL) were considered separately for each year.

Pairwise comparisons among growth, form and phenology traits both within and
among years (Figure 6) show that pairs of traits were more highly significantly
correlated in NW than in either NG or NE, particularly among growth traits and
among budburst and other traits. Relationships were often inconsistent among sites:
of the 105 pairwise comparisons among traits, 21 were not consistent (i.e. not
always either positive or negative, possibly due to the confounding effects of
genotype × environment interactions) among nurseries. Of the inconsistent
correlations, nearly all (N = 17) involved needle length or height increment (Figure 6).
Significantly associated growth traits were always positive, with the exception of
height increment in 2007 in NE and NW and in all years at NG (Figures 6).

Phenology traits were highly significantly positively correlated with one another and
were generally negatively correlated with traits related to form (with the exception of
slenderness and needle length in 2009). Considering only traits that were measured
in the same year (i.e. 2008 for phenology traits), days to budburst were negatively
correlated with height increment in NW, positively correlated in NE and not
associated in NG. Days to budburst were highly significantly negatively correlated
with all growth traits at NW, whereas correlations were either not significant or
positively significant for growth traits in NG and NE. In contrast, days to cessation of
growth were highly positively correlated with most growth traits in NG (with the
exception of height increment in 2007) but associations were much weaker and/or
negatively correlated in the other nurseries. Thus, in general, a long growing season
between budburst and growth cessation resulted in larger plants (Figure 6).

Figure 6. Pairwise correlations for traits for Scots pine trees growing in each nursery
(NE, NG, NW); colour indicates value of Pearson correlation coefficient. Significance
values: *, p 0.01-0.05; **, p 0.001-0.01; ***, p < 0.001.

For further analyses, traits within groups which were consistently highly significantly
positively correlated with one another were reduced to a single representative trait
(height increment 2008 to 2010, relative growth rate and absolute and increment
basal stem diameter were all represented by absolute height in 2010; needle length
in 2007 and 2008 were represented by number of buds; growth cessation was
represented by budburst)

Partitioning trait variance

There were highly significant differences among nurseries (i.e. environmental
variation over a large scale) as well among blocks within nurseries (environmental
variation over a small scale) for every trait (Table S1). Environmental variation (block
and nursery) accounted for more than half the total variance for budburst and needle
length, close to half for absolute height and around one quarter for the remaining
traits (slenderness, number of buds and height increment). Nursery accounted for a
greater proportion of variation than block for needle length, number of buds, height
increment and budburst (Figure 7), whereas the contribution of block to the total
variance was greater than nursery for slenderness and absolute height.

Figure 7. Proportion of variance in growth, form and phenology traits explained by
mixed model factors: environmental factors are coloured in shades of green (nursery
and block) and genetic/residual factors are coloured in shades of grey
(nursery*provenance, provenance, family, residual).
485
Discussion

This study highlights the profound impact that nursery environment can have on trait
variation in tree seedlings. The nurseries were exposed to the same nutrient, pot and
watering regimes, came from the same genetic background and experienced similar
average temperatures and monthly growing degree days. Differences in recorded
temperature among the nurseries occurred during the daytime within the growing
season, when daily temperature variance was much higher for trees growing in the
nursery in the east of Scotland compared to the nursery in the west. Seedlings in the
glasshouse nursery, in contrast, experienced high daily temperature variance (which
was also higher than in the neighbouring outdoor nursery), but temperatures were
also higher overall. The importance of the effect of warmer mean temperatures
(Morison and Morecroft 2006) and extremes of temperature (Niu *et al.* 2014) on plant
traits is increasingly well recognised, especially in the light of an anticipated increase
in extreme weather events due to climate change (Herring *et al.* 2022). However, the
contrasting effects of temperature variance and mean temperatures and their
comparative impact on the growth and development of plants and trees, is less well
understood.

In our study, all traits were strongly and significantly affected by the nursery
environment: both at a large spatial scale (i.e. among nurseries) but also (and
sometimes to a greater extent) at much finer spatial scales (i.e. among blocks within
nurseries). The large spatial effect was particularly pronounced between the
glasshouse and neighbouring outdoor nursery, highlighting the major impact that
nursery type (glasshouse vs outdoor nursery) has on growing plants. While the main
environmental difference among the nurseries adjacent (but of different types) to one
another was mean temperature, we cannot exclude the possibility that other abiotic
(e.g. wind speed, humidity) and/or biotic (e.g. symbionts, pests, pathogens) factors
also played a part in the differences we observed.

Growing the seedlings under high temperature variance (resulting in more days at
the extreme ends of the temperature range) but with the same mean temperature
may lead to lower rates of growth. Chiang, Bånkestad and Hoch (2020) reported
significantly taller seedlings when grown in fixed or sinusoidal temperature regimes
compared to the same species grown in pots outdoors with a naturally fluctuating
and high temperature variance, but only in spring. When grown in the summer, the
same seedlings grown in a fixed environment were reported to be significantly
shorter than those grown in the outdoor environment, possibly because the lower
temperatures that seedlings were exposed to in the outdoor environment in the

spring did not occur during summer. The extent of the difference between day time
and night time temperatures is also known to directly influence key plant traits
relating to growth and form, such as internode length and height (Myster and Moe
1995). Similarly, the allocation of resources to traits relating to tree form in this study
appeared to be more important in the nursery with both high variance and low mean
temperatures, indicating that a more conservative growth strategy was potentially
balanced by investment in traits relating to tree form (Climent *et al.* 2024), such as
number of buds.

The role of mean temperature on plant growth is better understood, with higher
mean temperatures generally positively associated with higher rates of growth
(measured as specific leaf area (Poorter *et al.* 2010). Our study supported this
finding, with seedlings grown in a nursery with warmer mean temperatures showing
higher rates of growth. Although the temperatures of our nurseries were not
controlled, and only a few nursery environments were studied, the environments are
analogous to those used in the forestry industry (i.e. outdoor nurseries and
glasshouses). We only recorded air temperature, but soil temperature may also have
been an important factor for the growth and development of the seedlings. The latter
is unlikely to have been consistent among the glasshouse and outdoor
environments: it has been reported that soil temperature inside pots tracks but lags
behind the air temperature until solar radiation falls directly onto pots, when the
temperature of the soil within the pots can rise to more than 20 °C higher than the air
temperature (Poorter *et al.* 2016) and that increased soil temperature is associated
with increased growth rate (Weih and Karlsson 1999) in saplings.

Growing Scots pine seedlings in nursery environments with different temperature
means, variances and photoperiods also had strong effects on phenological
variation. Budburst timing was strongly associated with growing degree days at all
sites but was not fully explained by differences in this environmental variable among
sites. This suggests a complex relationship between warm temperatures (i.e.
growing degree days) and cold temperatures (often referred to as the chilling
requirement), although the number of nurseries/years assessed in this study is
insufficient to enable firm conclusions to be reached on how these two variables
interact to result in the observed phenotype. Plant species generally cease growing
in response to decreasing photoperiod (Nitsch 1957; Singh *et al.* 2017) although
some may respond to decreasing temperature instead (Heide and Prestrud 2005). In
our study, trees growing in neighbouring nurseries, outside or within a glasshouse,
experienced the same photoperiod (although the light intensity through the glass
may have been different) but different mean temperatures. Whereas budburst
occurred much earlier in the glasshouse than in the neighbouring outdoor nursery
(reflecting the different cumulative growing degree days over the preceding period),
growth cessation occurred on average at a similar time which suggests photoperiod
to be the main driver.

The accuracy of using young seedlings to predict economic traits at harvest is of
major interest and importance in the genetic testing of progeny for forestry (Lee
2002; Hong, Fries and Wu 2015), but detailed early and late assessments of the
same tree seedlings are rarely available. Within these experimental trials, which
have since been transplanted to field sites in 2012, the same traits will continue to be
measured into maturity enabling age-age correlations from seedling to mature trees
to be estimated with high resolution (many traits will be measured annually).
Although the assessments presented here were confined to the early years of growth
this study nevertheless highlights the relative consistency of age-age correlations
among the different nursery environments, with the exception of height increment in
NG in the years before and after trees were moved outside. The latter showed that
plants experienced a shock as a result of the move outdoors which impacted their
subsequent growth. The well-documented phenomenon of 'transplantation shock' is
known to induce changes to growth and development (Close, Beadle and Brown
2005) whereby growth is impaired for a period following transfer. The accuracy of
age-age correlations should also, therefore, be reviewed in the context of the general
environment in which the trees are measured and the specific environment the trees
have been exposed to in the recent past.

The interactions among different traits were also affected by the nursery that plants
were grown in. For example, while timing (i.e. lateness) of growth cessation was
positively correlated with most growth traits in the protected glasshouse
environment, these relationships were weaker in outdoor environments, possibly
because these traits were not responding to a single climatic variable but to a
combination of many. Similarly, traits relating to form were inconsistent in their
relationship with growth traits among nurseries, with more positive associations
observed in the nurseries with higher temperature variance compared to the one with
low temperature variance.

Given the crucially important role of nurseries in the supply chain for forestry and
tree planting, the success of global tree planting efforts depends strongly on the
quality of plants they produce. Here we have shown the vital importance of properly
understanding the strength and persistence of nursery environment effects on tree
seedling growth and development. This knowledge has implications both for genetic
testing of material and also for operational tree production. For example, if analyses
do not also incorporate data relating to the early environment of the tested trees, an
important component of variation might be omitted and may lead to over- or under-
estimation of breeding values. Indeed, environmental variation over both small
(within sites, measured using blocks) and large (among sites) scales can have
significant effects on trait variation and should be carefully recorded and used as
covariates in subsequent analyses. For production forestry, although best practice
may be to target specific nurseries for the growth of particular species and/or for
planting in particular environments (Jaenicke, 1999), economic and practical
constraints often render practitioner choice of nursery environment relatively limited.

In this case, improved knowledge of the potential outcomes associated with nursery environment could help practitioners to better understand field traits as an outcome of the environment their trees experienced in the nursery. To this end we recommend routinely monitoring and making available nursery environmental data alongside plant provenance data on shipment from the nursery. The significant effect of nursery environment on all aspects of phenotype reported in this study highlights the cumulative and divergent effect that the early growing environment can have, an effect that may have particularly profound and persistent effects on long-lived species like trees. These findings will be investigated and tested further for the subset of trees that were transplanted to field locations in 2012, using measurements made over many subsequent years.

Conclusions

Despite the relatively low levels of environmental variation recorded among the three nurseries, there were significant effects on seedling mortality, variation in traits relating to growth, form and phenology and on interactions among traits. These findings have implications for the choice of nursery in which plants are raised given that effects may persist and affect subsequent performance in the field. Growers may wish to assess whether plants should be preferably raised in less benign nursery environments in order to minimise the shock they receive when transplanted. In any case, we are confident of the importance of measuring and collecting data at an early stage in order to compare trait variation throughout the lifetime of trees, from seedling through to maturity: an approach which would be of benefit for plants grown for both research and for operational purposes. The trees in this study have now been transplanted to three field environments (Beaton *et al.* 2022) and subsequent analysis explores the persistence of early environment carry over effects following transplantation (Perry et al., 2024b). Regular measurements on the trees since transplantation will also enable ontogenetic effects on intraspecific genetic variation to be characterised for a wide range of phenotypic traits and for age-age correlations from seedling to maturity to be estimated.

Funding

This work was supported by the newLEAF project (NE/V019813/1) under the UK's 'Future of UK Treescapes' programme, which was led by UKRI-NERC, with joint funding from UKRI-AHRC & UKRI-ESRC, and contributions from Defra and the Welsh and Scottish governments.

This work was also part of PROTREE, a project supported by grant BB/L012243/1, funded jointly by UKRI's BBSRC, NERC, & ESRC and Defra, the Forestry Commission and the Scottish Government, under the Tree Health and Plant Biosecurity Initiative

The James Hutton Institute contribution to this work was supported by Scottish
Government's Rural & Environment Science & Analytical Services Division, through
their Strategic Research Programmes (2006–2011, 2011–2016 and 2016–2022).

Acknowledgements

We thank Stewart Mackie and staff at Forest and Land Scotland (Selkirk Office), the
Forestry Commission, National Trust for Scotland, and Donald Barrie (Glensaugh)
for their help in locating, establishing and securing access to sites.

Conflict of interest statement

None declared

Data availability statement

The data underlying this article are available from the Environmental Information
Data Centre (EIDC, https://eidc.ac.uk), at https://doi.org/10.5285/29ced467-8e03-
4132-83b9-dc2aa50537cd. Temperature and temperature variances at each nursery
are available at https://doi.org/10.5285/81841d93-41e2-47a7-b15a-92d1e1cf07f7.

References

Aldhous JR, Mason WL. Forest Nursery Practice. Forest Research, Farnham,
1994:296.

Beaton J, Perry A, Cottrell J *et al.* Phenotypic trait variation in a long-term multisite
common garden experiment of Scots pine in Scotland. *Sci Data* 2022;**9**:671.

Benavides R, Carvalho B, Matesanz S *et al.* Phenotypes of *Pinus sylvestris* are more
coordinated under local harsher conditions across Europe. *J Ecol* 2021, DOI:
10.1111/1365-2745.13668.

Brancalion PHS, Holl KD. Guidance for successful tree planting initiatives. *J Appl*
Ecol 2020, DOI: 10.1111/1365-2664.13725.

Burdon RD, Bannister MH, Low CB. Genetic survey of Pinus radiata. 5: Between-
trait and age-age correlations for growth rate, morphology, and disease
resistance. *New Zealand Journal of Forestry Science* 1992;**22**:211–27.

Chiang C, Bånkestad D, Hoch G. Reaching natural growth: the significance of light
and temperature fluctuations in plant performance in indoor growth facilities.
Plants 2020;**9**, DOI: 10.3390/plants9101312.

Close DC, Beadle CL, Brown PH. The physiological basis of containerised tree
seedling 'transplant shock': a review. *Australian Forestry* 2005;**68**:112–20.

Climent, J., Alía, R., Karkkainen, K. *et al*. Trade-offs and Trait Integration in Tree
Phenotypes: Consequences for the Sustainable Use of Genetic Resources.
Current Forestry Reports 2024;**10**:196–222.

Dhar A, Balliet N, Hawkins CD *et al.* Bud flush phenology and nursery carryover
effect of paper birch provenances. *iForest* 2015;**8**:809–17.

Domisch T, Finér L, Lehto T. Effects of soil temperature on biomass and
carbohydrate allocation in Scots pine (Pinus sylvestris) seedlings at the
beginning of the growing season. *Tree Physiol* 2001;**21**:465–72.

Dong M, Fan Y, Wu Z *et al.* Age–age correlations and early selection for growth
traits in 40 half-sib families of Larix principis-rupprechtii. *Journal of Forestry*
Research 2018:1–7.

Donnelly K, Cavers S, Cottrell JE *et al.* Cryptic genetic variation and adaptation to
waterlogging in Caledonian Scots pine, Pinus sylvestris L. *Ecol Evol*
2018;**8**:8665–75.

Downs RJ, Borthwick HA. Effects of photoperiod on growth of trees. *Botanical*
Gazette 1956;**117**:310–26.

Falconer DS, Mackay TFC. *Introduction to Quantitative Genetics*. 4th ed. Essex, UK:
Longman, 1996.

Forestry Commission. *Tree Supply Report*. UK Government, 2023.

Forest Research. *The UK Forestry Standard*. Forest Research, Farnham, 2023:139.

Fukatsu E, Tamura A, Takahashi M *et al.* Efficiency of the indirect selection and the
evaluation of the genotype by environment interaction using Pilodyn for the
genetic improvement of wood density in *Cryptomeria japonica*. *J For Res*
2011;**16**:128–35.

Gebhardt‐Henrich SG, Van Noordwijk AJ. Nestling growth in the Great Tit I.
Heritability estimates under different environmental conditions. *J Evol Biol*
1991;**4**:341–62.

Gruffman L, Ishida T, Nordin A *et al.* Cultivation of Norway spruce and Scots pine on
organic nitrogen improves seedling morphology and field performance. *Forest*
Ecology and Management 2012;**276**:118–24.

Harrell Jr FE. *Hmisc: Harrell Miscellaneous*. R package, 2024.

Heide OM, Prestrud AK. Low temperature, but not photoperiod, controls growth
cessation and dormancy induction and release in apple and pear. *Tree Physiol*
2005;**25**:109–14.

Heiskanen J, Rikala R. Effect of Peat-based Container Media on Establishment of
Scots Pine, Norway Spruce and Silver Birch Seedlings after Transplanting in
Contrasting Water Conditions. *Scandinavian Journal of Forest Research*
2000;**15**:49–57.

Herbert R, Samuel S, Patterson G. *Using Native Stock for Planting Native Trees and*
Shrubs. Forestry Commission, 1999.

Herring SC, Christidis N, Hoell A *et al.* Explaining Extreme Events of 2020 from a
Climate Perspective. *Bull Amer Meteor Soc* 2022;**103**:S1–129.

Hoffmann AA, Merilä J. Heritable variation and evolution under favourable and

unfavourable conditions. *Trends Ecol Evol* 1999;**14**:96–101.

Holl KD, Brancalion PHS. Tree planting is not a simple solution. *Science*
2020;**368**:580–1.

Hong Z, Fries A, Wu HX. Age trend of heritability, genetic correlation, and efficiency
of early selection for wood quality traits in Scots pine. *Can J For Res*
2015;**45**:817–25.

Jaenicke H. Good tree nursery practices: practical guidelines for research nurseries.
World Agroforestry Centre; 1999.

Lee SJ. Selection of parents for the Scots pine breeding population in Britain.
Forestry 2002;**75**:293–303.

Luoranen J. Autumn versus spring planting: the initiation of root growth and
subsequent field performance of Scots pine and Norway spruce seedlings. *Silva*
Fenn 2018;**52**, DOI: 10.14214/sf.7813.

Marron N, Dillen SY, Ceulemans R. Evaluation of leaf traits for indirect selection of
high yielding poplar hybrids. *Environ Exp Bot* 2007;**61**:103–16.

Mason WL, Alía R. Current and future status of Scots pine (Pinus sylvestris L.)
forests in Europe. *For Syst* 2000;**9**:317–35.

Minitab. *Minitab Statistical Software*. Minitab, 2024.

Morison JIL, Morecroft MD eds. *Plant Growth and Climate Change*. Oxford, UK:
Blackwell Publishing Ltd, 2006.

Myster J, Moe R. Effect of diurnal temperature alternations on plant morphology in
some greenhouse crops—a mini review. *Sci Hortic* 1995;**62**:205–15.

Nerg A, Kainulainen P, Vuorinen M *et al.* Seasonal and geographical variation of
terpenes, resin acids and total phenolics in nursery grown seedlings of Scots
pine (*Pinus sylvestris* L.). *New Phytologist* 1994;**128**:703–13.

Nitsch JP. Growth response of woody plants to photoperiodic stimuli. *Proceedings*
American Society for Horticultural Science 1957;**10**:512–25.

Niu S, Luo Y, Li D *et al.* Plant growth and mortality under climatic extremes: An
overview. *Environ Exp Bot* 2014;**98**:13–9.

Osorio LF, White TL, Huber DA. Age-age and trait-trait correlations for Eucalyptus
grandis Hill ex Maiden and their implications for optimal selection age and
design of clonal trials. *Theor Appl Genet* 2003;**106**:735–43.

Pan Y, Birdsey RA, Fang J *et al.* A large and persistent carbon sink in the world's
forests. *Science* 2011;**333**:988–93.

Perry A, Brown AV, Cavers S *et al.* Has Scots pine (*Pinus sylvestris*) co-evolved
with *Dothistroma septosporum* in Scotland? Evidence for spatial heterogeneity
in the susceptibility of native provenances. *Evol Appl* 2016;**9**:982–93.

Perry, A.; Beaton, J. K.; Stockan, J. A. et al. Long-term multisite Scots pine trial,
Scotland: temperatures and temperature variances for tree nurseries, 2007-
2012. NERC EDS Environmental Information Data Centre. 2024a.
https://doi.org/10.5285/81841d93-41e2-47a7-b15a-92d1e1cf07f7

Perry A., Beaton, J. K., Stockan, J. A, et al., Significant and persistent carryover
effects in Scots pine. bioRxiv [preprint]. 2024b.
https://doi.org/10.1101/2024.08.23.609343

Poorter H, Fiorani F, Pieruschka R *et al.* Pampered inside, pestered outside?
Differences and similarities between plants growing in controlled conditions and
in the field. *New Phytol* 2016;**212**:838–55.

Poorter H, Niinemets U, Walter A *et al.* A method to construct dose-response curves
for a wide range of environmental factors and plant traits by means of a meta-
analysis of phenotypic data. *J Exp Bot* 2010;**61**:2043–55.

R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R Foundation for Statistical Computing, 2024.

Ramírez-Valiente JA, Solé-Medina A, Pyhäjärvi T *et al.* Selection patterns on early-
life phenotypic traits in Pinus sylvestris are associated with precipitation and
temperature along a climatic gradient in Europe. *New Phytol* 2021;**229**:3009–25.

Salmela MJ, Cavers S, Cottrell JE *et al.* Seasonal patterns of photochemical
capacity and spring phenology reveal genetic differentiation among native Scots
pine (Pinus sylvestris L.) populations in Scotland. *Forest Ecology and*
Management 2011;**262**:1020–9.

Selander J, Immonen A. Effect of fertilization and watering of Scots pine seedlings
on the feeding preference of the pine weevil (Hylobius abietis L.). *Silva Fenn*
1992;**26**, DOI: 10.14214/sf.a15637.

Singh RK, Svystun T, AlDahmash B *et al.* Photoperiod- and temperature-mediated
control of phenology in trees - a molecular perspective. *New Phytol*
2017;**213**:511–24.

Vapaavuori EM, Rikala R, Ryyppö A. Effects of root temperature on growth and
photosynthesis in conifer seedlings during shoot elongation. *Tree Physiol*
1992;**10**:217–30.

Villar-Salvador P, Ocaña L, Peñuelas J *et al.* Effect of water stress conditioning on
the water relations, root growth capacity, and the nitrogen and non-structural
carbohydrate concentration of Pinus halepensis Mill. (Aleppo pine) seedlings.
Ann For Sci 1999;**56**:459–65.

Visscher PM. On the sampling variance of interclass correlations and genetic
correlations . *Genetics* 1998;**149**:1605–14.

Weih M, Karlsson PS. The nitrogen economy of mountain birch seedlings:
implications for winter survival. *Journal of Ecology* 1999;**87**:211–9.

Westaway S, Grange I, Smith J *et al.* Meeting tree planting targets on the UK's path
to net-zero: A review of lessons learnt from 100 years of land use policies. *Land*
use policy 2023;**125**:106502.

Williams CG. Accelerated short-term genetic testing for loblolly pine families. *Can J*
For Res 1988;**18**:1085–9.

Wu HX, Yeh FC, Dhir NK *et al.* Genotype by environment interaction and genetic
correlation of greenhouse and field performance in *Pinus contorta* ssp. *latifolia.*
Silvae genetica 1997;**46**.

Zobel BJ, Talbert JT. *Applied Tree Improvement.* New York: John Wiley & Sons,
1984.