Tree nursery environments and their effect on early trait variation

Annika Perry[1*], Joan K. Beaton[2], Jenni A. Stockan[2], Glenn R. Iason[2], Joan E.

Cottrell[3] and Stephen Cavers[1]

[1]*UK Centre for Ecology & Hydrology (UKCEH), Bush Estate, Midlothian, EH26 0QB,*

UK

[2]*Ecological Sciences, James Hutton Institute, Craigiebuckler, Aberdeen, AB15 8QH,*

UK

[3]*Northern Research Station, Forest Research, Roslin, EH25 9SY, UK*

Corresponding author: annt@ceh.ac.uk

Abstract

Despite the necessity of using nurseries to raise young plants and trees prior to

transplantation into the landscape, not enough is known about the impact of their

associated climate on the growth and development of seedlings from initial

germination through to transplantation and beyond. Furthermore, the effect of

nursery environment on trait interactions and age-age correlations and their

subsequent relation to traits in mature trees is of significant importance and interest

to forestry practitioners but requires long-term experimental trials measured for

multiple traits in multiple years. A multisite progeny-provenance trial of the

economically and ecologically important species, Scots pine, was established in

three environmentally distinct nurseries in Scotland and measured for traits relating

to survival, growth, form and phenology while in the nurseries. Temperature variation

and photoperiod were the only uncontrolled environmental variables during this

period, and their effect on measured traits was found to be significant among

nurseries from the first growing season onwards. Furthermore, interactions among

traits were not always consistent among the different nurseries indicating that the

use of proxy traits to enable selection of desirable traits may be more or less

successful depending on the environment the trees are measured in. This study

represents the first in a series examining trait variation in Scots pine from seedlings

to mature trees, and highlights the importance of carefully considering the nursery

environment when growing trees for transplantation.

Introduction

Globally, forests function as the major terrestrial carbon sink (Pan *et al.* 2011) and

there is broad consensus that increasing the area of forest cover is vital to achieving

net zero targets (Westaway *et al.* 2023). If it happens, a greater level of forest cover

would also help tackle parallel environmental challenges like biodiversity loss, soil

stabilisation and watershed management (Holl and Brancalion 2020). The degree of

scaling up proposed is large, and has been exemplified in recent years by a

proliferation of tree-planting initiatives such as the Trillion Trees Campaign:

trilliontreecampaign.org, the Eden reforestation projects: edenprojects.org, and the

Bonn Challenge www.bonnchallenge.org. This is creating an increased need for

planting stock, which is most commonly supplied as nursery-grown saplings

(Brancalion and Holl 2020) and places the forest nursery in a pivotal position, even

as they continue to be required to supply the vast numbers of plants required for

traditional forestry, amenity and conservation planting. In Great Britain alone, it is

estimated that forest nurseries produced around 173M saplings for the year 2022-

2023 (Forestry Commission 2023). Therefore, now and in the future, nurseries will

play a major role in the supply chain for forestry and tree planting of all kinds and will

be key to the success of global tree planting efforts.

In their role as the primary location for raising young trees, choices made by

nurseries have been shown to profoundly influence size and development of young

trees, effects that may be observable long after transplantation. For example, the

choice of fertiliser (Gruffman *et al.* 2012), growing medium (Heiskanen and Rikala

2000), container type, root management, and watering regime (Villar-Salvador *et al.*

1999) during early life stages can affect root growth, seedling morphology, and

susceptibility to pests and pathogens (Selander and Immonen 1992). Even the

timing of transplantation from the nursery to the field can impact performance

(Luoranen 2018). Whilst soil, water and nutrients can be controlled quite precisely by

nursery managers, the geographic location of a nursery and any protection

measures used, such as glasshouse cover, determines other critical components of

its environment such as climate and photoperiod . As variables like temperature and

photoperiod can strongly affect the growth, development and biochemistry of

seedlings (Downs and Borthwick 1956; Vapaavuori, Rikala and Ryyppö 1992; Nerg

et al. 1994; Domisch, Finér and Lehto 2001) with effects lasting well beyond

transplantation(Dhar *et al.* 2015), the nursery environment can be hugely influential.

The use of locally sourced seed is an explicit requirement when planting native

species in the UK (Herbert, Samuel and Patterson 1999), but the location of the

nursery in which those plants are raised is rarely specified. For example, the

contemporary guiding document for UK forestry – The UK Forestry Standard (Forest

Research 2023) - does not comment on nursery management, nor connect nursery

practice to its recommendations on seed sourcing. Previous publications, such as

the Forestry Nursery Practice (Aldhous and Mason 1994), do consider site selection

factors, but largely from the point of view of maximising plant performance,

minimising risk and logistical considerations. In practice, choice of nursery is

generally governed by what the market offers and by economics, consequently the

nursery site may be environmentally distant from both the site of origin and planting

site. If the effect of the nursery environment on plant phenotype is large, then the

effect of local seed sourcing may be minimal in comparison. A quantitative

evaluation of the impact of the nursery environment on trait variance is therefore

potentially very important to seed sourcing policy, nursery practice and to the long-

term success of planting initiatives.

The impact of the early growing environment is also a critical factor to consider in

tree improvement and breeding programmes, particularly with regards to its effect on

survival and on variation in desirable traits. Traditional breeding practice, which

makes selections at around half rotation length (Zobel and Talbert 1984), is highly

inefficient for trees with long rotation times and there is much interest in methods for

screening trait variation at the nursery stage (Wu *et al.* 1997). However, trait values

prior to transplantation (ca. two years) rarely predict values in later years (Burdon,

Bannister and Low 1992; Hong, Fries and Wu 2015; Dong *et al.* 2018) (apart from

species with very short rotation lengths, e.g. *Eucalyptus grandis* in Colombia,

(Osorio, White and Huber 2003)). The use of 'proxy' traits to enable early selection

relies on the stability of trait-trait relationships among environments, which may

(Fukatsu *et al.* 2011) or may not be (Marron, Dillen and Ceulemans 2007)

identifiable. Significant variation in age-age and trait-trait correlations among

environments (Williams 1988; Osorio, White and Huber 2003) means that nursery

effects may confound selection based on early traits. Similarly, the issue of

genotype-environment interactions may become increasingly important as larger

numbers of trees are planted and on more challenging sites.

Pinus sylvestris L. (Scots pine) is a globally important tree species with a natural

range that extends from southern Spain to northern Finland, and from western

Scotland to far-eastern Russia. A foundation species with high conservation value, it

has been shown to exhibit high adaptive trait variation across the landscape it

inhabits at both regional and international scales (Salmela *et al.* 2011; Perry *et al.*

2016; Donnelly *et al.* 2018; Benavides *et al.* 2021; Ramírez-Valiente *et al.* 2021).

Pinus sylvestris is also an economically valuable timber production species and is

widely planted both within and beyond its natural range: it is estimated that it

occupies more than 20 percent of the productive forest area in the EU (Mason and

Alía 2000) and, within Great Britain, it is the second most abundantly grown species

in tree nurseries after Sitka spruce (Forestry Commission 2023). *Pinus sylvestris*

grows (both naturally and planted) across a vast and heterogeneous landscape. It

may therefore need to survive and thrive in environments with huge temperature

variance, or in others with relatively stable temperature profiles across seasons and

years.

Common garden trials are widely used in forestry to disentangle the genetic

component of phenotypic variation from the effects of environmental variation.

Extending this approach, but using the common garden design to control the genetic

composition of test populations, it is possible to use multiple common gardens in

different environments to examine the environmental component of phenotypic

variation. Nurseries, in their efforts to minimise environmental variation within a

single site and to standardise conditions for each plant, are effectively functional

common gardens. Here we compare multiple common garden nurseries, each

encapsulating the same genetic variation, to identify the contribution of the nursery

environment to phenotypic variation and estimate genotype by environment effects.

By doing so, we evaluate the importance of the nursery environment to longer term

development of the plant, and establish a baseline for interpreting field trial results in

the years to come. In addition, these results will be important in enabling age-age

correlations to be explored as trees mature in field trials.

For this study, we compared the performance of *P. sylvestris* seedlings in three

nursery environments. The plants had a common genetic background, collected

systematically from across the natural range in Scotland, and were grown under

standard soil, nutrient and watering regimes. We quantified the effect of the nursery

environment on early growth and development and survival and examined

relationships within and among traits.

Methods

Source of material

Seedling source and nursery environments are detailed in Beaton *et al.* (2022).

Briefly, seed was collected from 10 mother trees from each of 21 populations of *P.*

sylvestris from across the native range in Scotland in March 2007. Following

stratification, seeds were germinated at the James Hutton Institute, Aberdeen

(latitude 57.133214, longitude -2.158764) in June 2007 and transplanted into pots in

an unheated glasshouse. The full final collection consisted of 210 families (10

families from each of 21 populations) each consisting of 24 half-sibling progeny (total

5,040 individuals).

Nursery environment

After transfer into individual pots (the same compost and pots were used for all

plants) and with the intention of eventually using these plants to establish field based

provenance trials, 8 seedlings per family were moved to one of three nurseries (total

1,680 seedlings per nursery) in July 2007: an unheated nursery glasshouse (NG) at

the James Hutton Institute, Aberdeen; an outdoor nursery in the west of Scotland

(NW) at Inverewe Gardens, (latitude 57.775714, longitude -5.597181); an outdoor

nursery in the east of Scotland (NE) at the James Hutton Institute (location as

above). In each nursery, trees were arranged in 40 randomised blocks, where each

block contained two trees per family (total 42 trees). Watering was automatic in NG,

and manually, as required, for NE and NW. The watering regime ensured that water

was never limiting and waterlogging of seedlings was avoided. No artificial light was

used in any of the nurseries. In May 2010, the pots containing the seedlings from NG

were moved outdoors to Glensaugh in Aberdeenshire (latitude 56.893567, longitude

-2.535736).

Hourly temperature was recorded at each of the nursery sites using data loggers

from July 2007 until December 2010. Data loggers were suspended 50 cm above the

ground under an aluminium foil-covered funnel. Daily minimum, maximum and

average temperatures (between 0000-2300 each day) and the daily temperature

variances were calculated for each nursery over this period. Daily variance was also

partitioned into separate values for the day and night period, where day variance

was estimated between the hours of sunrise and sunset between 0000-2300 and

night variance was estimated between the hours of sunset and sunrise between

0000-2300. Sunset and sunrise for each day at each nursery were obtained from

www.timeanddate.com.

On the few occasions on which more than one hourly record per 24 hours was

missing, the daily temperatures were replaced with average daily temperature data

derived from the nearest weather stations to the nurseries (obtained from the Met

Office National Meteorological Archive for the period July 2007 to December 2010).

Weather stations were: Poolewe (0.7 miles from NW); Craibstone (4.3 miles from NE

and NG) and Glensaugh No 2 (0.1 miles from NG when moved outdoors to

Glensaugh). Where weather station data were missing (only relevant for Poolewe

over this period) they were replaced with measurements from the next closest

weather station: Aultbea No 2 (4.2 miles from NW). Due to the differences in the

location between the weather stations and the nurseries and the different methods of

recording temperature, data from weather stations were adjusted prior to inclusion in

the 'nursery temperatures' dataset by comparing daily temperatures across the

whole period. Missing logger temperatures were directly replaced, where possible,

by checking the weather station temperature at the missing time point and

calculating the mean logger temperature recorded at all other instances that the

same weather station temperature was recorded. In the few instances where there

were no direct replacements, replacement logger values were obtained by taking the

mean value for the three logger records either side of the missing logger data point

in a dataset which was ordered by the weather station temperatures. For missing

data in NG (prior to May 2010), replacement temperatures derived from NE were

used instead of weather station data. Where hourly data were missing, temperature

variance could not be recorded.

The growing season length in each year at each nursery was estimated as the

period bounded by a daily mean temperature >5 °C for >5 consecutive days and

daily mean temperature <5 °C for >5 consecutive days (after 1st July). During the

defined growing season period trees were assumed to be 'growing' whereas outwith

the defined growing season trees were assumed to be 'not growing'. The previously

described temperature variances were compared between these two different

periods. The growing degrees were estimated for each day in the period July 2007 to

December 2010 as the number of degrees above 5 °C. The cumulative growing

degree days (GDD) were estimated by summing the growing degrees for each

period leading up to phenology assessments at each nursery from 1 January in

2008. Chilling days were defined as the days in which the average maximum

temperature did not exceed 5 °C.

Phenotype assessments

Seedling phenotypes were assessed for a range of traits reflecting different life

history characteristics (Table 1): growth (annual increment in stem diameter and

height; absolute stem diameter and height; relative growth rate); form (current year

needle length, total number of buds; ratio of absolute height to absolute stem

diameter); phenology (timing of budburst and growth cessation); and survival.

Growth traits were assessed annually from 2007-2010. Annual increments in height

and base stem diameter were measured as the increase from the end of one

growing season to the end of the next (for comparisons of height increment across

multiple years, height in the first year of growth in 2007 is considered as the height

increment for this year), whilst absolute growth was measured at the end of the

growing season in 2010. Relative growth rate was estimated as the percentage

increase in height (increment) relative to the absolute height in the previous year.

Mortality was recorded each year from 2007 to 2010 and survival was recorded as

the percentage of surviving trees by 2010. Relative growth rate was estimated as the

percentage increase in height (increment) relative to the absolute height in the

previous year and was recorded for 2008 to 2010.

Form was measured annually from 2007-2009. Needle length was measured for

three needles per tree and a mean value was obtained. The total number of buds on

each seedling was counted in 2008 and 2009. The ratios of absolute height to

absolute stem diameter were recorded in 2010.

Phenology was assessed weekly during the spring and autumn of 2008. Budburst

was defined as the number of days from 31 March 2008 to the time when newly

emerged green needles were observed. The estimated proportion of trees which had

burst bud in each nursery was calculated at each assessment. In order to compare

budburst progression across the three sites, the point at which an estimated 50 % of

trees had undergone budburst (BB_{50}) was calculated as follows:

$$BB_{50} = (50 - B_{i<50}) - \frac{B_{i>50} - B_{i<50}}{A_{i>50} - A_{i<50}} + A_{i<50}$$

where $i<50$ is the ith assessment performed at each nursery where < 50 % of trees

had undergone budburst ($i<50$: NE = 2; NG = 3; NW = 2); $i>50$ is the ith assessment

where > 50 % of trees had undergone budburst ($i>50$: NE = 3; NG = 4; NW = 3); A is

the variable associated with each assessment performed at each nursery (A: number

of days after 31 March; or cumulative growing degree days since 1 January); and B

is the percentage of trees observed with budburst at each nursery at each

assessment.

Growth cessation was defined as the number of days from 10 September 2008 to

the time when no further height growth was observed. In order to compare the

growing season length at each nursery with the duration of growth observed in the

seedlings, 'growth duration' was calculated for each seedling as the number of days

between budburst and growth cessation.

Table 1. Traits measured in individual seedlings, grouped into 'Growth', 'Form',

'Phenology' and 'Survival'. Units for each trait are indicated, along with the years in

which data were collected.

Trait	Code	Unit	Years
Growth			
Stem diameter - absolute	DA	mm	2010
Stem diameter - increment	DI	mm	2009-2010
Height - absolute	HA	mm	2010
Height - increment	HI	mm	2007-2010
Relative growth rate	HR	%	2008-2010
Form			
Number of buds	Bu	Count	2008-2009
Height to stem diameter ratio	HD	Ratio	2010
Mean needle length	NL	mm	2007-2009
Phenology			
Budburst	BB	Days	2008
Growth cessation	GC	Days	2008
Growth duration	GD	Days	2008
Survival			
Survival	Su	%	2010
Mortality	M	0/1	2007-2010

Statistical analysis

Paired t tests were performed using R (R Core Team 2024) to compare nursery

means for annual climatic variables (mean daily temperature, daily variance, growing

degree days) and climatic variables measured within and outside the growing

season (day temperature variance and night temperature variance).

Traits and climatic variables were normalised using the minimum and maximum

value recorded for that trait/climatic variable for each nursery as follows:

$$x_{normalised} = \frac{x - \min(x)}{\max(x) - \min(x)}$$

Coefficient of variation, calculated as standard deviation divided by the mean, was

estimated for each trait in each nursery. Where traits were measured in multiple

years only the most recent year was used.

To evaluate relationships among traits, Pearson's correlation coefficients and

significance values were estimated for trees growing in each nursery using the

'Hmisc' package (Harrell Jr 2024) in R (R Core Team 2024).

Nested analyses of variance (ANOVA) were performed in Minitab version 21 (Minitab

2024) for all trees with nursery as a fixed effect and block nested within nursery as a

random effect. Nested ANOVA was also used to analyse the distribution of variance

among these traits. Tukey's post-hoc tests were performed to determine whether

mean values among nurseries were significantly different from one another for all

traits and for mean temperature, temperature variance (daily, day and night) and

growing degree days for each year. ANOVAs were repeated with all terms included

as random effects to identify their contribution to the total variance.

Narrow sense heritability (h^2), which is the proportion of total phenotypic variance

(V_P) explained by additive genetic effects (V_A) (Falconer and Mackay 1996), was

estimated using among family, block and residual variance (V_{fam}, V_{block} and V_{res}

respectively) from data pooled across populations and analysed separately for each

nursery:

$$h^2 = \frac{V_A}{V_P} = \frac{RV_{fam}}{V_{fam} + V_{block} + V_{res}}$$

271 where R is the relatedness of individuals within families (assumed to be half-siblings,

272 R = 4). Standard errors for heritability estimates were calculated as follows (Visscher

273 1998):

$$SE_{h^2} = R \sqrt{\frac{2(1 - \frac{h^2}{R})^2 \left[1 + (s-1)\frac{h^2}{R}\right]^2}{s(s-1)(f-1)}}$$

where R is the relatedness of trees within families, s is the mean number of offspring

per family and f is the mean number of families.

Results

Climatic variation among nurseries

Trees growing at each of the three nurseries experienced broadly similar mean daily

temperatures for each year between 2007 and 2010 which were not significantly

different (Figure 1) with the exception of NE and NG in both 2008 and 2009, where

mean daily temperature at NG was significantly warmer than at NE in both years and

for NG and NW in 2009. Trees at NG also experienced the fewest chilling days (total

119 between 2007 and 2010, compared with NE = 157 and NW = 123) and had

higher daily GDD throughout the majority of the year (Figure 1) compared to the

other two nurseries. In contrast, the daily temperature variance that trees were

exposed to was consistently and significantly lower for trees growing at NW than for

trees growing at either NG or NE (Figure 1). The daily temperature variance that

trees experienced in NG and NE was also significantly different, although the

magnitude of the difference was less than when comparing NW with either NE or NG

(Figure 1).

291
Figure 1. Climatic variation at the three nurseries for the period July 2007 to

December 2010 when P. sylvestris seedlings were growing in pots at the sites. Lines

indicate smoothed monthly means for each year. Boxplots indicate values for the

entire year: solid grey lines indicate the median value; the bottom and top of the

boxes indicate the first and third quartile; and the upper and lower whiskers extend to

the highest and lowest values within 1.5 times for interquartile range. No data are

included for daily temperature variance in 2007 due to the extent of missing hourly

data for this year. Significant differences among nurseries within each year are

indicated with asterisks (: p = 0.01-0.05; **: p = 0.001-0.01; ***: p = 0.0001-0.001;*

*****: p < 0.0001; ns: not significant. Units: mean temperature, °C; daily temperature*

variance, °C^2, growing degree days, number of degrees above 5 °C for each day.

A climatic profile was generated for each of the nurseries based on differences in

normalised values for mean daily temperature, temperature variance during the day

and night, GDD and chilling days (Figure 2a), highlighting differences among the

three nurseries summarised in the text above.

Interestingly, the differences in temperature variance were most pronounced among

nurseries and consistent among years during the daytime within the growing season

period (Figure S1). During this period, daytime variance was significantly higher at

NG than at NE or NW, and higher at NE than at NW in all years. In contrast, variance

during the night-time was lowest at NG compared to the outdoor nurseries in all

years outside the growing season (mean night temperature variance 2008-2010: NE

= 2.40 °C^2; NG = 1.35 °C^2; NW = 2.17 °C^2).

Trait variation among nurseries

Trees growing in NG had higher mean values for traits relating to growth than those

growing outdoors in NE and NW by 2010 (Figure 2b): they had thicker stems and

grew taller and faster, on average. They also had larger increments for both height

and stem diameter in 2010. Trees grown in NE were the smallest and thinnest with

the slowest growth rate and lowest increments on average in 2010. In contrast,

mean values for traits relating to form, and survival were lower, and phenology was

earlier, for trees growing in NG than those growing in one or both outdoor nurseries

(Figure 2b) with the exception of needle length. Trees growing at each nursery had

distinct profiles for mean trait values across the trait groups: trees growing at NE

generally had higher values for traits relating to form and survival, trees growing at

NG had higher values for traits relating to growth and trees growing at NW had

moderate values for traits relating to growth and form but also had the highest values

for phenology traits (indicating both late budburst initiation but also late growth

cessation). Coefficient of variation (CV) for each trait at each nursery were generally

the inverse of observed mean values (Figure 2c), for example: whereas trees

growing at NG had the highest mean values for all traits relating to growth, they had

the lowest CVs for growth traits, with the exception of relative growth rate. As a

result, the profiles of each nursery for CVs was quite different to their profiles for

mean trait values (Figures 2b and 2c): trees growing at NG showed greater variation

than the other nurseries for traits relating to phenology and, to a lesser extent, form;

trees growing at NW had moderate to high variation in growth traits compared to the

other nurseries, as well as for two of the three traits relating for form; and trees

growing at NE showed low overall levels of variation compared to trees growing at

the other two nurseries, but had distinctively high values for specific traits within each

group, namely needle length (form), growth cessation (phenology) and stem

diameter increment (growth). Comparing climatic and trait profiles for each nursery

allows simple relationships to be identified, for example: trees growing at NG and

experiencing higher values for climatic variables during the growing season than

trees at NE also have higher mean trait values and lower CVs for those traits,

whereas trees growing at NE which experience higher values for climatic variables

outside the growing season (more chilling days and higher nighttime temperature

variance) than NW have higher values for traits relating to form (with the exception of

needle length). Interestingly, differences among nurseries in daytime temperature

variance correspond inversely to differences in coefficient of variation (so, trees

growing in nurseries with less temperature variance have higher coefficients of

variation) for most growth traits.

Figure 2. Differences among nurseries (NE, NG and NW) based on normalised

variables, represented as profiles for a) climatic variables; b) mean trait values; and

c) coefficient of variation for trait values. Traits relating to form, growth, phenology (b

and c) and survival (b only) are indicated. For climatic profiles: 'Growing' refers to

variables recorded within the growing season; 'Not growing' refers to variables

recorded outside the growing season; 'Mean' is the mean average daily temperature;

'Chilling' is the mean number of days over the period 2007-2010 outside the growing

seasons where the mean maximum temperature <5 °C; 'Night variance' is the mean

variance in temperature during the night outside of the growing season; 'Day

variance' is the mean variance in temperature during the day within the growing

season; 'GDD' is the mean growing degree days within the growing season. For

climatic profiles, data from 2007 to 2010 are used with the exception of temperature

variances when data from 2007 are not included. For traits measured in multiple

years, only the most recent year is shown.

All trait means were significantly different among the three nurseries for each trait

and each year with the exception of height increment in 2009 and relative growth

rate in 2010, both among NG and NW, and growth cessation among NE and NG

(Figure S2). It is notable that the length of needles was considerably longer in all

nurseries in 2008 than in either 2007 or 2009 (Figure 3; Figure S3). Needles which

grew in 2008 were, on average, 3.93 (NE) to 4.59 (NW) times longer than those

which grew in 2007, and 2.11 (NW) to 3.21 (NE) times longer than needles which

grew in 2009.

Overall survival was much lower for trees in NW compared to the other two nurseries

(Figure S3) although this was due to a large spike in mortality in 2008 (Figure S4).

Mortality was highest/joint-highest at NW in three of the four assessment years and

was also extremely high in the other year. Mortality at the other outdoor site (NE)

was the lowest of all nurseries across all four years. By 2010, there were significant

differences among nurseries in seedling mortality. The highest levels of mortality in a

single year were observed in 2008 at NW (25.03 % annual mortality) which

accounted for over half of the seedling deaths across the period for this nursery. In

contrast, mortality in 2010 was low for all nurseries.

The association between trait variation and mortality was examined by comparing

the relative difference in trait means between trees that were alive and those that

died in the year after a trait was measured (Figure S5). Trait values were nearly

always lower (for traits relating to growth and form) or later (for traits relating to

phenology) for trees that died in the following year. Therefore, trees that died in the

following year were more likely to be shorter, thinner, grow less rapidly, form fewer

buds and shorter needles, and burst bud and stop growing later than trees which

remained alive (Figure S5). Exceptions to this were growth cessation in 2008 for

trees growing in NG, some growth traits (absolute and increment height and relative

growth rate) for trees growing in NW in 2008 and needle length in 2009 for trees

growing in NW and NE.

The relative difference in trait values between trees that were alive and those that

died increased as trees got older (although differences over more than one year can

only be compared for growth traits, as traits relating to phenology and form were not

measured in multiple years, with the exception of needle length, Figure S5).

However, the numbers of trees in the 'dead' group also reduced substantially over

this period, from an average of 221 trees in 2007 to 64 trees in 2008 and 10 trees in

2009 compared to those in the 'alive' group (average numbers of trees counted for

each trait: 1,313 in 2007; 1,198 in 2008; 1,191 in 2009).

Associations between nursery climate and trait variation

The timing of budburst in 2008 was significantly different among nurseries (Figure

S2) with budburst occurring, on average, 23 days earlier at NG than at either NE or

NW. The estimated number of days (since 31 March 2008) by which 50 % of the

trees had burst bud was 38.7 and 38.8 days in NE and NW, respectively, but only 16

days in NG (Figure S6). In contrast, over the same period, the differences among

nurseries in GDD was much lower (Figure S6, NE: 207.3; NG: 237.7; NW: 238.1).

Trees at NE required fewer cumulative GDD, and those at NW and NG required

more cumulative GDD, to complete budburst throughout the assessment period but

only by around 30 degree days. Trees at NW were exposed to two chilling days

during the budburst period (5th and 6th April 2008) but the environments were

otherwise similar among the two outdoor nurseries (NE and NW, Figure 1).

Despite significantly earlier initiation of budburst in NG, trees ceased growing at

around the same time as those in NE (Figure 2 and Figure S2) which was on

average about 13 days earlier than in NW. The duration of growth (period between

budburst and growth cessation) therefore varied on average between 166 and 189

days (NE and NG, respectively) with trees in NW actively growing for an average of

178 days each year. Mean growth duration was significantly positively associated

with growing season length at each site (Figure 5) although the amount of variation

in this trait at each site was substantial.

Figure 5. Linear regressions between growing season length (the period in days

bounded by a daily mean temperature >5 °C for >5 consecutive days and daily mean

temperature <5 °C for >5 consecutive days, after 1st July) and a) mean growth

duration in 2008; and b) mean relative growth rate. Mean growth duration: number

of days between budburst and growth cessation among trees at each site in 2008.

Error bars are one standard deviation either side of the mean. Mean relative growth

rate: mm per year as a proportion of its size at the end of the previous growing

season from 2007 to 2010.

There was a positive relationship between mean relative growth rate and the growing

season length at each site in each year (between 2007 and 2010, Figure 5), with a

possible threshold for growing season length (270 days) below which the growth rate

appeared to be limited. In-season performance might also be strongly associated

with the length of growing season in the previous year, due to provisioning of the bud

during the growing season, but these comparisons will be more powerful using field

rather than nursery data, due to the comparatively small number of datapoints

available for the latter.

Relationships within and among traits

Age-age correlations were performed for measurements of a given trait obtained in

pairs of years for all trees in each nursery site separately (Figure S7). The majority of

age-age correlations were significant across each pair of years in each of the

nurseries, with the exception of needle length (between years 2007 and 2008 in NG;

between years 2008 and 2009 in NE), diameter increment (between years 2009 and

2010 in NG), and height increment (between years 2007 and 2009, and 2007 and

2010 both in NG). Traits were consistently either positively or negatively correlated

among years, with the exception of height increment and needle length. Correlations

among height increments in different years for individuals in NG were negative when

2010 is included (in contrast to the same years in other nurseries for this trait) – this

is the year during which the pots were moved outdoors from the glasshouse. The

patterns of pairwise comparisons among the sites were more similar for the outdoor

sites (NE and NW) than NG. Correlations generally increased in significance as the

age of the trees increased and as the number of years between comparisons

decreased (Figure S7).

The number of traits measured across multiple years that were consistently

significantly associated (in the same direction) among years were reduced prior to

subsequent analyses, either to a single (most recent) assessment year (for traits for

which successive measurements were not independent: DA, HA) or to a mean value

among years (HR, DI). Traits which were not consistent among years (HI, NL) were

considered separately for each year.

Pairwise comparisons among growth, form and phenology traits both within and

among years (Figure S8) show that pairs of traits were more highly significantly

correlated in NW than in either NG or NE, particularly among growth traits and

among budburst and other traits. Relationships were often inconsistent among sites:

of the 105 pairwise comparisons among traits, 21 were not consistent (i.e. not

always either positive or negative) among nurseries. Of the inconsistent correlations,

nearly all (N = 17) involved needle length or height increment (Figure S8).

Significantly associated growth traits were always positive, with the exception of

height increment in 2007 in NE and NW and in all years at NG (Figures S7 and S8).

Phenology traits were highly significantly positively correlated with one another and

were generally negatively correlated with traits related to form (with the exception of

height to diameter ratio and needle length in 2009). Considering only traits that were

measured in the same year (i.e. 2008 for phenology traits), days to budburst were

negatively correlated with height increment in NW, positively correlated in NE and

not associated in NG. Days to budburst were highly significantly negatively

correlated with all growth traits at NW, whereas correlations were either not

significant or positively significant for growth traits in NG and NE. In contrast, days to

cessation of growth were highly positively correlated with most growth traits in NG

(with the exception of height increment in 2007) but associations were much weaker

and/or negatively correlated in the other nurseries. Thus, in general, a long growing

season between budburst and growth cessation resulted in larger plants.

For further analyses, traits within groups which were consistently highly significantly

positively correlated with one another were reduced to a single representative trait

(height increment 2008 to 2010, relative growth rate and absolute and increment

stem diameter were all represented by absolute height in 2010; needle length in

2007 and 2008 were represented by number of buds; growth cessation was

represented by budburst)

Partitioning trait variance

There were highly significant differences among nurseries (i.e. environmental

variation over a large scale) as well among blocks within nurseries (environmental

variation over a small scale) for every trait (Table S1). Nursery accounted for a

greater proportion of variation than block for needle length, number of buds, height

increment and budburst (Figure S9), whereas the contribution of block to the total

variance was greater than nursery for height to diameter ratio and absolute height.

Narrow sense heritability of growth traits was highest for trees growing in NG (Figure

6). For traits relating to form (specifically, bud counts and height to diameter ratio),

heritability values were lower for trees growing in NG compared to those growing in

NE and NG. Narrow sense heritability was highest overall for height increment at all

three nurseries (Figure 6) but was very low for the other growth trait (absolute height)

in all nurseries. Standard errors were extremely large, particularly in NW.

Figure 6. Estimated narrow sense heritability for each trait at each nursery (NG, NE,

NW) with associated standard error indicated by error bars. The acceptable range of

narrow sense heritability values is indicated by a white background (h^2 = 0 to 1) and

the area outside of this range is indicated by a grey background (h^2 > 1; h^2 < 0).

There is no value for narrow sense heritability for HD10 in NW due to zero

contribution of family to the total variance.

Discussion

This study highlights the profound impact that relatively small differences in the

temperature profiles of nurseries can have on trait variation in tree seedlings. The

two outdoor nurseries, only ca. 200 km apart, were exposed to the same nutrient, pot

and watering regimes and experienced similar average temperatures and monthly

growing degree days. The major difference between the sites occurred during the

daytime within the growing season, when daily temperature variance was much

higher for trees growing in the nursery in the east of Scotland compared to the

nursery in the west. Seedlings in the glasshouse nursery, in contrast, experienced

high daily temperature variance (which was also higher than in the neighbouring

outdoor nursery), but temperatures were also higher overall. The effects of warmer

mean temperatures (Morison and Morecroft 2006) and extremes of temperature (Niu

et al. 2014) on plant adaptive traits are increasingly well recognised, especially in the

light of an anticipated increase in extreme weather events due to climate change

(Herring *et al.* 2022). However, the contrasting effects of temperature variance and

mean temperatures and their comparative impact on the growth and development of

plants and trees, is less well understood.

In our study, all traits were strongly and significantly affected by the nursery

environment. The effect was particularly pronounced between the glasshouse and

neighbouring outdoor nursery, where the main difference was mean temperature,

compared to the two outdoor nurseries, where the main difference was daytime

temperature variance during the growing season. The distinct climatic profiles

generated for each nursery demonstrated the comparative importance of differences

in temperature mean and variance on adaptive traits.

Growing the seedlings under high temperature variance (resulting in more days at

the extreme ends of the temperature range) but with the same mean temperature led

to lower rates of growth: a finding supported by Chiang, Bånkestad and Hoch (2020),

who report significantly taller seedlings when grown in fixed or sinusoidal

temperature regimes compared to the same species grown in pots outdoors with a

naturally fluctuating and high temperature variance: but only within in spring. When

grown in the summer, the same seedlings grown in a fixed environment were

reported to be significantly shorter than those grown in the outdoor environment,

possibly because the lower temperatures that seedlings were exposed to in the

outdoor environment in the spring did not occur during summer (Chiang, Bånkestad

and Hoch 2020). The extent of the difference between day time and night time

temperatures is also known to directly influence key plant traits relating to growth

and form, such as internode length and height (Myster and Moe 1995). Similarly, the

allocation of resources to traits relating to tree form in this study appeared to be

more important in the nursery with both high variance and low mean temperatures,

indicating that a more conservative growth strategy was potentially balanced by

investment in traits relating to tree form (Banin *et al.* 2012), such as number of buds.

The role of mean temperature on plant growth is better understood, with higher

mean temperatures generally positively associated with higher rates of growth

(measured as specific leaf area (Poorter *et al.* 2010). Our study supported this

finding, with seedlings grown in a nursery with warmer mean temperatures showing

higher rates of growth. Although the temperatures of our nurseries were not

controlled, and only a few nursery environments were studied, the environments we

studied are analogous to those used in the forestry industry (i.e. outdoor nurseries

and glasshouses). We only recorded air temperature, but soil temperature mean and

variance may also have been important factors for the growth and development of

the seedlings. Soil temperature is unlikely to have been consistent among the

glasshouse and outdoor environments: it has been reported that soil temperature

inside pots tracks but lags behind the air temperature until solar radiation falls

directly onto pots, when the temperature of the soil within the pots can rise to more

than 20 °C higher than the air temperature (Poorter *et al.* 2016) and that increased

soil temperature is associated with increased growth rate (Weih and Karlsson 1999)

in saplings.

Growing Scots pine seedlings in nursery environments with different temperature

means, variances and photoperiods also had strong effects on phenological

variation. Budburst timing was strongly associated with growing degree days at all

sites, but was not fully explained by differences among sites. This suggests a

complex relationship between warm temperatures (i.e. growing degree days) and

cold temperatures (often referred to as the chilling requirement), although the

number of nurseries/years assessed in this study is too low to enable clear

conclusions to be reached. Plant species generally cease growing in response to

decreasing photoperiod (Nitsch 1957; Singh *et al.* 2017) although some may

respond to decreasing temperature instead (Heide and Prestrud 2005). In our study,

trees growing in neighbouring nurseries, outside or within a glasshouse, experienced

the same photoperiod but different mean temperatures. Whereas budburst occurred

much earlier in the glasshouse than in the neighbouring outdoor nursery (reflecting

the different cumulative growing degree days over the preceding period), growth

cessation occurred on average at a similar time.

The accuracy of using young seedlings to predict economic traits at harvest is of

major interest and importance in forestry (Lee 2002; Hong, Fries and Wu 2015), but

detailed early assessments of tree seedlings are rarely available. This study and

previous work indicate that we cannot make predictions based on trait correlations at

a very early age, but that age-age correlations strengthen with age. Within these

experimental trials, which will be transplanted to field sites, the same traits will

continue to be measured into maturity enabling age-age correlations from seedling to

mature trees to be estimated with high resolution (many traits will be measured

annually). This study highlights the relative consistency of age-age correlations

among the different nursery environments, with the exception of height increment in

NG in the years before and after trees were moved outside. The latter showed that

plants experienced a shock as a result of the move outdoors which impacted their

subsequent growth. The well-documented phenomenon of 'transplantation shock' is

known to induce changes to growth and development (Close, Beadle and Brown

2005) whereby growth is impaired for a period following transfer. The accuracy of

age-age correlations should also, therefore, be reviewed in the context of the general

environment the trees are measured and the specific environment the trees have

been exposed to in the recent past.

The interactions among different traits were also affected by the nursery that plants

were grown in. For example, while timing of growth cessation was positively

correlated with most growth traits in the protected glasshouse environment, these

relationships were weaker in outdoor environments, possibly because these traits

were not responding to a single climatic variable but to a combination of many.

Similarly, traits relating to form were inconsistent in their relationship with growth

traits among nurseries, with more positive associations observed in the nurseries

with higher temperature variance compared to the one with low temperature

variance. These findings will be investigated and tested further once trees are

transplanted to their field locations, as measurements can be made over many more

years.

The observation that narrow sense heritability values vary not only among traits but

also among different environments has been recognised in many systems (reviewed

by Hoffmann and Merilä (1999)). These findings have led to the proposal of various

hypotheses that aim to explain why the favourability of the conditions in which

organisms grow should lead to higher or lower heritability of quantitative traits. In our

study, it appears that growing seedlings in the warmer (and, presumably, more

favourable) environment of the glasshouse resulted in higher narrow sense

heritability values for growth and phenology traits compared to seedlings grown in

cooler conditions, although the errors in all estimates were large. This could be due

to the genetic potential being reached in the more favourable environment

(Gebhardt‑Henrich and Van Noordwijk 1991; Hoffmann and Merilä 1999). In

contrast, narrow sense heritability for traits relating to form had higher values for

seedlings grown in the cooler (i.e. putatively less favourable) conditions in the

outdoor nurseries, possibly because traits relating to form are more important for

long-term survival in unfavourable conditions than those relating to rapid growth

(Banin *et al.* 2012).

Conclusions

Despite the relatively low levels of environmental variation among the three

nurseries, there were significant effects on seedling mortality, variation in traits

relating to growth, form and phenology, interactions among traits, and on narrow

sense heritability. These findings have implications for the choice of nursery in which

plants are raised given that effects may persist and affect performance in the field.

Growers may wish to assess firstly whether raising plants in relatively benign nursery

environments is necessarily appropriate given the shock they receive when moved to

less benign conditions, and secondly whether the phenotypes produced by nursery

conditions benefit or hamper the plants on transfer to the field. In any case, we have

demonstrated the importance of measuring and collecting data at an early stage in

order to compare trait variation throughout the lifetime of trees, from seedling

through to maturity. The trees in this study have now been transplanted to three field

environments (Beaton *et al.* 2022) and subsequent analyses will explore ontogenetic

effects on intraspecific genetic variation in a wide range of phenotypic traits.

Funding

This work was supported by the newLEAF project (NE/V019813/1) under the UK's

'Future of UK Treescapes' programme, which was led by UKRI-NERC, with joint

funding from UKRI-AHRC & UKRI-ESRC, and contributions from Defra and the

Welsh and Scottish governments.

This work was also part of PROTREE, a project supported by grant BB/L012243/1,

funded jointly by UKRI's BBSRC, NERC, & ESRC and Defra, the Forestry

Commission and the Scottish Government, under the Tree Health and Plant

Biosecurity Initiative

The James Hutton Institute contribution to this work was supported by Scottish

Government's Rural & Environment Science & Analytical Services Division, through

their Strategic Research Programmes (2006–2011, 2011–2016 and 2016–2022).

Acknowledgements

We thank Stewart Mackie and staff at Forest and Land Scotland (Selkirk Office), the

Forestry Commission, National Trust for Scotland, and Donald Barrie (Glensaugh)

for their help in locating, establishing and securing access to sites.

Conflict of interest statement

None declared

Data availability statement

The data underlying this article are available from the Environmental Information

Data Centre (EIDC, https://eidc.ac.uk), at https://doi.org/10.5285/29ced467-8e03-4132-83b9-dc2aa50537cd

References

Aldhous JR, Mason WL. Forest Nursery Practice. Forest Research, Farnham, 1994:296.

Banin L, Feldpausch TR, Phillips OL *et al.* What controls tropical forest architecture? Testing environmental, structural and floristic drivers. *Global Ecology and Biogeography* 2012;**21**:1179–90.

Beaton J, Perry A, Cottrell J *et al.* Phenotypic trait variation in a long-term multisite common garden experiment of Scots pine in Scotland. *Sci Data* 2022;**9**:671.

Benavides R, Carvalho B, Matesanz S *et al.* Phenotypes of *Pinus sylvestris* are more coordinated under local harsher conditions across Europe. *J Ecol* 2021, DOI: 10.1111/1365-2745.13668.

Brancalion PHS, Holl KD. Guidance for successful tree planting initiatives. *J Appl Ecol* 2020, DOI: 10.1111/1365-2664.13725.

Burdon RD, Bannister MH, Low CB. Genetic survey of Pinus radiata. 5: Between-trait and age-age correlations for growth rate, morphology, and disease resistance. *New Zealand Journal of Forestry Science* 1992;**22**:211–27.

Chiang C, Bånkestad D, Hoch G. Reaching natural growth: the significance of light and temperature fluctuations in plant performance in indoor growth facilities. *Plants* 2020;**9**, DOI: 10.3390/plants9101312.

Close DC, Beadle CL, Brown PH. The physiological basis of containerised tree seedling 'transplant shock': a review. *Australian Forestry* 2005;**68**:112–20.

Dhar A, Balliet N, Hawkins CD *et al.* Bud flush phenology and nursery carryover

effect of paper birch provenances. *iForest* 2015;**8**:809–17.

Domisch T, Finér L, Lehto T. Effects of soil temperature on biomass and

carbohydrate allocation in Scots pine (Pinus sylvestris) seedlings at the

beginning of the growing season. *Tree Physiol* 2001;**21**:465–72.

Dong M, Fan Y, Wu Z *et al.* Age–age correlations and early selection for growth

traits in 40 half-sib families of Larix principis-rupprechtii. *Journal of Forestry*

Research 2018:1–7.

Donnelly K, Cavers S, Cottrell JE *et al.* Cryptic genetic variation and adaptation to

waterlogging in Caledonian Scots pine, Pinus sylvestris L. *Ecol Evol*

2018;**8**:8665–75.

Downs RJ, Borthwick HA. Effects of photoperiod on growth of trees. *Botanical*

Gazette 1956;**117**:310–26.

Falconer DS, Mackay TFC. *Introduction to Quantitative Genetics*. 4th ed. Essex, UK:

Longman, 1996.

Forestry Commission. *Tree Supply Report*. UK Government, 2023.

Forest Research. *The UK Forestry Standard*. Forest Research, Farnham, 2023:139.

Fukatsu E, Tamura A, Takahashi M *et al.* Efficiency of the indirect selection and the

evaluation of the genotype by environment interaction using Pilodyn for the

genetic improvement of wood density in *Cryptomeria japonica*. *J For Res*

2011;**16**:128–35.

Gebhardt‐Henrich SG, Van Noordwijk AJ. Nestling growth in the Great Tit I.

Heritability estimates under different environmental conditions. *J Evol Biol*

1991;**4**:341–62.

Gruffman L, Ishida T, Nordin A *et al.* Cultivation of Norway spruce and Scots pine on

organic nitrogen improves seedling morphology and field performance. *Forest*

Ecology and Management 2012;**276**:118–24.

Harrell Jr FE. *Hmisc: Harrell Miscellaneous*. R package, 2024.

Heide OM, Prestrud AK. Low temperature, but not photoperiod, controls growth

cessation and dormancy induction and release in apple and pear. *Tree Physiol*

2005;**25**:109–14.

Heiskanen J, Rikala R. Effect of Peat-based Container Media on Establishment of

Scots Pine, Norway Spruce and Silver Birch Seedlings after Transplanting in

Contrasting Water Conditions. *Scandinavian Journal of Forest Research*

2000;**15**:49–57.

Herbert R, Samuel S, Patterson G. *Using Native Stock for Planting Native Trees and*

Shrubs. Forestry Commission, 1999.

Herring SC, Christidis N, Hoell A *et al.* Explaining Extreme Events of 2020 from a

Climate Perspective. *Bull Amer Meteor Soc* 2022;**103**:S1–129.

Hoffmann AA, Merilä J. Heritable variation and evolution under favourable and

unfavourable conditions. *Trends Ecol Evol* 1999;**14**:96–101.

Holl KD, Brancalion PHS. Tree planting is not a simple solution. *Science*

2020;**368**:580–1.

Hong Z, Fries A, Wu HX. Age trend of heritability, genetic correlation, and efficiency

of early selection for wood quality traits in Scots pine. *Can J For Res*

2015;**45**:817–25.

Lee SJ. Selection of parents for the Scots pine breeding population in Britain.

Forestry 2002;**75**:293–303.

Luoranen J. Autumn versus spring planting: the initiation of root growth and

subsequent field performance of Scots pine and Norway spruce seedlings. *Silva*

Fenn 2018;**52**, DOI: 10.14214/sf.7813.

Marron N, Dillen SY, Ceulemans R. Evaluation of leaf traits for indirect selection of

high yielding poplar hybrids. *Environ Exp Bot* 2007;**61**:103–16.

Mason WL, Alía R. Current and future status of Scots pine (Pinus sylvestris L.)

forests in Europe. *For Syst* 2000;**9**:317–35.

Minitab. *Minitab Statistical Software*. Minitab, 2024.

Morison JIL, Morecroft MD eds. *Plant Growth and Climate Change*. Oxford, UK:

Blackwell Publishing Ltd, 2006.

Myster J, Moe R. Effect of diurnal temperature alternations on plant morphology in

some greenhouse crops—a mini review. *Sci Hortic* 1995;**62**:205–15.

Nerg A, Kainulainen P, Vuorinen M *et al.* Seasonal and geographical variation of

terpenes, resin acids and total phenolics in nursery grown seedlings of Scots

pine (*Pinus sylvestris* L.). *New Phytologist* 1994;**128**:703–13.

Nitsch JP. Growth response of woody plants to photoperiodic stimuli. *Proceedings*

American Society for Horticultural Science 1957;**10**:512–25.

Niu S, Luo Y, Li D *et al.* Plant growth and mortality under climatic extremes: An

overview. *Environ Exp Bot* 2014;**98**:13–9.

Osorio LF, White TL, Huber DA. Age-age and trait-trait correlations for Eucalyptus

grandis Hill ex Maiden and their implications for optimal selection age and

design of clonal trials. *Theor Appl Genet* 2003;**106**:735–43.

Pan Y, Birdsey RA, Fang J *et al.* A large and persistent carbon sink in the world's

forests. *Science* 2011;**333**:988–93.

Perry A, Brown AV, Cavers S *et al.* Has Scots pine (*Pinus sylvestris*) co-evolved

with *Dothistroma septosporum* in Scotland? Evidence for spatial heterogeneity

in the susceptibility of native provenances. *Evol Appl* 2016;**9**:982–93.

Poorter H, Fiorani F, Pieruschka R *et al.* Pampered inside, pestered outside?

Differences and similarities between plants growing in controlled conditions and

in the field. *New Phytol* 2016;**212**:838–55.

Poorter H, Niinemets U, Walter A *et al.* A method to construct dose-response curves

for a wide range of environmental factors and plant traits by means of a meta-

analysis of phenotypic data. *J Exp Bot* 2010;**61**:2043–55.

R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna,

Austria: R Foundation for Statistical Computing, 2024.

Ramírez-Valiente JA, Solé-Medina A, Pyhäjärvi T *et al.* Selection patterns on early-

life phenotypic traits in Pinus sylvestris are associated with precipitation and

temperature along a climatic gradient in Europe. *New Phytol* 2021;**229**:3009–25.

Salmela MJ, Cavers S, Cottrell JE *et al.* Seasonal patterns of photochemical

capacity and spring phenology reveal genetic differentiation among native Scots

pine (Pinus sylvestris L.) populations in Scotland. *Forest Ecology and*

Management 2011;**262**:1020–9.

Selander J, Immonen A. Effect of fertilization and watering of Scots pine seedlings

on the feeding preference of the pine weevil (Hylobius abietis L.). *Silva Fenn*

1992;**26**, DOI: 10.14214/sf.a15637.

Singh RK, Svystun T, AlDahmash B *et al.* Photoperiod- and temperature-mediated

control of phenology in trees - a molecular perspective. *New Phytol*

2017;**213**:511–24.

Vapaavuori EM, Rikala R, Ryyppö A. Effects of root temperature on growth and

photosynthesis in conifer seedlings during shoot elongation. *Tree Physiol*

1992;**10**:217–30.

Villar-Salvador P, Ocaña L, Peñuelas J *et al.* Effect of water stress conditioning on

the water relations, root growth capacity, and the nitrogen and non-structural

carbohydrate concentration of Pinus halepensis Mill. (Aleppo pine) seedlings.

Ann For Sci 1999;**56**:459–65.

Visscher PM. On the sampling variance of interclass correlations and genetic

correlations . *Genetics* 1998;**149**:1605–14.

Weih M, Karlsson PS. The nitrogen economy of mountain birch seedlings:

implications for winter survival. *Journal of Ecology* 1999;**87**:211–9.

Westaway S, Grange I, Smith J *et al.* Meeting tree planting targets on the UK's path

to net-zero: A review of lessons learnt from 100 years of land use policies. *Land*

use policy 2023;**125**:106502.

Williams CG. Accelerated short-term genetic testing for loblolly pine families. *Can J*

For Res 1988;**18**:1085–9.

Wu HX, Yeh FC, Dhir NK *et al.* Genotype by environment interaction and genetic

correlation of greenhouse and field performance in *Pinus contorta* ssp. *latifolia*.

Silvae genetica 1997;**46**.

Zobel BJ, Talbert JT. *Applied Tree Improvement*. New York: John Wiley & Sons,

1984.