Analysis of the effects of mating systems on lineage diversification across multiple genera

Kuangyi Xu[1*]

[1] Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks St, Toronto, ON, Canada M5S 3B2

[*] Correspondence: Kuangyi Xu, Email: kuangyi.xu@utoronto.ca

ORCID: https://orcid.org/0000-0002-3645-9580

Keywords: mating system, speciation, diversification, selfing, life form, hidden state, mixed mating

Summary

- The transition from outcrossing to self-fertilization is a major evolutionary trend in
plants, with selfing long hypothesized an evolutionary dead end. Recent theories suggest
that elevated extinction rates may be restricted to highly selfing rather than mixed-mating
populations. Previous analyses of the effects of mating systems on diversification found
mixed results, varying by focal clades and potentially suffering from several limitations.

- We collected data on three mating-system-related characters. We also collected life form
data to indirectly distinguish between mixed mating and highly selfing. We estimated
speciation and extinction rates for character states, and state transition rates in 27 genera.

- We find that outcrossing lineages diversify at higher rates than selfing lineages, while the
impact of mating systems on speciation is not consistent across genera. Among self-
compatible lineages, annuals/short-lived exhibit significantly lower diversification rates
than perennials. Models incorporating hidden states often provide a better fit than those
without.

- The results suggest that selfing lineages overall have lower diversification rates. This may
be primarily contributed by elevated extinction rates in highly selfing lineages, which
may also explain the prevalence of mixed-mating systems. However, the influences of
mating systems on diversification may be often driven by other factors correlated with
mating system transitions.

Introduction

The transition from outcrossing to self-fertilization is one of the major evolutionary

trends in flowering plants (Barrett, 2008), with selfing occurring in about 50% of species

(Goodwillie et al., 2005). However, it has long been hypothesized that selfing is an evolutionary

dead end (Igic & Busch, 2013). Transitions from selfing back to outcrossing are rarely observed

in nature (Barrett, 2013; Fujii et al., 2016), and self-incompatibility is hard to be regained once it

is lost (Igic et al., 2008). Compared to outcrossing lineages, selfing lineages are thought to suffer

higher extinction risks due to several microevolutionary effects of selfing (Hartfield et al., 2017).

Selfing increases homozygosity in the offspring, and reduces the effective population size

due to larger variance in the number of alleles an individual leaves to the next generation

(Caballero & Hill, 1992). Increased genome-wide homozygosity also reduces effective

recombination rates, which, in the presence of deleterious mutations, can lead to a further

reduction of the effective population size (background selection; Nordborg et al., 1996) and the

accumulation of deleterious mutations (Charlesworth et al., 1993; Xu, 2022). Moreover,

theoretical work and meta-analyses suggest that selfers tend to exhibit lower levels of additive

genetic variance in quantitative traits compared to outcrossers (Clo et al., 2019), although selfing

may temporarily reveal non-additive genetic variance in the short term (Clo et al., 2020; Clo &

Opedal, 2021).

Due to the reduced efficacy of selection in selfing populations, selfers are expected to

exhibit a higher ratio of nonsynonymous to synonymous divergence (D_n/D_s) or polymorphism

(π_n/π_s) compared to outcrossers (Glémin, 2007; Wright et al., 2008). Some studies did not

detect significant differences in these ratios when comparing sister species with contrasting

mating systems (Escobar et al., 2010). This is often attributed to the relatively recent origin of

selfing lineages, where inbreeding may not have persisted long enough to generate significant

differences. More recent studies, however, have found higher nonsynonymous-to-synonymous

ratios in selfers (Ness et al., 2012; Arunkumar et al., 2015; Wang et al., 2021).

The impacts of selfing on speciation remain poorly understood, as speciation can occur

through various processes (Coyne & Orr, 2004), and selfing can influence them in complex ways

(Marie-Orleach et al., 2024). Selfing may reduce gene flow between populations and contribute

to reinforcement (Castillo et al., 2016), but can also hinder the accumulation of alleles that drive

speciation by reducing effective population size and effective recombination rates. Recent

studies suggest that Dobzhansky-Muller incompatibilities might accumulate more rapidly in

selfing populations (Marie-Orleach et al., 2022; Marie-Orleach et al., 2024). Additionally,

transitions to selfing are often associated with a series of morphological changes (selfing

syndrome; Sicard & Lenhard, 2011), potentially contributing to reproductive isolation (Cutter,

2019). Furthermore, selfing may facilitate polyploid speciation by allowing polyploid species to

avoid mating with diploid individuals during the early stages of establishment (Rausch &

Morgan, 2005; Barringer, 2007).

The macroevolutionary effects of selfing on lineage diversification have been estimated

using state-dependent speciation and extinction (SSE) models (Helmstetter et al., 2023), but the

results vary across clades. Several studies have found that lineages with traits promoting selfing

tend to exhibit reduced rates of net diversification (speciation minus extinction) (Goldberg et al.,

2010; de Vos et al., 2014; Gamisch et al., 2015; Castillo et al., 2016; Freyman & Höhna, 2019).

However, opposite results have been observed in other clades (Castillo et al., 2016; Landis et al.,

2018). These studies also found mixed results for the effects of selfing on speciation, with selfing

enhancing speciation rates in some clades while inhibiting speciation in others.

Several potential limitations exist in previous diversification analyses. Most of these

studies were conducted at the family level, which may bias results toward genera with more

complete character state records. For instance, in the analyses of Solanaceae and Onagraceae,

over two-thirds and one-third of the included species are from *Solanum* and *Oenothera*,

respectively (Goldberg et al., 2010; Freyman & Höhna, 2019). Additionally, in many family-

level analyses, mating system polymorphism is primarily attributed to variation between genera,

while congeneric species often share identical mating system states. This can result in estimates

being confounded by unknown genus-specific traits. In fact, SSE models are prone to inferring

false positives; even if diversification rate variation is driven by other factors, a significant

dependency on the focal character may still be inferred if state transitions coincide with more or

less diverse parts of the tree (Rabosky & Goldberg, 2015). For example, the breakdown of self-

incompatibility may be often triggered by unfavorable environmental shifts, such as pollen

limitation (Busch & Schoen, 2008; Eckert et al., 2010; Encinas-Viso et al. 2020). In this case, it

may be pollen limitation, rather than the breakdown of self-incompatibility, that drives

extinction.

The problem of coincidental associations and clade dependency may be mitigated by

comparative analyses across multiple clades. For example, Sabath et al. (2016) showed that

dioecy neither consistently inhibits nor facilitates diversification across more than 30 genera.

Additionally, the issue of false positives has been explicitly treated by recently developed

methods, such as the HiSSE model (Hidden State Speciation and Extinction; Beaulieu &

O'Meara, 2016). HiSSE models extend BiSSE models by allowing speciation and extinction

rates to depend on both measured and hidden character states. HiSSE analysis in *Onagraceae*

found that, for all hidden states, self-compatible lineages consistently exhibit lower

diversification rates than self-incompatible lineages (Freyman & Höhna, 2019). However, HiSSE

analysis in Polemoniaceae found that diversification rates appear to be independent of whether

lineages are autogamous or cross-pollinated (Landis et al., 2018).

Importantly, early theoretical and empirical investigations often focus on differences

between predominantly outcrossing and predominantly selfing populations, while recent

progresses suggest that the rate of selfing is critical. Specially, a higher selfing rate reduces the

fixation probability of new adaptive mutations at a single locus only when alleles are dominant;

however, very high selfing rates significantly strengthens background selection, thereby

inhibiting fixation even for recessive alleles (Glémin & Ronfort, 2013). Additionally,

intermediate selfing rates are optimal in promoting rescue through the sweep of adaptive alleles

at multiple loci by preventing recombination from destroying haplotypes with multiple adaptive

alleles (Xu, 2023a). Populations with intermediate selfing rates may also avoid the rapid

accumulation of deleterious mutations that occurs in highly selfing populations (Xu, 2022). This

is because even a slight amount of effective recombination introduced by outcrossing is

sufficient to significantly slow down Muller's ratchet. Furthermore, compared to outcrossing

populations, populations with intermediate selfing rates tend to have lower genetic load and are

more likely to persist under pollen limitation (Busch & Delph, 2012; Roze, 2015; Xu, 2023b).

The above results suggest the hypothesis that elevated extinction rates may be restricted

to highly selfing, instead of intermediate selfing populations. This could partly explain why

intermediate selfing is common in nature, despite being predicted to be evolutionarily unstable,

while highly selfing, which is predicted to be evolutionarily stable, seldom occurs (Goodwillie et

al., 2005). Specifically, under the balance between lineage diversification and mating system

transitions, the much lower diversification rates in highly selfing lineages compared to mixed-

mating lineages could result in a low prevalence of highly selfing species and a substantial

proportion of mixed-mating lineages found in nature.

However, previous diversification analyses treated mating systems as binary-state

characters (e.g., self-incompatible/self-compatible). The binary-state classification provides no

information about the rate of selfing, since being self-compatible does not necessarily mean

selfing occurs at a high rate (Raduski et al., 2012). Therefore, testing the proposed hypothesis

necessitates diversification analyses that differentiate among predominantly outcrossing, mixed

mating, and highly selfing lineages.

To mitigate the limitations of previous analyses and clade dependency, we revisit the

question of the macroevolutionary consequences of selfing. We conducted BiSSE analyses

across 27 genera using a Bayesian framework. To reduce false positives, we also applied HiSSE

models to these genera and examined the effects of mating systems on diversification rates

across different hidden states.

To test the hypothesis that highly selfing lineages tend to have lower diversification rates

than those with intermediate selfing rates, it would be ideal to conduct diversification analyses

based on outcrossing rates. However, we did not identify appropriate genera with sufficient data

for this analysis. Therefore, we performed diversification analyses on the combination of self-

compatibility and life form states (annual/short-lived *versus* perennial). Given that annuals are

much more likely to be highly selfing compared to perennials (Aarssen, 2000; Ma et al., 2020),

we expect that self-compatible annuals will generally exhibit lower diversification rates and

higher extinction rates than self-compatible perennials.

Materials and methods

Data collection

We collected datasets for three characters related to the mating system: self-compatibility,

pollination system, and species-level outcrossing rate. The genus name were updated based on

the WFO Plant List (https://wfoplantlist.org). For the pollination system, we directly used the

dataset reported from Landis et al. (2018). The self-compatibility dataset was primarily obtained

by compiling datasets reported in previous works (Wagner et al., 2017; Grossenbacher et al.,

2017; Freyman & Höhna, 2019; Delaney & Igić, 2022; Krakos et al., 2022; Cascante-Marín et

al., 2023). For the dataset reported by Delaney & Igić (2022), we only included self-

compatibility data based on hand-pollination experiments, while sources marked as verbal

descriptions without experimental validation were considered unreliable and not used. In

addition, we included self-compatibility data for a few species that were previously missing

through a literature search. A species is classified as self-incompatible (SI) when the index of

self-compatibility is less than 0.2; otherwise, it is considered self-compatible (SC). In cases

where self-compatibility was measured or reported for multiple populations of a species, we

determined self-compatibility at the species level by selecting the self-compatibility status that

occurred in the majority of conspecific populations.

For the outcrossing rate dataset, we assembled data from three previous studies (Dumini

et al., 2009; Goodwillie et al., 2012; Whitehead et al., 2018). We also added estimations from

more recent publications. To do this, we focused exclusively on multi-locus estimates of

outcrossing rates, as these estimates are less influenced by selection compared to single-locus

estimates (Ritland, 2002; Bürkli et al., 2017). To identify relevant studies, we searched the Web

of Science up to November 12, 2023, for publications that cited at least one of five

methodological papers on multi-locus outcrossing rate estimations (Ritland, 1990; Ritland, 2002;

Ritland & Jain, 1981; David et al., 2007; Jones & Wang, 2010). For a species, when outcrossing

rate estimates were available for multiple populations or multiple flowering seasons, we

averaged these estimates to calculate a mean outcrossing rate at the species level. We recognized

that outcrossing rates often vary among conspecific populations. However, meta-analyses

indicate that this variation tends to be small in predominantly outcrossing and predominantly

selfing species (Whitehead et al., 2018), suggesting it may be valid to classify the mating system

of a species into discrete categories.

We classified the mating-system-related characters into binary states, with 0 representing

outcrossing and 1 representing selfing. Specifically, for self-compatibility, state 0 indicates self-

incompatible (SI) species, and state 1 indicates self-compatible (SC) species. For the pollination

system, state 0 and 1 correspond to cross-pollination and autogamy, respectively. For outcrossing

rates, state 0 represents species with mean outcrossing rates above 0.8, and state 1 represents

species with mean outcrossing rates below 0.8.

Two genera, *Solanum* and *Oenothera*, were found to be suitable to conduct diversification

analyses on the combination of self-compatibility and life form states. The life forms of *Solanum*

species were primarily determined using the dataset from Hilgenhof et al. (2023). Species

marked as shrub or tree in Hilgenhof et al. (2023) were classified as perennials. For species

marked as herb, we determined whether they are short-lived or perennial primarily based on
descriptions from Solanaceae Source (http://solanaceaesource.org) with additional literature
searching. Life forms of *Oenothera* species were determined based on descriptions in Dietrich &
Wagner (1988), Wagner et al. (2007), and Evans et al. (2011).

Phylogeny reconstruction

For genera selected from the pollination system dataset, we directly used the maximum
likelihood (ML) trees and bootstrap trees reported by Landis et al. (2018). For other genera,
phylogenies were reconstructed following the procedure described below.

1) For each genus, we selected outgroup species based on prior phylogenetic studies
(summarized in Table S1). Sequence data for all species within the genus, as well as the outgroup
species, were downloaded from GenBank and clustered using the R package *phylotaR* (Bennett
et al., 2018).

2) Within each cluster, sequences that were either twice as long or half as short as the
median sequence length were discarded. The best-represented sequence for each species was
selected using the "drop_by_rank" function in the *phylotaR* package with default settings.
Sequences within each cluster were then aligned using MAFFT version 7 (Katoh & Standley,
2013), and unreliable sites were removed using Gblocks (Castresana, 2000). Clusters containing
sequences from only few species were dropped. However, since previous studies suggest that
filtering genes based on missing data may reduce phylogenetic accuracy (Molloy & Warnow,
2018), most clusters were retained. The remaining clusters were concatenated into a partitioned
supermatrix for phylogenetic reconstruction.

3). For each genus, a ML tree and 100 bootstrap trees were reconstructed using raxmlGUI
2.0 (Edler et al., 2021) with an initial model test to select the best-fit model. To ultrametricize the
ML and 100 bootstrap trees, we employed the "chronos" function from the R package *ape*
(Paradis & Schliep, 2019). The smoothing parameter λ in the "chronos" function, which adjusts
the penalty for rate variation across branches (Sanderson, 2002), was determined through cross-
validation on the ML tree. The cross-validation was conducted using the "CV" function from the
R package *chronos* (https://github.com/josephwb/chronos), varying λ from 0.01 to 1000 at
intervals of 0.5 on a log10 scale. After ultrametricization, outgroups and species with missing
character state information were pruned for diversification analyses.

To reduce uncertainty caused by small sample sizes, we excluded genera from the

diversification analyses if their resulting phylogenies contained fewer than nine species or if

there were fewer than three species in each character state, unless the total number of species in

the genus was small. As a low sampling proportion reduces the confidence of SSE analyses

(Helmstetter et al., 2023), genera with excessively sampling fractions (i.e., the proportion of

species with character state data relative to the total number of species) lower than 5% were also

excluded from the analyses. Mating system character states, supermatrices, and phylogenies for

genera used in our diversification analysis are available at

https://doi.org/10.5281/zenodo.13265323. The number of species in each mating system state

and summary statistics of supermatrices for genera used in our diversification analyses are

available at Table S2.

Diversification analyses

We assessed the impacts of mating system states on lineage diversification using several

SSE models. To account for phylogenetic uncertainty, diversification analyses were performed

on 100 bootstrap trees for each genus. Briefly, we estimated diversification for outcrossing (state

0) and selfing (state 1) lineages using the BiSSE models (Maddison et al., 2007), implemented in

the R package *diversitree* (Fitzjohn, 2012). We also fitted a series of HiSSE models that include

hidden states using the R package *hisse* (Beaulieu & O'Meara, 2016). For genera with sufficient

character data, we additionally estimated diversification rates for the combination of self-

compatibility and life form states using MuSSE models (Multiple State Speciation and

Extinction), implemented in the *diversitree* package. Since estimations of extinction rates were

suggested to be unreliable (Rabosky, 2010), our analyses focused on the speciation and

diversification rates (speciation minus extinction).

All the SSE analyses were conducted by accounting for the sampling fraction of species

included in the analyses relative to the total number of species in a genus, which was determined

using the number of accepted species in the WFO Plant List. Since we lacked complete character

state information for all taxa in each genus, we assumed no state-dependent sampling bias, so

that the sampling fraction was the same for all character states.

BiSSE analyses

In a full BiSSE model, six parameters were estimated: the speciation rates for states 0 and

1 (λ_0, λ_1), the extinction rates for states 0 and 1 (μ_0, μ_1), and the transition rates between states 0

and 1 (q_{01}, q_{10}). For the self-compatibility character, we followed previous analyses (Goldberg

et al., 2010; Freyman & Höhna, 2019) and constrained the transition rate from state 0 (SI) to

state 1 (SC) to be 0 (i.e., $q_{10} = 0$). This constraint reflects the theoretical prediction and

observation that the breakdown of self-incompatibility is rarely reversible (Igic et al., 2008;

Barrett, 2013). When the character is pollination system or species-level outcrossing rate,

transitions between state 0 and state 1 may be reversible. Therefore, we implemented both the

full BiSSE model and a BiSSE model that assumes no transition from selfing (state 1) to

outcrossing (state 0). If the two models produced qualitatively different results for a genus (see

Table S3), the genus was considered sensitive to model assumptions and was dropped from the

BiSSE MCMC analysis.

For each tree, we implemented BiSSE models using Markov chain Monte Carlo

(MCMC) sampling (FitzJohn et al., 2009) to obtain the posterior probability distributions of each

parameter. MCMC chains were started at the maximum likelihood estimates of the model

parameters and were run for 2500 steps with the initial 20% steps discarded as the burn-in stage.

For each genus, we merged the post-burn-in MCMC steps for all 100 bootstrap trees, resulting in

a total of 200000 MCMC steps to account for phylogenetic uncertainty. Following Sabath et al.

(2015), We calculated the proportion of MCMC steps in which the posterior diversification rates

were higher for outcrossing lineages than for selfing lineages, denoted by $P(r_0 > r_1)$. Similar

quantities were calculated for speciation rates and extinction rates, denoted by $P(\lambda_0 > \lambda_1)$ and

$P(\mu_0 > \mu_1)$, respectively. These values should not significantly differ from 0.5 if the rate is

state-independent.

For each rate parameter, the prior distribution used in the MCMC sampling was assumed

to be exponential, with the mean being fp_0. Here p_0 is the initial rate estimate from the function

"starting.point.bisse" in the *diversitree* package, and f = 1, 2, 4 is a parameter that we varied to

examine whether the MCMC results were sensitive to the prior distributions (Sabath et al.,

2015). Genera were considered as unrobust and were excluded from the BiSSE analyses if the

$P(r_0 > r_1)$ values obtained under the three values of f differed by > 30% (see Table S3). In the

main text, we report the results obtained using f = 2.

To assess the significance of mating system on diversification across genera, we

combined $P(r_0 > r_1)$ values of robust genera. Following Sabath et al (2015), we used a one-

sample Wilcoxon rank-sum test. Under the null hypothesis of sate-independent diversification,

the mean value of $P(r_0 > r_1)$ is 0.5, as diversification rates in state 0 should be higher than in

state 1 for half of the MCMC steps. A similar test was conducted to evaluate the effects of

mating systems on speciation rates, using $P(\lambda_0 > \lambda_1)$ values.

MuSSE analyses

Our MuSSE analyses on the combination of self-compatibility and life form states were

similar to the BiSSE analyses described above. However, the MuSSE model included four

possible states: SI annual/short-lived (state 1), SI perennial (state 2), SC short-lived (state 3), and

SC perennial (state 4). As in the BiSSE analyses, we assumed no transitions from SC to SI states.

Transitions between life form states were reversible, as suggested from previous studies (Soltis et

al., 2013). We also assumed that simultaneous transitions between self-compatibility and life

form (e.g., from SI annual/short-lived to SC perennial) did not happen. Under these constraints,

the MuSSE model estimated 14 parameters: the speciation rates for each state ($\lambda_1, \lambda_2, \lambda_3, \lambda_4$),

the extinction rates for each state ($\mu_1, \mu_2, \mu_3, \mu_4$), transition rates in self-compatibility (q_{13}, q_{24}),

and transition rates in life form ($q_{12}, q_{21}, q_{34}, q_{43}$). MCMC sampling was run for 10000 steps,

with the initial 25% steps discarded as burn-in.

HiSSE analyses

For each genus, we fitted six SSE models from the *hisse* package (Beaulieu & O'Meara,

2016) to the ML tree and 100 bootstrap trees. Three models included only two mating system

states without hidden states, denoted as BiSSE. The other three models, labeled HiSSE,

incorporated two hidden states, A and B, resulting in four possible states: 0A, 1A, 0B, and 1B. In

all HiSSE models, we assumed no simultaneous transitions at both the mating system states and

hidden states (e.g., no transition from 0A to 1B). For both model categories, we considered three

types of model setups: a state-independent diversification model (BiSSE null, HiSSE CID2), a

state-dependent diversification model with no backward transitions from state 1 (selfing) to state

0 (outcrossing) (BiSSE irr, HiSSE irr), and a state-dependent diversification model with

reversible transitions between states 0 and 1 (BiSSE rev, HiSSE rev). It should be emphasized

that the BiSSE models from the *hisse* package here differ from those in the *diversitree* package

used in our MCMC analyses, as the model setups are not the same.

For each tree, we selected the best-fitting model from the six models based on the lowest

AICc values (rather than AIC values) to account for the small sample size. For each genus, we

summarized the best-fitting models across 100 bootstrap trees, and the best-fitted model for the

genus was determined based on the models with highest proportion. For all genera, we found this

model was the same as the best-fitting model identified from the ML tree.

When the model incorporating hidden states and state-dependency (HiSSE rev or HiSSE

irr) provided the best fit for a genus, we assessed whether outcrossing consistently gave higher or

lower diversification rates than selfing at both hidden states or not. To do this, we classified the

rate estimates from each bootstrap tree into two scenarios: 1) diversification rates for state 0

(outcrossing) are either consistently higher or lower than for state 1 (selfing) in both hidden

states ($r_{0A} > r_{1A}$ and $r_{0B} > r_{0B}$, or $r_{0A} < r_{1A}$ and $r_{0B} < r_{0B}$); 2) diversification rates for state 0

(outcrossing) are higher than for state 1 (selfing) in one hidden state but lower in the other

hidden state ($r_{0A} > r_{1A}$ and $r_{0B} < r_{1B}$, or $r_{0A} < r_{1A}$ and $r_{0B} > r_{1B}$). We recorded the occurrence

of the two scenarios in 100 bootstrap trees.

Results

Data description

After filtering for suitable genera, our diversification analyses include 27 genera,

summarized in Table S2. Among these, 21 herbaceous genera are from the self-incompatibility

dataset, which includes 9 genera in Asteraceae, 4 in Brassicaceae, 4 in Solanaceae, 2 in

Bromeliaceae, 1 in Onagraceae, and 1 in Fabaceae. Four genera in Polemoniaceae are used based

on the pollination system dataset. The outcrossing rate dataset provides 2 gymnosperm genera

(*Picea*, *Pinus*) suitable for analyses. In *Oenothera*, some SC species are functionally asexual due

to a unique genetic system known as permanent translocation heterozygosity (PTH), which

nearly suppresses meiotic recombination and segregation (Holsinger & Ellstrand, 1984). These

PTH species are thus dropped from the phylogeny used for diversification analyses.

In total, the diversification analysis included 360 species in state 0 and 453 species in

state 1, with 19 genera having sampling fractions greater than 10% (Fig. S1). Two genera,

Solanum and *Oenothera*, are suitable for performing diversification analyses on the combination

of self-compatibility and life form states. For *Solanum*, the MuSSE analysis includes no SI

annual/short-lived species, 73 SI perennial species, 25 SC annual/short-lived species, and 105 SC

perennial species. After excluding PTH species, the analysis in *Oenothera* includes 9 SI

annual/short-lived species, 31 SI perennial species, 21 SC annual/short-lived species, and 11 SC

perennial species.

BiSSE analysis on mating system states

Out of the 27 genera, BiSSE MCMC samplings in four genera (*Gilia*, *Leptosiphon*,

Nicotiana, *Picea*) show high sensitivity to the choice of prior distributions or transition

reversibility in mating systems (Table S3). Dropping these genera, our BiSSE analyses include

23 genera, covering 316 species in state 0 and 331 species in state 1. Fig. 1 presents the

distribution $P(r_0 > r_1)$ and $P(\lambda_0 > \lambda_1)$ values obtained from the 100 bootstrap trees for each

genus (detailed values are summarized in Table S4).

The results in Fig. 1(A) suggest that across genera, outcrossing (state 0) lineages have

higher diversification rates than selfing (state 1) lineages, as $P(r_0 > r_1)$ values significantly

deviate above 0.5 (*P*=0.002; one-sample Wilcoxon rank-sum test). In contrast, outcrossing

lineages do not consistently speciate at higher or lower rates compared to selfing lineages, as

$P(\lambda_0 > \lambda_1)$ values do not significantly deviate from 0.5 (*P*=0.286; one-sample Wilcoxon rank-

sum test). The above results are robust to different values of f used in the prior distributions

during MCMC sampling (Fig. S2).

Overall, the proportion of MCMC steps supporting higher diversification rates for

outcrossing lineages, $P(r_0 > r_1)$, is positively correlated with that for speciation rates,

$P(\lambda_0 > \lambda_1)$ (*r* = 0.45; *P*=0.032), while negatively correlated with the proportion supporting

higher extinction rates, $P(\mu_0 > \mu_1)$ (*r* = -0.46; *P*=0.026). These results suggest that the higher

diversification rates inferred for outcrossing lineages reflect both higher rates of speciation and

lower rates of extinction. However, it should be noted that the inferred extinction rates may be

unreliable (Rabosky, 2010).

Even if mating systems have no impact on diversification rates, genera with more

outcrossing species tend to show higher $P(r_0 > r_1)$ values. However, we find no significant

correlation between $P(r_0 > r_1)$ values and the fraction of outcrossing species included in the

diversification analysis (*r*=0.15; *P*=0.500). This result suggests that the finding of higher $P(r_0 >$

$r_1)$ values for outcrossing lineages is unlikely to be because outcrossing species happen to be

overrepresented in our focal genera.

Figure 1. The proportion of the posterior probabilities from BiSSE MCMC sampling that

supports higher diversification rates and speciation rates for state 0 (outcrossing) than state 1

(selfing), assuming no transition from state 1 to state 0. Colors represent different characters

related to mating systems. The p-value from one-sample Wilcoxon rank-sum test is 0.002 for

panel (A), and 0.111 for panel (B).

Diversification analyses on the combination of self-compatibility and life form

We performed MuSSE MCMC analysis on the combination of self-compatibility and life

form states in two genera, *Solanum* and *Oenothera*. For both genera, $P(r_0 > r_1)$ values from the

BiSSE MCMC analysis are close to 1 (see Table S4), suggesting that SI lineages diversify at

higher rates than SC lineages. Additional MuSSE analysis shows that within SC lineages, SC

annual/short-lived lineages tend to diversify at much lower rates than SC perennial lineages.

Figure 2. Posterior distribution of the diversification rates, speciation rates and transition rates

for the combination of self-compatibility and life form states in *Solanum*. Results were obtained

from MuSSE MCMC simulations on 100 bootstrap trees. Results are robust to different values of

f used in the prior distribution (Fig. S3).

In *Solanum*, SC annual/short-lived lineages exhibit much lower diversification rates than

the other two states (Fig. 2A). This pattern appears to be driven by a combination of lower

speciation rates (Fig. 2B) and higher extinction rates (Fig. S3). In contrast, SC perennial lineages

display only slightly lower diversification rates than SI perennial lineages (Fig. 2A), which is

likely driven by slightly higher extinction rates (Fig. S3) instead of difference in speciation rates

(Fig. 2B). Additionally, transition rates from SI to SC are quite frequent among perennials (q_{24} in

Fig. 2C). For life form evolution, the results show that the transition rates from SC perennials to

SC annuals/short-lived species are rarely reversed (q_{34}, q_{43} in Fig. 2C).

In *Oenothera*, SC lineages show lower diversification and speciation rates than SI

lineages in both annual/short-lived and perennial states (Figs. 3A, 3B). However, the differences

in diversification rates between SI and SC lineages are much more pronounced in the

annual/short-lived state than in the perennial state (Fig. 3A). Similar to the findings in *Solanum*,

SC annual/short-lived lineages exhibit the lowest diversification rates (Fig. 3A), perhaps due to

elevated extinction rates (Fig. S4). However, among SI species, SI annual/short-lived lineages

have higher diversification rates than SI perennials (Fig. 3A), mainly driven by higher speciation

rates (Figs. 3B, S4). Transition rates from SI to SC are higher in annual/short-lived lineages than

in perennial lineages (Fig. 3C). For transitions between life form states, the rate of transition

from annuals/short-lived to perennials is much higher than in the reverse direction in SI lineages

(compared q_{12}, q_{21} in Fig. 3D). However, among SC lineages, the rate of transition toward

perennials is greatly reduced, making the transition rates between life form states in both

directions comparable (compared q_{34}, q_{43} in Fig 3D).

Figure 3. Posterior distribution of the diversification rates, speciation rates and transition rates for the combination of self-compatibility and life form states in *Oenothera*. Results were obtained from MuSSE MCMC simulations on 100 bootstrap trees. Results are robust to different values of f used in the prior distribution (Fig. S4).

Diversification analysis with hidden states

Results from the MuSSE analysis indicate that the effects of selfing on lineage diversification may depend on the states of other characters. This conjecture is also supported by the HiSSE analysis. Out of the 27 focal genera, models incorporating hidden states are better fitted than models assuming no hidden states in 19 genera, as shown in Fig. 4A (detailed results are summarized in Table S5). Models assuming no backward transition from selfing to outcrossing (BiSSE irr, HiSSE irr) are best fitted in 12 genera (Fig. 4A). Models that allow reversible transitions between mating system states (BiSSE rev, HiSSE rev) are best fitting for the two genera from the outcrossing rate dataset (dark grey in Fig. 4A), suggesting that changes in the rate of selfing tends to be reversible. However, models allowing reversible transitions were also best fitted in 7 genera from the self-compatibility dataset. This may be due to overfitting and

may not accurately reflect reality, as most of these genera belong to Asteraceae and Brassicaceae,

where self-incompatibility systems are maintained by self-recognition mechanisms that should

rarely allow transitions from SC back to SI (Fujii et al., 2016). Models incorporating hidden

states and state-dependent diversification (HiSSE rev, HiSSE irr) are the best-fitting for 17

genera. However, no significant pattern is found regarding whether outcrossing lineages

consistently diversify at higher or lower rates compared to selfing lineages across both hidden

states (Fig. 4B).

Figure 4. (A) Distribution of the best-fitting model across 27 genera. (B) The fraction of 100

bootstrap trees of each genus for which the estimated diversification rates from the *hisse* package

for state 0 are consistently higher or lower than state 1 in both hidden states A and B. The

distribution of the fractions does not significantly deviate from 0.5 in one-sample Wilcoxon rank-

sum test, no matter whether the two HiSSE models were analyzed separately ($P = 0.478$ for

HiSSE irr; $P = 1.000$ for HiSSE rev) or combined ($P = 0.505$). Detailed results for each genus

are presented in Table S5.

Discussion

The current study investigated the effects of mating system states on lineage

diversification rates by performing phylogenetic-based diversification analyses to a dozen of

genera. The results show that outcrossing lineages tend to have higher net diversification rates

than selfing lineages, which is suggested to be attributed to both higher speciation rates and

lower extinction rates. Nevertheless, mating systems do not exhibit consistent effects on

speciation rates; selfing lineages display higher speciation rates in some genera while exhibiting

lower rates in others. This result aligns with the mixed outcomes observed in previous analyses.

This variability may stem from the fact that speciation events involve various mechanisms

(Coyne & Orr, 2004), and selfing can have complex effects on these mechanisms, as discussed in

the introduction, and reviewed by Marie-Orleach et al. (2024).

Most previous analyses were conducted at the family level, but the current genus-level

analyses reveal that within the same family, the effects of mating systems on diversification rates

can vary across genera. For instance, family-level analyses in Solanaceae reported lower

diversification rates in self-compatible (SC) lineages compared to self-incompatible (SI) lineages

(Goldberg et al., 2010). While similar results are observed in the genus *Solanum*, self-

compatibility shows no significant effects on diversification in *Nicotiana*. Since one-third of the

species included in Goldberg et al. (2010) are from *Solanum*, their results may be skewed by the

patterns observed in this genus.

We propose that highly selfing lineages tend to have much higher extinction rates than

mixed-mating lineages. Although we are not able to directly test this hypothesis due to

insufficient data, we find SC annual/short-lived lineages exhibit much lower diversification rates

than SC perennials in *Solanum*. Also, in *Oenothera*, diversification rate differences between SI

and SC lineages are more prominent in annuals/short-lived than in perennials. These results

indirectly support our hypothesis as annuals are more likely to be highly selfing than perennials

(Aarssen, 2000; Ma et al., 2020). A direct test of this hypothesis would require a diversification

analysis across a full range of selfing rates once outcrossing rates are measured in more species.

Additionally, this hypothesis can be indirectly examined by conducting diversification analyses

of quantitative traits related to selfing rates, such as anther-stigma distance and flower size

(Thomann et al., 2013).

As explained in the introduction, the above results may help explain the enigma of the

common occurrence of mixed-mating systems (Goodwillie et al., 2005). Additionally, our

analyses indirectly support the model prediction that mixed-mating systems tend to be

evolutionarily unstable (Lloyd, 1979; Lande & Schemske, 1985). Specifically, we find that

transition rates from SC perennials, which tend to exhibit mixed mating, to SC annuals/short-

lived plants, which are likely to be highly selfing, are higher than the reverse direction. However,

we also find relatively high transition rates from SI perennials to SC perennials, which suggests

that the prevalence of mixed mating systems may also be partly contributed by frequent

transitions from outcrossing lineages.

The analyses incorporating life form states suggest that variation in diversification rates

under different mating systems may often be partly driven by unmeasured factors associated with

mating system transitions. Indeed, we find that SSE models incorporating hidden states are better

fitted than those without hidden states for a majority of the studied genera. This may be partly

because transitions from outcrossing to selfing are often triggered by environmental changes,

such as colonization and pollen limitation (Busch & Schoen, 2008; Eckert et al., 2010; Pannell,

2016; Encinas-Viso et al., 2020). Nevertheless, how the impacts of selfing on diversification

rates depend on hidden states is not consistent across genera. In some genera, selfing consistently

promotes or inhibits diversification for both hidden states, but in other genera, selfing has

opposite effects on diversification in different hidden states.

Our studies also support the model prediction that self-fertilization tends to promote

evolution towards shorter lifespans (Lesaffre & Billiard, 2020). Among SC speciess in *Solanum*,

transitions from perennials to annuals/short-lived lineages are found to be nearly unidirectional.

Also, in *Oenothera*, although transition rates from perennials to annuals/short-lived are not

elevated in SC lineages, the rate of transitions from annuals/short-lived to perennials is greatly

reduced in SC lineages compared to SI lineages.

In short, the current analyses suggest that selfing lineages have lower diversification rates

and higher extinction rates, which may be mainly contributed in highly selfing, instead of mixed-

mating lineages. However, it should be emphasized that our analyses only reveal a correlation,

instead of causation, between selfing and low diversification rates. In fact, the impacts of selfing

on diversification are likely context-dependent, potentially interacting with other ecological or

genetic factors.

Acknowledgments

I would like to thank Stephen Wright and Mark Hibbins for comments and suggestions on the

analyses and a previous version of the manuscript. K.X. was supported by the EEB Postdoctoral

Fellowship offered by the Department of Ecology & Evolutionary Biology at the University of

Toronto.

Competing interests

The author declares no competing interests.

 Author contributions

 K.X. conceived and designed the study, conducted the analysis and drafted the article.

 Data availability

 Mating system character states, supermatrices and phylogenies of each genus are available at

 https://doi.org/10.5281/zenodo.13265323

 References

Aarssen, L. W. (2000). Why are most selfers annuals? A new hypothesis for the fitness benefit of selfing. *Oikos*, *89*(3), 606-612.

Arunkumar, R., Ness, R. W., Wright, S. I., & Barrett, S. C. (2015). The evolution of selfing is accompanied by reduced efficacy of selection and purging of deleterious mutations. *Genetics*, *199*(3), 817-829.

Barrett, S. C. (2008). Major evolutionary transitions in flowering plant reproduction: an overview. *International Journal of Plant Sciences*, *169*(1), 1-5.

Barrett, S. C. (2013). The evolution of plant reproductive systems: how often are transitions irreversible?. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1765), 20130913.

Barringer, B. C. (2007). Polyploidy and self-fertilization in flowering plants. *American Journal of Botany*, *94*(9), 1527-1533.

Beaulieu, J. M., & O'Meara, B. C. (2016). Detecting hidden diversification shifts in models of trait-dependent speciation and extinction. *Systematic biology*, *65*(4), 583-601.

Bennett, D. J., Hettling, H., Silvestro, D., Zizka, A., Bacon, C. D., Faurby, S., Vos, R. A., & Antonelli, A. (2018). phylotaR: An automated pipeline for retrieving orthologous DNA sequences from GenBank in R. *Life*, *8*(2), 20.

Bürkli, A., Sieber, N., Seppälä, K., & Jokela, J. (2017). Comparing direct and indirect selfing rate estimates: when are population-structure estimates reliable?. *Heredity*, *118*(6), 525-533.

Busch, J. W., & Delph, L. F. (2012). The relative importance of reproductive assurance and automatic selection as hypotheses for the evolution of self-fertilization. *Annals of botany*, *109*(3), 553-562.

Busch, J. W., & Schoen, D. J. (2008). The evolution of self-incompatibility when mates are limiting. *Trends in plant science*, *13*(3), 128-136.

Caballero, A., & Hill, W. G. (1992). Effects of partial inbreeding on fixation rates and variation of mutant genes. *Genetics*, *131*(2), 493-507.

Cascante-Marín, A., & Núñez-Hidalgo, S. (2023). A Review of Breeding Systems in the

Pineapple Family (Bromeliaceae, Poales). *The Botanical Review*, *89*(3), 308-329.

Castillo, D. M., Gibson, A. K., & Moyle, L. C. (2016). Assortative mating and self-fertilization

differ in their contributions to reinforcement, cascade speciation, and diversification. *Current*

Zoology, *62*(2), 169-181.

Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in

phylogenetic analysis. *Molecular biology and evolution*, *17*(4), 540-552.

Charlesworth, D., Morgan, M. T., & Charlesworth, B. (1993). Mutation accumulation in finite

outbreeding and inbreeding populations. *Genetics Research*, *61*(1), 39-56.

Clo, J., & Opedal, Ø. H. (2021). Genetics of quantitative traits with dominance under stabilizing

and directional selection in partially selfing species. *Evolution*, *75*(8), 1920-1935.

Clo, J., Gay, L., & Ronfort, J. (2019). How does selfing affect the genetic variance of

quantitative traits? An updated meta-analysis on empirical results in angiosperm

species. *Evolution*, *73*(8), 1578-1590.

Clo, J., Ronfort, J., & Abu Awad, D. (2020). Hidden genetic variance contributes to increase the

short-term adaptive potential of selfing populations. *Journal of Evolutionary Biology*, *33*(9),

1203-1215.

Coyne J. A., & Orr H. A. (2004). Speciation. Sinauer Associates, Sunderland, MA.

Cutter, A. D. (2019). Reproductive transitions in plants and animals: selfing syndrome, sexual

selection and speciation. *New Phytologist*, *224*(3), 1080-1094.

David, P., Pujol, B., Viard, F., Castella, V., & Goudet, J. (2007). Reliable selfing rate estimates

from imperfect population genetic data. *Molecular ecology*, 16(12), 2474-2487.

de Vos, J. M., Hughes, C. E., Schneeweiss, G. M., Moore, B. R., & Conti, E. (2014). Heterostyly

accelerates diversification via reduced extinction in primroses. *Proceedings of the Royal Society*

B: Biological Sciences, *281*(1784), 20140075.

Delaney, L. E., & Igić, B. (2022). The phylogenetic distribution and frequency of self-

incompatibility in Fabaceae. *International Journal of Plant Sciences*, *183*(1), 30-42.

Dietrich, W., & Wagner, W. L. (1988). Systematics of Oenothera section Oenothera subsection

Raimannia and subsection Nutantigemma (Onagraceae). *Systematic Botany Monographs*, 1-91.

Duminil, J., Hardy, O. J., & Petit, R. J. (2009). Plant traits correlated with generation time

directly affect inbreeding depression and mating system and indirectly genetic structure. *BMC*

Evolutionary Biology, *9*, 1-14.

Eckert, C. G., Kalisz, S., Geber, M. A., Sargent, R., Elle, E., Cheptou, P. O., ... & Winn, A. A.

(2010). Plant mating systems in a changing world. *Trends in ecology & evolution*, *25*(1), 35-43.

Edler, D., Klein, J., Antonelli, A., & Silvestro, D. (2021). raxmlGUI 2.0: a graphical interface

and toolkit for phylogenetic analyses using RAxML. *Methods in Ecology and Evolution*, *12*(2),

373-377.

Encinas-Viso, F., Young, A. G., & Pannell, J. R. (2020). The loss of self-incompatibility in a

range expansion. *Journal of Evolutionary Biology*, *33*(9), 1235-1244.

Escobar, J. S., Cenci, A., Bolognini, J., Haudry, A., Laurent, S., David, J., & Glémin, S. (2010).

An integrative test of the dead-end hypothesis of selfing evolution in Triticeae

(Poaceae). *Evolution*, *64*(10), 2855-2872.

Evans, M. E., Hearn, D. J., Theiss, K. E., Cranston, K., Holsinger, K. E., & Donoghue, M. J.

(2011). Extreme environments select for reproductive assurance: evidence from evening

primroses (Oenothera). *New Phytologist*, *191*(2), 555-563.

FitzJohn, R. G. (2012). Diversitree: comparative phylogenetic analyses of diversification in

R. *Methods in Ecology and Evolution*, *3*(6), 1084-1092.

FitzJohn, R. G., Maddison, W. P., & Otto, S. P. (2009). Estimating trait-dependent speciation and

extinction rates from incompletely resolved phylogenies. *Systematic biology*, *58*(6), 595-611.

Freyman, W. A., & Höhna, S. (2019). Stochastic character mapping of state-dependent

diversification reveals the tempo of evolutionary decline in self-compatible Onagraceae

lineages. *Systematic Biology*, *68*(3), 505-519.

Fujii, S., Kubo, K. I., & Takayama, S. (2016). Non-self-and self-recognition models in plant self-

incompatibility. *Nature plants*, *2*(9), 1-9.

Gagnon, E., Hilgenhof, R., Orejuela, A., McDonnell, A., Sablok, G., Aubriot, X., Giacomin, L.,

Gouvêa, Y., Bragionis, T., Stehmann, J.R., Bohs, L., Dodsworth, S., Martine, C., Poczai, P., &

Särkinen, T. (2022). Phylogenomic discordance suggests polytomies along the backbone of the

large genus Solanum. *American Journal of Botany*, *109*(4), 580-601.

Gamisch, A., Fischer, G. A., & Comes, H. P. (2015). Multiple independent origins of auto-

pollination in tropical orchids (Bulbophyllum) in light of the hypothesis of selfing as an

evolutionary dead end. *BMC Evolutionary Biology*, *15*, 1-18.

Glémin, S. (2007). Mating systems and the efficacy of selection at the molecular

level. *Genetics*, *177*(2), 905-916.

Glémin, S., & Ronfort, J. (2013). Adaptation and maladaptation in selfing and outcrossing

species: new mutations versus standing variation. *Evolution*, *67*(1), 225-240.

Goldberg, E. E., Kohn, J. R., Lande, R., Robertson, K. A., Smith, S. A., & Igić, B. (2010).

Species selection maintains self-incompatibility. *Science*, *330*(6003), 493-495.

Goodwillie, C., Kalisz, S., & Eckert, C. G. (2005). The evolutionary enigma of mixed mating

systems in plants: occurrence, theoretical explanations, and empirical evidence. *Annu. Rev. Ecol.*

Evol. Syst., *36*, 47-79.

Grossenbacher, D. L., Brandvain, Y., Auld, J. R., Burd, M., Cheptou, P. O., Conner, J. K., Grant,

A. G., Hovick, S. M., Pannell, J. R., Pauw, A. & Goldberg, E. E. (2017). Self-compatibility is

over-represented on islands. *New Phytologist*, *215*(1), 469-478.

Hartfield, M., Bataillon, T., & Glémin, S. (2017). The evolutionary interplay between adaptation

and self-fertilization. *Trends in Genetics*, *33*(6), 420-431.

Helmstetter, A. J., Zenil-Ferguson, R., Sauquet, H., Otto, S. P., Méndez, M., Vallejo-Marin, M.,

Schönenberger, J., Burgarella, C., Anderson, B., De Boer, H. and Glémin, S., & Käfer, J. (2023).

Trait-dependent diversification in angiosperms: Patterns, models and data. *Ecology*

Letters, *26*(4), 640-657.

Hilgenhof, R., Gagnon, E., Knapp, S., Aubriot, X., Tepe, E. J., Bohs, L., Giacomin, L. L.,

Gouvêa, Y. F., Martine, C.T., Orejuela, A. & Särkinen, T. (2023). Morphological trait evolution in

Solanum (Solanaceae): Evolutionary lability of key taxonomic characters. *Taxon*, *72*(4), 811-847.

Holsinger, K. E., & Ellstrand, N. C. (1984). The evolution and ecology of permanent

translocation heterozygotes. *The American Naturalist*, *124*(1), 48-71.

Igic, B., & Busch, J. W. (2013). Is self-fertilization an evolutionary dead end?. *New*

Phytologist, *198*(2), 386-397.

Igic, B., Lande, R., & Kohn, J. R. (2008). Loss of self-incompatibility and its evolutionary

consequences. *International Journal of Plant Sciences*, *169*(1), 93-104.

Jones, O. R., & Wang, J. (2010). COLONY: a program for parentage and sibship inference from

multilocus genotype data. *Molecular ecology resources*, *10*(3), 551-555.

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Molecular biology and evolution*, *30*(4), 772-780.

Krakos, K. N., Johnson, M. G., Hoch, P. C., Wagner, W. L., Huang, P., & Raven, P. H. (2022).

Molecular Phylogenetics Reveals Multiple Transitions to Self-Compatibility in a Primary

Subclade of Oenothera (Onagraceae) 1. *Annals of the Missouri Botanical Garden*, *107*(1), 405-

421.

Lande, R., & Schemske, D. W. (1985). The evolution of self-fertilization and inbreeding

depression in plants. I. Genetic models. *Evolution*, *39*(1), 24-40.

Landis, J. B., Bell, C. D., Hernandez, M., Zenil-Ferguson, R., McCarthy, E. W., Soltis, D. E., &

Soltis, P. S. (2018). Evolution of floral traits and impact of reproductive mode on diversification

in the phlox family (Polemoniaceae). *Molecular Phylogenetics and Evolution*, *127*, 878-890.

Lesaffre, T., & Billiard, S. (2020). The joint evolution of lifespan and self-fertilization. *Journal*

of Evolutionary Biology, *33*(1), 41-56.

Lloyd, D. G. (1979). Some reproductive factors affecting the selection of self-fertilization in

plants. *The American Naturalist*, *113*(1), 67-79.

Ma, Y., Barrett, S. C., Wang, F. Y., Deng, J. C., & Bai, W. N. (2021). Do annual and perennial

populations of an insect-pollinated plant species differ in mating system?. *Annals of*

Botany, *127*(7), 853-864.

Maddison WP, Midford PE, Otto SP. 2007. Estimating a binary character's effect on speciation

and extinction. *Systematic Biology* 56: 701–710.

Marie-Orleach, L., Brochmann, C., & Glémin, S. (2022). Mating system and speciation I:

Accumulation of genetic incompatibilities in allopatry. *PLoS Genetics*, *18*(12), e1010353.

Marie-Orleach, L., Glémin, S., Brandrud, M. K., Brysting, A. K., Gizaw, A., Gustafsson, A. L.

S., Rieseberg, L. H., Brochmann, C., & Birkeland, S. (2024). How Does Selfing Affect the Pace

and Process of Speciation?. *Cold Spring Harbor Perspectives in Biology*, a041426.

Molloy, E. K., & Warnow, T. (2018). To include or not to include: the impact of gene filtering on

species tree estimation methods. *Systematic biology*, *67*(2), 285-303.

Ness, R. W., Siol, M., & Barrett, S. C. (2012). Genomic consequences of transitions from cross-

to self-fertilization on the efficacy of selection in three independently derived selfing

plants. *BMC genomics*, *13*, 1-12.

Nordborg, M., Charlesworth, B., & Charlesworth, D. (1996). The effect of recombination on

background selection. *Genetics Research*, *67*(2), 159-174.

Pannell, J. R. (2016). Evolution of the mating system in colonizing plants. *Invasion Genetics:*

The Baker and Stebbins Legacy, 57-80.

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics*, *35*(3), 526-528.

Rabosky, D. L. (2010). Extinction rates should not be estimated from molecular phylogenies.

Evolution, *64*(6), 1816-1824.

Rabosky, D. L., & Goldberg, E. E. (2015). Model inadequacy and mistaken inferences of trait-

dependent speciation. *Systematic biology*, *64*(2), 340-355.

Raduski, A. R., Haney, E. B., & Igić, B. (2012). The expression of self-incompatibility in

angiosperms is bimodal. *Evolution*, *66*(4), 1275-1283.

Rausch, J. H., & Morgan, M. T. (2005). The effect of self-fertilization, inbreeding depression,

and population size on autopolyploid establishmen. *Evolution*, *59*(9), 1867-1875.

Ritland, K. (1990). A series of FORTRAN computer programs for estimating plant mating

systems. *J. Hered*. 81, 236–237.

Ritland, K. (2002). Extensions of models for the estimation of mating systems using n

independent loci. *Heredity*, *88*(4), 221-228.

Ritland, K., & Jain, S. (1981). A model for the estimation of outcrossing rate and gene

frequencies using n independent loci. Heredity, 47(1), 35-52.

Sabath, N., Goldberg, E. E., Glick, L., Einhorn, M., Ashman, T. L., Ming, R., Otto, S. P., Vamosi,

J. C., & Mayrose, I. (2016). Dioecy does not consistently accelerate or slow lineage

diversification across multiple genera of angiosperms. *New Phytologist*, *209*(3), 1290-1300.

Sanderson, M. J. (2002). Estimating absolute rates of molecular evolution and divergence times:

a penalized likelihood approach. *Molecular biology and evolution*, *19*(1), 101-109.

Sicard, A., & Lenhard, M. (2011). The selfing syndrome: a model for studying the genetic and

evolutionary basis of morphological adaptation in plants. *Annals of botany*, *107*(9), 1433-1443.

Soltis, D. E., Mort, M. E., Latvis, M., Mavrodiev, E. V., O'Meara, B. C., Soltis, P. S., Burleigh, J.

G., & Rubio de Casas, R. (2013). Phylogenetic relationships and character evolution analysis of

Saxifragales using a supermatrix approach. *American journal of botany*, *100*(5), 916-929.

Thomann, M., Imbert, E., Devaux, C., & Cheptou, P. O. (2013). Flowering plants under global

pollinator decline. *Trends in plant science*, *18*(7), 353-359.

Wagner, W. L., Hoch, P. C., & Raven, P. H. (2007). Revised classification of the

Onagraceae. *Systematic Botany Monographs*.

Wang, X. J., Barrett, S. C., Zhong, L., Wu, Z. K., Li, D. Z., Wang, H., & Zhou, W. (2021). The

genomic selfing syndrome accompanies the evolutionary breakdown of heterostyly. *Molecular*

biology and evolution, *38*(1), 168-180.

Whitehead, M. R., Lanfear, R., Mitchell, R. J., & Karron, J. D. (2018). Plant mating systems

often vary widely among populations. *Frontiers in Ecology and Evolution*, 6, 38.

Wright, S. I., Ness, R. W., Foxe, J. P., & Barrett, S. C. (2008). Genomic consequences of

outcrossing and selfing in plants. *International Journal of Plant Sciences*, *169*(1), 105-118.

Xu, K. (2022). Mutation accumulation in inbreeding populations under evolution of the selfing

rate. *Journal of Evolutionary Biology*, *35*(1), 23-39.

Xu, K. (2023a). Effects of selfing on multi-step adaptation. *Evolution*, *77*(2), 482-495.

Xu, K. (2023b) Population rescue through an increase of the selfing rate under pollen limitation:

plasticity vs. evolution. *The American Naturalist*, *202*(3), 337-350.