Assessing the combined effects of forest management and climate change

on carbon and water fluxes in European beech forests

Vincenzo Saponaro[1,2*], Miquel De Cáceres[3], Daniela Dalmonech[1,4], Ettore D'Andrea[4,5],

Elia Vangi[1], Alessio Collalti[1,4]

Affiliations:

1. Forest Modelling Lab., Institute for Agriculture and Forestry Systems in the
Mediterranean, National Research Council of Italy (CNR-ISAFOM), Perugia, Italy;
2. Department for Innovation in Biological, Agri-Food and Forest Systems (DIBAF),
University of Tuscia, Viterbo, Italy;
3. CREAF, E08193 Bellaterra (Cerdanyola del Vallès), Catalonia, Spain
4. National Biodiversity Future Center (NBFC), Palermo, Italy
5. Research Institute on Terrestrial Ecosystems, National Research Council of Italy (CNR-
IRET), Porano, Italy

*_Corresponding author_: vincenzo.saponaro@isafom.cnr.it

Abstract

The consequences of climate change continue to threaten European forests,

particularly for species located at the edges of their latitudinal and altitudinal ranges.

While extensively studied in Central Europe, European beech forests require further

investigation to understand how climate change will affect these ecosystems in

Mediterranean areas. Proposed silvicultural options increasingly aim at sustainable

management to reduce biotic and abiotic stresses and enhance these forest

ecosystems' resistance and resilience mechanisms. Process-based models (PBMs) can

help us to simulate such phenomena and capture early stress signals while considering

the effect of different management approaches. In this study, we focus on estimating

sensitivity of two state-of-the-art PBMs forest models by simulating carbon and water

fluxes at the stand level to assess productivity changes and feedback resulting from

different climatic forcings as well as different management regimes. We applied the 3D-CMCC-FEM and MEDFATE forest models for carbon (C) and water (H_2O) fluxes in in two sites of the Italian peninsula, Cansiglio in the north and Mongiana in the south, under managed vs. unmanaged scenarios and under current climate and different climatic scenarios (RCP4.5 and RCP8.5). To ensure confidence in the models' results, we preliminary evaluated their performance in simulating C and H_2O flux in three additional beech forests of the FLUXNET network along a latitudinal gradient spanning from Denmark to central Italy. The 3D-CMCC-FEM model achieved R^2 values of 0.83 and 0.86 with RMSEs of 2.53 and 2.05 for C and H_2O fluxes, respectively. MEDFATE showed R^2 values of 0.76 and 0.69 with RMSEs of 2.54 and 3.01. At the Cansiglio site in northern Italy, both models simulated a general increase in C and H_2O fluxes under the RCP8.5 climate scenario compared to the current climate. Still, no benefit in managed plots compared to unmanaged ones, as the site does not have water availability limitations, and thus, competition for water is low. At the Mongiana site in southern Italy, both models predict a decrease in C and H_2O fluxes and sensitivity to the different climatic forcing compared to the current climate; and an increase in C and H_2O fluxes when considering specific management regimes compared to unmanaged scenarios. Conversely, under unmanaged scenarios plots are simulated to experience first signals of mortality prematurely due to water stress (MEDFATE) and carbon starvation (3D-CMCC-FEM) scenarios. In conclusion, while management interventions may be considered a viable solution for the conservation of beech forests under future climate conditions at moister sites like Cansiglio, in drier sites like Mongiana conservation may not lie in management interventions alone.

Keywords: Climate change sensitivity, *Fagus sylvatica* L., Forest management sensitivity, Carbon fluxes, Water fluxes, Stress mitigation, Process-based models.

1. Introduction

Predicting the future evolution of European forests is essential to continue to benefit from the ecosystem services they provide for human well-being. Forests offer, for instance, climate change mitigation through their ability to store atmospheric carbon dioxide in biomass and soil (Augusto and Bo, 2022; Pan et al., 2024). In 2020, the European Green Deal prioritized the vital role of forests and the forestry sector in attaining sustainability objectives, such as promoting sustainable forest management, enhancing forest resilience, and climate change mitigation (European Commission, 2021). Technological advances and studies of forest ecosystem responses to management practices continue to promote the evolution of strategies that maintain or enhance forest ecosystem services, such as promoting biological diversity, water resources, soil protection, or carbon sequestration (Pukkala, 2016). Different forest management systems have been adopted in Europe over the years (e.g., clear-cutting or shelterwood) depending, among others, on the wood product desired, the stand age, and structure (Brunet et al., 2010).

Forest management can be a key element in mitigating the effects of climate warming, maintaining the current primary productivity and the current distribution of tree species, or altering forest composition to promote more suited and productive species (Nolè et al., 2015; Bosela et al., 2016). Indeed, the carbon sequestration capacity and productivity of forests are dependent, primarily, on species composition, site conditions as well as on stand age (Rötzer et al., 2010; Vangi et al., 2024a, b), which are affected by past and present forest management activities. According to Collalti et al. (2018) and Dalmonech et al. (2022), monospecific forests in Europe would appear unable to further increase current rates of carbon storage and biomass production under future climate scenarios, considering current management practices, but at the same time demonstrating that managing under Business as Usual (BAU) practices still

allows forests to accumulate biomass at higher rates compared to stands left to

develop undisturbed.

European beech (*Fagus sylvatica* L.) is an important deciduous tree species widely

distributed in Europe, from southern Scandinavia to Sicily and Spain to northwest

Turkey (Durrant et al., 2016). In Italy, according to the National Forest Inventory (INFC,

2015), beech forests cover a total area of 1,053,183 hectares, accounting for about 11.7%

of the country's overall forested land. European beech forests demonstrate

susceptibility to temperature and precipitation fluctuations. For instance, a warmer

environment and less precipitation are forcing shifts in distribution area or the onset of

loss of canopy greenness (Axer et al., 2021; Noce et al., 2017, 2023; Zuccarini et al.,

2023; Rezaie et al., 2018). According to Skrk et al. (2023), the decline in growth of the

beech forests primarily occurs in the dry and warm marginal conditions prevalent near

the geographical edge of its distribution with a sub-Mediterranean climatic regime,

posing a threat to the survival of beech populations in those areas. However, tree ring

analyses have also revealed an unexpected increase in growth in the south

Mediterranean region of Albania and Macedonia beech forests at the end of the 20[th]

century, challenging the presumed decline of forest ecosystems due to drought (Tegel

et al., 2014). Puchi et al. (2024) additionally shed light on the susceptibility to extreme

drought events of beech forests found at higher latitudes compared to those found at

lower latitudes in the Italian peninsula by highlighting an increase, for the latter, in

growth related to the abundance of precipitation. In this context, it is important to

minimize the uncertainty surrounding the response of the carbon, water, and energy

cycles within beech forest ecosystems, especially as they have been shown to adapt to

varying environmental drivers (Deb Burman et al., 2024).

Process-based models (PBMs) are useful tools for studying forest dynamics, as well as

water (H_2O) and carbon (C) use efficiency, and carbon stocks as key variables of forest

mitigation potential (Vacchiano et al., 2012; Pilli et al., 2022; Testolin et al., 2023;

Morichetti et al., 2024). Forest modelling has been widely used by forest ecologists for

tackling numerous applied research questions, and the field is continuously evolving to

improve process representation to achieve higher realism and predictive capacity

under warmer climate and forest management scenarios (Riviere et al., 2020; Kimmins

et al., 2008; Nolè et al., 2013; Maréchaux et al., 2021). By comparing the predictive

performance of different models under current environmental conditions, it is possible

to gain confidence in their predictions of future trends and make informed decisions in

forest ecosystem management and planning processes (Huber et al., 2013; Mahnken et

al., 2022).

The main goal of the present study is to evaluate the impact of forest management

regimes and climate change scenarios on European beech forests using two state-of-

the-science PBMs: 3D-CMCC-FEM (Collalti et al., 2014) and MEDFATE (De Cáceres et

al., 2023). More specifically, the study aims to provide deeper insights into the carbon

(C) and water (H_2O) fluxes of this species under varying management practices and

changing environmental conditions. The MEDFATE model is capable of simulating the

complex water dynamics linking the soil-vegetation-atmosphere continuum. The

performance of MEDFATE in simulating soil moisture dynamics and plant transpiration

has been extensively evaluated across various scales and different stand structures,

particularly in Mediterranean environments (De Cáceres et al., 2015, 2021; Sánchez-

Dávila et al., 2024). Complementarily, the 3D-CMCC-FEM model has been extensively

validated and shown to effectively capture the spatial and temporal variability of

carbon and water fluxes, while accounting for ecological heterogeneity and integrating

forest management practices across a wide range of scales (Collalti et al., 2018;

Dalmonech et al., 2022, 2024; Mahnken et al., 2022).

Since the study sites vary in terms of environmental factors that can affect gross

primary productivity (GPP), as well as latent heat (LE), which are the two variables

considered in this analysis, the use of two PBMs can provide the highest reliability in

capturing the complex dynamics of these variables under diverse environmental

conditions. By leveraging the strengths of both models, we can achieve a more robust

and comprehensive understanding of C and H_2O across the different sites. Specifically,

we tested: (i) to what extent different forest management options can influence C and

H_2O fluxes under the present-day climate; and, (ii) how harsher climate conditions may

affect the C and H_2O fluxes under different management options. To gain confidence

on the models predictive capacity we preliminary parameterized and evaluated

models' performances for C and H_2O fluxes at three forest stands dominated by beech

forests: the Sorø (DK-Sor), Hesse (FR-Hes), and Collelongo (IT-Col) sites, which are

included in the PROFOUND Database (PROFOUND DB) (Reyer et al., 2020a, b) and

makes part of the FLUXNET Network (Pastorello et al., 2020). To address the questions,

we assessed the C and H_2O fluxes at two target and independent beech forest sites in

Italy (Cansiglio and Mongiana) by simulating their development under various

management options and evaluating their (model) sensitivity to current and more

severe climate conditions.

2. Material and Methods
2.1 3D-CMCC-FEM model

The 3D-CMCC-FEM v.5.6 ('*Three-Dimensional – Coupled Model Carbon Cycle –*

Forest Ecosystem Module') (Collalti et al., 2024, and references therein; Marconi et al.,

2017; Dalmonech et al., 2022, 2024; Vangi et al., 2024a,b; Morichetti et al., 2024) is a C-

based, eco-physiological, biogeochemical and biophysical model. The model simulates

C and H_2O fluxes occurring within forest ecosystems daily, monthly, or annually,

depending on the processes to simulate, with a common spatial scale of one hectare (Collalti et al., 2016). Photosynthesis is simulated using the biochemical model of Farquhar–von Caemmerer–Berry (Farquhar et al., 1980), integrating the sunlit and shaded leaves of the canopy (De Pury and Farquhar, 1997). For the temperature dependence of the Michaelis-Menten coefficient for Rubisco and the CO_2 compensation point without mitochondrial respiration, the model adopts the parameterization described in Bernacchi et al. (2001, 2003). The net balance at the autotrophic level is represented by net primary production in eq 1:

$$NPP = GPP - R_a \qquad (1)$$

where R_a includes both maintenance respiration (R_m) and growth respiration (R_g). When R_m exceeds GPP, resulting in a negative NPP, the trees utilize their non-structural carbon reserves (NSC) (i.e., soluble sugars and starch, undistinguished) to meet the carbon demand (Merganičová et al., 2019; Collalti et al., 2020a). In deciduous trees, NSC is used to create new leaves during the bud-burst phase, replenishing during the growing season under favourable photosynthetic conditions, and finally remobilising to other tissues to prepare trees for dormancy at the end of the growth phase. The model assumes that NSC reserves are actively mobilized to meet metabolic demands during periods of stress or carbon deficits, such as drought. For instance, during periods of negative carbon balance, the model allocates stored NSC to sustain key physiological processes (e.g., maintenance respiration, leaf and fine root formation). The allocation scheme ensures that NSC replenishment is prioritized before supporting growth demands (i.e., wood growth), consistent with evidence showing that carbon flows are first directed toward restoring NSC reserves until critical thresholds are reached (Hartmann & Trumbore, 2016). Replenishment of non-structural carbon reserves is

essential to achieve the minimum safety threshold (i.e., 11% of sapwood dry mass for

deciduous trees; Schwalm and Ek, 2004). Failure to meet these thresholds may trigger

at first remobilization from leaves and fine root and subsequently to defoliation

mechanisms, while complete depletion of reserves (e.g., during prolonged stress

periods) could lead to the death of the entire cohort of trees through carbon

starvation.In 3D-CMCC-FEM stomatal conductance g_s is calculated using the Jarvis

equation (Jarvis, 1976). The equation includes a species-specific parameter $g_{s\,max}$ (i.e.,

maximum stomatal conductance) controlled by factors such as light, atmospheric CO_2

concentration, air temperature, soil water content, vapour pressure deficit (VPD), and

stand age (Collalti et al., 2019). According to Waring and Running (2007) and Monteith

and Unsworth (2008), the Penman-Monteith equation is used to calculate the latent

heat (LE) fluxes of evaporation as a function of incoming radiation, VPD, and

conductances at a daily scale, summing up the canopy, soil, and snow (if any) latent

heat flux expressed as $W\,m^{-2}$ or $MJ\,m^{-2}\,time^{-1}$

The 3D-CMCC-FEM accounts for forest stand dynamics, including growth, competition

for light, and tree mortality under different climatic conditions, considering both the

CO_2 fertilization effects and temperature acclimation (Collalti et al., 2018, 2019; Kattge

and Knorr, 2007). Several mortality routines are considered in the model, such as age-

dependent mortality, background mortality (stochastic mortality), self-thinning

mortality, and the aforementioned mortality due to carbon starvation. In addition to

mortality, biomass removal in 3D-CMCC-FEM results from forest management

practices, such as thinning and final harvest (Collalti et al., 2018; Dalmonech et al.,

2022; Testolin et al., 2023). The required model input data include stand age, average

DBH (Diameter at Breast Height), stand density, and tree height (Collalti et al., 2014).

The soil compartment is represented using one single bucket layer, in which the

available soil water (ASW, in mm) is updated every day considering the water inflows

(precipitation and, if provided, irrigation) and outflows (evapotranspiration, i.e., the sum

of evaporation from the soil and transpiration of the canopy). The remaining water

between these two opposite (in sign) fluxes that exceeds the site-specific soil water

holding capacity is considered lost as runoff. For a full 3D-CMCC-FEM description, see:

https://doi.org/10.32018/ForModLab-book-2024.

2.2 MEDFATE model

MEDFATE v.4.2.0 is an R-based modelling framework that allows the simulation of the

function of forest ecosystems, with a specific emphasis on drought impacts under

Mediterranean conditions (De Cáceres et al., 2021, 2023). MEDFATE calculates energy

balance, photosynthesis, stomatal regulation, and plant transpiration of gas exchange

separately at sub-daily steps. Like 3D-CMCC-FEM, MEDFATE also simulates

photosynthesis at the leaf level using the biochemical model of Farquhar–von

Caemmerer–Berry (Farquhar et al., 1980) for sunlit and shaded leaves (De Pury and

Farquhar, 1997). MEDFATE can simulate plant hydraulics and stomatal regulation

according to two different approaches: (a) steady-state plant hydraulics and

optimality-based stomatal regulation (Sperry et al., 1998; Sperry et al., 2017); and (b)

transient plant hydraulics including water compartments and empirical stomatal

regulation (Sureau-ECOS; Ruffault et al., 2022). In this work, we took the second

approach, i.e., Sureau-ECOS (Ruffault et al., 2022).

The hydraulic architecture of the Sureau-ECOS module comprises arbitrary soil layers,

where the rhizosphere containing coarse and fine root biomass is calculated for each

layer. The total root xylem conductance is determined by factors such as root length

(limited by soil depth), weight, and distribution across the different layers. In addition,

the resistance to water flow is dependent on two plant compartments (leaf and stem,

each composed of symplasm and apoplasm). Overall, plant conductance is defined by

the sum of resistances across the hydraulic network (i.e., soil, stem, and leaves), taking

into account processes such as plant capacitance effects (i.e., the variation of

symplasmic water reservoirs in the stem and leaves) and cavitation flows (i.e., water

released to the streamflow from cavitated cells to non-cavitated cells during

cavitation) (Hölttä et al., 2009). To withstand drought stress, adjacent conduits

(tracheids or vessels) and/or living cells (e.g., parenchyma) release water to the xylem

and may subsequently be refilled. In the event of embolization, cavitated xylem

conduits release their water to the non-cavitated parts of the xylem, which then

transfer it to adjacent compartments. Each element (roots, stem, leaves) of the

hydraulic network has a vulnerability curve $k\Psi$, that declines as water pressure

becomes more negative. The xylem vulnerability curve is modelled using a sigmoid

function, defined by the equation:

$$k\Psi = k_{max}/1 + exp^{\left(\frac{\text{slope}}{25}\right) \cdot \left(\frac{\Psi}{\Psi_{50}}\right)} \qquad (2)$$

where k_{max} is the maximum hydraulic conductance, $\Psi 50$ is the water potential

corresponding to 50% of conductance, and "$slope$" is the slope of the curve at that

point. The stem vulnerability curve can be used to determine the proportion of stem

conductance loss (PLC$_{stem}$) associated with vessel embolism. This embolism reduces

overall tree transpiration and photosynthesis. Plant hydraulic failure and tree death can

occur if the PLC$_{stem}$ exceeds the 50% threshold.

Gas exchange in the Sureau-ECOS module depends on stomatal conductance (which

depends on light, water availability, and air temperature) and leaf cuticular

conductance, which changes with leaf temperature due to changes in the permeability

of the epidermis. Stomatal regulation, unlike the 3D-CMCC-FEM, follows the Baldocchi

(1994) approach, which allows coupling leaf photosynthesis with water losses. In

addition, a multiplicative factor depending on leaf water potential is used to decrease

stomatal conductance under drought conditions, following a sigmoidal function similar

to stem vulnerability.

Soil water balance is computed daily. MEDFATE can consider an arbitrary number of

soil layers with varying depths in which the water movement within the soil follows a

dual–permeability model (Jarvis et al., 1991; Larsbo et al., 2005). Soil water content

(ΔV_{soil}, in mm) is calculated taking into account variables such as infiltration, capillarity

rise, deep drainage, saturation effect, evaporation from the soil surface, transpiration of

the herbaceous plant, and woody plant water uptake. A full MEDFATE description is

available at: https://emf-creaf.github.io/medfatebook/index.html.

2.3 Evaluation sites

Model evaluation was performed in three PROFOUND and FLUXNET Network

European beech sites, i.e., Sorø (DK–Sor, Denmark), Hesse (FR–Hes, France), and

Collelongo (IT–Col, Italy), in which we retrieved information on soil texture, soil depth,

and stand inventory data of forest structure for model initialization (Collalti et al., 2016;

Marconi et al., 2017; Reyer et al., 2020a,b; https://fluxnet.org/). These sites are

equipped with the Eddy Covariance towers (EC; Pastorello et al., 2020) for long–term

continuous monitoring of atmospheric carbon, water, and energy fluxes of the forests

(Fig. 1). The DK–Sor site is located in the forest Lille Bogeskov on the island of Zealand in

Denmark. FR–Hes is situated in the northeastern region of France and lies on the plain

at the base of the Vosges Mountains. IT–Col (Selva Piana stand) is a permanent

experimental plot installed in 1991 and situated in a mountainous area of the Abruzzo

region, centre of Italy.

The pedological characterization of soils exhibits distinct variations across the studied

sites. The soil at the DK–Sor site is predominantly classified as either Alfisols or

Mollisols. The FR–Hes site showcases an intermediary nature, displaying characteristics

akin to both luvisols and stannic luvisols. At the IT-Col site, the prevailing soil type is

identified as Humic alisols, according to the USDA soil classification system. Full details

of these sites are reported in Table 1.

The variables accounted for in the evaluation were obtained from the Fluxdata website

(http://fluxnet.fluxdata.org/data/fluxnet2015-dataset/)   from   the   FLUXNET2015

database (Pastorello et al., 2020). The variables considered are the daily GPP,

estimated from Net Ecosystem Exchange (NEE) measurements and quality checked

using the constant USTAR turbulence correction according to Papale et al. (2006) and

the Latent Heat flux (LE) with energy balance closure correction (i.e., 'LE_CORR')

(Pastorello et al., 2020).

2.4 Study sites

The two target sites considered in this study are Cansiglio and Mongiana Forests (Fig. 1)

(De Cinti et al., 2016). Each site consists of nine long-term monitored plots of differently

managed beech stands, with a spatial extension for each area above 3 hectares, for

about 27 hectares of the experimental area. Three different silvicultural treatments

were applied (see Fig. S1-S2). For each site, three of the nine plots considered were left

unmanaged (i.e., no cutting and leaving the stands to natural development), defined as

'Control' plots, three plots were managed following the historical shelterwood system

('Traditional'), and three with innovative cutting ('Innovative'). In Cansiglio, considering

the developmental stage of the stand was an establishment cut to open growing space

in the canopy for the establishment of regeneration. The 'Innovative' cutting consisted

of selecting a non-fixed number of scattered, well-shaped trees (the 'candidate trees')

and a thinning of neighbouring competitors to reduce competition and promote better

growth. In Mongiana, 'Traditional' silvicultural treatment was the first preparatory cut to

increase the vitality and health of the intended residual trees in the stand. The

'Innovative' option was the identification of 45-50 as 'candidate trees' per hectare and

removing only direct competitors.

The Cansiglio site is situated in a mountainous area in the Veneto region, northern Italy.

Mongiana site is located in a mountainous area in the Calabria region of southern Italy.

The latter shows higher mean annual temperature (MAT, C°) and lower mean annual

precipitation (MAP, mm year^{-1}) (i.e., drier conditions) than the Cansiglio site located at

higher latitudes (Table 1). Data on forest structure and soil texture were collected

during the field campaigns conducted in 2011 and 2019 (Cansiglio) and in 2012 and 2019

(Mongiana). At the Cansiglio site, soils are identified as Haplic luvisols, whereas at

Mongiana, the predominant soil classifications consist of Inceptisols and Entisols,

according to the USDA soil classification system. The variables analyzed in these sites,

like in the evaluation sites, were GPP and LE. A summary of these sites is reported in

Table 1.

2.5 Meteorological data

For the evaluation sites (i.e., DK-Sor, FR-Hes, IT-Col) observed meteorological data

were retrieved from the harmonized PROFOUND database (Reyer et al., 2020a, b) and

FLUXNET2015 database (Pastorello et al., 2020; https://data.icos-cp.eu/).

For the Mongiana and Cansiglio sites, meteorological data for 2010-2022 were

obtained at daily temporal resolution from the relevant region's Regional Environmental

Protection Italian Agencies (ARPAs), which are responsible for monitoring climate

variables with weather stations. The choice of thermo-pluviometric weather station

was based on the minimum distance from the study area (between 2 km and 9 km

away from the study sites, respectively) and on the data availability and integration

with other weather stations in the proximity, representing the best available and

obtainable meteorological observed data for these sites. The Bagnouls–Gaussen graph

(Fig. S3) shows the mean monthly precipitation (mm) and air temperature (°C)

recorded for every station inside the catchment.

Climate scenarios used as inputs for the two models at the Cansiglio and Mongiana

sites were from the COSMO-CLM simulation at a spatial resolution of approximately

2.2 km over Italy (Raffa et al., 2023).

The daily variables considered for 3D-CMCC-FEM were mean solar radiation (MJ m^{-2}

day^{-1}), maximum and minimum air temperature (°C), precipitation (mm day^{-1}), and the

mean relative air humidity (%). In contrast, the MEDFATE model uses mean solar

radiation, maximum and minimum air temperature, precipitation, the daily maximum

and minimum relative air humidity, and wind speed (m s^{-1}).

2.6 Modelling set-up

A set of parameters specific for *Fagus sylvatica* L. was provided as input to the model

3D-CMCC-FEM as described in Collalti et al. (2023) while for MEDFATE as in De

Cáceres et al. (2023). To remove any confounding factors related to parameterization,

the parameters related to photosynthesis and stomatal conductance were kept

constant between the two models (see Table 2). A complete list of parameters and

their values for both models as adopted in the present study can be found in

supplementary materials (Table S2).

We then used the LAI and Available Soil Water (AWS) values obtained from the 3D

CMCC-FEM outputs as input for running simulations with the MEDFATE model given

that the model function used here does not prognostically simulate LAI and Available

Soil Water (ASW). Precisely, here we used MEDFATE to simulate C and H_2O fluxes only

while considering plant hydraulics (De Cáceres et al., 2021), from the forest structure

predicted, in terms of LAI, by 3D-CMCC-FEM. For MEDFATE water balance, LAI values

determine the competition for light and also drive the competition for soil water, along

with the root distribution across soil layers.

2.7 Model evaluation

To evaluate the performance of the two models across the different sites examined and under varying environmental conditions, we first assessed GPP and LE daily fluxes along a latitudinal gradient at sites equipped with EC towers. The reliability of the two models subsequently allowed us to coherently simulate fluxes at sites where EC towers were absent. Both models were run for five years on the evaluation sites, with the simulation period determined by the availability of observed data provided, as already mentioned, from the PROFOUND database, specifically, from 2006 to 2010 at DK-Sor and IT-Col sites while for FR-Hes starting from 2014 to 2018. The performance metrics of the results of the evaluation for each site for the GPP and LE variables were the coefficient of determination (R^2), Root Mean Square Error (RMSE), and Mean Absolute Error (MAE).

2.8 Model application in managed sites

In the managed sites (i.e., Cansiglio and Mongiana), simulations were performed using Historical climate ('Hist') and, to analyse models' sensitivities to climate change, under two Representative Concentration Pathways 4.5 and 8.5 ('Moderate' and 'Hot Climate'), respectively. The 'Hist' climate was used to run simulations at the Cansiglio site from 2011 to 2022 and the Mongiana site from 2012 to 2022. In contrast, simulations using RCP4.5 and RCP8.5 climate ran accounting for the same period, that is, eleven years for the Cansiglio site and ten years for the Mongiana site, but considering the last years of the climate change scenarios (i.e., 2059-2070 and 2060-2070, respectively) to create harsher temperature and precipitation conditions, but with an increased atmospheric CO_2 concentration (in $\mu mol\ mol^{-1}$).

For each of the nine sampled areas, in the Cansiglio and Mongiana sites, we considered a representative area of one hectare for each type of plot: 'Control', 'Traditional', and

'Innovative'. At the beginning of the simulations, each site thus included a total of 9

plots, each one hectare in size—comprising three 'Control' plots, three 'Traditional'

plots, and three 'Innovative' plots. This setup resulted in a total of nine hectares being

simulated per site where the model 3D-CMCC-FEM removed a certain percentage of

the Basal Area (BA) according to the LIFE-ManFor project (see Table S1). 'Traditional'

and 'Innovative' cutting took place for the first time in 2012 (Cansiglio) and 2013

(Mongiana), respectively. Following preliminary results, since the Mongiana site

experienced a lighter thinning intensity compared to the Cansiglio site (refer to Table

S1), consequently, for the Mongiana site, we considered an alternative management

option involving the removal of 40% of the BA. This was done to evaluate whether a

more intensive management approach ('SM') could have influenced models' results on

GPP and LE fluxes related to the reduction in competition and enhanced water

availability.

3. Results

3.1 Model evaluation

The GPP at DK-Sor, FR-Hes, and IT-Col sites estimated from EC and simulated by 3D-

CMCC-FEM and MEDFATE are shown in Fig. 2. At the DK-Sor site, the 3D-CMCC-FEM

simulates a mean daily GPP of 5.14 gC m^{-2} day^{-1}, while MEDFATE 5.13 gC m^{-2} day^{-1};

and EC 5.54 gC m^{-2} day^{-1}; at the FR-Hes site, 3D-CMCC-FEM mean daily GPP of 6.18

gC m^{-2} day^{-1} compared to MEDFATE 4.82 gC m^{-2} day^{-1}, and EC 4.99 gC m^{-2} day^{-1};

lastly at the IT-Col site, 3D-CMCC-FEM mean daily GPP of 4.88 gC m^{-2} day^{-1}

compared to MEDFATE 4.19 gC m^{-2} day^{-1}; and EC 4.11 gC m^{-2} day^{-1}. Additionally, at

the DK-Sor site, the 3D-CMCC-FEM simulated a mean daily LE of 2.83 MJ m^{-2} day^{-1},

while the MEDFATE simulated a mean value of 2.22 MJ m^{-2} day^{-1}; and EC 3.19 MJ m^{-2}

day^{-1}; at the FR-Hes site, 3D-CMCC-FEM simulated a mean daily LE of 4.16 MJ m^{-2}

day^{-1} compared to MEDFATE 3.01 MJ m^{-2} day^{-1}; and EC 4.47 MJ m^{-2} day^{-1}; in the

end at the IT-Col site, 3D-CMCC-FEM simulated a mean daily LE of 2.02 MJ m^{-2} day^{-1}

compared to MEDFATE 2.57 MJ m^{-2} day^{-1}; while EC 3.93 MJ m^{-2} day^{-1}. The GPP

predicted by 3D-CMCC-FEM has shown higher values of R^2 (0.92) at DK-Sor and the

lowest value at FR-Hes site (R^2 = 0.76) whilst a value of R^2 = 0.83 at IT-Col site,

respectively. For the MEDFATE model, the highest predicted GPP value of R^2 (0.85) was

at DK-Sor, the lowest (R^2 = 0.68) at IT-Col, and at FR-Hes R^2 = 0.76, the same showed

for the 3D-CMCC-FEM model, respectively. Differently, the highest R^2 (0.89) value for

3D-CMCC-FEM considering LE predicted vs. observed was at FR-Hes site and almost

the same values for DK-Sor and IT-Col sites (R^2 = 0.85 and 0.84, respectively).

MEDFATE, for predicted vs. observed LE variable, has shown the highest R^2 (0.77) at IT-

Col site, lower R^2 (0.69) value at FR-Hes site and the lowest R^2 (0.62) value at DK-Sor

site, respectively. In general, both the Root Mean Square Error (RMSE) and Mean

Absolute Error (MAE) values in all sites were reasonably low, falling within the ranges

of 3.31 to 2.02 gC m^{-2} day^{-1} and 2.46 to 1.47 MJ m^{-2} day^{-1}, for both models and for

both the variables. In Fig. 3 and Table 3 the summary of the evaluation metrics

performance results.

3.2 Simulation results at Cansiglio

Fig. 4 shows the simulation results using the 3D-CMCC-FEM and MEDFATE models in

the Cansiglio site. For the 3D-CMCC-FEM, the 'Control' plot exhibited the lowest GPP

values under 'Hist' climate conditions, averaging 1681 gC m^{-2} year^{-1}. These values

increased slightly to 1982 gC m^{-2} year^{-1} under the RCP4.5 climate and further rose to

2204 gC m^{-2} year^{-1} under the RCP8.5 climate. Similarly, for plots managed with

'Traditional' methods, the trends were consistent with the 'Control' plot, showing

average GPP values of 1603, 1942, and 2141 gC m^{-2} year^{-1} under 'Hist', RCP4.5 and

RCP8.5 climate, respectively. However, 'Innovative' management showed lower GPP

fluxes across all three climate scenarios, with average values of 1534, 1882, and 2075

gC m^{-2} year^{-1} under 'Hist' RCP4.5 and RCP8.5 climate, respectively. The MEDFATE

model showed higher mean absolute GPP increases than the 3D-CMCC-FEM model

under RCP4.5 and RCP8.5 climates, respectively. Under the 'Hist' climate and all

treatments, the mean GPP values were about 1638 gC m^{-2} year^{-1}, whereas under the

RCP4.5 climate, they rose to 2516 gC m^{-2} year^{-1} and 2995 gC m^{-2} year^{-1} under

RCP8.5 climate. Analyzing in Fig. 4 the trends of LE for the 3D-CMCC-FEM model, these

trends closely follow those of GPP concerning management treatments. The 3D-

CMCC-FEM LE values for the 'Control' plots, similar to GPP, were lowest for the 'Hist'

climate with an average value over the simulation years of 1200 and 1501 MJ m^{-2}

year^{-1} for the RCP4.5 climate, and 1391 MJ m^{-2} year^{-1} for the RCP8.5 climate,

respectively. The LE of the 'Traditional' management predicts values of 1129 in the 'Hist'

climate, 1440 in the RCP4.5 climate, and 1338 MJ m^{-2} yr^{-1} in the RCP8.5 climate,

respectively. For the 'Innovative' management, the mean LE values were 1121 in the

'Hist' climate, 1403 for the RCP4.5 climate, and 1306 MJ m^{-2} yr^{-1} for the RCP8.5

climate, respectively. Similar to the GPP fluxes, the MEDFATE model simulated

reductions in LE fluxes among the treatments and higher values across the climates.

The mean LE value modelled in the 'Hist' climate, grouped by treatments (because of

slight differences among managements), was about 920, 1419 in the RCP4.5 climate,

and 1456 MJ m^{-2} yr^{-1} in the RCP8.5 climate, respectively (Fig. S12). MEDFATE

simulated a stem xylem conductance loss of approximately 40% in the seventh, eighth,

and twelfth years of simulation for the RCP8.5 climate scenario in the 'Control' plot. In

contrast, this loss was predicted only in the seventh year for the managed plots.

Conversely, near-zero or negligible stem embolism were simulated under the 'Hist' and

RCP4.5 climate scenarios. The 3D-CMCC-FEM simulated higher values, albeit in a small

percentage (i.e., between 8-10%) of NSC, increasing proportionally to the intensity of

basal area removed, better observable in the graph at the tree level.

3.3 Simulation results at Mongiana

Simulation results at the (drier) Mongiana site well depicted the differences with the

rainy Cansiglio site (Fig. 5). The 3D-CMCC-FEM model showed no significant

differences in the mean values of GPP among various management interventions under

'Hist' climate conditions, with a mean value of 2151 gC m^{-2} yr^{-1}. Compared to the

Cansiglio site, Mongiana exhibited lower average GPP values. Under RCP4.5 climate

conditions, the GPP for the 'Control' plot was 1864 gC m^{-2} yr^{-1}. In contrast, under the

current climate, the 'Traditional' and 'Innovative' management interventions yielded

higher average GPP values of 2115 gC m^{-2} yr^{-1} and 2086 gC m^{-2} yr^{-1}. The GPP values

under the more intensive management ('SM') and RCP4.5 climate decreased even

further than those of the 'Control' plot, with an average value of 1650 gC m^{-2} yr^{-1}.

Under the RCP8.5, no differences in GPP were observed among management

strategies, with values of about 1525 gC m^{-2} yr^{-1}. Moreover, under the RCP8.5, the

'Control' plot experienced complete mortality after five years of simulations. The

MEDFATE model predicted slightly higher average values of GPP in the 'Control' plots

(2099 gC m^{-2} yr^{-1}) compared to the managed plots (2087 gC m^{-2} yr^{-1},

encompassing both 'Traditional' and 'Innovative' of two management strategies), with

no significant differences observed among the management strategies and under 'Hist'

climate. Under the RCP4.5 and RCP8.5, the GPP values were 1608 gC m^{-2} yr^{-1} and

1935 gC m^{-2} yr^{-1}, respectively (Fig. S11). The PLC$_{stem}$ graph in Fig. 5 indicated very high

xylem embolism levels (i.e., reaching 100% every year) under RCP4.5 and RCP8.5

already in the first year of simulations. A pronounced embolism event was observed

under the 'Hist' climate in 2017, 2018, 2019, and 2022 in a 30-45% range for the 'Control' plots, while the managed plots experienced a maximum embolism of approximately 40% in 2017. Conversely, the 3D-CMCC-FEM model did not report any significant differences between managed and unmanaged plots for the LE. The average LE value for the 'Hist' climate was 1796 MJ m^{-2} yr^{-1}, which decreased to 1220 MJ m^{-2} yr^{-1} under the RCP4.5 and 1190 MJ m^{-2} yr^{-1} under the RCP8.5 in managed plots. As previously described, the 'Control' plot under the RCP8.5 experienced mortality in the sixth year of simulation. Similarly to the previously described GPP fluxes, the MEDFATE model reported a slight difference in LE fluxes between the 'Control' plot under historical climate conditions (1623 MJ m^{-2} yr^{-1}) and the managed plots (1603 MJ m^{-2} yr^{-1}). For the RCP4.5 and RCP8.5, the LE values were 1100 MJ m^{-2} yr^{-1} and 1089 MJ m^{-2} yr^{-1}, respectively (Fig. S12). The LE values of the 'Control' plots are not reported either for RCP4.5 or for RCP8.5 because of the mortality experienced for the simulation years.

4. Discussions

First, this study evaluated the performances of two different process-based models in simulating diverse beech stands across Europe, starting to the north of Europe and moving towards the south under different environmental conditions. Secondly, the study has focused on the models' sensitivity and the relative impacts of different management options and different climatic conditions in two independent beech forest stands in the north and south of the Italian peninsula.

4.1 Model evaluation

To assess the models' accuracy in predicting C and H$_2$O fluxes, we compared daily GPP and LE data obtained from the EC towers. Both models predicted daily GPP and LE

accurately and ensured a good range of general applicability of both models (Kramer et

al., 2002; Verbeeck et al., 2008). The 3D-CMCC-FEM model seems to slightly

overestimate GPP daily values along latitudinal gradients starting from the north (DK-

Sor) to the south (IT-Col), as already found in Collalti et al. (2016). MEDFATE, in

contrast, showed a slight overestimation of GPP only at IT-Col site. The LE predicted by

3D-CMCC-FEM is more accurate than MEDFATE prediction for DK-Sor and FR-Hes

sites but not in IT-Col site in which 3D-CMCC-FEM has shown to underestimate

compared to the observed EC values. For MEDFATE the underestimation of LE was

observed in all the evaluation sites.

The spread observed for the GPP and LE fluxes between the two models may be

attributed to the different assumptions that govern stomatal regulation since both

models use the Farquhar-von Caemmerer-Berry biochemical model to calculate

photosynthesis. The over or underestimation of the flows estimated by the models

both for GPP and the LE compared to the data observed from the EC towers can be

attributed either to the presence of the understory (although commonly sporadic in

mature beech stands), which was not considered in the simulations by both models

and to errors on daily measurement by EC technique (Loescher et al., 2006) or

because a not perfect fit in the modeled seasonality (i.e., the begin and the end of the

growing season) (Richardson et al., 2010). However, the overall leaf phenological

pattern of the European beech in these sites is well represented by the two models in

almost all of the years according to EC data as shown in supplementary materials (see

Fig. S4, S5, S6, S7, S8, S9, and Fig S10). It is important to note that we did not

specifically calibrate the models' parameters at each site separately. Instead, both

models were parameterized using existing values taken from the literature, therefore

with one single set of parameter values for all sites.

4.2 Climate change and forest management at the Cansiglio and Mongiana site

The pre-Alpine site of Cansiglio showed slight differences in the fluxes (i.e., GPP and LE) between the three different management practices and the three climate scenarios (i.e., no climate change, RCP4.5. and RCP8.5) used (Fig. S11–S12). Future climate is expected to be higher temperature if compared to the historical one, with MAT higher of about 3.93°C under RCP4.5 and 4.95°C under RCP8.5 for the Cansiglio site and 4.52°C under RCP4.5 and 5.42°C under RCP8.5 at the Mongiana site. Similarly, MAP is expected to be 510 mm lower under RCP4.5 and 602 mm under RCP8.5 at the Cansiglio site, respectively, while 902 mm lower under RCP4.5 and 914 mm under RCP8.5 at the Mongiana site, respectively.

Regarding management, the response of the 3D-CMCC-FEM to the removal of a percentage of the basal area from the stand led to a decrease in GPP in the 'Traditional' cutting and an even greater extent, in the 'Innovative' cutting compared to the 'Control' (i.e., no management). Similarly to 3D-CMCC-FEM, the MEDFATE model simulates slight differences in fluxes amount (e.g., lower values for 'Traditional' and 'Innovative' cutting than 'Control' plots) between the management regimes in the plots. These results align with those of Guillemot et al. (2014), who observed a slight decrease in GPP in managed compared to unmanaged temperate beech forests in France under different thinning regimes. However, differences were observed in both models under the three different climates used in the simulations. GPP increased from the 'Hist' climate to RCP4.5 and reached the maximum for RCP8.5, respectively. This suggests a plastic response (e.g., photosynthesis and stomatal response) of the stands, as simulated by models, to harsher conditions, indicating, potentially, a high drought acclimation capacity (Petrik et al., 2022) and increased GPP because of the early budburst, and a prolonged vegetative season (Peano et al., 2019), and the so-called 'atmospheric CO_2 fertilization' effect as also found by de Wergifosse et al. (2022) and

Reyer et al. (2013), especially in sites with no apparent water limitation both under current and projected future climate conditions. The anisohydric behavior of *Fagus sylvatica* L. results in prolonged stomatal opening relative to isohydric species, although Puchi et al. (2024) recently found large variability in European beech responses, maintaining prolonged photosynthetic activity. Still, this response is modulated by summer precipitation and the availability of soil water storage (Leuschner et al., 2021; Baudis et al., 2015). However, for high-altitude stands, growth could be negatively affected under warmer conditions, as suggested by Chmura et al. (2024). The LE results for the 3D-CMCC-FEM showed lower values over the simulation period for managed stands than unmanaged ones showing lesser sensitivity to forest management if compared to MEDFATE. Yet, under the RCP4.5, the LE values were higher compared to both the 'Hist' climate and the RCP8.5 one due to greater annual cumulative precipitation than the RCP8.5 and higher, on average, temperatures than the 'Hist' scenario (Fig. S12).

Conversely, the MEDFATE model was shown to be more sensitive to climate, with a clearer distinction between the 'Hist' climate, the RCP4.5 and RCP8.5 climates, with higher and nearly equal values in the harsher conditions (i.e., RCP4.5 and RCP8.5 climates), with slight differences in the management treatments as obtained by 3D-CMCC-FEM.

The Non-Structural Carbon (NSC) amount showed the highest values in 'Innovative' plots, followed by 'Traditional' plots, and the lowest values in 'Control' plots, suggesting a benefit in carbon stock accumulation with more carbon going for carbon biomass and less for reserve-replenishment for these stands under management interventions. Nevertheless, NSC levels remain nearly the same for the three climate scenarios throughout all the simulation years. It is important to note that MEDFATE simulated an initial loss of stem conductance under the climate scenarios, indicating a premature

onset of water stress for the stand. Although in RCP4.5 this is negligible, in RCP8.5

PLC_{stem} values reach a maximum xylem cavitation value of about 40% in the eighth

year of simulation for managed plots while for 'Control' plots in the eighth, ninth, and

twelfth years, highlighting potential benefits of management to reduce drought stress

because of less rain interception and canopy evaporation and transpiration (Giuggiola

et al., 2018; Schmied et al., 2023).

The GPP at the southern Apennine site of Mongiana showed a decrease under RCP4.5

and RCP8.5 scenarios when simulated by the 3D-CMCC-FEM model as a result of

harsher environmental conditions, as also resulted in the study by Yu et al. (2022), in

which the productivity and then the growth of European beech in southern regions are

expected to decrease as affected by more severe climate conditions such as decreased

precipitation and increased air temperature (Tognetti et al., 2019). Indeed, the increase

in air temperature, a reduction in soil water availability, and the rise in vapor pressure

deficit (VPD) lead to earlier stomatal closure, increased mesophyll resistance, and

elevated abscisic acid production (Kane and McAdam, 2023), all of which contribute to

a decrease in the carbon assimilation rate (Priwitzer et al., 2014; Grossiord et al., 2020).

Specifically, GPP is higher under 'Hist' climate conditions, decreases under the RCP4.5,

and ultimately reaches even lower values under the RCP8.5. Under the RCP8.5 at the

fifth year of simulation, the stand in the 'Control' plot is simulated to die due to carbon

starvation. The annual decline in NSC (Fig. 5) due to an imbalance between carbon

uptake (photosynthesis) and the demands for growth and respiration suggests that the

trees are unable to replenish their carbon reserves. The depletion of NSC reserves may

ultimately disrupt processes such as osmoregulation and phenology (Martínez-Vilalta et

al., 2016), potentially leading to stand defoliation and/or mortality. The management

options did not show changes in GPP under the 'Hist' climate. However, the increase of

GPP was observed under the RCP4.5 in the plots where 'Innovative' and 'Traditional'

cutting occurred, although no differences were observed between them. For instance,

the same increase in GPP was reported by Fibbi et al. (2019) for other European beech

forests under climate change scenarios in Italy. The thinning reduces the leaf area and

then the LAI and increases the soil water availability, which positively influences

stomatal conductance and carbon assimilation, providing an acclimation mechanism to

drought during periods of water scarcity (Lüttschwager and Jochheim, 2020; Diaconu

et al., 2017).

In contrast, the more intense cutting exhibited even lower GPP values than the 'Control'

plots. This is likely due to the overly intense thinning, which contrasts the microclimate

effects within this forest stand, reducing the potential to offset climate warming at the

local scale (Rita et al., 2021). Heavy thinning, on the other hand, can increase light

penetration, soil evaporation, and wind speed, thereby heightening tree sensitivity to

vapor pressure deficit under dry conditions (Schmied et al., 2023; Simonin et al., 2007).

LE decreased with the decrease in precipitation under the RCP4.5 and RCP8.5 climate

scenarios compared to the 'Hist' climate. There were no significant differences in LE

among the various management regimes. For the MEDFATE model, negligible or no

differences in GPP were observed under all the climates among various management

options. Although the GPP values estimated by the MEDFATE model under the RCP4.5

and RCP8.5 are similar to those obtained from the 3D–CMCC–FEM model, a closer

analysis of the daily outputs (data not shown) reveals that trees photosynthesize until

the end of July, after which they experience significant embolism (i.e., maximum value

of 100%), as indicated by the PLC_{stem} graph, indicating that the decrease in

precipitation led to summer soil moisture depletion and lethal drought stress levels.

Furthermore, the 'Control' plots experienced mortality even before reaching the

summer period. In recent decades, prolonged drought stress in Mediterranean

mountain regions has significantly reduced the productivity of beech forests, resulting

in a decline in Basal Area Increment (BAI) and overall growth (Piovesan et al., 2008). It

is also important to note that under 'Hist' climate conditions, the MEDFATE model

indicated a stem embolization loss ranging from approximately 10% to 45% during the

drought period (i.e., 2018-2020) in Europe as also highlighted in other study (Italiano et

al., 2024; Thom et al., 2023; Lombardi et al., 2023). The embolization was more

pronounced and long-lasting in the 'Control' plots than the managed ones. The same

trends were obtained for LE.

4.3 Uncertainties and factors influencing forest carbon and water dynamics

Although there is scientific evidence of the positive effects of CO_2 fertilization effect on

forest primary productivity, uncertainties remain regarding the long-term persistence

of this positive feedback and the level at which this may saturate (Sperlich et al., 2020;

Wang et al., 2020). Furthermore, the down-regulation of this fertilization effect on

photosynthesis is influenced by interannual variations in meteorological parameters, as

well as by interactions within the carbon and nitrogen cycles. These factors must be

carefully assessed to improve the accuracy of flux projections under future climatic

conditions (Zaehle et al., 2014). It is also worth highlighting that this study does not

account for biotic disturbances such as pest outbreaks and diseases, nor for abiotic

disturbances like extreme climatic events (e.g., heatwaves, late frosts, and wildfires).

These factors could significantly alter carbon and water fluxes (Yu et al., 2022b),

potentially depleting carbon reserves faster and reducing the capacity for carbon

sequestration by these forest ecosystems as well as possibly necessitating the

adoption of different management strategies (Langer and Bußkamp, 2023;

Margalef-Marrase et al., 2020). Another critical factor is the depth of the root zone and

soil, as well as its physico-chemical composition. For example, it has been found that

two-thirds of fine roots in European beech are within the top 30 cm of soil, while

coarse roots can extend beyond depths of 240 cm (Meier et al., 2017). Although

findings by Gessler et al. (2021) indicate that, unlike oak forests, European beech forests

cannot compensate for additional water uptake from deeper soil layers during drought

periods, Brinkmann et al. (2018) reported contrasting results. In this study, we analyzed

soil texture characteristics to a depth of 110 cm for Cansiglio and 40 cm for Mongiana.

However, the limited understanding of deeper layers may not fully capture the entire

soil water reservoir and its dynamics. Expanding knowledge of the deepest soil layers

is essential to better understand root development and, consequently, improve water

storage capacity and drought resilience in beech forests.

5. Conclusions

The two process-based models provide robust evidence for their application in

estimating fluxes, consistent with long-term EC tower measurements in European

beech forests. Despite the minimal parametrization effort to align the two models and

the avoidance of single site-specific parameters, reliable results can still be obtained,

as confirmed by the outputs from the Sorø, Hesse, and Collelongo sites. Regarding the

sub-Alpine Cansiglio site, although water limitation does not significantly impact fluxes

or the health of the forest under 'Moderate' climate conditions (RCP4.5), a potential

concern is the embolization predicted by the MEDFATE model under the 'Hot' climate

(RCP8.5) at this site, despite similar levels of precipitation. The high susceptibility of

European beech forests at the southern Apennine site of Mongiana to more severe (i.e.,

hotter and drier) climatic conditions could lead to the collapse of this forest ecosystem,

even with the application of management options to reduce competition. However, it is

crucial that these seasonal droughts are not prolonged or intense enough to exceed the

ecological limits of the European beech. To avoid that European beech forests may

necessitate of strategic and specifically designed management planning at the single

site level, including the ability to project (e.g., with forest models) and evaluate future

forest conditions for better management schemes. However, the ability of these forests

to survive or resist the impacts of climate change may not depend solely on density

reduction interventions. Prioritizing the exploration of alternative sustainable

management strategies to promote carbon sequestration in both above-ground

biomass and soil is crucial for enhancing climate change mitigation efforts. Additionally,

evaluating silvicultural plans such as the introduction of complementary species can

improve the resilience of vulnerable European beech ecosystems. A modeling

approach, similar to the one used in this study, offers a valuable tool for assessing

these alternative strategies and refining forestry adaptive management practices. By

integrating these approaches, we can strengthen the long-term sustainability of forests

while preserving the ecological balance of vulnerable regions.

CRediT authorship contribution statement

Vincenzo Saponaro: Conceptualization, Data curation, Formal analysis, Investigation,

Methodology, Resources, Software, Visualization, Writing – original draft, Writing –

review & editing. **Miquel De Càceres:** Conceptualization, Methodology, Software,

Supervision, Writing – review & editing. **Daniela Dalmonech:** Conceptualization,

Methodology, Software, Supervision, Writing – review & editing. **Ettore D'Andrea:**

Resources, Methodology, Data curation, Writing – review & editing. **Elia Vangi:**

Resources, Writing – review & editing. **Alessio Collalti:** Conceptualization,

Methodology, Resources, Software, Supervision, Project administration, Writing –

review & editing.

Data availability

Data will be made available on request.

Acknowledgements

We would like to thank the University of Tuscia, the CNR ISAFOM of Perugia and the

company ENI s.p.a. for the realization of this work. Special thanks to the Institute

Research Center for Ecological and Forestry Applications (CREAF) of Barcelona that

supported the research by the Spanish "Ministerio de Ciencia e Innovación"

(MCIN/AEI/10.13039/501100011033) (grant agreement No. PID2021-126679OB-I00).

The authors sincerely thank the ARPA Veneto and Calabria for providing the

meteorological data. The Fondazione Centro Euro-Mediterraneo sui Cambiamenti

Climatici (CMCC) for providing climate change scenarios. The Cost-Action PROFOUND

for providing both the stand ancillary data and the climate used in this work. D.D., E.V.

and A.C. has been partially supported by MIUR Project (PRIN 2020) "Unraveling

interactions between WATER and carbon cycles during drought and their impact on

water resources and forest and grassland ecosySTEMs in the Mediterranean climate

(WATERSTEM)" (Project number: 20202WF53Z), "WAFER" at CNR (Consiglio Nazionale

delle Ricerche) and by PRIN 2020 (cod. 2020E52THS) – Research Projects of National

Relevance funded by the Italian Ministry of University and Research entitled: "Multi-

scale observations to predict Forest response to pollution and climate change"

(MULTIFOR, project number: 2020E52THS). A.C. acknowledge also funding by the

project OptForEU Horizon Europe research and innovation programme under grant

agreement No. 101060554. D.D. and A.C. also acknowledge the project funded under

the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment

1.4 – Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18

December 2021 of Italian Ministry of University and Research funded by the European

Union – NextGenerationEU under award Number: Project code CN_00000033,

Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP B83C22002930006, Project title "National Biodiversity Future Centre – NBFC". This work used eddy covariance data acquired and shared by the FLUXNET community, including these networks: AmeriFlux, AfriFlux, AsiaFlux, CarboAfrica, CarboEuropeIP, CarboItaly, CarboMont, ChinaFlux, Fluxnet-Canada, GreenGrass, ICOS, KoFlux, LBA, NECC, OzFlux-TERN, TCOS-Siberia, and USCCC. The 3D-CMCC-FEM model code is publicly available and can be found on the GitHub platform at: https://github.com/Forest-Modelling-Lab/3D-CMCC-FEM. The MEDFATE model code is publicly available and can be found on the GitHub platform at: https://github.com/emf-creaf/medfate.

References

Augusto, L., Boča, A. (2022). Tree functional traits, forest biomass, and tree species diversity interact with site properties to drive forest soil carbon. Nat. Commun. 13, 1–12. https://doi.org/10.1038/s41467-022-28748-0

Axer, M., Schlicht, R., Kronenberg, R., Wagner, S. (2021). The Potential for Future Shifts in Tree Species Distribution Provided by Dispersal and Ecological Niches: A Comparison between Beech and Oak in Europe. Sustainab., 13, 13067. https://doi.org/10.3390/ su132313067

Baldocchi, D. (1994). An analytical solution for coupled leaf photosynthesis and stomatal conductance models. Tree Phys., 14(7-8–9), 1069–1079. https://doi.org/10.1093/treephys/14.7-8-9.1069

Baudis, M., Premper, T., Welk, E., & Bruelheide, H. (2015). The response of three Fagus sylvatica L. provenances to water availability at different soil depths. Ecol. Res., 30(5), 853–865.https://doi.org/10.1007/s11284-015-1287-x

Bernacchi, C. J., Singsaas, E. L., Pimentel, C., Portis, A. R., & Long, S. P. (2001). Improved temperature response functions for models of Rubisco-limited photosynthesis. Plant Cell & Env., 24(2), 253–259. https://doi.org/10.1111/j.1365-3040.2001.00668.x

Bernacchi, C. J., Calfapietra, C., Davey, P. A., Wittig, V. E., Scarascia-Mugnozza, G. E., Raines, C. A., & Long, S. P. (2003). Photosynthesis and stomatal conductance responses of poplars to free-air CO_2 enrichment (PopFACE) during the first growth cycle and immediately following coppice. New Phyt., 159(3), 609–621. https://doi.org/10.1046/j.1469-8137.2003.00850.x

Bosela, M., Štefančík, I., Petráš, R., Vacek, S. (2016). The effects of climate warming on the growth of European beech forests depend critically on thinning

strategy and site productivity. Agric. For. Meteorol. 222, 21–31. https://doi.org/10.1016/j.agrformet.2016.03.005

Bonan, G. B., Lawrence, P. J., Oleson, K. W., Levis, S., Jung, M., Reichstein, M., Lawrence, D. M., & Swenson, S. C. (2011). Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data. J. of Geoph. Res. Atm., 116(G2). https://doi.org/10.1029/2010jg001593

Brinkmann, N., Eugster, W., Buchmann, N., & Kahmen, A. (2018). Species-specific differences in water uptake depth of mature temperate trees vary with water availability in the soil. Plant Biol., 21(1), 71–81. https://doi.org/10.1111/plb.12907

Brunet, J.; Fritz, Ö.; Richnau, G. (2010). Biodiversity in European beech forests—A review with recommendations for sustainable forest management. Ecol. Bull. 53, 77–94, Available online: http://www.jstor.org/stable/41442021

Chmura, D. J., Banach, J., Kempf, M., Kowalczyk, J., Mohytych, V., Szeligowski, H., Buraczyk, W., & Kowalkowski, W. (2024). Growth and productivity of European beech populations show plastic response to climatic transfer at the north-eastern border of the species range. For. Eco. and Man., 565, 122043. https://doi.org/10.1016/j.foreco.2024.122043

Collalti, A., Perugini, L., Santini, M., Chiti, T., Nolè, A., Matteucci, G., Valentini, R. (2014). A process-based model to simulate growth in forests with complex structure: Evaluation and use of 3D-CMCC Forest Ecosystem Model in a deciduous forest in Central Italy. Ecol. Modell. 272, 362–378. https://doi.org/10.1016/j.ecolmodel.2013.09.016

Collalti, A., Marconi, S., Ibrom, A., Trotta, C., Anav, A., D'andrea, E., Matteucci, G., Montagnani, L., Gielen, B., Mammarella, I., Grünwald, T., Knohl, A., Berninger, F., Zhao, Y., Valentini, R., Santini, M. (2016). Validation of 3D-CMCC Forest Ecosystem Model (v.5.1) against eddy covariance data for 10 European forest sites. Geosci. Model Dev. 9, 479–504. https://doi.org/10.5194/gmd-9-479-2016

Collalti, A., Trotta, C., Keenan, T.F., Ibrom, A., Bond-Lamberty, B., Grote, R., Vicca, S., Reyer, C.P.O., Migliavacca, M., Veroustraete, F., Anav, A., Campioli, M., Scoccimarro, E., Šigut, L., Grieco, E., Cescatti, A., Matteucci, G. (2018). Thinning Can Reduce Losses in Carbon Use Efficiency and Carbon Stocks in Managed Forests Under Warmer Climate. J. Adv. Model. Earth Syst. 10, 2427–2452. https://doi.org/10.1029/2018MS001275

Collalti, A., Thornton, P.E., Cescatti, A., Rita, A., Borghetti, M., Nolè, A., Trotta, C., Ciais, P., Matteucci, G. (2019). The sensitivity of the forest carbon budget shifts across processes along with stand development and climate change. Ecol. Appl. 29, 1–18. https://doi.org/10.1002/eap.1837

Collalti, A., Tjoelker, M.G., Hoch, G., Mäkelä, A., Guidolotti, G., Heskel, M., Petit, G., Ryan, M.G., Battipaglia, G., Matteucci, G., Prentice, I.C. (2020). Plant respiration: Controlled by photosynthesis or biomass? Glob. Chang. Biol. 26, 1739–1753. https://doi.org/10.1111/gcb.14857

Collalti, A., Dalmonech, D., Marano, G., Vangi, E., Puchi, P., Grieco, E., Orrico, M. (2023). 3D-CMCC-FEM (Coupled Model Carbon Cycle). BioGeoChemical and

Biophysical Forest Ecosystem – User's Guide. CNR Edizioni; ISBN 978-88-8080-573-1 (electronic edition) https://doi.org/10.32018/3D-CMCC-FEM-2022;

Collalti, A. and Dalmonech, D. and Vangi, E. and Marano, G. and Puchi, P.F. and Morichetti, M. and Saponaro, V. and Orrico, M.R. and Grieco, E. (2024) Monitoring and Predicting Forest Growth and Dynamics. CNR Edizioni, Roma.

Collatz, G.J., Ball, J.T., Grivet, C. & Berry, J.A. (1991). Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: A model that includes a laminar boundary layer. Agric. For. Meteorol., 54, 107–136.

Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions: Forging a climate-resilient Europe – the new EU Strategy on Adaptation to Climate Change – COM(2021) 82 final.

Dalmonech, D., Marano, G., Amthor, J.S., Cescatti, A., Lindner, M., Trotta, C., Collalti, A. (2022). Feasibility of enhancing carbon sequestration and stock capacity in temperate and boreal European forests via changes to management regimes. Agric. For. Meteorol. 327, 109203. https://doi.org/10.1016/j.agrformet.2022.109203

Dalmonech, D., Vangi, E., Chiesi, M., Chirici, G., Fibbi, L., Giannetti, F., Marano, G., Massari, C., Nolè, A., Xiao, J., & Collalti, A. (2024). Regional estimates of gross primary production applying the Process-Based Model 3D-CMCC-FEM vs. Remote-Sensing multiple datasets. Europ. J. of Rem. Sens., 57(1). https://doi.org/10.1080/22797254.2023.2301657

De Cáceres, M., Martínez-Vilalta, J., Coll, L., Llorens, P., Casals, P., Poyatos, R., Pausas, J. G., & Brotons, L. (2015). Coupling a water balance model with forest inventory data to predict drought stress: the role of forest structural changes vs. climate changes. Agricul. For. Meteorol., 213, 77–90. https://doi.org/10.1016/j.agrformet.2015.06.012

De Cáceres, M., Mencuccini, M., Martin-StPaul, N., Limousin, J. M., Coll, L., Poyatos, R., Cabon, A., Granda, V., Forner, A., Valladares, F., & Martínez-Vilalta, J. (2021). Unravelling the effect of species mixing on water use and drought stress in Mediterranean forests: A modelling approach. Agricul. For. Meteorol., 296, 108233. https://doi.org/10.1016/j.agrformet.2020.108233

De Cáceres, M., Molowny-Horas, R., Cabon, A., Martínez-Vilalta, J., Mencuccini, M., García-Valdés, R., Nadal-Sala, D., Sabaté, S., Martin-StPaul, N., Morin, X., D'Adamo, F., Batllori, E., & Améztegui, A. (2023). MEDFATE 2.9.3: a trait-enabled model to simulate Mediterranean forest function and dynamics at regional scales. Geosc. Mod. Develop., 16(11), 3165–3201. https://doi.org/10.5194/gmd-16-3165-2023

De Cinti, B., Bombi, P., Ferretti, F., Cantiani, P., Di Salvatore, U., SimonÄiÄ, P., Kutnar, L., Äœeater, M., GarfÃ, V., Mason, F., & Matteucci, G. (2016). From the experience of LIFE+ ManFor C.BD to the Manual of Best Practices in Sustainable Forest Management. It. J. of Agrono., 11(s1), 1–175. https://doi.org/10.4081/ija.2016.789

De Pury, D.G.G., Farquhar, G.D., 1997. Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models. Plant, Cell Environ. 20, 537–557. https://doi.org/10.1111/j.1365-3040.1997.00094.x

De Wergifosse, L., André, F., Goosse, H., Boczon, A., Cecchini, S., Ciceu, A., Collalti, A., Cools, N., D'Andrea, E., De Vos, B., Hamdi, R., Ingerslev, M., Knudsen, M. A., Kowalska, A., Leca, S., Matteucci, G., Nord-Larsen, T., Sanders, T. G., Schmitz, A., Termonia, A., Vanguelova, E., Van Schaeybroeck, B., Verstraeten, A. Vesterdal, L., Jonard, M. (2022). Simulating tree growth response to climate change in structurally diverse oak and beech forests. Sci. Total Environ 806, 150422. https://doi.org/10.1016/j.scitotenv.2021.150422

Deb Burman, P. K., A. G. , P., Chakraborty, S., Tiwari, Y. K., Sarma, D., & Gogoi, N. (2024). Simulating the ecosystem-atmosphere carbon, water and energy fluxes at a subtropical Indian forest using an ecosystem model. Eco. Modell., 490, 110637. https://doi.org/10.1016/j.ecolmodel.2024.110637

Diaconu, D., Kahle, H. P., & Spiecker, H. (2017). Thinning increases drought tolerance of European beech: a case study on two forested slopes on opposite sides of a valley. European J. of For. Res., 136(2), 319–328. https://doi.org/10.1007/s10342-017-1033-8

Dufrene, E., Davi, H., François, C., Maire, G. L., Dantec, V. L., & Granier, A. (2005). Modelling carbon and water cycles in a beech forest. Eco. Modell., 185(2–4), 407–436. https://doi.org/10.1016/j.ecolmodel.2005.01.004

Durrant, H. T., de Rigo, D., Caudullo, G. (2016). Fagus sylvatica and other beeches in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T., Mauri, A. (Eds.), European Atlas of Forest Tree Species. Publ. Off. EU, Luxembourg, pp. e012b90+

Farquhar, G.D., Caemmerer, S., Berry, J.A. (1980). A biochemical model of photosynthetic CO_2 assimilation in leaves of C_3 species. Planta 149, 78–90–90.

Fibbi, L., Moriondo, M., Chiesi, M., Bindi, M., & Maselli, F. (2019). Impacts of climate change on the gross primary production of Italian forests. Ann. of For. Sc., 76(2). https://doi.org/10.1007/s13595-019-0843-x

Gessler, A., Bächli, L., Freund, E. R., Treydte, K., Schaub, M., Haeni, M., Weiler, M., Seeger, S., Marshall, J., Hug, C., Zweifel, R., Hagedorn, F., Rigling, A., Saurer, M., & Meusburger, K. (2021). Drought reduces water uptake in beech from the drying topsoil, but no compensatory uptake occurs from deeper soil layers. New Phyt., 233(1), 194–206. https://doi.org/10.1111/nph.17767

Giuggiola, A., Zweifel, R., Feichtinger, L. M., Vollenweider, P., Bugmann, H., Haeni, M., & Rigling, A. (2018). Competition for water in a xeric forest ecosystem – Effects of understory removal on soil micro-climate, growth and physiology of dominant Scots pine trees. For. Eco. and Manag., 409, 241–249. https://doi.org/10.1016/j.foreco.2017.11.002

Grossiord, C., Buckley, T. N., Cernusak, L. A., Novick, K. A., Poulter, B., Siegwolf, R. T. W., Sperry, J. S., & McDowell, N. G. (2020). Plant responses to rising vapor pressure deficit. New Phyt., 226(6), 1550–1566. https://doi.org/10.1111/nph.16485

Guillemot, J., Delpierre, N., Vallet, P., François, C., Martin-StPaul, N. K., Soudani, K., Nicolas, M., Badeau, V., & Dufrêne, E. (2014). Assessing the effects of management on forest growth across France: insights from a new functional–structural model. Ann. Bot., 114(4), 779–793. https://doi.org/10.1093/aob/mcu059

Hartmann, H., & Trumbore, S. (2016). Understanding the roles of nonstructural carbohydrates in forest trees – from what we can measure to what we want to know. New Phyt., 211(2), 386–403. https://doi.org/10.1111/nph.13955

Hölttä, T., Cochard, H., Nikinmaa, E., & Mencuccini, M. (2009). Capacitive effect of cavitation in xylem conduits: results from a dynamic model. Plant, Cell & Environment/Plant, Cell Environ., 32(1), 10–21. https://doi.org/10.1111/j.1365-3040.2008.01894.x

Jarvis, P. G. (1976). The Interpretation of the Variations in Leaf Water Potential and Stomatal Conductance Found in Canopies in the Field. Philosophical Transactions of the Royal Society B: Biological Sciences, 273(927), 593–610. https://doi.org/10.1098/rstb.1976.0035

Jarvis, N.J., Jansson, P-E., Dik, P.E., Messing, I. (1991). Modelling water and solute transport in macroporous soil. I. Model description and sensitivity analysis. J. Soil Sc., 42, 59–70.

Kane, C., & McAdam, S. (2023). Abscisic acid driven stomatal closure during drought in anisohydric Fagus sylvatica. J. Pl. Hydr., 9, 002. https://doi.org/10.20870/jph.2023.002

Kattge, J., Knorr, W., (2007). Temperature acclimation in a biochemical model of photosynthesis: A reanalysis of data from 36 species. Plant, Cell Environ. 30, 1176–1190. https://doi.org/10.1111/j.1365-3040.2007.01690.x

Kimmins, (Hamish) J. P., Blanco, J. A., Seely, B., Welham, C., & Scoullar, K. (2008). Complexity in modelling forest ecosystems: How much is enough? For. Eco. Manag., 256(10), 1646–1658. https://doi.org/10.1016/j.foreco.2008.03.011

Kramer, K., Leinonen, I., Bartelink, H., Berbigier, P., Borghetti, M., Bernhofer, C., Cienciala, E., Dolman, A., Froer, O., Gracia, A., Granier, A., Grünwald, T., Hari, P., Jans, W., Kellomäki, S., Loustau, D., Magnani, F., Markkanen, T., Matteucci, G., Mohren, G.M.J., Moors, E., Nissinen, A., Peltola, H. Sabaté, S., Sanchez, A. (2002). Evaluation of six process-based forest growth models using eddy-covariance measurements of CO_2 and H_2O fluxes at six forest sites in Europe. Glob. Ch. Biol., 8(3), 213–230. https://doi.org/10.1046/j.1365-2486.2002.00471.x

Langer, G. J., & Bußkamp, J. (2023). Vitality loss of beech: a serious threat to Fagus sylvatica in Germany in the context of global warming. Journal of Plant Diseases and Protection, 130(5), 1101–1115. https://doi.org/10.1007/s41348-023-00743-7

Larsbo, M., Roulier, S., Stenemo, F., Kasteel, R., Jarvis, N. (2005). An Improved Dual-Permeability Model of Water Flow and Solute Transport in the Vadose Zone. Vadose Zone Journal, 4, 398–406.

Leuning, R. (2002). Temperature dependence of two parameters in a photosynthesis model. Plant, Cell Environ., 25, 1205–1210.

Leuschner, C., Schipka, F., & Backes, K. (2021). Stomatal regulation and water potential variation in European beech: challenging the iso/anisohydry concept. Tree Phys., 42(2), 365–378. https://doi.org/10.1093/treephys/tpab104

Loescher, H. W., Law, B. E., Mahrt, L., Hollinger, D. Y., Campbell, J., & Wofsy, S. C. (2006). Uncertainties in, and interpretation of, carbon flux estimates using the eddy covariance technique. J. of Geoph. Res. Atm., 111(D21). https://doi.org/10.1029/2005jd006932

Lombardi, D., Micalizzi, K., & Vitale, M. (2023). Assessing carbon and water fluxes in a mixed Mediterranean protected forest under climate change: An integrated bottom – up and top – down approach. Ecol. Infor., 78, 102318. https://doi.org/10.1016/j.ecoinf.2023.102318

Lüttschwager, D., & Jochheim, H. (2020). Drought Primarily Reduces Canopy Transpiration of Exposed Beech Trees and Decreases the Share of Water Uptake from Deeper Soil Layers. Forests, 11(5), 537. https://doi.org/10.3390/f11050537

Huber, M. O., Eastaugh, C. S., Gschwantner, T., Hasenauer, H., Kindermann, G., Ledermann, T., Lexer, M. J., Rammer, W., Schörghuber, S., & Sterba, H. (2013). Comparing simulations of three conceptually different forest models with National Forest Inventory data. Environ. Modell. Soft., 40, 88–97. https://doi.org/10.1016/j.envsoft.2012.08.003

INFC (2015). Inventario Nazionale delle Foreste e dei Serbatoi Forestali di Carbonio. Ministero delle Politiche Agricole Alimentari e Forestali, Ispettorato Generale – Corpo Forestale dello Stato, CRA – Istituto Sperimentale per l'Assestamento Forestale e per l'Alpicoltura. [online] URL: https://www.inventarioforestale.org/it/statistiche_infc/

Italiano, S. S., Camarero, J. J., Borghetti, M., Colangelo, M., Rita, A., & Ripullone, F. (2024). Drought legacies in mixed Mediterranean forests: Analysing the effects of structural overshoot, functional traits and site factors. Sci. Total Environ 927, 172166. https://doi.org/10.1016/j.scitotenv.2024.172166

Mahnken, M., Cailleret, M., Collalti, A., Trotta, C., Biondo, C., D'Andrea, E., Dalmonech, D., Marano, G., Mäkelä, A., Minunno, F., Peltoniemi, M., Trotsiuk, V., Nadal- Sala, D., Sabaté, S., Vallet, P., Aussenac, R., Cameron, D. R., Bohn, F. J., Grote, R., Augustynczik, A. L. D., Yousefpour, R., Huber, N., Bugmann, H., Merganičová, K., Merganic, J., Valent, P., Lasch-Born, P., Hartig, F., Vega del Valle, I. D., Volkholz, J., Gutsch, M., Matteucci, G., Krejza, J., Ibrom, A., Meesenburg, H., Rötzer, T., van der Maaten-Theunissen, M., van der Maaten, E., & Reyer, C. P. O. (2022). Accuracy, realism and general applicability of European forest models. Glob. Chan. Biol., 28(23), 6921–6943. https://doi.org/10.1111/gcb.16384

Marconi S., Chiti T., Nolè A., Valentini R., Collalti A. (2017). The role of Respiration in estimation of the net Carbon cycle: coupling soil Carbon dynamics and canopy turnover in a novel version of 3D-CMCC Forest Ecosystem Model, Forests, 8: 220. https://doi.org/10.3390/f8060220

Maréchaux, I., Langerwisch, F., Huth, A., Bugmann, H., Morin, X., Reyer, C. P., Seidl, R., Collalti, A., De Paula, M. D., Fischer, R., Gutsch, M., Lexer, M. J., Lischke, H., Rammig, A., Rödig, E., Sakschewski, B., Taubert, F., Thonicke, K., Vacchiano, G., & Bohn, F. J. (2021). Tackling unresolved questions in forest ecology: The past and future role of simulation models. Ecol. and Evo., 11(9), 3746–3770. https://doi.org/10.1002/ece3.7391

Margalef-Marrase, J., Pérez-Navarro, M. Á., & Lloret, F. (2020b). Relationship between heatwave induced forest die-off and climatic suitability in multiple tree species. Glob. Chan. Biol., 26(5), 3134–3146. https://doi.org/10.1111/gcb.15042

Martínez-Vilalta, J., Sala, A., Asensio, D., Galiano, L., Hoch, G., Palacio, S., Piper, F. I., & Lloret, F. (2016). Dynamics of non-structural carbohydrates in terrestrial plants: a global synthesis. Ecol. Monog., 86(4), 495–516. https://doi.org/10.1002/ecm.1231

Medlyn, B. E., Badeck, F. W., De Pury, D. G. G., Barton, C. V. M., Broadmeadow, M., Ceulemans, R., De Angelis, P., Forstreuter, M., Jach, M. E., Kellomäki, S., Laitat, E., Marek, M., Philippot, S., Rey, A., Strassemeyer, J., Laitinen, K., Liozon, R., Portier, B., Roberntz, P., Wang, K., Jarvis, P. G. (1999). Effects of elevated [CO2] on photosynthesis in European forest species: a meta-analysis of model parameters. Plant Cell Environ., 22(12), 1475–1495. https://doi.org/10.1046/j.1365-3040.1999.00523.x

Meier, I. C., Knutzen, F., Eder, L. M., Müller-Haubold, H., Goebel, M., Bachmann, J., Hertel, D., & Leuschner, C. (2017). The Deep Root System of Fagus sylvatica on Sandy Soil: Structure and Variation Across a Precipitation Gradient. Ecosystems, 21(2), 280–296. https://doi.org/10.1007/s10021-017-0148-6

Merganičová, K., Merganič, J., Lehtonen, A., Vacchiano, G., Sever, M. Z. O., Augustynczik, A. L. D., Grote, R., Kyselová, I., Mäkelä, A., Yousefpour, R., Krejza, J., Collalti, A., & Reyer, C. P. O. (2019). Forest carbon allocation modelling under climate change. Tree Phy., 39(12), 1937–1960. https://doi.org/10.1093/treephys/tpz105

Monteith, J. L. and M. H. Unsworth (2008). Principles of Environmental Physics, 3rd Edition. Burlington, MA, Academic Press https://doi.org/10.3832/ifor1802-009

Morichetti, M., Vangi, E., & Collalti, A. (2024). Predicted Future Changes in the Mean Seasonal Carbon Cycle Due to Climate Change. Forests, 15(7), 1124. https://doi.org/10.3390/f15071124

Noce, S., Collalti, A., & Santini, M. (2017). Likelihood of changes in forest species suitability, distribution, and diversity under future climate: The case of Southern Europe. Ecol. and Evo., 7(22), 9358–9375. https://doi.org/10.1002/ece3.3427

Noce, S., Cipriano, C., & Santini, M. (2023). Altitudinal shifting of major forest tree species in Italian mountains under climate change. Front. in For. and Glob. Chan., 6. https://doi.org/10.3389/ffgc.2023.1250651

Nolè, A., Collalti, A., Magnani, F. et al. Assessing temporal variation of primary and ecosystem production in two Mediterranean forests using a modified 3-PG

model. Ann. For. Sc. 70, 729–741 (2013). https://doi.org/10.1007/s13595-013-0315-7

Nolè, A. et al. (2015). The Role of Managed Forest Ecosystems: A Modeling Based Approach. In: Valentini, R., Miglietta, F. (eds) The Greenhouse Gas Balance of Italy. Environmental Science and Engineering(). Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-32424-6_5

Papale, D., Reichstein, M., Aubinet, M., Canfora, E., Bernhofer, C., Kutsch, W. L., Longdoz, B., Rambal, S., Valentini, R., Vesala, T., & Yakir, D. (2006). Towards a standardized processing of Net Ecosystem Exchange measured with eddy covariance technique: algorithms and uncertainty estimation. Biogeosciences, 3(4), 571–583. https://doi.org/10.5194/bg-3-571-2006

Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Reichstein, M., Ribeca, A., van Ingen, C., Vuichard, N., Zhang, L., Amiro, B., Ammann, C., Arain, M.A., Ardö, J., Arkebauer, T., Arndt, S.K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Belelli Marchesini, L., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T.A., Blanken, P.D., Bohrer, G., Boike, J., Bolstad, P.V., Bonal, D., Bonnefond, J.-M., Bowling, D.R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns, S.P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T.R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B.D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P.S., D'Andrea, E., da Rocha, H., Dai, X., Davis, K.J., De Cinti, B., de Grandcourt, A., De Ligne, A., De Oliveira, R.C., Delpierre, N., Desai, A.R., Di Bella, C.M., di Tommasi, P., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H.A.M., Eugster, W., Ewenz, C.M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., Gielen, B., Gioli, B., Gitelson, A., Goded, I., Goeckede, M., Goldstein, A.H., Gough, C.M., Goulden, M.L., Graf, A., Griebel, A., Gruening, C., Grünwald, T., Hammerle, A., Han, S., Han, X., Hansen, B.U., Hanson, C., Hatakka, J., He, Y., Hehn, M., Heinesch, B., Hinko-Najera, N., Hörtnagl, L., Hutley, L., Ibrom, A., Ikawa, H., Jackowicz-Korczynski, M., Janouš, D., Jans, W., Jassal, R., Jiang, S., Kato, T., Khomik, M., Klatt, J., Knohl, A., Knox, S., Kobayashi, H., Koerber, G., Kolle, O., Kosugi, Y., Kotani, A., Kowalski, A., Kruijt, B., Kurbatova, J., Kutsch, W.L., Kwon, H., Launiainen, S., Laurila, T., Law, B., Leuning, R., Li, Y., Liddell, M., Limousin, J.-M., Lion, M., Liska, A.J., Lohila, A., López-Ballesteros, A., López-Blanco, E., Loubet, B., Loustau, D., Lucas-Moffat, A., Lüers, J., Ma, S., Macfarlane, C., Magliulo, V., Maier, R., Mammarella, I., Manca, G., Marcolla, B., Margolis, H.A., Marras, S., Massman, W., Mastepanov, M., Matamala, R., Matthes, J.H., Mazzenga, F., McCaughey, H., McHugh, I., McMillan, A.M.S., Merbold, L., Meyer, W., Meyers, T., Miller, S.D., Minerbi, S., Moderow, U., Monson, R.K., Montagnani, L., Moore, C.E., Moors, E., Moreaux, V., Moureaux, C., Munger, J.W., Nakai, T., Neirynck, J., Nesic, Z., Nicolini, G., Noormets, A., Northwood, M., Nosetto, M., Nouvellon, Y., Novick, K., Oechel,

W., Olesen, J.E., Ourcival, J.-M., Papuga, S.A., Parmentier, F.-J., Paul-Limoges, E., Pavelka, M., Peichl, M., Pendall, E., Phillips, R.P., Pilegaard, K., Pirk, N., Posse, G., Powell, T., Prasse, H., Prober, S.M., Rambal, S., Rannik, Ü., Raz-Yaseef, N., Rebmann, C., Reed, D., Resco de Dios, V., Restrepo-Coupe, N., Reverter, B.R., Roland, M., Sabbatini, S., Sachs, T., Saleska, S.R., Sánchez-Cañete, E.P., Sanchez-Mejia, Z.M., Schmid, H.P., Schmidt, M., Schneider, K., Schrader, F., Schroder, I., Scott, R.L., Sedlák, P., Serrano-Ortíz, P., Shao, C., Shi, P., Shironya, I., Siebicke, L., Šigut, L., Silberstein, R., Sirca, C., Spano, D., Steinbrecher, R., Stevens, R.M., Sturtevant, C., Suyker, A., Tagesson, T., Takanashi, S., Tang, Y., Tapper, N., Thom, J., Tomassucci, M., Tuovinen, J.-P., Urbanski, S., Valentini, R., van der Molen, M., van Gorsel, E., van Huissteden, K., Varlagin, A., Verfaillie, J., Vesala, T., Vincke, C., Vitale, D., Vygodskaya, N., Walker, J.P., Walter-Shea, E., Wang, H., Weber, R., Westermann, S., Wille, C., Wofsy, S., Wohlfahrt, G., Wolf, S., Woodgate, W., Li, Y., Zampedri, R., Zhang, J., Zhou, G., Zona, D., Agarwal, D., Biraud, S., Torn, M., & Papale, D. (2020). The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data. Scientific Data, 7(1). https://doi.org/10.1038/s41597-020-0534-3

Pan, Y., Birdsey, R. A., Phillips, O. L., Houghton, R. A., Fang, J., Kauppi, P. E., Keith, H., Kurz, W. A., Ito, A., Lewis, S. L., Nabuurs, G. J., Shvidenko, A., Hashimoto, S., Lerink, B., Schepaschenko, D., Castanho, A., & Murdiyarso, D. (2024). The enduring world forest carbon sink. Nature, 631(8021), 563–569. https://doi.org/10.1038/s41586-024-07602-x

Peano D., Materia S., Collalti A., Alessandri A., Anav A., Bombelli A., Gualdi S. (2019). Global variability of simulated and observed vegetation growing season. Journal of Geophysical Research, Biogeosciences, 124, 3569–3587, https://doi.org/10.1029/2018JG004881

Petrik, P., Petek-Petrik, A., Kurjak, D., Mukarram, M., Klein, T., Gömöry, D., Střelcová, K., Frýdl, J., & Konôpková, A. (2022). Interannual adjustments in stomatal and leaf morphological traits of European beech (Fagus sylvatica L.) demonstrate its climate change acclimation potential. Plant Biol., 24(7), 1287–1296. https://doi.org/10.1111/plb.13401

Pietsch, S. A., Hasenauer, H., & Thornton, P. E. (2005). BGC-model parameters for tree species growing in central European forests. For. Eco. Manag., 211(3), 264–295.https://doi.org/10.1016/j.foreco.2005.02.046

Pilli, R., Alkama, R., Cescatti, A., Kurz, W. A., & Grassi, G. (2022). The European forest carbon budget under future climate conditions and current management practices. Biogeosciences, 19(13), 3263–3284. https://doi.org/10.5194/bg-19-3263-2022

Piovesan, G., Biondi, F., Di Filippo, A., Alessandrini, A., & Maugeri, M. (2008). Drought- driven growth reduction in old beech (Fagus sylvatica L.) forests of the central Apennines, Italy. Glob. Chan. Biol., 14(6), 1265–1281. https://doi.org/10.1111/j.1365-2486.2008.01570.x

Priwitzer, T., Kurjak, D., Kmeť, J., Sitková, Z., & Leštianska, A. (2014). Photosynthetic response of European beech to atmospheric and soil drought. Central

European Forestry Journal, 60(1), 32–38. https://doi.org/10.2478/forj-2014-0003

Puchi, P. F., Dalmonech, D., Vangi, E., Battipaglia, G., Tognetti, R., & Collalti, A. (2024). Contrasting patterns of water use efficiency and annual radial growth among European beech forests along the Italian peninsula. Sc. Rep., 14(1). https://doi.org/10.1038/s41598-024-57293-7

Pukkala, T. (2016). Which type of forest management provides most ecosystem services? For. Ecos., 3(1). https://doi.org/10.1186/s40663-016-0068-5

Raffa, M., Adinolfi, M., Reder, A. et al. (2023). Very High Resolution Projections over Italy under different CMIP5 IPCC scenarios. Sci. Data 10, 238 . https://doi.org/10.1038/s41597-023-02144-9

Reyer, C., Lasch-Born, P., Suckow, F., Gutsch, M., Murawski, A., & Pilz, T. (2013). Projections of regional changes in forest net primary productivity for different tree species in Europe driven by climate change and carbon dioxide. Ann. For. Sc., 71(2), 211–225. https://doi.org/10.1007/s13595-013-0306-8

Reyer, C., Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-Born, P., Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders, T., Kolari, P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., Matteucci, G., Collalti, A., D'Andrea, E., Foltýnová, L., Krejza, J., Ibrom, A., Pilegaard, K., Loustau, D., Bonnefond, J.-M., Berbigier, P., Picart, D., Lafont, S., Dietze, M., Cameron, D., Vieno, M., Tian, H., Palacios-Orueta, A., Cicuendez, V., Recuero, L., Wiese, K., Büchner, M., Lange, S., Volkholz, J., Kim, H., Weedon, G., Sheffield, J., Vega del Valle, I., Suckow, F., Horemans, J., Martel, S., Bohn, F., Steinkamp, J., Chikalanov, A., Mahnken, M., Gutsch, M., Trotta, C., Babst, F., & Frieler, K. (2020a). The PROFOUND database for evaluating vegetation models and simulating climate impacts on European forests version V. 0.3. GFZ Data Services. https://doi.org/10.5880/PIK.2020.006

Reyer, C. P. O., Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-Born, P., Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders, T. G. M., Kolari, P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., Collalti, A., Trotta, C., Matteucci, G., D'Andrea, E., Foltýnová, L., Krejza, J., Ibrom, A., Pilegaard, K., Loustau, D., Bonnefond, J.-M., Berbigier, P., Picart, D., Lafont, S., Dietze, M., Cameron, D., Vieno, M., Tian, H., Palacios-Orueta, A., Cicuendez, V., Recuero, L., Wiese, K., Büchner, M., Lange, S., Volkholz, J., Kim, H., Horemans, J. A., Bohn, F., Steinkamp, J., Chikalanov, A., Weedon, G. P., Sheffield, J., Babst, F., Vega del Valle, I., Suckow, F., Martel, S., Mahnken, M., Gutsch, M., & Frieler, K. (2020b). The PROFOUND database for evaluating vegetation models and simulating climate impacts on European forests. Earth System Science Data, 12(2), 1295–1320. https://doi.org/10.5194/essd-12-1295-2020

Rezaie, N., D'Andrea, E., Bräuning, A., Matteucci, G., Bombi, P., & Lauteri, M. (2018). Do atmospheric CO2 concentration increase, climate and forest management affect iWUE of common beech? Evidences from carbon isotope analyses in tree rings. Tree Phys., 38(8), 1110–1126. https://doi.org/10.1093/treephys/tpy025

Richardson, A.D., Black, T.A., Ciais, P., Delbart, N., Friedl, M.A., Gobron, N., Hollinger, D.Y., Kutsch, W.L., Longdoz, B., Luyssaert, S., Migliavacca, M., Montagnani, L., Munger, J.W., Moors, E., Piao, S., Rebmann, C., Reichstein, M., Saigusa, N., Tomelleri, E., Vargas, R., & Varlagin, A. (2010). Influence of spring and autumn phenological transitions on forest ecosystem productivity. Philos. Trans. – Royal Society. Biological Sciences, 365(1555), 3227–3246. https://doi.org/10.1098/rstb.2010.0102

Rita, A., Bonanomi, G., Allevato, E., Borghetti, M., Cesarano, G., Mogavero, V., Rossi, S., Saulino, L., Zotti, M., & Saracino, A. (2021). Topography modulates near-ground microclimate in the Mediterranean Fagus sylvatica treeline. Sc. Rep.s, 11(1). https://doi.org/10.1038/s41598-021-87661-6

Riviere, M., Caurla, S., & Delacote, P. (2020). Evolving Integrated Models From Narrower Economic Tools: the Example of Forest Sector Models. Environmental Model. & Assess., 25(4), 453–469. https://doi.org/10.1007/s10666-020-09706-w

Rötzer, T., Dieler, J., Mette, T., Moshammer, R., & Pretzsch, H. (2010). Productivity and carbon dynamics in managed Central European forests depending on site conditions and thinning regimes. Forestry, 83(5), 483–496. https://doi.org/10.1093/forestry/cpq031

Ruffault, J., Pimont, F., Cochard, H., Dupuy, J.-L. & Martin-StPaul, N. (2022). SurEau-Ecos v2.0: A trait-based plant hydraulics model for simulations of plant water status and drought-induced mortality at the ecosystem level. Geosc. Mod. Develop., 15, 5593–5626.

Skrk, N., Martinez del Castillo, E., Serrano-Notivoli, R., Luis, M. De, Novak, K., Merela, M., Cufar, K. (2023). Spatial and temporal variation of Fagus sylvatica growth in marginal areas under progressive climate change. Dendrochronologia 81. https://doi.org/10.1016/j.dendro.2023.126135

Sánchez-Dávila, J., De Cáceres, M., Vayreda, J., & Retana, J. (2024). Regional patterns and drivers of modelled water flows along environmental, functional, and stand structure gradients in Spanish forests. Hyd. and Earth Sys. Sc., 28(13), 3037–3050. https://doi.org/10.5194/hess-28-3037-2024

Schmied, G., Pretzsch, H., Ambs, D., Uhl, E., Schmucker, J., Fäth, J., Biber, P., Hoffmann, Y. D., Šeho, M., Mellert, K. H., & Hilmers, T. (2023). Rapid beech decline under recurrent drought stress: Individual neighborhood structure and soil properties matter. Forest Ecology and Management, 545, 121305. https://doi.org/10.1016/j.foreco.2023.121305

Schwalm, C.R. and Ek, A.R. (2004). A process-based model of forest ecosystems driven by meteorology. Ecol. Modell., 179, 317–348. https://doi.org/10.1016/j.ecolmodel.2004.04.016

Simonin, K., Kolb, T., Montes-Helu, M., & Koch, G. (2007). The influence of thinning on components of stand water balance in a ponderosa pine forest stand during and after extreme drought. Agricul. For. Met., 143(3–4), 266–276. https://doi.org/10.1016/j.agrformet.2007.01.003

Sperlich, D., Nadal-Sala, D., Gracia, C., Kreuzwieser, J., Hanewinkel, M., & Yousefpour, R. (2020). Gains or Losses in Forest Productivity under Climate

Change? The Uncertainty of CO_2 Fertilization and Climate Effects. Climate, 8(12), 141. https://doi.org/10.3390/cli8120141

Sperry, J. S., Adler, F. R., Campbell, G. S., & Comstock, J. P. (1998). Limitation of plant water use by rhizosphere and xylem conductance: results from a model. Plant, Cell & Environment/Plant, Cell Environ., 21(4), 347–359. https://doi.org/10.1046/j.1365-3040.1998.00287.x

Sperry, J.S., Venturas, M.D., Anderegg, W.R.L., Mencuccini, M., Mackay, D.S., Wang, Y., et al. (2017). Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost. Plant, Cell Environ., 40, 816–830. https://doi: 10.1111/pce.12852.

Tegel, W., Seim, A., Hakelberg, D., Hoffmann, S., Panev, M., Westphal, T., Büntgen, U. (2014). A recent growth increase of European beech (Fagus sylvatica L.) at its Mediterranean distribution limit contradicts drought stress. Eur. J. For. Res. 133, 61–71. https://doi.org/10.1007/s10342-013-0737-7

Testolin, R., Dalmonech, D., Marano, G., Bagnara, M., D'Andrea, E., Matteucci, G., Noce, S., Collalti, A. (2023). Simulating diverse forest management options in a changing climate on a Pinus nigra subsp. laricio plantation in Southern Italy. Sci. Total Environ. 857, 159361. https://doi.org/10.1016/j.scitotenv.2022.159361

Thom, D., Buras, A., Heym, M., Klemmt, H. J., & Wauer, A. (2023). Varying growth response of Central European tree species to the extraordinary drought period of 2018 – 2020. Agricul. For. Meteorol., 338, 109506. https://doi.org/10.1016/j.agrformet.2023.109506

Tognetti, R., Lasserre, B., Di Febbraro, M., & Marchetti, M. (2019). Modeling regional drought-stress indices for beech forests in Mediterranean mountains based on tree-ring data. Agricul. For. Meteorol.,, 265, 110–120. https://doi.org/10.1016/j.agrformet.2018.11.015

Vacchiano, G., Magnani, F., & Collalti, A. (2012). Modeling Italian forests: state of the art and future challenges. IForest, 5(1), 113–120. https://doi.org/10.3832/ifor0614-005

Vangi, E., Dalmonech, D., Morichetti, M., Grieco, E., Giannetti, F., D'Amico, G., Nakhavali, M., Chirici, G., & Collalti, A. (2024a). Stand Age and Climate Change Effects on Carbon Increments and Stock Dynamics. Forests, 15(7), 1120. https://doi.org/10.3390/f15071120

Vangi, E., Dalmonech, D., Cioccolo, E., Marano, G., Bianchini, L., Puchi, P. F., Grieco, E., Cescatti, A., Colantoni, A., Chirici, G., & Collalti, A. (2024b). Stand age diversity (and more than climate change) affects forests' resilience and stability, although unevenly. J. of Environ. Man., 366, 121822. https://doi.org/10.1016/j.jenvman.2024.121822

Verbeeck, H., Samson, R., Granier, A. A., Montpied, P., & Lemeur, R. (2008). Multi-year model analysis of GPP in a temperate beech forest in France. Ecol. Modell., 210(1–2), 85–103. https://doi.org/10.1016/j.ecolmodel.2007.07.010

Wang, S., Zhang, Y., Ju, W., Chen, J. M., Ciais, P., Cescatti, A., Sardans, J., Janssens, I. A., Wu, M., Berry, J. A., Campbell, E., Fernández-Martínez, M., Alkama, R., Sitch, S., Friedlingstein, P., Smith, W. K., Yuan, W., He, W., Lombardozzi, D., ... Peñuelas, J. (2020). Recent global decline of CO_2 fertilization effects on

vegetation photosynthesis. Science, 370(6522), 1295–1300. https://doi.org/10.1126/science.abb7772

Waring, R. H. and S. W. Running (2007). Forest Ecosystems: Analysis at Multiple Scales. San Francisco, CA, Elsevier Academic Press.

Yu, X., Orth, R., Reichstein, M., Bahn, M., Klosterhalfen, A., Knohl, A., Koebsch, F., Migliavacca, M., Mund, M., Nelson, J. A., Stocker, B. D., Walther, S., & Bastos, A. (2022). Contrasting drought legacy effects on gross primary productivity in a mixed versus pure beech forest. Biogeosciences, 19(17), 4315–4329. https://doi.org/10.5194/bg-19-4315-2022

Zaehle, S., Medlyn, B. E., De Kauwe, M. G., Walker, A. P., Dietze, M. C., Hickler, T., Luo, Y., Wang, Y., El-Masri, B., Thornton, P., Jain, A., Wang, S., Warlind, D., Weng, E., Parton, W., Iversen, C. M., Gallet-Budynek, A., McCarthy, H., Finzi, A., . . . Norby, R. J. (2014). Evaluation of 11 terrestrial carbon–nitrogen cycle models against observations from two temperate Free-Air CO2 Enrichment studies. New Phyt., 202(3), https://doi.org/10.1111/nph.12697

Zuccarini, P., Delpierre, N., Mariën, B., Peñuelas, J., Heinecke, T., Campioli, M. (2023). Drivers and dynamics of foliar senescence in temperate deciduous forest trees at their southern limit of distribution in Europe. Agricul. For. Meteorol. 342, 109716. https://doi.org/10.1016/j.agrformet.2023.109716

Fig 1. Map of the study sites. Triangles represent sites for validating fluxes, while the squares represent sites for management investigation.

1319

1320
1321 Fig 2. Daily mean variations of GPP (gC m⁻² day⁻¹) and LE (MJ m⁻² day⁻¹) estimated from the direct
1322 micrometeorological eddy covariance measurements (GPP-Obs and LE-Obs) and models' simulation
1323 (GPP-3D-CMCC-FEM, LE-3D-CMCC-FEM and, GPP-MEDFATE, LE-MEDFATE) during the evaluation
1324 period at the DK-Sor, IT-Col and FR-Hes at the Beech forest in 2006-2010 and 2014-2018, respectively.
1325
1326
1327

1328

1329

1330

1331

Fig 3. Scatter plots and linear regressions of GPP (gC m⁻² day⁻¹) and LE (MJ m⁻² day⁻¹) of the models versus the direct micrometeorological eddy covariance measurements (Obs) at the Sorø (DK–Sor; 2006–2010 period), Collelongo (IT–Col; 2006–2010 period) and Hesse (FR–Hes; 2014–2018 period).

1335

1336

Fig 4. Comparative analysis between models output at the Cansiglio site. The top-left panel displays the
PLC_{stem} as modelled by MEDFATE, while the top-right panel shows the modelled LAI for 3D-CMCC-FEM
(and used by MEDFATE). The middle-up panels (left and right) present annual GPP (gC m^{-2} year^{-1}) as
modelled by the MEDFATE and 3D-CMCC-FEM, respectively. The middle-down panels (left and right)
depict annual LE (MJ m^{-2} yr^{-1}) modelled by the MEDFATE and 3D-CMCC-FEM, respectively. The
bottom panels (left and right) depict the annual minimum of NSC concentration (%) at the stand and tree
level, respectively, as modelled by the 3D-CMCC-FEM. Different plot management strategies are
represented by distinct line styles: solid lines for 'Control' plots ('no management'), dotted lines for
'Innovative' plots, and dashed lines for 'Traditional' plots (Shelterwood). Climate scenarios are indicated
by line colours: black for 'Hist' climate data (2010-2022), orange and blue for RCP4.5 and RCP8.5 climate
(2059-2070), respectively.
1348

1349

Fig 5. Comparative analysis between models output at the Mongiana site. The top-left panel displays the
percent loss of PLC_{stem} as modelled by MEDFATE while the top-right panel shows the modelled LAI for
3D-CMCC-FEM (and used by MEDFATE). The middle-up panels (left and right) present annual GPP (gC
m^{-2} $year^{-1}$) as modelled by the MEDFATE and 3D-CMCC-FEM, respectively. The middle-down panels
(left and right) depict annual LE (MJ m^{-2} yr^{-1}) modelled by the MEDFATE and 3D-CMCC-FEM,
respectively. The bottom panels (left and right) depict the annual minimum of NSC concentration (%) at
the stand and tree level, respectively, as modelled by the 3D-CMCC-FEM. Different plot management
strategies are represented by distinct line styles: solid lines with circles for 'Control' plots ('no
management'), dotted lines with squares for 'Innovative' plots, dashed lines with stars for 'Traditional'
plots (Shelterwood) and dash-dotted lines with triangles for 'SM' management. Climate scenarios are
indicated by line colours: black for 'Hist' climate data (2010-2022), orange and blue for RCP4.5 and
RCP8.5 climate (2060-2070), respectively.

SITE DESCRIPTION					
	Evaluation sites			Test sites	
Variable	DK-Sor	FR-Hes	IT-Col	Cansiglio	Mongiana
Coordinates (WGS84)	55°49'N, 11°64' E	48°66'N, 7°08'E	41°85'N, 13°59'E	46°02'N, 12°22'E	38°29'N, 16°14'E
Country	Denmark	France	Italy	Italy	Italy
Altitude (m a.s.l.)	40	305	1500	1300	1300
Area (ha)	1	1	1	27	27
MAT (°C)	8.52	10.27	6.95	6.44	11.01
MAP (mm)	818	853	1075	2219	1701
Slope (%)	–	5	35	12	10
Aspect (°)	0	0	252	135	135
Stand age (yr)	90	45	118	120	90
Summer prec (J-J-A)(mm)	292	205	120	493	141

1367   Table 1. Characteristics of the study sites. The age of the stands refers to 2010. The mean annual
1368   temperature (MAT) and mean annual precipitation (MAP) for DK-Sor, FR-Hes and IT-Col refer to the
1369   period evaluated (i.e., 2006-2010 for the Sorø and Collelongo site and 2014-2018 for the Hesse site)
1370   while for Cansiglio and Mongiana from 2010 to 2022. The sum of precipitation in summer refers to June
1371   (J), July (J) and August (A) for the same period.

1372

PARAMETERS AND VARIABLES		
Name	**Value**	**Unit**
$g_{s\,max}$	0.006 Pietsch et al. (2005)	$m\,s^{-1}$
J_{max}	–160 De Cáceres et al. (2023)	$\mu mol\ photons\ m^{-2}\,s^{-1}$
V_{cmax}	–95 De Cáceres et al. (2023)	$\mu mol\ CO_2\ m^{-2}\,s^{-1}$
LAI	from 3D–CMCC–FEM to MEDFATE	$m^2\,m^{-2}$
ASW	from 3D–CMCC–FEM to MEDFATE	mm

1376   Table 2. Parameters and variables set for both models during the simulations.

	3D–CMCC–FEM						MEDFATE					
	GPP			LE			GPP			LE		
	R^2	RMSE	MAE	R^2	RMSE	MAE	R^2	RMSE	MAE	R^2	RMSE	MAE
DK–Sor	0.91	2.17	1.65	0.85	2.02	1.58	0.85	2.52	1.97	0.62	3.09	2.39
FR–H	0.76	3.30	2.46	0.89	2.09	1.47	0.76	2.80	2.21	0.69	3.31	2.37

es												
IT – Col	0.83	2.09	1.56	0.84	2.05	1.57	0.68	2.32	1.87	0.77	2.64	1.90

1381 Table 3. The correlation coefficient (R^2), the Root Mean Square Error (RMSE) and the Mean Absolute Error
1382 (MAE) for, the GPP (gC m^{-2}day^{-1}) and LE (MJ m^{-2} day^{-1}) of the daily simulations at DK–Sor, IT–Col, and
1383 FR–Hes sites performed from both models 3D–CMCC–FEM and MEDFATE in the beech forest stands.
1384
1385