Assessing models' sensitivity to the effects of forest management and climate change on carbon and water fluxes in European beech forests

Vincenzo Saponaro[1,2*], Miquel De Cáceres[3], Daniela Dalmonech[1,4], Ettore D'Andrea[4,5], Elia Vangi[1], Alessio Collalti[1,4]

Affiliations:

1. Forest Modelling Lab., Institute for Agriculture and Forestry Systems in the Mediterranean, National Research Council of Italy (CNR–ISAFOM), Perugia, Italy;
2. Department for Innovation in Biological, Agri–Food and Forest Systems (DIBAF), University of Tuscia, Viterbo, Italy;
3. CREAF, E08193 Bellaterra (Cerdanyola del Vallès), Catalonia, Spain
4. National Biodiversity Future Center (NBFC), Palermo, Italy
5. Research Institute on Terrestrial Ecosystems, National Research Council of Italy (CNR–IRET), Porano, Italy

Corresponding author: vincenzo.saponaro@isafom.cnr.it

Highlights

- Modelled carbon and water fluxes under different climates and management regimes

- Different climates increases fluxes in the north and decreases them in the south

- 3D-CMCC–FEM and MEDFATE satisfactorily predicted productivity and latent heat

- 3D-CMCC–FEM predicts carbon starvation, MEDFATE predicts stem embolism in the south

- High thinning intensity of the stand in the south negatively affected carbon fluxes

Abstract

The consequences of climate change continue to threaten European forests,

particularly for species located at the edges of their latitudinal and altitudinal ranges.

While extensively studied in Central Europe, European beech forests require further

investigation to understand how climate change will affect these ecosystems in

Mediterranean areas. Proposed silvicultural options increasingly aim at sustainable

management to reduce biotic and abiotic stresses and enhance these forest

ecosystems' resistance and resilience mechanisms. Process-based models (PBMs) can

help us to simulate such phenomena and capture early stress signals while considering

the effect of different management approaches. In this study, we focus on estimating

sensitivity of two state-of-the-art PBMs forest models by simulating carbon and water

fluxes at the stand level to assess productivity changes and feedback resulting from

different climatic forcings. Utilizing 3D-CMCC-FEM and MEDFATE models, we

simulated and analyzed carbon (C) and water (H_2O) fluxes in diverse forest plots under

managed vs. unmanaged scenarios and under current climate and different climatic

forcings (RCP4.5 and RCP8.5), in two sites, on the Italian peninsula, Cansiglio in the

north and Mongiana in the south. To ensure confidence in the models' results, we first

evaluated their performance in simulating C and H_2O flux in three additional beech

forests along a latitudinal gradient spanning from Denmark to central Italy. The results

from both models for C and H_2O flux assessment showed generally good model

accuracy. At the Cansiglio site in northern Italy, both models simulated a general

increase in C and H_2O fluxes under the RCP8.5 climate scenario compared to the

current climate. Still, no benefit in managed plots compared to unmanaged ones, as the

site does not have water availability limitations, and thus, competition for water is low.

At the Mongiana site in southern Italy, both models simulate a decrease in C and H_2O

fluxes and sensitivity to the different climatic forcings compared to the current climate,

with an increase in C and H_2O fluxes considering specific management regimes

compared to unmanaged scenarios. Conversely, in both models, under unmanaged

scenarios, plots are simulated to experience first signals of mortality prematurely due

to water stress (MEDFATE) and carbon starvation (3D-CMCC-FEM) scenarios. In

conclusion, while management interventions may be considered a viable solution for

the conservation of beech forests under future climate conditions at moister sites like

Cansiglio, in drier sites like Mongiana may not lie in management interventions alone

but rather in the establishment of synergistic mechanisms with other species.

Keywords: Climate change sensitivity, *Fagus sylvatica* L., Forest management
sensitivity, Carbon fluxes, Water fluxes, Stress mitigation, Process-based models.

1. Introduction

Predicting the future evolution of European forests is essential to continue to benefit

from the ecosystem services they provide for human well-being. Forests offer, for

instance, climate change mitigation through their ability to store atmospheric carbon

dioxide in biomass and soil (Augusto and Bo, 2022; Pan et al., 2024). In 2020, the

European Green Deal prioritized the vital role of forests and the forestry sector in

attaining sustainability objectives, such as promoting sustainable forest management,

enhancing forest resilience, and climate change mitigation (European Commission,

2021). Technological advances and studies of forest ecosystem responses to

management practices continue to promote the evolution of strategies that maintain or

enhance forest ecosystem services, such as promoting biological diversity, water

resources, soil protection, or carbon sequestration (Pukkala, 2016). Different forest

management systems have been adopted in Europe over the years (e.g., clear-cutting

or shelterwood) depending, among others, on the wood product desired, the stand

age, and structure (Brunet et al., 2010).

Forest management can be a key element in mitigating the effects of climate warming, maintaining the current primary productivity and the current distribution of tree species, or altering forest composition with more suited and productive species (Bosela et al., 2016). Indeed, the carbon sequestration capacity and productivity of forests are dependent, primarily, on species composition, site conditions as well as on stand age (Rötzer et al., 2010; Vangi et al., 2024a, b), which are affected by past and present forest management activities. According to Collalti et al. (2018) and Dalmonech et al. (2022), monospecific forests in Europe would appear unable to further increase the current rates of carbon storage and biomass production in future climate scenarios, considering current management practices, but at the same time demonstrating that managing under Business as Usual (BAU) practices still allows forests to accumulate biomass at higher rates compared to stands left to develop undisturbed.

European beech (*Fagus sylvatica* L.) is an important deciduous tree species widely distributed in Europe, from southern Scandinavia to Sicily and Spain to northwest Turkey (Durrant et al., 2016). In Italy, according to the National Forest Inventory (INFC, 2015), beech forests cover a total area of 1,053,183 hectares, accounting for about 11.7% of the country's overall forested land. European beech forests demonstrate susceptibility to temperature and precipitation fluctuations. For instance, a warmer environment and less precipitation are forcing shifts in distribution area or the onset of loss of canopy greenness (Axer et al., 2021; Noce et al., 2017, 2023; Zuccarini et al. 2023; Rezaie et al. 2018). According to Skrk et al. (2023), the decline in growth of the beech forests primarily occurs in the dry and warm marginal conditions prevalent near the geographical edge of its distribution with a sub-Mediterranean climatic regime, posing a threat to the survival of beech populations in those areas. However, tree ring analyses have also revealed an unexpected increase in growth in the south Mediterranean region of Albania and Macedonia beech forests at the end of the 20th

century, challenging the presumed suppression of forest ecosystems due to drought (Tegel et al., 2014). Puchi et al. (2024) additionally shed light on the susceptibility to extreme drought events of beech forests found at higher latitudes compared to those found at lower latitudes in the Italian peninsula by highlighting an increase, for the latter, in growth related to the abundance of precipitation. In this context, it is important to minimize the uncertainty surrounding the response of the carbon, water, and energy cycles within beech forest ecosystems, especially as they have been shown to adapt to varying environmental drivers (Deb Burman et al., 2024).

Process-based models (PBMs) are useful tools for studying forest dynamics, such as growth and mortality, as well as water (H_2O) and carbon (C) use efficiency, and carbon stocks as key variables of forest mitigation potential (Vacchiano et al., 2012; Pilli et al., 2022; Testolin et al., 2023; Morichetti et al., 2024). For many years, forest modelling has been widely used by forest ecologists for tackling numerous applied research questions, and the field is continuously evolving to include increasingly complex processes to improve model predictions of forest ecosystem responses to changing climates (Riviere et al., 2020; Kimmins et al., 2008; Maréchaux et al., 2021). By comparing the predictive performance of different models under current environmental conditions, it is possible to gain confidence in their predictions of future trends and make informed decisions in forest ecosystem management and planning processes (Huber et al., 2013; Mahnken et al., 2022).

The main goal of the present study is to investigate the sensitivity of two state-of-the-science PBMs: 3D-CMCC-FEM (Collalti et al., 2014) and MEDFATE (De Cáceres et al., 2023). First under different forest management regimes and climate change scenarios for European beech forests in the Mediterranean area, seeking further insights into the C and H_2O fluxes of this species under different management practices and changing environmental conditions. The study sites vary in terms of environmental factors that

can affect gross primary productivity (GPP), as well as latent heat (LE), which are the

two variables considered in this analysis. Specifically, we tested: (i) to what extent

different forest management options can influence C and H_2O fluxes under the

present-day climate; and, (ii) how harsher climate conditions may affect the C and H_2O

fluxes under different management options. To address these questions we

parameterized and evaluated model performance for C and H_2O fluxes at three forest

stands dominated by beech forests: the Sorø (DK-Sor), Hesse (FR-Hes), and Collelongo

(IT-Col) sites, which are included in the PROFOUND Database (PROFOUND DB) (Reyer

et al., 2020a, b). Subsequently, we assessed the C and H_2O fluxes at two target and

independent beech forest sites in Italy (Cansiglio and Mongiana) by simulating their

development under various management options and evaluating their (model)

sensitivity to current and more severe climate conditions.

2. Material and Methods

2.1 3D-CMCC-FEM model

The 3D-CMCC-FEM v.5.6 ('*Three-Dimensional – Coupled Model Carbon Cycle –*

Forest Ecosystem Module') (Collalti et al., 2024 (and references therein); Marconi et al.,

2017; Dalmonech et al., 2022, 2024; Vangi et al., 2024a, 2024b; Morichetti et al., 2024)

is an eco-physiological, biogeochemical and biophysical model. The model simulates C

and H_2O fluxes occurring within forest ecosystems daily, monthly, or annually,

depending on the processes to simulate, with a common spatial scale of one hectare

(Collalti et al., 2016). Photosynthesis is simulated using the biochemical model of

Farquhar–von Caemmerer–Berry (Farquhar et al., 1980), integrating the sunlit and

shaded leaves of the canopy (De Pury and Farquhar, 1997). For the temperature

dependence of the Michaelis-Menten coefficient for Rubisco and the CO_2

compensation point without mitochondrial respiration, the model adopts the

parameterization described in Bernacchi et al. (2001, 2003). The net balance at the

autotrophic level is represented by net primary production in eq 1:

167                          $NPP = GPP - R_a$ (1)

where R_a includes both maintenance respiration (R_m) and growth respiration (R_g).

When R_m exceeds GPP, resulting in a negative NPP, the trees utilize their non-structural

carbon reserves (NSC) (i.e., soluble sugars and starch) to meet the carbon demand

(Collalti et al., 2020a; Merganikova et al., 2019). In deciduous trees, NSC is used to

create new leaves during the bud-burst phase, replenishes during the growing season

under favourable photosynthetic conditions, and finally remobilized in tissues to

prepare for dormancy at the end of the growth phase. The replenishment of non-

structural carbon reserves is prioritized to reach the minimum safety threshold (i.e., 11%

of sapwood dry mass for deciduous trees). Failure to consume almost all reserves may

trigger defoliation mechanisms, or in the case of complete depletion (e.g., during

prolonged stress periods), it may lead to the death of the entire cohort of trees (i.e.,

carbon starvation). The stomatal conductance g_s is calculated using the Jarvis equation

(Jarvis, 1976). The equation includes a species-specific parameter $g_{s\ max}$ (i.e., maximum

stomatal conductance) controlled by some factors such as light, atmospheric CO_2

concentration, air temperature, soil water content, vapour pressure deficit (VPD), and

stand age. According to Waring and Running (2007) and Monteith and Unsworth

(2008), the Penman-Monteith equation is used to calculate the latent heat (LE) fluxes of

evaporation as a function of incoming radiation, VPD, and conductances at a daily

scale, summing up the canopy, soil, and snow (if any) latent heat expressed as Wm^{-2}.

The 3D-CMCC-FEM accounts for forest stand dynamics, including growth, competition

for light, and tree mortality under different climatic conditions, considering both CO_2

fertilization effects and temperature acclimation (Collalti et al., 2018, 2019; Kattge and

Knorr, 2007). Several mortality routines are considered in the model, such as age-

dependent mortality, background mortality (stochastic mortality), self-thinning

mortality, and the aforementioned mortality due to carbon starvation. In addition to

mortality, biomass removal in 3D-CMCC-FEM results from forest management

practices, such as thinning and final harvest (Collalti et al., 2018, Dalmonech et al.,

2022; Testolin et al., 2023). The required model input data include stand age, average

DBH (Diameter at Breast Height), stand density, and tree height (Collalti et al., 2014).

The soil compartment is represented using one single bucket layer, in which the

available soil water (ASW, in mm) is updated every day considering the water inflows

(precipitation and, if provided, irrigation) and outflows (evapotranspiration, i.e., the sum

of evaporation from the soil and transpiration of the canopy). The remaining water

between these two opposite (in sign) fluxes that exceeds the site-specific soil water

holding capacity is considered lost as runoff. For a full 3D-CMCC-FEM description, see:

https://doi.org/10.32018/ForModLab-book-2024.

2.2 MEDFATE model

MEDFATE v.4.2.0 is an R-based modelling framework that allows the simulation of the

function and dynamics of forest ecosystems, with a specific emphasis on drought

impacts under Mediterranean conditions (De Cáceres et al., 2021, 2023). MEDFATE

calculates energy balance, photosynthesis, stomatal regulation, and plant transpiration

of gas exchange separately at sub-daily steps. Like 3D-CMCC-FEM, MEDFATE also

simulates photosynthesis at the leaf level using the biochemical model of Farquhar–von

Caemmerer–Berry (Farquhar et al., 1980) for sunlit and shaded leaves (De Pury and

Farquhar, 1997). MEDFATE can simulate plant hydraulics and stomatal regulation

according to two different approaches: (a) steady-state plant hydraulics and

optimality-based stomatal regulation (Sperry et al., 1998; Sperry et al., 2017); and (b)

transient plant hydraulics including water compartments and empirical stomatal

regulation (Sureau-ECOS; Ruffault et al., 2022). In this work, we took the second

approach, i.e., Sureau-ECOS (Ruffault et al., 2022).

The hydraulic architecture of the Sureau-ECOS module comprises arbitrary soil layers.

The rhizosphere contains coarse and fine root biomass calculated for each soil layer.

The total root xylem conductance is determined by factors such as root length (limited

by soil depth), weight, and distribution across the different layers. In addition, the

resistance to water flow is dependent on two plant compartments (leaf and stem, each

composed of symplasm and apoplasm). Overall, plant conductance is defined by the

sum of resistances across the hydraulic network (i.e., soil, stem, and leaves), taking into

account processes such as plant capacitance effects (i.e., the variation of symplasmic

water reservoirs in the stem and leaves) and cavitation flows (i.e., water released to the

streamflow from cavitated cells to non-cavitated cells during cavitation) (Hölttä et al.,

2009). It also considers cuticular transpiration of the stem and leaf flows. Each element

(roots, stem, leaves) of the hydraulic network has a vulnerability curve $k(\Psi)$, that

declines as water pressure becomes more negative. The xylem vulnerability curve is

modelled using a sigmoid function, defined by the equation:

235                $$k(\Psi) = k_{max}/1 + exp^{(slope/25)\cdot(\Psi - \Psi 50)} \qquad (2)$$

where k_{max} is the maximum hydraulic conductance, $\Psi 50$ is the water potential

corresponding to 50% of conductance, and "slope" is the slope of the curve at that

point.

The stem vulnerability curve can be used to determine the proportion of stem

conductance loss (PLC_{stem}) associated with vessel embolism. This embolism reduces

overall tree transpiration and photosynthesis. Plant hydraulic failure and tree death can

occur if the PLC_{stem} exceeds the 50% threshold.

Gas exchange in the Sureau-ECOS module depends on stomatal conductance (which

depends on light, water availability, and air temperature) and leaf cuticular

conductance, which changes with leaf temperature due to changes in the permeability

of the epidermis. Stomatal regulation, unlike the 3D-CMCC-FEM, follows the Baldocchi

(1994) approach, which allows coupling leaf photosynthesis with water losses. In

addition, a multiplicative factor depending on leaf water potential is used to decrease

stomatal conductance under drought conditions, following a sigmoidal function similar

to stem vulnerability.

Soil water balance is computed daily. MEDFATE can consider an arbitrary number of

soil layers with varying depths in which the water movement within the soil follows a

dual-permeability model (Jarvis et al., 1991, Larsbo et al., 2005). Soil water content

(ΔV_{soil}, in mm) is calculated taking into account variables such as infiltration, capillarity

rise, deep drainage, saturation effect, evaporation from the soil surface, transpiration of

the herbaceous plant, and woody plant water uptake. A full MEDFATE description is

available at: https://emf-creaf.github.io/medfatebook/index.html.

2.3 Evaluation sites

Model evaluation was performed in three PROFOUND European beech sites, i.e., Sorø

(DK-Sor, Denmark), Hesse (FR-Hes, France), and Collelongo (IT-Col, Italy), in which we

retrieved information on soil texture, soil depth, and stand inventory data of forest

structure for model initialization (Reyer et al., 2020a, b). Moreover, these sites are

equipped with the Eddy Covariance towers (EC; Pastorello et al., 2020) for long-term

continuous monitoring of atmospheric carbon, water, and energy fluxes of the forests

(Fig. 1). The DK-Sor site is located in the forest Lille Bogeskov on the island of Zealand in

Denmark. FR-Hes is situated in the northeastern region of France and lies on the plain

at the base of the Vosges Mountains. IT–Col (Selva Piana stand) is a permanent

experimental plot installed in 1991 and situated in a mountainous area of the Abruzzo

region, the centre of Italy.

The pedological characterization of soils exhibits distinct variations across the studied

sites. The soil at the DK–Sor site is predominantly classified as either Alfisols or

Mollisols. The FR–Hes site showcases an intermediary nature, displaying characteristics

akin to both luvisols and stannic luvisols. At the IT–Col site, the prevailing soil type is

identified as Humic alisols, according to the USDA soil classification system. Full details

of these sites are reported in Table 1.

The variables accounted for in the evaluation were obtained from the Fluxdata website

(http://fluxnet.fluxdata.org/data/fluxnet2015–dataset/)   from   the   FLUXNET2015

database (Pastorello et al., 2020). The variables considered are the daily GPP,

estimated from Net Ecosystem Exchange (NEE) measurements and quality checked

using the constant USTAR turbulence correction according to Papale et al. (2006) and

the Latent Heat flux (LE) with energy balance closure correction (i.e., 'LE_CORR')

(Pastorello et al., 2020).

2.4 Study sites

The two target sites considered in this study are Cansiglio and Mongiana Forests (Fig. 1)

(De Cinti et al., 2016). Each site consists of nine long–term monitored plots of differently

managed beech stands, with a spatial extension for each area above 3 hectares, for

about 27 hectares of the experimental area. Three different treatments were applied

(see Fig. S1–S2). For each site, three of the nine plots considered were left unmanaged

(i.e., no cutting and leaving the stands to natural development), defined as 'Control'

plots, three plots were managed following the historical shelterwood system

('Traditional'), and three with innovative cutting ('Innovative'). In Cansiglio, considering

the developmental stage of the stand was an establishment cut to open growing space

in the canopy for the establishment of regeneration. The 'Innovative' cutting consisted

of selecting a non-fixed number of scattered, well-shaped trees (the 'candidate trees')

and a thinning of neighbouring competitors to reduce competition and promote better

growth. In Mongiana, 'Traditional' silvicultural treatment was the first preparatory cut to

increase the vitality and health of the intended residual trees in the stand. The

'Innovative' option was the identification of 45–50 as 'candidate trees' per hectare and

removing only direct competitors.

The Cansiglio site is situated in a mountainous area in the Veneto region, northern Italy.

Mongiana site is located in a mountainous area in the Calabria region of southern Italy.

The latter shows higher mean annual temperature (MAT, C°) and lower mean annual

precipitation (MAP, mm year^{-1}) (i.e., drier conditions) than the Cansiglio site located at

higher latitudes (Table 1). Data on forest structure and soil texture were collected

during the field campaigns conducted in 2011 and 2019 (Cansiglio) and in 2012 and 2019

(Mongiana). At the Cansiglio site, soils are identified as Haplic luvisols, whereas at

Mongiana, the predominant soil classifications consist of Inceptisols and Entisols,

according to the USDA soil classification system. The variables analyzed in these sites,

like in the evaluation sites, were GPP and LE. A summary for these sites is reported in

Table 1.

317
Fig 1. Map of the study sites. Red dots represent sites for validating fluxes, while the blue dot signifies
sites designated for management investigation.

Table 1. Characteristics of the study sites. The age of the stands refers to 2010. The mean annual temperature (MAT) and mean annual precipitation (MAP) for DK–Sor, FR–Hes and IT–Col refer to the period evaluated (i.e., 2006–2010 for the Sorø and Collelongo site and 2014–2018 for the Hesse site) while for Cansiglio and Mongiana from 2010 to 2022. The sum of precipitation in summer refers to June (J), July (J) and August (A) for the same period.

SITE DESCRIPTION					
	Evaluation sites			Managed sites	
Variable	DK–Sor	FR–Hes	IT–Col	Cansiglio	Mongiana
Coordinates (WGS84)	55°49'N, 11°64' E	48°66'N, 7°08'E	41°85'N, 13°59'E	46°02'N, 12°22'E	38°29'N, 16°14'E
Country	Denmark	France	Italy	Italy	Italy
Altitude (m a.s.l.)	40	305	1500	1300	1300
Area (ha)	1	1	1	27	27
MAT (°C)	8.52	10.27	6.95	6.44	11.01
MAP (mm)	818	853	1075	2219	1701
Slope (%)	–	5	35	12	10
Aspect (°)	0	0	252	135	135
Stand age (yr)	90	45	118	120	90
Summer prec (J–J–A)(mm)	292	205	120	493	141

2.5 Meteorological data

For the evaluation sites (i.e., DK–Sor, FR–Hes, IT–Col) observed meteorological data were retrieved from the harmonized PROFOUND database (Reyer et al., 2020a, b) and Fluxnet database (https://data.icos-cp.eu/).

For the Mongiana and Cansiglio sites, meteorological data for 2010–2022 were obtained at daily temporal resolution from the relevant region's Regional Environmental Protection Agencies (ARPAs), which are responsible for monitoring climate variables with weather stations. The choice of thermo-pluviometric weather station was based on the minimum distance from the study area (between 2 km and 20 km away from the study sites, respectively) and on the data availability and integration with other

weather stations in the proximity. The Bagnouls–Gaussen graph (Fig. S3) shows the

mean monthly precipitation (mm) and air temperature (°C) recorded for every station

inside the catchment.

Climate scenarios used as inputs for two models at the Cansiglio and Mongiana sites,

were from the COSMO–CLM simulation at a resolution of approximately 2.2 km over

Italy (Raffa et al., 2023).

The daily variables considered for 3D-CMCC-FEM were mean solar radiation (MJ m^{-2}

day^{-1}), maximum and minimum air temperature (°C), precipitation (mm day^{-1}), and the

mean relative air humidity (%). In contrast, the MEDFATE model uses mean solar

radiation, maximum and minimum air temperature, precipitation, the daily maximum

and minimum relative air humidity, and wind speed (m s^{-1}).

2.6 Modelling set-up

A set of parameters specific for *Fagus sylvatica* L. was provided as input to the model

3D-CMCC-FEM as described in Collalti et al. (2023) while for MEDFATE as in De

Cáceres et al. (2023). To remove any confounding factors related to parameterization,

the parameters related to photosynthesis and stomatal conductance were kept

constant between the two models (see Table 2). The maximum stomatal conductance

(g$_{s\ max}$) was set for 3D-CMCC-FEM according to Pietsch et al. (2005) as in BIOME BGC

model. In 3D-CMCC-FEM, the maximum RuBisCO carboxylation rate (V$_{cmax}$) and the

maximum electron transport rate for the RuBP regeneration (J$_{max}$) at 25°C are corrected

for leaf temperature according to Medlyn et al. (1999) and soil water content as in

Bonan et al. (2011). In MEDFATE, the V$_{cmax}$ and J$_{max}$ at 25°C are modified according to

Leuning (2002) leaf temperature dependence and modelled according to Medlyn et al.

(1999) and Collatz et al. (1991).

We then used the LAI and Available Soil Water (AWS) values obtained from the 3D-

CMCC-FEM outputs as input for running simulations with the MEDFATE model. This

allowed us to isolate and evaluate the specific effects of the processes of interest, such

as the influence of NSC on stomatal conductance and photosynthesis, compared to the

impact of hydraulic constraints on these same physiological processes.

The LAI values were forced annually from the 3D-CMCC-FEM to the MEDFATE model

since it is currently not possible to prescribe LAI values in the 3D-CMCC-FEM model

(i.e., it is calculated prognostically). Precisely, here we used MEDFATE to simulate C

and H_2O fluxes only while considering plant hydraulics (De Cáceres et al., 2021), from

the forest structure predicted by 3D-CMCC-FEM. For MEDFATE water balance, LAI

values determine the competition for light and also drive the competition for soil water,

along with the root distribution across soil layers.

Table 2. Parameters and variables set for both models during the simulations.

PARAMETERS AND VARIABLES		
Name	Value	Unit
$g_{s\ max}$	0.006 Pietsch et al. (2005)	m s^{-1}
J_{max}	-160 De Cáceres et al. (2023)	μmol photons m^{-2} s^{-1}
V_{cmax}	-95 De Cáceres et al. (2023)	μmol CO$_2$ m^{-2} s^{-1}
LAI	from 3D-CMCC-FEM to MEDFATE	m^2 m^{-2}
ASW	from 3D-CMCC-FEM to MEDFATE	mm

393
394

2.7 Model evaluation

Both models were run for five years on the evaluation sites, with the simulation period

determined by the availability of observed data provided, as already mentioned, from

the PROFOUND database, specifically, from 2006 to 2010 at DK-Sor and IT-Col sites

while for FR-Hes starting from 2014 to 2018. The performance metrics of the results of

the evaluation for each site for the GPP and LE variables were the coefficient of

determination (R^2), Root Mean Square Error (RMSE), and Mean Absolute Error (MAE).

2.8 Model application in managed sites

In the managed sites (i.e., Cansiglio and Mongiana), simulations were performed using Historical climate ('Hist') and, to analyse models' sensitivities to climate change, under two Representative Concentration Pathways 4.5 and 8.5 ('Moderate' and 'Hot Climate'), respectively. The 'Hist' climate was used to run simulations at the Cansiglio site from 2011 to 2022 and the Mongiana site from 2012 to 2022. In contrast, simulations using RCP4.5 and RCP8.5 climate ran accounting for the same period, that is, 11 years for the Cansiglio site and ten years for the Mongiana site, but considering the last years of the climate change scenarios (i.e., 2059-2070 and 2060-2070, respectively) to create harsher temperature and precipitation conditions, but with an increased atmospheric CO_2 concentration (in µmol mol^{-1}).

For each of the nine sampled areas, in the Cansiglio and Mongiana sites, we considered a representative area of one hectare for each type of plot: 'Control', 'Traditional' , and 'Innovative'. At the beginning of the simulations, each site thus included a total of 9 plots, each one hectare in size—comprising three 'Control' plots, three 'Traditional' plots, and three 'Innovative' plots. This setup resulted in a total of nine hectares being simulated per site where the model 3D-CMCC-FEM removed a certain percentage of the Basal Area (BA) according to the LIFE-ManFor project (see Table S2). 'Traditional' and 'Innovative' cutting took place for the first time in 2012 (Cansiglio) and 2013 (Mongiana), respectively. Following preliminary results, since the Mongiana site experienced a lighter thinning intensity compared to the Cansiglio site (refer to Table S2), consequently, for the Mongiana site, we considered an alternative management option involving the removal of 40% of the BA. This was done to evaluate whether a more intensive management approach ('SM') could have influenced models' results on

Gross Primary Productivity (GPP) and Latent Heat (LE) fluxes related to the reduction in

competition and enhancing water availability.

3. Results

3.1 Model evaluation

The daily gross primary productivity (GPP) at DK-Sor, FR-Hes, and IT-Col sites was

estimated from EC and simulated by 3D-CMCC-FEM and MEDFATE, are shown in Fig.

2. At DK-Sor site, the 3D-CMCC-FEM simulates a mean daily GPP of 5.14 gC m^{-2} day^{-1},

while MEDFATE 5.13 gC m^{-2} day^{-1}; and EC 5.54 gC m^{-2} day^{-1}; at the FR-Hes site, 3D-

CMCC-FEM mean daily GPP of 6.18 gC m^{-2} day^{-1} compared to MEDFATE 4.82 gC m^{-2}

day^{-1}, and EC 4.99 gC m^{-2} day^{-1}; lastly at the IT-Col site, 3D-CMCC-FEM mean daily

GPP of 4.88 gC m^{-2} day^{-1} compared to MEDFATE 4.19 gC m^{-2} day^{-1}; and EC 4.11 gC m^{-2}

day^{-1}; Additionally, at the DK-Sor site, the 3D-CMCC-FEM mean daily LE to the

atmosphere of 2.83 MJ m^{-2} day^{-1}, while the MEDFATE mean value of 2.22 MJ m^{-2} day^{-1};

and EC 3.19 MJ m^{-2} day^{-1}; at the FR-Hes site, 3D-CMCC-FEM mean daily LE of 4.16 MJ

m^{-2} day^{-1} compared to MEDFATE 3.01 MJ m^{-2} day^{-1}; and EC 4.47 MJ m^{-2} day^{-1}; in the

end at the IT-Col site, 3D-CMCC-FEM mean daily LE of 2.02 MJ m^{-2} day^{-1} compared to

MEDFATE 2.57 MJ m^{-2} day^{-1}; while EC 3.93 MJ m^{-2} day^{-1}. The GPP predicted by 3D-

CMCC-FEM has shown higher values of R^2 (0.92) at DK-Sor and the lowest value at FR-

Hes site (R^2 = 0.76) whilst a value of R^2 = 0.83 at IT-Col site, respectively. For the

MEDFATE model, the GPP predicted highest value of R^2 (0.85) was at DK-Sor the

lowest (R^2 = 0.68) at IT-Col, and at FR-Hes R^2 = 0.76 the same showed for the 3D-

CMCC-FEM model, respectively. Differently, the highest R^2 (0.89) value for 3D-CMCC-

FEM considering LE predicted vs. observed was at FR-Hes site and almost the same

values for DK-Sor and IT-Col sites (R^2 = 0.85 and 0.84, respectively). MEDFATE, for

predicted vs. observed LE variable, has shown the highest R^2 (0.77) at IT–Col site, lower

R^2 (0.69) value at FR–Hes site and the lowest R^2 (0.62) value at DK–Sor site,

respectively. In general, both the Root Mean Square Error (RMSE) and Mean Absolute

Error (MAE) values in all sites were reasonably low, falling within the ranges of 3.31 to

2.02 gC m^{-2} day^{-1} and 2.46 to 1.47 MJ m^{-2} day^{-1}, for both models and for both the

variables. In Fig. 3 and Table 3 the summary of the evaluation metrics performance

results.

Fig 2. Daily mean variations of GPP (gC m⁻² day⁻¹) and LE (MJ m⁻² day⁻¹) estimated from the direct micrometeorological eddy covariance measurements (GPP-Obs and LE-Obs) and models' simulation (GPP-3D-CMCC-FEM, LE-3D-CMCC-FEM and, GPP-MEDFATE, LE-MEDFATE) during the evaluation period at the DK-Sor, IT-Col and FR-Hes at the Beech forest in 2006–2010 and 2014–2018, respectively.

469 Fig 3. Scatter plots and linear regressions of GPP (gC m⁻² day⁻¹) and LE (MJ m⁻² day⁻¹) of the models

470 versus the direct micrometeorological eddy covariance measurements (Obs) at the Sorø (DK-Sor;

471 2006–2010 period), Collelongo (IT-Col; 2006–2010 period) and Hesse (FR-Hes; 2014–2018 period).

492 Table 3. The correlation coefficient (R²), the Root Mean Square Error (RMSE) and the Mean Absolute Error
493 (MAE) for, the GPP (gC m⁻² day⁻¹) and LE (MJ m⁻² day⁻¹) of the daily simulations at DK–Sor, IT–Col, and
494 FR–Hes sites performed from both models 3D–CMCC–FEM and MEDFATE in the beech forest stands.

| | 3D–CMCC–FEM | | | | | | MEDFATE | | | | | |
| | GPP | | | LE | | | GPP | | | LE | | |
	R^2	RMSE	MAE	R^2	RMSE	MAE	R^2	RMSE	MAE	R^2	RMSE	MAE
DK–Sor	0.91	2.17	1.65	0.85	2.02	1.58	0.85	2.52	1.97	0.62	3.09	2.39
FR–Hes	0.76	3.30	2.46	0.89	2.09	1.47	0.76	2.80	2.21	0.69	3.31	2.37
IT–Col	0.83	2.09	1.56	0.84	2.05	1.57	0.68	2.32	1.87	0.77	2.64	1.90

495
496
497 **3.2 Simulation results at Cansiglio**
498
Fig. 4 shows the simulation results using the 3D–CMCC–FEM and MEDFATE models in

the Cansiglio site. For the 3D–CMCC–FEM, the 'Control' plot exhibited the lowest GPP

values under 'Hist' climate conditions, averaging 1681 gC m⁻² year⁻¹. These values

increased slightly to 1982 gC m⁻² year⁻¹ under the RCP4.5 climate and further rose to

2204 gC m⁻² year⁻¹ under the RCP8.5 climate. Similarly, for plots managed with

'Traditional' methods, the trends were consistent with the 'Control' plot, showing

average GPP values of 1603, 1942, and 2141 gC m⁻² year⁻¹ under 'Hist', RCP4.5 and

RCP8.5 climate, respectively. However, 'Innovative' management showed lower GPP

fluxes across all three climate scenarios, with average values of 1534, 1882, and 2075

gC m⁻² year⁻¹ under 'Hist' RCP4.5 and RCP8.5 climate, respectively. The MEDFATE

model showed higher mean absolute GPP increases than the 3D–CMCC–FEM model

under RCP4.5 and RCP8.5 climates, respectively. Under the 'Hist' climate and all treatments, the mean GPP values were about 1638 gC m^{-2} $year^{-1}$, whereas under the RCP4.5 climate, they rose to 2516 gC m^{-2} $year^{-1}$ and 2995 gC m^{-2} $year^{-1}$ under RCP8.5 climate. Analyzing in Fig. 4 the trends of LE for the 3D-CMCC-FEM model, these trends closely follow those of GPP concerning management treatments. The 3D-CMCC-FEM LE values for the 'Control' plots, similar to GPP, were lowest for the 'Hist' climate with an average value over the simulation years of 1200 and 1501 MJ m^{-2} $year^{-1}$ for the RCP4.5 climate, and 1391 MJ m^{-2} $year^{-1}$ for the RCP8.5 climate, respectively. The LE of the 'Traditional' management predicts values of 1129 in the 'Hist' climate, 1440 in the RCP4.5 climate, and 1338 MJ m^{-2} yr^{-1} in the RCP8.5 climate, respectively. For the 'Innovative' management, the mean LE values were 1121 in the 'Hist' climate, 1403 for the RCP4.5 climate, and 1306 MJ m^{-2} yr^{-1} for the RCP8.5 climate, respectively. Similar to the GPP fluxes, the MEDFATE model simulated reductions in LE fluxes among the treatments and higher values across the climates. The mean LE value modelled in the 'Hist' climate, grouped by treatments (because of slight differences among managements), was about 920, 1419 in the RCP4.5 climate, and 1456 MJ m^{-2} yr^{-1} in the RCP8.5 climate, respectively. MEDFATE simulated a stem xylem conductance loss of approximately 40% in the seventh, eighth, and twelfth years of simulation for the RCP8.5 climate scenario in the 'Control' plot. In contrast, this loss was predicted only in the seventh year for the managed plots. Conversely, near-zero or negligible stem embolism were simulated under the 'Hist' and RCP4.5 climate scenarios. The 3D-CMCC-FEM simulated higher values, albeit in a small percentage (i.e., between 8-10%) of NSC, increasing proportionally to the intensity of basal area removed, better observable in the graph at the tree level.

Fig 4. Comparative analysis between models output at the Cansiglio site. The top-left panel displays the
PLC$_{stem}$ as modelled by MEDFATE, while the top-right panel shows the modelled LAI for 3D-CMCC-FEM
(and used by MEDFATE). The middle-up panels (left and right) present annual GPP (gC m^{-2} year^{-1}) as
modelled by the MEDFATE and 3D-CMCC-FEM, respectively. The middle-down panels (left and right)
depict annual LE (MJ m^{-2} yr^{-1}) modelled by the MEDFATE and 3D-CMCC-FEM, respectively. The bottom
panels (left and right) depict the annual minimum of NSC concentration (%) at the stand and tree level,
respectively, as modelled by the 3D-CMCC-FEM. Different plot management strategies are represented
by distinct line styles: solid lines for 'Control' plots ('no management'), dotted lines for 'Innovative' plots,
and dashed lines for 'Traditional' plots (Shelterwood). Climate scenarios are indicated by line colours:
black for 'Hist' climate data (2010-2022), orange and blue for RCP4.5 and RCP8.5 climate (2059-2070),
respectively.

3.3 Simulation results at Mongiana

Simulation results at the (drier) Mongiana site well depicted the differences with the

rainy Cansiglio site (Fig. 5). The 3D-CMCC-FEM model showed no significant

differences in the mean values of GPP among various management interventions under

'Hist' climate conditions, with a mean value of 2151 gC m^{-2} yr^{-1}. Compared to the

Cansiglio site, Mongiana exhibited lower average GPP values. Under RCP4.5 climate

conditions, the GPP for the 'Control' plot was 1864 gC m^{-2} yr^{-1}. In contrast, under the

current climate, the 'Traditional' and 'Innovative' management interventions yielded

higher average GPP values of 2115 gC m^{-2} yr^{-1} and 2086 gC m^{-2} yr^{-1}. The GPP values

under the more intensive management ('SM') and RCP4.5 climate decreased even

further than those of the 'Control' plot, with an average value of 1650 gC m^{-2} yr^{-1}.

Under the RCP8.5, no differences in GPP were observed among management

strategies, with values of about 1525 gC m^{-2} yr^{-1}. Moreover, under the RCP8.5, the

'Control' plot experienced complete mortality after five years of simulations. The

MEDFATE model predicted slightly higher average values of GPP in the 'Control' plots

(2099 gC m^{-2} yr^{-1}) compared to the managed plots (2087 gC m^{-2} yr^{-1}, encompassing

both 'Traditional' and 'Innovative' of two management strategies), with no significant

differences observed among the management strategies and under 'Hist' climate.

Under the RCP4.5 and RCP8.5, the GPP values were 1608 gC m^{-2} yr^{-1} and 1935 gC m^{-2}

yr^{-1}, respectively. The PLC$_{stem}$ graph in Fig. 5 indicated very high xylem embolism levels

(i.e., reaching 100% every year) under RCP4.5 and RCP8.5 already in the first year of

simulations. A pronounced embolism event was observed under the 'Hist' climate in

2017, 2018, 2019, and 2022 in a 30-45% range for the 'Control' plots, while the

managed plots experienced a maximum embolism of approximately 40% in 2017.

Conversely, the 3D-CMCC-FEM model did not report any significant differences

between managed and unmanaged plots for the LE. The average LE value for the 'Hist'

climate was 1796 MJ m^{-2} yr^{-1}, which decreased to 1220 MJ m^{-2} yr^{-1} under the RCP4.5

and 1190 MJ m^{-2} yr^{-1} under the RCP8.5 in managed plots. As previously described, the

'Control' plot under the RCP8.5 experienced mortality in the sixth year of simulation.

Similarly to the previously described GPP fluxes, the MEDFATE model reported a slight

difference in LE fluxes between the 'Control' plot under historical climate conditions

(1623 MJ m^{-2} yr^{-1}) and the managed plots (1603 MJ m^{-2} yr^{-1}). For the RCP4.5 and

RCP8.5, the LE values were 1100 MJ m^{-2} yr^{-1} and 1089 MJ m^{-2} yr^{-1}, respectively. The LE

values of the 'Control' plots are not reported neither for RCP4.5 nor for RCP8.5 because

of the mortality experienced for the simulation years.

Fig 5. Comparative analysis between models output at the Mongiana site. The top-left panel displays the
percent loss of PLC_{stem} as modelled by MEDFATE while the top-right panel shows the modelled LAI for
3D-CMCC-FEM (and used by MEDFATE). The middle-up panels (left and right) present annual GPP (gC
m^{-2} $year^{-1}$) as modelled by the MEDFATE and 3D-CMCC-FEM, respectively. The middle-down panels
(left and right) depict annual LE (MJ m^{-2} yr^{-1}) modelled by the MEDFATE and 3D-CMCC-FEM,
respectively. The bottom panels (left and right) depict the annual minimum of NSC concentration (%) at
the stand and tree level, respectively, as modelled by the 3D-CMCC-FEM. Different plot management
strategies are represented by distinct line styles: solid lines with circles for 'Control' plots ('no
management'), dotted lines with squares for 'Innovative' plots, dashed lines with stars for 'Traditional'
plots (Shelterwood) and dash-dotted lines with triangles for 'SM' management. Climate scenarios are
indicated by line colours: black for 'Hist' climate data (2010-2022), orange and blue for RCP4.5 and
RCP8.5 climate (2060-2070), respectively.

4. Discussion

First, this study evaluated the performance of two different process-based models in

simulating different beech stands across Europe, starting to the north of Europe and

moving towards the south under different environmental conditions. Secondly, the

study focused on models' sensitivity to management and different management

options and considering different climatic conditions in two specific beech forest

stands in the north and south of the Italian peninsula.

4.1 Model evaluation

To assess the models' accuracy in predicting C and H_2O fluxes, we compared the GPP

and the LE data obtained from the EC towers. Both models predicted GPP and LE

accurately and ensured a good range of general applicability of both models (Kramer et

al., 2002, Verbeeck et al., 2008). The 3D-CMCC-FEM model seems to slightly

overestimate GPP daily values along latitudinal gradients starting from the north (DK-

Sor) to the south (IT-Col), as already found in Collalti et al. (2016). MEDFATE, in

contrast, showed a slight overestimation of GPP only at IT-Col site. The LE predicted by

3D-CMCC-FEM is more accurate than MEDFATE prediction for DK-Sor and FR-Hes

sites but not in IT-Col site in which 3D-CMCC-FEM has shown to underestimate

compared to the observed EC values. For MEDFATE the underestimation of LE was

observed in all the evaluation sites.

The spread observed for the GPP and LE fluxes between the two models may be

attributed to the different assumptions that govern stomatal regulation since both

models use the Farquhar-von Caemmerer-Berry biochemical model to calculate

photosynthesis. The over or underestimation of the flows estimated by the models

both for GPP and the LE compared to the data observed from the EC towers can be

attributed either to the presence of the understory (although commonly sporadic in

mature beech stands), which was not considered in the simulations by both models

and to errors on daily measurement by EC technique (Loescher et al., 2006) or

because a not perfect fit in the modeled seasonality (i.e., the begin and the end of the

growing season) (Richardson et al., 2010). However, the leaf phenological pattern of

the European beech in these sites is well represented by the two models in almost all

of the years according to EC data as shown in supplementary materials (see Fig. S4, S5,

S6, S7, S8, S9, and Fig S10). It is important to note that we did not specifically calibrate

the model parameters from the eddy covariance data for each site. Instead, as Dufrêne

et al. (2005) already did, both models were parameterized using existing literature

values and with one set of parameter values for all sites.

4.2 Climate change and forest management at the Cansigio and Mongiana site

The pre-Alpine site of Cansiglio showed slight differences in the fluxes (i.e., GPP and LE)

between three different management practices and the three climates used. Future

climate is expected to be higher temperature if compared to the historical one, with

MAT higher of about 3.93°C under RCP4.5 and 4.95°C under RCP8.5 for the Cansiglio

site and 4.52°C under RCP4.5 and 5.42°C under RCP8.5 at the Mongiana site. Similarly,

MAP is expected to be 510 mm lower under RCP4.5 and 602 mm lower under RCP8.5

at the Cansiglio site, while 902 mm lower under RCP4.5 and 914 mm lower under

RCP8.5 at the Mongiana site.

Regarding management, the response of the 3D-CMCC-FEM to the removal of a

percentage of the basal area from the stand led to a decrease in GPP in the 'Traditional'

cutting and an even greater extent, in the 'Innovative' cutting compared to the 'Control'

(i.e., no management). Similarly to 3D-CMCC-FEM, the MEDFATE model simulates

slight differences in fluxes amount (e.g., lower values for 'Traditional' and 'Innovative'

cutting than 'Control' plots) between the management regimes in the plots. These

results align with those of Guillemot et al. (2014), who observed a slight decrease in

GPP in managed compared to unmanaged temperate beech forests in France under

different thinning regimes. However, differences were observed in both models under

the three different climates used in the simulations. GPP increased from the 'Hist'

climate to RCP4.5 and reached the maximum for RCP8.5, respectively. This suggests a

plastic response (e.g., photosynthesis and stomatal response) of the stands, as

simulated by models, to harsher conditions, indicating, potentially, a high drought

acclimation capacity (Petrik et al., 2022) and increased GPP because of the so-called

'atmospheric CO_2 fertilization' effect as also found by de Wergifosse et al. (2022) and

Reyer et al. (2013), especially in sites with no apparent water limitation both under

current and projected future climate conditions. The anisohydric behavior of *Fagus*

sylvatica L. results in prolonged stomatal opening relative to isohydric species,

although Puchi et al. (2024) recently found large variability in European beech

responses, maintaining prolonged photosynthetic activity, though this response is

modulated by summer precipitation and the availability of soil water storage

(Leuschner et al., 2021; Baudis et al., 2015). However, for high-altitude stands, growth

could be negatively affected under warmer conditions, as suggested by Chmura et al.

(2024). The LE results for the 3D-CMCC-FEM showed lower values over the simulation

period for managed stands than unmanaged ones showing lesser sensitivity to forest management if compared to MEDFATE. However, under the RCP4.5, the LE values were higher compared to both the 'Hist' climate and the RCP8.5 one due to greater annual cumulative precipitation than the RCP8.5 and higher, on average, temperatures than the 'Hist' scenario.

Conversely, the MEDFATE model was shown to be more sensitive to climate, with a clearer distinction between the 'Hist' climate, the RCP4.5 and RCP8.5 climates, with higher and nearly equal values in the harsher conditions (i.e., RCP4.5 and RCP8.5 climates), with slight differences in the management treatments as obtained by 3D-CMCC-FEM. The Non-Structural carbon (NSC) amount showed the highest values in 'Innovative' plots, followed by 'Traditional' plots, and the lowest values in 'Control' plots, suggesting a benefit in carbon stock accumulation with more carbon going for carbon biomass and less for reserve-replenishment for these stands under management interventions. However, NSC levels remain nearly the same for the three climate scenarios throughout all the simulation years. It is important to note that MEDFATE simulated an initial loss of stem conductance under the climate scenarios, indicating the onset of water stress for the stand. Although in RCP4.5 this is negligible, in RCP8.5 PLC_{stem} values reach a maximum xylem cavitation value of about 40% in the eighth year of simulation for managed plots while for 'Control' plots in the eighth, ninth, and twelfth years, highlighting potential benefits of management to reduce drought stress (Giuggiola et al., 2018; Schmied et al., 2023).

The GPP at the southern Apennine site of Mongiana showed a decrease under RCP4.5 and RCP8.5 scenarios when simulated by the 3D-CMCC-FEM model as a result of harsher environmental conditions, as also resulted in the study by Yu et al. (2022), in which the productivity and then the growth of European beech in southern regions are expected to decrease as affected by more severe climate conditions such as decreased

precipitation and increased in air temperature (Tognetti et al., 2019). Indeed, the

increase in air temperature, a reduction in soil water availability, and the rise in vapor

pressure deficit (VPD) lead to earlier stomatal closure, increased mesophyll resistance,

and elevated abscisic acid production (Kane and McAdam, 2023), all of which

contribute to a decrease in the carbon assimilation rate (Priwitzer et al., 2014; Grossiord

et al., 2020). Specifically, GPP is higher under 'Hist' climate conditions, decreases under

the RCP4.5, and ultimately reaches even lower values under the RCP8.5. Under the

RCP8.5 at the fifth year of simulation, the stand in the 'Control' plot is simulated to die

due to carbon starvation. The annual decline in NSC (Fig. 5) due to an imbalance

between carbon uptake (photosynthesis) and the demands for growth and respiration

suggests that the trees are unable to replenish their carbon reserves. The depletion of

NSC reserves may ultimately disrupt processes such as osmoregulation and phenology

(Martínez-Vilalta et al., 2016), potentially leading to stand defoliation and/or mortality.

The management options did not show changes in GPP under the 'Hist' climate.

However, the increase of GPP was observed under the RCP4.5 in the plots where

'Innovative' and 'Traditional' cutting occurred, although no differences were observed

between them. For instance, the same increase in GPP was reported by Fibbi et al.

(2019) for beech forest under climate change scenarios in Italy. The thinning reduces

the LAI and increase the soil water availability, which positively influence stomatal

conductance and carbon assimilation, providing an acclimation mechanism to drought

during periods of water scarcity (Lüttschwager and Jochheim, 2020; Diaconu et al.,

2017).

In contrast, the more intense cutting exhibited even lower GPP values than the 'Control'

plots. This is likely due to the overly intense thinning, which contrasts the microclimate

effects within this forest stand, reducing the potential to offset climate warming at the

local scale (Rita et al., 2021). Heavy thinning, on the other hand, can increase light

penetration, soil evaporation, and wind speed, thereby heightening tree sensitivity to

vapor pressure deficit under dry conditions (Schmied et al., 2023; Simonin et al., 2007).

LE decreased with the decrease in precipitation under the RCP4.5 and RCP8.5 climate

scenarios compared to the 'Hist' climate. There were no significant differences in LE

among the various management regimes. For the MEDFATE model, negligible or no

differences in GPP were observed under all the climates among various management

options. Although the GPP values estimated by the MEDFATE model under the RCP4.5

and RCP8.5 are similar to those obtained from the 3D-CMCC-FEM model, a closer

analysis of the daily outputs (data not shown) reveals that trees photosynthesize until

the end of July, after which they experience significant embolism (i.e., maximum value

of 100%), as indicated by the PLC_{stem} graph, indicating that the decrease in

precipitation led to summer soil moisture depletion and lethal drought stress levels.

Furthermore, the 'Control' plots experienced mortality even before reaching the

summer period. In recent decades, prolonged drought stress in Mediterranean

mountain regions has significantly reduced the productivity of beech forests, resulting

in a decline in Basal Area Increment (BAI) and overall growth (Piovesan et al., 2008). It

is also important to note that under 'Hist' climate conditions, the MEDFATE model

indicated a stem embolization loss ranging from approximately 10% to 45% during the

drought period (i.e., 2018-2020) in Europe (Italiano et al., 2024; Thom et al., 2023,

Lombardi et al., 2023). The embolization was more pronounced and long-lasting in the

'Control' plots than the managed ones. The same trends were obtained for LE.

5. Conclusion

These two process-based models provide robust evidence for their application in

estimating fluxes, consistent with long-term EC tower measurements in European

beech forests. Despite the minimal parametrization effort to align the two models and

the avoidance of site-specific parameters, reliable results can still be obtained, as

confirmed by the outputs from the Sorø, Hesse, and Collelongo sites. Regarding the

sub-Alpine Cansiglio site, although water limitation does not significantly impact fluxes

or the health of the forest under Moderate climate conditions (RCP4.5), a potential

concern is the embolization predicted by the MEDFATE model under the Hot climate

(RCP8.5) at this site, despite similar levels of precipitation. The high susceptibility of

beech forests at the southern Apennine site of Mongiana to more severe (i.e., hotter

and drier) climatic conditions could lead to the collapse of this forest ecosystem, even

with the application of management options to reduce competition. This necessitates

strategic management planning, including the ability to project (e.g., with forest

models) and evaluate future forest conditions for better management schemes (Taylor

et al., 2009). However, the ability of these forests to survive or resist the impacts of

climate change may not depend solely on density reduction interventions. Prioritizing

the exploration of alternative sustainable management strategies to promote carbon

sequestration in both above-ground biomass and soil is crucial for enhancing climate

change mitigation efforts. Additionally, evaluating silvicultural plans such as the

introduction of complementary species can improve the resilience of vulnerable beech

ecosystems. A modeling approach, similar to the one used in this study, offers a

valuable tool for assessing these alternative strategies and refining forestry

management practices. By integrating these approaches, we can strengthen the long-

term sustainability of forests while preserving the ecological balance of vulnerable

regions.

CRediT authorship contribution statement

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

We would like to thank the University of Tuscia, the CNR ISAFOM of Perugia and the company ENI s.p.a. for the realization of this work. Special thanks to the Institute Research Center for Ecological and Forestry Applications (CREAF) of Barcelona that supported the research by the Spanish "Ministerio de Ciencia e Innovación" (MCIN/AEI/10.13039/501100011033) (grant agreement No. PID2021-126679OB-I00). The authors sincerely thank the ARPA Veneto and Calabria for providing the meteorological data. The Fondazione Centro Euro-Mediterraneo sui Cambiamenti

Climatici (CMCC) for providing climate change scenarios. The Cost-Action PROFOUND for providing both the stand ancillary data and the climate used in this work. D.D., E.V. and A.C. has been partially supported by MIUR Project (PRIN 2020) "Unraveling interactions between WATER and carbon cycles during drought and their impact on water resources and forest and grassland ecosySTEMs in the Mediterranean climate (WATERSTEM)" (Project number: 20202WF53Z), "WAFER" at CNR (Consiglio Nazionale delle Ricerche) and by PRIN 2020 (cod. 2020E52THS) – Research Projects of National Relevance funded by the Italian Ministry of University and Research entitled: "Multi-scale observations to predict Forest response to pollution and climate change" (MULTIFOR, project number: 2020E52THS). D.D., E.V., A.C. acknowledge also funding by the project OptForEU Horizon Europe research and innovation programme under grant agreement No. 101060554. D.D. and A.C. also acknowledge the project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 – Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU under award Number: Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP B83C22002930006, Project title "National Biodiversity Future Centre – NBFC". This work used eddy covariance data acquired and shared by the FLUXNET community, including these networks: AmeriFlux, AfriFlux, AsiaFlux, CarboAfrica, CarboEuropeIP, CarboItaly, CarboMont, ChinaFlux, Fluxnet-Canada, GreenGrass, ICOS, KoFlux, LBA, NECC, OzFlux-TERN, TCOS-Siberia, and USCCC. The 3D-CMCC-FEM model code is publicly available and can be found on the GitHub platform at: https://github.com/Forest-Modelling-Lab/3D-CMCC-FEM. The MEDFATE model code is publicly available and can be found on the GitHub platform at: https://github.com/emf-creaf/medfate.

References

Augusto, L., Boča, A. (2022). *Tree functional traits, forest biomass, and tree species diversity interact with site properties to drive forest soil carbon.* Nat. Commun. 13, 1–12. https://doi.org/10.1038/s41467-022-28748-0

Axer, M., Schlicht, R., Kronenberg, R., Wagner, S. (2021). *The Potential for Future Shifts in Tree Species Distribution Provided by Dispersal and Ecological Niches: A Comparison between Beech and Oak in Europe.* Sustainability, 13, 13067. https://doi.org/10.3390/ su132313067

Baldocchi, D. (1994). *An analytical solution for coupled leaf photosynthesis and stomatal conductance models.* Tree Physiology, 14(7-8-9), 1069–1079. https://doi.org/10.1093/treephys/14.7-8-9.1069

Baudis, M., Premper, T., Welk, E., & Bruelheide, H. (2015). *The response of three Fagus sylvatica L. provenances to water availability at different soil depths.* Ecological Research, 30(5), 853–865. https://doi.org/10.1007/s11284-015-1287-x

Bernacchi, C. J., Singsaas, E. L., Pimentel, C., Portis, A. R., & Long, S. P. (2001). *Improved temperature response functions for models of Rubisco-limited photosynthesis.* Plant Cell & Environment, 24(2), 253–259. https://doi.org/10.1111/j.1365-3040.2001.00668.x

Bernacchi, C. J., Calfapietra, C., Davey, P. A., Wittig, V. E., Scarascia-Mugnozza, G. E., Raines, C. A., & Long, S. P. (2003). *Photosynthesis and stomatal conductance responses of poplars to free-air CO2 enrichment (PopFACE) during the first growth cycle and immediately following coppice.* New Phytologist, 159(3), 609–621. https://doi.org/10.1046/j.1469-8137.2003.00850.x

Bosela, M., Štefančík, I., Petráš, R., Vacek, S. (2016). *The effects of climate warming on the growth of European beech forests depend critically on thinning strategy and site productivity.* Agric. For. Meteorol. 222, 21–31. https://doi.org/10.1016/j.agrformet.2016.03.005

Brunet, J.; Fritz, Ö.; Richnau, G. (2010). *Biodiversity in European beech forests—A review with recommendations for sustainable forest management.* Ecol. Bull. 53, 77–94, Available online: http://www.jstor.org/stable/41442021

Bonan, G. B., Lawrence, P. J., Oleson, K. W., Levis, S., Jung, M., Reichstein, M., Lawrence, D. M., & Swenson, S. C. (2011). *Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data.* Journal of Geophysical Research Atmospheres, 116(G2). https://doi.org/10.1029/2010jg001593

Chmura, D. J., Banach, J., Kempf, M., Kowalczyk, J., Mohytych, V., Szeligowski, H., Buraczyk, W., & Kowalkowski, W. (2024). *Growth and productivity of European beech populations show plastic response to climatic transfer at the north-eastern border of the species range.* Forest Ecology and Management, 565, 122043. https://doi.org/10.1016/j.foreco.2024.122043

Collalti, A., Perugini, L., Santini, M., Chiti, T., Nolè, A., Matteucci, G., Valentini, R. (2014). *A process-based model to simulate growth in forests with complex structure: Evaluation and use of 3D-CMCC Forest Ecosystem Model in a deciduous forest in Central Italy.* Ecol. Modell. 272, 362–378. https://doi.org/10.1016/j.ecolmodel.2013.09.016

Collalti, A., Marconi, S., Ibrom, A., Trotta, C., Anav, A., D'andrea, E., Matteucci, G., Montagnani, L., Gielen, B., Mammarella, I., Grünwald, T., Knohl, A., Berninger, F., Zhao, Y., Valentini, R., Santini, M. (2016). *Validation of 3D-CMCC Forest Ecosystem*

Model (v.5.1) against eddy covariance data for 10 European forest sites. Geosci.
Model Dev. 9, 479–504. https://doi.org/10.5194/gmd-9-479-2016
Collalti, A., Biondo, C., Buttafuoco, G., Maesano, M., Caloiero, T., Lucà, F., Pellicone, G.,
Ricca, N., Salvati, R., Veltri, A., Mugnozza, G. S., & Matteucci, G. (2017). *Simulation,*
calibration and validation protocols for the model 3D-CMCC-CNR-FEM: a case
study in the Bonis' watershed (Calabria, Italy). Forest@, 14(4), 247–256.
https://doi.org/10.3832/efor2368-014
Collalti, A., Trotta, C., Keenan, T.F., Ibrom, A., Bond-Lamberty, B., Grote, R., Vicca, S.,
Reyer, C.P.O., Migliavacca, M., Veroustraete, F., Anav, A., Campioli, M.,
Scoccimarro, E., Šigut, L., Grieco, E., Cescatti, A., Matteucci, G. (2018). *Thinning*
Can Reduce Losses in Carbon Use Efficiency and Carbon Stocks in Managed
Forests Under Warmer Climate. J. Adv. Model. Earth Syst. 10, 2427–2452.
https://doi.org/10.1029/2018MS001275
Collalti, A., Thornton, P.E., Cescatti, A., Rita, A., Borghetti, M., Nolè, A., Trotta, C., Ciais,
P., Matteucci, G. (2019). *The sensitivity of the forest carbon budget shifts across*
processes along with stand development and climate change. Ecol. Appl. 29, 1–
18. https://doi.org/10.1002/eap.1837
Collalti, A., Tjoelker, M.G., Hoch, G., Mäkelä, A., Guidolotti, G., Heskel, M., Petit, G., Ryan,
M.G., Battipaglia, G., Matteucci, G., Prentice, I.C. (2020). *Plant respiration:*
Controlled by photosynthesis or biomass? Glob. Chang. Biol. 26, 1739–1753.
https://doi.org/10.1111/gcb.14857
Collalti, A., Dalmonech, D., Marano, G., Vangi, E., Puchi, P., Grieco, E., Orrico, M. (2023).
3D-CMCC-FEM (Coupled Model Carbon Cycle). BioGeoChemical and Biophysical
Forest Ecosystem – User's Guide. CNR Edizioni; ISBN 978-88-8080-573-1
(electronic edition) https://doi.org/10.32018/3D-CMCC-FEM-2022;
Collalti, A. and Dalmonech, D. and Vangi, E. and Marano, G. and Puchi, P.F. and
Morichetti, M. and Saponaro, V. and Orrico, M.R. and Grieco, E. (2024) *Monitoring*
and Predicting Forest Growth and Dynamics. CNR Edizioni, Roma.
Collatz, G.J., Ball, J.T., Grivet, C. & Berry, J.A. (1991). *Physiological and environmental*
regulation of stomatal conductance, photosynthesis and transpiration: A model
that includes a laminar boundary layer. Agricultural and Forest Meteorology, 54,
107–136.
Communication from the Commission to the European Parliament, the Council, the
European Economic and Social Committee and the Committee of the Regions:
Forging a climate-resilient Europe – the new EU Strategy on Adaptation to
Climate Change – COM(2021) 82 final.
Dalmonech, D., Marano, G., Amthor, J.S., Cescatti, A., Lindner, M., Trotta, C., Collalti, A.
(2022). *Feasibility of enhancing carbon sequestration and stock capacity in*
temperate and boreal European forests via changes to management regimes.
Agric. For. Meteorol. 327, 109203. https://doi.org/10.1016/j.agrformet.2022.109203
Dalmonech, D., Vangi, E., Chiesi, M., Chirici, G., Fibbi, L., Giannetti, F., Marano, G.,
Massari, C., Nolè, A., Xiao, J., & Collalti, A. (2024). *Regional estimates of gross*
primary production applying the Process-Based Model 3D-CMCC-FEM vs.
Remote-Sensing multiple datasets. European Journal of Remote Sensing, 57(1).
https://doi.org/10.1080/22797254.2023.2301657
De Cáceres, M., Mencuccini, M., Martin-StPaul, N., Limousin, J. M., Coll, L., Poyatos, R.,
Cabon, A., Granda, V., Forner, A., Valladares, F., & Martínez-Vilalta, J. (2021).
Unravelling the effect of species mixing on water use and drought stress in
Mediterranean forests: A modelling approach. Agricultural and Forest
Meteorology, 296, 108233. https://doi.org/10.1016/j.agrformet.2020.108233
De Cáceres, M., Molowny-Horas, R., Cabon, A., Martínez-Vilalta, J., Mencuccini, M.,
García-Valdés, R., Nadal-Sala, D., Sabaté, S., Martin-StPaul, N., Morin, X.,

D'Adamo, F., Batllori, E., & Améztegui, A. (2023). *MEDFATE 2.9.3: a trait-enabled model to simulate Mediterranean forest function and dynamics at regional scales.* Geoscientific Model Development, 16(11), 3165–3201. https://doi.org/10.5194/gmd-16-3165-2023

De Cinti, B., Bombi, P., Ferretti, F., Cantiani, P., Di Salvatore, U., SimonÄ iÄ, P., Kutnar, L., Äœeater, M., GarfÃ, V., Mason, F., & Matteucci, G. (2016). *From the experience of LIFE+ ManFor C.BD to the Manual of Best Practices in Sustainable Forest Management.* Italian Journal of Agronomy, 11(s1), 1–175. https://doi.org/10.4081/ija.2016.789

De Pury, D.G.G., Farquhar, G.D., 1997. *Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models.* Plant, Cell Environ. 20, 537–557. https://doi.org/10.1111/j.1365-3040.1997.00094.x

De Wergifosse, L., André, F., Goosse, H., Boczon, A., Cecchini, S., Ciceu, A., Collalti, A., Cools, N., D'Andrea, E., De Vos, B., Hamdi, R., Ingerslev, M., Knudsen, M. A., Kowalska, A., Leca, S., Matteucci, G., Nord-Larsen, T., Sanders, T. G., Schmitz, A., Termonia, A., Vanguelova, E., Van Schaeybroeck, B., Verstraeten, A. Vesterdal, L., Jonard, M. (2022). *Simulating tree growth response to climate change in structurally diverse oak and beech forests.* The Science of the Total Environment, 806, 150422. https://doi.org/10.1016/j.scitotenv.2021.150422

Deb Burman, P. K., A. G. , P., Chakraborty, S., Tiwari, Y. K., Sarma, D., & Gogoi, N. (2024). *Simulating the ecosystem-atmosphere carbon, water and energy fluxes at a subtropical Indian forest using an ecosystem model.* Ecological Modelling, 490, 110637. https://doi.org/10.1016/j.ecolmodel.2024.110637

Diaconu, D., Kahle, H. P., & Spiecker, H. (2017). *Thinning increases drought tolerance of European beech: a case study on two forested slopes on opposite sides of a valley.* European Journal of Forest Research, 136(2), 319–328. https://doi.org/10.1007/s10342-017-1033-8

Dufrene, E., Davi, H., François, C., Maire, G. L., Dantec, V. L., & Granier, A. (2005). *Modelling carbon and water cycles in a beech forest.* Ecological Modelling, 185(2–4), 407–436. https://doi.org/10.1016/j.ecolmodel.2005.01.004

Durrant, H. T., de Rigo, D., Caudullo, G. (2016). *Fagus sylvatica and other beeches in Europe: distribution, habitat, usage and threats.* In: San-Miguel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T., Mauri, A. (Eds.), *European Atlas of Forest Tree Species.* Publ. Off. EU, Luxembourg, pp. e012b90+

Farquhar, G.D., Caemmerer, S., Berry, J.A. (1980). *A biochemical model of photosynthetic CO_2 assimilation in leaves of C_3 species.* Planta 149, 78-90–90.

Fibbi, L., Moriondo, M., Chiesi, M., Bindi, M., & Maselli, F. (2019). *Impacts of climate change on the gross primary production of Italian forests.* Annals of Forest Science, 76(2). https://doi.org/10.1007/s13595-019-0843-x

Giuggiola, A., Zweifel, R., Feichtinger, L. M., Vollenweider, P., Bugmann, H., Haeni, M., & Rigling, A. (2018). *Competition for water in a xeric forest ecosystem – Effects of understory removal on soil micro-climate, growth and physiology of dominant Scots pine trees.* Forest Ecology and Management, 409, 241–249. https://doi.org/10.1016/j.foreco.2017.11.002

Grossiord, C., Buckley, T. N., Cernusak, L. A., Novick, K. A., Poulter, B., Siegwolf, R. T. W., Sperry, J. S., & McDowell, N. G. (2020). *Plant responses to rising vapor pressure deficit.* New Phytologist, 226(6), 1550–1566. https://doi.org/10.1111/nph.16485

Guillemot, J., Delpierre, N., Vallet, P., François, C., Martin-StPaul, N. K., Soudani, K., Nicolas, M., Badeau, V., & Dufrêne, E. (2014). *Assessing the effects of management on forest growth across France: insights from a new functional–structural model.* Annals of Botany, 114(4), 779–793. https://doi.org/10.1093/aob/mcu059

Hölttä, T., Cochard, H., Nikinmaa, E., & Mencuccini, M. (2009). *Capacitive effect of cavitation in xylem conduits: results from a dynamic model.* Plant, Cell & Environment/Plant, Cell and Environment, 32(1), 10–21. https://doi.org/10.1111/j.1365-3040.2008.01894.x

Jarvis, P. G. (1976). *The Interpretation of the Variations in Leaf Water Potential and Stomatal Conductance Found in Canopies in the Field.* Philosophical Transactions of the Royal Society B: Biological Sciences, 273(927), 593–610. doi:10.1098/rstb.1976.0035

Jarvis, N.J., Jansson, P-E., Dik, P.E., Messing, I. (1991). *Modelling water and solute transport in macroporous soil.* I. Model description and sensitivity analysis. Journal of Soil Science, 42, 59–70.

Kane, C., & McAdam, S. (2023). *Abscisic acid driven stomatal closure during drought in anisohydric Fagus sylvatica.* Journal of Plant Hydraulics, 9, 002. https://doi.org/10.20870/jph.2023.002

Kattge, J., Knorr, W., (2007). Temperature acclimation in a biochemical model of photosynthesis: A reanalysis of data from 36 species. Plant, Cell Environ. 30, 1176–1190. https://doi.org/10.1111/j.1365-3040.2007.01690.x

Kimmins, (Hamish) J. P., Blanco, J. A., Seely, B., Welham, C., & Scoullar, K. (2008). *Complexity in modelling forest ecosystems: How much is enough?* Forest Ecology and Management, 256(10), 1646–1658. https://doi.org/10.1016/j.foreco.2008.03.011

Kramer, K., Leinonen, I., Bartelink, H., Berbigier, P., Borghetti, M., Bernhofer, C., Cienciala, E., Dolman, A., Froer, O., Gracia, A., Granier, A., Grünwald, T., Hari, P., Jans, W., Kellomäki, S., Loustau, D., Magnani, F., Markkanen, T., Matteucci, G., Mohren, G.M.J., Moors, E., Nissinen, A., Peltola, H. Sabaté, S., Sanchez, A. (2002). *Evaluation of six process-based forest growth models using eddy-covariance measurements of CO2 and H2O fluxes at six forest sites in Europe.* Global Change Biology, 8(3), 213–230. https://doi.org/10.1046/j.1365-2486.2002.00471.x

Larsbo, M., Roulier, S., Stenemo, F., Kasteel, R., Jarvis, N. (2005). *An Improved Dual-Permeability Model of Water Flow and Solute Transport in the Vadose Zone.* Vadose Zone Journal, 4, 398–406.

Leuning, R. (2002). *Temperature dependence of two parameters in a photosynthesis model.* Plant, Cell and Environment, 25, 1205–1210.

Leuschner, C., Schipka, F., & Backes, K. (2021). *Stomatal regulation and water potential variation in European beech: challenging the iso/anisohydry concept.* Tree Physiology, 42(2), 365–378. https://doi.org/10.1093/treephys/tpab104

Loescher, H. W., Law, B. E., Mahrt, L., Hollinger, D. Y., Campbell, J., & Wofsy, S. C. (2006). *Uncertainties in, and interpretation of, carbon flux estimates using the eddy covariance technique.* Journal of Geophysical Research Atmospheres, 111(D21). https://doi.org/10.1029/2005jd006932

Lombardi, D., Micalizzi, K., & Vitale, M. (2023). *Assessing carbon and water fluxes in a mixed Mediterranean protected forest under climate change: An integrated bottom – up and top – down approach.* Ecological Informatics, 78, 102318. https://doi.org/10.1016/j.ecoinf.2023.102318

Lüttschwager, D., & Jochheim, H. (2020). *Drought Primarily Reduces Canopy Transpiration of Exposed Beech Trees and Decreases the Share of Water Uptake from Deeper Soil Layers.* Forests, 11(5), 537. https://doi.org/10.3390/f11050537

Huber, M. O., Eastaugh, C. S., Gschwantner, T., Hasenauer, H., Kindermann, G., Ledermann, T., Lexer, M. J., Rammer, W., Schörghuber, S., & Sterba, H. (2013). *Comparing simulations of three conceptually different forest models with National Forest Inventory data.* Environmental Modelling & Software, *40*, 88–97. https://doi.org/10.1016/j.envsoft.2012.08.003

INFC (2015). Inventario Nazionale delle Foreste e dei Serbatoi Forestali di Carbonio.
Ministero delle Politiche Agricole Alimentari e Forestali, Ispettorato Generale –
Corpo Forestale dello Stato, CRA – Istituto Sperimentale per l'Assestamento
Forestale   e   per   l'Alpicoltura.   [online]   URL:
https://www.inventarioforestale.org/it/statistiche_infc/

Italiano, S. S., Camarero, J. J., Borghetti, M., Colangelo, M., Rita, A., & Ripullone, F.
(2024). *Drought legacies in mixed Mediterranean forests: Analysing the effects of*
structural overshoot, functional traits and site factors. The Science of the Total
Environment, 927, 172166. https://doi.org/10.1016/j.scitotenv.2024.172166

Mahnken, M., Cailleret, M., Collalti, A., Trotta, C., Biondo, C., D'Andrea, E., Dalmonech,
D., Marano, G., Mäkelä, A., Minunno, F., Peltoniemi, M., Trotsiuk, V., Nadal-Sala, D.,
Sabaté, S., Vallet, P., Aussenac, R., Cameron, D. R., Bohn, F. J., Grote, R.,
Augustynczik, A. L. D., Yousefpour, R., Huber, N., Bugmann, H., Merganičová, K.,
Merganic, J., Valent, P., Lasch-Born, P., Hartig, F., Vega del Valle, I. D., Volkholz, J.,
Gutsch, M., Matteucci, G., Krejza, J., Ibrom, A., Meesenburg, H., Rötzer, T., van der
Maaten-Theunissen, M., van der Maaten, E., & Reyer, C. P. O. (2022). *Accuracy,*
realism and general applicability of European forest models. Global Change
Biology, 28(23), 6921–6943. https://doi.org/10.1111/gcb.16384

Maréchaux, I., Langerwisch, F., Huth, A., Bugmann, H., Morin, X., Reyer, C. P., Seidl, R.,
Collalti, A., De Paula, M. D., Fischer, R., Gutsch, M., Lexer, M. J., Lischke, H.,
Rammig, A., Rödig, E., Sakschewski, B., Taubert, F., Thonicke, K., Vacchiano, G., &
Bohn, F. J. (2021). *Tackling unresolved questions in forest ecology: The past and*
future role of simulation models. Ecology and Evolution, 11(9), 3746–3770.
https://doi.org/10.1002/ece3.7391

Martínez-Vilalta, J., Sala, A., Asensio, D., Galiano, L., Hoch, G., Palacio, S., Piper, F. I., &
Lloret, F. (2016). *Dynamics of non‐structural carbohydrates in terrestrial plants: a*
global   synthesis.   Ecological   Monographs,   86(4),   495–516.
https://doi.org/10.1002/ecm.1231

Medlyn, B. E., Badeck, F. W., De Pury, D. G. G., Barton, C. V. M., Broadmeadow, M.,
Ceulemans, R., De Angelis, P., Forstreuter, M., Jach, M. E., Kellomäki, S., Laitat, E.,
Marek, M., Philippot, S., Rey, A., Strassemeyer, J., Laitinen, K., Liozon, R., Portier, B.,
Roberntz, P., Wang, K., Jarvis, P. G. (1999). *Effects of elevated [CO2] on*
photosynthesis in European forest species: a meta‐analysis of model parameters.
Plant Cell & Environment, 22(12), 1475–1495. https://doi.org/10.1046/j.1365–
3040.1999.00523.x

Merganičová, K., Merganič, J., Lehtonen, A., Vacchiano, G., Sever, M. Z. O.,
Augustynczik, A. L. D., Grote, R., Kyselová, I., Mäkelä, A., Yousefpour, R., Krejza, J.,
Collalti, A., & Reyer, C. P. O. (2019). *Forest carbon allocation modelling under*
climate   change.   Tree   Physiology,   39(12),   1937–1960.
https://doi.org/10.1093/treephys/tpz105

Monteith, J. L. and M. H. Unsworth (2008). *Principles of Environmental Physics, 3rd*
Edition. Burlington, MA, Academic Press https://doi.org/10.3832/ifor1802–009

Morichetti, M., Vangi, E., & Collalti, A. (2024). *Predicted Future Changes in the Mean*
Seasonal Carbon Cycle Due to Climate Change. Forests, 15(7), 1124.
https://doi.org/10.3390/f15071124

Noce, S., Collalti, A., & Santini, M. (2017). *Likelihood of changes in forest species*
suitability, distribution, and diversity under future climate: The case of Southern
Europe.   Ecology   and   Evolution,   7(22),   9358–9375.
https://doi.org/10.1002/ece3.3427

Noce, S., Cipriano, C., & Santini, M. (2023). *Altitudinal shifting of major forest tree*
species in Italian mountains under climate change. Frontiers in Forests and Global
Change, 6. https://doi.org/10.3389/ffgc.2023.1250651
Papale, D., Reichstein, M., Aubinet, M., Canfora, E., Bernhofer, C., Kutsch, W. L.,
Longdoz, B., Rambal, S., Valentini, R., Vesala, T., & Yakir, D. (2006). *Towards a*
standardized processing of Net Ecosystem Exchange measured with eddy
covariance technique: algorithms and uncertainty estimation. Biogeosciences,
3(4), 571–583. https://doi.org/10.5194/bg-3-571-2006
Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.W., Poindexter,
C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Reichstein, M.,
Ribeca, A., van Ingen, C., Vuichard, N., Zhang, L., Amiro, B., Ammann, C., Arain,
M.A., Ardö, J., Arkebauer, T., Arndt, S.K., Arriga, N., Aubinet, M., Aurela, M.,
Baldocchi, D., Barr, A., Beamesderfer, E., Belelli Marchesini, L., Bergeron, O.,
Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T.A., Blanken, P.D.,
Bohrer, G., Boike, J., Bolstad, P.V., Bonal, D., Bonnefond, J.-M., Bowling, D.R.,
Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns, S.P., Buysse,
P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T.R., Cleverly, J.,
Collalti, A., Consalvo, C., Cook, B.D., Cook, D., Coursolle, C., Cremonese, E., Curtis,
P.S., D'Andrea, E., da Rocha, H., Dai, X., Davis, K.J., De Cinti, B., de Grandcourt, A.,
De Ligne, A., De Oliveira, R.C., Delpierre, N., Desai, A.R., Di Bella, C.M., di Tommasi,
P., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A.,
Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H.A.M., Eugster, W., Ewenz, C.M.,
Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G.,
Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., Gielen, B., Gioli, B.,
Gitelson, A., Goded, I., Goeckede, M., Goldstein, A.H., Gough, C.M., Goulden, M.L.,
Graf, A., Griebel, A., Gruening, C., Grünwald, T., Hammerle, A., Han, S., Han, X.,
Hansen, B.U., Hanson, C., Hatakka, J., He, Y., Hehn, M., Heinesch, B., Hinko-Najera,
N., Hörtnagl, L., Hutley, L., Ibrom, A., Ikawa, H., Jackowicz-Korczynski, M., Janouš,
D., Jans, W., Jassal, R., Jiang, S., Kato, T., Khomik, M., Klatt, J., Knohl, A., Knox, S.,
Kobayashi, H., Koerber, G., Kolle, O., Kosugi, Y., Kotani, A., Kowalski, A., Kruijt, B.,
Kurbatova, J., Kutsch, W.L., Kwon, H., Launiainen, S., Laurila, T., Law, B., Leuning,
R., Li, Y., Liddell, M., Limousin, J.-M., Lion, M., Liska, A.J., Lohila, A., López-
Ballesteros, A., López-Blanco, E., Loubet, B., Loustau, D., Lucas-Moffat, A., Lüers, J.,
Ma, S., Macfarlane, C., Magliulo, V., Maier, R., Mammarella, I., Manca, G., Marcolla,
B., Margolis, H.A., Marras, S., Massman, W., Mastepanov, M., Matamala, R.,
Matthes, J.H., Mazzenga, F., McCaughey, H., McHugh, I., McMillan, A.M.S., Merbold,
L., Meyer, W., Meyers, T., Miller, S.D., Minerbi, S., Moderow, U., Monson, R.K.,
Montagnani, L., Moore, C.E., Moors, E., Moreaux, V., Moureaux, C., Munger, J.W.,
Nakai, T., Neirynck, J., Nesic, Z., Nicolini, G., Noormets, A., Northwood, M., Nosetto,
M., Nouvellon, Y., Novick, K., Oechel, W., Olesen, J.E., Ourcival, J.-M., Papuga, S.A.,
Parmentier, F.-J., Paul-Limoges, E., Pavelka, M., Peichl, M., Pendall, E., Phillips, R.P.,
Pilegaard, K., Pirk, N., Posse, G., Powell, T., Prasse, H., Prober, S.M., Rambal, S.,
Rannik, Ü., Raz-Yaseef, N., Rebmann, C., Reed, D., Resco de Dios, V., Restrepo-
Coupe, N., Reverter, B.R., Roland, M., Sabbatini, S., Sachs, T., Saleska, S.R.,
Sánchez-Cañete, E.P., Sanchez-Mejia, Z.M., Schmid, H.P., Schmidt, M., Schneider,
K., Schrader, F., Schroder, I., Scott, R.L., Sedlák, P., Serrano-Ortíz, P., Shao, C., Shi,
P., Shironya, I., Siebicke, L., Šigut, L., Silberstein, R., Sirca, C., Spano, D.,
Steinbrecher, R., Stevens, R.M., Sturtevant, C., Suyker, A., Tagesson, T., Takanashi,
S., Tang, Y., Tapper, N., Thom, J., Tomassucci, M., Tuovinen, J.-P., Urbanski, S.,
Valentini, R., van der Molen, M., van Gorsel, E., van Huissteden, K., Varlagin, A.,
Verfaillie, J., Vesala, T., Vincke, C., Vitale, D., Vygodskaya, N., Walker, J.P., Walter-
Shea, E., Wang, H., Weber, R., Westermann, S., Wille, C., Wofsy, S., Wohlfahrt, G.,

Wolf, S., Woodgate, W., Li, Y., Zampedri, R., Zhang, J., Zhou, G., Zona, D., Agarwal, D., Biraud, S., Torn, M., & Papale, D. (2020). *The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data.* Scientific Data, 7(1). https://doi.org/10.1038/s41597-020-0534-3

Pan, Y., Birdsey, R. A., Phillips, O. L., Houghton, R. A., Fang, J., Kauppi, P. E., Keith, H., Kurz, W. A., Ito, A., Lewis, S. L., Nabuurs, G. J., Shvidenko, A., Hashimoto, S., Lerink, B., Schepaschenko, D., Castanho, A., & Murdiyarso, D. (2024). *The enduring world forest carbon sink. Nature*, 631(8021), 563–569. https://doi.org/10.1038/s41586-024-07602-x

Petrik, P., Petek-Petrik, A., Kurjak, D., Mukarram, M., Klein, T., Gömöry, D., Střelcová, K., Frýdl, J., & Konôpková, A. (2022). *Interannual adjustments in stomatal and leaf morphological traits of European beech (Fagus sylvatica L.) demonstrate its climate change acclimation potential.* Plant Biology, 24(7), 1287–1296. https://doi.org/10.1111/plb.13401

Pietsch, S. A., Hasenauer, H., & Thornton, P. E. (2005). *BGC-model parameters for tree species growing in central European forests.* Forest Ecology and Management, 211(3), 264–295. https://doi.org/10.1016/j.foreco.2005.02.046

Pilli, R., Alkama, R., Cescatti, A., Kurz, W. A., & Grassi, G. (2022). *The European forest carbon budget under future climate conditions and current management practices.* Biogeosciences, 19(13), 3263–3284. https://doi.org/10.5194/bg-19-3263-2022

Piovesan, G., Biondi, F., Di Filippo, A., Alessandrini, A., & Maugeri, M. (2008). *Drought-driven growth reduction in old beech (Fagus sylvatica L.) forests of the central Apennines, Italy.* Global Change Biology, 14(6), 1265–1281. https://doi.org/10.1111/j.1365-2486.2008.01570.x

Priwitzer, T., Kurjak, D., Kmeť, J., Sitková, Z., & Leštianska, A. (2014). *Photosynthetic response of European beech to atmospheric and soil drought.* Central European Forestry Journal, 60(1), 32–38. https://doi.org/10.2478/forj-2014-0003

Puchi, P. F., Dalmonech, D., Vangi, E., Battipaglia, G., Tognetti, R., & Collalti, A. (2024). *Contrasting patterns of water use efficiency and annual radial growth among European beech forests along the Italian peninsula.* Scientific Reports, 14(1). https://doi.org/10.1038/s41598-024-57293-7

Pukkala, T. (2016). *Which type of forest management provides most ecosystem services?* Forest Ecosystems/Forest Ecosystems, 3(1). https://doi.org/10.1186/s40663-016-0068-5

Raffa, M., Adinolfi, M., Reder, A. et al. (2023). *Very High Resolution Projections over Italy under different CMIP5 IPCC scenarios.* Sci Data 10, 238 . https://doi.org/10.1038/s41597-023-02144-9

Reyer, C., Lasch-Born, P., Suckow, F., Gutsch, M., Murawski, A., & Pilz, T. (2013). *Projections of regional changes in forest net primary productivity for different tree species in Europe driven by climate change and carbon dioxide.* Annals of Forest Science, 71(2), 211–225. https://doi.org/10.1007/s13595-013-0306-8

Reyer, C., Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-Born, P., Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders, T., Kolari, P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., Matteucci, G., Collalti, A., D'Andrea, E., Foltýnová, L., Krejza, J., Ibrom, A., Pilegaard, K., Loustau, D., Bonnefond, J.-M., Berbigier, P., Picart, D., Lafont, S., Dietze, M., Cameron, D., Vieno, M., Tian, H., Palacios-Orueta, A., Cicuendez, V., Recuero, L., Wiese, K., Büchner, M., Lange, S., Volkholz, J., Kim, H., Weedon, G., Sheffield, J., Vega del Valle, I., Suckow, F., Horemans, J., Martel, S., Bohn, F., Steinkamp, J., Chikalanov, A., Mahnken, M., Gutsch, M., Trotta, C., Babst, F., & Frieler, K. (2020a). *The PROFOUND database for evaluating vegetation models and simulating climate*

impacts on European forests version V. 0.3. GFZ Data Services. https://doi.org/10.5880/PIK.2020.006

Reyer, C. P. O., Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-Born, P., Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders, T. G. M., Kolari, P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., Collalti, A., Trotta, C., Matteucci, G., D'Andrea, E., Foltýnová, L., Krejza, J., Ibrom, A., Pilegaard, K., Loustau, D., Bonnefond, J.-M., Berbigier, P., Picart, D., Lafont, S., Dietze, M., Cameron, D., Vieno, M., Tian, H., Palacios-Orueta, A., Cicuendez, V., Recuero, L., Wiese, K., Büchner, M., Lange, S., Volkholz, J., Kim, H., Horemans, J. A., Bohn, F., Steinkamp, J., Chikalanov, A., Weedon, G. P., Sheffield, J., Babst, F., Vega del Valle, I., Suckow, F., Martel, S., Mahnken, M., Gutsch, M., & Frieler, K. (2020b). *The PROFOUND database for evaluating vegetation models and simulating climate impacts on European forests.* Earth System Science Data, 12(2), 1295–1320. https://doi.org/10.5194/essd-12-1295-2020

Rezaie, N., D'Andrea, E., Bräuning, A., Matteucci, G., Bombi, P., & Lauteri, M. (2018). *Do atmospheric CO2 concentration increase, climate and forest management affect iWUE of common beech? Evidences from carbon isotope analyses in tree rings.* Tree Physiology, 38(8), 1110–1126. https://doi.org/10.1093/treephys/tpy025

Richardson, A.D., Black, T.A., Ciais, P., Delbart, N., Friedl, M.A., Gobron, N., Hollinger, D.Y., Kutsch, W.L., Longdoz, B., Luyssaert, S., Migliavacca, M., Montagnani, L., Munger, J.W., Moors, E., Piao, S., Rebmann, C., Reichstein, M., Saigusa, N., Tomelleri, E., Vargas, R., & Varlagin, A. (2010). *Influence of spring and autumn phenological transitions on forest ecosystem productivity.* Philosophical Transactions – Royal Society. Biological Sciences, 365(1555), 3227–3246. https://doi.org/10.1098/rstb.2010.0102

Rita, A., Bonanomi, G., Allevato, E., Borghetti, M., Cesarano, G., Mogavero, V., Rossi, S., Saulino, L., Zotti, M., & Saracino, A. (2021). *Topography modulates near-ground microclimate in the Mediterranean Fagus sylvatica treeline.* Scientific Reports, 11(1). https://doi.org/10.1038/s41598-021-87661-6

Riviere, M., Caurla, S., & Delacote, P. (2020). *Evolving Integrated Models From Narrower Economic Tools: the Example of Forest Sector Models.* Environmental Modeling & Assessment, 25(4), 453–469. https://doi.org/10.1007/s10666-020-09706-w

Rötzer, T., Dieler, J., Mette, T., Moshammer, R., & Pretzsch, H. (2010). *Productivity and carbon dynamics in managed Central European forests depending on site conditions and thinning regimes.* Forestry: An International Journal of Forest Research, 83(5), 483–496. https://doi.org/10.1093/forestry/cpq031

Ruffault, J., Pimont, F., Cochard, H., Dupuy, J.-L. & Martin-StPaul, N. (2022). *SurEau-Ecos v2.0: A trait-based plant hydraulics model for simulations of plant water status and drought-induced mortality at the ecosystem level.* Geoscientific Model Development, 15, 5593–5626.

Skrk, N., Martinez del Castillo, E., Serrano-Notivoli, R., Luis, M. De, Novak, K., Merela, M., Cufar, K. (2023). *Spatial and temporal variation of Fagus sylvatica growth in marginal areas under progressive climate change.* Dendrochronologia 81. https://doi.org/10.1016/j.dendro.2023.126135

Schmied, G., Pretzsch, H., Ambs, D., Uhl, E., Schmucker, J., Fäth, J., Biber, P., Hoffmann, Y. D., Šeho, M., Mellert, K. H., & Hilmers, T. (2023). *Rapid beech decline under recurrent drought stress: Individual neighborhood structure and soil properties matter.* Forest Ecology and Management, 545, 121305. https://doi.org/10.1016/j.foreco.2023.121305

Simonin, K., Kolb, T., Montes-Helu, M., & Koch, G. (2007). *The influence of thinning on components of stand water balance in a ponderosa pine forest stand during and*

after extreme drought. Agricultural and Forest Meteorology, 143(3–4), 266–276. https://doi.org/10.1016/j.agrformet.2007.01.003

Sperry, J. S., Adler, F. R., Campbell, G. S., & Comstock, J. P. (1998). *Limitation of plant water use by rhizosphere and xylem conductance: results from a model.* Plant, Cell & Environment/Plant, Cell and Environment, 21(4), 347–359. https://doi.org/10.1046/j.1365-3040.1998.00287.x

Sperry, J.S., Venturas, M.D., Anderegg, W.R.L., Mencuccini, M., Mackay, D.S., Wang, Y., et al. (2017). *Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost.* Plant, Cell & Environment, 40, 816–830. https://doi: 10.1111/pce.12852.

Taylor, A. R., Chen, H. Y. H., & VanDamme, L. (2009). *A Review of Forest Succession Models and Their Suitability for Forest Management Planning.* Forest Science, 55(1), 23–36. https://doi.org/10.1093/forestscience/55.1.23

Tegel, W., Seim, A., Hakelberg, D., Hoffmann, S., Panev, M., Westphal, T., Büntgen, U. (2014). *A recent growth increase of European beech (Fagus sylvatica L.) at its Mediterranean distribution limit contradicts drought stress.* Eur. J. For. Res. 133, 61–71. https://doi.org/10.1007/s10342-013-0737-7

Testolin, R., Dalmonech, D., Marano, G., Bagnara, M., D'Andrea, E., Matteucci, G., Noce, S., Collalti, A. (2023). *Simulating diverse forest management options in a changing climate on a Pinus nigra subsp. laricio plantation in Southern Italy.* Sci. Total Environ. 857, 159361. https://doi.org/10.1016/j.scitotenv.2022.159361

Thom, D., Buras, A., Heym, M., Klemmt, H. J., & Wauer, A. (2023). *Varying growth response of Central European tree species to the extraordinary drought period of 2018 – 2020.* Agricultural and Forest Meteorology, 338, 109506. https://doi.org/10.1016/j.agrformet.2023.109506

Tognetti, R., Lasserre, B., Di Febbraro, M., & Marchetti, M. (2019). *Modeling regional drought-stress indices for beech forests in Mediterranean mountains based on tree-ring data.* Agricultural and Forest Meteorology, 265, 110–120. https://doi.org/10.1016/j.agrformet.2018.11.015

Vacchiano, G., Magnani, F., & Collalti, A. (2012). *Modeling Italian forests: state of the art and future challenges. IForest, 5*(1), 113–120. https://doi.org/10.3832/ifor0614-005

Vangi, E., Dalmonech, D., Cioccolo, E., Marano, G., Bianchini, L., Puchi, P. F., Grieco, E., Cescatti, A., Colantoni, A., Chirici, G., & Collalti, A. (2024a). *Stand age diversity and climate change affects forests' resilience and stability, although unevenly.* bioRxiv (Cold Spring Harbor Laboratory). https://doi.org/10.1101/2023.07.12.548709

Vangi, E., Dalmonech, D., Cioccolo, E., Marano, G., Bianchini, L., Puchi, P. F., Grieco, E., Cescatti, A., Colantoni, A., Chirici, G., & Collalti, A. (2024b). *Stand age diversity (and more than climate change) affects forests' resilience and stability, although unevenly.* Journal of Environmental Management, 366, 121822. https://doi.org/10.1016/j.jenvman.2024.121822

Verbeeck, H., Samson, R., Granier, A. A., Montpied, P., & Lemeur, R. (2008). *Multi-year model analysis of GPP in a temperate beech forest in France.* Ecological Modelling, 210(1–2), 85–103. https://doi.org/10.1016/j.ecolmodel.2007.07.010

Waring, R. H. and S. W. Running (2007). *Forest Ecosystems: Analysis at Multiple Scales.* San Francisco, CA, Elsevier Academic Press.

Yu, X., Orth, R., Reichstein, M., Bahn, M., Klosterhalfen, A., Knohl, A., Koebsch, F., Migliavacca, M., Mund, M., Nelson, J. A., Stocker, B. D., Walther, S., & Bastos, A. (2022b). *Contrasting drought legacy effects on gross primary productivity in a mixed versus pure beech forest.* Biogeosciences, 19(17), 4315–4329. https://doi.org/10.5194/bg-19-4315-2022

Zuccarini, P., Delpierre, N., Mariën, B., Peñuelas, J., Heinecke, T., Campioli, M. (2023). *Drivers and dynamics of foliar senescence in temperate deciduous forest trees at*

1285   *their southern limit of distribution in Europe.* Agric. For. Meteorol. 342, 109716.
1286   https://doi.org/10.1016/j.agrformet.2023.109716