Phthalate exposure influences mating behavior and sperm morphology in an aquatic

ecotoxicology model system

Bryan Lamberto Guevara[1,2], Nadia Patel[3]*, Yi Tu[4,5]*, Maurine Neiman[1]

[1]University of Iowa, Department of Biology, Iowa City, IA, USA

[2]Indiana University, Department of Biology, Bloomington, IN, USA

[3]Cedar Falls High School, Cedar Falls, IA, USA

[4]The Bishop's School, La Jolla, CA, USA

[5]Rice University, Houston, TX, USA

*Contributed equally

Corresponding author: brygueva@indiana.edu

Keywords: *Potamopyrgus antipodarum*, water quality, plastics, snail, reproduction,

pollution, endocrine-disrupting chemical, hormones

Abstract

Phthalates are a group of chemicals used to make plastics more durable, found in

applications from cosmetics, lubricating oils, and flooring to soap, shampoo, and

hairspray (CDC, 2021). Phthalates are also now known to be endocrine disruptors with

connections to adverse reproductive outcomes in animals, including humans. Here, we

evaluate the potential effects of a widely used phthalate ester, dimethyl phthalate

(DMP), on male reproduction in a freshwater snail. DMP is found in industrial

applications like solid rocket propellant as well as consumer products such as insect

repellents and plastics. While there is some evidence that DMP negatively affects

reproduction, especially in females, we still know very little about potential DMP effects

on males. We addressed this important knowledge gap by testing the effects of DMP on

Potamopyrgus antipodarum, a prosobranch snail native to New Zealand. These snails

are very sensitive to water conditions and environmental chemicals, including

endocrine-disrupting compounds, and are thus rising in prominence as water-quality

sentinels and ecotoxicology models. We exposed experimental groups of male *P.*

antipodarum to one of three different concentrations of DMP and characterized mating

behavior and sperm morphology as a function of DMP exposure. Differences in these

traits were primarily observed between the males in the control versus the High (10^{-6} M)

DMP concentration group. As DMP exposure levels increased, we found that mating

frequency ultimately decreased by more than 69% and that sperm morphology was

increasingly altered relative to control males. Altogether, study outcomes suggest DMP

exposure in male animals could have negative effects on reproduction, with particular

relevance in aquatic and marine environments that are especially likely to harbor

leached endocrine-disrupting chemicals.

Introduction

Plastic waste is a major global issue, harming everything from marine organisms like

turtles and seabirds to terrestrial organisms such as elephants and cattle. While plastics

are capable of deteriorating, this process takes much longer than for organic matter.

This extended process of decomposition leads to the formation of microplastics that can

make their way into our rivers, streams, and waterways, ultimately endangering humans

as well as aquatic and marine ecosystems (Ragusa et al. 2021). For example, we now

know that microplastics can be found in all stages of human life, from prenatal

development (Ragusa et al. 2021) to late adulthood.

In addition to posing physical harm, many microplastics also harbor endocrine-

disrupting chemicals (EDCs) that are added to commercial plastics as plasticizers

(D'Angelo & Meccariello 2021). There is emerging evidence that EDCs can damage

human reproductive health as they accumulate in the body (Cuenca 2020). EDCs leach

when they dissociate from the rest of the polymer as a consequence of environmental

stressors such as temperature and pH changes (D'Angelo & Meccariello 2021).

Consumer plastics are commonly exposed to UV radiation, humidity, and abrasion,

which can exacerbate EDC leaching in marine environments (Chen et al. 2019).

Once leached, EDCs can enter human systems orally, topically, and via

inhalation, and can then affect the endocrine system by, for example, mimicking natural

hormones or blocking hormone access to intended receptors and signaling mechanisms

(Diamanti-Kandarakis et al. 2009). In humans, EDCs have been linked to abnormalities

in sex organs, certain cancers, metabolic issues, immune function, and decreased

sperm quality and overall fertility in both males and females (Pan et al. 2024).

Considered broadly, the growing environmental prevalence of EDCs has the capacity to

alter sex ratios, impair immune function, disrupt reproductive health, and ultimately

cause population declines (Windsor et al. 2018).

A class of EDCs known as phthalates are of particular concern as they are not

chemically bound to the plastic polymer, increasing the likelihood of leaching when

faced with environmental stress. Phthalates have been used since the 1930s as

plasticizers for PVC plastics, medical instruments, children's toys, and personal care

products. Phthalates are also found in a variety of different commercial products, with

the implication that phthalates are more likely to bioaccumulate and at a greater rate

than other common commercially used EDCs (Heudorf et al. 2007). The negligent

production, use, and disposal practices of plastic-based products are the major source

of phthalates in aquatic systems via runoff and leaching (Metcalfe et al. 2022). While we

still know relatively little about consequences of phthalates for humans, increased

phthalate exposure has been associated with decreased testosterone levels and

increased levels of estradiol in male humans, which could in turn reduce male fertility

(Zhu et al. 2022).

Dimethyl phthalate (DMP) is a phthalate ester (PAE) found in various consumer

plastics. Most notably, DMP is a main component of plastic #1, which is primarily

polyethylene terephthalate (PET). DMP, despite its similarities to other more toxic

PAEs, has not been found to be toxic during development (Field et al. 1993, NTP 1989).

We do know from animals experiencing experimental oral exposure to DMP that this

exposure can translate into relatively minor decreases in kidney function (U.S. EPA

1987, U.S. DHHS 1993). What remains largely uncharacterized is whether

environmental DMP influences reproduction. The few studies of which we are aware

have produced mixed results. For example, while Gray et al. (2000) found that perinatal

exposure to DMP did not affect testis weight, areola size, or frequency of reproductive

malformations in mice, Mei et al. (2019) found that prolonged exposure to DMP for

female mice decreased the secretion of follicle-stimulating hormones and increased the

levels of the estradiol and luteinizing hormones that are critical regulators of the

menstrual cycle in mammals. There is an especially obvious knowledge gap regarding

whether and how DMP alters male reproductive traits, and especially after the postnatal

period.

Using a freshwater snail ecotoxicology model to test DMP effects on sperm

Potamopyrgus antipodarum is a New Zealand freshwater snail that inhabits a wide range of

freshwater and brackish habitats and is commonly located in more disturbed areas (Loo et al.

2007, Ponder 1988, Zaranko et al. 1997). This snail is also a global invader (reviewed in Geist

et al. 2024). The widespread distribution, broad habitat use, availability of asexual lineages

(e.g., Möller et al. 1996), and documented sensitivity to environmentally relevant

concentrations of EDCs (Duft et al. 2003, 2007, Jobling et al. 2003, Stange & Oehlmann 2012,

Lecomte et al. 2013) make *P. antipodarum* a powerful ecotoxicology model and aquatic

sentinel (e.g., Geiß et al. 2017, Subba et al. 2021). Most of the studies demonstrating

sensitivity to EDCs and other toxicants in *P. antipodarum* emphasize their effects on female

reproductive output or fecundity. By contrast, we know little about the potential for EDCs

(including DMP) to affect reproductive function in male *P. antipodarum*.

A previous study in mice found evidence that motility and activity in sperm

experiencing relatively short-term exposure to phthalates was reduced, a consequence

of decreased sperm capacity to generate ATP when exposed to phthalates. This short-

term phthalate exposure also resulted in abnormal capacitation, the penultimate step in

maturation of mammalian spermatozoa (Amjad et al. 2021). A study of marine mussels

exposed to dibutyl phthalate (DBP), di(2-ethylhexyl) phthalate (DEHP), and DMP

reported increased expression of glutathione (GSH) and other antioxidant enzymes that

protect cells against reactive oxygen species, relative to unexposed individuals. GSH

acted as the primary antioxidant in the mussels exposed to DMP. This influx of GSH

ultimately disturbed the metabolomic balance in these mussels, providing a mechanistic

basis for a connection between DMP exposure and sperm function (Gu et al. 2021).

These previous studies set the stage to investigate whether and how EDCs affect male

P. antipodarum, which can in turn provide insights into how these common

environmental pollutants affect male function across taxa.

In accordance with Dalsenter et al. (2006), who found increases in intromission

latency and number of intromissions for ejaculation for male mice with exposure of 500

mg/kg per day of another phthalate, we hypothesize that DMP will depress male *P.*

antipodarum mating behavior. Using a rat model, Parks et al. (2000) and Drake et al.

(2009) saw decreased fetal testosterone synthesis and disruption of testosterone

production in male reproductive development when the rats were exposed to DEHP and

DBP, respectively, leading us to hypothesize that phthalate exposure will alter sperm

production in *P. antipodarum* via its consequences for testosterone signaling. We

evaluated these hypotheses by comparing the mating behavior and sperm morphology

of male *P. antipodarum* that we experimentally exposed to one of three different

concentrations of DMP, predicting that mating frequency and duration will decrease and

sperm morphology will become increasingly deviant relative to controls as DMP

concentration increases.

Methods

Snail selection

Potamopyrgus antipodarum males used in this study had been cultured under

laboratory conditions for several years at the University of Iowa following descent from a

diverse set of sexual snails collected from multiple New Zealand lake populations.

These snail cultures are housed in 20-liter tanks filled with carbon-filtered tap water in a

16^0 C room with a 16:8-hr light:dark cycle and a relative humidity of 64%. Snails were

haphazardly selected from these tanks. Males were then distinguished from females by

visualizing each snail under a NIKONSMZ800 stereo microscope with a magnification

ranging from 10x to 63x and then by turning each individual on its back and waiting for

the snail to right itself, which reveals the penis on the right side of the male body. We

also measured the length of each snail at this time, choosing only snails that were =\>

3.0 mm from apex to aperture, typically used as the lower size threshold for adulthood

in this species (e.g., Larkin et al. 2016).

Experimental Design

Each of the 32 males that we used for this experiment was placed individually into a 16

oz. glass Mason jar filled with 300 mL of carbon-filtered tap water. While we did not

monitor water quality aside from regular quantification of chlorine levels, all water for all

treatments came from the same source at the same time, allowing us to exclude water

quality differences across treatments other than our focal DMP treatments as the source

of any treatment effects. We applied one of three DMP treatments to each of 24 snails

(8 snails per treatment), with the remaining 8 snails serving as controls (no DMP). The

DMP-exposed snails were housed in water containing 10^{-10} M DMP ("low" treatment),

10^{-8} M DMP ("intermediate" treatment), or 10^{-6} M DMP ("high" treatment). The snails in

the control group did not receive any experimental DMP exposure. Because we were

not aware of any other studies that manipulated DMP exposure in snails, we based our

DMP exposure levels on concentrations used for another common household EDC,

bisphenol A, in a study of its effects on embryo production in *P. antipodarum* (Jobling et

al. 2003). We broadened the range of concentrations employed by Jobling et al (2003)

in order to increase our likelihood of detecting treatment effects. The snails were

exposed to these DMP concentrations for a five-week period prior to mating trials and

dissection for sperm characterization. We used this timeline because spermatogenesis

generally takes 2 – 5 weeks under optimal conditions in small gastropods (de Freitas

Tallarico et al. 2004, Rodrigues et al. 2021). Following standard *P. antipodarum* culture

conditions (e.g., Stork et al. 2022), each jar of snails was supplied 260 µL of a solution

containing 50 mL carbon-filtered tap water, 0.096 g of freshwater fish fry food, and 0.01

g of chalk 3x/week.

Mating Trials

Following the procedures for mating trials used in a previous study of *P. antipodarum*

mating behavior (Stork et al. 2022), an individual trial consisted of one male and two

female *P. antipodarum* placed into a 100-mm petri dish. Male snails were marked with

nail polish on their shells to readily differentiate them from females. The shallow

structure and relatively small size of the petri dish has been found to be an effective

means of ensuring that snails come in contact during the two hours of the trial along

with facilitating clear video analysis of behavior (Stork et al. 2022). We used a two-hour

time period for the trial following Neiman & Lively (2005), who showed that *P.*

antipodarum copulation is almost always concluded within minutes to an hour. At the

start of the two-hour period, the two females were placed equidistant from the male.

Next, all three snails were allowed to freely move around the dish for two hours. We

filmed the petri dish from above using a 1080p Logitech Webcam and the time-lapse

software VideoVelocity (v3.7.2104; CandyLabs 2022). Following Nelson & Neiman

(2011) and Stork et al. (2022), we defined mating attempts as a physical interaction

involving one snail mounting another for more than 15 seconds (Nelson & Neiman

2011, Stork et al. 2022). We used these recordings to quantify the frequency of mating

attempts per male and the duration of each attempt.

Dissections & Microscopic Visualization

We first dissected each male by using fine-tipped forceps to break the shell at its widest

point, enabling careful extraction of the whole body and leaving the sperm duct intact.

We then identified and removed the sperm duct, wet mounted the sperm duct on a

microscope slide, and sealed the slide with acrylic paint to retain moisture. We

visualized sperm under a Leica DMi8 automated microscope using brightfield

technology. Following a similar study of *P. antipodarum* sperm by Jalinsky et al. (2020),

we took images of ~10 sperm per snail at 1000x and used these images and ImageJ

software (Schneider et al. 2012) to measure head length, width, circumference, and

sperm tail length (Figure 1).

Figure 1. Sperm structure and microscopic imaging and measurement in *P.*

antipodarum. Black bar scaled to 30 μm. Yellow segmented line represents the

measuring tool used for measuring sperm parameters. Top left panel depicts isolated

sperm at 10000x with red circle surrounding sperm selected for measurement. Top

right, bottom left, and bottom right panel depict how sperm head circumference, width,

and length, respectively, were measured in ImageJ.

Statistical Analyses

We used Shapiro-Wilks tests to assess normality for each dataset (mating frequency,

mating duration, sperm head length, sperm head width, sperm head circumference,

sperm head length, sperm tail length), and found that each of these datasets

represented significant departures from normality ($p < 1.01e^{-05}$). Accordingly, we used

nonparametric approaches for all subsequent analyses: Kruskal-Wallis tests for the

comparisons of mating attempt frequency, mating attempt duration, and sperm head

length, width, and circumference, and tail length across the four DMP treatment groups.

We also compared each of the possible pairwise sets from the Kruskal-Wallis test using

the post-hoc Dunn test. All data analyses were performed with R (R Core Team, 2024)

via janitor (v.2.2.0; Firke 2023), dplyr (v1.1.4; Wickham et al. 2023), and FSA (v0.9.5;

Ogle et al. 2023). Data were visualized with ggplot2 (Wickham 2016) and ggstatsplot

(Patil 2021).

Results

The mating behavioral trials revealed that mating frequency differed across the

treatment groups ($X^2 = 15.4$, df = 3, $p = 0.001$), with mating frequency in the control

group significantly higher than both the intermediate and high DMP treatment groups

(Figure 2, Supplementary Figure 1, Supplementary Table 1). We did not observe any

significant differences between treatment groups for mating duration (X^2 = 5.62, df = 3,

p = 0.132; Supplementary Table 1).

There were no significant differences across the four DMP treatment groups for

sperm head circumference and head length (Table 1, Figure 3, Supplementary Figure 2,

Supplementary Table 2). The treatment groups did differ in sperm head width and tail

length (Supplementary Table 2). Post-hoc pairwise analysis revealed that this outcome

was driven by the significantly shorter sperm head width in the high DMP group vs.

each of the other 3 DMP treatments. There were no significant differences between the

intermediate and low, intermediate and control, and control and low DMP treatment

groups for sperm head width. Snails from the control treatment had significantly longer

tails than snails from the intermediate and high DMP treatments. Additionally, snails

from the low DMP treatment had longer tails than snails exposed to high DMP. These

results are summarized in Supplementary Table 3.

Table 1. Mean and standard deviation (SD) of sperm metrics from 32 *P. antipodarum*

males across the four treatment groups in micrometers (μm).

DMP Treatment	Head Circumference (SD)	Head Length (SD)	Head Width (SD)	Tail Length (SD)
Control	9.91 (0.79)	4.16 (0.40)	1.11 (0.14)	114.31 (16.45)
Low	9.98 (0.59)	4.26 (0.45)	1.07 (0.09)	113.46 (16.49)
Intermediate	9.88 (0.66)	4.25 (0.32)	1.08 (0.12)	106.75 (14.30)
High	10 (0.83)	4.20 (0.35)	1.06 (0.19)	107.01 (17.19)

Figure 2. Increasing DMP concentration is associated with decreased mating frequency and duration in male *P. antipodarum*. Comparisons of mean (A) frequency of male mating across the 4 DMP treatment groups and (B) time males spent mating. Error bars represent standard error. Statistically significant different pairwise comparisons are denoted by different versus shared letters above error bars.

Figure 3. Sperm morphology as a function of DMP concentration. Comparisons of

mean sperm (A) head circumference, (B) head length, (C) head width, and (D) tail

length across varying levels of DMP. Error bars represent standard error. Statistically

significant different pairwise comparisons are denoted by different versus shared letters

above error bars.

Discussion

We used an increasingly prominent ecotoxicology model system, *Potamopyrgus*

antipodarum, to evaluate potential consequences for male reproduction of exposure to a

common household EDC, dimethyl phthalate (DMP). Our study is novel in comparing

mating behavior and sperm morphology across various levels of DMP concentration in a

freshwater mollusk, which allows us to observe the short-term effects that exposure to

EDCs may impose on male reproduction.

Our experiment revealed that increased exposure to DMP resulted in a

significant decrease in male mating frequency and a trend towards a reduction in mating

duration. Increased DMP concentration was also associated with significantly shorter

sperm tails and head widths. While the relationship between sperm morphology and

fertility is a contentious topic, there is evidence from multiple animal systems that do

demonstrate that sperm morphology is associated with reproductive hormone levels

(Zhao et al. 2020) and the extent of fragmentation of sperm nuclear DNA (Gu et al.

2021), which are in turn linked to fertility (Benchaib et al. 2003). To give a specific

example relevant to our own work, Oppliger et al. (2003) and Radwan (1996) showed

that sperm tail length is a critical predictor of motility and ability to successfully fertilize.

Alterations to the sperm head shape or size may be indicative of DNA damage, as

revealed by studies indicating that expression of GSH, an antioxidant that protects cells

against damage from reactive oxygen species, is increased when exposed to phthalates

(Gu et al. 2021). Studies in biological models such as the pig also demonstrate that

sperm morphology affects sperm motility characteristics (Gil et al. 2009).

Our results are consistent with a scenario where DMP exposure could indeed

influence male fertility in freshwater organisms like *P. antipodarum* and perhaps in

other, more distantly related animals like humans. Because, to the best of our

knowledge, no studies have gone beyond showing increased frequency of

morphological abnormalities in sperm produced by males exposed to EDCs (Hauser et

al. 2003, Pérez-Herrera et al. 2008, Dziewirska et al. 2018), whether these

morphological and behavioral changes we observed will influence can affect male

fertility remains to be seen. Fruitful next steps could include assessing the expression

and release of antioxidants in *P. antipodarum* when exposed to phthalates or other

EDCs, which may reveal DNA damage and provide insight regarding the ability of

damaged sperm ability to fertilize. Direct observations of motility as a function of

changes in *P. antipodarum* sperm tail length would provide further insight into potential

links between phthalate exposure and fertility.

These effects of DMP exposure were evident when DMP treatments were

confined to *adult* snails is notable in light of the fact that phthalate exposure early in

development can cause testicular dysgenesis syndrome (TDS) in humans (Hlisníková et

al. 2020). With this point in mind, it would be valuable to experimentally expose juvenile

P. antipodarum males to phthalates to determine whether gonadal structure and

function are affected and whether and how these consequences of exposure affect

sperm morphology and function. To date, and to the best of our knowledge, the

potential for contaminant-induced abnormalities in mature sperm has not been studied

in any marine or aquatic invertebrate species.

It is possible that at least some of the mating and sperm-related consequences of

DMP exposure for adult snails might be a function of stress. Indeed, chronic stress is

known to lower testosterone levels and damage reproductive organs in humans (Xiong

et al. 2022). In humans, lifestyle factors such as obesity, prolonged exposure to mobile

phone radiation, and occupational stress have been shown to significantly damage

sperm DNA and result in immature sperm (Radwan et al. 2016). Mental and physical

stress are also known to enhance the presence of free radicals, ultimately leading to

oxidative stress that can in turn result in reduced sperm count and sperm motility in

humans (Nowicka-Bauer & Nixon 2020). Thermal stress also decreases mating

frequency and inhibits flight in *Drosophila mojavensis,* but not courtship behaviors such

as wing waving (Patton & Krebs 2011). At least some of these phenomena linked to

thermal stress could be linked to the activation of heat shock protein (HSP; and

specifically HSP70). Indeed, high levels of HSP70 have been reported in infertile men

relative to fertile men (Erata et al. 2008, Kumar & Singh 2022), with Erata et al. (2008)

concluding that HSP70 expression increased in these infertile men as a defense

mechanism against sperm apoptosis.

Together, our study suggests that DMP exposure for adult male animals can

manifest in altered mating behavior and sperm morphology. These results are

significant in being documented in an ecotoxicology and aquatic sentinel model system,

suggesting that the endocrine-disrupting chemicals that leach from plastics into aquatic

and marine systems may be disrupting male fertility in the organisms that inhabit these

environments.

Implications for Conservation

We provide new insight into the potential threat that plastic pollution poses for the

reproductive health of aquatic organisms. We revealed that exposure to a common

EDC, generally assumed to be relatively harmless for humans, alters mating behavior

and sperm morphology in freshwater snails. Because *P. antipodarum* serves as an

aquatic sentinel and ecotoxicology model system, our results suggest that plastic

pollution, through the leaching of EDCs into water bodies, could disrupt reproduction of

various aquatic organisms. Over extended periods of time, the leaching of these

chemicals into our water bodies may lead to population decline for organisms living in

exposed habitats, with potential cascading effects on biodiversity and ecosystem

services.

These findings highlight the need for conservation efforts to prioritize plastic

pollution control and monitoring for EDCs to protect the reproductive health of aquatic

organisms and to ensure long term sustainability of the ecosystems they inhabit.

Ethics and Integrity Statements

Data Availability Statement: We will make all datasets publicly available upon

publication in Dryad.

Ethics and permit approval statement: We have the appropriate permit from the Iowa

Department of Natural Resources to house and culture *P. antipodarum* in the Neiman

lab.

Funding statement: We gratefully acknowledge funding from Linda and Rick Maxson,

the Belin-Blank Center, the National Science Foundation UI-LSAMP program, and the

Iowa Sciences Academy.

Conflict of interest disclosure: We have no conflicts of interest to declare.

Permission to reproduce material from other sources: Not applicable.

Acknowledgments

We gratefully acknowledge help and generosity with respect to microscopy from Dr.

Sarit Smolikove and her lab group, funding from Linda and Rick Maxson, the Belin-

Blank Center, the National Science Foundation UI-LSAMP program, and the Iowa

Sciences Academy, feedback and input from the Neiman lab, and snail care from

Morgan Anderson. We are also grateful for constructive feedback from several

anonymous peer reviewers.

References

Amjad, S., Rahman, M. S., Pang, W.-K., Ryu, D.-Y., Adegoke, E. O., Park, Y.-J., &

Pang, M.-G. (2021). Effects of phthalates on the functions and fertility of mouse

spermatozoa. *Toxicology*, *454*, 152746. https://doi.org/10.1016/j.tox.2021.152746

Benchaib, M., Braun, V., Lornage, J., Hajd, S., Salle, B., Lejeune, H., & Guérin, J. F.

(2003). Sperm DNA fragmentation decreases the pregnancy rate in an assisted

reproductive technique. *Human Reproduction*, *18*(5), 1023–1028.

https://doi.org/10.1093/humrep/deg228

CandyLabs. (2022). VideoVelocity Time-Lapse Capture Studio [Computer software].

Vancouver, Canada.

Chen, Q., Allgeier, A., Yin, D., & Hollert, H. (2019). Leaching of endocrine disrupting

chemicals from marine microplastics and mesoplastics under common life stress

conditions. *Environment International*, *130*, 104938.

https://doi.org/10.1016/j.envint.2019.104938

Cuenca, L. L. (2020). *Analysis of the Molecular Mechanisms Underpinning Germline*

Dysfunction Stemming from Exposure to Environmental Chemicals in C.

elegans (Doctoral dissertation, Harvard University).

Dalsenter, P. R., Santana, G. M., Grande, S. W., Andrade, A. J., & Araújo, S. L. (2006).

Phthalate affect the reproductive function and sexual behavior of male Wistar rats.

Human & Experimental Toxicology, *25*(6), 297–303.

https://doi.org/10.1191/0960327105ht624oa

D'Angelo, S., & Meccariello, R. (2021). Microplastics: A threat for male fertility.

International Journal of Environmental Research and Public Health, *18*(5), 2392.

https://doi.org/10.3390/ijerph18052392

De Freitas Tallarico, L., Okazaki, K., Kawano, T., De Bragança Pereira, C. A., &

Nakano, E. (2004). Dominant lethal effect of 60Co gamma radiation in *Biomphalaria*

glabrata (SAY, 1818). *Mutation Research/Genetic Toxicology and Environmental*

Mutagenesis, *561*(1–2), 139–145. https://doi.org/10.1016/j.mrgentox.2004.05.002

Diamanti-Kandarakis, E., Bourguignon, J.-P., Giudice, L. C., Hauser, R., Prins, G. S.,

Soto, A. M., Zoeller, R. T., & Gore, A. C. (2009). Endocrine-Disrupting Chemicals:

An Endocrine Society scientific statement. *Endocrine Reviews*, *30*(4), 293–342.

https://doi.org/10.1210/er.2009-0002

Drake, A. J., Van Den Driesche, S., Scott, H. M., Hutchison, G. R., Seckl, J. R., &

Sharpe, R. M. (2009). Glucocorticoids amplify dibutyl phthalate-induced disruption

of testosterone production and male reproductive development. *Endocrinology*,

150(11), 5055–5064. https://doi.org/10.1210/en.2009-0700

Duft, M., Schmitt, C., Bachmann, J., Brandelik, C., Schulte-Oehlmann, U., & Oehlmann,

J. (2007). Prosobranch snails as test organisms for the assessment of endocrine

active chemicals—an overview and a guideline proposal for a reproduction test

with the freshwater mudsnail *Potamopyrgus antipodarum*. *Ecotoxicology*, *16*(1),

169–182. https://doi.org/10.1007/s10646-006-0106-0

Duft, M., Schulte-Oehlmann, U., Weltje, L., Tillmann, M., & Oehlmann, J. (2003).

Stimulated embryo production as a parameter of estrogenic exposure via

sediments in the freshwater mudsnail *Potamopyrgus antipodarum*. *Aquatic*

Toxicology, *64*(4), 437–449. https://doi.org/10.1016/S0166-445X(03)00102-4

Dziewirska, E., Radwan, M., Wielgomas, B., Klimowska, A., Radwan, P., Kałużny, P., ...

& Jurewicz, J. (2019). Human semen quality, sperm DNA damage, and the level of

urinary concentrations of 1N and TCPY, the biomarkers of nonpersistent

insecticides. *American journal of men's health*, *13*(1), 1557988318816598.

Erata, G. Ö., Koçak Toker, N., Durlanık, Ö., Kadıoğlu, A., Aktan, G., & Aykaç Toker, G.

(2008). The role of heat shock protein 70 (Hsp 70) in male infertility: Is it a line of

defense against sperm DNA fragmentation? *Fertility and Sterility*, *90*(2), 322–327.

https://doi.org/10.1016/j.fertnstert.2007.06.021

Field, E. A., Price, C. J., Sleet, R. B., George, J. D., Marr, M. C., Myers, C. B., Schwetz,

B. A., & Morrissey, R. E. (1993). Developmental toxicity evaluation of diethyl and

dimethyl phthalate in rats. *Teratology*, *48*(1), 33–44.

https://doi.org/10.1002/tera.1420480107

Firke S (2023). _janitor: Simple Tools for Examining and Cleaning Dirty Data_. R

package version 2.2.0, <https://CRAN.R-project.org/package=janitor>.

Gil, M. C., García-Herreros, M., Barón, F. J., Aparicio, I. M., Santos, A. J., & García-

Marín, L. J. (2009). Morphometry of porcine spermatozoa and its functional

significance in relation with the motility parameters in fresh

semen. *Theriogenology*, *71*(2), 254-263.

Geiß, C., Ruppert, K., Askem, C., Barroso, C., Faber, D., Ducrot, V., Holbech, H.,

Hutchinson, T. H., Kajankari, P., Kinnberg, K. L., Lagadic, L., Matthiessen, P.,

Morris, S., Neiman, M., Penttinen, O.-P., Sanchez-Marin, P., Teigeler, M., Weltje,

L., & Oehlmann, J. (2017). Validation of the OECD reproduction test guideline with

the New Zealand mudsnail *Potamopyrgus antipodarum* using trenbolone and

prochloraz. *Ecotoxicology*, *26*(3), 370–382. https://doi.org/10.1007/s10646-017-

1770-y

Gray, L. E., Jr., Ostby, J., Furr, J., Price, M., Veeramachaneni, D. N. R., & Parks, L.

(2000). Perinatal exposure to the phthalates DEHP, BBP, and DINP, but not DEP,

DMP, or DOTP, alters sexual differentiation of the male rat. *Toxicological Sciences*,

58(2), 350–365. https://doi.org/10.1093/toxsci/58.2.350

Gu, Y.-Y., Wei, Q., Wang, L.-Y., Zhang, Z.-M., Zhang, X.-Q., Sun, A.-L., Chen, J., & Shi,

X.-Z. (2021). A comprehensive study of the effects of phthalates on marine

mussels: Bioconcentration, enzymatic activities and metabolomics. *Marine Pollution*

Bulletin, *168*, 112393. https://doi.org/10.1016/j.marpolbul.2021.112393

Hauser, R., Chen, Z., Pothier, L., Ryan, L., & Altshul, L. (2003). The relationship

between human semen parameters and environmental exposure to polychlorinated

biphenyls and p, p'-DDE. *Environmental health perspectives*, *111*(12), 1505-1511.

Hlisníková, H., Petrovičová, I., Kolena, B., Šidlovská, M., & Sirotkin, A. (2020). Effects

and mechanisms of phthalates' action on reproductive processes and reproductive

Health: A literature review. *International Journal of Environmental Research and*

Public Health, *17*(18), 6811. https://doi.org/10.3390/ijerph17186811

Heudorf, U., Mersch-Sundermann, V., & Angerer, J. (2007). Phthalates: Toxicology and

exposure. *International Journal of Hygiene and Environmental Health*, *210*(5), 623–

634. https://doi.org/10.1016/j.ijheh.2007.07.011

Jalinsky, J., Logsdon, J. M., & Neiman, M. (2020). Male phenotypes in a female

framework: Evidence for degeneration in sperm produced by male snails from

asexual lineages. *Journal of Evolutionary Biology*, *33*(8), 1050–1059.

https://doi.org/10.1111/jeb.13632

Jobling, S., Casey, D., Rodgers-Gray, T., Oehlmann, J., Schulte-Oehlmann, U.,

Pawlowski, S., Baunbeck, T., Turner, A. P., & Tyler, C. R. (2003). Comparative

responses of molluscs and fish to environmental estrogens and an estrogenic

effluent. *Aquatic Toxicology*, *65*(2), 205–220. https://doi.org/10.1016/S0166-

445X(03)00134-6

Kumar, N., & Singh, A. K. (2022). Impact of environmental factors on human semen

quality and male fertility: A narrative review. *Environmental Sciences Europe*, *34*(1),

6. https://doi.org/10.1186/s12302-021-00585-w

Larkin, K., Tucci, C., & Neiman, M. (2016). Effects of polyploidy and reproductive mode

on life history trait expression. *Ecology and Evolution*, *6*(3), 765–778.

https://doi.org/10.1002/ece3.1934

Lecomte, J.-B., Benoît, H. P., Ancelet, S., Etienne, M.-P., Bel, L., & Parent, E. (2013).

Compound Poisson-gamma vs. Delta-gamma to handle zero-inflated continuous

data under a variable sampling volume. *Methods in Ecology and Evolution*, *4*(12),

1159–1166. https://doi.org/10.1111/2041-210X.12122

Loo, S. E., Nally, R. M., & Lake, P. S. (2007). Forecasting New Zealand Mudsnail

Invasion Range: Model Comparisons Using Native and Invaded Ranges.

Ecological Applications, *17*(1), 181–189. https://doi.org/10.1890/1051-

0761(2007)017[0181:FNZMIR]2.0.CO;2

Mei, Y., Rongshuang, M., Ruizhi, Z., Hongyuan, H., Qiyue, T., & Shuhua, Z. (2019).

Effects of dimethyl phthalate (DMP) on serum sex hormone levels and apoptosis in

C57 female mice. *International Journal of Endocrinology and Metabolism*, *17*(2),

e82882. https://doi.org/10.5812/ijem.82882

Metcalfe, C. D., Bayen, S., Desrosiers, M., Muñoz, G., Sauvé, S., & Yargeau, V. (2022).

An introduction to the sources, fate, occurrence and effects of endocrine disrupting

chemicals released into the environment. *Environmental research, 207,* 112658.

https://doi.org/10.1016/j.envres.2021.112658

Møller, V., Forbes, V. E., & Depledge, M. H. (1996). Population responses to acute and

chronic cadmium exposure in sexual and asexual estuarine

gastropods. *Ecotoxicology, 5,* 313-326.

Nelson, A. E., & Neiman, M. (2011). Persistent copulation in asexual female

Potamopyrgus antipodarum: Evidence for male control with size-based

preferences. *International Journal of Evolutionary Biology*, *2011*(1), 439046.

https://doi.org/10.4061/2011/439046

Nowicka-Bauer, K., & Nixon, B. (2020). Molecular changes induced by oxidative stress

that impair human sperm motility. *Antioxidants*, *9*(2), Article 2.

https://doi.org/10.3390/antiox9020134

NTP (National Toxicology Program). (1989). Developmental toxicity evaluation of

dimethyl phthalate (CAS No. 131-11-3) administered to CD rats on gestational days

6 through 15. Prepared by Research Triangle Institute, Research Triangle Park, NC

for National Toxicology Program, Research Triangle Park, NC.

Ogle DH, Doll JC, Wheeler AP, Dinno A (2023). _FSA: Simple Fisheries Stock

Assessment Methods_. R package version 0.9.5, <https://CRAN.R-

project.org/package=FSA>.

Pan, J., Liu, P., Yu, X., Zhang, Z., & Liu, J. (2024). The adverse role of endocrine

disrupting chemicals in the reproductive system. *Frontiers in Endocrinology*, *14*,

1324993. https://doi.org/10.3389/fendo.2023.1324993

Parks, L. G., Ostby, J. S., Lambright, C. R., Abbott, B. D., Klinefelter, G. R., Barlow, N.

J., & Gray Jr., L. E. (2000). The plasticizer diethylhexyl phthalate induces

malformations by decreasing fetal testosterone synthesis during sexual

differentiation in the male rat. *Toxicological Sciences*, *58*(2), 339–349.

https://doi.org/10.1093/toxsci/58.2.339

Patil, I. (2021). *Visualizations with statistical details: The 'ggstatsplot' approach. Journal*

of Open Source Software, 6(61), 3167, doi:10.21105/joss.03167

Patton, Z. J., & Krebs, R. A. (2001). The effect of thermal stress on the mating behavior

of three *Drosophila* species. *Physiological and Biochemical Zoology*, *74*(6), 783–

788. https://doi.org/10.1086/323327

Pérez-Herrera, N., Polanco-Minaya, H., Salazar-Arredondo, E., Solís-Heredia, M. J.,

Hernández-Ochoa, I., Rojas-García, E., ... & Quintanilla-Vega, B. (2008).

PON1Q192R genetic polymorphism modifies organophosphorous pesticide effects

on semen quality and DNA integrity in agricultural workers from southern

Mexico. *Toxicology and applied pharmacology, 230*(2), 261-268.

Ponder, W. F. (1988). *POTAMOPYRGUS ANTIPODARUM*—A MOLLUSCAN

COLONISER OF EUROPE AND AUSTRALIA. *Journal of Molluscan Studies*,

54(3), 271–285. https://doi.org/10.1093/mollus/54.3.271

Radwan, J. (1996). Intraspecific variation in sperm competition success in the bulb mite:

A role for sperm size. *Proceedings of the Royal Society of London. Series B:*

Biological Sciences, *263*(1372), 855–859. https://doi.org/10.1098/rspb.1996.0126

Radwan, M., Jurewicz, J., Merecz-Kot, D., Sobala, W., Radwan, P., Bochenek, M., &

Hanke, W. (2016). Sperm DNA damage—The effect of stress and everyday life

factors. *International Journal of Impotence Research*, *28*(4), 148–154.

https://doi.org/10.1038/ijir.2016.15

Ragusa, A., Svelato, A., Santacroce, C., Catalano, P., Notarstefano, V., Carnevali, O.,

... & Giorgini, E. (2021). Plasticenta: First evidence of microplastics in human

placenta. *Environment international*, *146*, 106274.

R Core Team (2024). _R: A Language and Environment for Statistical Computing_. R

Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Rodrigues, C. C., Caixeta, M. B., & Rocha, T. L. (2021). New insights into the

gametogenesis of *Biomphalaria glabrata* (Mollusca, Gastropoda, Pulmonata):

implications for histopathological assessment. *Brazilian Archives of Biology and*

Technology, *64*, e21200574. https://doi.org/10.1590/1678-4324-2021200574

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25

years of image analysis. *Nature Methods*, *9*(7), 671–675. doi:10.1038/nmeth.2089

Stange, D., & Oehlmann, J. (2012). Identification of oestrogen-responsive transcripts in

Potamopyrgus antipodarum. Journal of Molluscan Studies, *78*(4), 337–342.

https://doi.org/10.1093/mollus/eys019

Stork, S., Jalinsky, J., & Neiman, M. (2022). Evidence for stronger discrimination

between conspecific and heterospecific mating partners in sexual *vs.* asexual

female freshwater snails. *PeerJ*, *10*, e14470. https://doi.org/10.7717/peerj.14470

Subba, M., Keough, M. J., Kellar, C., Long, S., Miranda, A., & Pettigrove, V. J. (2021).

Potamopyrgus antipodarum has the potential to detect effects from various land

use activities on a freshwater ecosystem. *Environmental Pollution*, *287*, 117563.

https://doi.org/10.1016/j.envpol.2021.117563

U.S. Environmental Protection Agency. Health and Environmental Effects Profile for

Phthalate Acid Alkyl, Aryl, and Alkyl/Aryl Esters. ECAO-CIN-P188. Environmental

Criteria and Assessment Office, Office of Health and Environmental Assessment,

Office of Research and Development, Cincinnati, OH. 1987.

U.S. Department of Health and Human Services. Hazardous Substances Data Bank

(HSDB, online database). National Toxicology Information Program, National

Library of Medicine, Bethesda, MD. 1993.

Wickham, H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York,

2016.

Wickham H, François R, Henry L, Müller K, Vaughan D (2023). _dplyr: A Grammar of

Data Manipulation_. R package version 1.1.4, <https://CRAN.R-

project.org/package=dplyr>.

Windsor, F. M., Ormerod, S. J., & Tyler, C. R. (2018). Endocrine disruption in aquatic

systems: Up-scaling research to address ecological consequences. *Biological*

Reviews, *93*(1), 626-641.

Xiong, X., Wu, Q., Zhang, L., Gao, S., Li, R., Han, L., Fan, M., Wang, M., Liu, L., Wang,

X., Zhang, C., Xin, Y., Li, Z., Huang, C., & Yang, J. (2022). Chronic stress inhibits

testosterone synthesis in Leydig cells through mitochondrial damage via Atp5a1.

Journal of Cellular and Molecular Medicine, *26*(2), 354–363.

https://doi.org/10.1111/jcmm.17085

Zaranko, D. T., Farara, D. G., & Thompson, F. G. (1997). Another exotic mollusc in the

Laurentian Great Lakes: The New Zealand native Potamopyrgus antipodarum

(Gray 1843) (Gastropoda, Hydrobiidae). *Canadian Journal of Fisheries and*

Aquatic Sciences, *54*(4), 809–814. https://doi.org/10.1139/f96-343

Zhao, W., Jing, J., Shao, Y., Zeng, R., Wang, C., Yao, B., & Hang, D. (2020).

Circulating sex hormone levels in relation to male sperm quality. *BMC Urology*, *20*,

1–7. https://doi.org/10.1186/s12894-020-00674-7

Zhu, Y., Han, X., Wang, X., Ge, T., Liu, H., Fan, L., Li, L., Su, L., & Wang, X. (2022).

Effect of the phthalates exposure on sex steroid hormones in the US population.

Ecotoxicology and Environmental Safety, *231*, 113203.

https://doi.org/10.1016/j.ecoenv.2022.113203