1 **Urinary Sodium Wasting and Disrupted Collecting Duct Function in Mice with**
2 **dRTA-Causing *SLC4A1* Mutations**
3

4 Priyanka Mungara[1,2], Kristina MacNaughton[1,2], AKM Shahid Ullah[,1,2], Grace
5 Essuman[1,2], Forough Chelangarimiyandoab[1,2], Rizwan Mumtaz[4], J. Christopher
6 Hennings[4], Christian A. Hübner[4,5], Dominique Eladari[6,7,8], R. Todd Alexander[1,2,3],
7 Emmanuelle Cordat[1,2*]
8

[1]Department of Physiology, University of Alberta, Edmonton, AB, Canada
[2]Membrane Protein Disease Research Group, University of Alberta, Edmonton, AB,
Canada
[3]Department of Pediatric, University of Alberta, Edmonton, AB, Canada
[4]Institute of Human Genetics, Jena University Hospital, Friedrich Schiller University, Jena,
Germany
[5] Center for rare diseases, Jena University Hospital, Friedrich Schiller University, Jena,
Germany
[6]Service de Médecine de Précision des Maladies Rénales et Métaboliques, CHU Amiens
Picardie ; Faculté de Médecine d'Amiens, Université de Picardie Jules Verne, Amiens,
France
[7]Institut National de la Santé et de la Recherche Médicale U970, Centre de recherche
PARCC (Paris centre de recherche cardiovasculaire), Paris, France
[8]French Clinical Research Infrastructure Network (F-CRIN): INI-CRCT, Vandœuvre-lès-
Nancy, France
24

25 *To whom correspondence should be addressed: Department of Physiology, Membrane
26 Protein Disease Research Group, Room 7-34, Medical Sciences Building, University of
27 Alberta, Edmonton, Alberta, T6G 2H7, Canada, Tel: 780-492-0209, Fax : 780-492-8915,
28 e-mail : cordat@ualberta.ca

29
30
31
32

Abstract

Distal renal tubular acidosis (dRTA) results in metabolic acidosis due to impaired urinary acidification and an unexplained urinary sodium-wasting phenotype. Here, we report the generation and characterization of a novel dRTA mutant mouse line, Ae1 L919X knockin (KI). Homozygous L919X KI mice exhibit typical dRTA features including a reduced ability to acidify urine in response to an acid load. This renal acidification defect was associated with a reduced number of type A intercalated cells. To assess whether these mice exhibit urinary sodium-wasting as seen in some dRTA patients, homozygous KI L919X and the previously described R607H KI mice were fed a salt-depleted acid diet. Here, in line with human patients, both mouse strains exhibited urinary sodium loss. Additionally, we identified increased expression of tight junction proteins claudin-4 and -10b, suggesting a compensatory paracellular pathway in the loop of Henle. Consistent with data from human patients, L919X KI mice displayed a milder phenotype than R607H KI mice. Our findings reveal that both mouse strains are appropriate models for dRTA with a urinary salt-wasting phenotype with a compensatory up-regulation of the paracellular pathway in the ascending limb of the loop of Henle.

Introduction

Renal tubular acidosis is a group of disorders in which dysregulation of acid-base homeostasis occurs due to improper bicarbonate reabsorption or non-volatile acid excretion by the nephron [1–5]. The inherited or acquired distal renal tubular acidosis (dRTA) impairs distal proton secretion and consequently renal acid excretion[2,4,6–9]. Patients with this disease present with alkaline urine, hyperchloremic metabolic acidosis [2,5–9,11], hypokalemia[10], nephrocalcinosis / kidney stones[11], and renal sodium-wasting[12] with urinary-concentrating defects[5]. Notably, in some patients with inherited dRTA, correction of the metabolic acidosis does not prevent the renal sodium wasting and impaired renal salt handling, consistent with an ion tubular nephropathy[13].

dRTA affects the collecting duct (CD), a primary regulator of acid base homeostasis and electrolyte and fluid status, which finetunes the final urine composition[6]. The CD is composed of principal (PCs) and intercalated (ICs) cells. PCs mediate electrogenic salt and water reabsorption via the epithelial sodium channel (ENaC) and aquaporin 2 (AQP2)[14,15]. ICs regulate acid-base homeostasis via type A (A-ICs) and type B (B-IC)[16]. A-ICs secrete protons via an apical vacuolar proton ATPase (V-ATPase) and promote reabsorption of bicarbonate via the basolateral kidney anion exchanger 1 (kAE1). B-ICs facilitate bicarbonate secretion through the apical chloride/bicarbonate exchanger pendrin and reabsorb protons via a basolateral V-ATPase. The B-ICs also contribute to electroneutral salt reabsorption through the coupled activity of pendrin and the sodium-dependent chloride/bicarbonate exchanger (NDCBE)[17–19]. At least another intercalated cell type exists, characterized by apical expression of both pendrin and the V-ATPase, which is called non-A non-B ICs. The role of non-A, non-B cells is unclear.

Inherited dRTA may arise from autosomal dominant or recessive mutations in the *SLC4A1* gene encoding kAE1[20–23]. One such dominant mutation is the AE1 R589H, which is located on the cytosolic side of transmembrane domain 6 in the gate domain of kAE1[24–26]. *In vitro* studies have revealed retention in the endoplasmic reticulum[23,27] and apical mistargeting[28] of the mutant protein, and a 20-50 % reduction in its anion exchange activity compared to wild-type (WT) kAE1[23]. However, the mutant protein is able to hetero-oligomerize with WT kAE1[27,29], thereby resulting in a slightly reduced chloride/bicarbonate exchange at the cell surface. AE1 R901X is another common autosomal dominant mutation, which causes a cytosolic C-terminal tail truncation. In *in vitro* studies, the R901X mutant has normal transport activity, but is retained intracellularly a apically mistargeted[27,29–31], similarly to the R589H mutant. *In vivo* data from transgenic mice[32,33]

and from dRTA patients' kidney biopsies[34,35] indicate that a defective (or a loss of) kAE1
decreases the abundance of ICs. The kAe1 R607H knock in (KI) mouse model
(orthologous to human dRTA R589H mutation) recapitulates classical metabolic
acidosis[32]. In this model, R607H KI/KI mice exhibited proper kAe1 basolateral membrane
targeting but decreased staining intensity for both kAe1 and the B1 subunit of V-ATPase,
and decreased Ae1 mRNA and protein abundance. In addition, the remaining ICs of these
mutant mice were larger in size and accumulated the autophagy marker p62. Together,
these results are consistent with a reduced number of A-ICs in the R607H KI/KI mice[32].
96
To date, the R607H KI mouse is the only dominant mouse model available. In the
current work, we report the generation and characterization of a second dRTA mutant
mouse model, carrying the autosomal dominant L919X KI mutation, orthologous to the
dominant R901X dRTA human mutation. This second mouse model exhibits the same
inability to acidify urine at baseline as the R607H KI mouse line[32], but overall has a milder
phenotype. Additionally, after salt restriction in acid-loaded mice, we found that both
L919X KI/KI and R607H KI/KI mice display a persistent urinary sodium loss as seen in
dRTA patients. The objectives of this work were to 1) characterize the kAe1 L919X KI
mouse line, 2) determine whether the two mouse models accurately reflect the urinary
ion loss seen in dRTA patients, and 3) begin investigating the mechanism causing urinary
sodium loss in these kAe1 mutant transgenic mice.
108
109
110

Concise Methods

Generation of the L919X KI mice. In Germany, animal experiments were approved by the Thüringer Landesamt für Verbraucherschutz (TLV) under the license 02-016/14. In Canada, animal protocols were approved by the University of Alberta's Animal Care and Use Committee (AUP #1277) and were in accordance with the National and institutional Animal care guidelines. L919X KI mice were generated by homologous recombination similar to Ae1 R607H KI mice[33] as described in Supplemental Material.

Metabolic Cage Experimental Design and Urine and Serum Analysis R607H or L919X mice and WT littermates were provided a "24 hour Salt Depletion or a "14 day Salt Depletion with Acid Load" and placed in metabolic cages (Tecniplast) as described in Supplemental Material. Body weight, chow and water consumption, urine (under mineral oil), and feces were collected every 24 hours. Urine and serum analysis was completed using i-STAT Chem8+ cartridge chip (Abbott Laboratories) and ion chromatography (Dionex Aquion Ion Chromatography System, Thermo Fisher Scientific Inc.).

Whole Kidney Tissues Analysis. Immediately after collection, a half kidney was either incubated in RNAlater (ThermoFisher) to use for reverse transcription quantitative PCR or a quarter of kidney was decapsulated and processed for immunoblotting as described in Supplemental Material (**Supplementary Tables 1 & 2**)**.** Alternatively, kidneys retrogradely perfused with PFA (4 %) were flash frozen in liquid nitrogen cooled isopentane, and cryosections processed for Masson-Goldner stain, Von Kossa staining or further processed for immunostaining as detailed in the Supplemental Material.

Detailed methods are listed in the Supplemental Material.

Results

Generation of Ae1 L919X Knock in Mice and Confirmation of Decreased numbers of A-ICs.

We generated mice carrying the orthologous mouse Ae1 L919X mutation by homologous recombination (**Figure 1 A**). Homozygous mice were viable, developed normally and were fertile. Neither gross kidney malformations nor nephrocalcinosis were observed (**Supplementary Figure 1 A & B**), similar to the Ae1 R607H mutant mice[33]. RT-qPCR analysis on whole kidneys from WT and L919X KI/KI mice at steady state confirmed a significant decrease in Ae1 mRNA abundance in the mutant animals (**Figure 1 B**). Immunostaining for kAe1 in the cortex and medulla of WT, heterozygous, and homozygous L919X KI mice (**Figure 1 C & D**) revealed that similar to WT mice, the kAe1 protein predominantly localized to the basolateral membrane, indicating normal membrane targeting of the mutant protein (**Figure 1 C**). This finding is also consistent with previous findings for R607H KI mice[32]. Additionally, staining intensity of kAe1 in L919X KI/KI mice was markedly reduced compared to WT and heterozygous L919X KI mice, both in the cortex and the medulla. Conversely, signals for pendrin did not exhibit genotype-dependent differences (**Figure 1 C & D**). Collectively, these results suggest a diminished abundance of A-ICs in the L919X KI mouse model, consistent with Ae1 R607H KI mice observations.

L919X KI/KI and R607H KI/KI mice excrete an alkaline urine coupled with less urinary ammonium than WT littermates.

To validate the L919X KI/KI mice as a model that recapitulates human dRTA, we assessed plasma and urine parameters at baseline (**Figure 2, Supplementary Table 3).** L919X KI/KI mice had alkaline urine compared to WT littermates (**Figure 2 A).** In agreement, they also excreted significant less urinary ammonium (**Figure 2 B**). Therefore, the Ae1 L919X KI mice display the characteristic inability to properly acidify urine as seen in dRTA human patients[5]. There was no difference in urine osmolality between genotypes (**Figure 2 C**) and the L919X KI/KI mice excreted sodium, chloride, and potassium comparably to WT mice (**Figure 2 D - F).** There were no differences in plasma ion concentrations or pH in the L919X KI/KI compared to WT mice (**Supplementary Table 3).** We completed a baseline analysis of the R607H KI/KI mice as well (**Figure 2 G-L and Supplementary Table 4**). We confirmed that R607H KI/KI mice displayed the same phenotype as L919X KI/KI mice at baseline (**Figure 2 G - L**). These mutant mice had alkaline urine and decreased ammonium excretion (**Figure 2 G,**

H), as previously shown[33]. There was no difference in urine osmolality (**Figure 2 I**) and mutant mice also excreted sodium, chloride, and potassium to a similar extent as WT mice (**Figure 2 J - K**). Overall, this baseline characterization supports an abnormal urinary acidification in both dRTA mouse lines at baseline, consistent with the hallmark characteristics of human dRTA patients[1].

Acute sodium depletion confirms impaired urinary acidification and potassium excretion in dRTA mutant mice compared to WT.

To assess whether dRTA mutant mice show a salt-losing phenotype similar to dRTA patients, we challenged R607H KI/KI, L919X KI/KI mice or WT littermates with a sodium and chloride-depleted diet for 24 hours (**Protocol 1***)* (**Supplementary Figure 2, Supplementary Tables 5 & 6**). Both L919X and R607H KI/KI mice consistently excreted urine with a higher pH compared to WT littermates (**Supplementary Figure 2 A & F**). Both mutant mice strains had no significant differences in urine osmolality between genotypes, although there was trend towards lower osmolality for L919X KI/KI mice (**Supplementary Figure 2 B & G).** Both L919X KI/KI and R607H KI/KI mice reduced urinary sodium and chloride excretion appropriately, to a similar extent as WT littermates (**Supplementary Figure 2 C-D, H-I**). Interestingly, L919X KI/KI mice excreted less urinary potassium while there was only a trend in R607H KI/KI mice (**Supplementary Figure 2 E, J**). No significant plasma difference was found between genotypes for each strain, except that R607H KI/KI mice continued to have increased plasma potassium compared to WT littermates (**Supplementary Tables 5 & 6**). Together these results provide preliminary evidence of renal electrolyte mishandling under acute sodium-depleted conditions for both L919X and R607H KI/KI mice, suggesting a defective coupling of potassium secretion to sodium and chloride reabsorption in the CDs of both strains.

Salt depleted acid loading results in hypernatremia and hyperchloremia, as well as increased renin levels in both dRTA mutant strains.

To further challenge the dRTA mutant mice, we fed WT and mutant mice a sodium-depleted acid diet (Supplementary Material, **Protocol 2***).* This diet was designed to recreate a study by Sebastien et al. revealing an urinary sodium wasting phenotype in dRTA patients on a salt-restricted diet for 8-10 days[12]. Following this challenge, only R607H KI/KI mice displayed metabolic acidosis from the acid load, with decreased plasma bicarbonate and pH (**Supplementary Table 8)**. L919X KI mice only showed a trend for acidosis (p = 0.068) (**Supplementary Table 7)**, suggesting a more severe phenotype in R607H KI/KI than in L919X KI/KI mice. Notably, in both L919X and R607H

KI/KI mutant mice, plasma sodium and chloride concentrations were significantly increased compared to WT littermates and compared to steady state **(Figure 3 A & B, G &H, Supplementary Tables 7 & 8**), suggesting a dehydrated state in KI mice. Given the significant hypernatremia and hyperchloremia in both mutant animals, we wondered whether the renin-angiotensin-aldosterone system (RAAS) or dehydration was activated in mutant mice. Indeed, we observed a higher renin mRNA abundance in homozygous R607H KI mice with a similar trend in L919X KI/KI mice (**Figure 3 C, I**). We next quantified water consumption (**Figure 3 E, K**), urine osmolality (**Figure 3 D, J**), and plasma hemoglobin and hematocrit levels for both strains (**Supplementary Tables 7 & 8**). While no difference in urine osmolality was observed compared to WT littermates (**Figure 3 D, J**), homozygous R607H and L919X mice had reduced water consumption (**Figure 3 E, K**), despite increased hemoglobin concentration in homozygous R607H mice (**Supplementary Table 8**). These results suggest an abnormal salt handling with activation of the RAAS and/or potential dehydration in these mice.

Salt-depleted acid loading reveals a salt wasting nephropathy in homozygous L919X and R607H KI mice.

Similarly to the sodium depletion diet alone, both homozygous L919X and R607H KI mice acidified their urine, although not to the same extent as their WT littermates (**Figure 4 A & 6 A**). R607H KI/KI mice excreted significantly less ammonium compared to WT (**Figure 5 B**), while L919X KI/KI mice (**Figure 4 B**) showed no difference compared to their controls, although both genotypes increased overall ammonium excretion at day 14 compared to day 0 (**Figure 4 B & 5 B**). The salt-depleted acid diet also revealed a pronounced urinary sodium-losing phenotype in homozygous L919X and R607H KI mice (**Figure 4 C&F, Figure 5 C&F**), in agreement with observations from dRTA patients under similar physiological conditions[12]. Distinct phenotypes were observed concerning renal potassium and chloride handling (**Figure 4 & 5 D, E, G, H**). L919X KI/KI mice exhibited urinary chloride loss with no difference in potassium excretion compared to WT mice (**Figure 4 D & E**). In contrast, R607H KI/KI mice displayed lower urinary potassium and chloride excretion (**Figure 5 D & E**). These experiments demonstrate that in mice, both the Ae1 L919X and R607H mutation lead to urinary sodium loss and inability to maximally acidify their urine, thus reproducing major features observed in patients suffering with type 1 dRTA.

Paracellular proteins claudin 4 and claudin 10b are upregulated in dRTA mutant mice.

Given the alkaline urine and the urinary sodium loss after salt depleted acid
loading, we quantified mRNA and protein levels of specific markers of the CD, the thick
ascending limb (TAL) and the distal convoluted tubule (DCT) in KI/KI mice and the
respective WT littermates (**Figure 6, Supplementary Figures 3 - 4)**.

We found increased mRNA and protein levels of the paracellular pores claudin 4
and claudin 10b, in both L919X KI/KI and R607H KI/KI mice (**Figure 6 A-H**). Claudin-4 is
a paracellular chloride pore and a sodium blocker expressed in the thin ascending limb
of Henle's loop, the distal tubule and CD[41–43], while claudin 10b is a sodium pore in the
TAL[44–46]. These findings suggest a common compensatory mechanism via the
paracellular pathway in the loop of Henle of both mutant strains.

The salt-depleted acid challenge also revealed a downregulation of A-IC and B-IC
marker gene expression in R607H KI/KI mice, with similar trends in L919X KI/KI mice
(**Supplementary Figure 3 & 4 A - I)**. There was no clear trend for principal cell markers
between KI/KI mice and WT littermates. In agreement with decreased numbers of ICs
reported for R607H KI/KI mice[32], kAe1 and B1-ATPase mRNA or protein abundances in
whole kidney lysate were significantly reduced compared to WT littermates
(**Supplementary Figure 3 & 4 A, C, K)**. Messengers for pendrin and AE4 were
significantly reduced in R607H KI/KI mice, with a similar trend in L919X KI/KI mice
(**Supplementary Figure 3 & 4 C, E**). Increased NDCBE mRNA abundance was observed
in R607H KI/KI mice with a similar trend in L919X KI/KI mice (**Supplementary Figure 3
& 4 D**).

Given the higher abundance of the TAL-specific claudin-10b in both mutant mouse lines
(**Figure 6 C, D, G, H)**, we investigated gene expression of various transporters from prior
nephron segments (**Supplementary Figure 3 & 4 H – J, M)**. The mRNA abundance of
the DCT sodium-chloride cotransporter (NCC) was significantly lower in L919X KI/KI mice
with a similar trend in R607H KI mice (**Supplementary Figure 3 & 4 J**). We finally
assessed the mRNA abundance of the sodium/potassium/chloride cotransporter NKCC2
and sodium/proton exchanger NHE3 in whole kidneys, with the caveat that NHE3 is also
abundantly expressed in the proximal tubule. These levels did not differ between L919X
KI/KI and WT counterparts, but were significantly decreased in R607H KI/KI mice
compared to WT littermates (**Supplementary Figure 3 & 4 H, I**). Overall, these results
highlight paracellular adaptations in both KI/KI mice, potentially in an attempt to maintain
salt homeostasis.

Discussion

In this study, we characterized a second dRTA mouse model that carries the murine Ae1 L919X mutation orthologous to human dominant R901X dRTA mutation. In agreement with the previously published R607H KI/KI model, at baseline, the L919X KI/KI mice were not able to acidify their urine to the same extent as WT littermates, mimicking the main physiological defect observed in dRTA patients. In perfused kidneys from homozygous L919X and R607H KI mice, we observed decreased Ae1 gene expression reflecting a loss of A-ICs. Overall, our results support that both L919X and R607H KI/KI mouse lines reproduce dRTA and thus are good models to investigate the underlying pathophysiology.

Our results also reveal a more pronounced phenotype in R607H KI/KI compared to L919X KI/KI mice with slight differences in abundance of mRNA of Henle's loop and CD markers. One study reporting dominant dRTA mutations in the *SLC4A1* gene found that despite phenotypic similarities, the R901X mutation caused a milder phenotype than the R598H. This includes a lower plasma bicarbonate concentration and a more severe hypokalemia in patients with the R589H mutation compared to R901X[22]. Our results confirm this trend with significant differences observed in R607H KI/KI mice versus WT littermates but only similar trends in the L919X KI mice.

We challenged homozygous Ae1 L919X and R607H KI/KI mice and the respective controls with an acute salt-depleted diet or a salt-depleted diet in combination with an acid challenge to assess whether these mice develop a sodium wasting phenotype as shown for some dRTA patients[13]. Both dRTA mutant mice exhibit renal sodium loss compared to WT controls. Both KI/KI mice had elevated plasma sodium and chloride levels with increased abundance of renal renin mRNA suggestive of a volume contracted state despite a similar volume of water consumed and urine osmolality. Importantly, we found an increase in claudin 4 and claudin 10b gene and protein expression in both homozygous L919X and R607H KI mice compared to WT. A previous study feeding a salt restricted diet to a B1-H⁺-ATPase knockout mouse model elucidated one mechanism: loss of *Atp6v1b1* impaired sodium reabsorption via pendrin/NDCBE in B-ICs. These mice also had increased urinary ATP and prostaglandin E2 (PGE2) release (known sodium movement inhibitors), which decreased ENaC expression on PCs further contributing to sodium wasting[15,47]. Consistent with these findings, our experiments showed decreased pendrin and Ae4 gene expression levels, while NDCBE gene expression was increased in R607H KI/KI mice. We also observed increased γ-ENaC (SCNN1G) mRNA and protein

abundance in R607H KI/KI, but no significant difference in renal outer medullary potassium channel ROMK gene expression. These trends were also observed in L919X KI/KI mice, with the additional finding of decreased NCC mRNA abundance.

Both mouse lines displayed a similar alteration in their paracellular pathway in Henle's loop as an increase in claudin-4[6,45,4] and claudin-10b[42] mRNA abundance was observed after acid challenge with salt depletion. Claudin-4 functionally interacts with kAe1, as expressing the exchanger in immortalized inner medullary collecting duct cells increases transepithelial permeability to both sodium and chloride in a mechanism dependent in part on claudin-4 and with-no-lysine kinase 4[48,49]. Although present in tight junction, claudin 10b also colocalizes with the sodium/potassium ATPase and chloride channels in the basolateral infoldings of the TAL[45]. While its role there remains unclear, it may play a crucial role in facilitating high electrolyte flow, and preventing basolateral water influx. Therefore, increased expression of claudin 10b may represent a compensatory mechanism to enhance urine concentration and regulate electrolyte balance.

The upregulation of claudins in Henle's loop points towards a possible compensatory role of this segment in urinary acidification and sodium reabsorption. The ascending limb of Henle's loop is a site for urinary acidification via the combined action of apical NHE3 and basolateral NBCn1[50,51]. As shown by our experiments, mutant mice still acidify their urine although they are IC-depleted, which points towards a possible contribution of this segment to urinary acidification, in addition to sodium and chloride reabsorption. The function of acid-base transporters is tightly coupled to that of salts in the TAL, through apical NHE3 and the basolateral electroneutral sodium bicarbonate co-transporter, NBCn1[49, 50]. NBCn1 facilitates urine acidification during metabolic acidosis[50], promoting ammonium reabsorption across the renal tubules, rapidly restoring arterial blood pH and facilitating its secretion by the CD[51]. Previous work reported that R607H KI/KI mice have increased NCBn1 protein abundance at steady state compared to WT mice[33]. The potential upregulation of this protein in the KI/KI mice lines could provide an explanation for their ability to effectively regulate plasma pH and acidify their urine to some extent despite the loss of IC. Additionally, NKCC2 and ROMK in the TAL may cause a net chloride reabsorption creating a lumen-positive transepithelial potential driving paracellular sodium reabsorption through claudin 10b.

After the salt-depleted acid challenge, we detected a significantly decreased
mRNA abundance of NHE3 and NKCC2 in homozygous R607H KI mice compared to
WT. The relevance of this finding may be limited as both NHE3 and NKCC2 proteins are
post-translationally regulated. NKCC2 regulation occurs via membrane trafficking,
phosphorylation, and protein-protein interactions[52]. NHE3 is regulated through
phosphorylation, and its activity can be inhibited by aldosterone[53,54], which may be
relevant in the context of salt depletion.
360
Overall, our results show that both L919X and R607H and KI/KI mice recapitulate
dRTA symptoms, including the urinary sodium loss seen in patients. Our results reveal
potentially relevant physiological mechanisms for this complex disease and point to a
potential compensatory acidification and salt reabsorption in earlier nephron segments.
Therefore, these mice provide an excellent tool to further investigate the
pathophysiological mechanisms resulting in this disease.
367
368

Acknowledgments

This study was funded by the Canadian Institutes of Health Research

(PJT#168871) and the Kidney Foundation of Canada (2020KHRG-666615) to E.C.

C.A.H. was supported by a grant of the DFG (HU 800/7-2). D.E. is supported by a grant

from the French National Research Agency (ANR-14-CE12-0013-01), and by the National

Center for Precision Diabetic Medicine – PreciDIAB. (ANR-18-IBHU-0001). P.M. received

a Canada Graduate Scholarship-Master's graduate studentship from the Canadian

Institutes of Health Research, a University of Alberta Graduate Recruitment Scholarship

and a Walter H. Johns Graduate Fellowship from the University of Alberta. A.K.M.S.U.

received an NSERC CREATE graduate studentship, G.E. received a Graduate Student

Engagement Scholarship, a Faculty of Medicine and Dentistry Delnor Scholarship and a

Faculty of Medicine and Dentistry 75[th] Anniversary award. F. C. was awarded an

University of Alberta Graduate Recruitment Scholarship.

References

1. Mohebbi, N. & Wagner, C. A. Pathophysiology, diagnosis and treatment of

inherited distal renal tubular acidosis. *J. Nephrol.* **31**, 511–522 (2018).

2. Alper, S. L. Familial renal tubular acidosis. *J Nephrol* **23 Suppl 1**, S57-76 (2010).

3. Batlle, D., Ghanekar, H., Jain, S. & Mitra, A. Hereditary distal renal tubular

acidosis: new understandings. *Annu Rev Med* **52**, 471–484 (2001).

4. Giglio, S., Montini, G., Trepiccione, F., Gambaro, G. & Emma, F. Distal renal

tubular acidosis: a systematic approach from diagnosis to treatment. *Journal of*

Nephrology **34**, 2073–2083 (2021).

5. Wagner, C. A. *et al.* The pathophysiology of distal renal tubular acidosis. *Nat.*

Rev. Nephrol. 2023 196 **19**, 384–400 (2023).

6. Fry, A. C. & Karet, F. E. Inherited renal acidoses. *Physiol.* **22**, 202–211 (2007).

7. Karet, F. E. Inherited distal renal tubular acidosis. *J Am Soc Nephrol* **13**, 2178–

2184 (2002).

8. Laing, C. M., Toye, A. M., Capasso, G. & Unwin, R. J. Renal tubular acidosis:

developments in our understanding of the molecular basis. *Int J Biochem Cell Biol*

37, 1151–1161 (2005).

9. Unwin, R. J. & Capasso, G. The renal tubular acidoses. *J R Soc Med* **94**, 221–

225 (2001).

10. Aronson, P. S. & Giebisch, G. Effects of pH on potassium: New explanations for

old observations. *Journal of the American Society of Nephrology* **22**, 1981–1989

(2011).

11. Alonso-Varela, M., Gil-Peña, H. & Santos, F. Incomplete distal renal tubular

acidosis in children. *Acta Paediatr.* **109**, 2243–2250 (2020).

12. Sebastian, A., McSherry, E. & Morris Jr., R. C. Impaired renal conservation of

sodium and chloride during sustained correction of systemic acidosis in patients

with type 1, classic renal tubular acidosis. *J Clin Invest* **58**, 454–469 (1976).

13. Sebastian, A., Mcsherry, E. & Moymis, R. C. Impaired Renal Conservation of

Sodium and Chloride during Sustained Correction of Systemic Acidosis in

Patients with Type 1, Classic Renal Tubular Acidosis.

14. Lashhab, R., Ullah, A. K. M. S. & Cordat, E. Renal collecting duct physiology and

pathophysiology. *Biochem. Cell Biol.* 1–9 (2018). doi:10.1139/bcb-2018-0192

15. Roy, A., Al-bataineh, M. M. & Pastor-Soler, N. M. Collecting duct intercalated cell

function and regulation. *Clin. J. Am. Soc. Nephrol.* **10**, 305–24 (2015).

16. Saxena, V. *et al.* Kidney intercalated cells are phagocytic and acidify internalized

uropathogenic Escherichia coli. *Nat. Commun.* **12**, (2021).

17. Purkerson, J. M., Heintz, E. V., Nakamori, A. & Schwartz, G. J. Insights into

acidosis-induced regulation of SLC26A4 (pendrin) and SLC4A9 (AE4)

transporters using three-dimensional morphometric analysis of β-intercalated

cells. *Am. J. Physiol. - Ren. Physiol.* **307**, F601 (2014).

18. Peña-Münzenmayer, G., George, A. T., Shull, G. E., Melvin, J. E. & Catalán, M.

A. Ae4 (Slc4a9) is an electroneutral monovalent cation-dependent Cl-/HCO3-

exchanger. *J. Gen. Physiol.* **147**, 423–436 (2016).

19. SM, W., JW, V. & CA, R. The Renal Physiology of Pendrin-Positive Intercalated

Cells. *Physiol. Rev.* **100**, 1119–1147 (2020).

20. Bruce, L. J. *et al.* Familial distal renal tubular acidosis is associated with mutations

in the red cell anion exchanger (Band 3, AE1) gene. *J Clin Invest* **100**, 1693–1707

(1997).

21.   Bruce, L. J. *et al.* Band 3 mutations, renal tubular acidosis and South-East Asian

ovalocytosis in Malaysia and Papua New Guinea: loss of up to 95% band 3

transport in red cells. *Biochem J* **350 Pt 1**, 41–51 (2000).

22.   Karet, F. E. *et al.* Mutations in the chloride-bicarbonate exchanger gene AE1

cause autosomal dominant but not autosomal recessive distal renal tubular

acidosis. *Proc Natl Acad Sci U S A* **95**, 6337–6342 (1998).

23.   Jarolim, P. *et al.* Autosomal dominant distal renal tubular acidosis is associated in

three families with heterozygosity for the R589H mutation in the AE1 (Band 3) Cl-

/HCO3/- exchanger. *J. Biol. Chem.* **273**, 6380–6388 (1998).

24.   Arakawa, T. *et al.* Crystal structure of the anion exchanger domain of human

erythrocyte band 3. *Science (80-.).* **350**, 680–684 (2015).

25.   Zhekova, H. R. *et al.* CryoEM structures of anion exchanger 1 capture multiple

states of inward- and outward-facing conformations. *Commun. Biol.* **5**, 1–13

(2022).

26.   Reithmeier, R. A. F. *et al.* Band 3, the human red cell chloride/bicarbonate anion

exchanger (AE1, SLC4A1), in a structural context. *Biochimica et Biophysica Acta

- Biomembranes* **1858**, 1507–1532 (2016).

27.   Cordat, E. *et al.* Dominant and recessive distal renal tubular acidosis mutations of

kidney anion exchanger 1 induce distinct trafficking defects in MDCK cells. *Traffic*

7, 117–128 (2006).

28.   Toye, A. M., Banting, G. & Tanner, M. J. A. Regions of human kidney anion

exchanger 1 (kAE1) required for basolateral targeting of kAE1 in polarized kidney cells: Mis-targeting explains dominant renal tubular acidosis (dRTA). *J. Cell Sci.* **117**, 1399–1410 (2004).

29. Quilty, J. A. A., Cordat, E. & Reithmeier, R. A. F. A. Impaired trafficking of human kidney anion exchanger (kAE1) caused by hetero-oligomer formation with a truncated mutant associated with distal renal tubular acidosis. *Biochem J* **368**, 895–903 (2002).

30. Toye, A. M., Bruce, L. J., Unwin, R. J., Wrong, O. & Tanner, M. J. Band 3 Walton, a C-terminal deletion associated with distal renal tubular acidosis, is expressed in the red cell membrane but retained internally in kidney cells. *Blood* **99**, 342–7. (2002).

31. Cordat, E., Li, J. & Reithmeier, R. A. F. Carboxyl-terminal truncations of human anion exchanger impair its trafficking to the plasma membrane. *Traffic* **4**, 642–651 (2003).

32. Mumtaz, R. *et al.* Intercalated Cell Depletion and Vacuolar H+-ATPase Mistargeting in an Ae1 R607H Knockin Model. *J. Am. Soc. Nephrol.* **28**, 1507–1520 (2017).

33. Stehberger, P. A. *et al.* Distal renal tubular acidosis in mice lacking the AE1 (band3) Cl-/HCO3- exchanger (slc4a1). *J Am Soc Nephrol* **18**, 1408–1418 (2007).

34. DeFranco, P. E., Haragism, L., Schmitz, P. G., Bastani, B. & Li, J. P. Absence of vacuolar H+-ATPase pump in the collecting duct of a patient with hypokalemic distal renal tubular acidosis and Sjogren's syndrome. *J. Am. Soc. Nephrol.* **6**, 295–301 (1995).

35.   Vichot, A. A. *et al.* Loss of kAE1 expression in collecting ducts of end-stage

kidneys from a family with SLC4A1 G609R-associated distal renal tubular

acidosis. *Clin. Kidney J.* **10**, 135–140 (2017).

36.   Plain, A. *et al.* Claudin-12 knockout mice demonstrate reduced proximal tubule

calcium permeability. *Int. J. Mol. Sci.* **21**, 1–18 (2020).

37.   Grinstein, M., Dingwall, H. L., Shah, R. R., Capellini, T. D. & Galloway, J. L. A

robust method for RNA extraction and purification from a single adult mouse

tendon. *PeerJ* **2018**, 1–17 (2018).

38.   Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using

real-time quantitative PCR and the 2-ΔΔCT method. *Methods* **25**, 402–408

(2001).

39.   Kim, E. Y. *et al.* Altered regulation of renal acid base transporters in response to

ammonium chloride loading in rats. *Korean J. Physiol. Pharmacol.* **16**, 91–95

(2012).

40.   López-Cayuqueo, K. I. *et al.* A mouse model of pseudohypoaldosteronism type II

reveals a novel mechanism of renal tubular acidosis. *Kidney Int.* **94**, 514–523

(2018).

41.   Chambrey, R. & Trepiccione, F. Relative Roles of Principal and Intercalated Cells

in the Regulation of Sodium Balance and Blood Pressure. *Current Hypertension*

Reports **17**, (2015).

42.   Hou, J., Rajagopal, M. & Yu, A. S. Claudins and the kidney. *Annu Rev Physiol* **75**,

479–501 (2013).

43.   Leiz, J. & Schmidt-Ott, K. M. Claudins in the renal collecting duct. *International*

Journal of Molecular Sciences **21**, (2020).

44.   Yu, A. S. L. Claudins and the Kidney. *J. Am. Soc. Nephrol.* **26**, 11–19 (2015).

45.   Quintanova, C. *et al.* Unrecognized role of claudin-10b in basolateral membrane

infoldings of the thick ascending limb. *Ann. N. Y. Acad. Sci.* **1517**, 266–278

(2022).

46.   Hou, J., Renigunta, A., Yang, J. & Waldegger, S. Claudin-4 forms paracellular

chloride channel in the kidney and requires claudin-8 for tight junction localization.

Proc. Natl. Acad. Sci. **107**, 18010–18015 (2010).

47.   Gueutin, V. *et al.* Renal β-intercalated cells maintain body fluid and electrolyte

balance. *J. Clin. Invest.* **123**, 4219–4231 (2013).

48.   Lashhab, R., Essuman, G., Chavez-Canales, M., Alexander, R. T. & Cordat, E.

Expression of the kidney anion exchanger 1 affects WNK4 and SPAK

phosphorylation and results in claudin-4 phosphorylation. *Heliyon* **9**, e22280

(2023).

49.   Lashhab, R. *et al.* The kidney anion exchanger 1 affects tight junction properties

via claudin-4. *Sci. Rep.* **9**, 3099 (2019).

50.   de Bruijn, P. I. A. *et al.* Furosemide-induced urinary acidification is caused by

pronounced H+ secretion in the thick ascending limb. *Am. J. Physiol. Renal*

Physiol. **309**, F146–F153 (2015).

51.   Olsen, J. S. M. *et al.* NBCn1 Increases NH4 + Reabsorption Across Thick

Ascending Limbs, the Capacity for Urinary NH4 + Excretion, and Early Recovery

from Metabolic Acidosis. *J. Am. Soc. Nephrol.* **32**, 852–865 (2021).

52.   Ares, G. R., Caceres, P. S. & Ortiz, P. A. Molecular regulation of NKCC2 in the

thick ascending limb. *American Journal of Physiology - Renal Physiology* **301**,

F1143 (2011).

53.   Girardi, A. C. C. & di Sole, F. Deciphering the mechanisms of the Na+/H+

exchanger-3 regulation in organ dysfunction. *American Journal of Physiology -*

Cell Physiology **302**, 1569–1587 (2012).

54.   Watts, B. A., George, T. & Good, D. W. Aldosterone inhibits apical NHE3 and

HCO_3^- absorption via a nongenomic ERK-dependent pathway in medullary thick

ascending limb. *Am. J. Physiol. Physiol.* **291**, F1005–F1013 (2006).

 Figure Legends

Figure 1: Generation and characterization of the L919X KI mice. (**A**)Targeting and

screening strategy with murine Ae1 locus (top) and targeted locus (middle) with locus

after removal of the selection cassette by Cre recombination. Right, Southern blot result

and sequencing validating the introduction of the mutation to the locus. (**B**) Graph

representing the gene expression of Ae1 in perfused whole kidney from WT or L919X

KI/KI mice. (C) immunofluorescence images from confocal microscopy of WT,

heterozygous or homozygous L919X KI/KI mice medullary or cortical kidney sections

stained with anti-pendrin (red) or anti-AE1 (green) antibodies. Scale bar corresponds to

50 µm. (D) Quantification of A-IC (Ae1 positive) and B-IC (pendrin positive) in the cortex

and medulla of WT or homozygous Ae1 L919X KI mice. Error bars correspond to means

± SEM, n = 4-6 kidney sections, ***$P < 0.001$ using Two-way ANOVA.

Figure 2: At baseline, both R607H and L919X KI/KI mice exhibit an alkaline urine and

decreased urinary ammonium secretion. Urinary pH (**A**), ammonium (**B**), osmolality (**C**),

sodium/creatinine ratio (**C**), chloride/creatinine ratio (**D**) and potassium/creatinine ratio (**E**)

comparison between WT or L919X KI/KI mice. Urinary pH (**G**), ammonium (**H**), osmolality

(**I**), sodium/creatinine ratio (**J)**, chloride/creatinine ratio (**K**) and potassium/creatinine ratio

(**L**) comparison between WT or R607H KI/KI mice. Error bars correspond to means ±

SEM, **$P < 0.01$, ****$P < 0.0001$ using Student's t-test or Mann-Whitney test.

Figure 3: After a salt-depleted acid diet, both R607H and L919X KI/KI mice are

hypernatremic and hyperchloremic. Plasma sodium (**A**) and chloride (**B**) concentrations

from L919X KI/KI mice or WT littermates. (**C**) Renin gene expression in L919X KI/KI mice

or WT littermates. Urine osmolality (**D**) and water consumed (**E**) by L919X KI/KI mice or

WT littermates over the 14 day-time course of the diet. Plasma sodium (**G**) and chloride

(**H**) concentrations from R607H KI/KI mice or WT littermates. (**I**) Renin gene expression

in R607H KI/KI mice or WT littermates. Urine osmolality (**J**) and water consumed (**K)** by

R607H KI/KI mice or WT littermates over the 14 day-time course of the diet. Error bars

correspond to means ± SEM, *P < 0.05, ***P < 0.001, ****P < 0.0001 using Student's t-

test or two-way ANOVA with Tukey's multiple comparison test.

Figure 4: After a salt-depleted acid diet, L919X KI/KI mice produce an alkaline urine and

waste urinary sodium and chloride. (**A**) Day 14 (left) and trend (right) over the 14

experimental days urine pH in L919X KI/KI mice or WT littermates. (**B**) Day 14 (left) and

trend (right) urinary ammonium/creatinine ratio in L919X KI/KI mice or WT littermates.

Day 14 urinary sodium/creatinine ratio (**C**), sodium/chloride ratio (**D**) and

potassium/creatinine ratio (**E**). (**F-H**) trend of sodium/creatinine, sodium/chloride and

potassium/creatinine ratios over the time course of the experiment in L919X KI/KI mice

or WT littermates. Error bars correspond to means ± SEM, *P < 0.05, **P < 0.01, ***P <

0.001, ****P < 0.0001 using Student's t-test or two-way ANOVA with Tukey's multiple

comparison test.

Figure 5: After a salt-depleted acid diet, R607H KI/KI mice produce an alkaline urine and

waste urinary sodium and chloride. (**A**) Day 14 (left) and trend (right) over the 14

experimental days urine pH in R607H KI/KI mice or WT littermates. (**B**) Day 14 (left) and

trend (right) urinary ammonium/creatinine ratio in R607H KI/KI mice or WT littermates.

Day 14 urinary sodium/creatinine ratio (**C**), sodium/chloride ratio (**D**) and

potassium/creatinine ratio (**E**). (**F-H**) trend of sodium/creatinine, sodium/chloride and

potassium/creatinine ratios over the time course of the experiment in R607H KI/KI mice

or WT littermates. Error bars correspond to means ± SEM, *P < 0.05, **P < 0.01, ***P <

0.001, ****P < 0.0001 using Student's t-test or two-way ANOVA with Tukey's multiple

comparison test.

Figure 6: Claudin-4 and claudin-10b are upregulated in both L919X and R607H KI/KI

mice after a salt depleted acid diet. Claudin-4 gene expression (**A & E**) and protein

abundance (**B & F**) with a representative immunoblot image (right). Claudin-10b gene

expression (**C & G**) and claudin-10 protein abundance (**D & H**) with a representative

immunoblot image (right). Error bars correspond to means ± SEM, *P < 0.05, **P < 0.01,

using Student's t-test or two-way ANOVA with Tukey's multiple comparison test.

 Supplementary Table 1: Primers and Probes Used for qPCR on Mouse Genes

Name	Forward	Reverse	Probe
AE4 NM_N172830	5'-CTCTGTGTCTTTGAGCTTTGC-3'	5'-TCTGCCTATGGTTGCTTCTG-3'	5'-/56-FAM/TGTTTCCTC/ZEN/CTGTGCCTGGGTATTG/3IABKFQ/-3'
alpha-ENaC NM_011324.2	5'-CAGTGATGTCCCTGTCAAGAA-3'	5'-CTTGGGCTTAGGGTAGAAGATG-3'	5'- /56-FAM/TGA TCA AGA /ZEN/AGT GTGGCT GTG CCT /3IABKFQ/ -3'
Atp6v1b1 NM_134157.1	5'-CTTCTCACACTGGTAGGCAAG-	5'-AGTCAGATTTCGAGCAGAATGG-3'	5'-/56-FAM/CCGCTCAAT/ZEN/CGTAGGGTCATTGGC/3IABkFQ/-3'
Cldn10b NM_N021386.1	5'-GCAGCGATCATTAGTCCTCTAC-3'	5'-CCACAGCCACTTATTTTGCC-3'	5-/56-FAM/A ATCGGTA A/ZEN/CGCAGATCTCCACAGG/3IABkFQ/-3'
Cld4 NM_N009903.1	5-'CAGGACTGCCAAGGAGATTC-3'	5-'AACACTTTCTCAGCCCTCTG-3'	5-'/56-FAM/CATAGACGC/ZEN/CATCGCTCAGCCTC/3IABkFQ/-3*
gamma-ENaC NM_N011326.1	5'-GTCAGAGGTGTCATTTGAGCA-3'	5'-GAGAACGAGAAGGGAAAGGC-3'	5'-/56-FAM/TCGGAAGCG/ZEN/GAAAATCAGGGGAA/3IABKFQ/-3'
kAE1 NM_009902.4	5'-AAGAGGTCAAGGAACAGCG-3'	5'-GTACAGGAAGATGCCGAAGAG-3'	5-/56-FAM/CCCACAAGC/ZEN/ACAGAGACCAGGAG/3IABkFQ/-3'
NCC NM_019415	5'-CAGATCAACAGGATGGACGAAG-3'	5'-GCAATCATGTCCTCAAACCG-3'	5'-/56-FAM/TCCCGACAT/ZEN/CAACCAGA AGCCC/3IABkFQ/-3'
NDCBE NM_021530.2	5'-ATGGGAAGATGGTGAGGAAC-3'	5'-ACGTGCTCTTTTGGTCCTG-3'	5'-/56-FAM/AAGGTGCTG/ZEN/GAGACGATGAAGGG/3IABKFQ/-3'
NHE3 NM_N001081060.1	5'-CCAAACAGCAAGAAATCCAGG-3'	5'-TCCCTCTATGGTGTCTTCCTC-3'	5'-/56-FAM/CCA ATTTC/ZEN/AGTTCACCCATCA AGCCAC/3IABKFQ/-3'
NKCC2 NM_183354	5'-CAAGGAACATTACCAAGCC-3'	5'-CACCAGTTTGATTGAACTCC-3*	5'-/56-FAM/CAAGAAAGA/ZEN/CGGCAACATCAGCAGC/3IABKFQ/-3'
Pendrin NM_011867	5'-TGATGGAGGCAGAGATGAATG-3'	5'-CAGAAAACACTGAGAGACTG-3'	5-/56-FAM/ATGTTCAGG/ZEN/ATGAGGCCATGCGTAG/3IABkFQ/-3'
Renin	TaqMan Gene Expression Assay (Applied Biosystems) REF# 4331182		

RPLP0 Mm.PT.58.43894205	5'-TTATA ACCCTGA AGGCTCGAC-3'	5'-CGCTTGTACCCATTG ATGATG-3'	5'-/56-FAM/AGGCCCTGC/ZEN/ACTC TCGCTT/3IABkFQ/-3'

597 **Supplementary Table 2**: Antibodies Used for Immunoblotting

Name	Reference	Reference	Dilution
Rb anti-Claudin-10	Primary	Antibodies.com	1:1000
Rb anti-Claudin-4	Primary	Invitrogen	1:1000
Rb anti-ENaC gamma	Primary	StressMarq Biosciences Inc	1:5000
Direct-Blot HRP anti-β Actin	Primary	BioLegend	1:10000

598
599

600
601
602 **Supplementary Table 3.** Baseline Plasma Characterization of WT and L919X Ae1 KI/KI Mice.
603

Parameter	Units	WT (n=10)	L919X AE1 KI/KI (=10)	p-value
Na^+	mM	147.5 ± 0.563	146.9 ± 0.623	0.4327
Cl^-	mM	118.8 ± 0.964	119.6 ± 1.536	0.6644
K^+	mM	4.633 ± 0.167	4.610 ± 0.196	0.7029
HCO_3^-	mM	18.38 ± 0.563	19.95 ± 0.511	0.0930
pH		7.289 ± 0.002	7.249 ± 0.009	0.1186
TCO_2	mmHg	19.50 ± 0.749	21.30 ± 0.517	0.0636
BUN	mg/dl	22.70 ± 1.382	19.80 ± 0.987	0.1050
Glucose	mg/dl	181 ± 13.11	198.3 ±10.58	0.3181
HCT	%	36.80 ± 0.554	36 ± 0.789	0.4174
pCO2	mmHg	38.99 ± 2.856	45.75 ± 1.595	0.0525
AnGAP	mM	14.6 ± 0.618	13.22 ± 1.325	0.1197
Hb	g/dl	12.53 ± 0.188	12.23 ± 0.269	0.3729
BEecf	mM	-8.100 ±0.781	-7.100 ± 0.504	0.2963

604 *Data is presented as mean ± SEM. Student's T-Test or Mann-Whitney test used where*
605 *appropriate.*

606
607 **Supplementary Table 4.** Baseline Plasma Characterization of WT and R607H Ae1 KI/KI Mice.
608

Parameter	Units	WT (n=11)	R607H AE1 KI/KI (n=9)	p-value
Na^+	mM	146.4 ± 0.472	147.1 ± 0.586	0.3148
Cl^-	mM	118.2 ±1.110	118.5 ± 0.619	0.1409
K^+	**mM**	**4.5 ± 0.142**	**4.920 ± 0.077***	**0.0207**
HCO_3^-	mM	18.66 ±0.627	19.07 ±0.463	0.6142
pH		7.256 ± 0.014	7.258 ± 0.013	0.9102
TCO_2	mmHg	19.91 ± 0.653	20.20 ± 0.533	0.7370
BUN	**mg/dl**	**21.09 ± 1.351**	**23.60 ± 0.921***	**0.0438**
Glucose	mg/dl	196.3 ± 7.065	197.3 ± 12.31	0.9417
HCT	%	37.73 ± 1.221	38.60 ± 0.340	0.5178
pCO2	mmHg	42.05 ± 1.467	42.72 ± 1.410	0.7417
AnGAP	mM	13.27 ± 1.538	14.60 ± 0.600	0.9863
Hb	g/dl	12.82 ± 0.414	13.14 ± 0.117	0.4826
BEecf	mM	-8.545 ± 0.731	-8.100 ± 0.586	0.5666

609 *Data is presented as mean ± SEM. Student's T-Test or Mann-Whitney test used where*
610 *appropriate. **Significant data is bolded.***

611
612

Supplementary Table 5. Plasma Characterization of WT and L919X Ae1 KI/KI Mice Following
24 hours of Salt Restriction

Parameter	Units	WT (n=10)	L919X AE1 KI/KI (n=10)	p-value
Na^+	mM	147.9 ± 0.690	147.7 ± 0.578	0.8268
Cl^-	mM	117.6 ± 0.897	120.9 ± 1.329	0.0543
K^+	mM	4.980 ± 0.158	5.180 ± 0.464	0.5661
HCO_3^-	mM	20.59 ± 0.613	20.40 ± 0.825	0.8554
pH		7.240 ± 0.021	7.248 ± 0.033	0.8469
TCO_2	mmHg	22.20 ± 0.593	21.80 ± 0.854	0.7048
BUN	mg/dl	30.10 ± 4.365	23.30 ± 0.989	0.1249
Glucose	mg/dl	164.5 ± 12.69	157.1 ± 9.719	0.6490
HCT	%	40.70 ± 0.667	38.90 ± 0.690	0.1152
pCO2	mmHg	47.95 ± 1.127	47.74 ± 3.411	0.9540
AnGAP	mM	14.90 ± 0.795	11.89 ± 1.369	0.0677
Hb	g/dl	13.84 ± 0.225	13.22 ± 0.237	0.0741
BEecf	mM	-6.900 ± 0.971	-6.800 ± 1.073	0.9457

*Data is presented as mean ± SEM. Student's T-Test or Mann-Whitney test used where appropriate. **Significant data is bolded.***

Supplementary Table 6. Plasma Characterization of WT and R607H Ae1 KI/KI Mice Following
24 hours of Salt Restriction

Parameter	Units	WT (n=12)	R607H AE1 KI/KI (n=14)	p-value
Na^+	mM	146.3 ± 0.890	148.1 ± 0.275	0.8166
Cl^-	mM	121.5 ± 2.062	122.9 ± 0.769	0.5196
K^+	**mM**	**4.475 ± 0.100**	**4.843 ± 0.100**	**0.0162**
HCO_3^-	mM	19.04 ± 0.965	18.39 ± 0.540	0.5437
pH		7.203 ± 0.021	7.215 ± 0.014	0.6389
TCO_2	mmHg	20.50 ± 1.004	19.86 ± 0.592	0.9305
BUN	mg/dl	23.75 ± 1.473	25.00 ± 1.267	0.5236
Glucose	mg/dl	154.5 ± 11.59	182.6 ± 9.245	0.0667
HCT	%	37.50 ± 2.183	37.86 ± 0.710	0.1096
pCO2	mmHg	48.33 ± 2.356	45.75 ± 1.901	0.3983
AnGAP	mM	10.25 ± 2.394	11.64 ± 0.608	0.1453
Hb	g/dl	12.76 ± 0.741	12.86 ± 0.246	0.0994
BEecf	mM	-9.000 ± 1.206	-9.357 ± 0.608	0.7847

*Data is presented as mean ± SEM. Student's T-Test or Mann-Whitney test used where appropriate. **Significant data is bolded.***

Supplementary Table 7. Plasma Characterization of WT and L919X Ae1 KI/KI Mice Following
Salt Depleted Acid Load.

Parameter	Units	WT (n=14)	L919X AE1 KI/KI (n=14)	p-value
Na$^+$	**mM**	**149.4 ± 0.644**	**156.4 ± 1.152**	**< 0.0001**
Cl$^-$	**mM**	**126.9 ± 2.087**	**135.1 ± 0.900**	**0.0140**
K$^+$	mM	4.777 ± 0.154	4.669 ± 0.170	0.6430
HCO$_3^-$	mM	15.63 ± 0.940	13.88 ± 0.470	0.1164
pH		7.170 ± 0.030	7.131 ± 0.014	0.0683
TCO$_2$	**mmHg**	**17.62 ± 0.789**	**15.15 ± 0.504**	**0.0147**
BUN	**mg/dl**	**23.75 ± 0.954**	**33.62 ± 1.723**	**< 0.0001**
Glucose	mg/dl	148.9 ± 12.93	131.92 ± 5.054	0.2452
HCT	%	37.38 ± 1.274	37.46 ± 1.264	0.9713
pCO2	mmHg	42.84 ± 2.581	41.72 ± 1.632	0.7216
AnGAP	mM	13.33 ± 1.694	13.09 ± 0.948	0.9042
Hb	g/dl	12.70 ± 0.432	12.74 ± 0.430	0.9713
BEecf	mM	-12.93 ± 1.294	-15.31 ± 0.603	0.1169

*Data is presented as mean ± SEM. Student's T-Test or Mann-Whitney test used where appropriate. **Significant data is bolded.***

Supplementary Table 8. Plasma Characterization of WT and R607H Ae1 KI/KI Mice Following Salt Depleted Acid Load.

Parameter	Units	WT (n=14)	R607H AE1 KI/KI (n = 18)	p-value
Na⁺	**mM**	**145.2 ± 0.882**	**156.4 ± 0.903**	**< 0.0001**
Cl⁻	**mM**	**126.8 ± 1.959**	**138.6 ± 0.536**	**< 0.0001**
K⁺	mM	4.557 ± 0.229	4.406 ± 0.185	0.6069
HCO₃⁻	**mM**	**14.15 ± 0.921**	**11.64 ± 0.775**	**0.0448**
pH		**7.192 ± 0.025**	**7.085 ± 0.020**	**0.0016**
TCO₂	mmHg	15.29 ± 0.963	12.83 ± 0.797	0.0572
BUN	**mg/dl**	**22.50 ± 1.725**	**34.17 ± 3.618**	**0.0026**
Glucose	**mg/dl**	**163.2 ± 11.93**	**118.8 ± 7.284**	**0.0024**
HCT	%	34.00 ± 2.176	32.56 ± 2.297	0.8434
pCO2	mmHg	46.21 ± 1.429	38.56 ± 2.400	0.4397
AnGAP	mM	10.08 ± 2.065	13.43 ± 1.462	0.6004
Hb	**g/dl**	**11.56 ± 0.739**	**13.55 ± 0.189**	**0.0489**
BEecf	**mM**	**-14.07 ± 1.273**	**-18.22 ± 0.967**	**0.0129**

Data is presented as mean ± SEM. Student's T-Test or Mann-Whitney test used where appropriate. **Significant data is bolded.**

Figure 1 Mungara et al.

Figure 4 Mungara et al.

Figure 6 Mungara et al.