Predictive power of different *Akkermansia* phylogroups in clinical response to PD-1

blockade against non-small cell lung cancer

Peixin Fan[1], Mi Ni[1], Yu Fan[1], Magdalena Ksiezarek[1], Gang Fang[1*]

[1]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New

York, NY, USA.

[*]Correspondence to Gang Fang, gang.fang@mssm.edu

Summary

Immune checkpoint blockade has emerged as a promising form of cancer therapy. However, only some patients respond to checkpoint inhibitors, while a significant proportion of patients do not, calling for the discovery of reliable biomarkers. Recent studies reported the importance of the gut microbiome in the clinical response to PD-1 blockade against advanced non-small cell lung cancer (NSCLC), highlighting Akkermansia muciniphila as a candidate biomarker. Motivated by the genomic and phenotypic differences across Akkermansia muciniphila strains and Akkermansia (Akk) phylogroups (AmIa, AmIb, AmII, AmIII and AmIV), we analyzed fecal metagenomic sequencing data from three publicly available NSCLC cohorts (n=425). Encouragingly, we found that patients' responses to PD-1 blockade are significantly different across different Akk phylogroups, highlighting the relatively stronger association between AmIa and positive response than the other phylogroups. We built a machine learning model based on Akk gene profiles, which improved the predictive power and shed light on a group of Akk genes that may be associated with the response to PD-1 blockade. In summary, our study underlines the benefits of high-resolution analysis of Akk genomes in the search for biomarkers that may improve the prediction of patients' responses to cancer immunotherapy.

Key words:

Non-small cell lung cancer (NSCLC), immunotherapy, *Akkermansia muciniphila*, phylogroup

Introduction

Lung cancer is the third most common type of cancer in the US and the leading cause of cancer-related mortality worldwide[1,2]. Non-small cell lung cancer (NSCLC) is the most prevalent form of lung cancer (84%). The emergence of immunotherapy, especially the immune checkpoint blockade, has revolutionized the cancer treatments with great promise, especially for NSCLC[3,4,5]. However, only some patients experience durable responses to immune checkpoint inhibitors (ICIs), while others do not respond. For example, although NSCLC has a relatively higher response rate to ICIs than other cancer types, only ~20% of NSCLC patients respond favorably to ICIs as monotherapy[6]. Therefore, it is urgent to understand critical factors that influence the response to immunotherapy and discover predictive biomarkers that may guide personalized medicine.

Several promising biomarkers have been discovered for predicting responses to immunotherapy and have been incorporated into clinical practice, including the expression of immune checkpoint proteins (e.g., PD-1 or PD-L1), pre-existing immune responses, tumor mutation burden, and tumor-associated proteins[6,7,8,9,10]. From a recent study, bronchoalveolar lavage fluid analysis of cellular profiles or secreted molecules can help identify patients with NSCLC who respond to ICIs[11]. In addition, recent studies reported an association between patients' gut microbiome and their responses to immunotherapies[12,13]. Animal model-based studies found that antitumor activity of ICIs was lower in microbiome-depleted mice such as germ-free mice and mice treated with antibiotics, compared to specific pathogen-free mice[14,15]. Patient cohort studies showed significant differences in gut microbiome composition between NSCLC, renal cell carcinoma, and melanoma patients who responded to anti-PD-1 therapy and those who did not[15,16,17,18]. Fecal microbiota transplantation (FMT) was shown to influence the

ICIs treatment of refractory melanoma and colorectal cancer in clinical trials[19,20], suggesting

some specific gut microbes may serve as non-invasive predictive biomarkers.

For NSCLC, recent studies reported that *Akkermansia muciniphila*, a mucin-degrading

gut commensal bacterium, was associated with better response to PD-1-based immunotherapy,

and that it can enhance ICIs treatment in mice[21,22,23]. Functionally, *A. muciniphila* is associated

with beneficial effects on host metabolism and homeostatic immunity[24,25]. It can also directly

trigger intestinal adaptive immune responses through inducing T cell-dependent IgG1 and IgA[23].

Immunogenic activity of lipid (a diacyl phosphatidylethanolamine with two branched chains) and

peptides (e.g., Amuc_1100 and P9) produced by *A. muciniphila* were observed *in vitro*[26,27].

Notably, several studies have demonstrated the genomic and phenotypic differences across *A.*

muciniphila strains, which initiated further classification into *Akkermansia* (Akk) phylogroups

(primarily AmIa, AmIb, AmII, AmIII and AmIV). Among these phylogroups, only AmIa and AmIb

represent *A. muciniphila,* while the remaining are being actively characterized as potential new

Akkermansia species[28,29,30,31]. Different Akk phylogroups were shown to have significant

phenotypic differences that are associated with different host-microbe interactions: growth rates

in mucin, binding to epithelial surfaces, and activation of innate immune responses[28,29,30,31,32]. A

recent study also identified correlations between Akkermansia species and subspecies and

human health outcomes[31]. Consistently, two different Akk strains showed different capacities to

modulate differentiation of regulatory T cells (Tregs) in mice[15]. Based on these genetic and

phenotypic variations among Akk strains, we hypothesized that NSCLC patients carrying

different phylogroups of Akk in their gut microbiome have different responses to

immunotherapy.

To test this hypothesis, we first determined the distribution of Akk phylogroups in NSCLC

metagenomics datasets of 425 samples from three publicly available cohorts[15,22], and evaluated

the predictive power of different Akk phylogroups for patients' responses to PD-1 blockade.

Furthermore, we performed a rigorous analysis and highlighted a subset of genes in the Akk

pangenome that are enriched in patient responders.

Results

Akk phylogroups across three cohorts of NSCLC patients with ICI treatment

We performed a meta-analysis across three publicly available NSCLC cohorts (*C1*:

Routy et al. Discovery, n = 60; *C2*: *Routy* et al. Validation, n = 27; *C3 Derosa* et al. n = 338) that

used fecal shotgun metagenomics to study the associations between Akk and response to PD-1

blockade[13,20]. All raw shotgun metagenomic sequencing data were processed consistently using

the same bioinformatics pipeline (STAR Methods).

We first re-evaluated the associations between the presence/absence of Akk (Akk$^+$ or

Akk$^-$, determined by MetaPhlAn[33], Methods) and response to PD-1 blockade against NSCLC.

The sequencing yield (Gb) per sample did not have a significant difference between Akk$^+$ and

Akk$^-$ samples across three cohorts (P_{C1} = 0.98, P_{C2} = 0.53, P_{C3} = 0.18, Student's t test, Fig. S1).

Consistent with previous studies, the Akk$^+$ group had higher (compared to the Akk$^-$ group)

proportion of progression-free patients [complete response (CR), partial response (PR), or

stable disease (SD), Fig. S2A-C] and objective response (CR and PR, Supplementary Fig. 2D-

F) in all cohorts. Noteworthy, in the *C2* cohort, all the 5 patients with CR or PR were

characterized as Akk$^+$ group (P = 0.0009, Fig. S2E). We also evaluated the predictive power of

Akk relative abundance, instead of Akk presence/absence, and found significant association

with better response (CR, PR, SD) (P = 0.025) in the *C1* cohort but not in *C2* or *C3*. These

analyses supported the use of the presence/absence of Akk in the following analyses, which is

consistent with a recent study that reported bacteria presence/absence is more robust than

bacterial relative abundance in microbiome-based association studies[34]. Overall, NSCLC

patients carrying Akk (presence vs. absence) had a higher response rate to PD-1 blockade,

consistently across the three cohorts.

To distinguish Akk phylogroups in the meta-analysis, we first built a comprehensive Akk

pangenome including diverse Akk strains (216 Akk isolate genomes from the NCBI database)

that include five known Akk phylogroups (Fig. S3A; STAR Methods). Out of the 231 Akk$^+$

samples across the three cohorts, 24 (out of 30), 8 (out of 12) and 122 (out of 189) samples

were assigned with an Akk phylogroup (Methods), respectively (Fig. 1A-B, Fig. S3B-D, C1: 10

AmIa, 10 AmIb, and 4 AmII; C2: 4 AmIa, 3 AmIb, and 1 AmII; C3: 58 AmIa, 37 AmIb, 21 AmII,

and 6 AmIV). Consistent across the three cohorts, AmIa and AmIb are the two most prevalent

Akk phylogroups, representing typical *A. muciniphila* species that are the most common Akk

species found in the human microbiome[26], followed by AmII and AmIV.

Akk phylogroups have different predictive power for patient response to PD-1 blockade

We compared patients' response rate between NSCLC patients carrying different Akk

phylogroups using a logistic regression model (with covariates; STAR Methods) and found that

AmIa is associated with better response (than other phylogroups) to PD-1 blockade across all

the three NSCLC cohorts (Fig. 2). Specifically, the rates of responders (CR, PR or SD) in the

AmIa group were 80% in *C1* (P_{AmIa} = 0.037, Fig. 2A), 75% in *C2* (P_{AmIa} = 0.075, Fig. 2B), and

58.62% in *C3* (P_{AmIa} = 0.016, Fig. 2C), respectively. We also evaluated the logistic regression

models that included Akk phylogroup vs. Akk presence/absence (Akk$^{+/-}$, not stratifying

phylogroups) to predict objective response (CR or PR) using Akaike information criterion

(AIC)[35], which measures the relative prediction error between regression models (the smaller

value the better model). Consistently across the three cohorts, the logistic regression models

that included Akk phylogroup have better accuracy and AIC than the models that used Akk

species: C1: AIC = 50.538 (Akk phylogroup) vs 52.907 (Akk$^{+/-}$); *C2*: AIC = -16.065 (Akk

phylogroup) vs 20.375 (Akk$^{+/-}$); *C3*: AIC = 299.9 (Akk phylogroup) vs 305.19 (Akk$^{+/-}$). We further

compared patient survival across different Akk phylogroups. In the *C1* cohort, 60% of patients

carrying AmIa had progression-free survival (PFS) longer than 3 months (vs. 43.75% of the

other patients, P_{AmIa} = 0.036, Fig. 2D). In the *C2* cohort, 75% of patients carrying AmIa had PFS

longer than 3 months (vs. 33.33% of AmIb patients, P_{AmIa} = 0.0005, Fig. 2E). Consistently, in the

C3 cohort, AmIa patients had longer overall survival (OS) than patients not carrying AmIa based

on Cox proportional-Hazards regression analysis (P_{AmIa} = 0.0371, Fig. 2F). It is worth noting that

both AmIa and AmIb are prevalent in the three NSCLC cohorts, yet AmIa is associated with

better patients' response and survival than AmIb, consistently across the three cohorts.

Derosa et al. reported in the *C3* cohort that tumor microenvironment differed between

Akk$^+$ patients and Akk$^-$ patients, highlighting several differentially expressed genes[22]. To

evaluate if Akk phylogroups and patients' differential response and survival rate correlate with

patients' tumor microenvironment, we examined the gene expression levels of 28 immune-

related genes between patients with AmIa and AmIb phylogroups. Interestingly, 21 of the 28

genes ($P_{binomial}$ = 0.006) that were expressed higher in Akk$^+$ patients than Akk$^-$ patients showed

greater expression level in patients carrying AmIa compared to those in AmIb, with 5 of them

showing significantly difference (P_{adj} < 0.05, Fig. 2G). The *TDO2* gene, which is the interferon

fingerprint and associated with checkpoint pathway, showed the largest difference between

AmIa and AmIb group (LogFC = 3.79, P_{adj} = 0.008). *CXCL10* (LogFC = 1.95, P_{adj} = 0.049),

related to type II interferon signaling, *VCAM1* (LogFC = 1.94, P_{adj} = 0.015), related to leukocyte

migration, *RB1* (LogFC = 1.69, P_{adj} = 0.033) related to tumor marker, and *CMKLR1* (LogFC =

1.62, P_{adj} = 0.048), related to dendritic cell and macrophage also had higher expression in AmIa

than AmIb. These analyses suggest that the tumor microenvironment genes associated with

Akk$^+$ patients have more pronounced changes in their expression level in the patients carrying

the AmIa phylogroup than the AmIb, consistent with their stronger anti-tumor immune response

to ICI.

Taken together, these analyses suggest that the presence of AmIa phylogroup in

NSCLC patients' gut microbiome is associated with better responses than the other Akk

phylogroups to PD-1 blockage, survival and tumor microenvironment.

Akk gene-based prediction of response to anti-PD1 blockade

Building on the observations above that NSCLC patients carrying AmIa were associated

with better response than the other phylogroups, we further hypothesized that AmIa genomes

may be enriched for certain genes that are positively associated with the response to ICI. It is

also worth noting that not all the patients carrying AmIa responded to PD-1 blockade while

some patients with other Akk phylogroups did respond to PD-1 blockade. This heterogeneity

within a phylogroup is not unexpected considering the genomic differences across strains within

each phylogroup. Therefore, we further performed an Akk gene-level analysis to better

understand the different predictive power between Akk phylogroups for patients' response to

PD-1 blockage.

We built a machine-learning model (model I) with patients' Akk phylogroup, patients'

metadata (covariates: age, gender, antibiotics, sequencing depth, which are available in all the

three cohorts) to predict response (CR/PR/SD or PD/D). We compared the accuracy of model I

with another model (model II) built with the addition of presence or absence of Akk genes.

Considering the sample sizes of the three cohorts, we trained the two models using all the

available Akk$^+$ samples with defined phylogroups in the combined *C1* and *C2* cohort (46 Akk$^+$

samples including 14 additional samples collected at a later time point from the cohorts), and

the *C3* cohort (122 Akk$^+$ samples), respectively. For the combined cohort (*C1* and *C2*), the

leave-one-out cross-validation (LOOCV) accuracy by the model I using random forest had an

area under the receiver operating characteristic (AUROC) value of 0.76 (Fig. 3A), while the

LOOCV of model II had an AUROC of 0.86 (Fig. 3B). For the *C3* cohort, the LOOCV of models I

and II had AUROCs of 0.49 and 0.57 (with respect to the 54.92% responders in *C3*),

respectively (Fig. 3C-D). The improvements in LOOCV accuracy of the training model with the

addition of Akk gene feature (consistent between the combined *C1*/*C2* cohort and the *C3*

cohort) indicate the association between certain Akk genes and patients' response to PD-1

blockade can provide better predictive power than Akk phylogroups itself.

To investigate which genes contribute significantly to improving the predictive power and

their distribution across different phylogroups, we combined the 168 Akk$^+$ patients with defined

Akk phylogroup across the three cohorts and constructed a model (Model III). The training

model with patients' metadata, sequencing depth, and presence and absence matrix of all the

4,576 pan-Akk genes, selected using strict criteria to avoid ambiguous mapping of genes from

other species to Akk (STAR Methods), reach the accuracy of 0.71 (Fig. 4B), which is higher

than the accuracy solely based on Akk phylogroups (55.0%, Fig. 4A). 73 Akk genes (variable

importance score (VIS) > 0.5, present > 1% of at least one Akk phylogroup) showed significantly

different between responders and non-responders ($P_{chi\text{-}square}$ < 0.05, Table S1), with 35 enriched

in the responders (Fig. 4C) and 38 enriched in the non-responders (Fig. 4D), respectively.

Among the 35 Akk genes enriched in the responders, 22 genes had highest prevalence in AmIa,

followed by 7 genes in AmIb, 3 genes in AmIV, 2 gene in AmII, and 1 gene had highest

prevalence in both AmIa and AmII (Fig. 4C). These responder-associated genes include

multiple carbohydrate-active enzymes (CAZy) encoding genes, such as glycosyltransferase

family 1 protein (*CXU07_01840*), alpha-1,2-fucosyltransferase (*Amuc_2087*), and glycosyl

transferase group 1 (*Amuc_2082*). The distribution of these CAZy encoding genes was much

higher in AmIa followed by AmIb phylogroup and rarely identified in AmII or AmIV phylogroups.

For the 38 Akk genes enriched in non-responders over responders, 18 genes had highest

prevalence in AmIb, followed by 16 genes in AmII and 4 genes in AmIa (Fig. 4D). These genes

encode several bacteriophage related proteins, such as bacteriophage tail assembly protein-like

protein (*Amuc_1355*) and phage major capsid protein which had the highest prevalence in AmII,

as well as phage portal protein, which was only detected in AmIb.

Besides, nonresponder-associated Akk genes also include certain CAZy encoding gene,

such as glycosyltransferase 2-like domain-containing protein (CXU19_11355), which is

frequently detected in AmII but not in other Akk phylogroups. These results suggest a link

between carbohydrate utilization of gut *Akkermansia* and response to immunotherapy of hosts,

likely reflecting variation in intestinal mucosal environments, particularly glycoside diversity

between responders and nonresponders, which probably results from their distinct immune and

metabolic status, that favor growth of different Akk phylogroups[36,37]. The patients dominated by

AmI phylogroup may also benefit from bacterial glycoside fermentation end products, such as

short-chain fatty acids, that enhance their response to PD-1 blockade[38,39]. AmIa and AmIb

containing the most and second most genes enriched in responders is consistent with the

observation that showed patients with AmIa and AmIb are more likely (than the other

phylogroups) to be responders. In contrast, AmIb and AmII harboring more genes enriched in

non-responders, and particularly AmII had few genes enriched in responders, which is

consistent with the observation that patients with AmII are more likely to be non-responders.

Several additional genes that had higher prevalence in responders also showed varied

distribution across Akk phylogroups. For example, Amuc_1677, a gene enriched in responders,

was reported to encode an enzyme involved in the (S)-acetolacetate to 2-methyl-2-oxovaleric

acid step in the branched-chain amino acids to phosphatidylethanolamine pathway[27]. The

frequency of Amuc_1677 across the Akk phylogroups is 51.85% in AmIa, 17.24% in AmIb,

absent in AmII and AmIV. Among the patients with either AmIa or AmIb, those carrying Akk with

an Amuc_1677 gene had a higher chance to be responders (Fig. 4E). Another gene enriched in

responders, Amuc_2164 (glycoside hydrolase family 18, a predicted mucin binding chitinase) is

only in strains from the AmI phylogroup (83% in AmIa, 99.14% in AmIb) but absent in the other

phylogroup. These data indicate AmI strains, encode certain Akk genes that are associated with

better response to PD-1 blockage, while AmII strains lack beneficial genes or might harbor

certain genes that are negatively associated with patients' response to PD-1 blockade. It is

worth noting these genes represent association but not causality, and their functionality in the

context of responses to PD-1 blockage needs to be further explored in the future studies.

To summarize, we found that the model that incorporate Akk genes has enhanced

performance in the prediction of patients' response to PD-1 blockade. Importantly, the gene-

level association analysis (with patient response and with Akk phylogroups) helped elucidate the

difference in the predictive power of different Akk phylogroups in clinical response to PD-1

blockade against NSCLC.

Discussion

In this study, we assessed the difference between Akk phylogroups with respect to their

predictive capacity in responses to PD-1 blockade in NSCLC patients. Through the delineation

of Akk phylogroups and an extensive meta-analysis across three previously published cohorts,

we unveiled diverse Akk phylogroups among NSCLC's patients, highlighting the better power of

the AmIa phylogroup, consistently across three independent cohorts, in predicting responses to

immunotherapy compared to the other phylogroups. Also consistently, we observed stronger

association between AmIa and patient survival and the RNA expression of patients' genes

relevant to tumor microenvironment. Using a machine learning model predicting patients'

response based on Akk gene profiles, we searched for the genes associated with AmIa's

superior predictive capabilities, and highlighted a number of genes predominantly prevalent in

the AmIa phylogroup and associated with patients' response to PD-1 blockade. Some of these

genes are also present in other phylogroups, e.g. AmIb, but less prevalent compared to AmIa.

This is consistent with the observation that some patients carrying AmIb are also responders. In

addition, we found a few genes that are predominantly prevalent in non-responders and AmII,

which are consistent with the worse response of patients carrying AmII. These revelations

underscore the importance of discerning Akk phylogroups and Akk genes in disease association

analysis, which may help the development of innovative microbiome-centric diagnostic tools and

personalized medicine approaches for PD-1 blockage in NSCLC patients.

Our integration and analysis of the three cohorts also emphasized some limitations and

the need for further validations through larger, independent studies, considering the complexity

of NSCLC and patients' response to PD-1 blockage. First, although our analytical models used

three co-variants shared across the three cohorts (age, gender, antibiotics), we think the

collection and incorporation of additional covariates is paramount in these analyses, given their

potential to introduce confounding effects. For instance, cohorts C1 and C2 predominantly

comprised patients who underwent chemotherapy before PD-1 blockade, whereas cohort C3

had a higher proportion of patients treated with radiotherapy before PD-1 blockade. These

variations might explain the relatively better predictive power of AmIa in C1 and C2 than in C3,

even though the three cohorts consistently highlight the importance of AmIa. Second, while this

study was focused on the microbiome, we also recognize the substantial importance of host

factors such as ECOG performance status, PD-L1 expression, tumor gene expression data, in

predicting patients' response to PD-1 blockage[22]. Although ECOG performance status, PD-L1

expression, and tumor gene expression data are available in the C3 cohort, they were not part

of the other two cohorts. Given the complexity of NSCLC and patients' response to PD-1

blockage, future development in this field necessitates studies with a holistic approach,

integrating both microbiome and host gene expression data to decipher their relationships and

their joint predictive power[23,40,41]. Besides, due to the limitation of annotated Akk gene based on

our current knowledge, the functions of response-associated sequence clusters need to be

further explored.

While Akk has persistently demonstrated a link to responses in NSCLC patients

undergoing PD-1 blockade across varied cohorts, several other bacterial species like

Eubacterium hallii, *Bifidobacterium adolescentis*, and *Parabacteroides distasonis* have also

been reported, albeit with less consistency across cohorts[22,42,43]. This discrepancy might arise

from the possibly more subtle yet legitimate associations these bacteria have with PD-1

blockade responses or due to the inherent patient heterogeneity observed across diverse

cohorts. Indeed, the complexity of responses to PD-1 blockade in NSCLC is multifaceted,

interweaving numerous factors from both the human host and the gut microbiome. While Akk is

a significant player in inducing intestinal adaptive immune responses during homeostasis, it's

plausible that other bacterial species also play integral roles in this biological puzzle. With this

rationale, we made an attempt to assess the predictive capabilities of additional bacterial

species, probing their association with responses to PD-1 blockade (Supplementary Text).

Monoglobus pectinilyticus exhibited a consistent positive association with responders across all

the three NSCLC cohorts (Supplementary Text and Fig. S4), although future work is imperative

to validate this finding on large and independent cohorts as well as the complementarity

between Akk and additional bacterial species.

Beyond its association with responses to PD-1 blockade, Akk has also been implicated

in a spectrum of other diseases, including obesity, type 2 diabetes, and inflammatory bowel

disease, among others[44,45,46]. The insights from this study also suggest the use of similar

analytical strategies to investigate the predictive capabilities of distinct Akk phylogroups in

relation to these other diseases and their respective treatment outcomes. Given the recognition

of Akk as a promising candidate in the development of next-generation probiotics, it is important

to attain a nuanced understanding of the various Akk phylogroups, their specific genes, and

genomic variations[47] across a range of diseases, which could unravel transformative insights

and facilitate the advancement of innovative therapeutic interventions and proactive preventive

measures.

Figure legend

Figure 1. NSCLC patients carrying different phylogroup of Akk varied in response to ICI treatment. (A) The Akk phylogroups carried by NSCLC patients from three cohorts. **(B)** The AmIa and AmIb phylogroup carried by NSCLC patients from three cohorts. The three cohorts are *Routy* Discovery cohort (n = 60), *Routy* Validation cohort (n = 27), and *Derosa* cohort (n = 338, *P* = 0.016).

Figure 2. **NSCLC patients carrying different phylogroup of Akk varied in response to ICI treatment. (A to C)** Correlations between phylogroup of Akk in the stool of NSCLC patients and response to PD-1 blockade. AmIa group had higher proportion of patients being CR, PR, or SD in *Routy* Discovery cohort (n = 60, *P* = 0.037) (**A**), *Routy* Validation cohort (n = 27, *P* = 0.075) (**B**), and *Derosa* cohort (n = 338, *P* = 0.016) (**C**). In *Routy* Validation cohort, AmIa group showed significantly higher proportion of patients being CR or PR ($P = 9.04 \times 10^{-8}$). CR; complete response. PR; partial response. SD; stable disease. PD; progressive disease. D; death. Data were analyzed using generalized linear model. (**D**-**F**) Correlations between phylogroup of Akk in stool of NSCLC patients and survival length. AmIa group had higher proportion of patients which had progression-free survival (PFS) longer than 3 months in *Routy* Discovery cohort (n = 60, *P* = 0.036) (**D**) and *Routy* Validation cohort (n = 27, *P* = 0.0005) (**E**), and had longer overall survival in *Derosa* cohort (**F**). Data were analyzed using generalized linear model. **G**, Bar graph showing most immune-related genes that were expressed at higher level in lung tumor of patients carrying Akk had higher expression level in AmIa group. Differential expression analysis was conducted using voom method. Akk-/un represents for the samples that are either

detected as Akk negative using MetaPhlAn 3 or detected as Akk positive in MetaPhIAN 3 but

not in PanPhlan*, $P < 0.05$; #, $0.05 < P < 0.1$.

Figure 3. Akk gene-based model improved performance to predict response to PD-1

blockade. (A) The area under the receiver operating characteristic (AUROC) of the random

forest training model to predict response using Akk phylogroup, patients' metadata, and

sequencing depth in the combined C1 and C2 cohort (46 Akk$^+$ samples). **(B)** The AUROC of the

random forest training model to predict response with Akk gene presence or absence

information together with patients' metadata, and sequencing depth in the combined C1 and C2

cohort (46 Akk$^+$ samples). **(C)** The AUROC of the random forest training model to predict

response using Akk phylogroup, patients' metadata, and sequencing depth in the C3 cohort

(122 Akk$^+$ samples). **(D)** The AUROC of the random forest training model to predict response

with Akk gene presence or absence information together with patients' metadata, and

sequencing depth in the C3 cohort (122 Akk$^+$ samples).

Figure 4. Potential Akk genes that are associated with response to PD-1 blockade.

(A) The AUROC of the random forest training model to predict response using Akk phylogroup,

patients' metadata, and sequencing depth in the combined C1, C2, and C3 cohort (168 Akk$^+$

samples). **(B)** The AUROC of the random forest training model to predict response with

presence or absence information of Akk genes together with patients' metadata, and

sequencing depth in the combined C1, C2, and C3 cohort (168 Akk$^+$ samples). (**C-D)** The

heatmaps showing the distribution of response-associated Akk genes across Akk phylogroups.

Only Akk genes with prevalence higher than 1% in at least one phylogroup were included in the

heatmaps **(C)** The distribution of responder-associated Akk genes ($P_{Chi-square} < 0.05$) with

variable importance score (VIC) higher than 0.5 in the random forest model across different Akk

phylogroups. Only Akk genes present in more than 1% of at least 1 phylogroup were included in

the heatmap. **(D)** The distribution of nonresponder-associated Akk genes ($P_{\text{Chi-square}} < 0.05$) with

VIC higher than 0.5 in the random forest model across different Akk phylogroups. Only Akk

genes present in more than 1% of at least 1 phylogroup were included in the heatmap. **(e)** The

bargraph showing the prevalence of responders and nonresponders of the patients in AmIa or

AmIb that carry or not carry Amuc_1677 gene.

STAR Methods

Public populations of NSCLC patients

Raw gut metagenomic sequencing data of three populations (C1: *Routy* et al. Discovery, n = 60;

C2: *Routy* et al. Validation, n = 27; C3: *Derosa* et al. n = 338) were downloaded from the

Sequence Read Archive (SRA) under the BioProject accession PRJEB22863

(https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJEB22863) and PRJNA751792

(https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA751792), mainly from two recently

published papers[15,22]. The raw tumor RNA sequencing data were downloaded from the SRA

under accession the NCBI accession GSE182328

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE182328) from *Derosa* et al. (2022)[22].

Identification of Akk presence and Akk phylogroup in the stool metagenomes

The KneadData tool (v0.10.0) was used to perform quality control on raw metagenomic

sequencing data. Briefly, adapters and low-quality reads were removed using Trimmomatic

(v0.39), and repetitive sequences were trimmed using the TRF (v4.09.1). The human reads

were detected by mapping remaining reads to the human genome (hg37) by bowtie2 (v.2.2.5),

and removed. *Akkermansia muciniphila* presence and relative abundances of bacterial taxa

were determined and estimated by MetaPhlAn 3 (v3.0) with ChocoPhlAn 3 database[32].

PanPhlan pipeline (v3.0) was applied to profile *A. muciniphila* strains within stool metagenomes.

A total of 216 available *A. muciniphila* isolate genomes (fasta file) were download from NCBI

and were used to build a comprehensive pangenome with UniRef90 ID using PanPhlan

pangenome exporter (https://github.com/SegataLab/PanPhlAn_pangenome_exporter). Due to

high genetic diversity within *A. muciniphila* strains, this species has been suggested to be

classified into four species-level phylogroups, and we considered the UniRef genes families-

level pangenome as *Akkermansia* (Akk) pangenome. Then each metagenomic sample was

mapped against the generated Akk custom pangenome (panphlan_map.py). The coverages of

all gene positions were detected and extracted using samtools (v1.16), and then integrated to a

gene family coverage profile for each sample. Pangenome-related Akk strains were predicted to

exist if coverage curves display a satisfying plateau using the least stringent threshold of

panphlan_profile.py (--min_coverage 1, --left_max 1.70, --right_min 0.30). Principal coordinates

analysis (PCoA) of Jaccard distance was performed on UniRef90 ID presence and absence

binary matrix of metagenomic samples using ggplot2 package in the R (v4.3) to distinguish

phylogroup of Akk strains according to cluster with references.

Evaluation of correlations between Akk presence/absence, Akk phylogroup and

response to ICI treatment

Generalized linear model was applied to evaluate the effects of Akk presence/absence or

specific Akk phylogroup on the response. The patients were classified as responders (complete

response (CR), partial response (PR), and stable disease (SD)) and non-responders

(progressive disease (PD) and death (D)) according to the previous study[15]. Akk presence or

Akk phylogroup with sequencing depth (number of raw reads) and patients' metadata (age,

gender, time point, ECOG performance status, PD-L1 expression if available) were included as

independent variables. Responders, objective response (CR and PR), or survival lengths were

served as dependent variables. Akaike information criterion (AIC) and accuracy were used to

compare the quality of models.

Tumor RNA sequencing data analysis

To investigate relationships between the expression of immune-related genes in tumors and

Akk phylogroup the patients were carrying, the differences in Akk-associated immune-related

genes reported by previous study was analyzed using the published tumor RNA sequencing

data[14]. The low-quality reads and adapter were removed from raw RNA sequencing data using

Trim Galore (v0.6.6). The remaining reads were mapped to human GRCh38 cDNA reference

transcriptome using STAR (v2.7.2a). The number of reads mapping to each gene was counted

using featureCounts (v2.0.1). The differentially expressed genes (DEGs) between patients

carrying AmIa (n = 3) and AmIb (n = 5) were analyzed in R using voom package (v3.38.14).

Construction and evaluation of the machine-learning model based on Akk genes profile

To determine whether Akk gene profiles could enhance the prediction of response to PD-1

blockade, the random forest models were built using R caret package (v 6.0-94). The patients

from the three cohorts with Akk phylogroup assigned were included in this analysis. Additionally,

to increase the sample size, an additional 14 fecal metagenomic samples from the C1 cohort

collected after the 2[nd] (T1- 1 month), 4[th] (T2- 2 months) and 12[th] (T3- 6 months) injection of anti-

PD-1/PD-L1 mAb and with Akk phylogroup assigned were included[22].

The training models were constructed with leave one out cross validation to distinguish

responders and non-responders. The features used for model building consist of sequencing

depth (number of raw reads), patient metadata (age, gender, usage of antibiotics), and Akk

profile (phylogroup, presence and absence of Akk only genes). The UniRef genes that only exist

in *Akkermansia* were considered as Akk only genes. Briefly, the metadata for the full UniRef90

gene set was retrieved from the UniRef website (http://www.uniprot.org/uniref), which includes

the organism/species information. All the Akk-related genes was extracted by using the

Organism tag ("*Akkermansia*"). These Akk genes were classified into different categories based

on the number of homologous. The genes that only existed in Akk were pulled out as Akk only

genes. The models were optimized by tuning the mtry (number of randomly selected features to

be sampled at each iteration) and the ntree (number of decision tree). The area under the

receiver operating characteristic (AUROC) and accuracy were used to evaluate the

performance of the training models. The variable importance scores were extracted for each

feature in the models, and distribution of critical Akk genes with high score (> 0.5) with

significant difference present in responders and non-responders ($P_{chi\text{-}square} < 0.05$) were

determined across phylogroups.

References

1. Centers for Disease Control and Prevention. Lung Cancer Statistics.

https://www.cdc.gov/cancer/lung/statistics/index.htm#:~:text=Lung%20cancer%20is%20the%20th

ird,any%20other%20type%20of%20cancer. 2023.

2. Centers for Disease Control and Prevention. An Update on Cancer Deaths in the United States.

https://www.cdc.gov/cancer/dcpc/research/update-on-cancer-deaths/index.htm.2020.

3. Zhou F, Qiao M, Zhou C. The cutting-edge progress of immune-checkpoint blockade in lung cancer.

Cell. Mol. Immunol. 2021; **18**: 279-293.

4. Chiang AC, Herbst RS. Frontline immunotherapy for NSCLC—the tale of the tail. *Nat. Rev. Clin. Oncol.*

2020;**17**: 73-74.

5. MacManus M, Hegi-Johnson F. Overcoming immunotherapy resistance in NSCLC. *Lancet Oncol.*

2022; **23**: 191-193.

6. Doroshow DB, Bhalla S, Beasley MB, Sholl LM, Kerr KM, Gnjatic S, Wistuba II, Rimm DL, Tsao MS,

Hirsch FR. PD-L1 as a biomarker of response to immune-checkpoint inhibitors. *Nat. Rev. Clin.*

Oncol. 2021; **18**: 345-362.

7. Kambayashi Y, Fujimura T, Hidaka T, Aiba S. Biomarkers for predicting efficacies of anti-PD1

antibodies. *Front. Med. (Lausanne).* 2019*,* **6**: 174.

8. Liu Y, Altreuter J, Bodapati S, Cristea S, Wong CJ, Wu CJ, Michor F. Predicting patient outcomes after

treatment with immune checkpoint blockade: A review of biomarkers derived from diverse data

modalities. *Cell Genom.* 2024; **4**: 100444.

9. Lu S, Stein JE, Rimm DL, Wang DW, Bell JM, Johnson DB, Sosman JA, Schalper KA, Anders RA,

Wang H et al. Comparison of Biomarker Modalities for Predicting Response to PD-1/PD-L1

Checkpoint Blockade: A Systematic Review and Meta-analysis. *JAMA Oncol.* 2019; **5**: 1195-

1204.

10. Ren D, Hua Y, Yu B, Ye X, He Z, Li C, Wang J, Mo Y, Wei X, Chen Y et al. Predictive biomarkers and

mechanisms underlying resistance to PD1/PD-L1 blockade cancer immunotherapy. *Mol. Cancer.*

2020; **19**: 31.

11. Masuhiro K, Tamiya M, Fujimoto K, Koyama S, Naito Y, Osa A, Hirai T, Suzuki H, Okamoto N,

Shiroyama T et al. Bronchoalveolar lavage fluid reveals factors contributing to the efficacy of PD-

1 blockade in lung cancer. JCI Insight. 2022; **7**: e157915.

12. Zhou CB, Zhou YL, Fang JY. Gut microbiota in cancer immune response and immunotherapy. *Trends*

Cancer. 2021; **7**: 647-660 (2021).

13. Simpson RC, Shanahan ER, Batten M, Reijers ILM, Read M, Silva IP, Versluis JM, Ribeiro R,

Angelatos AS, Tan J et al. Diet-driven microbial ecology underpins associations between cancer

immunotherapy outcomes and the gut microbiome. *Nat. Med.* 2022; **28**: 2344-2352.

14. Iida N, Dzutsev A, Stewart CA, Smith L, Bouladoux N, Weingarten RA, Molina DA, Salcedo R, Back

T, Cramer S et al. Commensal bacteria control cancer response to therapy by modulating the

tumor microenvironment. *Science* 2013; **342**: 967-970.

15. Routy B, Le Chatelier E, Derosa L, Duong CPM, Alou MT, Daillère R, Fluckiger A, Messaoudene M,

498   Rauber C, Roberti MP et al. Gut microbiome influences efficacy of PD-1-based immunotherapy

499   against epithelial tumors. *Science.* 2018; **359**: 91-97.

500   16. Vernocchi P, Gili T, Conte F, Del Chierico F, Conta G, Miccheli A, Botticelli A, Paci P, Caldarelli G,

501   Nuti M et al. Network analysis of gut microbiome and metabolome to discover microbiota-linked

502   biomarkers in patients affected by non-small cell lung cancer. *Int. J. Mol. Sci.* 2020; **21**: 8730.

503   17. Deluce J, Maleki Vareki S, Fernandes R. The role of gut microbiome in immune modulation in

504   metastatic renal cell carcinoma. *Ther. Adv. Med. Oncol.* 2022; **14**: 17588359221122714.

505   18. Gopalakrishnan V, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpinets TV, Prieto PA, Vicente D,

506   Hoffman K, Wei SC et al. Gut microbiome modulates response to anti–PD-1 immunotherapy in

507   melanoma patients. *Science.* 2018; **359**: 97-103.

508   19. Routy B, Lenehan JG, Miller WH Jr, Jamal R, Messaoudene M, Daisley BA, Hes C, Al KF, Martinez-

509   Gili L, Punčochář M et al. Fecal microbiota transplantation plus anti-PD-1 immunotherapy in

510   advanced melanoma: a phase I trial. *Nat. Med.* 2023; **29**: 2121-2132.

511   20. Wang Y, Wiesnoski DH, Helmink BA, Gopalakrishnan V, Choi K, DuPont HL, Jiang ZD, Abu-Sbeih H,

512   Sanchez CA, Chang CC et al. Fecal microbiota transplantation for refractory immune checkpoint

513   inhibitor-associated colitis. *Nat. Med.* 2018; **24**: 1804-1808.

514   21. Zheng Y, Wang T, Tu X, Huang Y, Zhang H, Tan D, Jiang W, Cai S, Zhao P, Song R et al. Gut

515   microbiome affects the response to anti-PD-1 immunotherapy in patients with hepatocellular

516   carcinoma. *J. Immunother. Cancer.* 2019; **7**: 193.

517   22. Derosa L, Routy B, Thomas AM, Iebba V, Zalcman G, Friard S, Mazieres J, Audigier-Valette C, Moro-

518   Sibilot D, Goldwasser F et al. Intestinal Akkermansia muciniphila predicts clinical response to PD-

519   1 blockade in patients with advanced non-small-cell lung cancer. *Nat. Med.* 2022; **28**: 315-324.

520   23. Ansaldo E, Slayden LC, Ching KL, Koch MA, Wolf NK, Plichta DR, Brown EM, Graham DB, Xavier

521   RJ, Moon JJ et al. *Akkermansia muciniphila* induces intestinal adaptive immune responses during

522   homeostasis. *Science.* 2019; **364**: 1179-1184.

523   24. Yoon HS, Cho CH, Yun MS, Jang SJ, You HJ, Kim JH, Han D, Cha KH, Moon SH, Lee K, et al.

524   *Akkermansia muciniphila* secretes a glucagon-like peptide-1-inducing protein that improves

glucose homeostasis and ameliorates metabolic disease in mice. *Nat. Microbiol.* 2021; **6**: 563-

573.

25. Kim SM, Park S, Hwang SH, Lee EY, Kim JH, Lee GS, Lee G, Chang DH, Lee JG, Hwang J et al.

Secreted *Akkermansia muciniphila* threonyl-tRNA synthetase functions to monitor and modulate

immune homeostasis. *Cell Host Microbe.* 2023; **31**: 1021-1037.

26. Jian H, Liu Y, Wang X, Dong X, Zou X. *Akkermansia muciniphila* as a next-generation probiotic in

modulating human metabolic homeostasis and disease progression: a role mediated by gut–

liver–brain axes? *Int. J. Mol. Sci.* 2023; **24**, 3900.

27. Bae M, Cassilly CD, Liu X, Park SM, Tusi BK, Chen X, Kwon J, Filipčík P, Bolze AS, Liu Z et al.

Akkermansia muciniphila phospholipid induces homeostatic immune responses. *Nature.* 2022;

608: 168-173.

28. Guo X, Li S, Zhang J, Wu F, Li X, Wu D, Zhang M, Ou Z, Jie Z, Yan Q et al. Genome sequencing of

39 *Akkermansia muciniphila* isolates reveals its population structure, genomic and functional

diversity, and global distribution in mammalian gut microbiotas. *BMC Genomics. 2017*; **18**: 1-2.

29. Becken B, Davey L, Middleton DR, Mueller KD, Sharma A, Holmes ZC, Dallow E, Remick B, Barton

GM, David LA et al. Genotypic and phenotypic diversity among human isolates of *Akkermansia*

muciniphila. *Mbio.* 2021; **12**: e00478-21.

30. Karcher N, Nigro E, Punčochář M, Blanco-Míguez A, Ciciani M, Manghi P, Zolfo M, Cumbo F, Manara

S, Golzato D et al. Genomic diversity and ecology of human-associated *Akkermansia* species in

the gut microbiome revealed by extensive metagenomic assembly. *Genome Biol.* 2021; **22**: 209.

31. Mueller KD, Panzetta ME, Davey L, McCann JR, Rawls JF, Flores GE, Valdivia RH. Pangenomic

analysis identifies correlations between *Akkermansia* species and subspecies and human health

outcomes. *Microbiome Research Reports.* 2024; **3**: 33.

32. Xing J, Li X, Sun Y, Zhao J, Miao S, Xiong Q, Zhang Y, Zhang G. Comparative genomic and

functional analysis of *Akkermansia muciniphila* and closely related species. *Genes Genomics.*

2019; **41**, 1253-1264.

33. F. Beghini, L.J. McIver, A. Blanco-Míguez, L. Dubois, F. Asnicar, S. Maharjan, A. Mailyan, P. Manghi,

M. Scholz, A.M. Thomas, M. Valles-Colomer, G. Weingart, Y. Zhang, M. Zolfo, C. Huttenhower,

E.A. Franzosa, N. Segata. Integrating taxonomic, functional, and strain-level profiling of diverse

microbial communities with bioBakery 3. *Elife* **10**, e65088 (2021).

34. Beghini F, McIver LJ, Blanco-Míguez A, Dubois L, Asnicar F, Maharjan S, Mailyan A, Manghi P,

Scholz M, Thomas AM et al. Host phenotype classification from human microbiome data is mainly

driven by the presence of microbial taxa. *PLOS Comput. Biol.* 2022; **18**: e1010066.

35. Palmqvist S, Tideman P, Cullen N, Zetterberg H, Blennow K; Alzheimer's Disease Neuroimaging

Initiative; Dage JL, Stomrud E, Janelidze S, Mattsson-Carlgren N et al. Prediction of future

Alzheimer's disease dementia using plasma phospho-tau combined with other accessible

measures. *Nat. Med.* 2021; **27**: 1034-1042.

36. Kudelka MR, Stowell SR, Cummings RD, Neish AS. Intestinal epithelial glycosylation in homeostasis

and gut microbiota interactions in IBD. *Nat. Rev. Gastroenterol. Hepatol.* 2020; **17**: 597-617.

37. Moran AP, Gupta A, Joshi L. Sweet-talk: role of host glycosylation in bacterial pathogenesis of the

gastrointestinal tract. *Gut.* 2011; **60**: 1412-1425.

38. Bergstrom K, Xia L. The barrier and beyond: Roles of intestinal mucus and mucin-type O-

glycosylation in resistance and tolerance defense strategies guiding host-microbe symbiosis. *Gut*

Microbes. 2022; **14**: 2052699.

39. Shuoker B, Pichler MJ, Jin C, Sakanaka H, Wu H, Gascueña AM, Liu J, Nielsen TS, Holgersson J,

Nordberg Karlsson E et al. Sialidases and fucosidases of *Akkermansia muciniphila* are crucial for

growth on mucin and nutrient sharing with mucus-associated gut bacteria. *Nat. Commun.* 2023;

14: 1833.

40. Fidelle M, Rauber C, Alves Costa Silva C, Tian AL, Lahmar I, de La Varende AM, Zhao L,

Thelemaque C, Lebhar I, Messaoudene M et al. A microbiota-modulated checkpoint directs

immunosuppressive intestinal T cells into cancers. *Science.* 2023; **380**: eabo2296.

41. Park JS, Gazzaniga FS, Wu M, Luthens AK, Gillis J, Zheng W, LaFleur MW, Johnson SB, Morad G,

Park EM et al. Targeting PD-L2-RGMb overcomes microbiome-related immunotherapy

resistance. *Nature.* 2023; **617**: 377-385.

42. Sivan A, Corrales L, Hubert N, Williams JB, Aquino-Michaels K, Earley ZM, Benyamin FW, Lei YM,
Jabri B, Alegre ML et al. Commensal Bifidobacterium promotes antitumor immunity and facilitates
anti-PD-L1 efficacy. *Science.* 2015; **350**: 1084-1089.

43. Mao J, Wang D, Long J, Yang X, Lin J, Song Y, Xie F, Xun Z, Wang Y, Wang Y et al. Gut microbiome
is associated with the clinical response to anti-PD-1 based immunotherapy in hepatobiliary
cancers. *J. Immunother. Cancer.* 2021; **9**: e003334.

44. Depommier C, Everard A, Druart C, Plovier H, Van Hul M, Vieira-Silva S, Falony G, Raes J, Maiter D,
Delzenne NM et al. Supplementation with *Akkermansia muciniphila* in overweight and obese
human volunteers: a proof-of-concept exploratory study. *Nat. Med.* 2019; **25**: 1096-1103.

45. Forslund K, Hildebrand F, Nielsen T, Falony G, Le Chatelier E, Sunagawa S, Prifti E, Vieira-Silva S,
Gudmundsdottir V, Pedersen HK et al. Disentangling type 2 diabetes and metformin treatment
signatures in the human gut microbiota. *Nature.* 2015; **528**: 262-266.

46. Zhang T, Ji X, Lu G, Zhang F. The potential of *Akkermansia muciniphila* in inflammatory bowel
disease. *Appl. Microbiol. Biotechnol.* 2021; **105**, 5785-5794.

47. Liu R, Zou Y, Wang WQ, Chen JH, Zhang L, Feng J, Yin JY, Mao XY, Li Q, Luo ZY et al. Gut
microbial structural variation associates with immune checkpoint inhibitor response. *Nat.*
Commun. 2023; **14**: 7421.

Acknowledgements

We thank Edward Mead, Yanchun Zhang and other members in the Fang Lab for their helpful

feedback. This work was supported by R35 GM139655 (G.F.) from the National Institutes of

Health. G.F. is a Hirschl Research Scholar by Irma T. Hirschl/Monique Weill-Caulier Trust, and

a Nash Family Research Scholar. This work was also supported in part through the

computational resources and staff expertise provided by the Department of Scientific Computing

at the Icahn School of Medicine at Mount Sinai.

Author Contributions

P.F. performed all the data analyses. P.F. and G.F. designed the meta-analyses across the three cohorts. M.N., Y.F. and M.K. assisted in the Akk pangenome analysis. G.F. conceived and supervised the project. P.F. and G.F. wrote the manuscript with input and comments from all co-authors.

Competing Interests

The authors declare no competing interests.

Figure 1

Figure 2

Figure 3

Figure 4