Distinct checkpoint and homolog biorientation pathways

regulate meiosis I in *Drosophila* oocytes

Joanatta G. Shapiro, Neha Changela, Janet K. Jang, Jay N.

Joshi, and Kim S. McKim

Waksman Institute and Department of Genetics, Rutgers, the State University of
New Jersey, Piscataway, New Jersey, United States of America.
8

Condensed title: Checkpoint and biorientation pathways in oocytes

Correspondence: Kim S. McKim (mckim@waksman.rutgers.edu)

Corresponding author: Kim S. McKim
190 Frelinghuysen Road,
Piscataway, NJ-08854
848-445-1164
mckim@waksman.rutgers.edu
17

Abstract

Mitosis and meiosis have two mechanisms for regulating the accuracy of

chromosome segregation: error correction and the spindle assembly checkpoint (SAC).

We have investigated the function of several checkpoint proteins in meiosis I of

Drosophila oocytes. Evidence of a SAC response by several of these proteins is found

upon depolymerization of microtubules by colchicine. However, unattached

kinetochores or errors in biorientation of homologous chromosomes does not induce a

SAC response. Furthermore, the metaphase I arrest does not depend on SAC genes,

suggesting the APC is inhibited even if the SAC is silenced. Two SAC proteins, ROD of

the ROD-ZW10-Zwilch (RZZ) complex and MPS1, are also required for the biorientation

of homologous chromosomes during meiosis I, suggesting an error correction function.

Both proteins aid in preventing or correcting erroneous attachments and depend on

SPC105R for localization to the kinetochore. We have defined a region of SPC105R,

amino acids 123-473, that is required for ROD localization and biorientation of

homologous chromosomes at meiosis I. Surprisingly, ROD removal, or "streaming", is

independent of the dynein adaptor Spindly and is not linked to the stabilization of end-

on attachments. Instead, meiotic RZZ streaming appears to depend on cell cycle stage

and may be regulated independently of kinetochore attachment or biorientation status.

We also show that dynein adaptor Spindly is also required for biorientation at meiosis I,

and surprisingly, the direction of RZZ streaming.

Author Summary

The Spindle Assembly Checkpoint (SAC) is known to delay cell cycle

progression until chromosomes are properly attached to microtubules. Meiotic cells

often have modified cell cycle phases, and natural arrest points such as metaphase I in

Drosophila. We show that in *Drosophila* oocytes, the SAC is sensitive to loss of

microtubules, but not sensitive to a variety of kinetochore attachment errors. Thus, the

function of the SAC appears to be limited to monitoring oocyte spindle assembly, and

not required for accurate chromosome segregation. However, two of the SAC genes,

rod and *Mps1*, are required for the biorientation of homologous chromosomes during

meiosis I, suggesting an error correction function. Rod is part of the RZZ complex and is

notable for its property of streaming off the kinetochores. However, our results show

that streaming off the kinetochore may not contribute to RZZ regulation of microtubule

attachments, and only be associated with SAC function. Instead, the establishment of

stable end-on attachments may occur while RZZ is still present at kinetochore. We

suggest that RZZ interacts with multiple motors to promote bidirectional movement of

kinetochores along microtubules, which allows chromosomes to find and attach to the

correct pole.

Introduction

Errors during meiotic chromosome segregation can result in inviable gametes

leading to spontaneous abortions and infertility, or viable offspring with chromosomal

abnormalities. Several mechanisms exist in dividing cells to prevent aneuploidy

resulting from improper kinetochore-microtubule (KT-MT) attachments. The spindle

assembly checkpoint (SAC) delays anaphase onset until attachments form between

kinetochores and microtubules [1, 2]. Unattached kinetochores generate a signal

involving SAC proteins such as MAD1, MAD2, BUB1, and BUB3 that have a global

effect on all chromosomes and cell cycle progression. In contrast, error correction (EC)

is specific to individual kinetochores whereupon interactions between microtubules and

properly bi-oriented kinetochores are stabilized while those with improperly oriented

kinetochores are weakened [3]. Multiple models have been proposed to explain how

improperly oriented kinetochores are identified, involving both tension-dependent and -

independent mechanisms, and how they are corrected, including phosphorylation of

kinetochore proteins to destabilize attachments [4, 5].

Error rates are higher in human oocytes than in spermatocytes, which has been

attributed to several causes [6-8]. One hypothesis is that the SAC is weak or inefficient

[7, 9-11]. Indeed, Xenopus oocytes may not have a functional SAC [12, 13]. The

observation that mammalian oocytes progress through anaphase with segregation

errors may be a result of a weak checkpoint [14, 15]. The reason for a weak SAC in

oocytes is not known, although possible contributing factors include the large size of

oocytes or the lack of centrosomes [9, 16-19]. The SAC may also have other functions

in oocytes, such as regulating the development of the oocytes and the timing of meiotic

progression [10].[1, 2].

Drosophila oocytes are an excellent model for studying meiosis in females, but

the SAC has not been examined. *Drosophila* oocytes have kinetochores that include the

KMN complex (KNL1-MIS12-NDC80) [20]. Biorientation of homologous chromosomes

involves types of kinetochore-microtubule attachments, lateral attachments that depend

on KNL1/SPC105R and end-on attachments that depend on NDC80 [21, 22]. To

understand the contributions of SAC and EC to biorientation in *Drosophila* oocytes, we

characterized the SAC response during the first meiotic division, investigated how the

SAC proteins are recruited to the kinetochores, and tested several SAC proteins for

functions in meiotic EC. Our data suggest that the SAC is not sensitive to kinetochore

attachment status and might be silenced prior to the metaphase I arrest. Furthermore,

among seven SAC proteins, only two, ROD of the ROD-ZW10-Zwilch (RZZ) complex

and MPS1, are required for biorientation of meiotic homologous chromosomes at

metaphase I.

We examined the role of RZZ in *Drosophila* oocytes because it has not been

studied before in meiosis. Previous studies in *C. elegans* embryos and *Drosophila*

neuroblasts suggest that RZZ prevents errors by inhibiting premature stabilization of

interactions between the ends of microtubules and NDC80 [23, 24]. RZZ also has the

property of streaming off the kinetochore, which could alleviate the inhibition of end-on

attachments [24-26]. We have shown that the RZZ complex is required for accurate

homolog biorientation in meiosis I, which could indicate a role for the RZZ complex in

delaying the formation of end-on attachments. We also show that Spindly, a dynein

adaptor for RZZ, is also required for homolog biorientation in meiosis I. However, RZZ

streaming in oocytes is not required for, nor does it depend on, the formation of end-on

attachments. We propose that RZZ streams in response to the silencing of the SAC

prior to the metaphase I arrest. Biorientation, however, depends on an interaction

between RZZ, Spindly, and possibly other motor activities, that regulate chromosome

movement and attachment status.

Results

Analysis of the SAC protein localization in oocytes

The localization of SAC proteins was examined in *Drosophila* oocytes. Mature

Drosophila oocytes, known as stage 14, arrest at metaphase I with the chromosomes

clustered together in a single mass, or karyosome, and the centromeres of homologous

chromosomes oriented towards opposite poles. Several SAC proteins were detected at

kinetochores in stage 14 oocytes, including BUBR1, BUB3, MPS1 and ROD (Figure

1A). In addition to its kinetochore localization, ROD localized to the spindle region

between the pole and the centromeres (Figure 1). This redistribution of ROD from the

kinetochores to the spindle is termed "streaming" and has been previously observed in

mitotic cells [25, 27, 28]. Another RZZ subunit, ZW10 tagged with HA, overlapped with

ROD^{GFP} staining, validating that ROD^{GFP} provides an accurate representation of the

RZZ complex's localization (Figure S 1A).

In contrast to the above proteins, we did not detect MAD1 in stage 14 oocytes

(Figure 1A). Therefore, we attempted to increase SAC signaling by incubating stage 14

oocytes in colchicine for 60 minutes to remove nearly all microtubules. Following this

treatment, robust kinetochore localization of MAD1 was observed in all oocytes (Figure

1B). Because colchicine induced a change in MAD1 localization, we tested if the

localization of other checkpoint protein was increased by loss of microtubules. BUBR1

and MPS1 intensity increased in colchicine, suggesting their levels are sensitive to the

SAC (Figure 1C). We did not, however, observe a SAC response with BUB3

localization.

A more dramatic response was observed with ROD. In colchicine-treated

oocytes, ROD expanded around the kinetochores (Figure 1B). In some cases, ROD

expanded to surround the karyosome and individual kinetochore foci could no longer be

observed. This behavior of the RZZ complex likely corresponds to expansion of the

fibrous corona, observed in miotic cells that lack kinetochore-microtubule (KT-MT)

attachments [25, 29]. This expansion is thought to facilitate microtubule capture by

increasing the surface area of the kinetochore which can contact the microtubules.

Thus, in *Drosophila* meiotic oocytes, the fibrous corona expands when microtubules are

absent.

Activation of the SAC is also associated with elevated Aurora B activity [30]. In

Drosophila oocytes, the chromosomal passenger complex (CPC) is recruited to the

chromosomes but then moves to the central spindle in prometaphase I oocytes [31].

CPC proteins are not detected at the meiotic centromeres in fixed samples [31, 32]. In

colchicine-treated oocytes, however, the CPC component INCENP localized to the

chromosomes and the centromeres (Figure 1B, C). These results suggest that, like

MAD1, colchicine treatment causes increased localization of the CPC to the

centromeres.

Effect of kinetochore-microtubule attachments on the SAC

response

In all colchicine-treated oocytes, MAD1 localization was consistent and observed

on all kinetochores, suggesting a global SAC response to the absence of microtubules.

To investigate the mechanism of SAC activation, we examined MAD1 localization in

oocytes where end-on attachments or tension was disrupted. NDC80 is required for

end-on attachments, so nearly all attachments in *Ndc80^RNAi* oocytes are lateral [22].

MAD1 was not detected in *Ndc80^RNAi* oocytes unless they were treated with colchicine,

suggesting that the SAC is insensitive to the loss of end-on attachments (Figure 2A-C).

We also examined MAD1 in a *mei-218* mutant (*mei-218^null*), which has a high error rate

and lacks tension on the kinetochores due to the absence of chiasmata [33]. MAD1 was

not detected in *mei-218^null* oocytes unless they were treated with colchicine, suggesting

that loss of tenson does not induce a SAC response (Figure 2A, B, D). The absence of

MAD1 in the untreated *mei-218^null* oocytes is consistent with the prior observation that

oocytes lacking chiasma bypass the metaphase I arrest and precociously enter

anaphase in stage 14 oocytes [33]. These results show that the SAC is only activated in

the absence of microtubules and does not respond to the lack of stable attachments,

biorientation errors, or tension.

Localization of checkpoint proteins in Drosophila oocytes

depends on SPC105R

The outer kinetochore protein KNL1/SPC105R is involved in the recruitment of

several checkpoint proteins in mitotic cells [34, 35]. Therefore, we investigated the role

of SPC105R, the *Drosophila* homolog of KNL1, in the recruitment of MAD1, ROD,

MPS1, and the CPC to kinetochores in *Drosophila* oocytes. To test if SPC105R is

required for the recruitment of INCENP and MAD1, we treated *Spc105R^RNAi* oocytes

with colchicine. Colchicine-induced kinetochore localization of both INCENP and MAD1

depended on SPC105R (Figure S 2A-D). However, both still localized to the

chromosomes. In the absence of SPC105R, MAD1 moved to the pericentric regions

adjacent to the centromeres, while INCENP localized to all chromatin and occasionally

to regions adjacent to the chromatin. The observation that MAD1 localized in colchicine-

treated oocytes depleted of SPC105R suggests that the oocyte can respond to the loss

of microtubules even in the absence of the outer kinetochore.

We also examined localization of INCENP and MAD1 in *Spc105R* deletion

mutants to investigate the domains of SPC105R required for the SAC response (Figure

S 3). Because *Spc105R* is an essential gene, we used a combination of *Spc105R^{RNAi}*

and RNAi-resistant transgenes to examine mutants in oocytes. The control for each

mutant was an RNAi-resistant wild-type transgene, *Spc105R^B* (Figure S 3). MAD1

kinetochore localization was reduced in mutants such as *Spc105R^C* or *Spc105R^{ΔM}* that

deleted motifs (MELT, KI, and ExxEED) in the central region of SPC105R (Figure S 2A,

B). INCENP kinetochore localization was also lost from the kinetochore in the majority

of *Spc105R^C* or *Spc105R^{ΔM}* oocytes, but due to the diffuse chromatin localization, this

was not reflected in significant differences in intensity (Figure S 2C, D). These results

suggest that the MELT-like, KI-like, and ExxEED repeats are involved in recruiting

MAD1 and the CPC to the kinetochore as part of the SAC response.

MPS1 localization depends on SPC105R, as shown by its drastic reduction in

Spc105R^{RNAi} oocytes (Figure S 4A, B). *Ndc80^{RNAi}* oocytes also had a decrease in MPS1

localization, but this was less severe than *Spc105R^{RNAi}* oocytes, indicating that the

majority of MPS1 is recruited by SPC105R (Figure S 4A, C). Similarly, the *Spc105R^C*,

Spc105R^{ΔM}, and *Spc105R^{ΔExxEED}* mutants had significant decreases in MPS1

localization, but not to the same degree as the *Spc105R^{RNAi}* oocytes (Figure S 4B, D).

Each of these mutants retains the C-terminal domain of SPC105R that recruits NDC80

[22]. Thus, these mutants may recruit more MPS1 than *Spc105R^{RNAi}* oocytes because

NDC80 recruits some MPS1. Therefore, these results suggest that MPS1 is recruited by

domains within both SPC105R and NDC80.

ROD localization also depends on SPC105R, as shown by its reduction in

Spc105R^{RNAi} oocytes (Figure 3A, B). In comparison to the other SAC proteins,

recruitment of RZZ depends on a smaller and specific domain of SPC105R (Figure S 3,

Figure 3, D). Oocytes with a deletion of everything except the KT binding domain

(*Spc105R^{C}*) or a deletion of all the sequences between the N-terminal and C-terminal

domains (MELT-like, KI-like, and ExxEED, termed *Spc105R^{ΔM}*) had almost no ROD^{GFP}

localization, similar to *Spc105R^{RNAi}* oocytes. Furthermore, a deletion of the MELT-KI

(*Spc105R^{ΔMELT-KI}*), but not the ExxEED domain, had reduced ROD localization. These

results suggest that ROD is recruited by the MELT-KI region. Smaller deletions of only

the KI and MELT domains (*Spc105R^{ΔKI}* and *Spc105R^{ΔMELT}*) resulted in significantly

reduced ROD^{GFP} localization at the kinetochores, although these intensities were higher

than in the *Spc105R^{ΔMELT-KI}* deletion. This suggests that ROD is recruited by both the

MELT and KI regions.

We also tested if the localization of ROD and MPS1 depended on other SAC

proteins. ROD (Figure 4A, B) and MPS1 (Figure 4C, D) localization was significantly

reduced in both *Bub3^{RNAi}* and *BubR1^{RNAi}* oocytes. The more significant reductions that

were observed with *Bub3* knockdown could be due to less efficient *BubR1^{RNAi}* (see

Methods), or redundancy between BUBR1 and BUB1. We also tested the effect of

Cyclin A knockdown because it may regulate MPS1 localization [36]. *CycA^{RNAi}* oocytes

had a significant reduction in MPS1 localization, although not as strong as *Bub3^{RNAi}*,

and a less significant reduction in ROD localization. Furthermore, MPS1 and ROD

localization was inter-dependent, although there was a milder reduction of ROD in

Mps1^{RNAi}. These results suggest ROD and MPS1 localization depends on both SAC

proteins like BUB3 and Cyclin A.

RZZ and MPS1, but not other SAC proteins, are required for

meiosis I biorientation

To determine which SAC genes are required for biorientation during meiosis I,

we used fluorescent in-situ hybridization (FISH) probes against the pericentromeric

regions of the 2nd, 3rd, and X chromosomes. Biorientation was defined as two

homologous centromeres at opposite ends of the karyosome (Figure 5A). A mono-

orientation defect was defined as the two homologous centromeres being in the same

half of the karyosome. Oocytes from females expressing *rod^{RNAi}* or *Mps1^{RNAi}* had

increased chromosome mono-orientation (Figure 5B). The results with *Mps1* are

consistent with previous studies showing nondisjunction using fertile allele combinations

of *Mps1* [37, 38]. We also tested *CycA^{RNAi}* oocytes, and consistent with a previous

report [39], found an elevated level of biorientation defects (Figure 5B).

In contrast, *BubR1^{RNAi}*, *Bub1^{RNAi}, and Bub3^{RNAi}* oocytes had insignificant

frequencies of mono-orientation (Figure 5B). Similarly, we have previously shown that

flies with knock down of *Bub1* by RNAi are fertile and do not have increased

nondisjunction [31]. BUBR1 is required for sister chromatid cohesion protection in

Drosophila, however, this may have not been detected by our FISH assay at metaphase

I [22, 40]. Mutants of *mad1* and *mad2* are viable and fertile [41, 42], so we used genetic

crosses to measure the frequency of nondisjunction (see Methods). Neither mutant had

a significant increase in nondisjunction (Table 1). We also used a balancer chromosome

to suppress crossing over on the X chromosome (*Bwinscy*), thus testing the role of

mad1 and *mad2* in an achiasmate system [43]. Only a mild increase in nondisjunction

was observed. These results suggest that most of the SAC proteins are not required for

biorientation of homologous chromosomes at metaphase I, while the RZZ complex and

MPS1 may be involved in error correction in addition to their roles in the meiotic SAC.

Stage 14 oocytes arrest in metaphase I [33, 44]. If the metaphase I arrest

depends on the SAC, we would predict that SAC mutants would precociously bypass

this arrest and have increased oocytes with split karyosomes. However, this was not

observed in most oocytes, suggesting the metaphase I arrest does not depend on SAC

activity (Figure 5C, D). The most striking exceptions were *rod^{RNAi}* and the *mad2* mutant.

The split karyosome phenotype is less than observed in mutants lacking crossing over

[22, 33], suggesting these genes could have a minor role in the metaphase I arrest, or in

the case of Rod, the precocious anaphase could result from the bi-orientation defects

(see below).

ROD streaming is independent of end-on attachments

MSP1 and ROD may have two distinct functions, one in the SAC and another for

biorientation. A striking feature of RZZ is its streaming off the kinetochores, but how

streaming contributes to kinetochore attachments and biorientation in meiosis is not

known. A small fraction of wild-type oocytes have no ROD streaming, which are likely

oocytes in early prometaphase [45]. If streaming is in response to proper attachment of

kinetochores to microtubules, we predicted that RZZ would not stream if errors were

present, or if end-on attachments could not form. To determine if biorientation errors

blocked streaming, we examined ROD in *mei-218^{null}* oocytes, in which 90% of

crossovers are absent, resulting in frequent chromosome segregation errors [46]. In the

absence of crossovers, we still observed precocious anaphase and ROD streaming

(Figure 6A,B). Thus, streaming occurs even in the absence of tension and in the

presence of attachment errors.

To determine if attachment status regulates streaming, we examined ROD in

Ndc80^RNAi oocytes, which lack end-on attachments [21]. Despite the lack of end-on

attachments, ROD streamed in all *Ndc80^RNAi* oocytes, and there was no effect on ROD

accumulation at the kinetochores (Figure 6A-C). Furthermore, oocytes lacking protein

phosphatase 2A-B56 (PP2A-B56), which is also required for the creation of end-on

attachments [47], still exhibited ROD streaming (Figure 6A-C). Indeed, there may even

be a reduction of ROD at the kinetochore in *PP2A-B56^RNAi* oocytes. These results

demonstrate that end-on attachments are not required for the initiation of RZZ

streaming.

ROD is not required for lateral or end-on attachments

Microtubule nucleation from the kinetochore could depend on RZZ [48]. Thus, the

biorientation defects observed in the *rod^RNAi* oocytes could be caused by the inability to

create lateral or end-on attachments, or the creation of inaccurate attachments. If ROD

is necessary for making end-on attachments, we would expect that oocytes depleted of

rod alone would have lateral attachments. However, most attachments in *rod^RNAi*

oocytes were end-on (Figure 7A, B). Interestingly, there was an increase of lateral

attachments in *rod^RNAi* compared to wild-type oocytes. This could be explained if *rod^RNAi*

oocytes are delayed in making end-on attachments, possibly due to frequent

biorientation errors. To test whether ROD was necessary for forming lateral

attachments, we knocked down both *rod* and *Ndc80* (*rod^RNAi*, *Ndc80^RNAi*). Similar to

Ndc80^RNAi, we observed that *rod^RNAi*, *Ndc80^RNAi* oocytes had primarily lateral

attachments, suggesting that ROD is not required for making lateral attachments

(Figure 7A, B).

As noted above, *rod^RNAi* oocytes have an elevated frequency of separated

karyosomes. This phenotype was not observed in *rod^RNAi*, *Ndc80^RNAi* oocytes,

suggesting that separated karyosome phenotype is caused by the formation of end-on

attachments is required for the (Figure 7A, C).

ROD localization to the kinetochore may delay, but does not

prevent, end-on attachments

Because ROD is not required for making lateral or end-on attachments, we

reasoned that it could be required for regulating the transition from lateral to end-on

attachments. This would align with a model proposed in mitotic cells in which high levels

of RZZ at the kinetochores inhibit NDC80 from prematurely establishing end-on

attachments [23, 24]. In this model, RZZ streaming allows for the establishment of end-

on attachments. If this was true in meiosis, we would expect oocytes in which ROD was

exclusively on kinetochores and did not stream to have lateral attachments. This ROD

behavior was observed in two *Spc105R* mutants. A deletion of the ExxEED repeats

(*Spc105R^ΔExxEED*) or the N-terminal region (*Spc105R^ΔN*) had higher ROD intensities at

the kinetochores (Figure 3) and less streaming than in *Spc105R^B* oocytes (Figure 7D,

E). However, oocytes with ROD only on kinetochores still exhibited end-on attachments

(Figure 7D). These results suggest that the N-terminal region and ExxEED repeats of

SPC105R may regulate streaming, and that RZZ streaming is not required for the

creation of end-on attachments in *Drosophila* oocytes.

To further test whether end-on attachments are inhibited when the RZZ complex

is localized to the kinetochore, we stably targeted RZZ to the kinetochore. The rationale

for this experiment is that the recruitment of the RZZ components are interdependent

[49]. We fused *Zw10* to the KT-binding C-terminal domain of *Spc105R* (*Zw10-*

Spc105R^C), along with three copies of the HA epitope at the C-terminal end (Figure S

3). The ZW10-SPC105R^C fusion protein was targeted to the kinetochores (Figure 7F).

When *Zw10-Spc105R^C* was combined with an shRNA targeting the endogenous

Spc105R (*Spc105R^RNAi*), loss of endogenous SPC105R was confirmed cytologically

using an antibody against the first 400 amino acids of SPC105R, which is absent in the

fusion construct (Figure S 5A). In addition, ROD localized to the kinetochores and did

not stream in *Zw10-Spc105R^C* oocytes, indicating that the RZZ complex was restricted

to the kinetochores (Figure S 5B). ROD^GFP did not stream even when there was

endogenous SPC105R present at the kinetochores. Thus, the ZW10-SPC105R^C fusion

protein prevented endogenous RZZ streaming from kinetochores. *Zw10-Spc105R^C*

oocytes exhibited an increase in lateral attachments compared to *Spc105R^C* oocytes

(Figure 7G). However, the majority of attachments were still end-on (~60%), indicating

that RZZ may delay, but does not prevent, end-on attachments. Therefore, in

agreement with the two *Spc105R* mutants (Figure 7D), streaming of RZZ is not required

to establish end-on attachments in *Drosophila* oocytes. Indeed, end-on attachments

occur while RZZ is localized to the kinetochore.

Spindly regulates ROD streaming

Streaming is proposed to be dependent on the microtubule motor protein Dynein

[50]. To further examine the role of RZZ streaming and the interaction with Dynein in

meiosis, we investigated two dynein-associated proteins previously implicated in RZZ

removal from the *Drosophila* kinetochores: Spindly [51], and NudE [52]. We performed

FISH on stage 14 oocytes to determine whether these two genes are required for

chromosome biorientation (Figure 8A, B). There was only a slight increase in

chromosome mono-orientation in *nudE^RNAi* oocytes compared to wild-type (9% and 0%,

respectively). However, 35% of chromosomes were mono-oriented in *Spindly^RNAi*

oocytes, suggesting that Spindly is necessary for accurate meiotic chromosome

segregation.

To determine whether these proteins regulate streaming, we characterized ROD

behavior in oocytes depleted of *Spindly* or *nudE* (Figure 8C-E). Both *Spindly^RNAi* and

nudE^RNAi oocytes exhibited ROD streaming (Figure 8D). This was surprising for Spindly,

which has been shown to be required for streaming in *Drosophila* S2 cells [51].

However, *Spindly^RNAi* oocytes also exhibited unusual behavior, notably ROD streaming

in the central spindle region between homologous centromeres in approximately 50% of

oocytes (Figure 8E, Figure S 6A). Thus, Spindly may be required for proper streaming

towards the spindle pole. To confirm that ROD was being efficiently removed and not

accumulating at the kinetochores, we measured the intensity of ROD in *Spindly^RNAi* and

nudE^RNAi oocytes (Figure 8F). Both RNAi oocytes had ROD levels similar to those in

wild-type oocytes, indicating that ROD can still stream from kinetochores without NudE

or Spindly. Therefore, while neither protein is required for RZZ streaming, Spindly may

be required for RZZ to stream towards the spindle poles.

RZZ is proposed to recruit Spindly to the kinetochore, which is followed by

streaming of both Spindly and RZZ [51, 53]. To examine the localization of Spindly on

the meiotic spindle, we used previously described *Spindly-GFP* fusion transgenes [54].

Oocytes expressing wild-type *Spindly* had weak localization throughout the spindle

(Figure S 6B). We also examined *Spindly* mutants with deletions of domains implicated

in streaming: the N-terminus (a deletion of amino acids 1-228) and the Spindly Box (a

deletion of amino acids 229-250). Both mutants had a dominant sterility phenotype

when expressed in oocytes using *matα*, suggesting that these mutants interfere with the

activity of the wild-type Spindly protein. In contrast to the wild-type oocytes, which had

weak localization throughout the spindle, both mutants had pronounced localization

near the centromeres. Spindly$^{\Delta N}$ localized to the kinetochores and spindle, while

Spindly$^{\Delta SB}$ primarily localized to the kinetochores (Figure S 6B, C). These results

suggest that the Spindly Box domain is important for removal of Spindly from the

kinetochores. The robust localization of these mutants was then used to test if ROD is

required for Spindly localization, as shown in mitotic cells [51]. Indeed, Spindly was

absent in *rod^{RNAi}* oocytes, indicating ROD is required for Spindly localization in oocyte

meiosis (Figure S 6C, D).

Discussion

The function of the SAC in oocytes

We analyzed the localization and meiotic phenotypes of several SAC proteins in

Drosophila oocytes. Increased localization of MAD1, BUBR1, INCENP, and ROD at the

kinetochores following loss of microtubules indicated *Drosophila* oocytes have a

functioning SAC. This SAC response was observed even in the absence of a

kinetochore. However, a SAC response was not observed following loss of microtubule

attachments to the kinetochores, the lack of tension, or the presence of multiple

biorientation errors. Thus, the function of the SAC in oocytes may be to promote spindle

assembly, rather than proper microtubules attachments to kinetochores [28].

Only two of the SAC proteins we tested were required for biorientation of

homologous chromosomes at meiosis I, MPS1 [37, 38] and ROD, suggesting that the

SAC is not required for error correction. Similarly, *C. elegans* may correct attachment

errors using a mechanism that does not rely on the SAC [55]. The SAC may have a

more prominent role in chromosome segregation in mouse oocytes that arrest in

meiosis II, where increased aneuploidy has been observed with loss of SAC proteins

[56, 57]. However, the evidence comes from a variety of assays that may not be

comparable to our FISH experiments, and one study concluded that Mad2 is

dispensable for accurate meiotic chromosome segregation except in response to stress

[58].

In most cases, the loss of SAC proteins in *Drosophila* oocytes does not result in

progression into anaphase (Figure 5) or a rise in APC activity [59]. These observations

suggest the SAC may not regulate the APC in metaphase I-arrested oocytes. Instead,

the metaphase I arrest in *Drosophila* oocytes may be maintained by a SAC-independent

mechanism relying on chiasmata and cohesion, with progression into anaphase I

depending on cohesion release. This would explain why *Drosophila* oocytes lacking

chiasmata and tension bypass metaphase I arrest and progress into anaphase I [33,

60].

Furthermore, several observations suggest that the SAC is silenced in oocytes,

possibly to allow for developmental regulation of the metaphase I arrest, with

progression into anaphase occurring only after oocytes enter the oviduct.

Developmentally regulated SAC silencing would explain why ROD streams in wild-type

metaphase I-arrested oocytes, even in the absence of end-on attachments or the

presence of multiple biorientation errors in *Ndc80^{RNAi}* and *PP2A-B56^{RNAi}* oocytes [61].

Similarly, MAD2 localizes to centromeres in early (stage 13) but not late (stage 14)

stage oocytes [45], and the metaphase I spindle has some anaphase-like

characteristics such as high concentrations of the CPC on the central spindle [62].

Yeast meiotic cells silence their SAC to prevent prolonged arrest, indicating this may be

a conserved feature of meiotic cells [63].

Recruitment of checkpoint proteins during meiosis

Our results are consistent with the known role of KNL1/SPC105R in the

recruitment of checkpoint proteins to the kinetochore [1]. We previously showed that

BUBR1 is recruited by both the MELT-KI and C-terminal regions of SPC105R [22].

MAD1, INCENP, and probably BUB3 depend on the MELT-KI and ExxEED repeat

regions of SPC105R, which correspond to the repeated MELT motifs found in

vertebrate KNL1. In the absence of SPC105R, recruitment of MAD1 shifts to the peri-

centromeric regions. Thus, some components of the SAC do not depend on SPC105R,

and there is a SPC105R-independnet SAC response. Human cells may also have two

SAC pathways, and one is active in the absence of KNL1 [64]. Thus, centromere

components may recruit SAC proteins. In *Drosophila*, this could be facilitated by ZW10,

which interacts with and is recruited by centromere protein CAL1 [65].

Multiple functions and recruitment factors for MPS1 and RZZ

MPS1 was originally discovered with the mutation *ald* as being required for

meiotic chromosome segregation [66], and subsequent studies have characterized its

role in *Drosophila* meiosis [37, 38] and mouse oocytes [67]. While MPS1 has been

shown to be recruited by NDC80 [68-71], two observations show that a significant

amount of MPS1 is also recruited by SPC105R. First, we observed a more severe

decrease in MPS1 localization upon *Spc105R* depletion than *Ndc80* depletion. Second,

we observed a decrease in MPS1 localization in *Spc105R* mutants that still recruit

NDC80. These data suggest that MPS1 is recruited by both NDC80 and SPC105R in

Drosophila oocytes. The SPC105R-dependent MPS1 could function in the SAC while

NDC80-dependent MPS1 could regulate attachments [68, 72].

Like MPS1, the RZZ complex has roles in error correction and in the SAC. Like

MPS1, our data suggest there are two separate RZZ populations: one that is recruited

by BUB3 and BUBR1 and is involved in the SAC and a second that is recruited by the

MELT-KI domains of SPC105R and is important for EC. The SAC population of ROD

that depends on BUB3 could recruit other checkpoint proteins like MAD1 and MAD2

[73, 74]. These conclusions are based on the observation that unlike *rod*[RNAi], *Bub3*[RNAi]

and *BubR1*[RNAi] oocytes do not have biorientation defects, suggesting that BUB3 and

BUBR1 do not recruit the population of RZZ which is involved in error correction.

Furthermore, there is a substantial amount of ROD^{GFP} remaining after *Bub3* and *BubR1*

are knocked down, arguing that they do not recruit the total population of RZZ at

kinetochores. However, the *Spc105R$^{\Delta MELT-KI}$* mutant (deleting amino acids 154-840)

abolishes nearly all ROD^{GFP} localization and does have biorientation defects [22].

Therefore, the MELT-KI domains of SPC105R are likely involved in recruiting the

populations of RZZ involved in EC, independently of BUB3.

The MELT-KI region of SPC105R is highly conserved [75]. Indeed, our results

refine observations made in HeLa cells showing that the N-terminal domain of KNL1

recruits RZZ [76, 77]. In *Drosophila* cell lines, a region within the KI-like domain (amino

acids 266-384) and BUBR1 are required for RZZ recruitment to kinetochores [78].

Interestingly, this study also showed that the recruitment of RZZ by SPC105R could

promote chromosome movement towards the spindle poles. Like this study, we found

the KI domain recruits the majority of RZZ. It remains to be determined, however, why

both the MELT and KI domains are required to recruit RZZ and for biorientation in

meiosis I [22].

ROD and MPS1 localization was reduced in *CycA^{RNAi}* oocytes, and *CycA^{RNAi}*

oocytes have biorientation defects in *Drosophila* oocytes [39]. Indeed, Cyclin A has

been shown to regulate KT-MT attachments in mammalian cells [79, 80] and in

Drosophila in mitotic cells, Cyclin A/CDK1 activates Polo kinase, which has recently

been shown to regulate Spindly and prevent it from inhibiting RZZ [24]. One explanation

for these results is that Cyclin A is required for the functions of ROD and MPS1

associated with ensuring the accuracy of biorientation.

The role of RZZ localization in attachment stabilization

Previous studies in *C. elegans, Drosophila,* and human mitotic cells have

suggested a two-part model for how RZZ regulates KT-MT attachments [23, 24, 81, 82].

First, RZZ interacts with NDC80 to inhibit end-on attachments. Second, Spindly-

dependent RZZ streaming alleviates RZZ inhibition to allow for the formation of end-on

attachments. Three of our observations are consistent with this hypothesis. First,

knockdown of *Spindly* resulted in a decrease in end-on and an increase in lateral

attachments. This would be expected if Spindly negatively regulates RZZ. Second,

Zw10 fused to the C terminus of *Spc105R* led to complete RZZ retention on the

kinetochores and increased lateral attachments. Third, we observed an increase in

karyosome separation in *rod^{RNAi}* oocytes, which could be due to precocious stabilization

of end-on attachments [22, 83]. These results suggest that the RZZ complex at the

kinetochore has a role in preventing or delaying end-on attachments.

However, some of our results are not consistent with this model. Oocytes

expressing *Zw10-Spc105R^{C}* primarily had end-on attachments. A similar observation

was made with two mutants (*Spc105R^{ΔExxEED}* and *Spc105R^{ΔN}*) that had reduced ROD

streaming but still formed end-on attachments. Therefore, if KT-localized RZZ inhibits

end-on attachments, this can be overridden in oocytes. Finally, ROD streaming was

observed in the absence of end-on attachments. Thus, physical removal of RZZ (or

streaming) may not be required for the formation of end-on microtubule attachments,

and alternative mechanisms for regulating this population of RZZ may exist. For

example, a conformational change in RZZ may affect its regulation of attachments [23].

Understanding the relationship between the localization of the RZZ complex and its

effects on microtubule attachments is crucial for analyzing its role in promoting accurate

chromosome segregation during female meiosis.

The role of Spindly in meiotic RZZ streaming and homolog

biorientation

Spindly has a conserved role as a Dynein adaptor [84], and we have shown that

RZZ recruits Spindly to kinetochores in *Drosophila* oocytes. Spindly has been shown in

multiple organisms to recruit Dynein, resulting in removal of the RZZ complex from the

kinetochores to stabilize microtubule attachments [24, 53, 81, 85]. This activity has been

used to explain why depletion of RZZ can suppress defects in mitotic cells depleted of

Spindly [23, 24, 81]. However, the role of Spindly in meiosis has not been reported.

Spindly is required for biorientation of homologous chromosomes during oocyte

meiosis. The failure to remove RZZ from the kinetochores could be the reason for the

meiotic biorientation defect. Contrary to the expectations from data in mitotic cells

described above, however, knocking down *Spindly* did not abolish ROD streaming.

Instead, we found that ROD displayed abnormal behaviors in oocytes depleted of

Spindly, such as streaming in the opposite direction and localizing in the central spindle

region between kinetochores. Therefore, it is likely that other proteins are involved in the

removal of RZZ from the kinetochores in oocytes.

In the absence of Spindly in HeLa cells, alternative mechanisms are activated to

remove kinetochore proteins [86]. One candidate is CENP-E, which is a plus-end

directed motor required for the fibrous corona [87], and is required for homolog

biorientation in *Drosophila* oocytes [21]. Given that RZZ can move SPC105R oligomers

[88], and our observation that streaming off the kinetochore may not affect attachment

status, an important role of RZZ may be to move kinetochores. Rather than being

necessary for the direct removal of RZZ from the kinetochores, Spindly, CENP-E, and

RZZ might form a complex capable of bidirectional movement of the kinetochores [89].

Understanding how RZZ regulates biorientation may depend on reconciling the relative

importance of two activities, RZZ moving kinetochores with motors like Dynein and

CENP-E, and RZZ blocking or delaying end-on attachments.

Materials and Methods

Drosophila Genotypes and Expression of Transgenes and

shRNAs

The transgenes and mutants used in this study are listed in Table 2. The

localization of SAC proteins was examined in *Drosophila* oocytes using GFP fusion

transgenes, most under the control of their own promoter. To examine MPS1, we used

a construct used extensively in cell culture but modified for making transgenics

($P\{EGFP\text{-}Mps1.F\} = Mps1^{GFP}$) [90, 91]. To visualize the RZZ complex in oocytes, we

used a construct where ROD was tagged with Green Fluorescent Protein (GFP) at the

N-terminus and under the control of its own promoter ($P\{gEGFP\text{-}rod\} = rod^{GFP}$) [25]

(Figure S 1A). Expression of either transgene resulted in very low levels (~0.25%) of X-

chromosome nondisjunction (Table S1), which was comparable to wild-type strains [92].

This is consistent with previous studies showing that this transgene can rescue the

lethality of *rod* mutants [25, 93].

The UAS/GAL4 system was utilized to drive expression of many of the

transgenes in oocytes. *Matα* was used to induce expression following DNA replication

and early pachytene and throughout meiotic prophase.

A fusion between the C-terminal domain of *Spc105R* and the entire coding region

of *Zw10* was generated by GenScript in pENTR4. The C-terminal fragment of SPC105R

has been characterized previously and is sufficient for kinetochore localization [22]. The

fusion gene was transferred into pPWH, which fused the coding region to a 3XHA tag at

547   the C-terminal end. This was injected into embryos for germline transformation by

548   Model Systems Injections.

RNA Extraction and Quantification of RNAi Knockdown

Among three shRNA lines targeting *rod*, one (*P{Val22:rod-153}1attP40, =*

rod$^{RNAi)}$) was chosen because it had the strongest phenotype and most severe mRNA

knockdown (4% of the wild-type). Expression of *rod^{RNAi}* using *matα* produced sterile

females and ubiquitous expression caused lethality. To confirm the efficacy of the

rod^{RNAi} in oocytes, we examined ROD^{GFP} in oocytes expressing the *rod* shRNA. The

intensity of ROD^{GFP} at the kinetochores in *rod^{RNAi}* oocytes was reduced to nearly

background levels (Figure S 1B, C), indicating that the *rod* shRNA provides a strong

knockdown of *rod*. Similarly, the intensity of MPS1^{GFP} at the kinetochores in *Mps1^{RNAi}*

oocytes was significantly reduced (Figure S 1D, E).

RT-qPCR was performed to determine knockdown of mRNAs for each shRNA.

After the parental flies were crossed, mature female offspring with the correct genotype

were collected into yeasted vials for three days. These flies were then ground in 1X

phosphate-buffered saline (PBS) and filtered through meshes to isolate the stage 14

oocytes. 1 mL of TRIzol reagent was added and RNA was extracted per the

manufacturer's instructions (Thermo Fisher Scientific). Taqman RT-qPCR was

performed to quantify RNAi knockdowns. 2 μg of isolated RNA was reverse transcribed

into cDNA using the High Capacity cDNA Reverse Transcription Kit (Applied

Biosystems). Afterwards, qPCR was performed using a TaqMan Assay (Thermo

Scientific) with four replicates per reaction. The efficacy of RNAi knockdowns has been

published for *Ndc80 GL00625* = 6% remaining*, Spc105R GL00392* = 13% remaining

[21], *Bub1 GL00151* = 2% remaining [31], *BubR1 GLV21065* = 22% remaining, *BubR1*

GL00236= 27% remaining, and *Bub3 HMS00789* = 0.7% remaining.

Measuring X-Chromosome Nondisjunction

To measure nondisjunction of the X chromosome, virgin females with the desired

genotype were crossed to males with a dominant *Bar* mutation on the Y chromosome

($yw/B^S Y$). This mutation led to a visible bar-eyed phenotype where the eye was thinner

than wild-type eyes. The normal segregation of chromosomes would lead to female

offspring (XX) with wild-type eyes and males (XY) with bar eyes. If a nondisjunction

event occurred in the X chromosome, the following sex chromosome combinations were

possible: XO, XXY, XXX, and YO. XO and XXY were viable and resulted in normal-eyed

males and bar-eyed females, respectively. XXX and YO were lethal. To calculate

nondisjunction rates while taking into account the lethal genotypes, the following

equation was used: $\frac{\# \; flies \; with \; nondisjunction \times 2}{(\# \; flies \; with \; nondisjunction \times 2) \; + \; (\# \; of \; normal \; flies)}$. This rate determined

whether a particular genotype resulted in increased meiotic nondisjunction.

Cytology of Stage 14 Oocytes

Cytology of stage 14 oocytes was performed as previously described [94]. After

the parental flies were crossed, mature female offspring with the correct genotypes

were collected into yeasted vials for one to two days. These flies were then broken open

in a blender while in 1x modified Robb's buffer and filtered through meshes to isolate

the stage 14 oocytes. If a colchicine treatment was needed, oocytes were incubated

with 150 µM colchicine or 0.5% EtOH in modified Robb's buffer for one hour to remove

all microtubules. The oocytes were fixed with 5% formaldehyde and heptane before

being rinsed in 1X PBS. The oocytes were then rolled between the frosted surface of a

glass slide and a cover slip to remove the membranes and nutated for two hours in

PBS/1% Triton X-100 to make them permeable to antibody staining. Afterwards,

oocytes were washed in PBS/0.05% Triton X-100 and blocked for one hour in blocking

buffer (0.5% BSA and 0.1% Tween-20 in 1X PBS). Subsequently, primary antibodies

were added to the oocytes while the secondary antibodies were added to *Drosophila*

embryos to remove any nonspecific binding. After nutating overnight at 4°C, the oocytes

were washed in blocking buffer four times at room temperature for 15 minutes. The

secondary antibodies were added and the oocytes were nutated for 3-4 hours at room

temperature. The oocytes were then washed in blocking buffer with Hoechst33342 (10

µg/mL) to stain the DNA. The oocytes were washed twice more in blocking buffer and

then were ready for mounting and imaging.

The primary antibodies used in this paper were tubulin monoclonal antibody

DM1A at 1:50 conjugated to FITC (Sigma-Aldrich), tubulin E7 monoclonal antibody at

1:150 [95], rat anti-INCENP at 1:600 [96], guinea pig anti-CENP-C at 1:1000 [97], rat

anti-tubulin at 1:300 (Millipore), rabbit anti-Spc105R (1:4000) [98], rabbit anti-GFP

(1:200) (Thermo Fisher), rat anti-HA (Roche, 1:50). Multiply preabsorbed secondary

antibodies were from Jackson ImmunoResearch, and were conjugated to Alexa 647,

Cy3, Alexa 594 or Alexa 488.

To visualize pairs of homologous chromosomes, fluorescent *in situ* hybridization

(FISH) was performed. After oocytes fixation, 2X SSC was added and the membranes

were removed by rolling the oocytes between glass slides. The oocytes were then

nutated in increasing concentrations of formamide solution (20%, 40%, and 50%) before

being added to the hybridization solution and probes. The probes were synthesized by

IDT, and recognize the 359 bp repeats on the X chromosome (labeled with AlexaFluor

594), the AACAC repeats on the second chromosome (labeled with Cy3), or the dodeca

repeats on the third chromosome (labeled with Alexa 647). This was followed by

incubation in 80°C for 20 minutes and then 37°C overnight. The next day, the oocytes

were washed in 50% formamide solution twice at 37°C and 20% formamide once at

room temperature. The oocytes were then rinsed three times in 2X SSCT, once in 2X

PBST, and nutated in blocking buffer for 4 hours at room temperature. Afterwards,

mouse anti-tubulin-FITC antibody (Sigma) was added (1:50) and the oocytes were

nutated overnight at room temperature. The next day, the oocytes were washed in

blocking buffer with Hoechst33342 (10 µg/mL). After washing twice more with blocking

buffer, the oocytes were mounted and imaged.

Imaging Oocytes, Image Analysis, and Statistical Analysis

Oocytes were mounted on a glass slide in SlowFade Gold (Invitrogen) and

images were taken with a 63x lens on a Leica SP8 or Stellaris microscope. Protein

intensities were quantified using Imaris image analysis software (Bitplane). To measure

the intensity of proteins at kinetochores, spots marking the kinetochores and random

areas of the background were created. The average intensity of the protein of interest at

the background was subtracted from the intensity at each kinetochore to generate the

normalized intensity value. GraphPad Prism software was used to graph data and

perform statistical analysis. All statistical tests and sample sizes are reported in the

figure legends.

Data availability statement

All the data underlying this study are available in the published article and its

online supplementary material.

Acknowledgements

We thank Marina Druzhinina for technical assistance, Christian Lehner, and

Rager Karess for providing Drosophila stocks and antibodies. Stocks obtained from the

Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this study. The

E7 tubulin monoclonal antibody was obtained from the Developmental Studies

Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of

Iowa, Department of Biology, Iowa City, IA 52242. Image acquisition and analysis was

made possible by the Waksman Institute Shared Imaging Core Facility and The Human

Genetics Institute Imaging Core Facility at Rutgers. This work was supported by NIH

grant GM101955 to K.S.M.

Table 1: Nondisjunction in *mad* mutants

Genotype	XX	XY	XXY	XO	% NDJ
mad2[EY21687]*/Df*	3857	3558	6	17	0.6
mad1[1]*/Df*	805	769	1	6	1.0
Bwinscy/+; mad2[EY21687]*/Df*	357	263	4	12	4.9
Bwinscy/+; mad1[1]*/mad1*[1]	427	321	9	4	3.4
Bwinscy/+; mad1[1]*/+*	588	375	2	4	1.2
Bwinscy/+; mad2[EY21687]*/+*	509	301	2	2	1.0

657   *mad2* Df = *Df(3L)Exel6107* and *mad1* Df = *Df(2R)ED1791*

662   Table 2. *Drosophila* genotypes used in this work.

Genotype	Abbreviation	Reference
P{w+mC=matalpha-GAL-VP16}V37	*matα*	[99]
P{Val22:rod-153}attP40	*rod^{RNAi}*	This paper
P{gEGFP-rod}	*rod^{GFP}*	[25]
P{EGFP-Mps1.F}	*Mps1^{GFP}*	[90]
P{Mad1-GFP}	*Mad1^{GFP}*	[42]
P{BubR1.GFP}	*BubR1^{GFP}*	[73]
P{gEGFP-Bub3}III.3	*Bub3^{GFP}*	[100]
P{w+mC=UASp-Spc105RB}	*Spc105R^B*	[22]
P{UASp-Spindly.RR.GFP}	*Spindly^{GFP}*	[54]
P{w+mC=UASp-Zw10-Spc105RC}	*Zw10-Spc105R^C*	This paper
P{TRiP.GL00392}attP2	*Spc105R^{RNAi}*	[101]
P{TRiP.GL00625}attP40	*Ndc80^{RNAi}*	[101]
P{TRiP.HMS01283}attP2	*Spindly^{RNAi}*	[101]
P{TRiP.HMS01868}attP40	*nudE^{RNAi}*	[101]
P{TRiP.GL00151}attP2	*Bub1^{RNAi}*	[101]
P{wrd^{GL00671}wdb^{HMS01864}}/TM3	*PP2A-B56^{RNAi}*	[101]
P{TRiP.GL00236}attP2	*BubR1^{RNAi}*	[101]
P{TRiP.HMS00789}attP2	*Bub3^{RNAi}*	[101]
Mad1^1	*Mad1*	[42]

Mad2^{EY21687}	*mad2*	[41]
mei-218⁶	*mei-218^{null}*	[102]
Mps1^{ald}	*Mps1^{ald}*	[66]
Mps1^{C3}	*Mps1^{C3}*	[37]

663
664

Figure Legends

Figure 1. **Localization of SAC proteins in oocytes**.

(A, B) Oocytes were incubated for one hour in Robb's buffer with 0.5% EtOH (A) or 250
μM colchicine (B). BUBR1, BUB3, MPS1, ROD, and MAD1 were detected using GFP-
tagged transgenes, while INCENP was detected using an anti-INCENP antibody (all
proteins shown in green), DNA is in blue, CENP-C in white, and tubulin in red. Single
channel images show SAC protein localization. All images are maximum intensity
projections of z stacks. Scale bars represent 5 μm. (C) Quantification of SAC protein
intensity at kinetochores, normalized to background signal (n for EtOH and colchicine =
313 and 174 for BUBR1, 67 and 83 for BUB3, 219 and 356 for MPS1, 139 and 184 for
ROD, 138 and 185 for MAD1, and 74 and 200 for INCENP). Error bars show mean ±
s.d.; ****P<0.0001, ns=no significance (unpaired two-tailed t test).
678

Figure 2. **Induction of MAD1 localization is independent of tension and attachment
status**.

(A, B) MAD1^{GFP} localization in wild-type, $Ndc80^{RNAi}$, or mei-218^{null} oocytes that were
either untreated (A) or treated with 250 μM colchicine (B). MAD1^{GFP} is shown in green,
DNA in blue, CENP-C in white, and tubulin in red. Single channel images (bottom) show
MAD1^{GFP} with karyosome outlines in dashed blue in (A). All images are maximum
intensity projections of z stacks. Scale bars represent 5 μm. (C, D) Quantification of
MAD1^{GFP} intensity at kinetochores, normalized to background GFP signal (n =106 and
127 kinetochores for (C) and 158 and 311 kinetochores for (D)). Error bars show mean
± s.d.; ****P<0.0001 (unpaired two-tailed t test).
689

Figure 3. **ROD localization depends on the MELT-KI domain of SPC105R**.

(A) ROD^{GFP} localization in control and an $Spc105R^{RNAi}$ oocytes, with ROD^{GFP} in green,
DNA in blue, CENP-C in white, and tubulin in red. Single channel images (bottom) show
ROD^{GFP}. (B) Quantification of ROD^{GFP} intensity at kinetochores normalized to
background GFP signal in indicated oocytes (from left to right, n = 142, 169, 144, 144,
167, 138, 206, 159, 173, and 146 kinetochores). Error bars show mean ± s.d.;
****P<0.0001, ***P=0.0006, ns=no significance (unpaired two-tailed t test). All $Spc105R$
mutants are in an $Spc105R^{RNAi}$ background targeting the endogenous $Spc105R$. (C)
Schematic of SPC105R^{B} domains in $Drosophila$ $melanogaster$. (D) ROD^{GFP} localization
in the indicated $Spc105R$ mutant oocytes, with ROD^{GFP} in green, DNA in blue, CENP-C
in white, and tubulin in red. Single channel images (bottom) show ROD^{GFP}. All images
are maximum intensity projections of z stacks. Scale bars represent 5 μm.
702

Figure 4. **ROD and MPS1 localization depend on SAC genes**.

(A) ROD^{GFP} localization in wild-type, *Bub3^{RNAi}*, *BubR1^{RNAi}*, *Mps1^{RNAi}*, and *CycA^{RNAi}*
oocytes with ROD^{GFP} in green, DNA in blue, CENP-C in white, and tubulin in red. Single
channel images (bottom) show ROD^{GFP}. (B) Quantification of ROD^{GFP} intensity at
kinetochores, normalized to background GFP signal in indicated oocytes (from left to
right, n = 142, 170, 66, 76, and 155 kinetochores). Error bars show mean ± s.d.;
****P<0.0001, *P≤0.0123 (unpaired two-tailed t test). (C) MPS1^{GFP} localization in wild-
type, *Bub3^{RNAi}*, *BubR1^{RNAi}*, *rod^{RNAi}*, and *CycA^{RNAi}* oocytes with MPS1^{GFP} in green, DNA
in blue, CENP-C in white, and tubulin in red. Single channel images (bottom) show
MPS1^{GFP}. (D) Quantification of MPS1^{GFP} intensity at kinetochores, normalized to
background GFP signal in indicated oocytes (from left to right, n = 455, 97, 149, 180,
and 152 kinetochores). Error bars show mean ± s.d.; ****P<0.0001 (unpaired two-tailed
t test). All images are maximum intensity projections of z stacks. Scale bars represent 5
μm.
717

Figure 5. **Meiotic chromosome alignment in SAC RNAi oocytes**.

(A) Representative images depicting chromosome orientation in wild-type, *rod^{RNAi}*, and
Mps1^{RNAi} oocytes. Images show DNA (blue), tubulin (green), the X chromosome
(magenta), the second chromosome (red), and the third chromosome (white). Tubulin
and FISH probes are also shown in a separate channel, with examples of mono-
oriented chromosomes circled. All images are maximum intensity projections of z
stacks. Scale bars represent 5 μm. (B) Percent of chromosomes per oocyte with the
indicated orientation (from left to right, n = 132, 65, 114, 94, 65, 66, 87, 39, and 30 pairs
of chromosomes). ****P<0.0001 (Fisher's exact test). (C) Examples of wild-type, *mad1^1*,
and *mad2^{EY21687}* mutant oocytes, with tubulin in green, DNA in blue, CENP-C in red, and
INCENP in white. Single channel images show DNA. (D) Percent of oocytes with the
specified karyosome morphology (from left to right, n=114, 102, 103, 41, 31, 32, and 31
oocytes). **P=0.0020, *P=0.0375 (Fisher's exact test).
731

Figure 6. **ROD streams in the absence of tension and end-on attachments**.

(A) ROD^{GFP} localization in wild-type, *mei-218^{null}*, *Ndc80^{RNAi}*, and *PP2A-B56^{RNAi}* oocytes
with ROD^{GFP} in green, DNA in blue, CENP-C in white, and tubulin in red. Single channel
images (bottom) show ROD^{GFP}. All images are maximum intensity projections of z
stacks. Scale bars represent 5 μm. (B) Percent of oocytes with ROD^{GFP} streaming in
indicated genotypes (from left to right, n = 25, 15, 22, and 16 oocytes), ns=no
significance (Fisher's exact test). (C) Quantification of ROD^{GFP} intensity at kinetochores,
normalized to background GFP signal in indicated oocytes (from left to right, n = 142,
139, and 92 kinetochores). Error bars show mean ± s.d.; *P=0.0202 (unpaired two-tailed
t test).
742

Figure 7. **ROD is not required for microtubule attachments**.

(A) Wild-type; *rod^RNAi*; *Ndc80^RNAi*; and *rod^RNAi*, *Ndc80^RNAi* oocytes with tubulin in green,
DNA in blue, CENP-C in white, and INCENP in red. *Rod^RNAi* oocyte on the left has an
intact karyosome and on the right a split karyosome. (B) Percent of kinetochores per
oocyte with lateral kinetochore-microtubule attachments in indicated genotypes (from
left to right, n = 40, 46, 30, 20, and 31 oocytes). Error bars show s.d.; ****P<0.0001,
***P=0.0007, ns=no significance (unpaired two-tailed t test). (C) Percent of oocytes with
the specified karyosome morphology in the indicated genotypes (from left to right, n =
41, 46, 22, 20, and 31 oocytes). ***P=0.0002, *P=0.0260 (Fisher's exact test). (D)
ROD^GFP localization in *Spc105R^B*, *Spc105R^ΔN*, and *Spc105R^ΔExxEED* oocytes with
ROD^GFP in green, DNA in blue, CENP-C in white, and tubulin in red. Single channel
images (bottom) show ROD^GFP. All *Spc105R* mutants are in an *Spc105R^RNAi*
background targeting the endogenous *Spc105R*. (E) Percent of oocytes with ROD^GFP
streaming in indicated genotypes (from left to right, n = 25, 20, 20, and 28 oocytes);
**P=0.0012 (Fisher's exact test). (F) *Spc105R^C* and *Zw10-Spc105R^C* oocytes (both in
an *Spc105R^RNAi* background) with DNA in blue, CENP-C in white, tubulin in red, and
ZW10-SPC105R^C in red (right). Single channel images (bottom) show CENP-C (left) or
ZW10-SPC105R^C (right). (G) Percent of kinetochores per oocyte with lateral
kinetochore-microtubule attachments in indicated genotypes (n = 42 and 51 oocytes).
Error bars show s.d.; ****P<0.0001 (unpaired two-tailed t test). All *Spc105R^C* and *Zw10-*
Spc105R^C oocytes are in an *Spc105R^RNAi* background targeting the endogenous
Spc105R. All images are maximum intensity projections of z stacks. Scale bars
represent 5 µm.
766

Figure 8. **Requirement for Spindly in meiosis and interactions with ROD.**

(A) Representative images depicting chromosome orientation in wild-type, *Spindly^RNAi*,
and *nudE^RNAi* oocytes. Images show DNA (blue), tubulin (green), the X chromosome
(magenta), the second chromosome (red), and the third chromosome (white). Tubulin
and FISH probes are also shown in a separate channel, with examples of mono-
oriented chromosomes circled. (B) Percent of chromosomes per oocyte with the
indicated orientation (from left to right, n = 86, 66, and 54 pairs of chromosomes);
****P<0.0001, **P=0.0076 (Fisher's exact test). (C) ROD^GFP localization in wild-type,
Spindly^RNAi, and *nudE^RNAi* oocytes with ROD^GFP in green, DNA in blue, CENP-C in white,
and tubulin in red. Single channel images (bottom) show ROD^GFP. Arrow points to
abnormal streaming in the *Spindly^RNAi* oocyte. See also Figure S 6A. (D) Percent of
oocytes with ROD^GFP streaming in indicated genotypes (from left to right, n = 25, 41,
and 10 oocytes); ns=no significance (Fisher's exact test). (E) Percent of oocytes that
abnormal streaming, defined as ROD^GFP in the central spindle, in wild-type or
Spindly^RNAi oocytes (n=24 and 43 oocytes); ***P=0.0002 (Fisher's exact test). Abnormal
streaming is defined as having ROD^GFP in the central region between the centromeres.
(F) Quantification of ROD^GFP intensity at kinetochores, normalized to background GFP
signal in indicated oocytes (from left to right, n = 142, 256, and 72 kinetochores. Error
bars show mean ± s.d.; ns=no significance (unpaired two-tailed t test). All images are
maximum intensity projections of z stacks. Scale bars represent 5 µm.
787

Figure S 1. **Validation of *rod* and *mps1* reagents.**

(A) Wild-type oocyte with ROD^{GFP} in green, ZW10^{HA} in red, DNA in blue, and tubulin in
white. Single channel images show ROD^{GFP} (middle) and ZW10^{HA} (right). (B) ROD^{GFP}
localization in wild-type and *rod*^{RNAi} oocytes with ROD^{GFP} in green, DNA in blue, CENP-
C in white, and tubulin in red. Single channel images (right) show ROD^{GFP}. (C)
Quantification of ROD^{GFP} intensity at kinetochores, normalized to background GFP
signal in wild-type and *rod*^{RNAi} oocytes (n = 142 and 126 kinetochores). Error bars show
mean ± s.d.; ****P<0.0001 (unpaired two-tailed t test). (D) MPS1^{GFP} localization in wild-
type and *Mps1*^{RNAi} oocytes with MPS1^{GFP} in green, DNA in blue, CENP-C in white, and
tubulin in red. Single channel images (right) show MPS1^{GFP}. (E) Quantification of
MPS1^{GFP} intensity at kinetochores, normalized to background GFP signal in wild-type
and *Mps1*^{RNAi} oocytes (n = 75 and 71 kinetochores). Error bars show mean ± s.d.;
****P<0.0001 (unpaired two-tailed t test). All images are maximum intensity projections
of z stacks. Scale bars represent 5 µm.
803
Figure S 2. **MAD1 and INCENP localization depends on SPC105R.**

Control (*Spc105R*^{B}) or mutant oocytes were incubated for one hour in 250 µM
colchicine. (A) MAD1^{GFP} (green) localization in indicated genotypes, with DNA in blue,
CENP-C in white, and tubulin in red. Single channel images (bottom) show MAD1^{GFP}.
(B) Quantification of MAD1^{GFP} intensity at kinetochores, normalized to background GFP
signal (from left to right, n = 91, 256, 224, 174, 212, 208, and 236 kinetochores). Error
bars show mean ± s.d.; ****P<0.0001, ***P=0.0006 (unpaired two-tailed t test). (C)
INCENP localization (green) in indicated genotypes with DNA in blue, CENP-C in white,
and tubulin in red. Single channel images (bottom) show INCENP. (D) Quantification of
INCENP presence at kinetochores (from left to right, n = 16, 29, 11, 31, 20, 18, 21, and
11 oocytes). Error bars show mean ± s.d.; ****P<0.0001, *P=0.02 (Fisher's exact test).
All images are maximum intensity projections of z stacks. Scale bars represent 5 µm. All
Spc105R mutants are in an *Spc105R*^{RNAi} background targeting the endogenous
Spc105R.
818
Figure S 3. **Structure of SPC105R and mutant variants.**

A schematic of the *Spc105R* mutants used in this study. The coordinates on the
schematic represent the first amino acid of each domain. The N-terminal includes SLRK
and RISF. Following this is a domain with three MELT-like motifs, a region that contains
two KI-like repeats, a central domain containing repeats with the consensus ExxEED,
and the C-terminal region containing coiled-coil motifs.
825
Figure S 4. **MPS1 localization depends on SPC105R and NDC80.**

(A) MPS1^{GFP} localization in wild-type, *Spc105R*^{RNAi}, and *Ndc80*^{RNAi} oocytes, with
MPS1^{GFP} in green, DNA in blue, CENP-C in white, and tubulin in red. Single channel
images (bottom) show MPS1^{GFP}. (B) Quantification of MPS1^{GFP} intensity at
kinetochores, normalized to background GFP signal in indicated oocytes (from left to

right, n = 389, 195, 139, 160, 165, 198, 128, and 143 kinetochores). Error bars show
mean ± s.d.; ****P<0.0001 (unpaired two-tailed t test). (C) Quantification of MPS1^{GFP}
intensity at kinetochores, normalized to background GFP signal in wild-type and
Ndc80^{RNAi} oocytes (n=114 and 148 kinetochores). Error bars show mean ± s.d.;
****P<0.0001 (unpaired two-tailed t test). (D) MPS1^{GFP} localization in the indicated
Spc105R mutants with MPS1^{GFP} in green, DNA in blue, CENP-C in white, and tubulin in
red. All mutants are in an *Spc105R^{RNAi}* background targeting the endogenous *Spc105R*.
Single channel images (bottom) show MPS1^{GFP}. All images are maximum intensity
projections of z stacks. Scale bars represent 5 µm.
840

Figure S 5. **ROD and SPC105R localization in *Zw10-Spc105R^C* oocytes.**

(A) SPC105R localization in *Zw10-Spc105R^C*, either in the presence or absence of
Spc105R^{RNAi}. SPC105R (green) was detected using an antibody which recognizes the
N-terminal regions of SPC105R and does not detect SPC105R^C. DNA is in blue, CENP-
C in white, and tubulin in red. All images are maximum intensity projections of z stacks.
(B) ROD^{GFP} localization in oocytes expressing *Zw10-Spc105R^C,* either in the presence
or absence of *Spc105R^{RNAi}*. ROD^{GFP} is in green, DNA in blue, CENP-C in white, and
tubulin in red. Scale bars represent 5 µm.
849

Figure S 6. **Localization of Spindly variants and dependence of Spindly on ROD.**

(A) Examples of normal (left) and abnormal (right) ROD^{GFP} streaming in *Spindly^{RNAi}*
oocytes, with ROD^{GFP} in green, DNA in blue, CENP-C in white, and tubulin in red.
Single channel images (bottom) show ROD^{GFP}. Arrow points to abnormal streaming,
which is defined as having ROD^{GFP} in the central region between the centromeres. (B)
Spindly^{GFP} localization in wild-type or mutants of *Spindly*. Spindly^{GFP} is in green, DNA in
blue, CENP-C in white, and tubulin in red. (C) Localization of Spindly$^{\Delta SB.GFP}$ in *rod^{RNAi}*
oocytes, with Spindly^{GFP} in green, DNA in blue, CENP-C in white, and tubulin in red.
Spindly^{GFP} and CENP-C are shown below the merged images. All images are maximum
intensity projections of z stacks. Scale bars represent 5 µm. (D) Quantification of
Spindly^{GFP} intensity at kinetochores in the indicated oocytes (from left to right, n = 123,
164, 83, and 75 kinetochores). Error bars show mean ± s.d.; ****P<0.0001 (unpaired
two-tailed t test).
863

References

1. McAinsh AD, Kops G. Principles and dynamics of spindle assembly checkpoint signalling. Nat Rev Mol Cell Biol. 2023;24(8):543-59. Epub 20230324. doi: 10.1038/s41580-023-00593-z. PubMed PMID: 36964313.
2. Lara-Gonzalez P, Pines J, Desai A. Spindle assembly checkpoint activation and silencing at kinetochores. Seminars in cell & developmental biology. 2021;117:86-98. Epub 20210629. doi: 10.1016/j.semcdb.2021.06.009. PubMed PMID: 34210579; PubMed Central PMCID: PMCPMC8406419.
3. Krenn V, Musacchio A. The Aurora B Kinase in Chromosome Bi-Orientation and Spindle Checkpoint Signaling. Front Oncol. 2015;5:225. Epub 2015/11/04. doi: 10.3389/fonc.2015.00225. PubMed PMID: 26528436; PubMed Central PMCID: PMCPMC4607871.
4. Barbosa J, Sunkel CE, Conde C. The Role of Mitotic Kinases and the RZZ Complex in Kinetochore-Microtubule Attachments: Doing the Right Link. Front Cell Dev Biol. 2022;10:787294. Epub 20220128. doi: 10.3389/fcell.2022.787294. PubMed PMID: 35155423; PubMed Central PMCID: PMCPMC8832123.
5. Lampson MA, Grishchuk EL. Mechanisms to Avoid and Correct Erroneous Kinetochore-Microtubule Attachments. Biology (Basel). 2017;6(1). Epub 2017/01/10. doi: 10.3390/biology6010001. PubMed PMID: 28067761; PubMed Central PMCID: PMCPMC5371994.
6. Kitajima TS. Mechanisms of kinetochore-microtubule attachment errors in mammalian oocytes. Dev Growth Differ. 2018;60(1):33-43. Epub 2018/01/11. doi: 10.1111/dgd.12410. PubMed PMID: 29318599.
7. Mihajlovic AI, FitzHarris G. Segregating Chromosomes in the Mammalian Oocyte. Curr Biol. 2018;28(16):R895-R907. Epub 2018/08/22. doi: 10.1016/j.cub.2018.06.057. PubMed PMID: 30130515.
8. Webster A, Schuh M. Mechanisms of Aneuploidy in Human Eggs. Trends Cell Biol. 2017;27(1):55-68. Epub 2016/10/25. doi: 10.1016/j.tcb.2016.09.002. PubMed PMID: 27773484.
9. Charalambous C, Webster A, Schuh M. Aneuploidy in mammalian oocytes and the impact of maternal ageing. Nat Rev Mol Cell Biol. 2023;24(1):27-44. Epub 20220906. doi: 10.1038/s41580-022-00517-3. PubMed PMID: 36068367.
10. Touati SA, Wassmann K. How oocytes try to get it right: spindle checkpoint control in meiosis. Chromosoma. 2016;125(2):321-35. Epub 20150811. doi: 10.1007/s00412-015-0536-7. PubMed PMID: 26255654.
11. Wu T, Gu H, Luo Y, Wang L, Sang Q. Meiotic defects in human oocytes: Potential causes and clinical implications. Bioessays. 2022;44(12):e2200135. Epub 20221007. doi: 10.1002/bies.202200135. PubMed PMID: 36207289.
12. Liu D, Shao H, Wang H, Liu XJ. Meiosis I in Xenopus oocytes is not error-prone despite lacking spindle assembly checkpoint. Cell Cycle. 2014;13(10):1602-6. Epub 20140319. doi: 10.4161/cc.28562. PubMed PMID: 24646611; PubMed Central PMCID: PMCPMC4050165.
13. Shao H, Li R, Ma C, Chen E, Liu XJ. Xenopus oocyte meiosis lacks spindle assembly checkpoint control. J Cell Biol. 2013;201(2):191-200. Epub 20130408. doi: 10.1083/jcb.201211041. PubMed PMID: 23569212; PubMed Central PMCID: PMCPMC3628510.
14. Gui L, Homer H. Spindle assembly checkpoint signalling is uncoupled from chromosomal position in mouse oocytes. Development. 2012;139(11):1941-6. PubMed PMID: 22513372.

15.   Lane SI, Yun Y, Jones KT. Timing of anaphase-promoting complex activation in mouse
oocytes is predicted by microtubule-kinetochore attachment but not by bivalent alignment or
tension. Development. 2012;139(11):1947-55. PubMed PMID: 22513370.
16.   Kyogoku H, Kitajima TS. The large cytoplasmic volume of oocyte. J Reprod Dev.
2023;69(1):1-9. Epub 20221126. doi: 10.1262/jrd.2022-101. PubMed PMID: 36436912; PubMed
Central PMCID: PMCPMC9939283.
17.   Radford SJ, Nguyen AL, Schindler K, McKim KS. The chromosomal basis of meiotic
acentrosomal spindle assembly and function in oocytes. Chromosoma. 2017;126(3):351-64.
Epub 2016/11/12. doi: 10.1007/s00412-016-0618-1. PubMed PMID: 27837282; PubMed Central
PMCID: PMCPMC5426991.
18.   Blengini CS, Schindler K. Acentriolar spindle assembly in mammalian female meiosis
and the consequences of its perturbations on human reproduction†. Biol Reprod.
2022;106(2):253-63. doi: 10.1093/biolre/ioab210. PubMed PMID: 34791041; PubMed Central
PMCID: PMCPMC8862719.
19.   Wartosch L, Schindler K, Schuh M, Gruhn JR, Hoffmann ER, McCoy RC, et al. Origins
and mechanisms leading to aneuploidy in human eggs. Prenat Diagn. 2021;41(5):620-30. Epub
20210322. doi: 10.1002/pd.5927. PubMed PMID: 33860956; PubMed Central PMCID:
PMCPMC8237340.
20.   Singleton MR. Getting to the heart of an unusual kinetochore. Open Biol.
2016;6(4):160040. doi: 10.1098/rsob.160040. PubMed PMID: 27249344; PubMed Central
PMCID: PMCPMC4852463.
21.   Radford SJ, Hoang TL, Głuszek AA, Ohkura H, McKim KS. Lateral and End-On
Kinetochore Attachments Are Coordinated to Achieve Bi-orientation in Drosophila Oocytes.
PLoS Genet. 2015;11(10):e1005605. doi: 10.1371/journal.pgen.1005605. PubMed PMID:
26473960.
22.   Joshi JN, Changela N, Mahal L, Jang J, Defosse T, Wang LI, et al. Meiosis-specific
functions of kinetochore protein SPC105R required for chromosome segregation in Drosophila
oocytes. Mol Biol Cell. 2024:mbcE24020067. Epub 20240612. doi: 10.1091/mbc.E24-02-0067.
PubMed PMID: 38865189.
23.   Cheerambathur DK, Gassmann R, Cook B, Oegema K, Desai A. Crosstalk between
microtubule attachment complexes ensures accurate chromosome segregation. Science.
2013;342(6163):1239-42. Epub 2013/11/16. doi: 10.1126/science.1246232. PubMed PMID:
24231804; PubMed Central PMCID: PMCPMC3885540.
24.   Barbosa J, Martins T, Bange T, Tao L, Conde C, Sunkel C. Polo regulates Spindly to
prevent premature stabilization of kinetochore-microtubule attachments. EMBO J.
2020;39(2):e100789. Epub 2019/12/19. doi: 10.15252/embj.2018100789. PubMed PMID:
31849090; PubMed Central PMCID: PMCPMC6960449.
25.   Basto R, Scaerou F, Mische S, Wojcik E, Lefebvre C, Gomes R, et al. In vivo dynamics
of the rough deal checkpoint protein during Drosophila mitosis. Curr Biol. 2004;14(1):56-61.
PubMed PMID: 14711415.
26.   Williams BC, Gatti M, Goldberg ML. Bipolar spindle attachments affect redistributions of
ZW10, a Drosophila centromere/kinetochore component required for accurate chromosome
segregation. J Cell Biol. 1996;134(5):1127-40. doi: 10.1083/jcb.134.5.1127. PubMed PMID:
8794856; PubMed Central PMCID: PMCPMC2120981.
27.   Scaërou F, Starr DA, Piano F, Papoulas O, Karess RE, Goldberg ML. The ZW10 and
Rough Deal checkpoint proteins function together in a large, evolutionarily conserved complex
targeted to the kinetochore. J Cell Sci. 2001;114(Pt 17):3103-14. doi: 10.1242/jcs.114.17.3103.
PubMed PMID: 11590237.
28.   Kops G, Gassmann R. Crowning the Kinetochore: The Fibrous Corona in Chromosome
Segregation. Trends Cell Biol. 2020;30(8):653-67. Epub 2020/05/11. doi:
10.1016/j.tcb.2020.04.006. PubMed PMID: 32386879.

964 29. Pereira C, Reis RM, Gama JB, Celestino R, Cheerambathur DK, Carvalho AX, et al.
965 Self-Assembly of the RZZ Complex into Filaments Drives Kinetochore Expansion in the
966 Absence of Microtubule Attachment. Current biology : CB. 2018;28(21):3408-21.e8. Epub
967 2018/11/13. doi: 10.1016/j.cub.2018.08.056. PubMed PMID: 30415699; PubMed Central
968 PMCID: PMCPMC6224608.
969 30. Marston AL, Wassmann K. Multiple Duties for Spindle Assembly Checkpoint Kinases in
970 Meiosis. Front Cell Dev Biol. 2017;5:109. Epub 20171213. doi: 10.3389/fcell.2017.00109.
971 PubMed PMID: 29322045; PubMed Central PMCID: PMCPMC5733479.
972 31. Wang LI, DeFosse T, Jang JK, Battaglia RA, Wagner VF, McKim KS. Borealin directs
973 recruitment of the CPC to oocyte chromosomes and movement to the microtubules. J Cell Biol.
974 2021;220(6). Epub 2021/04/10. doi: 10.1083/jcb.202006018. PubMed PMID: 33836043.
975 32. Radford SJ, Jang JK, McKim KS. The Chromosomal Passenger Complex is required for
976 Meiotic Acentrosomal Spindle Assembly and Chromosome Bi-orientation. Genetics.
977 2012;192:417-29. PubMed PMID: 22865736.
978 33. McKim KS, Jang JK, Theurkauf WE, Hawley RS. Mechanical basis of meiotic
979 metaphase arrest. Nature. 1993;362(6418):364-6. PubMed PMID: 8455723.
980 34. Saurin AT. Kinase and Phosphatase Cross-Talk at the Kinetochore. Front Cell Dev Biol.
981 2018;6:62. Epub 2018/07/05. doi: 10.3389/fcell.2018.00062. PubMed PMID: 29971233;
982 PubMed Central PMCID: PMCPMC6018199.
983 35. Caldas GV, DeLuca JG. KNL1: bringing order to the kinetochore. Chromosoma.
984 2014;123(3):169-81. doi: 10.1007/s00412-013-0446-5. PubMed PMID: 24310619; PubMed
985 Central PMCID: PMCPMC4032621.
986 36. Hayward D, Alfonso-Pérez T, Cundell MJ, Hopkins M, Holder J, Bancroft J, et al. CDK1-
987 CCNB1 creates a spindle checkpoint-permissive state by enabling MPS1 kinetochore
988 localization. J Cell Biol. 2019;218(4):1182-99. Epub 20190123. doi: 10.1083/jcb.201808014.
989 PubMed PMID: 30674582; PubMed Central PMCID: PMCPMC6446832.
990 37. Gilliland WD, Hughes SE, Cotitta JL, Takeo S, Xiang Y, Hawley RS. The multiple roles
991 of mps1 in Drosophila female meiosis. PLoS Genet. 2007;3(7):e113. doi:
992 10.1371/journal.pgen.0030113. PubMed PMID: 17630834; PubMed Central PMCID:
993 PMCPMC1914070.
994 38. Gilliland WD, Wayson SM, Hawley RS. The meiotic defects of mutants in the Drosophila
995 mps1 gene reveal a critical role of Mps1 in the segregation of achiasmate homologs. Curr Biol.
996 2005;15(7):672-7. doi: 10.1016/j.cub.2005.02.062. PubMed PMID: 15823541.
997 39. Bourouh M, Dhaliwal R, Rana K, Sinha S, Guo Z, Swan A. Distinct and Overlapping
998 Requirements for Cyclins A, B and B3 in Drosophila Female Meiosis. G3 (Bethesda, Md. 2016.
999 doi: 10.1534/g3.116.033050. PubMed PMID: 27652889; PubMed Central PMCID:
1000 PMCPMC5100870.
1001 40. Malmanche N, Owen S, Gegick S, Steffensen S, Tomkiel JE, Sunkel CE. Drosophila
1002 BubR1 is essential for meiotic sister-chromatid cohesion and maintenance of synaptonemal
1003 complex. Curr Biol. 2007;17(17):1489-97. PubMed PMID: 17702574.
1004 41. Buffin E, Emre D, Karess RE. Flies without a spindle checkpoint. Nat Cell Biol.
1005 2007;9(5):565-72. doi: 10.1038/ncb1570. PubMed PMID: 17417628.
1006 42. Emre D, Terracol R, Poncet A, Rahmani Z, Karess RE. A mitotic role for Mad1 beyond
1007 the spindle checkpoint. J Cell Sci. 2011;124(Pt 10):1664-71. Epub 20110421. doi:
1008 10.1242/jcs.081216. PubMed PMID: 21511728.
1009 43. Hughes SE, Miller DE, Miller AL, Hawley RS. Female Meiosis: Synapsis,
1010 Recombination, and Segregation in Drosophila melanogaster. Genetics. 2018;208(3):875-908.
1011 Epub 2018/03/01. doi: 10.1534/genetics.117.300081. PubMed PMID: 29487146; PubMed
1012 Central PMCID: PMCPMC5844340.

44.   Theurkauf WE, Hawley RS. Meiotic spindle assembly in Drosophila females: behavior of
nonexchange chromosomes and the effects of mutations in the nod kinesin-like protein. J Cell
Biol. 1992;116(5):1167-80. PubMed PMID: 1740471.
45.   Costa MFA, Ohkura H. The molecular architecture of the meiotic spindle is remodeled
during metaphase arrest in oocytes. J Cell Biol. 2019;218(9):2854-64. Epub 2019/07/07. doi:
10.1083/jcb.201902110. PubMed PMID: 31278080; PubMed Central PMCID:
PMCPMC6719438.
46.   Carpenter AT, Sandler L. On recombination-defective meiotic mutants in Drosophila
melanogaster. Genetics. 1974;76(3):453-75. PubMed PMID: 4208856.
47.   Jang JK, Gladstein AC, Das A, Shapiro JG, Sisco ZL, McKim KS. Multiple pools of PP2A
regulate spindle assembly, kinetochore attachments, and cohesion in Drosophila oocytes. J Cell
Sci. 2021;134(14):jcs254037. Epub 2021/06/24. doi: 10.1242/jcs.254037. PubMed PMID:
34160620.
48.   Wu J, Larreategui-Aparicio A, Lambers MLA, Bodor DL, Klaasen SJ, Tollenaar E, et al.
Microtubule nucleation from the fibrous corona by LIC1-pericentrin promotes chromosome
congression. Curr Biol. 2023;33(5):912-25 e6. Epub 20230130. doi: 10.1016/j.cub.2023.01.010.
PubMed PMID: 36720222; PubMed Central PMCID: PMCPMC10017265.
49.   Menant A, Karess RE. Mutations in the Drosophila rough deal gene affecting RZZ
kinetochore function. Biol Cell. 2020;112(10):300-15. Epub 20200720. doi:
10.1111/boc.201900105. PubMed PMID: 32602944.
50.   Wojcik E, Basto R, Serr M, Scaërou F, Karess R, Hays T. Kinetochore dynein: its
dynamics and role in the transport of the Rough deal checkpoint protein. Nature cell biology.
2001;3(11):1001-7. Epub 2001/11/21. doi: 10.1038/ncb1101-1001. PubMed PMID: 11715021.
51.   Griffis ER, Stuurman N, Vale RD. Spindly, a novel protein essential for silencing the
spindle assembly checkpoint, recruits dynein to the kinetochore. J Cell Biol. 2007;177(6):1005-
15. doi: 10.1083/jcb.200702062. PubMed PMID: 17576797; PubMed Central PMCID:
PMCPMC2064361.
52.   Wainman A, Creque J, Williams B, Williams EV, Bonaccorsi S, Gatti M, et al. Roles of
the Drosophila NudE protein in kinetochore function and centrosome migration. J Cell Sci.
2009;122(Pt 11):1747-58. Epub 2009/05/07. doi: 10.1242/jcs.041798. PubMed PMID:
19417004; PubMed Central PMCID: PMCPMC2684831.
53.   Gassmann R, Essex A, Hu JS, Maddox PS, Motegi F, Sugimoto A, et al. A new
mechanism controlling kinetochore-microtubule interactions revealed by comparison of two
dynein-targeting components: SPDL-1 and the Rod/Zwilch/Zw10 complex. Genes Dev.
2008;22(17):2385-99. doi: 10.1101/gad.1687508. PubMed PMID: 18765790; PubMed Central
PMCID: PMCPMC2532926.
54.   Clemente GD, Hannaford MR, Beati H, Kapp K, Januschke J, Griffis ER, et al.
Requirement of the Dynein-Adaptor Spindly for Mitotic and Post-Mitotic Functions in Drosophila.
J Dev Biol. 2018;6(2). Epub 2018/04/05. doi: 10.3390/jdb6020009. PubMed PMID: 29615558;
PubMed Central PMCID: PMCPMC6027351.
55.   Davis-Roca AC, Muscat CC, Wignall SM. Caenorhabditis elegans oocytes detect meiotic
errors in the absence of canonical end-on kinetochore attachments. J Cell Biol.
2017;216(5):1243-53. Epub 20170329. doi: 10.1083/jcb.201608042. PubMed PMID: 28356326;
PubMed Central PMCID: PMCPMC5412562.
56.   Jones KT, Lane SI. Molecular causes of aneuploidy in mammalian eggs. Development.
2013;140(18):3719-30. doi: 10.1242/dev.090589. PubMed PMID: 23981655.
57.   Touati SA, Buffin E, Cladiere D, Hached K, Rachez C, van Deursen JM, et al. Mouse
oocytes depend on BubR1 for proper chromosome segregation but not for prophase I arrest.
Nat Commun. 2015;6:6946. Epub 2015/04/22. doi: 10.1038/ncomms7946. PubMed PMID:
25897860; PubMed Central PMCID: PMCPMC4439927.

58.   Qiao JY, Zhou Q, Xu K, Yue W, Lei WL, Li YY, et al. Mad2 is dispensable for accurate
chromosome segregation but becomes essential when oocytes are subjected to environmental
stress. Development. 2023;150(14). Epub 20230724. doi: 10.1242/dev.201398. PubMed PMID:
37485540.
59.   Batiha O, Swan A. Evidence that the spindle assembly checkpoint does not regulate
APC(Fzy) activity in Drosophila female meiosis. Genome. 2012;55(1):63-7. Epub 20111223.
doi: 10.1139/g11-079. PubMed PMID: 22196012.
60.   Jang JK, Messina L, Erdman MB, Arbel T, Hawley RS. Induction of metaphase arrest in
Drosophila oocytes by chiasma-based kinetochore tension. Science. 1995;268(5219):1917-9.
61.   !!! INVALID CITATION !!! [22, 46].
62.   !!! INVALID CITATION !!! [31, 58].
63.   !!! INVALID CITATION !!! [60].
64.   Silió V, McAinsh AD, Millar JB. KNL1-Bubs and RZZ Provide Two Separable Pathways
for Checkpoint Activation at Human Kinetochores. Dev Cell. 2015;35(5):600-13. Epub
2015/12/15. doi: 10.1016/j.devcel.2015.11.012. PubMed PMID: 26651294.
65.   Pauleau AL, Bergner A, Kajtez J, Erhardt S. The checkpoint protein Zw10 connects
CAL1-dependent CENP-A centromeric loading and mitosis duration in Drosophila cells. PLoS
Genet. 2019;15(9):e1008380. Epub 20190925. doi: 10.1371/journal.pgen.1008380. PubMed
PMID: 31553715; PubMed Central PMCID: PMCPMC6779278.
66.   O'Tousa J. Meiotic chromosome behavior influenced by mutation-altered disjunction in
Drosophila melanogaster females. Genetics. 1982;102:503-24.
67.   Hached K, Xie SZ, Buffin E, Cladiere D, Rachez C, Sacras M, et al. Mps1 at
kinetochores is essential for female mouse meiosis I. Development. 2011;138(11):2261-71.
PubMed PMID: 21558374.
68.   Sarangapani KK, Koch LB, Nelson CR, Asbury CL, Biggins S. Kinetochore-bound Mps1
regulates kinetochore-microtubule attachments via Ndc80 phosphorylation. J Cell Biol.
2021;220(12). Epub 20211014. doi: 10.1083/jcb.202106130. PubMed PMID: 34647959;
PubMed Central PMCID: PMCPMC8641409.
69.   Hiruma Y, Sacristan C, Pachis ST, Adamopoulos A, Kuijt T, Ubbink M, et al. CELL
DIVISION CYCLE. Competition between MPS1 and microtubules at kinetochores regulates
spindle checkpoint signaling. Science. 2015;348(6240):1264-7. Epub 20150611. doi:
10.1126/science.aaa4055. PubMed PMID: 26068855.
70.   Ji Z, Gao H, Yu H. CELL DIVISION CYCLE. Kinetochore attachment sensed by
competitive Mps1 and microtubule binding to Ndc80C. Science. 2015;348(6240):1260-4. doi:
10.1126/science.aaa4029. PubMed PMID: 26068854.
71.   Kemmler S, Stach M, Knapp M, Ortiz J, Pfannstiel J, Ruppert T, et al. Mimicking Ndc80
phosphorylation triggers spindle assembly checkpoint signalling. EMBO J. 2009;28(8):1099-
110. Epub 20090319. doi: 10.1038/emboj.2009.62. PubMed PMID: 19300438; PubMed Central
PMCID: PMCPMC2683709.
72.   Hayward D, Roberts E, Gruneberg U. MPS1 localizes to end-on microtubule-attached
kinetochores to promote microtubule release. Curr Biol. 2022;32(23):5200-8 e8. Epub
20221116. doi: 10.1016/j.cub.2022.10.047. PubMed PMID: 36395767.
73.   Buffin E, Lefebvre C, Huang J, Gagou ME, Karess RE. Recruitment of Mad2 to the
kinetochore requires the Rod/Zw10 complex. Curr Biol. 2005;15(9):856-61. PubMed PMID:
15886105.
74.   Kops GJ, Kim Y, Weaver BA, Mao Y, McLeod I, Yates JR, 3rd, et al. ZW10 links mitotic
checkpoint signaling to the structural kinetochore. J Cell Biol. 2005;169(1):49-60. doi:
10.1083/jcb.200411118. PubMed PMID: 15824131; PubMed Central PMCID:
PMCPMC1351127.

75.   Tromer E, Snel B, Kops GJ. Widespread Recurrent Patterns of Rapid Repeat Evolution
in the Kinetochore Scaffold KNL1. Genome Biol Evol. 2015;7(8):2383-93. Epub 2015/08/09. doi:
10.1093/gbe/evv140. PubMed PMID: 26254484; PubMed Central PMCID: PMCPMC4558858.
76.   Caldas GV, Lynch TR, Anderson R, Afreen S, Varma D, DeLuca JG. The RZZ complex
requires the N-terminus of KNL1 to mediate optimal Mad1 kinetochore localization in human
cells. Open Biol. 2015;5(11). doi: 10.1098/rsob.150160. PubMed PMID: 26581576; PubMed
Central PMCID: PMCPMC4680571.
77.   Zhang G, Lischetti T, Hayward DG, Nilsson J. Distinct domains in Bub1 localize RZZ and
BubR1 to kinetochores to regulate the checkpoint. Nat Commun. 2015;6:7162. Epub 20150602.
doi: 10.1038/ncomms8162. PubMed PMID: 26031201; PubMed Central PMCID:
PMCPMC4458899.
78.   McGory JM, Verma V, Barcelos DM, Maresca TJ. Multimerization of a disordered
kinetochore protein promotes accurate chromosome segregation by localizing a core dynein
module. J Cell Biol. 2024;223(3). Epub 20240105. doi: 10.1083/jcb.202211122. PubMed PMID:
38180477; PubMed Central PMCID: PMCPMC10770731.
79.   Davydenko O, Schultz RM, Lampson MA. Increased CDK1 activity determines the timing
of kinetochore-microtubule attachments in meiosis I. J Cell Biol. 2013;202(2):221-9. doi:
10.1083/jcb.201303019. PubMed PMID: 23857768; PubMed Central PMCID:
PMCPMC3718970.
80.   Kabeche L, Compton DA. Cyclin A regulates kinetochore microtubules to promote
faithful chromosome segregation. Nature. 2013;502(7469):110-3. doi: 10.1038/nature12507.
PubMed PMID: 24013174; PubMed Central PMCID: PMCPMC3791168.
81.   Amin MA, McKenney RJ, Varma D. Antagonism between the dynein and Ndc80
complexes at kinetochores controls the stability of kinetochore-microtubule attachments during
mitosis. J Biol Chem. 2018;293(16):5755-65. Epub 2018/02/25. doi: 10.1074/jbc.RA117.001699.
PubMed PMID: 29475948; PubMed Central PMCID: PMCPMC5912454.
82.   Sacristan C, Ahmad MUD, Keller J, Fermie J, Groenewold V, Tromer E, et al. Dynamic
kinetochore size regulation promotes microtubule capture and chromosome biorientation in
mitosis. Nat Cell Biol. 2018;20(7):800-10. Epub 2018/06/20. doi: 10.1038/s41556-018-0130-3.
PubMed PMID: 29915359; PubMed Central PMCID: PMCPMC6485389.
83.   Wang LI, Das A, McKim KS. Sister centromere fusion during meiosis I depends on
maintaining cohesins and destabilizing microtubule attachments. PLoS Genet.
2019;15(5):e1008072. Epub 2019/06/01. doi: 10.1371/journal.pgen.1008072. PubMed PMID:
31150390; PubMed Central PMCID: PMCPMC6581285.
84.   Barbosa J, Conde C, Sunkel C. RZZ-SPINDLY-DYNEIN: you got to keep 'em separated.
Cell Cycle. 2020;19(14):1716-26. Epub 20200616. doi: 10.1080/15384101.2020.1780382.
PubMed PMID: 32544383; PubMed Central PMCID: PMCPMC7469663.
85.   Barisic M, Sohm B, Mikolcevic P, Wandke C, Rauch V, Ringer T, et al. Spindly/CCDC99
is required for efficient chromosome congression and mitotic checkpoint regulation. Mol Biol
Cell. 2010;21(12):1968-81. Epub 2010/04/30. doi: 10.1091/mbc.E09-04-0356. PubMed PMID:
20427577; PubMed Central PMCID: PMCPMC2883941.
86.   Gassmann R, Holland AJ, Varma D, Wan X, Civril F, Cleveland DW, et al. Removal of
Spindly from microtubule-attached kinetochores controls spindle checkpoint silencing in human
cells. Genes & development. 2010;24(9):957-71. Epub 2010/05/05. doi: 10.1101/gad.1886810.
PubMed PMID: 20439434; PubMed Central PMCID: PMCPMC2861194.
87.   Wu J, Raas MWD, Alcaraz PS, Vos HR, Tromer EC, Snel B, et al. A farnesyl-dependent
structural role for CENP-E in expansion of the fibrous corona. J Cell Biol. 2024;223(1). Epub
20231107. doi: 10.1083/jcb.202303007. PubMed PMID: 37934467; PubMed Central PMCID:
PMCPMC10630089.
88.   McGory JM, Verma V, Barcelos DM, Maresca TJ. Multimerization of a disordered
kinetochore protein promotes accurate chromosome segregation by localizing a core dynein

module. J Cell Biol. 2024;223(3):e202211122. Epub 20240105. doi: 10.1083/jcb.202211122. PubMed PMID: 38180477; PubMed Central PMCID: PMCPMC10770731.

89. Cmentowski V, Ciossani G, d'Amico E, Wohlgemuth S, Owa M, Dynlacht B, et al. RZZ-Spindly and CENP-E form an integrated platform to recruit dynein to the kinetochore corona. Embo j. 2023;42(24):e114838. Epub 20231120. doi: 10.15252/embj.2023114838. PubMed PMID: 37984321; PubMed Central PMCID: PMCPMC10711656.

90. Althoff F, Karess RE, Lehner CF. Spindle checkpoint-independent inhibition of mitotic chromosome segregation by Drosophila Mps1. Mol Biol Cell. 2012;23(12):2275-91. Epub 20120502. doi: 10.1091/mbc.E12-02-0117. PubMed PMID: 22553353; PubMed Central PMCID: PMCPMC3374747.

91. Fischer MG, Heeger S, Hacker U, Lehner CF. The mitotic arrest in response to hypoxia and of polar bodies during early embryogenesis requires Drosophila Mps1. Curr Biol. 2004;14(22):2019-24. doi: 10.1016/j.cub.2004.11.008. PubMed PMID: 15556864.

92. Zeng Y, Li H, Schweppe NM, Hawley RS, Gilliland WD. Statistical analysis of nondisjunction assays in Drosophila. Genetics. 2010;186(2):505-13. doi: 10.1534/genetics.110.118778. PubMed PMID: 20660647; PubMed Central PMCID: PMCPMC2954469.

93. Défachelles L, Hainline SG, Menant A, Lee LA, Karess RE. A maternal effect rough deal mutation suggests that multiple pathways regulate Drosophila RZZ kinetochore recruitment. J Cell Sci. 2015;128(6):1204-16. Epub 20150122. doi: 10.1242/jcs.165712. PubMed PMID: 25616898; PubMed Central PMCID: PMCPMC4359924.

94. Radford SJ, McKim KS. Techniques for Imaging Prometaphase and Metaphase of Meiosis I in Fixed Drosophila Oocytes. Journal of visualized experiments : JoVE. 2016;116(116):e54666. doi: 10.3791/54666. PubMed PMID: 27842371.

95. Chu DT, Klymkowsky MW. The appearance of acetylated alpha-tubulin during early development and cellular differentiation in Xenopus. Dev Biol. 1989;136(1):104-17. doi: 10.1016/0012-1606(89)90134-6. PubMed PMID: 2680681.

96. Wu C, Singaram V, McKim KS. mei-38 is required for chromosome segregation during meiosis in Drosophila females. Genetics. 2008;180(1):61-72. PubMed PMID: 18757915.

97. Fellmeth JE, Jang JK, Persaud M, Sturm H, Changela N, Parikh A, et al. A dynamic population of prophase CENP-C is required for meiotic chromosome segregation. PLoS Genet. 2023;19(11):e1011066. Epub 20231129. doi: 10.1371/journal.pgen.1011066. PubMed PMID: 38019881.

98. Schittenhelm RB, Chaleckis R, Lehner CF. Intrakinetochore localization and essential functional domains of Drosophila Spc105. EMBO J. 2009;28(16):2374-86. doi: 10.1038/emboj.2009.188. PubMed PMID: 19590494; PubMed Central PMCID: PMCPMC2735171.

99. Sugimura I, Lilly MA. Bruno inhibits the expression of mitotic cyclins during the prophase I meiotic arrest of Drosophila oocytes. Dev Cell. 2006;10(1):127-35. PubMed PMID: 16399084.

100. Schittenhelm RB, Heeger S, Althoff F, Walter A, Heidmann S, Mechtler K, et al. Spatial organization of a ubiquitous eukaryotic kinetochore protein network in Drosophila chromosomes. Chromosoma. 2007;116(4):385-402. PubMed PMID: 17333235.

101. Ni JQ, Zhou R, Czech B, Liu LP, Holderbaum L, Yang-Zhou D, et al. A genome-scale shRNA resource for transgenic RNAi in Drosophila. Nature methods. 2011;8(5):405-7. PubMed PMID: 21460824.

102. Liu H, Jang JK, Graham J, Nycz K, McKim KS. Two genes required for meiotic recombination in Drosophila are expressed from a dicistronic message. Genetics. 2000;154(4):1735-46.

1212   Figure 1

1213

1214
1215

1216   Figure 2

1217

1218

1219   Figure 3

1220

1221

1222   Figure 4

1223

1224

1225   Figure 5

1228   Figure 6

1229

1230

1231　　　　　Figure 7

Figure 8