Inferring tumour microenvironment ecosystems from scRNA-seq atlases

Chengxin Yu[1, 2], Michael J Geuenich[1, 2], Sabrina Ge[3, 8], Gun-Ho Jang[6], Tan Tiak Ju[1, 2], Amy Zhang[6], Grainne M O'Kane[6, 8, 9], Faiyaz Notta[3, 6, 8], Kieran R Campbell[1, 2, 4, 5, 6, 7, *]

[1] Lunenfeld-Tanenbaum Research Institute, Sinai Health System, Toronto, ON M5G 1X5, Canada

[2] Department of Molecular Genetics, University of Toronto, Toronto, ON M5S 1A8, Canada

[3] Department of Medical Biophysics, University of Toronto, Toronto, ON M5G 1L7, Canada

[4] Department of Statistical Sciences, University of Toronto, Toronto, ON M5S 3G3, Canada

[5] Department of Computer Science, University of Toronto, Toronto, ON M5T 3A1, Canada

[6] Ontario Institute of Cancer Research, Toronto, ON M5G 1M1, Canada

[7] Vector Institute, Toronto, ON M5G 1M1, Canada

[8] Princess Margaret Cancer Centre, Toronto, ON M5G 2M9, Canada

[9] Department of Medical Oncology, St. Vincents University Hospital and University College,Dublin

[*] Correspondence: kierancampbell@lunenfeld.ca

Accurate inference of granular cell states that co-occur within the tumour microenvironment (TME) is central to defining pro- and anti-tumour environments. Here, we describe how ecosystems of cell populations can be robustly inferred from nonspatial single-cell RNA-seq (scRNA-seq) atlases. Leveraging a unique discovery-validation setup across eight scRNA-seq datasets profiling pancreatic ductal adenocarcinoma (PDAC), we show highly consistent co-occurrence of fine-grained cell states across patients and characterize the positive predictive value of such analyses. Building on this, we develop a novel probabilistic model to quantify multi-cellular ecosystems directly from such atlas-scale scRNA-seq datasets. By mapping these ecosystems to spatial transcriptomics data, we demonstrate that such ecosystems represent bona fide spatial variation within individual tumours despite being learned from nonspatial data. Importantly, through mapping these ecosystems across two large clinical cohorts, we show they are more predictive of therapy response than any individual cell state and are associated with specific tumour somatic mutations. Together, this work lays the foundation for inferring reproducible multicellular ecosystems directly from large nonspatial scRNA-seq atlases.

Introduction

Recent advances in expression profiling technologies have significantly accelerated our ability
to quantify cellular phenotypes at single-cell and spatial resolutions. Single cell RNA-seq[1,2]
(scRNA-seq) enables transcriptome-wide gene expression measurements at single cell
resolution and has led to multiple discoveries of cell types and transcriptional states associated
with patient outcomes and therapy response[3–8]. In tandem, spatial expression profiling
technologies[9–12] that quantify gene or protein expression at (sub-) cellular resolution while
retaining the spatial location of each measurement have led to multiple studies that associate
tumour spatial architecture with patient outcomes[4,5,13,14]. Of note, significant attention has been
given to using these technologies for the *de novo* identification of cell types and states that co-
occur within tissues[15] and form multi-cellular communities (or ecosystems)[16–18]. These
ecosystems form multi-cellular networks whose presence in tissues has been linked to clinical
outcomes such as treatment response across a range of pathologies[13,19]. Importantly, much of
our understanding of tissue dynamics leading to target discovery comes from reasoning about
cellular co-occurrence within the TME.
48
However, reliable quantification of cell state co-occurrence within individual tumours leading to
cellular ecosystems remains challenging. In particular, the ability to robustly identify recurrent
coexisting cell states or gene programs in specific clinical scenarios relies on having sufficient
independent samples, particularly for rare or transient cell states. Spatial protein technologies
are sufficiently cost-efficient to profile the required number of tissue samples, but can only
quantify 40-60 markers simultaneously, limiting the ability to infer granular cell states and
signaling pathways. On the other hand, spatial transcriptomic technologies can measure 100s of
mRNA species or even the whole transcriptome simultaneously, but are currently too expensive
to profile at scale and suffers resolution or RNA contamination problems[20]. In contrast, scRNA-
seq can quantify single-cell whole transcriptome expression measurements with quantified cell
state abundance largely reflecting tissue composition[21–23]. While scRNA-seq nominally suffers
from the same expense per sample as spatial transcriptomics technologies, the push by the
community over the past decade through initiatives such as the Human Cell Atlas[24] and Human
Tumour Analysis Network[25] has resulted in unprecedented data sizes, comprising more than 12
tumour atlases[4,26–31] by mid-2024 alone. This leads to the tantalizing possibility of quantifying
cell state co-occurrence and cellular ecosystems directly from large multi-cohort scRNA-seq
cancer atlases. However, whether this can lead to reproducible, clinically relevant insights
remains largely unknown.
67
To help answer this question, we focus on pancreatic ductal adenocarcinoma (PDAC) due to
both the availability of multiple independent primary tumour scRNA-seq datasets studies[4,32–38],
combined with its complex microenvironment that is characteristically difficult to target. Existing

studies of the PDAC TME have highlighted diverse cellular heterogeneity across all lineages[39–

43]. Additionally, single-cell profiling of PDAC tumours has highlighted previous underappreciated

heterogeneity within individual tumours cells themselves, with the basal-like/classical[44] and later

hybrid subtypes[40] being shown to co-exist within single tumours[45,46]. Despite this, few efforts

have sought to explicitly quantify the co-occurrence of granular cell states within single tumours.

Oh et al. curated scRNA-seq PDAC from four studies to infer cell type co-occurrence using

clustering and interactome analysis[41], but due to having relatively few samples did not include

validation or highly granular cell states. In contrast, Luca et al. built a machine learning

framework EcoTyper to discover multicellular communities across solid tumours from bulk RNA-

seq data, which is well-powered at the patient level but may lack the ability to uncover granular

cell states *de novo*. Therefore, there is an outstanding question for whether granular PDAC cell

states can be robustly inferred and validated across patient cohorts, and whether cell state co-

occurrence provides insights into the immune dynamics in tumours.

In this work, we present a comprehensive platform to dissect the extent to which cell state co-

occurrence and multi-cellular ecosystems can be reproducibly inferred in the PDAC tumour

microenvironment from atlas-scale scRNA-seq (**Fig. 1A**). We collate eight independent datasets

encompassing >150 samples and >500,000 cells to create a unique dataset-level discovery-

validation setup. In doing so, the existence of known and *de novo* granular cellular subtypes in

the TME can be independently verified, along with their ability to co-occur within and across cell

types and form cellular ecosystems. We map 50 cell type-specific gene programs across 8 cell

types and confirm the existence of known subtypes such as exhausted cytotoxic T cell while

discovering novel cellular variation such as multiple reproducible inflammatory cancer

associated fibroblast (CAF) subtypes. Next, we evaluate the extent to which co-occurrence of

these granular cell state programs is reproducible within and between cell types, finding

consistent co-occurrence across major cell types. We subsequently formalize a probabilistic

model for quantifying multicellular ecosystems from atlas-scale scRNA-seq and find them to be

reproducible across our datasets. Interestingly, by mapping these ecosystems to spatial

transcriptomics data, we demonstrate that such ecosystems represent bona fide spatial

variation within individual tumours despite being learned from nonspatial data. Finally, we map

the validated gene programs and ecosystems to two bulk datasets and find that certain

ecosystems are associated with patient outcome along with specific KRAS mutations.

104   Results

105   Identifying reproducible cellular subtype heterogeneity in the PDAC TME

To systematically quantify cell states from the PDAC tumour microenvironment, we compiled 8

previously generated scRNA-seq datasets (**Supplementary Fig. 1A** and **Methods**,

Supplementary Table 1)[4,32–37,47] comprising 578,000 total cells. To exclude possible

technology-induced bias, all selected scRNA-seq data were generated using the 10x Genomics
platform. We selected primary PDAC tumour samples from each patient cohort and split the
datasets into a discovery group (4 datasets, n=74 tumour samples) and a validation group (4
datasets, n=89 tumour samples). Raw count data preprocessing was performed in line with
previous studies (**Supplementary Fig. 1B**, **Methods**). An automated workflow was developed
to systematically annotate single cells using a public multi-cohort PDAC atlas as reference
(**Supplementary Fig. 2A** and **Methods**)[47]. Eight major cell types were identified in our curated
atlas (**Fig. 1B**). To validate the automatically assigned cell type labels, we examined known
marker expressions in each dataset (**Supplementary Fig. 2B**).
118
To quantify fine-grained cellular subtypes in the TME, we extracted cell type-specific gene
programs independently for the discovery and the validation datasets. Specifically, we applied
integrative non-negative matrix factorization (iNMF)[48] that learns a set of gene programs defined
by the expression pattern of a set of genes along with their activities in each single cell
(**Supplementary Fig. 3A**, **Methods**). The inherent plasticity of the cells of the TME motivated
our use of NMF rather than sub-clustering[5,32,49,50], which allows each cell to have a continuous
activation of multiple gene programs compared to clustering approaches that force each cell to
belong to a distinct cluster, following recent work in this area[51].
127
Given concerns around whether granular cell subclusters from scRNA-seq reflect noise or
overclustering[52], we set out to identify a subset of cell type-specific high-confidence gene
programs found in both our discovery and validation datasets. We first employed the Hungarian
algorithm[53] to obtain optimal one-to-one mapping between gene programs from the discovery
and the validation datasets (**Supplementary Fig. 3B, C**). Next, we merged mapped gene
programs with similar gene weights to reduce redundancy (**Methods: Program validation,
Program merge**) (**Supplementary Figure 3D**). Lastly, we removed any gene programs likely
corresponding to ambient RNA programs contaminating[54] by manual examination of the top
program markers (**Methods**). After applying this procedure, we obtained 10 high-confidence
gene programs in epithelial cells, 5 high-confidence gene programs in each of fibroblasts and
dendritic cells, 11 in monocytes/macrophages, 8 in CD8 T/NK cells, 3 in CD4 T cells, 4 in B
cells, and 4 in endothelial cells (**Fig. 1D**). In total, we obtained 50 cell type-specific gene
programs that capture both the overall TME landscape and the granular phenotypic
heterogeneity (**Supplementary Fig. 4**). Overall, our approach allowed us to identify high-
confidence recurrent cellular subtypes in PDAC TMEs for downstream analysis.
143
We next contextualized the high-confidence cell type-specific gene programs to distinguish
potentially novel PDAC TME cell states from known ones. We took a multi-faceted approach
including marker gene inspection, GSEA, and literature review (**Methods**). Importantly, our
unique discovery-validation setup that independently discovered reproducible programs in

multiple datasets confers high confidence on their manifestation in patients. While this
recapitulated many existing cell states as previously reported in single-cell profiling of
PDAC[32,39,44,55]—including basal and classical epithelial subtypes as well as inflammatory and
myofibroblastic cancer associated fibroblasts (CAFS)—we discovered certain high-confidence
gene programs previously undescribed.
153
In the epithelial compartment, we discovered a previously undescribed single-cell KRAS
amplification epithelial program marked by upregulation of *KRAS* and *BNIP3L*/*NIX*—previously
shown to be a driver of glycolysis and disease progression in pancreatic cancer[56] (**Fig. 1E**). In
the fibroblast compartment, we identified two distinct inflammatory cancer associated fibroblast
(iCAF) subsets and two myofibroblastic CAF (myCAF) subsets for the first time (**Fig. 1F**). The
classical pathway (CCP) iCAF program showed strong upregulation of *C7* and the classical
pathway genes *C1S* and *C1R*. The CCP-iCAF program was also enriched for a neurotropic
gene set linked to tumour growth, stress adaptation, and metastasis in PDAC[5,57]
(**Supplementary Fig. 5A**). In contrast the *adhesive iCAF* program was highly enriched for an
adhesion gene signature including strong upregulation of the gene *SERPINE1* that encodes for
the protein plasminogen activator inhibitor-1 (*PAI-1*) (**Fig. 1F**). The *ACSL3-PAI-1* signaling axis
supports PDAC progression, links to fibrosis, immunosuppression, and prognosis[58]. We found
the CCP iCAF program and the adhesive iCAF program to be present in a mutually exclusive
pattern across patients within the dataset (**Supplementary Fig. 5B**). Two myCAF programs
were identified based on CAF subtype gene loadings in fibroblast gene programs
(**Supplementary Fig. 5C**). The *TPM2+* myCAF-1 program was highlighted by the high loading
of *MIF*[56] (**Fig. 1F**). The *VMP1+* myCAF-2 program was identified as a potential novel myCAF
subtype. Although VMP1 has been associated with PanIN formation[59] and patient survival in
pancreatic neuroendocrine tumours[60], its role in PDAC CAFs is still unclear.
173
Finally, since intra- and inter-tumour heterogeneity is closely coupled with treatment resistance
and therapy response across cancers[6,30,61,62], we investigated whether heterogeneity in the
granular cell states depicted by our gene programs was conserved between our discovery and
validation datasets. We calculated gene program inter-patient heterogeneity and intra-patient
heterogeneity **(Methods)** and found inter- and intra-patient program loading heterogeneity
varied greatly across cell types and within cell types (**Supplementary Fig. 6A, B**). Gene
programs of CD4 T cells, CD8 T/NK cells, epithelial cells, and macrophages showed wide
ranges of intra-patient heterogeneity across patient cohorts (**Supplementary Fig. 6B**). Across
cell types, we found inter-patient heterogeneity tended to agree between the discovery and the
validation datasets, except for T cell programs and macrophage programs (**Supplementary**
Fig. 6B). In short, we showed in single-cell data for the first time that both intra- and inter-patient
cell state heterogeneity is conserved between discovery datasets and validation datasets (intra-
patient heterogeneity: $\rho = 0.67$, $p = 1 \times 10^{-08}$, inter-patient heterogeneity: $\rho = 0.44$, $p = 0.00053$).

This suggests that the abundance of granular gene programs within the TME can serve as a
robust estimate of heterogeneity and help guide target discovery efforts that specifically attempt
to cover the entire phenotypic space.
190

Co-occurring cell type-specific gene programs are highly reproducible in PDAC

We next evaluated the extent to which gene program co-occurrence within individual tumours
could be reproducibly quantified to understand if fine-grained cellular subtype co-occurrence
can be robustly estimated from scRNA-seq data. We computed co-occurrence of all possible
gene program pairs (N=1770 total, **Methods**) separately in the discovery and the validation
groups (**Fig. 2A**). To estimate this co-occurrence, we computed the Spearman correlation of the
activation of each pair of gene programs across samples. This lets us estimate both the co-
occurrence of gene programs within major cell types (such as if exhausted and proliferative CD8
T cells co-occur) as well as the co-occurrence of gene programs between major cell types (such
as if exhausted CD8 T cells and inflammatory cancer associated fibroblasts co-occur). We
quantified reproducibility of these estimates by correlating the co-occurrence in discovery and
validation settings, finding highly concordant estimates (ρ=0.46, $p<2x10^{-16}$) (**Fig. 2B**). We next
used the aggregated program pair co-occurrence data to derive the positive predictive value
(PPV) treating the validation co-occurrence as the ground truth and discovery as the prediction.
We found that overall, the program co-occurrence PPV 80.6% for intra-cell-type co-occurrence
and 74.7% for inter-cell-type co-occurrence (**Fig. 2B**). In addition, most programs saw a PPV
above 50% when examined individually (**Fig. 2E**), allowing for high confidence interpretation of
co-occurrence relationships.
209

Next, we examined the reproducibility of cell state co-occurrence within and between major cell
types. For co-occurrence of programs from the same cell type, we found highest reproducibility
in fibroblasts (ρ=0.77) followed by monocyte/macrophage (ρ=0.59) (**Fig. 2C**). Gene program co-
occurrence within T cells, DCs, and epithelial cells exhibited lower reproducibility between
discovery and validation (ρ=0.37 to 0.5), while endothelial cells and B cells saw the lowest
reproducibility across datasets (ρ=0.086 and 0.22 respectively). Nevertheless, B cell programs
had positive co-occurrence in discovery (ρ=0.2-0.81) and validation (ρ=0.38-0.8), demonstrating
that the overall effect size was consistent (**Fig. 2C**). We found all cell types exhibit significantly
reproducible inter-cell-type program co-occurrence, most notably once again in Fibroblasts
(ρ=0.47) and Monocyte/Macrophage (ρ=0.49) (**Fig. 2D**). Indeed, on a program-by-program
basis we found most to have significantly reproducible inter-cell type co-occurrence
relationships (**Fig. 2E**).
222

We next focused on the co-occurrence of specific cellular subtypes that are either significantly
reproducible between discovery and validation or *a priori* known to hold clinical significance. The
co-occurrence of regulatory CD4 T cells and exhausted CD8 T cells in the PDAC

microenvironment was well known from mouse studies[63], but is yet to be demonstrated in
humans. We found that the CD4 T_{reg} program consistently co-occurred with the CD8 T cell
exhaustion program in patient tumours (**Fig. 2F**). Also, the CD4 T_{reg} program co-occurred with
the *CD163*+ alternatively activated macrophage program and the endothelial angiogenesis state
program marked by *TIE1*, *INSR*, and *NOTCH4*. This angiogenesis program has been found to
be an ECM remodeling and angiogenic signature in lung and breast cancers[64,65] and likely
interact with TAMs and regulatory T cells in breast cancer to promote an immunosuppressive
phenotype in the microenvironment[65]. This T_{reg}-T_{exh}-Mac_{alt} co-occurrence pattern is spatially
confirmed in multiple cancers[66], suggesting our subtype program co-occurrence may be
associated with spatial ecosystems in the PDAC.
236
Finally, we investigated whether known multi-cellular structures in the TME were consistent with
our co-occurrence results. Specifically, we focused on Tertiary lymphoid structures (TLSs),
localized immune aggregates within the TME that can lead to production of tumour-specific
antibodies. TLSs in PDAC are usually composed of proliferating, activated, germinal center
(GC) B cells, antigen presenting cells, and T cells and their presence is predictive of patient
survival[67]. Therefore, we examined the co-occurrence pattern for our germinal center B cell
program. Indeed, the B cell activated and cycling programs are among the top programs that
co-occur with the GC program (**Fig. 2G**). In addition, we found significant co-occurrence of the
activation and memory program of CD8 T cell programs with the GC B cells (ρ=0.3 in the
validation group) along with mature and plasmacytoid dendritic cells (ρ=0.45). Examination of
the patient sample level loadings of the TLS related cell state programs suggests variable
presence of TLS in PDAC tumours (**Supplementary Fig. 7A, B**). These results suggest that in
our datasets, TLS presence and phenotype could be read out from scRNA-seq. Our results
imply co-occurrence of granular cellular subtypes from scRNA-seq can be meaningfully
quantified from large scale atlases, and complex immune cell clusters can be identified and
characterized without the need for additional spatial transcriptomics or proteomics.
253
A probabilistic model to infer recurrent multi-cellular ecosystems in the TME
Beyond the pairwise co-occurrence of gene programs, multi-cellular ecosystems play a
fundamental role in the TME, as these systems often act in concert, and their function is highly
synergistic[68]. For this reason, we asked whether it was possible to infer ecosystems
reproducibly from scRNA-seq. However, there are no existing computational tools to do this
from scRNA-seq using a set of biologically meaningful gene programs as input. Therefore, we
developed a novel probabilistic model (**Methods**) that can take gene programs representing
cellular subtypes and infer multiple multi-cellular ecosystems, reporting each both in terms of
the subtypes that co-occur as well as the ecosystem abundance within each patient sample.
Importantly, our model carefully decouples the fact that programs can co-vary within the same
cell (intrinsic variation) but also between cells (extrinsic variation) (**Methods**). Using maximum a

posteriori inference (**Methods**), we fitted the ecosystem model separately on the discovery and
validation groups using all the 50 cell type specific gene programs. We quantified four
ecosystems, a number which is guided by sub-TMEs found with histology[46].
268
We first contrasted the ecosystems found in the discovery and validation datasets to both
validate the overall methodology and generalizability of the findings. We characterized the
ecosystems by examination of the abundance of gene programs within each (**Fig. 3A**). We
found significant positive correlation in 3/4 ecosystems between discovery and validation (**Fig.**
3B). We identified a Classical ecosystem enriched for corresponding epithelial programs,
endothelial cells, iCAFs, and tumour associated macrophages. A Basal-like ecosystem was
enriched for epithelial EMT programs, myCAF programs, and alternatively activated
macrophage programs. We termed the third ecosystem immune activated as it was enriched
with functional immune cell programs related to anti-tumour response, such as the memory CD4
T cell program and the effector CD8 T cell program. Notably, the TLS related programs were
also enriched in the immune activated ecosystem (**Fig. 3A**). For confirmation, we found the
classical and basal-like ecosystems in individual tumour samples were significantly anti-
correlated in both discovery and validation cohorts (ρ = -0.43 and ρ = -0.51 respectively)
(**Supplementary Fig. 8A**).
283
We next investigated whether our tissue ecosystems represent spatial variation within the
tumour rather than simply per-patient co-occurrence. Leveraging 24 PDAC 10x Visium spatial
transcriptomics slides across 16 patients[4], we scored individual spots for each ecosystem (**Fig.**
3C, F, Methods)**.** We computed Moran's I—a measure of spatial autocorrelation—for each
ecosystem, then shuffled gene program weights to create 100 randomized ecosystems. We
found the observed spatial autocorrelation from our scRNA-seq inferred ecosystems was higher
on average than random across all three ecosystems, and significantly so for two (Basal-like: p
= 0.024, immune activated: p < 0.0001, **Fig. 3D**). We also compared our ecosystems to an
unsupervised learning approach for TME community discovery by generating consensus multi-
program factors from spot-wise program scores using NMF. We compared the spot-wise spatial
loadings of the factors with our ecosystem by computing a correlation coefficient. We found the
NMF factors were not able to reproduce the Basal and immune activated ecosystem, while only
the NMF factor 3 was showing moderate agreement with the classical ecosystem (**Fig. 3E**).
Indeed, examining the spatial presence of the ecosystems inferred from scRNA-seq across four
Visium slides shows pronounced spatial heterogeneity (**Fig. 3F**).
299

Associating ecotypes with tumour genetics and clinical outcomes
To quantify the association between both our gene programs and ecosystems with tumour
genetics and clinical outcomes, we curated clinical and genome sequencing data for two
cohorts of PDAC tumour samples with paired bulk RNA-seq data, which we refer to as the Chan

cohort[67] (N=183 patients) and TCGA pancreatic cancer cohort[69] (N=151 patients). We scored
each sample for each of the gene programs we identified (**Methods**)[70] and subsequently
associated each sample with each ecotype by quantifying weighted combinations of gene
programs (**Methods**). We first examined associations between the novel gene programs we
identified and overall patient survival. Interestingly, the KRAS amplification epithelial program
was significantly associated with poor patient survival in both the TCGA cohort and the Chan
cohort (**Fig. 4A**). Importantly, the presence of this program was more predictive of patient
survival than well the established classical and basal gene expression subtypes (**Fig. 4B**).
Together, this implies such workflows that exploit high-confidence gene programs identified
from single-cell RNA-seq data and validated via discovery-validation splits may be more
clinically relevant than existing subtypes identified from bulk profiling approaches.
315
Next, we examined links between the multicellular ecosystems and patient response to therapy.
Interestingly, the Basal-like ecosystem was positively associated with progressive disease after
chemotherapy in both the Chan cohort (p<0.01) and the TCGA cohort (p=0.40)
(**Supplementary Fig. 8B, 9**), and more significantly associated with progressive disease than
any individual gene program including all Basal epithelial programs in the Chan cohort (**Fig. 4C,**
D). Similarly, the immune activation ecosystem was positively associated with partial remission
and negatively with progressive disease in the TCGA cohort (p<0.05) more significantly than
any individual gene program (**Fig. 4C, D**). Similarly, this ecosystem was negatively associated
with progressive disease in the Chan cohort, albeit not statistically significantly, (p=0.11)
(**Supplementary Fig. 8B, 9**). These differences in statistical power between cohorts likely
reflect differences in sample preparation: the Chan cohort underwent laser capture
microdissection for the tumour epithelium, making it better placed to capture tumour associated
ecosystems (Basal-like), while the TCGA cohort was not tumour enriched, which given the low
cellularity of PDAC tumours (average = 35%) makes it more adept at capturing immune
populations.
331
Finally, we contextualized the link between multicellular ecosystems and tumour genetics.
PDAC tumours carrying *KRAS*-G12D mutations are associated with poorer prognosis[71], and
strategies targeting specific *KRAS* neoantigens are being developed[72]. However, the
mechanisms through which different mutations affect the microenvironment in human PDAC
remains unclear. We obtained *KRAS* variant calls for the HTAN Zhou (WUSTL) patient dataset[4]
in our validation cohort and fitted a linear mixed model between ecosystem presence and *KRAS*
variant status (**Methods**). We found that compared to *KRAS*-G12D, tumours with a *KRAS*-
G12V/G12R mutations had a significantly higher presence of the immune activated ecosystem,
while tumours with *KRAS*-G12R mutations had a reduced presence of the basal-like ecosystem
(**Fig. 4E**). Importantly, we retrieved similar *KRAS* variant calls in the bulk Chan patient
cohort[69,70] and found consistent effect sizes for all associations (**Supplementary Fig. 8C**,

Supplementary Table 3), validating the overall findings. The decreased presence of the
immune activated ecosystem in *KRAS*-G12D PDAC suggests differences in anti-tumour
immune response may underline prognosis and effectiveness of immunotherapy[71,72]. Together,
these results suggest that the specific driver mutations in PDAC have a significant effect on the
phenotypes and cell communities in the TME.
348

Discussion

Transcriptome profiling of tumour samples has historically been used to identify molecular
markers that could facilitate cancer diagnosis, prognosis prediction and identification of
molecular candidates to target therapeutically[73–76]. The advent of single cell technologies has
facilitated the identification of the relative distribution of cell populations specific for certain
cancer types, predicting clinical outcomes and uncovering specific cellular subtypes as
targets[77–80]. However, it is becoming increasingly clear that focusing on the expression pattern
of a handful of genes or the presence/absence of certain cell populations is not sufficiently
informative to gain useful biological insights into the whole TME, even though this information
could be leveraged to design targeted therapies aimed at tackling the combined action of
multiple cell types. In this context, spatial information is extremely useful to identify co-
localization of different cell types, whose crosstalk requires a certain three-dimensional
organization to synergistically alter tumour growth and survival. However, a substantial number
of samples are needed to robustly identify co-occurrence of interacting cells across patients,
especially considering that the cellular subtypes of interest are potentially rare. Therefore, the
current cost and/or limitations of spatial transcriptomics and proteomics limit the identification of
reproducible co-existing cell states with prognostic value[79]. By contrast, the number of scRNA-
seq datasets from TMEs that contain transcriptome-wide expression quantification at single-cell
resolution is increasing exponentially, allowing, in theory, to identify co-occurrence of granular
cell states and associate them with clinical outcomes.
369

Here we have undertaken a unique discovery-validation analysis on the largest PDAC scRNA-
seq atlas to-date to quantify the extent to which granular cellular subtypes and ecosystems can
be reliably inferred. Our 50 reproducible cell type specific gene programs captured previously
known heterogeneity such as classical and basal tumour transcriptional but also novel subtypes
such as two iCAF subtypes as well as a novel *KRAS amplification* epithelial cell program
significantly associated with survival. We further revealed for the first time that co-occurrence of
gene programs can be reproducibly quantified from large scRNA-seq datasets. We further
identified a TLS pattern centered around the GC B cell program, recapitulating a spatial
structure important for anti-tumour immune response in multiple cancers including PDAC[81]. We
introduced a new model for inferring ecosystems in TME and found reproducible ecosystems,

which were validated in spatial transcriptomic data and associated with both driver gene

mutants as well as response to therapy.

Overall, we show that large scRNA-seq atlas can be employed to not only identify TME cellular

subtypes and ecosystems with high confidence, but that these have profound biological and

clinical significance, as the frequency of co-occurrence of certain programs and the presence of

certain ecosystems are associated with patient survival. These analyses suggest that in the

future, scRNA-seq datasets will be used to study the broad interplay among pro-tumorigenic cell

populations in the TME. This approach may eventually expand the clinical usability of single cell

data, offering a new lens to develop innovative therapeutic strategies targeting PDAC.

Despite these findings, our study had multiple limitations. While our discovery group was well

balanced between four datasets, the validation cohort was heavily weighted towards one cohort,

which may have limited the cell state program validation process though we note that, since

there was no overlap in samples, it still functions as a well-powered validation cohort. In

addition, the lack of well-standardized and curated clinical metadata across the single-cell

cohorts limited the ability to directly associate cell states and ecosystems with clinical

covariates, instead necessitating mapping them to bulk datasets with standardized metadata.

Moving forward, we identified several venues to extend the project. Future spatial profiling of the

PDAC microenvironment at cellular resolution in large cohorts will help to create a more

comprehensive understanding of the cell state co-occurrence related to therapy resistance and

disease progression. Integrating T cell receptor (TCR) or B cell receptor (BCR) sequencing

analysis will help to delineate relationships between lymphocyte clonality with functional cell

states, potentially revealing specific anti-tumour T cell clones and effects of therapy on the

tumour immune cell population. In the long term, our analysis workflow may be adapted to

multiple cancer types, leveraging the ever-increasing tumour single-cell data space to create a

pan-cancer microenvironment cell state atlas, and identifying clinically relevant cellular subtypes

across cancers. The increasing characterization of the TME at single-cell level will enable the

identification of concomitant presence of multicellular ecosystems in various types of cancer.

Our approach unlocks a new way to analyse scRNA-seq data that will not only facilitate

prognosis and assessment of therapeutic response, but will provide valuable information to

design specific therapies aimed at rewiring the milieu compatible with the co-existence of pro-

tumorigenic cellular subtypes and ecosystems.

Methods

Data curation

A total of 8 publicly available PDAC single-cell RNA-seq (scRNA-seq) datasets, which include 214 samples, were curated[4,32–37,47]. Most datasets were available for downloading through the Gene Expression Omnibus database. One dataset was available on author consent, in which case we contacted the authors for permission. Each dataset contains varying types of samples. We excluded samples which are of the following nature: metastasis, normal pancreas, peripheral blood, and adjacent normal tissue. In total, 163 samples labeled as primary PDAC tumour were retained for downstream analysis.

After sample filtering, we performed in house pre-processing and quality control given all datasets were available in the CellRanger output format. Doublet scores for single cells were calculated and their doublet status was called using doubletFinder_v3 from DoubletFinder v.2.0.3[82]. Cells labeled as doublets were removed from each sample. After doublet removal, cells with mitochondrial RNA content percentage greater than 20%, detected number of genes lower than 500, and total transcript counts lower than 1000 were removed as low quality cells. Finally, the UMI counts were normalized by per-cell size factor and log2-transformed with logNormCounts from scuttle v.1.8.4[83].

Cell annotation unification

We performed re-annotation to unify the cell annotation in each patient cohort in our curated dataset. First, we obtained a single-cell PDAC atlas built from divergent and heterogeneous datasets[81]. The 'celltype2' annotation from the atlas was selected as the reference labels after comparison of resolution and quality of all annotation levels in the atlas. Then, we performed unbiased cell type recognition at the single cell level using SingleR from SingleR v.2.0.0[84]. Briefly, SingleR first trains a classifier on the PDAC atlas reference dataset with reference labels by leveraging differential expression between labels, then the labels were assigned to each single cell in the query dataset using the classifier combined with iterative fine-tuning. We ran SingleR for each patient cohort in the curated dataset independently with the following settings: excluding ribosomal and mitochondrial genes in the reference, selecting top $500 * (2/3)^{log_2(N)}$ (N = # unique genes) DE genes with ranking by log-fold change, aggregating the reference into pseudo-bulk samples with default settings, and enabling fine-tuning with default settings. After annotation by SingleR, quality checks were performed by examining the SingleR predicted scores for each label in single cells in each cohort, the distribution of delta between assigned label score and median score across labels, and the distribution of scores for each label (**Supplementary Fig. 1C**). Final validation was done by examining known cell type marker expressions in each assigned cell type. Finally, the SingleR pruned.label column was selected and cell type labels were assigned as the unified annotation for single cells. To facilitate

453   downstream analysis, endocrine cells and neural cells were removed from the curated dataset,

454   and acinar and ductal cells were both labeled as epithelial cells. The eight consolidated cell type

455   labels were then shortened for readability.

Discovery-Validation data split

We first subset the full curated dataset by cell annotation, splitting it into eight cell type subsets:

Epithelial cell, Endothelial cell, Fibroblast, Monocyte/Macrophage, mDC, CD4 T cell, CD8 T/NK

cell, and B cell. We then split each cell type subset into two 4-cohort groups, one as the

discovery group and the other as the validation group. The discovery group contained 74

samples from the Peng, Steele, Werba, and Chan patient cohorts, while the validation group

contains 89 samples from the Chen, Lee, Lin, and Zhou patient cohorts. In total, the discovery-

validation split resulted in eight discovery-validation pairs of cell type subsets, with each pair

containing single cells assigned to one cell type.

Reproducible analysis pipeline

468   We built a comprehensive analysis pipeline to generate reproducible results from our curated

469   dataset and reduce human-introduced variations in analyses across the cell types and between

470   the discovery group and the validation group. We used Snakemake v.7.25.0[85] for parameter and

471   input-output control for all analysis scripts and visualization steps.

Defining a set of robust lineage-specific cell state gene programs

We carried out integrative NMF (iNMF) using LIGER[82] for the discovery and validation group of

patient cohorts across cell types independently to generate cell lineage-specific gene programs

that capture the phenotypic heterogeneity within each cell type in the PDAC microenvironment.

LIGER runs iNMF to decompose the expression matrices from multiple batches (patient

cohorts) and remove batch effect at the same time by maximizing a basis matrix shared

between cohorts while minimizing the multiple cohort-specific basis matrices that represent

patient cohort level biases. We defined the batches to be patient cohorts to focus on capturing

heterogeneity present across tumours through cell lineage-specific programs. We followed the

suggested workflow of LIGER to perform iNMF. Importantly, the following steps were performed

after the discovery-validation split as well as an 8-way cell type split within the discovery and the

validation group, leading to 16 independent gene program extraction steps, 8 independent

program validation steps, and 8 independent program merging steps.

Program extraction

First, we selected variable genes from non-ribosomal/mitochondrial protein-coding genes using

selectGenes from rliger v.1.0.0[83] with default settings. We then created a liger object using the

raw counts matrix, removed genes with zero expression, and normalized and scaled gene

expression following suggested workflow from rliger documentation before performing a subset

to the variable genes. Finally, after filtering to variable genes, we fitted iNMF on the expression

matrix using optimizeALS. We set rank (k) to 20, 10, 5, 10, 10, 20, 10, and 10 for program

extraction from epithelial cell, fibroblast, endothelial cell, CD4 T cell, CD8 T/NK cell,

monocyte/macrophage, mDC, and B cell respectively. We set lambda to 5 for all cell types as

we did not find it to be a major factor affecting extraction results. The gene loadings in each

program were obtained as the W matrix from the fitted liger object, and the program loadings in

single-cells were obtained as the H.norm matrix from the fitted liger object.

Program validation

Our program validation process was performed between discovery and validation for each cell

type. To start, we transformed the discovery and validation gene loading (W) matrix with log1p

in R. We then computed the spearman correlation and Minkowski distance (with p = 1) between

all possible pairs of discovery and validation programs within the cell type. The Hungarian

algorithm was used to generate a one-to-one relationship between discovery and validation

programs by permuting the rows of the distance and correlation matrices to minimize and

maximize the trace of the matrices respectively. After assigning the one-to-one partner of each

discovery program, we rejected any discovery programs whose gene loading correlation with

their validation program partner was below 0.5, or if the gene loading distance with their

validation program partner was above 25,000. We further confirm validated programs by

permutation test. Gene loadings in each validated program were permuted to generate a 1000-

sample test set and the correlation and distance to the validation program partner were

calculated for fitting a null distribution. A discovery program is finalized as a validated program if

the real distance and correlation are laid beyond 1% of the null distribution.

Program merge

We collapsed validated programs which were too similar to each other given NMF does not

guarantee orthogonality between the extracted factors. We looked at all possible pairs of

validated programs within each cell type and marked any pair whose gene loading correlation

was above 0.4 for merging. The merging step leveraged simple linear models. For a pair of

validated programs, e.g. sig-1 and sig-2, marked for merging, we first computed their explained

expression y by multiplying program loading in cell and gene loading in program respectively.

Then, we obtained the merged program's gene loading x by taking the mean of gene loadings in

sig-1 and sig-2. Finally, the single-cell loading of the merged program was obtained by fitting

lm(y ~ 0 + x) and pulling the coefficients. After merging, the newly merged program was

included for downstream analysis while sig-1 and sig-2 were excluded.

Program rescoring in the validation group

Using the robust set of validated and merged cell type specific programs, we rescored the

validation group single cells to get single-cell program loadings that can be compared with

program loadings in the discovery group in downstream analysis. The rescoring step was
performed for each cell type and each patient cohort independently, leading to 32 rescoring
steps.
534
To start, gene expression in a validation group patient cohort was normalized using
runSeuratNormalizeData from singleCellTK v.2.8.0[86] with default settings. Then, for rescoring
for programs of, for example, cell type A, we subset an expression matrix to cells annotated as
cell type A and variable genes used in the program extraction step. Finally, we performed non-
negative least squares to fit a simple linear model to the subset validation expression with cell
type A program gene loadings, and pulled out the estimated optimal coefficient matrix as the
rescored program loading in single-cells in validation. We used fcnnls from NMF v.0.26[87] with
default settings for this step.
543
Program contextualization with known cell states
We combined multiple approaches to get a comprehensive understanding of the biological
meaning of our validated cell type-specific programs. We curated multiple PDAC focused
malignant, stromal, and immune cell state marker lists from literature (**Supplementary Table 4**).
To associate the programs with the known cell states, we selected the top 50 loaded genes of
each program and performed a hypergeometric test with the cell state marker lists from its
respective cell type, using GeneOverlap v.1.34.0[88] in R. The human reference hg19 genome[89]
was used as the background for significance testing. We made a preliminary biological
interpretation of the programs based on significance (P-value < 1e-3) and high Jaccard index
with cell state marker lists. Then, the programs were further subjected to GO and KEGG
overrepresentation analysis using the clusterProfiler v.4.6.2 package[90]. Specifically, the
biological process subcategory of GO and KEGG PATHWAY gene sets were chosen for the
analysis. The top ten enriched gene sets in each category were used to help correct or confirm
our initial program interpretation. After the first correction, we also applied GSEA to the
programs with gene sets mined from the Curated Cancer Cell Atlas (3CA)[91] to resolve any
remaining uncertainty. We ran fgsea from fgsea v.1.24.0[92] with default settings, while setting
nPermSimple to 10,000, as suggested by the documentation. We used the top 10 positively
enriched 3CA gene programs to guide a second program interpretation correction step.
Programs that were not associated with known cell states previously were further characterized
after this analysis. Finally, we completed a manual check across programs by looking at the top
10 loaded genes, focusing on the programs that were yet to be contextualized and searching in
the literature for novel evidence of their biological meaning in PDAC. Any novel programs
remained were named by their top loaded genes.
567

Ambient RNA program removal

Due to technical limitation of droplet-based single-cell RNA-seq experiments, some reads associated with a cell barcode may originate from ambient RNA[54], contributing to background noises that are picked up by the granular cell state gene programs. Therefore, we denoted gene programs in a cell lineage that are marked by lineage markers from other lineages to be ambient RNA programs and removed such ambient programs from downstream analysis.

Cell state co-occurrence

To study co-occurrence between programs in the PDAC microenvironment, we constructed sample level program loading profiles by averaging program loadings across single-cells. We then calculated the correlation across samples between programs both within cell type and across cell types. By comparing program loading correlations between the discovery group and the validation group, we define program pairs whose loading correlation was above 0.4 and differed by less than 0.25 in discovery and validation as high confidence pairs that tend to co-occur with each other.

Scoring RNA-seq samples for cell state programs and ecosystems

To score bulk PDAC tumour samples for our cell state gene programs, we performed gene set variation analysis (GSVA) on each bulk sample using gsva from GSVA[93] ver. 1.46.0, with the top-50 loaded genes in each gene programs, and used the program enrichment scores or downstream analysis. To score bulk PDAC tumour samples for our ecosystems, we performed multiplication between the sample-by-program enrichment score matrix and the program-by-ecosystem abundance matrix, to obtain a sample-by-ecosystem enrichment score matrix.

Association between ecosystems and KRAS variants

KRAS variant information was obtained from published data and from collaborators. To account for the patient-repeated measurements in the Chan cohort, we fitted a linear mixed model with the *lmerTest* package in R with the KRAS mutation status as the fixed effect and the patient as the random effect. For the TCGA cohort to estimate effect of KRAS variant on ecosystem enrichment scores we used a simple linear model while controlling for the expression subtype (Basal-like or Classical) of the tumour.

Association between gene programs, ecosystems, and clinical outcomes

To associate the cell type-specific gene programs and multicellular ecosystems with clinical outcomes, we mapped scored RNA-seq samples in the Chan cohort and the TCGA cohort to metadata and clinical data. We fitted univariate linear regression models to each gene program-clinical outcome pair and each ecosystem-clinical outcome pair, and retrieved the t-statistics as estimate of association. We further fitted univariate Cox Proportional-Hazard ratio models to

606 obtain hazard ratios for each program and ecosystem. For survival analysis, we used
607 consolidated survival data from each cohort to plot KM curves to stratify patients.
608

609 **Cell state co-occurrence ecosystems discovery model**
610 To further dissect the relationships between programs, we created a statistical model to
611 deconvolve program co-occurrence and discover multi-cellular population ecosystems.
612

When considering why programs are (anti-) correlated in single-cells, there can be two reasons.
Firstly, within single cells, programs may be correlated due to gene regulation, epigenetics, etc.
We defined correlations resulting from these mechanisms cell intrinsic program correlations.
Secondly, within patients, cells with a high program of one type may be present with (or
exclude) cells with a different program. This could be due to paracrine signaling. We defined
these correlations cell extrinsic program correlations. If, for example, there is positive correlation
between loading of two programs only at the patient level, but not at the single cell level, then
the correlation is dominated by cell extrinsic program correlation, and vice versa for cell intrinsic
program correlation. We note that in both scenarios, the overall correlation between the
programs is the same, even if the mechanisms and interpretation are different. So, it is
important to distinguish the two types of correlation in our model.
624

Model data definition
We start by defining L as the number of ecosystems we want to get, C as the number of cell
types, N as the total number of cells, K as the total number of cell state programs, and P as the
total number of patient samples. Therefore, we have:

$$\text{The set of all cell types: } \boldsymbol{C} = \{C_1, C_2, C_3, \dots, C_c\}$$

$$\sum_{i=1}^{C} K_{C_i} = K$$

$$\sum_{i=1}^{C} N_{C_i} = N$$

$$\text{The set of all patients: } \boldsymbol{P} = \{1, 2, 3, \dots, P\}$$

$$\sum_{i=1}^{C} P_{C_i} = P$$

$$0 < P_{C_i} < P \quad \forall\, C_i \in \boldsymbol{C}$$

Then, for each cell type C_i, we have K_{C_i} programs across N_{C_i} single-cells, and each single-cell
belongs to one of P_{C_i} patients.
637

638 Next, we define some cell-type-specific variables. For each cell type C_i, we have a matrix $\boldsymbol{Y}_{C_i}$
639 storing the cell state program loadings in single-cells:

$$Y_{C_i} = \begin{pmatrix} y_{11} & y_{12} & \cdots & y_{1K_{C_i}} \\ y_{21} & y_{22} & \cdots & y_{2K_{C_i}} \\ \vdots & \vdots & \ddots & \vdots \\ y_{N_{C_i}1} & y_{N_{C_i}2} & \cdots & y_{N_{C_i}K_{C_i}} \end{pmatrix}$$

In Y_{C_i} each row represent one K_{C_i}-dimensional vector for a single-cell. This vector stores the loadings of each cell state program for cell type C_i in the cell. Of note, each single-cell is assigned to one cell type only, and thus only shows up in one of the Y_{C_i} matrices.

To further indicate the patient sample each single-cell belongs to, we introduce a sample indicator vector for each cell type C_i:

$$x_{C_i} = \left(x_1, x_2, x_3, \dots, x_{N_{C_i}} \right) \text{ where } x_i \in \{1, 2, 3, \dots, P_{C_i}\} \ \forall i \in \{1, 2, 3, \dots, N_{C_i}\}$$

Model parameter definition

We first introduce the ecosystem parameters by defining:

1. The ecosystem loading matrix $H = (h_{ij})_{P \times L}$
2. The ecosystem basis matrix $W = (w_{ij})_{L \times K}$
3. The patient level program loading estimate matrix considering extrinsic covariance $\Phi = (\phi_{ij})_{P \times K} = H \times W$

Where each row w_l of W is a K-dimensional vector of program weights in ecosystem l, and each row h_p of H is an L-dimensional vector of ecosystem weights in sample p.

With the ecosystem related parameters defined, we introduce:

1. The patient level program loading estimate matrix considering intrinsic + extrinsic covariance $M = (\mu_{ij})_{P \times K}$
2. The cell-type-specific patient level program loading sub-matrices $M_{C_i} = M_{1:P, j_{C_i} : j_{C_i} + K_{C_i}}$

Where j_{C_i} is the index for the first program of cell type C_i in M, specifically:

$$j_{C_1} = 0$$

$$j_{C_i} = j_{C_{i-1}} + K_{C_{i-1}} \ \forall i \in \{2, 3, 4, \dots, C\}$$

We lastly explicitly define, for each cell type C_i, the cell intrinsic program covariance matrix as:

$$\Sigma_{intrinsic_{C_i}} = \begin{pmatrix} Var(program_1) & \cdots & Cov(program_1, program_{K_{C_i}}) \\ \vdots & \ddots & \vdots \\ Cov(program_{K_{C_i}}, program_1) & \cdots & Var(program_{K_{C_i}}) \end{pmatrix}$$

Model definition

First, we let

$$y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \quad where$$
$$\boldsymbol{y} \in \mathbb{R}^k, \boldsymbol{\mu} \in \mathbb{R}^k, \boldsymbol{\Sigma} = (\sigma_{ij})_{k \times k}$$

We then consider cell extrinsic and intrinsic program correlations separately.

Cell extrinsic program correlations are correlations in the means of the program loadings for each patient. We thus introduce P mean vectors $\boldsymbol{\mu}_p$ (one $\boldsymbol{\mu}_p$ per patient) that are each K-dimensional and encode the mean program loading of patient p. Of note, Then, we can write:

$$\boldsymbol{\mu}_p \sim \mathcal{N}(\boldsymbol{\phi}_p, \boldsymbol{\Sigma}_{extrinsic})$$

Here, $\boldsymbol{\phi}_p$ is the prior on the patient mean program loading, and $\boldsymbol{\Sigma}_{extrinsic}$ is the program by program $K \times K$ covariance matrix of cell extrinsic program correlations.

Cell intrinsic program correlations are the correlation of the programs conditioned on the mean program loadings of each patient. We thus introduce N cell level program loading vectors $\boldsymbol{y}_{n_{C_i}}$ that are each K_{C_i}-dimensional and encode the program loadings for cell n_{C_i} where $C_i \in \boldsymbol{C}$. Therefore, we can write:

$$\boldsymbol{y}_{n_{C_i}} \sim \mathcal{N}(\boldsymbol{\mu}_{x_{n_{C_i}}}, \boldsymbol{\Sigma}_{intrinsic_{C_i}})$$

Here, $x_{n_{C_i}}$ selects the patient sample corresponding to cell n, so $\boldsymbol{\mu}_{x_{n_{C_i}}}$ corresponds to the mean program loadings of the patient corresponding to cell n_{C_i}. $\boldsymbol{\Sigma}_{intrinsic_{C_i}}$ is the $K \times K$ covariance matrix that denotes the program correlation after conditioning on the patient means. Using Gaussians allows us to interpret the total covariance as the sum of the intrinsic and extrinsic covariance, i.e. $\boldsymbol{\Sigma}_{total_{C_i}} = \boldsymbol{\Sigma}_{extrinsic} + \boldsymbol{\Sigma}_{intrinsic_{C_i}}$.

Together, we can write the hierarchical model:

$$\boldsymbol{\phi}_p \sim \mathcal{N}(0, 10) \ \forall p \in \boldsymbol{P}$$
$$\boldsymbol{\Sigma}_{intrinsic_{C_i}} \sim InverseWishart(\cdot) \ \forall C_i \in \boldsymbol{C}$$
$$\boldsymbol{\Sigma}_{extrinsic} \sim InverseWishart(\cdot)$$
$$\boldsymbol{\mu}_p \sim \mathcal{N}(\boldsymbol{\phi}_p, \boldsymbol{\Sigma}_{extrinsic})$$
$$\boldsymbol{y}_{n_{C_i}} \sim \mathcal{N}(\boldsymbol{\mu}_{x_{n_{C_i}}}, \boldsymbol{\Sigma}_{intrinsic_{C_i}})$$

Here,

$$\boldsymbol{\Phi} = \begin{bmatrix} \boldsymbol{\phi}_1 \\ \boldsymbol{\phi}_2 \\ \vdots \\ \boldsymbol{\phi}_P \end{bmatrix} \quad \boldsymbol{M} = \begin{bmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \\ \vdots \\ \boldsymbol{\mu}_P \end{bmatrix} \quad \boldsymbol{Y}_{C_i} = \begin{bmatrix} \boldsymbol{y}_1 \\ \boldsymbol{y}_2 \\ \vdots \\ \boldsymbol{y}_{n_{C_i}} \end{bmatrix}$$

Target function

The target function of the model:

$$\mathbb{P}(\boldsymbol{\Phi}, \boldsymbol{M} \mid \boldsymbol{Y}) \propto \mathbb{P}(\boldsymbol{W}, \boldsymbol{H}, \boldsymbol{M}, \boldsymbol{\Sigma} \mid \boldsymbol{Y}) \propto \mathbb{P}(\boldsymbol{Y} \mid \boldsymbol{M}, \boldsymbol{\Sigma})\mathbb{P}(\boldsymbol{M} \mid \boldsymbol{W}, \boldsymbol{H})\mathbb{P}(\boldsymbol{W})\mathbb{P}(\boldsymbol{H})\mathbb{P}(\boldsymbol{\Sigma})$$

The joint posterior is then:

$$\mathbb{P}(\boldsymbol{\Phi}, \boldsymbol{M} \mid \boldsymbol{Y}) \propto \left(\prod_{i=1}^{C} \mathbb{P}(\boldsymbol{\Sigma}_i)\mathbb{P}(\boldsymbol{Y}_i \mid \boldsymbol{M}, \boldsymbol{\Sigma}_i)\right) \cdot \left(\prod_{i=1}^{P} \mathbb{P}(\boldsymbol{M} \mid \boldsymbol{W}, \boldsymbol{H}_i) \cdot \mathbb{P}(\boldsymbol{H}_i)\right) \cdot \left(\prod_{i=1}^{L} \mathbb{P}(\boldsymbol{W}_i)\right) \propto$$

$$\left(\prod_{i=1}^{C}\prod_{n=1}^{N_i} \mathbb{P}(\boldsymbol{\Sigma}_i)\mathbb{P}(\boldsymbol{Y}_n \mid \boldsymbol{M}_{x_n}, \boldsymbol{\Sigma}_i)\right) \cdot \left(\prod_{i=1}^{P} \mathbb{P}(\boldsymbol{M} \mid \boldsymbol{W}, \boldsymbol{H}_i) \cdot \mathbb{P}(\boldsymbol{H}_i)\right) \cdot \left(\prod_{i=1}^{L} \mathbb{P}(\boldsymbol{W}_i)\right)$$

To encourage the model to find more distinct ecosystems, we add an orthogonality term to the model target function. The upper triangular matrix of the transpose cross product of the ecosystem basis matrix $\boldsymbol{W}$ is used to calculate a mean as a proxy for ecosystem similarity. This term is weighted by a hyperparameter λ and minimized in the target function. Therefore, using maximum a posteriori inference, we maximize the new target function including the log joint posterior and the orthogonality term:

$$\sum_{i=1}^{C}\left(log\mathbb{P}(\boldsymbol{\Sigma}_i) + \sum_{n=1}^{N_i} log\mathbb{P}(\boldsymbol{Y}_n \mid \boldsymbol{M}_{x_n}, \boldsymbol{\Sigma}_i)\right) + \sum_{i=1}^{P}(log\mathbb{P}(\boldsymbol{M} \mid \boldsymbol{W}, \boldsymbol{H}_i) + log\mathbb{P}(\boldsymbol{H}_i)) + \sum_{i=1}^{L} log\mathbb{P}(\boldsymbol{W}_i) +$$

$$\lambda \cdot Mean\big(upper\_tri(\boldsymbol{W}^{\mathsf{T}}\boldsymbol{W})\big)$$

Data preparation for the ecosystem discovery model

To increase the power of our model, we removed patient samples which contained fewer than 20 single-cells. To speed up the inference process, we sampled down each patient sample to 1500 single-cells. For each cell state programs, we scaled the program loadings in single-cells across all cells to get z-scores as input to the model.

Model initialization and optimization

We coded the ecosystem discovery model using the statistical programing language Stan. We used cmdstan_model from the R package cmdStanR ver. 0.5.3[94] to compile the model. Then, we used optimize from cmdStanR to initialize and fit the model to our data, with settings: algorithm="lbfgs", tol_param=1e⁻⁷, and iter=8000. The model parameters were initialized as specified below.

Model initialization was provided for $\boldsymbol{\Sigma}_{intrinsic}$ for each cell type and for $\boldsymbol{W}$ and $\boldsymbol{H}$. We performed NMF on the sample level cell state program loading mean matrix with rank=L to obtain the initial values of the ecosystem matrices. Covariance between program loadings within respective cell types were input as the initial values of $\boldsymbol{\Sigma}_{intrinsic}$.

PDAC Visium data

Visium data was downloaded from Synapse[95] with author permission. Tumour sample slides from primary tumours were selected. Preprocessing was done using the SpatialData[96] library. Low quality spots containing more than 20% hemoglobin genes and fewer than 1000 detected genes were filtered. Non-protein coding genes, mitochondrial genes, ribosomal genes, and hemoglobin genes were excluded from downstream analysis. Raw UMI counts were log-normalized by calling computePooledFactors followed by logNormCounts with default parameters.

Scoring programs and ecosystems in Visium

Top 20 genes ranked by weight in each gene program were used to compute spot-wise scores in Visium PDAC samples. Since each spot on the slide is effectively a mini bulk sample, we leveraged gene set variation analysis (GSVA) to obtain an enrichment score for each program in each spot of each Visium slide. Next, program scores in each spot were scaled to 0-1 range. Consensus weighing factors for each ecosystem was obtained by removing programs whose weights had opposite signs in discovery vs. validation. We then obtained spot-wise ecosystem scores as the sum of the program scores weighted by the consensus weighing factors. Randomized ecosystem baselines were generated by shuffling the consensus weighing factors.

Concatenated NMF

To obtained consensus NMF factors. All Visium spots were concatenated to create one spot x program score matrix. A single NMF model was fitted on the pooled matrix, and the expression was decomposed into three factors common across images.

Spatial autocorrelation

To measure spatial autocorrelation, we computed the Moran's Index for the inferred ecosystems, the randomized ecosystems, and the concatenated NMF factors in each Visium image. We used Squidpy[97] graph algorithms with default parameters for Moran's I computation.

Funding

This work was supported by funding from CIHR project grant PJT175270 (KC), NSERC Discovery grant RGPIN-2020-04083 (KC), and support from the Data Science Institute at University of Toronto (CY). This research was undertaken, in part, thanks to funding from the Canada Research Chairs Program.

Data and code availability

All data and code required to reproduce this study can be found at
https://zenodo.org/records/13332977 and https://github.com/camlab-bioml/PDAC_cellular_niche_analyses.

Contributions

Project conception: CY, KC. Data curation: CY, SG, TTJ, AZ. Data analysis and coding: CY, MG, SG, GHJ, TTJ, AZ. Result interpretation and manuscript writing: CY, KC. Project direction: GO, FN, KC.

Acknowledgements

We thank Life Science Editors for editing services (www.lifescienceeditors.com).

Figures

Figure 1: Learning multicellular ecosystems directly from scRNA-seq atlases. A. Overview
of discovery-validation platform for identifying co-occurrence and ecosystems of cell type
specific gene programs in the PDAC TME. **B**. UMAP representations for the discovery (top) and
validation (bottom) datasets colored by cohorts (left) and annotated cell type (right). DC, Myeloid
Dendritic Cells. Mono/Mac, Monocyte/Macrophage. **C**. Overview of cell type composition for
samples (in rows) in discovery group (top) and in validation group (bottom). **D**, Overview of
numbers of candidate gene programs extracted from the discovery group, numbers of gene
programs validated between discovery and validation, and numbers of consolidated gene
programs after merging the redundant ones in each cell type. **E-F**. Heatmaps showing weights
of top ranked genes in selected gene programs from epithelial cells (**E**) and fibroblasts (**F**).
799

800

Figure 2: Pairwise co-occurrences of gene programs are highly reproducible. A. Two
examples of gene program co-occurrence patterns. The program co-occurrence scores for each
program pair are calculated in the discovery and the validation group separately for comparison
between datasets. **B.** Program co-occurrence agreement between discovery and validation.
Each dot denotes one gene program pair. Co-occurrence per program pair was measured as
Spearman correlations between sample level loading means. **C.** Within cell type gene program
co-occurrence agreement between discovery and validation. **D.** Across cell type gene program
co-occurrence agreement between discovery and validation. **E.** Gene programs ranked by co-
occurrence agreement between discovery and validation. Programs whose co-occurrence
patterns are more similar between discovery and validation have higher co-occurrence

25

reproducibility. Asterisk signs denote gene programs whose co-occurrence with other programs

is significantly reproducible between discovery and validation. The grey bars below indicate

empirical positive predictive value for co-occurrence of the respective gene programs. **F-G,**

Sample level co-occurrence between the CD4 regulatory T cell gene program and other

programs (**F**), and between the germinal center B cell program and other programs (**G**).

Figure 3: Multicellular ecosystems discovered by a probabilistic model reflect spatial

variation. A. Gene program presence in validated ecosystems. **B**. Correlations of ecosystems

between discovery and validation datasets. **C.** Workflow to validate scRNA atlas ecosystems

using Visium data. **D**. Spatial autocorrelation of validated scRNA-seq inferred ecosystems

compared to randomly generated ecosystems. **E**. Spot-wise correlation between scRNA

ecosystems and NMF factors computed using spot-wise gene program loadings. **F**. Examples of

ecosystem loadings in tumor samples.

Figure 4: Multicellular ecosystems and cell type-specific programs are associated with patient outcomes and tumour genetics. **A**. The KRAS amplification epithelial program predicts patient survival in the Chan cohort (left) and the TCGA cohort (right). **B**. Dichotomization of patient survival by expression subtypes in the Chan and TCGA PDAC cohort. **C**. Association between ecosystems/programs and therapy outcome in the Chan cohort. **D.** Association between ecosystems/programs and therapy outcome in the TCGA cohort. **E**. Relationships between KRAS mutant and ecosystem loading in tumour samples. '*': p-value < 0.05; '**': p-value < 0.01, '***': p-value < 0.001 (all Benjamini-Hochberg FDR corrected).

References

1. Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nat. Commun.* **8**, 14049 (2017).

2. Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell* **161**, 1202–1214 (2015).

3. Zhang, S. *et al.* Single cell transcriptomic analyses implicate an immunosuppressive tumor microenvironment in pancreatic cancer liver metastasis. *Nat. Commun.* **14**, 5123 (2023).

4. Cui Zhou, D. *et al.* Spatially restricted drivers and transitional cell populations cooperate with the microenvironment in untreated and chemo-resistant pancreatic cancer. *Nat. Genet.* **54**, 1390–1405 (2022).

5. Hwang, W. L. *et al.* Single-nucleus and spatial transcriptome profiling of pancreatic cancer identifies multicellular dynamics associated with neoadjuvant treatment. *Nat. Genet.* **54**, 1178–1191 (2022).

6. McAndrews, K. M. *et al.* Identification of functional heterogeneity of carcinoma-associated fibroblasts with distinct IL6-mediated therapy resistance in pancreatic cancer. *Cancer Discov.* **12**, 1580–1597 (2022).

7. Chijimatsu, R. *et al.* Establishment of a reference single-cell RNA sequencing dataset for human pancreatic adenocarcinoma. *iScience* **25**, 104659 (2022).

8. Falcomatà, C. *et al.* Selective multi-kinase inhibition sensitizes mesenchymal pancreatic cancer to immune checkpoint blockade by remodeling the tumor microenvironment. *Nat. Cancer* **3**, 318–336 (2022).

9. Giesen, C. *et al.* Highly multiplexed imaging of tumor tissues with subcellular resolution by mass cytometry. *Nat. Methods* **11**, 417–422 (2014).

10. Angelo, M. *et al.* Multiplexed ion beam imaging of human breast tumors. *Nat. Med.* **20**,

436–442 (2014).

11. Janesick, A. *et al.* High resolution mapping of the tumor microenvironment using integrated

single-cell, spatial and in situ analysis. *Nat. Commun.* **14**, 8353 (2023).

12. He, S. *et al.* High-plex imaging of RNA and proteins at subcellular resolution in fixed tissue

by spatial molecular imaging. *Nat. Biotechnol.* **40**, 1794–1806 (2022).

13. Jackson, H. W. *et al.* The single-cell pathology landscape of breast cancer. *Nature* **578**,

615–620 (2020).

14. Kim, S. *et al.* Integrative analysis of spatial and single-cell transcriptome data from human

pancreatic cancer reveals an intermediate cancer cell population associated with poor

prognosis. *Genome Med.* **16**, 20 (2024).

15. Moncada, R. *et al.* Integrating microarray-based spatial transcriptomics and single-cell

RNA-seq reveals tissue architecture in pancreatic ductal adenocarcinomas. *Nat.*

Biotechnol. **38**, 333–342 (2020).

16. Storrs, E. P. *et al.* High-dimensional deconstruction of pancreatic cancer identifies tumor

microenvironmental and developmental stemness features that predict survival. *NPJ*

Precis. Oncol. **7**, 105 (2023).

17. Luca, B. A. *et al.* Atlas of clinically distinct cell states and ecosystems across human solid

tumors. *Cell* **184**, 5482-5496.e28 (2021).

18. Wu, S. Z. *et al.* A single-cell and spatially resolved atlas of human breast cancers. *Nat.*

Genet. **53**, 1334–1347 (2021).

19. Sorin, M. *et al.* Single-cell spatial landscapes of the lung tumour immune microenvironment.

Nature **614**, 548–554 (2023).

20. Ni, Z. *et al.* SpotClean adjusts for spot swapping in spatial transcriptomics data. *Nat.*

Commun. **13**, 2971 (2022).

21. Pijuan-Sala, B. *et al.* A single-cell molecular map of mouse gastrulation and early

organogenesis. *Nature* **566**, 490–495 (2019).

22. Smillie, C. S. *et al.* Intra- and inter-cellular rewiring of the human colon during ulcerative

colitis. *Cell* **178**, 714-730.e22 (2019).

23. Büttner, M., Ostner, J., Müller, C. L., Theis, F. J. & Schubert, B. scCODA is a Bayesian

model for compositional single-cell data analysis. *Nat. Commun.* **12**, 6876 (2021).

24. Rozenblatt-Rosen, O. *et al.* Building a high-quality Human Cell Atlas. *Nat. Biotechnol.* **39**,

149–153 (2021).

25. Rozenblatt-Rosen, O. *et al.* The Human Tumor Atlas Network: Charting tumor transitions

across space and time at single-cell resolution. *Cell* **181**, 236–249 (2020).

26. Terekhanova, N. V. *et al.* Epigenetic regulation during cancer transitions across 11 tumour

types. *Nature* **623**, 432–441 (2023).

27. Heiser, C. N. *et al.* Molecular cartography uncovers evolutionary and microenvironmental

dynamics in sporadic colorectal tumors. *Cell* **186**, 5620-5637.e16 (2023).

28. Johnson, B. E. *et al.* An omic and multidimensional spatial atlas from serial biopsies of an

evolving metastatic breast cancer. *Cell Rep. Med.* **3**, 100525 (2022).

29. Yanagawa, J. *et al.* Single-cell characterization of pulmonary nodules implicates

suppression of immunosurveillance across early stages of lung adenocarcinoma. *Cancer*

Res. **83**, 3305–3319 (2023).

30. Glasner, A. *et al.* Conserved transcriptional connectivity of regulatory T cells in the tumor

microenvironment informs new combination cancer therapy strategies. *Nat. Immunol.* **24**,

1020–1035 (2023).

31. Chan, J. M. *et al.* Signatures of plasticity, metastasis, and immunosuppression in an atlas

of human small cell lung cancer. *Cancer Cell* **39**, 1479-1496.e18 (2021).

32. Chan-Seng-Yue, M. *et al.* Transcription phenotypes of pancreatic cancer are driven by

genomic events during tumor evolution. *Nat. Genet.* **52**, 231–240 (2020).

33. Peng, J. *et al.* Single-cell RNA-seq highlights intra-tumoral heterogeneity and malignant

progression in pancreatic ductal adenocarcinoma. *Cell Res.* **29**, 725–738 (2019).

34. Steele, N. G. *et al.* Multimodal mapping of the tumor and peripheral blood immune

landscape in human pancreatic cancer. *Nature Cancer* **1**, 1097–1112 (2020).

35. Werba, G. *et al.* Single-cell RNA sequencing reveals the effects of chemotherapy on human

pancreatic adenocarcinoma and its tumor microenvironment. *Nat. Commun.* **14**, 797

(2023).

36. Lee, J. J. *et al.* Elucidation of tumor-stromal heterogeneity and the ligand-receptor

interactome by single-cell transcriptomics in real-world pancreatic cancer biopsies. *Clin.*

Cancer Res. **27**, 5912–5921 (2021).

37. Lin, W. *et al.* Single-cell transcriptome analysis of tumor and stromal compartments of

pancreatic ductal adenocarcinoma primary tumors and metastatic lesions. *Genome Med.*

12, 80 (2020).

38. Chen, K. *et al.* Single-cell RNA-seq reveals dynamic change in tumor microenvironment

during pancreatic ductal adenocarcinoma malignant progression. *eBioMedicine* **66**, (2021).

39. Öhlund, D. *et al.* Distinct populations of inflammatory fibroblasts and myofibroblasts in

pancreatic cancer. *J. Exp. Med.* **214**, 579–596 (2017).

40. Zhang, T., Ren, Y., Yang, P., Wang, J. & Zhou, H. Cancer-associated fibroblasts in

pancreatic ductal adenocarcinoma. *Cell Death Dis.* **13**, 1–11 (2022).

41. Caronni, N. *et al.* IL-1β+ macrophages fuel pathogenic inflammation in pancreatic cancer.

Nature **623**, 415–422 (2023).

42. LaRue, M. M. *et al.* Metabolic reprogramming of tumor-associated macrophages by

collagen turnover promotes fibrosis in pancreatic cancer. *Proceedings of the National*

Academy of Sciences **119**, e2119168119 (2022).

43. Cassetta, L. & Pollard, J. W. Targeting macrophages: therapeutic approaches in cancer.

Nat. Rev. Drug Discov. **17**, 887–904 (2018).

44. Moffitt, R. A. *et al.* Virtual microdissection identifies distinct tumor- and stroma-specific

subtypes of pancreatic ductal adenocarcinoma. *Nat. Genet.* **47**, 1168–1178 (2015).

45. Oh, K. *et al.* Coordinated single-cell tumor microenvironment dynamics reinforce pancreatic

cancer subtype. *Nat. Commun.* **14**, 1–13 (2023).

46. Grünwald, B. T. *et al.* Spatially confined sub-tumor microenvironments in pancreatic cancer.

Cell **184**, 5577-5592.e18 (2021).

47. Chen, K. *et al.* Immune profiling and prognostic model of pancreatic cancer using

quantitative pathology and single-cell RNA sequencing. *J. Transl. Med.* **21**, 210 (2023).

48. Welch, J. D. *et al.* Single-Cell Multi-omic Integration Compares and Contrasts Features of

Brain Cell Identity. *Cell* **177**, 1873-1887.e17 (2019).

49. Raghavan, S. *et al.* Microenvironment drives cell state, plasticity, and drug response in

pancreatic cancer. *Cell* **184**, 6119-6137.e26 (2021).

50. Elyada, E. *et al.* Cross-species single-cell analysis of pancreatic ductal adenocarcinoma

reveals antigen-presenting cancer-associated fibroblasts. *Cancer Discov.* **9**, 1102–1123

(2019).

51. Kunes, R. Z., Walle, T., Land, M., Nawy, T. & Pe'er, D. Supervised discovery of

interpretable gene programs from single-cell data. *Nat. Biotechnol.* **42**, 1084–1095 (2024).

52. Kiselev, V. Y., Andrews, T. S. & Hemberg, M. Challenges in unsupervised clustering of

single-cell RNA-seq data. *Nat. Rev. Genet.* **20**, 273–282 (2019).

53. Kuhn, H. W. The Hungarian method for the assignment problem. *Nav. Res. Logist. Q.* **2**,

83–97 (1955).

54. Janssen, P. *et al.* The effect of background noise and its removal on the analysis of single-

cell expression data. *Genome Biol.* **24**, 140 (2023).

55. Helms, E., Onate, M. K. & Sherman, M. H. Fibroblast heterogeneity in the pancreatic tumor

microenvironment. *Cancer Discov.* **10**, 648–656 (2020).

56. Humpton, T. J. *et al.* Oncogenic KRAS induces NIX-mediated mitophagy to promote

pancreatic cancer. *Cancer Discov.* **9**, 1268–1287 (2019).

57. Shi, D. D. *et al.* Therapeutic avenues for cancer neuroscience: translational frontiers and

clinical opportunities. *Lancet Oncol.* **23**, e62–e74 (2022).

58. Rossi Sebastiano, M. *et al.* ACSL3-PAI-1 signaling axis mediates tumor-stroma cross-talk

promoting pancreatic cancer progression. *Sci. Adv.* **6**, eabb9200 (2020).

59. Loncle, C. *et al.* The pancreatitis-associated protein VMP1, a key regulator of inducible

autophagy, promotes KrasG12D-mediated pancreatic cancer initiation. *Cell Death Dis.* **7**,

e2295–e2295 (2016).

60. Gao, H. *et al.* The expression of versican and its role in pancreatic neuroendocrine tumors.

Pancreatology **20**, 142–147 (2020).

61. Li, K. *et al.* Myeloid-derived suppressor cells as immunosuppressive regulators and

therapeutic targets in cancer. *Signal Transduction and Targeted Therapy* **6**, 1–25 (2021).

62. Bärthel, S., Falcomatà, C., Rad, R., Theis, F. J. & Saur, D. Single-cell profiling to explore

pancreatic cancer heterogeneity, plasticity and response to therapy. *Nat. Cancer* **4**, 454–

467 (2023).

63. Gulhati, P. *et al.* Targeting T cell checkpoints 41BB and LAG3 and myeloid cell

CXCR1/CXCR2 results in antitumor immunity and durable response in pancreatic cancer.

Nat. Cancer **4**, 62–80 (2023).

64. Goveia, J. *et al.* An Integrated Gene Expression Landscape Profiling Approach to Identify

Lung Tumor Endothelial Cell Heterogeneity and Angiogenic Candidates. *Cancer Cell* **37**,

421 (2020).

65. Geldhof, V. *et al.* Single cell atlas identifies lipid-processing and immunomodulatory

endothelial cells in healthy and malignant breast. *Nat. Commun.* **13**, 1–19 (2022).

66. Nieto, P. *et al.* A single-cell tumor immune atlas for precision oncology. *Genome Res.* **31**,

1913–1926 (2021).

67. Fridman, W. H. *et al.* B cells and tertiary lymphoid structures as determinants of tumour

immune contexture and clinical outcome. *Nat. Rev. Clin. Oncol.* **19**, 441–457 (2022).

68. Gorgun, C. *et al.* Synergistic role of three dimensional niche and hypoxia on conservation of

cancer stem cell phenotype. *Int. J. Biol. Macromol.* **90**, 20–26 (2016).

69.   Liu, J. *et al.* An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality

Survival Outcome Analytics. *Cell* **173**, 400-416.e11 (2018).

70.   Cancer Genome Atlas Research Network. Electronic address:

andrew_aguirre@dfci.harvard.edu & Cancer Genome Atlas Research Network. Integrated

Genomic Characterization of Pancreatic Ductal Adenocarcinoma. *Cancer Cell* **32**, 185-

203.e13 (2017).

71.   Till, J. E. *et al.* Circulating KRAS G12D but not G12V is associated with survival in

metastatic pancreatic ductal adenocarcinoma. *Nat. Commun.* **15**, 1–12 (2024).

72.   Leidner, R. *et al.* Neoantigen T-cell receptor gene therapy in pancreatic cancer. *N. Engl. J.*

Med. **386**, 2112–2119 (2022).

73.   Uhlen, M. *et al.* A pathology atlas of the human cancer transcriptome. *Science* **357**, (2017).

74.   Marisa, L. *et al.* Gene expression classification of colon cancer into molecular subtypes:

Characterization, validation, and prognostic value. *PLoS Med.* **10**, e1001453 (2013).

75.   Győrffy, B., Surowiak, P., Budczies, J. & Lánczky, A. Online survival analysis software to

assess the prognostic value of biomarkers using transcriptomic data in non-small-cell lung

cancer. *PLoS One* **8**, e82241 (2013).

76.   Calon, A. *et al.* Stromal gene expression defines poor-prognosis subtypes in colorectal

cancer. *Nat. Genet.* **47**, 320–329 (2015).

77.   Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in primary

glioblastoma. *Science* **344**, 1396–1401 (2014).

78.   Zilionis, R. *et al.* Single-Cell Transcriptomics of Human and Mouse Lung Cancers Reveals

Conserved Myeloid Populations across Individuals and Species. *Immunity* **50**, 1317-

1334.e10 (2019).

79.   Savas, P. *et al.* Single-cell profiling of breast cancer T cells reveals a tissue-resident

memory subset associated with improved prognosis. *Nat. Med.* **24**, 986–993 (2018).

80. van der Leun, A. M., Thommen, D. S. & Schumacher, T. N. CD8+ T cell states in human

cancer: insights from single-cell analysis. *Nat. Rev. Cancer* **20**, 218–232 (2020).

81. Kang, J. *et al.* Systematic dissection of tumor-normal single-cell ecosystems across a

thousand tumors of 30 cancer types. *Nat. Commun.* **15**, 1–17 (2024).

82. McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-

Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-337.e4

(2019).

83. McCarthy, D. J., Campbell, K. R., Lun, A. T. L. & Wills, Q. F. Scater: pre-processing, quality

control, normalization and visualization of single-cell RNA-seq data in R. *Bioinformatics* **33**,

1179–1186 (2017).

84. Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a

transitional profibrotic macrophage. *Nat. Immunol.* **20**, 163–172 (2019).

85. Mölder, F. *et al.* Sustainable data analysis with Snakemake. *F1000Res.* **10**, 33 (2021).

86. Wang, Y. *et al.* Interactive analysis of single-cell data using flexible workflows with SCTK2.

Patterns (N Y) **4**, 100814 (2023).

87. Gaujoux, R. & Seoighe, C. A flexible R package for nonnegative matrix factorization. *BMC*

Bioinformatics **11**, 367 (2010).

88. Shen, L. GeneOverlap: An R package to test and visualize gene overlaps. *R Package*

(2016).

89. Nurk, S. *et al.* The complete sequence of a human genome. *Science* **376**, 44–53 (2022).

90. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.

Innovation (Camb.) **2**, 100141 (2021).

91. Gavish, A. *et al.* Hallmarks of transcriptional intratumour heterogeneity across a thousand

tumours. *Nature* **618**, 598–606 (2023).

92. Korotkevich, G. *et al.* Fast gene set enrichment analysis. *bioRxiv* 060012 (2021)

doi:10.1101/060012.

93. Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for

microarray and RNA-seq data. *BMC Bioinformatics* **14**, 7 (2013).

94. Jonah Gabry and Rok Češnovar and Andrew Johnson and Steve Bronder. cmdstanr: R

Interface to "CmdStan." https://mc-stan.org/cmdstanr/ (2023).

95. de Bruijn, I. *et al.* Sharing data from the Human Tumor Atlas Network through standards,

infrastructure and community engagement. *Nat. Methods* **22**, 664–671 (2025).

96. Marconato, L. *et al.* SpatialData: an open and universal data framework for spatial omics.

Nat. Methods **22**, 58–62 (2025).

97. Palla, G. *et al.* Squidpy: a scalable framework for spatial omics analysis. *Nat. Methods* **19**,

171–178 (2022).

1054