CARD8 inflammasome activation during HIV-1 cell-to-cell transmission

Jessie Kulsuptrakul[1,2], Michael Emerman[2,*], Patrick S. Mitchell[3,4*]
[1]Molecular and Cellular Biology Graduate Program, University of Washington, Seattle, WA 98195, USA

[2]Divisions of Human Biology and Basic Sciences, Fred Hutchinson Cancer Center, Seattle, WA 98109, USA

[3]Department of Microbiology, University of Washington, Seattle, WA 98109, USA

[4]Howard Hughes Medical Institute, University of Washington, Seattle, WA, 98109, USA

*Corresponding authors: memerman@fredhutch.org, psmitche@uw.edu

Abstract

Our previous work demonstrated that CARD8 detects HIV-1 infection by sensing the enzymatic

activity of the HIV protease, resulting in CARD8-dependent inflammasome activation

(Kulsuptrakul et al., 2023). CARD8 harbors a motif in its N-terminus that functions as a HIV

protease substrate mimic, permitting innate immune recognition of HIV-1 protease activity,

which when cleaved by HIV protease triggers CARD8 inflammasome activation. Here, we

sought to understand CARD8 responses in the context of HIV-1 cell-to-cell transmission via a

viral synapse. We observed that cell-to-cell transmission of HIV-1 between infected T cells and

primary human monocyte-derived macrophages induces CARD8 inflammasome activation in a

manner that is dependent on viral protease activity and largely independent of the NLRP3

inflammasome. Additionally, to further evaluate the viral determinants of CARD8 sensing, we

tested a panel of HIV protease inhibitor resistant clones to establish how variation in HIV

protease affects CARD8 activation. We identified mutant HIV-1 proteases that differentially

cleave and activate CARD8 compared to wildtype HIV-1, thus indicating that natural variation in

HIV protease affects not only the cleavage of the viral Gag-Pol polyprotein but also likely

impacts innate sensing and inflammation.

INTRODUCTION

HIV-1 disease progression is characterized by chronic inflammation, immune activation,

CD4+ T cell depletion and eventual destruction of the immune system and susceptibility to

opportunistic infections. The primary cellular targets of HIV-1 are activated CD4+ T helper cells,

specialized CD4+ T cell subtypes such as Th17 cells (Brenchley et al., 2008; Gosselin et al.,

2009; Rodriguez-Garcia et al., 2014), central memory cells (Chun et al., 1997b, 1997a, 1995),

and macrophages (Collman et al., 1990, 1989). Chronic immune activation is primarily caused

by rapid depletion of mucosal Th17 cells responsible for maintaining gut epithelial barrier

integrity (Brenchley et al., 2008, 2006). In addition to inflammation induced by circulating

microbial ligands, inflammation can also originate from HIV-infected cells through activation of

innate immune sensors that form cytosolic immune complexes known as inflammasomes.

Inflammasome activation ultimately results in activation of pro-inflammatory caspases including

caspase 1 (CASP1). Active CASP1 processes inflammatory cytokines and activates the pore-

forming protein gasdermin D (GSDMD), which forms small pores in the plasma membrane and

initiates a lytic form of cell death known as pyroptosis and the release of mature inflammatory

cytokines interleukin (IL)-1β and IL-18 (Broz and Dixit, 2016; Fink and Cookson, 2005).

In prior work, we and others showed that the inflammasome-forming sensor CARD8

senses HIV-1 infection through the detection of HIV-1 protease (HIV^{PR}) activity (Clark et al.,

2022; Kulsuptrakul et al., 2023; Wang et al., 2021). While the canonical function of HIV^{PR} is to

cleave viral polyproteins during virion maturation, active HIV^{PR} is also released into the host cell,

which is sensed by CARD8 via HIV^{PR} cleavage of its N-terminus and subsequent inflammasome

activation. In this way, the CARD8 N-terminus functions as a "molecular tripwire" to recognize

the enzymatic activity of HIV^{PR} and other viral proteases (Castro and Daugherty, 2023; Nadkarni

et al., 2022; Tsu et al., 2023). Moreover, HIV^{PR} cleavage of CARD8 occurs rapidly after infection

such that HIV^{PR} inhibitors and fusion inhibitors, but not reverse transcriptase (RT) inhibitors can

prevent CARD8 inflammasome activation, implying that CARD8 detects HIV-1 viral protease

activity of virion-packaged or "incoming" HIV[PR] upon virion fusion (Kulsuptrakul et al., 2023;

Wang et al., 2024, 2021). Interestingly, CARD8 inflammasome activation in resting CD4+ T cells

results in pyroptosis but not the release of pro-inflammatory cytokines IL-1β or IL-18 (Wang et

al., 2024), suggesting that CARD8 inflammasome activation in T cells does not directly

contribute to chronic inflammation. Here, we address whether or not CARD8 may influence HIV-

1 pathogenesis through the maturation and release of IL-1β from infected macrophages.

HIV-1 can be transmitted from one cell to another via two main mechanisms: "cell-free"

infection through binding of free HIV-1 virions to target cells, and cell-to-cell infection whereby

infected cells directly transfer virus to an uninfected target cell via the formation of a transient

viral synapse (Chen et al., 2007; Galloway et al., 2015; Iwami et al., 2015). Cell-to-cell

transmission of HIV-1 has been reported between multiple HIV-1 target cell types including

between active and resting CD4+ T cells (Agosto et al., 2018; Martin et al., 2010) and between

CD4+ T cells and macrophages (Baxter et al., 2014; Dupont and Sattentau, 2020; Lopez et al.,

2019). Cell-to-cell transmission delivers a large influx of virus to target cells, resulting in a high

multiplicity of infection (MOI) (Agosto et al., 2015; Del Portillo et al., 2011; Duncan et al., 2013;

Russell et al., 2013), which has been proposed to enhance viral fitness by overwhelming host

restriction factors including Tetherin/BST-2 (Jolly et al., 2010; Zhong et al., 2013), SAMHD1 (Xie

et al., 2019), and TRIM5α (Richardson et al., 2008), and evading adaptive immune responses

including broadly neutralizing antibodies (Abela et al., 2012; Dufloo et al., 2018). Cell-to-cell

spread of HIV-1 is thus an important consideration in studying CARD8 inflammasome activation.

Here, we investigated both host and viral determinants of CARD8 inflammasome

activation upon HIV-1 infection. We evaluated CARD8 sensing of HIV[PR] during cell-to-cell

transmission of HIV-1 from T cell lines to myeloid cells in both immortalized and primary cell

models of infection. We found that CARD8 inflammasome activation occurs in the context of

cell-to-cell transmission from both SUPT1 cells, a T cell lymphoma cell line, to THP-1 cells, an

acute myeloid leukemia cell line, and from primary CD4+ T cells to primary monocyte-derived

macrophages. We also observed that HIV-triggered CARD8 inflammasome activation is largely

independent of the NLRP3 inflammasome, which has previously been implicated in innate

sensing of HIV-1 (Bandera et al., 2018; Chivero et al., 2017; Hernandez et al., 2013; Leal et al.,

2020; Mamik et al., 2017; Zhang et al., 2021). Our findings suggest that CARD8 sensing of

HIV^{PR} activity during cell-to-cell transmission of HIV-1 to macrophages, leading to robust

secretion of IL-1β, may be a source of inflammatory cytokines that promote pathogenic chronic

inflammation and disease progression. In addition, we also show that natural variation in HIV^{PR}

due to resistance to protease inhibitors also affects CARD8 cleavage and subsequent

inflammasome activation. Our results extend the role of incoming HIV^{PR} on CARD8-dependent

inflammasome activation of inflammasome responses as a function of cell type, mode of

transmission, and virus evolution in response to antiviral therapy.

96
RESULTS

Cell-to-cell transmission of HIV-1 induces CARD8 inflammasome activation

Our previous work investigating HIV-dependent CARD8 inflammasome activation used

the cationic polymer DEAE-dextran, which is a common reagent used to enhance viral infection

in cell culture (Bailey et al., 1984). However, we found that DEAE-dextran could induce

inflammasome activation in the absence of viral infection in some "wildtype" (WT) THP-1 cell

stocks (see **Supplemental Note**). These results prompted us to establish other models of HIV-1

infection and subsequent inflammasome activation that lack cationic polymers. Thus, we

designed an *in vitro* coculture infection system to mimic HIV-1 cell-to-cell transmission by

infecting SUPT1 cells, a T-cell lymphoma line (i.e., donor cells) and then mixing them with

uninfected THP-1 cells (i.e., target cells). We opted for SUPT1 cells as the viral producer cell

line because they are permissive to HIV-1 infection, and unlike THP-1 cells, SUPT1 cells do not

respond to a known CARD8 inflammasome activator, ValboroPro (VbP), as assayed by both IL-

1β secretion and cell death, indicating that SUPT1 cells do not have a functional CARD8

inflammasome pathway (**Figure 1A**). This allowed us to infer that inflammasome outputs (e.g.

IL-1β secretion) in our coculture system occur upon cell-to-cell transmission of HIV-1 from

Figure 1. HIV-1 cell-to-cell infection induces inflammasome activation. (**A**) SUPT1 or THP-1 cells were primed with Pam3CSK4 (500ng/mL) overnight then treated with 5μM ValboroPro (VbP) for 24 hours then assessed for IL-1β secretion and cell death via propidium iodide (PI) uptake. %PI positive was normalized to mock control. (**B**) (left) Schematic illustrating experimental setup for SUPT1:THP-1 cell coculture either with (bottom) or without (top) a transwell. (right) SUPT1 cell were either mock infected or infected with HIV-1$_{LAI}$ then cocultured with primed WT THP-1 cells 20 hours post infection. Mock- or HIV-1$_{LAI}$-infected SUPT1 cells were either mixed with the THP-1 cells or put in a transwell with a virus-permeable membrane as shown in panel B (left). Supernatant in the cell-to-cell condition and in the supernatant outside of the transwell were sampled and measured for infectious HIV virions via reverse transcriptase (RT) assay or (**C**) IL-1β secretion 3 days after starting the coculture. Dotted line indicates limit of detection (LoD). Datasets represent mean ± SD (A: n=2; B,C n=4 biological replicates). One-way ANOVA with (B) Tukey's or (C) Dunnett's test using GraphPad Prism 10. ns = not significant, *p<0.05,**p<0.01, ***p<0.001, ****p<0.0001.

SUPT1 cells to the CARD8-competent THP-1 cells.

We found that coculture of THP-1 cells with HIV-1$_{LAI}$-infected SUPT1 cells (**Figure 1B**)

but not mock-infected SUPT1 cells results in robust inflammasome activation as indicated by IL-

1β secretion, suggesting that our coculture system, which lacks DEAE-dextran, can induce HIV-

dependent inflammasome activation via cell-to-cell infection (**Figure 1C**). To further test this

assumption, we prevented cell-to-cell contact using a virus-permeable transwell with a 0.4μm

pore insert (**Figure 1B,** left). We verified that there were equivalent amounts of infectious virus

in the cell-to-cell condition versus the lower chamber of the transwell condition by measuring RT

activity in the supernatant (**Figure 1B,** right). Despite equivalent amounts of infectious virus in

both conditions, we observed that HIV-1$_{LAI}$-infected SUPT1 cells (upper chamber) cocultured

with THP-1 cells (lower chamber) did not lead to detectable IL-1β secretion (**Figure 1C**). These

data suggest that our SUPT1:THP-1 coculture system can trigger HIV-dependent

inflammasome activation in a manner dependent on cell-to-cell contact.

We next assessed the role of CARD8 and other inflammasome sensors during cell-to-

cell transmission of HIV-1. We cocultured mock- or HIV-1$_{LAI}$-infected SUPT1 cells with either WT

or *CARD8* KO THP-1 cells and compared inflammasome activation by measuring levels of

secreted IL-1β. HIV-1$_{LAI}$-infected SUPT1 cells cocultured with WT but not *CARD8* KO THP-1

cells resulted in a significant increase in IL-1β (**Figure 2A**). These results suggest that CARD8

is the primary sensor that drives inflammasome activation in HIV-1 cell-to-cell transmission to

THP-1 cells. Since the NLRP3 inflammasome has previously been implicated in HIV-dependent

inflammasome activation (Bandera et al., 2018; Chivero et al., 2017; Hernandez et al., 2013;

Leal et al., 2020; Mamik et al., 2017; Zhang et al., 2021), we also assessed the effects of the

NLRP3 inflammasome-specific inhibitor MCC950 (Coll et al., 2015; Primiano et al., 2016) on

inflammasome activation in our coculture system. Treatment with MCC950 or the caspase 1

(CASP1) inhibitor VX765 (Wannamaker et al., 2007) were sufficient to abrogate inflammasome

activation induced by the ionophore nigericin, a well-characterized NLRP3 agonist (**Figure 2–**

figure supplement 1A). However, in the HIV-1 coculture system, MCC950 treatment had only

a modest effect on inflammasome activation while VX765 and the HIV^{PR} inhibitor lopinavir

(LPV), which prevents CARD8 cleavage by HIV-1^{PR} (Kulsuptrakul et al., 2023; Wang et al.,

2021), completely abrogated IL-1β secretion (**Figure 2B**). We observed similar results during

HIV-1$_{LAI}$ and HIV-1$_{LAI-VSVG}$ cell-free infection of THP-1 cells in the presence of DEAE-dextran

(**Figure 2– figure supplement 1B**). Taken together, these findings indicate that HIV-dependent

Figure 2. HIV-1 cell-to-cell transmission induces CARD8-dependent activation largely independent of NLRP3. (A) SUPT1 cells were either mock-infected or infected with HIV-1$_{LAI}$ for 18-20 hours prior to coculture with wildtype (WT) or *CARD8* knockout (KO) THP-1 cells. The coculture was harvested 72 hours later to probe for IL-1β secretion in the coculture supernatant via IL-1R reporter assay. THP-1 cells were primed with Pam3CSK4 (500ng/mL) for 16-24 hours prior to coculture. SUPT1 cells were infected with HIV-1$_{LAI}$ such that 30% of the cells were positive for intracellular p24^{gag} after 18-20 hours. **(B)** SUPT1 cells were either mock or HIV-1$_{LAI}$-infected as in (A) for 18-20 hours then incubated in DMSO, lopinavir (LPV), MCC950, or VX765 at 0.01%, 5μM, 10μM, or 1μg/mL, respectively, for 15 minutes prior to coculturing with primed WT THP-1 cells. The coculture was assessed for subsequent inflammasome activation after 72 hours as in (A). Dotted line indicates limit of detection (LoD). Datasets represent mean ± SD (n=3 biological replicates). Two-way ANOVA with Dunnett's test using GraphPad Prism 10. ns = not significant, *p<0.05,**p<0.01, ***p<0.001, ****p<0.0001.

145 inflammasome activation via cell-to-cell transmission is CARD8-dependent and largely NLRP3-

146 independent.

Figure 2– figure supplement 1. HIV-dependent inflammasome activation is largely NLRP3-independent. (**A**) Wildtype THP-1 cells were pre-treated with either DMSO, lopinavir (LPV), MCC950, or VX765 at 0.01%, 5μM, 10μM, or 1μg/mL, respectively, for 15 minutes prior to 4-hour treatment with 5μg/mL nigericin. Subsequent inflammasome activation was assessed via (A, left) IL-1β secretion via IL-1R reporter assay and (A, right) cell death via propidium iodide (PI) dye uptake. (**B**) Wildtype THP-1 cells were pre-treated with indicated inhibitors as in (A) then infected with either HIV-1$_{LAI}$ or VSV-G pseudotyped HIV-1$_{LAI}$ (HIV-1$_{LAI-VSVG}$) in the presence of 10μg/mL DEAE-dextran such that both HIV-1$_{LAI}$-infected and HIV-1$_{LAI-VSVG}$ -infected cells were ~30% positive for intracellular p24^{gag} after 24 hours by flow cytometry. Subsequent inflammasome activation was assessed 24 hours post infection via IL-1β secretion and cell death as in (A). Dotted line indicates limit of detection (LoD). Datasets represent mean ± SD (A: n=3, B: n=2 biological replicates). One-way (A) or Two-way (B) ANOVA with Dunnett's test using GraphPad Prism 10. ns = not significant, *p<0.05,**p<0.01, ***p<0.001, ****p<0.0001.

CARD8, but not NLRP3, **is required for inflammasome activation during HIV-1 cell-to-cell**

transmission into primary monocyte-derived macrophages

We next examined inflammasome activation upon HIV-1 cell-to-cell transmission in

primary human monocyte-derived macrophages (MDMs). Previously, we had observed that

CARD8 could sense active HIV-1^{PR} released into the host cytosol following viral fusion, which

we refer to as "incoming" HIV-1^{PR} in our cell-free infection system in THP-1 cells using DEAE-

dextran and spinoculation (Kulsuptrakul et al., 2023). Thus, we assessed the importance of viral

entry by coculturing MDMs from three independent donors with mock-, HIV-1$_{LAI}$-, or HIV-1$_{NL4.3-}$

$_{BaL}$-infected SUPT1 cells expressing CCR5 (SUPT1-CCR5) and assayed for inflammasome

activation (**Figure 3A**). HIV-1$_{LAI}$ is a CXCR4 tropic strain unable to infect macrophages whereas

HIV-1$_{NL4.3-BaL}$ uses CCR5 as a co-receptor which is a requirement for infection of macrophages.

We observed inflammasome activation, as measured by IL-1β secretion, in MDMs cocultured

with HIV-1$_{NL4.3-BaL}$-infected SUPT1 cells but not in MDMs cocultured with mock- or HIV-1$_{LAI}$-

infected SUPT1-CCR5 cells (**Figure 3A**). This demonstrates that HIV-dependent inflammasome

activation can occur in MDMs during cell-to-cell infection in a manner dependent on viral entry.

To further ascertain if this inflammasome activation was CARD8-dependent and driven by

incoming HIV^{PR} during SUPT1:MDM cell-to-cell transmission, we investigated the effects of

different inhibitors on inflammasome activation in MDM cocultures with HIV-1$_{NL4.3-BaL}$-infected

SUPT1s. We observed that IL-1β secretion was abrogated by treatment with lopinavir, an HIV-1

protease inhibitor, and VX765, a CASP1 inhibitor, indicating that inflammasome activation in

MDM cocultures is dependent on HIV^{PR} and CASP1, respectively (**Figure 3A**). In addition, we

used an RT inhibitor, nevirapine (NVP), to prevent synthesis of *de novo* translated HIV^{PR}, and

thus any CARD8-dependent IL-1β secretion would only be due to incoming HIV^{PR} in the

presence of NVP. Indeed, we observed HIV-dependent inflammasome activation in the

presence of NVP that was added at the time of coculture, indicating that incoming HIV^{PR} is

sufficient to elicit an inflammasome response (**Figure 3A**). Lastly, MDM cocultures treated with

MCC950, an inhibitor of the NLRP3 inflammasome, had no effect on IL-1β secretion (**Figure**

3A), and we observed similar inflammasome activation results regardless of whether the MDMs

were primed with TLR1/2 agonist Pam3CSK4 versus TLR4 agonist lipopolysaccharide (LPS)

(**Figure 3– figure supplement 1A**). Thus, cell-to-cell contact of infected cells with primary

monocyte-derived macrophages can elicit an inflammasome response in a manner that is

dependent on viral entry, CASP1, and incoming HIV^{PR}, and independent from NLRP3.

To further assess the timing of this inflammasome activation, we conducted a

SUPT1:MDM time course coculture experiment with 3 additional donors, assaying IL-1β

secretion at 4, 24, 48, and 72 hours post coculture in the presence and absence of NVP. We

observed inflammasome activation as determined by IL-1β secretion by 24 hours post coculture

in all donors that persisted at a similar level at 48 and 72 hours post coculture (**Figure 3B**). As

expected, we observed donor-to-donor variation in the extent to which IL-1β secretion occurred

following HIV-1 infection. However, per donor, HIV-1-driven IL-1β levels were comparable to

that of VbP-induced inflammasome activation (**Figure 3–figure supplement 1B**). Moreover,

adding NVP had no effect on IL-1β secretion in HIV-1 infected MDM cocultures (**Figure 3B**). To

verify that NVP was functional, we assayed the supernatant of the mock and NVP-treated

cocultures from **Figure 3B** at 48 hours post-coculture for infectious virions via an assay for RT

activity and observed lower RT activity in NVP-treated MDM donors (**Figure 3–figure**

supplement 1C). Unlike the 'cell-free' infection conditions in which we previously observed an

increase in IL-1β levels 4h post-infection, we did not detect measurable differences in IL-1β

secretion at this early timepoint following the establishment of SUPT1:MDM coculture.

Nevertheless, the data are consistent with incoming HIV^{PR} being responsible for inflammasome

activation during cell-to-cell transmission of HIV because the induction of IL-1β persists in the

presence of NVP which would block any *de novo* synthesis of new *gag/pol* products (**Figure 3A**

and B). Taken together, these data suggest that in the context of cell-to-cell transmission,

CARD8 is likely the inflammasome-forming sensor that detects HIV-1 infection via incoming

HIV^{PR} activity in primary monocyte-derived macrophages.

To specifically address the role of CARD8 in HIV-1 induced inflammasome activation in

MDMs, we genetically edited MDMs by isolating monocytes from five donors and

electroporating them with Cas9 RNPs complexed with three unique sgRNAs per gene targeting

AAVS1, a safe harbor locus, *CARD8* or *NLRP3* (only 2 donors for *NLRP3* KO). Edited MDMs

were then differentiated for 6 days prior to evaluating KO efficiency and initiating cocultures with

HIV-1-infected SUPT1 cells. To verify KO efficiency, we immunoblotted with an antibody that

detects the CARD8 C-terminus in *CARD8* KO MDMs relative to the *AAVS1* KO control and

observed a marked reduction of the full-length and FIIND-processed CARD8 protein in all 5

donors (**Figure 3C**). In addition, we confirmed *AAVS1*, *CARD8*, and *NLRP3* KO at the genetic

level via Synthego ICE analysis (Conant et al., 2022), measuring >85% KO efficiency (**Figure**

3C). We also observed robust inflammasome activation upon treatment with CARD8

inflammasome activator VbP as measured by IL-1β secretion in *AAVS1* KO MDMs from 2 of the

3 donors, which was completely abrogated in *CARD8* KO MDMs, confirming functional loss of

CARD8 (**Figure 3– figure supplement 1D**). We then cocultured either *AAVS1* KO, *CARD8* KO,

or *NLRP3* KO MDMs with mock or HIV-1$_{NL4.3-BaL}$-infected SUPT1-CCR5 cells at a 1:1 ratio and

measured inflammasome activation via IL-1β secretion 48 hours post-coculture. In all 5 donors,

we observed significant reduction in inflammasome activation in *CARD8* KO cocultures relative

to the *AAVS1* KO control (**Figure 3D**). Consistent with our findings with SUPT1:THP-1

coculture, we observed no difference in inflammasome activation when coculturing infected

SUPT1 cells with *AAVS1* KO vs *NLRP3* KO MDMs, suggesting that the inflammasome

activation was also largely NLRP3-independent (**Figure 3D**). Taken together, these data

demonstrate that CARD8 is required for inflammasome activation in MDMs during HIV-1 cell-to

cell transmission.

Figure 3. Cell-to-cell HIV infection induces CARD8-dependent inflammasome activation in monocyte-derived macrophages (MDMs). (**A**) MDMs from 3 independent donors were cocultured with SUPT1 cells expressing CCR5 (SUPT1-CCR5) that were mock-, HIV-1$_{LAI}$- or HIV-1$_{NL4.3-BaL}$-infected then assayed for inflammasome activation 48 hours post-coculture for IL-1β secretion. Fifteen minutes before starting the coculture, SUPT1-CCR5 cells infected with HIV-1$_{NL4.3-BaL}$ were pre-treated with either DMSO, lopinavir (5μM), nevirapine (50μM), MCC950 (10μM) or VX765 (1μg/mL), inhibiting HIV-1 protease (HIV^{PR}), HIV-1 reverse transcriptase (HIV^{RT}), NLRP3, or caspase 1 (CASP1), respectively. (**B**) MDMs from 3 independent donors were cocultured with SUPT1-CCR5 cells infected with HIV-1$_{NL4.3-BaL}$ in either the presence or absence of nevirapine (NVP). Supernatant was harvested at 4, 24, 48, or 72 hours to assay for IL-1β secretion. (**C**) MDMs from 5 independent donors were knocked out (KO) for *AAVS1 CARD8,* or *NLRP3* using a Synthego gene KO kit then immunoblotted using an anti-CARD8 antibody or anti-vinculin. Full-length and FIIND-processed CARD8 intermediates are marked with a purple arrow. Table between CARD8 and vinculin blot shows Synthego gene KO% scores for each donor KO line. (**D**) *AAVS1*, *CARD8* or *NLRP3* KO MDM lines from (C) were primed with Pam3CSK4 (500ng/mL) overnight and then cocultured with SUPT1-CCR5 cells mock-, or HIV-1$_{NL4.3-BaL}$-infected then assayed for inflammasome activation 48 hours post-coculture for IL-1β secretion. For all SUPT1:MDM experiments, SUPT1-CCR5 cells were infected with HIV-1$_{LAI}$ or HIV-1$_{NL4.3-BaL}$ such that 5-20% of cells were positive for intracellular p24^{gag} after 20 hours. IL-1 levels shown were normalized to the SUPT1 mock-infected coculture control. Dotted line indicates limit of detection (LoD). Datasets represent mean ± SD (A: n=3 independent donors, B: n=2 biological replicates for each donor, D: n=3 technical replicates per donor). One-way ANOVA with (A)Tukey's or (D) Sidak's test or (B) two-way ANOVA with Tukey's test using GraphPad Prism 10. ns = not significant, *p<0.05,**p<0.01, ***p<0.001, ****p<0.0001.

Figure 3– figure supplement 1: Cell-to-cell HIV infection induces CARD8-dependent activation in monocyte-derived macrophages (MDMs). (**A**) MDMs from donor 6 were primed with either 500ng/mL Pam3CSK4 or 5µg/mL LPS then cocultured with SUPT1-CCR5 cells that had been infected with HIV-1$_{NL4.3-BaL}$ 24 hours prior to coculture. Each coculture was started in the presence of DMSO, nevirapine (NVP) or lopinavir (LPV). Supernatant was harvested 72 hours post coculture to assay for IL-1β secretion. (**B**) MDMs from the same 3 independent blood donors assayed in Figure 3B were primed overnight with Pam3CSK4 then treated with 10µM VbP for 24 hours before assaying for IL-1β secretion via IL-1 reporter assay. (**C**) Supernatant from SUPT1:MDM coculture experiment done in Figure 3B was harvested at 48 hours post coculture to assay for infectious virions via reverse transcriptase (RT) assay. (**D**) *AAVS1* or *CARD8* KO MDMs from donor 7-9 assayed in Figure 3C were primed and treated with VbP for 24 hours then assayed for IL-1β secretion. IL-1 levels from VbP treatment were normalized to untreated mock control. Dotted line indicates limit of detection (LoD). Datasets represent mean ± SD (A: n=2 biological replicates for one donor, B-C: n=2 technical replicates for each independent donor, D: n=3 technical replicates per donor). Two-way ANOVA with (A) Sidak's or (C) Dunnett's test (using GraphPad Prism 10. ns = not significant, *p<0.05,**p<0.01, ***p<0.001, ****p<0.0001.

226
227

Figure 4. Primary CD4+ T cell:MDM coculture elicits CARD8-dependent inflammasome activation. (A) CD4+ T cells from a blood donor were isolated, activated, and either mock infected or infected with HIV-1$_{NL4.3-BaL}$ for 3 days such that ~10% of cells were positive for intracellular p24^{gag}. MDMs were primed with Pam3CSK4 then cocultured with mock or HIV-1 infected primary CD4 T cells in the presence or absence of lopinavir (LPV), nevirapine (NVP), or VX765, inhibiting HIV protease, reverse transcriptase, or caspase 1, respectively. Supernatants were harvest 3 days post coculture to assay for IL-1β secretion via IL-1 reporter assay. (**B**) CD4+ T cells from donor 12 and MDMs from donor 14 were cocultured as in A in the presence or absence of LPV. Supernatant was harvested at 4, 24 and 48 hours post coculture to probe for IL-1β secretion. (**C**) *AAVS1* or *CARD8* MDM KOs were immunoblotted using an anti-CARD8 antibody or anti-vinculin. Full-length and FIIND-processed CARD8 intermediates are marked with a purple arrow. (**D**) *AAVS1* or *CARD8* KO MDMs from (C) were cocultured with CD4+ T cells infected with HIV-1$_{NL4.3-BaL}$ then assayed for IL-1β secretion 48 hours post coculture. The donor 12 cocultures consisted of autologous CD4s and MDMs whereas the MDMs from donors 13-15 were cocultured with donor 12 CD4s. Dotted line indicates limit of detection (LoD). Datasets represent mean ± SD (A,D: n=2 technical replicates for each donor, B: n=3 technical replicates for one donor). (A) Two-way ANOVA with Tukey's test (D) One-way ANOVA with Sidak's test using GraphPad Prism 10. ns = not significant, *p<0.05,**p<0.01, ***p<0.001, ****p<0.0001.

Coculture of HIV-1 infected primary CD4+ T cells with primary MDMs elicits CARD8-dependent inflammasome activation

We next investigated inflammasome activation in the context of cell-to-cell infection using primary CD4+ T cells as donor cells, rather than SUPT1 cells, and primary MDMs as target cells. Mock or HIV-1$_{NL4.3-BaL}$ infected CD4+ T cells were cocultured with MDMs in the presence or absence of various inhibitors (as in **Figure 3A**). We assessed inflammasome activation via IL-1β secretion 72 hours post coculture and observed inflammasome activation when coculturing with HIV-infected T cells but not mock infected T cells or cocultures treated with LPV or VX765, demonstrating that inflammasome activation is driven by HIV^{PR} and CASP1 (**Figure 4A**) and consistent with our findings using the SUPT1:MDM coculture (**Figure 3**). We also observed that inflammasome activation persisted in NVP-treated cocultures, suggesting that incoming protease is also important for inflammasome activation in the context of cell-to-cell transmission of HIV-1 from primary CD4+ T cells to primary MDMs (**Figure 4A**). To confirm the potency of LPV and NVP, we assayed for infectious virions in the supernatant of these CD4:MDM cocultures and detected a dramatic decrease in RT activity in the presence of either of these drugs, indicating that these drugs were efficacious at this dose (**Figure 4– figure supplement 1A**). We also conducted a time course experiment with a CD4:MDM coculture from an independent donor in the presence or absence of LPV. Similar to the SUPT1:MDM time course (**Figure 3B**), we were able to detect elevated levels of IL-1β by 24 hours post coculture, which was again strictly dependent on the enzymatic activity of the viral protease as LPV treatment completely inhibited IL-1β secretion (**Figure 4B**).

To interrogate the specific role of CARD8 in this primary CD4:MDM coculture system, we generated *AAVS1* (as a control) or *CARD8* KO MDMs and cocultured the MDMs with either mock or HIV-infected primary CD4+ T cells then assayed for inflammasome activation 48 hours post coculture. KO efficiency was confirmed via immunoblot and functional response to VbP (in one of the two donors) (**Figure 4C, Figure 4–figure supplement 1B**). We detected a significant

decrease in inflammasome activation when infected T cells were cocultured with *CARD8* KO

versus *AAVS1* KO MDMs (**Figure 4D**), indicating that CARD8 is required for inflammasome

activation in MDMs during HIV-1 transmission from primary CD4+ T cells to MDMs. Taken

together, our data indicate that CARD8 plays a pivotal role in sensing and responding to HIV-1

cell-to-cell infection between primary CD4 T cells and macrophages.

Protease inhibitor resistant strains of HIV-1 differentially cleave and activate CARD8

The consequences of CARD8 inflammasome activation on viral replication have been

challenging to assess given that viral fitness is intrinsically linked to viral protease processing of

the viral polyprotein such that inhibiting HIV^{PR} also prevents viral replication. In an attempt to

Figure 4– figure supplement 1. Primary CD4 T cell:MDM coculture elicits CARD8-dependent inflammasome activation. (**A**) Supernatant was harvested 72 hours post coculture from coculture described in Figure 4A then assayed for infectious virions via (RT) transcriptase assay. (**B**) *AAVS1* or *CARD8* KO MDMs from donor 12 were primed with Pam3CSK4 then treated with VbP for 24 hours and probed for IL-1β secretion. (n=2 technical replicates for each donor). (A) Two-way ANOVA with Tukey's test (B) One-way ANOVA with Sidak's test using GraphPad Prism 10. ns = not significant, *p<0.05,**p<0.01, ***p<0.001, ****p<0.0001.

circumvent this issue, we surveyed a panel of multi-HIV[PR] inhibitor-resistant (PI-R) infectious

molecular clones of HIV-1 (Varghese et al., 2013). This panel of PI-R molecular clones vary in

resistance to HIV protease inhibitors including nelfinavir (NFV), fosamprenavir (FPV), saquinavir

(SQV), indinavir (IDV), atazanavir (ATV), lopinavir (LPV), tipranavir (TPV), and darunavir (DRV).

Each molecular clone encodes 4 to 11 mutations in HIV[PR] as well as various compensatory

HIV[gag] mutations (Varghese et al., 2013) (**Table S1**).

We initially tested if PI-R HIV-1 proviruses differentially cleave CARD8 by co-transfecting

HEK293T cells with an expression plasmid encoding an N-terminal mCherry tagged human

CARD8 and either wildtype HIV-1$_{LAI}$ or PI-R HIV-1 proviruses. HIV-1$_{LAI}$ protease cleaves CARD8

between phenylalanine (F) 59 and F60 (Wang et al 2021), resulting in a ~33kDa product

(**Figure 5A,** top). By quantifying the 33kDa CARD8 cleavage product with each HIV-1 provirus,

we identified a PI-R clone that exhibited similar efficiency at cleaving CARD8 to HIV-1$_{LAI}$ (i.e., PI-

R1), PI-R clones that were markedly less efficient at cleaving CARD8 than HIV-1$_{LAI}$ (i.e., PI-R2,

PI-R3, PI-R5, PI-R9, and PI-R10) and two PI-R clones, PI-R12 and PI-R13, that were more

efficient at cleaving CARD8 than HIV-1$_{LAI}$ (**Figure 5A,** top**, Table S1**). Of note, all PI-R

proviruses had similar levels of HIV[PR] activity for HIV[gagpol] polyprotein processing from p55[gag] to

p24[gag] as indicated by the ratio of p24[gag]/p55[gag] quantified from the anti-p24[gag] immunoblot

(**Figure 5A,** middle). These results indicate that naturally occurring HIV-1 protease mutations

can influence host targets like CARD8.

We next assessed if PI-R clones exhibiting reduced (PI-R2 and -9) or increased (PI-R12

and -13) cleavage of CARD8 relative to HIV-1$_{LAI}$ (**Figure 5A and B**, and **Table S1**) resulted in

differential inflammasome activation. HEK293T cells endogenously express CARD8 but lack the

downstream components (i.e., CASP1, GSDMD, and IL-1β/IL18) of the inflammasome pathway.

Thus, we reconstituted the inflammasome pathway in HEK293T cells by co-transfection of

human caspase 1, human pro-IL-1β, and either empty vector, HIV-1$_{LAI}$ or representative PI-R

proviruses then quantified CASP1-dependent processing of pro-IL-1β as a readout of CARD8

inflammasome activation as in (Tsu et al., 2023). Consistent with the observed differences in

CARD8 cleavage by PI-R clones (**Figure 5A**), we found that PI-R2 and PI-R9, which exhibited

less CARD8 cleavage than HIV-1$_{LAI}$, also induced lower IL-1β levels than HIV-1$_{LAI}$ (**Figure 5B**).

Similarly, PI-R12 and PI-R13, which demonstrated enhanced CARD8 cleavage, elicited higher

IL-1β levels than HIV-1$_{LAI}$ (**Figure 5B**). However, these PI-R clones, relative to the LAI strain,

may have distinct protease substrate specificity, variable efficiency/kinetics in viral assembly,

gag dimerization, and other factors which may also influence CARD8 inflammasome activation.

We next assessed inflammasome activation by the PI-R clones in a cell-to-cell transmission

model using HEK293T cells as donor cells rather than SUPT1 cells and either WT or *CARD8*

KO THP-1 cells as the target line at a 1:1 ratio. We opted to overexpress the HIV-1$_{LAI}$ or the PI-

R proviruses in HEK293T cells rather than infecting SUPT1 cells due to dramatic variability in

replication kinetics between PI-R strains. In these HEK293T:THP-1 cocultures, we observed

that cell-to-cell transmission of PI-R2 and PI-R9 resulted in lower IL-1β levels while PI-R12 and

PI-R13 resulted in higher IL-1β levels compared to HIV-1$_{LAI}$, respectively (**Figure 5C**),

consistent with our findings from CARD8 cleavage (**Figure 5A**) and reconstituted inflammasome

assays (**Figure 5B**). Our findings suggest that HIV-dependent inflammasome activation is under

genetic control of the viral protease in a manner that can be increased or decreased with

naturally occurring mutations induced by drug resistance.

Figure 5. Protease inhibitor resistant strains of HIV-1 differentially cleave and activate CARD8.
(**A**) HEK293T cells were transfected with a construct encoding CARD8 with an N-terminal mCherry tag (mCherry-CARD8) and indicated HIV-1 proviral constructs. Protease inhibitor-resistant (PI-R) clones of HIV-1 are a subset of a panel expressing prototypical multidrug resistant HIV-1 protease (HIV[PR]) in an NL4.3 backbone (Table S1). Top: Immunoblotting using anti-mCherry antibody to detect mCherry-CARD8. The full-length (mC-CARD8) and FIIND-processed bands are indicated as well as the HIV[PR] cut product. The band at ~45 kDa is the result of cleavage by the 20S proteasome (Hsiao et al., 2022). % CARD8 cleavage was calculated by quantifying the HIV[PR] cut band relative to the HIV-1[LAI] control using BioRad Image Lab 6. Middle: Immunoblotting with an anti-p24[gag] antibody showing HIV-1[gag] cleavage products p41[gag] and p24[gag], and/or full-length HIV-1[gag], p55[gag]. %p24/p55 was calculated from the ratio of p24[gag] versus p55[gag] product by quantifying the volume of the p24[gag] bands versus the p55[gag] band relative to the HIV-1[LAI] control using BioRad Image lab 6. Bottom: Immunoblotting with an anti-vinculin antibody to detect vinculin as a loading control. (**B**) HEK293T cells were transfected with human caspase 1 and human pro-IL-1β, and either carrier vector or indicated HIV-1 proviruses then probed for IL-1β secretion 24 hours post-transfection via IL-1R reporter assay. (**C**) HEK293T cells were transfected with indicated HIV-1 proviruses (300ng). 24 hours post-transfection either wildtype (WT) or *CARD8* knockout (KO) THP-1s were overlayed on the transfected HEK293T cells in a 1:1 ratio. THP-1s were primed with Pam3CSK4 overnight prior to coculture. Supernatants were harvested 24 hours post coculture to assay for IL-1β secretion as in (B). Dotted line indicates limit of detection (LoD). Datasets represent mean ± SD (n=4 biological replicates). p-Values were determined by two-way ANOVA with Dunnett's test using GraphPad Prism 10. ns = not significant, *p<0.05,**p<0.01, ***p<0.001, ****p<0.0001.

Table S1: Protease inhibitor resistance mutations and relative CARD8 cleavage.

Clone name	Reported PI-resistance mutations in HIV^{PR}	HIV^{gag} mutation	Strongest PI-R[1]	% CARD8 cleavage (relative to HIV$_{LAI}$)	Additional amino acid changes in HIV^{PR} relative to NL4.3
HIV-1$_{LAI}$	wildtype	wildtype	wildtype	100	37S
CA126802 (PI-R1)	11I, 32I, 33F, 46I, 47V, 54M, 58E, 73S, 84V, 89V, 90M	431V (NC/p1) 437N (NC/p1) 453LF (p1/p6)	FPV, LPV, TPV, DRV	51	10I, 12K, 13V, 20V, 35G, 36I, 37D, 57K, 63P, 64V, 66V, 7IV
CA122805 (PI-R2)	**10F, 33F, 43T, 46L, 54V, 82A, 84V, 90M**	**431V (NC/p1) 532S (p1/p6)**	**SQV**	**7**	16A,19I, 20R, 35D, 36L, 37D, 55R, 57K, 60E, 62V, 63P, 71V, 93L
CA126805 (PI-R3)	33F, 43T, 46I, 48V, 50V, 54S, 82A	437N (NC/p1) 449VF (p1/p6) PTAP insertion (p6)	SQV, IDV, LPV	**8**	not determined*
CA96457 (PI-R5)	48V, 53L, 54V, 82A, 90M	436R (NC/p1)	SQV	11	10I, 37D, 63P, 71T, 77I, 93L
CA96458 (PI-R9)	**10F, 30N, 33F, 43T, 84V, 88D, 90M**	**431V (NC/p1)**	**NFV, SQV**	**24**	15V, 35D, 36L, 37E, 60E, 62V, 63P
CA50834-1 (PI-R10)	24I, 46L, 54V, 76V, 82A	431V (NC/p1)	IDV, LPV	38	10I, 14R, 35D, 36I, 37E, 63P,71V
CA96451 (PI-R12)	**32I, 33F, 43T, 46I, 47V, 54M, 73S, 82A, 89V, 90M**	**437N (NC/p1) PTAP insertion (p6)**	**FPV, LPV**	**192**	10V, 12V, 13V, 15V, 20M, 60E, 61N, 62V, 63P, 67Y, 69K, 71I, 72L, 77I
CA20392-1 (PI-R13)	**24I, 46L, 54V, 82A**	**431V (NC/p1)**	**LPV**	**206**	10I, 14R, 35D, 36I, 37E, 63P, 71V

Table S1 shows the protease inhibitor-resistant (PI-R) clones assayed in **Figure 5** with corresponding mutations in HIV protease (HIV^{PR}) and HIV^{gag}. [1]These clones were previously cloned and assayed for PI-R in (Varghese et al., 2013). The PI-R subset used in Figure 5B are bolded and highlighted in red or green and denote either hypo- or hyper-active CARD8 cleavage, respectively. The last column reports additional amino acid changes in the PI-R clones that were observed via whole plasmid Oxford Nanopore sequencing. *We were unable to sequence verify PI-R3 due to poor plasmid quality. NFV-nelfinavir; FPV-fosamprenavir; SQV- saquinavir; IDV- indinavir; LPV- lopinavir; TPV- tipranavir; DRV- darunavir. The consensus subtype B sequence can be found on the Stanford HIV Drug Resistance Database (HIVDB) ("Stanford - HIV Drug Resistance Database," n.d.). Relative CARD8 cleavage was determined by quantifying band volume of the CARD8 cleavage product in BioRad Image Lab 6 and comparing to cleavage with HIV-1$_{LAI}$.

DISCUSSION

We demonstrate that cell-to-cell transmission of HIV-1 from T cells to myeloid cells in both

immortalized and primary cell coculture models of infection yields CARD8-dependent

inflammasome activation via incoming HIV^{PR}. This inflammasome activation occurs in a largely

NLRP3-independent manner. In addition, we identified protease inhibitor resistant strains of

HIV-1 that differentially cleave and activate the CARD8 inflammasome. Thus, HIV^{PR} mutants

selected for their resistance to different protease inhibitors also affect their ability to cleave host

proteins including the inflammasome-forming sensor CARD8.

CARD8 is the primary inflammasome-forming sensor of HIV-1 infection

Previously, both the NLRP3 and IFI16 inflammasomes have been implicated as innate

sensors of HIV-1 infection and drivers of CD4+ T cell depletion using blood and lymphoid-

derived CD4+ T cells, respectively, and cell-to-cell transmission was reported to be crucial for

IFI16 sensing of abortive HIV transcripts (Doitsh et al., 2014; Galloway et al., 2015; Monroe et

al., 2014; Zhang et al., 2021). However, the mechanism of NLRP3 inflammasome activation in

response to HIV-1 remains elusive. Similarly, there have been reports that IFI16 is not an

inflammasome-forming sensor, and instead a nuclear transcriptional regulator of antiviral genes

including type I interferons and RIG-I (Hornung et al., 2009; Jiang et al., 2021; Thompson et al.,

2014), suggesting that there may be other mechanisms of CD4+ T cell depletion and HIV-

dependent inflammasome activation at play. Indeed, CARD8, which is expressed and functional

in naïve and memory CD4+ and CD8+ T cells (Linder et al., 2020), was recently shown to be

required for pyroptosis in primary human blood- and lymphoid-derived CD4+ T cells and

humanized mouse models (Wang et al., 2024), implicating CARD8 as a major driver of CD4+ T

cell depletion during HIV-1 infection. In this study and our prior work (Kulsuptrakul et al., 2023),

we demonstrate that CARD8 is also the primary innate sensor during HIV-1 infection in myeloid

cell types during cell-to-cell transmission. However, our present study does not rule out the

possibility that under certain conditions or in certain cell types, NLRP3 inflammasome activation

may occur, for example following GSDMD pore formation following CARD8 inflammasome

activation and play a more profound role in promoting HIV-dependent inflammation.

Nonetheless, these data along with other recent work (Wang et al., 2024) strongly suggest that

CARD8 is a major innate sensor of HIV-1 infection.

Protease inhibitor resistance mutations and inflammatory disease

Given the important role of HIV^{PR} in replication, early combination antiretroviral therapy for

people living with HIV (PLWH) included protease inhibitors along with RT inhibitors. However,

resistance mutations to protease inhibitors quickly arose in PLWH through mutations around the

HIV^{PR} active site allowing for polyprotein processing and viral maturation while avoiding drug

inhibition. Despite typically having poor overall viral fitness due to less efficient polyprotein

processing and replication relative to wildtype HIV-1 in the absence of protease inhibitors, these

mutant drug-resistant HIV-1 strains can persist in PLWH on antiviral therapy, posing a major

threat to controlling disease progression (De Luca, 2006; Martinez-Picado et al., 1999; Prado et

al., 2002; Resch et al., 2002). To compensate for mutations in HIV^{PR} that change its substrate

specificity, HIV^{gag} sometimes evolves mutations around HIV^{PR} cleavage sites to permit proper

polyprotein processing (Varghese et al., 2013). Here, we identified multiple HIV^{PR} inhibitor-

resistant strains of HIV-1 that can differentially cleave and activate the CARD8 inflammasome

(**Figure 5, Table S1**). As the degree of inflammation is a better predictor of disease progression

in untreated individuals than viral load (Deeks et al., 2004; Giorgi et al., 1999), we speculate that

differential CARD8 inflammasome activation could influence disease progression for PLWH

harboring HIV^{PR} resistance mutations that cleave CARD8 more or less efficiently. More broadly,

we suggest that host targets of viral proteases like CARD8 may influence the selection of viral

variants during treatment with antiviral protease inhibitor monotherapies.

Viral protease influx activates the CARD8 inflammasome

In this study, we found that HIV-dependent CARD8 inflammasome activation during cell-

free infection requires a cationic polymer like DEAE-dextran to facilitate efficient viral infection

(see **Supplemental Note**). Despite being infected with the same amount of virus and exhibiting

similar percent infection 24 hours post-infection, as measured by intracellular p24^{gag}, with and

without DEAE-dextran, we hypothesize that DEAE-dextran during cell-free infection may

increase the total viral dose that enters cells consisting of both infectious particles and non-

infectious particles that may nonetheless contain active protease, leading to more efficient viral

protease influx to trigger CARD8 sensing. Hence, the percentage of p24^{gag} positive cells after 24

hours may be an underestimate of total amount of viral entry in the DEAE-dextran condition. We

speculate that a considerable influx of incoming HIV^{PR} may be necessary to induce CARD8

inflammasome activation.

We also observed that DEAE-dextran can trigger inflammasome activation in some THP-1

cell lines (see **Supplemental Note**), prompting us to assay for inflammasome activation upon

HIV-1 cell-to-cell transmission with target cells at a 1:1 ratio in the absence of cationic polymer.

We found that inflammasome activation following cell-to-cell transmission of HIV-1 could be

detected by 24 hours (**Figure 3B**), which is delayed relative to our detection of CARD8

inflammasome activation 2 hours post cell-free HIV-1 infection in the presence of DEAE-dextran

(Kulsuptrakul et al., 2023). Nonetheless, we still hypothesize that this inflammasome activation

is driven by active incoming viral protease because treatment with an RT inhibitor has no effect

on inflammasome activation, which implies that *de novo* protease production is not necessary

(**Figure 3A and B, Figure 4A and B**). Similarly, we observed that IL-1β levels do not increase

after plateauing 24 hours after establishing the coculture (**Figure 3B, Figure 4B**), suggesting

that secondary infection does not further amplify inflammasome activation. We infer that this is

also likely a product of the efficiency of viral entry and the necessity for multiple virions infecting

at the same time to deliver a sufficient amount of active HIV^{PR} for cytosolic CARD8 sensing. We

postulate that under certain physiological conditions, cell-to-cell transmission can cause CARD8

inflammasome activation when there is an influx of active incoming HIV^{PR} across the viral

synapse. Taken together, we speculate that both cell-free infection facilitated by cationic

polymer or cell-to-cell transmission can achieve sufficient levels of active HIV^{PR} influx to activate

the CARD8 inflammasome.

Macrophages have been reported to be primarily infected through phagocytosis of

infected CD4+ T cells or cell-to-cell transmission (Dupont and Sattentau, 2020; Martínez-

Méndez et al., 2017; Orenstein, 2000). We demonstrate that unlike CD4+ T cells, which are

rapidly depleted by HIV-1 infection and do not release IL-1β or IL-18 (Linder et al., 2020),

primary macrophages release pro-inflammatory cytokines in response to HIV^{PR} during cell-to-

cell infection from primary CD4+ T cells and T cell lines (**Figure 3, Figure 4**), thus representing

a potential source of sustained IL-1β and subsequent chronic immune activation. In addition to

promoting chronic immune activation, HIV-dependent IL-1β release from macrophages may

also contribute to HIV-1 pathogenesis by activating nearby CD4+ T cells, rendering them

susceptible to becoming infected with HIV-1, and thus indirectly promoting CD4+ T cell

depletion. Collectively with our prior work (Kulsuptrakul et al., 2023), our findings provide further

evidence that CARD8 inflammasome activation is driven by incoming HIV^{PR} under conditions

where multiple virions may enter cells, and thus could be a potential driver of HIV-1

pathogenesis by promoting chronic immune activation.

METHODS

Plasmids and Reagents

pMD2.G used for HIV-1$_{LAI-VSVG}$ production was a gift from Didier Trono (Addgene). HIV-1$_{LAI}$ has

been previously described (Peden et al., 1991). The following reagents were obtained through

the NIH HIV Reagent Program, Division of AIDS, NIAID, NIH: lopinavir (LPV), nevirapine (NVP),

Human Immunodeficiency Virus 1 (HIV-1) NL4-3 BaL Infectious Molecular Clone (p20-36) (HIV-

$1_{NL4.3-BaL}$), ARP-11442, contributed by Dr. Bruce Chesebro (Chesebro et al., 1992, 1991; Toohey

et al., 1995; Walter et al., 2005), and Panel of Multi-Protease Inhibitor Resistant Infectious

Molecular Clones, HRP-12740, contributed by Dr. Robert Shafer (Varghese et al., 2013). Mutant

HIV^{PR} sequences were amplified from clinically-derived viral cDNA encoding protease genes

with resistance to multiple PRis then cloned into an NL4.3 backbone with overhangs including

the 3' end of gag with the gag cleavage site and the 5' end of RT as previously described

(Varghese et al., 2013). CARD8 variant constructs were cloned as previously described

(Kulsuptrakul et al., 2023). VX765 and MCC950 were sourced from Invivogen (cat: inh-vx765i-1

and inh-mcc, respectively).

Cell culture

SUPT1 (ATCC) and THP-1 cells (JK and ATCC) were cultured in RPMI (Invitrogen) with 10%

FBS, 1% penicillin/streptomycin antibiotics, 10 mM HEPES, 0.11 g/L sodium pyruvate, 4.5 g/L

D-glucose and 1% Glutamax. JK THP-1 cells were used for all experiments in this manuscript

and our previous work unless explicitly stated (see **Supplemental Note**) (Kulsuptrakul et al.,

2023). Primary monocytes were cultured in RPMI (Invitrogen) with 10% FBS, and 1%

penicillin/streptomycin antibiotics and differentiated in the presence of 20ng/mL GM-CSF

(Peprotech cat: 300-03) and 20ng/mL M-CSF (Peprotech cat: 300-25). Primary CD4+ T cells

were cultured in RPMI (Invitrogen) with 10% FBS, 1% penicillin/streptomycin antibitiocs and

100U/mL IL-2. HEK293T (ATCC) lines were cultured in DMEM (Invitrogen) with 10% FBS and

1% penicillin/streptomycin antibiotics. All lines routinely tested negative for mycoplasma bacteria

(Fred Hutch Specimen Processing & Research Cell Bank). SUPT1 and THP-1 cell lines were

authenticated by STR profiling analysis (Fred Hutch Genomics core and TransnetYX, Inc.) (see

Supplemental Note).

HIV-1$_{LAI}$, HIV-1$_{LAI-VSVG}$, and HIV-1$_{NL4.3-BaL}$ production

293T cells were seeded at 2-3×10^5 cells/well in six-well plates the day before transfection using

TransIT-LT1 reagent (Mirus Bio LLC) at 3 µL transfection reagent/well as previously described

(OhAinle et al., 2018). For HIV-1 production, 293Ts were transfected with 1 µg/well HIV$_{LAI}$ or

HIV-1$_{NL4.3-BaL}$ proviral DNA or 1 µg/well HIV$_{LAI}$ Δenv DNA and 500 ng/well pMD2.G for HIV-1$_{LAI}$,

HIV-1$_{NL4.3-BaL}$, and HIV-1$_{LAI-VSVG}$, respectively. One day post-transfection, media was replaced.

Two days post-transfection, viral supernatants were collected and filtered through a 20 µm filter

and aliquots were frozen at –80°C. HIV-1$_{LAI}$, HIV-1$_{NL4.3-BaL}$ and HIV-1$_{LAI-VSVG}$ proviruses were

previously described (Bartz and Vodicka, 1997; Gummuluru et al., 2003; Peden et al., 1991).

Cell-free and cell-to-cell coculture HIV-1 infection

Cell-free infections with HIV-1$_{LAI-VSVG}$ were done as previously described (Kulsuptrakul et al.,

2023). Subsequent cell death was assessed by incubating in media containing propidium iodide

dye (10µg/mL) for 5 minutes at room temperature then washed once with PBS before fixing with

BD CytoFix/Cytoperm (cat:BDB554714) and staining for intracellular p24^{gag} (Beckman Coulter

cat#: 6604665) for flow cytometry. In the HIV-1 cell-to-cell transmission system, SUPT1

expressing CCR5 (SUPT1-CCR5) cells were spinoculated at 1100g for 30min with either HIV-

1$_{LAI}$ or HIV-1$_{NL4.3-BaL}$ in the presence of 10µg/mL DEAE-dextran. SUPT1-CCR5 cells were

lentiviral transduced to express CCR5 (Dingens et al., 2017). After 24 hours, mock or HIV-1

infected SUPT1-CCR5 cells were washed three times in PBS such that DEAE-dextran and cell-

free virus were removed before starting coculture with THP-1 cells or MDMs. THP-1 cells and

MDMs were seeded at 5 x 10^5 cells/well and primed with 500ng/mL Pam3CSK4 (Invivogen) for

16-24 hours before coculture. Mock or infected SUPT1 cells were seeded at 5 x 10^5 cells/well.

Cultured supernatants from coculture were harvested 48 hours after starting the coculture for

the IL-1R reporter assay, which was previously described (Kulsuptrakul et al., 2023).

Transwell coculture HIV-1 infection

SUPT1 cells were spinoculated at 1100g for 30min with HIV-1$_{LAI}$ in the presence of 10µg/mL

DEAE-dextran. After 24 hours, mock or HIV-1 infected SUPT1 cells were washed 3 times in

PBS and either mixed in a 24-well with THP-1 cells or placed in a transwell insert above target

THP-1 cells at a concentration of 5 x 10^5 infected SUPT1 cells and 2.5 x 10^5 THP-1 cells per

well. THP-1 cells were primed overnight with 500ng/mL Pam3CSK4 before starting coculture.

The transwell insert has a 0.4µm membrane at the bottom of the well (ThinCert™ Tissue

Culture Inserts, Sterile, Greiner Bio-One cat:665640), allowing virus to pass out of the transwell

but not the infected cell. Reverse transcriptase (RT) activity in viral supernatants was measured

using the RT activity assay as previously described (Roesch et al., 2018; Vermeire et al., 2012).

A stock of HIV-1$_{LAI}$ virus was titered multiple times, aliquoted at −80°C and used as the standard

curve in all assays.

Monocyte-derived macrophage isolation, differentiation, and editing

Primary monocytes were isolated via negative selection using the EasySep™ Human Monocyte

Isolation Kit (Easy Sep, 1x10^9) (Stem Cell Technologies) according to the manufacturer's

protocols from PBMCs collected from blood donors. Upon isolation, monocytes were seeded at

1 x 10^6 cell/mL and differentiated for 5 days in the presence of media containing 20ng/mL GM-

CSF (Peprotech cat: 300-03) and 20ng/mL M-CSF (Peprotech cat: 300-25), changing media

every other day. For edited MDMs, isolated monocytes were electroporated in cuvettes (100µL)

with 2.5-5 x 10^6 cells/nucleofection in the presence of pre-complexed Cas9-RNPs (300pmol

sgRNA: 100pmol Cas9) in Lonza P2 buffer using pulse code DK-100. RNPs were complexed

with sgRNA from the Synthego gene KO kit, which includes 3 sgRNAs per gene. Thus, each

sgRNA was present at a 1:1 ratio with Cas9 (QB3 MacroLab or Synthego SpCas9 2NLS

Nuclease). A table of sgRNAs used for *AAVS1, CARD8,* or *NLRP3* KO can be found in **Table**

S2 below. After nucleofection, cells were supplemented with 900µL of prewarmed media and

allowed to recover for 30 minutes at 37°C before counting and seeding at ~1-1.5x10^6 cells/mL

for differentiation. Media was changed 24 hours post nucleofection then differentiated for 5 more

days before characterizing knockout efficiency and conducting coculture experiments.

Table S2:

sgRNA	Sequence
CARD8 sgRNA1	CUCUGCAGUGACAUCAAACA
CARD8 sgRNA2	UGACGAUUGCGUUUGGUUCC
CARD8 sgRNA3	AGCGUUUGGUUCCCCACUGC
AAVS1 sgRNA 1	GUUAAUGUGGCUCUGGUUCU
AAVS1 sgRNA 2	ACCCCACAGUGGGGCCACUA
AAVS1 sgRNA 3	CCUUCCUAGUCUCCUGAUAU
NLRP3 sgRNA 1	GCUCAGAAUGCUCAUCAUCG
NLRP3 sgRNA 2	GAUGAUGUUGGACUGGGCAU
NLRP3 sgRNA 3	CAAGGCUCACCUCCCGACAG

CD4+ T cell isolation, infection, and coculture

Primary CD4+ T cells were isolated via positive selection using the EasySep^{TM} Release Human

CD4 Positive selection kit (Stem Cell Technologies Cat: 17752) according to the manufacturer's

instructions from PBMCs collected from blood donors and seeded at 2.5x10^6 cells/mL in the

presence of 100U/mL IL-2. T cells were activated 24 hours post-isolation with CD3/CD28

activation beads (Miltenyi Biotech Cat: 130-091-441). Activation beads were removed according

to the manufacturer's protocols 24 hours later for infection. For infection, T cells were

suspended at 1-1.5x10^6 cells/mL in 15mL conical tubes containing 8µg/mL polybrene, 100U/mL

IL-2, and HIV-1$_{NL4.3-BaL}$ then spinoculated at 1100g at 30°C for 90 min. Three days post-infection,

CD4s were assessed for intracellular p24^{gag} via flow cytometry (~10% infected) then washed

thrice with PBS before coculturing with MDMs. CD4:MDM coculture RPMI media was

supplemented with 100U/mL IL-2, 20ng/mL GM-CSF, 20ng/mL M-CSF, and 500ng/mL

Pam3CSK4.

CARD8 cleavage assay

HEK293T cells were seeded at $1–1.5 \times 10^5$ cells/well in 24-well plates the day before

transfection using TransIT-LT1 reagent at 1.5 μL transfection reagent/well (Mirus Bio LLC). One

hundred ng of indicated constructs encoding an N-terminal mCherry-tagged CARD8 were co-

transfected into HEK293T cells with empty vector ('–'), HIV_{LAI} or PI-R provirus. To normalize

HIV^{gag} expression between HIV-1$_{LAI}$ and the PI-R clones, which are in a different vector

backbone, 400ng of HIV-1$_{LAI}$ and 200ng of all PI-R clones were transfected. All conditions were

normalized with empty vector to contain the same amount of DNA. Cytoplasmic lysates were

harvested 24 hours post-transfection and immunoblotted as previously described (Kulsuptrakul

et al., 2023).

HEK reconstitution assay

HEK293T cells, which endogenously express CARD8, were seeded at 2.25×10^5 cell/well in 24-

well plates the day before transfection using TransIT-LT1 reagent at 1.5uL transfection

reagent/well (Mirus Bio LLC). Functional inflammasomes were reconstituted by transfecting in

5ng human CASP1 and 100ng human IL-1β. To assess the effects of different viral proteases

on inflammasome activation, HIV-1$_{LAI}$ or PI-R clones were co-transfected in with CASP1 and IL-

1β. As with the CARD8 cleavage assay, a higher amount of 250ng HIV-1$_{LAI}$ was added relative

to the PI-R clones, which were all added at 125ng, to normalize HIV^{gag} expression between the

different vector backbones. All conditions were normalized with empty vector to contain the

same amount of DNA. Cultured supernatant was harvested 24 hours post-transfection to assay

for IL-1β secretion via IL-1R reporter assay.

Acknowledgements

We thank everyone in the Emerman and Mitchell labs for helpful feedback on the project, Terry

Hafer and Marisa Yonemitsu for critical reading of the manuscript, Liang Shan and his lab

members for discussions and sharing of unpublished results and protocols, and the Fred

Hutchinson Shared Resources Genomics, Flow Cytometry, and Specimen Processing &

Research Cell Bank cores. LPV (HRP-9481), NVP (HRP-4666) and p24^{gag} antibody (ARP-3537)

were provided by the AIDS Reagent Program, Division of AIDS, NIAID, NIH. PSM is an HHMI

Freeman Hrabowski Scholar and is supported by grants from the National Institutes of Health

(NIH) (DP2 AI 154432-01) and the Mallinckrodt Foundation to PSM. ME is supported by NIH

grant DP1 DA051110. JK is supported by the University of Washington Cellular and Molecular

Biology Training Grant (T32 GM007270).

References

Abela IA, Berlinger L, Schanz M, Reynell L, Günthard HF, Rusert P, Trkola A. 2012. Cell-Cell
Transmission Enables HIV-1 to Evade Inhibition by Potent CD4bs Directed Antibodies.
PLOS Pathog **8**:e1002634. doi:10.1371/journal.ppat.1002634
Agosto LM, Herring MB, Mothes W, Henderson AJ. 2018. HIV-1-Infected CD4+ T Cells
Facilitate Latent Infection of Resting CD4+ T Cells through Cell-Cell Contact. *Cell Rep*
24:2088–2100. doi:10.1016/j.celrep.2018.07.079
Agosto LM, Uchil PD, Mothes W. 2015. HIV cell-to-cell transmission: effects on pathogenesis
and antiretroviral therapy. *Trends Microbiol* **23**:289–295. doi:10.1016/j.tim.2015.02.003
Bailey CA, Miller DK, Lenard J. 1984. Effects of DEAE-dextran on infection and hemolysis by
VSV. Evidence that nonspecific electrostatic interactions mediate effective binding of
VSV to cells. *Virology* **133**:111–118. doi:10.1016/0042-6822(84)90429-x
Bandera A, Masetti M, Fabbiani M, Biasin M, Muscatello A, Squillace N, Clerici M, Gori A,
Trabattoni D. 2018. The NLRP3 Inflammasome Is Upregulated in HIV-Infected
Antiretroviral Therapy-Treated Individuals with Defective Immune Recovery. *Front*
Immunol **9**.
Bartz SR, Vodicka MA. 1997. Production of High-Titer Human Immunodeficiency Virus Type 1
Pseudotyped with Vesicular Stomatitis Virus Glycoprotein. *Methods* **12**:337–342.
doi:10.1006/meth.1997.0487
Baxter AE, Russell RA, Duncan CJA, Moore MD, Willberg CB, Pablos JL, Finzi A, Kaufmann
DE, Ochsenbauer C, Kappes JC, Groot F, Sattentau QJ. 2014. Macrophage infection via
selective capture of HIV-1-infected CD4+ T cells. *Cell Host Microbe* **16**:711–721.
doi:10.1016/j.chom.2014.10.010
Brenchley JM, Paiardini M, Knox KS, Asher AI, Cervasi B, Asher TE, Scheinberg P, Price DA,
Hage CA, Kholi LM, Khoruts A, Frank I, Else J, Schacker T, Silvestri G, Douek DC.
2008. Differential Th17 CD4 T-cell depletion in pathogenic and nonpathogenic lentiviral
infections. *Blood* **112**:2826–2835. doi:10.1182/blood-2008-05-159301
Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, Kazzaz Z, Bornstein E,
Lambotte O, Altmann D, Blazar BR, Rodriguez B, Teixeira-Johnson L, Landay A, Martin
JN, Hecht FM, Picker LJ, Lederman MM, Deeks SG, Douek DC. 2006. Microbial
translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med*
12:1365–1371. doi:10.1038/nm1511

Broz P, Dixit VM. 2016. Inflammasomes: mechanism of assembly, regulation and signalling. *Nat*
Rev Immunol **16**:407–420. doi:10.1038/nri.2016.58
Castro LK, Daugherty MD. 2023. Tripping the wire: sensing of viral protease activity by CARD8
and NLRP1 inflammasomes. *Curr Opin Immunol* **83**:102354.
doi:10.1016/j.coi.2023.102354
Chen P, Hübner W, Spinelli MA, Chen BK. 2007. Predominant mode of human
immunodeficiency virus transfer between T cells is mediated by sustained Env-
dependent neutralization-resistant virological synapses. *J Virol* **81**:12582–12595.
doi:10.1128/JVI.00381-07
Chesebro B, Nishio J, Perryman S, Cann A, O'Brien W, Chen IS, Wehrly K. 1991. Identification
of human immunodeficiency virus envelope gene sequences influencing viral entry into
CD4-positive HeLa cells, T-leukemia cells, and macrophages. *J Virol* **65**:5782–5789.
doi:10.1128/JVI.65.11.5782-5789.1991
Chesebro B, Wehrly K, Nishio J, Perryman S. 1992. Macrophage-tropic human
immunodeficiency virus isolates from different patients exhibit unusual V3 envelope
sequence homogeneity in comparison with T-cell-tropic isolates: definition of critical
amino acids involved in cell tropism. *J Virol* **66**:6547–6554. doi:10.1128/JVI.66.11.6547-
6554.1992
Chivero ET, Guo M-L, Periyasamy P, Liao K, Callen SE, Buch S. 2017. HIV-1 Tat Primes and
Activates Microglial NLRP3 Inflammasome-Mediated Neuroinflammation. *J Neurosci*
37:3599–3609. doi:10.1523/JNEUROSCI.3045-16.2017
Chun T-W, Carruth L, Finzi D, Shen X, DiGiuseppe JA, Taylor H, Hermankova M, Chadwick K,
Margolick J, Quinn TC, Kuo Y-H, Brookmeyer R, Zeiger MA, Barditch-Crovo P, Siliciano
RF. 1997a. Quantification of latent tissue reservoirs and total body viral load in HIV-1
infection. *Nature* **387**:183–188. doi:10.1038/387183a0
Chun T-W, Finzi D, Margolick J, Chadwick K, Schwartz D, Siliciano RF. 1995. In vivo fate of
HIV-1-infected T cells: Quantitative analysis of the transition to stable latency. *Nat Med*
1:1284–1290. doi:10.1038/nm1295-1284
Chun T-W, Stuyver L, Mizell SB, Ehler LA, Mican JAM, Baseler M, Lloyd AL, Nowak MA, Fauci
AS. 1997b. Presence of an inducible HIV-1 latent reservoir during highly active
antiretroviral therapy. *Proc Natl Acad Sci* **94**:13193–13197.
doi:10.1073/pnas.94.24.13193
Clark KM, Kim JG, Wang Q, Gao H, Presti RM, Shan L. 2022. Chemical inhibition of DPP9
sensitizes the CARD8 inflammasome in HIV-1-infected cells. *Nat Chem Biol* 1–9.
doi:10.1038/s41589-022-01182-5
Coll RC, Robertson AAB, Chae JJ, Higgins SC, Muñoz-Planillo R, Inserra MC, Vetter I, Dungan
LS, Monks BG, Stutz A, Croker DE, Butler MS, Haneklaus M, Sutton CE, Núñez G, Latz
E, Kastner DL, Mills KHG, Masters SL, Schroder K, Cooper MA, O'Neill LAJ. 2015. A
small-molecule inhibitor of the NLRP3 inflammasome for the treatment of inflammatory
diseases. *Nat Med* **21**:248–255. doi:10.1038/nm.3806
Collman R, Godfrey B, Cutilli J, Rhodes A, Hassan NF, Sweet R, Douglas SD, Friedman H,
Nathanson N, Gonzalez-Scarano F. 1990. Macrophage-tropic strains of human
immunodeficiency virus type 1 utilize the CD4 receptor. *J Virol* **64**:4468–4476.
doi:10.1128/jvi.64.9.4468-4476.1990
Collman R, Hassan NF, Walker R, Godfrey B, Cutilli J, Hastings JC, Friedman H, Douglas SD,
Nathanson N. 1989. Infection of monocyte-derived macrophages with human
immunodeficiency virus type 1 (HIV-1). Monocyte-tropic and lymphocyte-tropic strains of
HIV-1 show distinctive patterns of replication in a panel of cell types. *J Exp Med*
170:1149–1163. doi:10.1084/jem.170.4.1149

Conant D, Hsiau T, Rossi N, Oki J, Maures T, Waite K, Yang J, Joshi S, Kelso R, Holden K,
Enzmann BL, Stoner R. 2022. Inference of CRISPR Edits from Sanger Trace Data.
CRISPR J **5**:123–130. doi:10.1089/crispr.2021.0113
De Luca A. 2006. The impact of resistance on viral fitness and its clinical implications In: Geretti
AM, editor. Antiretroviral Resistance in Clinical Practice. London: Mediscript.
Deeks SG, Kitchen CMR, Liu L, Guo H, Gascon R, Narváez AB, Hunt P, Martin JN, Kahn JO,
Levy J, McGrath MS, Hecht FM. 2004. Immune activation set point during early HIV
infection predicts subsequent CD4+ T-cell changes independent of viral load. *Blood*
104:942–947. doi:10.1182/blood-2003-09-3333
Del Portillo A, Tripodi J, Najfeld V, Wodarz D, Levy DN, Chen BK. 2011. Multiploid Inheritance
of HIV-1 during Cell-to-Cell Infection ▽. *J Virol* **85**:7169–7176. doi:10.1128/JVI.00231-11
Dingens AS, Haddox HK, Overbaugh J, Bloom JD. 2017. Comprehensive mapping of HIV-1
escape from a broadly neutralizing antibody. *Cell Host Microbe* **21**:777-787.e4.
doi:10.1016/j.chom.2017.05.003
Doitsh G, Galloway NL, Geng X, Yang Z, Monroe KM, Zepeda O, Hunt PW, Hatano H, Sowinski
S, Muñoz-Arias I, Greene WC. 2014. Pyroptosis drives CD4 T-cell depletion in HIV-1
infection. *Nature* **505**:509–514. doi:10.1038/nature12940
Dufloo J, Bruel T, Schwartz O. 2018. HIV-1 cell-to-cell transmission and broadly neutralizing
antibodies. *Retrovirology* **15**:51. doi:10.1186/s12977-018-0434-1
Duncan CJA, Russell RA, Sattentau QJ. 2013. High multiplicity HIV-1 cell-to-cell transmission
from macrophages to CD4+ T cells limits antiretroviral efficacy. *AIDS* **27**:2201.
doi:10.1097/QAD.0b013e3283632ec4
Dupont M, Sattentau QJ. 2020. Macrophage Cell-Cell Interactions Promoting HIV-1 Infection.
Viruses **12**:492. doi:10.3390/v12050492
Fink SL, Cookson BT. 2005. Apoptosis, Pyroptosis, and Necrosis: Mechanistic Description of
Dead and Dying Eukaryotic Cells. *Infect Immun* **73**:1907–1916.
doi:10.1128/IAI.73.4.1907-1916.2005
Galloway NLK, Doitsh G, Monroe KM, Yang Z, Muñoz-Arias I, Levy DN, Greene WC. 2015.
Cell-to-Cell Transmission of HIV-1 Is Required to Trigger Pyroptotic Death of Lymphoid-
Tissue-Derived CD4 T Cells. *Cell Rep* **12**:1555–1563. doi:10.1016/j.celrep.2015.08.011
Giorgi JV, Hultin LE, McKeating JA, Johnson TD, Owens B, Jacobson LP, Shih R, Lewis J,
Wiley DJ, Phair JP, Wolinsky SM, Detels R. 1999. Shorter Survival in Advanced Human
Immunodeficiency Virus Type 1 Infection Is More Closely Associated with T Lymphocyte
Activation than with Plasma Virus Burden or Virus Chemokine Coreceptor Usage. *J*
Infect Dis **179**:859–870. doi:10.1086/314660
Gosselin A, Monteiro P, Chomont N, Diaz-Griffero F, Said EA, Fonseca S, Wacleche V, El-Far
M, Boulassel M-R, Routy J-P, Sekaly R-P, Ancuta P. 2009. Peripheral Blood
CCR4+CCR6+ and CXCR3+CCR6+ CD4+ T Cells Are Highly Permissive to HIV-1
Infection. *J Immunol* **184**:1604–1616. doi:10.4049/jimmunol.0903058
Gummuluru S, Rogel M, Stamatatos L, Emerman M. 2003. Binding of Human Immunodeficiency
Virus Type 1 to Immature Dendritic Cells Can Occur Independently of DC-SIGN and
Mannose Binding C-Type Lectin Receptors via a Cholesterol-Dependent Pathway. *J*
Virol **77**:12865–12874. doi:10.1128/JVI.77.23.12865-12874.2003
Hernandez JC, Latz E, Urcuqui-Inchima S. 2013. HIV-1 Induces the First Signal to Activate the
NLRP3 Inflammasome in Monocyte-Derived Macrophages. *Intervirology* **57**:36–42.
doi:10.1159/000353902
Hornung V, Ablasser A, Charrel-Dennis M, Bauernfeind F, Horvath G, Caffrey DR, Latz E,
Fitzgerald KA. 2009. AIM2 recognizes cytosolic dsDNA and forms a caspase-1
activating inflammasome with ASC. *Nature* **458**:514–518. doi:10.1038/nature07725
Hsiao JC, Neugroschl AR, Chui AJ, Taabazuing CY, Griswold AR, Wang Q, Huang H-C, Orth-
He EL, Ball DP, Hiotis G, Bachovchin DA. 2022. A ubiquitin-independent proteasome

pathway controls activation of the CARD8 inflammasome. *J Biol Chem* **298**:102032.
doi:10.1016/j.jbc.2022.102032
Iwami S, Takeuchi JS, Nakaoka S, Mammano F, Clavel F, Inaba H, Kobayashi T, Misawa N,
Aihara K, Koyanagi Y, Sato K. 2015. Cell-to-cell infection by HIV contributes over half of
virus infection. *eLife* **4**:e08150. doi:10.7554/eLife.08150
Jiang Z, Wei F, Zhang Y, Wang T, Gao W, Yu S, Sun H, Pu J, Sun Y, Wang M, Tong Q, Gao C,
Chang K-C, Liu J. 2021. IFI16 directly senses viral RNA and enhances RIG-I
transcription and activation to restrict influenza virus infection. *Nat Microbiol* **6**:932–945.
doi:10.1038/s41564-021-00907-x
Jolly C, Booth NJ, Neil SJD. 2010. Cell-Cell Spread of Human Immunodeficiency Virus Type 1
Overcomes Tetherin/BST-2-Mediated Restriction in T cells. *J Virol* **84**:12185–12199.
doi:10.1128/jvi.01447-10
Kulsuptrakul J, Turcotte EA, Emerman M, Mitchell PS. 2023. A human-specific motif facilitates
CARD8 inflammasome activation after HIV-1 infection. *eLife* **12**:e84108.
doi:10.7554/eLife.84108
Leal VNC, Reis EC, Fernandes FP, Pontillo A. 2020. NLRP3 inflammasome activation profile in
CD4+ T and CD19+ B lymphocytes from HIV-infected patients. *J Immunol* **204**:225.8.
doi:10.4049/jimmunol.204.Supp.225.8
Linder A, Bauernfried S, Cheng Y, Albanese M, Jung C, Keppler OT, Hornung V. 2020. CARD8
inflammasome activation triggers pyroptosis in human T cells. *EMBO J* **39**:e105071.
doi:10.15252/embj.2020105071
Lopez P, Koh WH, Hnatiuk R, Murooka TT. 2019. HIV Infection Stabilizes Macrophage-T Cell
Interactions To Promote Cell-Cell HIV Spread. *J Virol* **93**:e00805-19.
doi:10.1128/JVI.00805-19
Mamik MK, Hui E, Branton WG, McKenzie BA, Chisholm J, Cohen EA, Power C. 2017. HIV-1
Viral Protein R Activates NLRP3 Inflammasome in Microglia: implications for HIV-1
Associated Neuroinflammation. *J Neuroimmune Pharmacol* **12**:233–248.
doi:10.1007/s11481-016-9708-3
Martin N, Welsch S, Jolly C, Briggs JAG, Vaux D, Sattentau QJ. 2010. Virological Synapse-
Mediated Spread of Human Immunodeficiency Virus Type 1 between T Cells Is
Sensitive to Entry Inhibition. *J Virol* **84**:3516–3527. doi:10.1128/jvi.02651-09
Martínez-Méndez D, Rivera-Toledo E, Ortega E, Licona-Limón I, Huerta L. 2017. Monocyte-
lymphocyte fusion induced by the HIV-1 envelope generates functional heterokaryons
with an activated monocyte-like phenotype. *Exp Cell Res* **352**:9–19.
doi:10.1016/j.yexcr.2017.01.014
Martinez-Picado J, Savara AV, Sutton L, D'Aquila RT. 1999. Replicative fitness of protease
inhibitor-resistant mutants of human immunodeficiency virus type 1. *J Virol* **73**:3744–
3752. doi:10.1128/JVI.73.5.3744-3752.1999
Monroe KM, Yang Z, Johnson JR, Geng X, Doitsh G, Krogan NJ, Greene WC. 2014. IFI16 DNA
Sensor Is Required for Death of Lymphoid CD4 T-cells Abortively Infected with HIV.
Science **343**:428–432. doi:10.1126/science.1243640
Nadkarni R, Chu WC, Lee CQE, Mohamud Y, Yap L, Toh GA, Beh S, Lim R, Fan YM, Zhang
YL, Robinson K, Tryggvason K, Luo H, Zhong F, Ho L. 2022. Viral proteases activate the
CARD8 inflammasome in the human cardiovascular system. *J Exp Med* **219**:e20212117.
doi:10.1084/jem.20212117
OhAinle M, Helms L, Vermeire J, Roesch F, Humes D, Basom R, Delrow JJ, Overbaugh J,
Emerman M. 2018. A virus-packageable CRISPR screen identifies host factors
mediating interferon inhibition of HIV. *eLife* **7**:e39823. doi:10.7554/eLife.39823
Orenstein JM. 2000. In Vivo Cytolysis and Fusion of Human Immunodeficiency Virus Type 1-
Infected Lymphocytes in Lymphoid Tissue. *J Infect Dis* **182**:338–342.
doi:10.1086/315640

Peden K, Emerman M, Montagnier L. 1991. Changes in growth properties on passage in tissue
culture of viruses derived from infectious molecular clones of HIV-1LAI, HIV-1MAL, and
HIV-1ELI. *Virology* **185**:661–672. doi:10.1016/0042-6822(91)90537-L
Prado JG, Wrin T, Beauchaine J, Ruiz L, Petropoulos CJ, Frost SDW, Clotet B, D'Aquila RT,
Martinez-Picado J. 2002. Amprenavir-resistant HIV-1 exhibits lopinavir cross-resistance
and reduced replication capacity. *AIDS Lond Engl* **16**:1009–1017.
doi:10.1097/00002030-200205030-00007
Primiano MJ, Lefker BA, Bowman MR, Bree AG, Hubeau C, Bonin PD, Mangan M, Dower K,
Monks BG, Cushing L, Wang S, Guzova J, Jiao A, Lin L-L, Latz E, Hepworth D, Hall JP.
2016. Efficacy and Pharmacology of the NLRP3 Inflammasome Inhibitor CP-456,773
(CRID3) in Murine Models of Dermal and Pulmonary Inflammation. *J Immunol*
197:2421–2433. doi:10.4049/jimmunol.1600035
Resch W, Ziermann R, Parkin N, Gamarnik A, Swanstrom R. 2002. Nelfinavir-resistant,
amprenavir-hypersusceptible strains of human immunodeficiency virus type 1 carrying
an N88S mutation in protease have reduced infectivity, reduced replication capacity, and
reduced fitness and process the Gag polyprotein precursor aberrantly. *J Virol* **76**:8659–
8666. doi:10.1128/jvi.76.17.8659-8666.2002
Richardson MW, Carroll RG, Stremlau M, Korokhov N, Humeau LM, Silvestri G, Sodroski J,
Riley JL. 2008. Mode of Transmission Affects the Sensitivity of Human
Immunodeficiency Virus Type 1 to Restriction by Rhesus TRIM5α. *J Virol* **82**:11117–
11128. doi:10.1128/jvi.01046-08
Rodriguez-Garcia M, Barr FD, Crist SG, Fahey JV, Wira CR. 2014. Phenotype and susceptibility
to HIV infection of CD4+ Th17 cells in the human female reproductive tract. *Mucosal*
Immunol **7**:1375–1385. doi:10.1038/mi.2014.26
Roesch F, OhAinle M, Emerman M. 2018. A CRISPR screen for factors regulating SAMHD1
degradation identifies IFITMs as potent inhibitors of lentiviral particle delivery.
Retrovirology **15**:26. doi:10.1186/s12977-018-0409-2
Russell RA, Martin N, Mitar I, Jones E, Sattentau QJ. 2013. Multiple proviral integration events
after virological synapse-mediated HIV-1 spread. *Virology* **443**:143–149.
doi:10.1016/j.virol.2013.05.005
Stanford - HIV Drug Resistance Database. n.d. https://hivdb.stanford.edu/page/release-
notes/#consensusbsequences
Thompson MR, Sharma S, Atianand M, Jensen SB, Carpenter S, Knipe DM, Fitzgerald KA,
Kurt-Jones EA. 2014. Interferon γ-inducible Protein (IFI) 16 Transcriptionally Regulates
Type I Interferons and Other Interferon-stimulated Genes and Controls the Interferon
Response to both DNA and RNA Viruses. *J Biol Chem* **289**:23568–23581.
doi:10.1074/jbc.M114.554147
Toohey K, Wehrly K, Nishio J, Perryman S, Chesebro B. 1995. Human immunodeficiency virus
envelope V1 and V2 regions influence replication efficiency in macrophages by affecting
virus spread. *Virology* **213**:70–79. doi:10.1006/viro.1995.1547
Tsu BV, Agarwal R, Gokhale NS, Kulsuptrakul J, Ryan AP, Fay EJ, Castro LK, Beierschmitt C,
Yap C, Turcotte EA, Delgado-Rodriguez SE, Vance RE, Hyde JL, Savan R, Mitchell PS,
Daugherty MD. 2023. Host-specific sensing of coronaviruses and picornaviruses by the
CARD8 inflammasome. *PLOS Biol* **21**:e3002144. doi:10.1371/journal.pbio.3002144
Varghese V, Mitsuya Y, Fessel WJ, Liu TF, Melikian GL, Katzenstein DA, Schiffer CA, Holmes
SP, Shafer RW. 2013. Prototypical Recombinant Multi-Protease-Inhibitor-Resistant
Infectious Molecular Clones of Human Immunodeficiency Virus Type 1. *Antimicrob*
Agents Chemother **57**:4290–4299. doi:10.1128/AAC.00614-13
Vermeire J, Naessens E, Vanderstraeten H, Landi A, Iannucci V, Van Nuffel A, Taghon T,
Pizzato M, Verhasselt B. 2012. Quantification of reverse transcriptase activity by real-

time PCR as a fast and accurate method for titration of HIV, lenti- and retroviral vectors.
PloS One **7**:e50859. doi:10.1371/journal.pone.0050859

Walter BL, Wehrly K, Swanstrom R, Platt E, Kabat D, Chesebro B. 2005. Role of low CD4 levels
in the influence of human immunodeficiency virus type 1 envelope V1 and V2 regions on
entry and spread in macrophages. *J Virol* **79**:4828–4837. doi:10.1128/JVI.79.8.4828-
4837.2005

Wang Q, Clark KM, Tiwari R, Raju N, Tharp GK, Rogers J, Harris RA, Raveendran M, Bosinger
SE, Burdo TH, Silvestri G, Shan L. 2024. The CARD8 inflammasome dictates HIV/SIV
pathogenesis and disease progression. *Cell* **187**:1223-1237.e16.
doi:10.1016/j.cell.2024.01.048

Wang Q, Gao H, Clark KM, Mugisha CS, Davis K, Tang JP, Harlan GH, DeSelm CJ, Presti RM,
Kutluay SB, Shan L. 2021. CARD8 is an inflammasome sensor for HIV-1 protease
activity. *Science* **371**:eabe1707. doi:10.1126/science.abe1707

Wannamaker W, Davies R, Namchuk M, Pollard J, Ford P, Ku G, Decker C, Charifson P, Weber
P, Germann UA, Kuida K, Randle JCR. 2007. (S)-1-((S)-2-{[1-(4-amino-3-chloro-phenyl)-
methanoyl]-amino}-3,3-dimethyl-butanoyl)-pyrrolidine-2-carboxylic acid ((2R,3S)-2-
ethoxy-5-oxo-tetrahydro-furan-3-yl)-amide (VX-765), an orally available selective
interleukin (IL)-converting enzyme/caspase-1 inhibitor, exhibits potent anti-inflammatory
activities by inhibiting the release of IL-1beta and IL-18. *J Pharmacol Exp Ther* **321**:509–
516. doi:10.1124/jpet.106.111344

Xie M, Leroy H, Mascarau R, Woottum M, Dupont M, Ciccone C, Schmitt A, Raynaud-Messina
B, Vérollet C, Bouchet J, Bracq L, Benichou S. 2019. Cell-to-Cell Spreading of HIV-1 in
Myeloid Target Cells Escapes SAMHD1 Restriction. *mBio* **10**:10.1128/mbio.02457-19.
doi:10.1128/mbio.02457-19

Zhang C, Song J-W, Huang H-H, Fan X, Huang L, Deng J-N, Tu B, Wang K, Li J, Zhou M-J,
Yang C-X, Zhao Q-W, Yang T, Wang L-F, Zhang J-Y, Xu R-N, Jiao Y-M, Shi M, Shao F,
Sékaly R-P, Wang F-S. 2021. NLRP3 inflammasome induces CD4$^+$ T cell loss in
chronically HIV-1–infected patients. *J Clin Invest* **131**. doi:10.1172/JCI138861

Zhong P, Agosto LM, Ilinskaya A, Dorjbal B, Truong R, Derse D, Uchil PD, Heidecker G, Mothes
W. 2013. Cell-to-Cell Transmission Can Overcome Multiple Donor and Target Cell
Barriers Imposed on Cell-Free HIV. *PLOS ONE* **8**:e53138.
doi:10.1371/journal.pone.0053138

822
823

Supplemental Note

There have been 4 different sublines of THP-1 cells previously characterized (Kasai et al., 2022). Using short tandem repeat (STR) profiling, we were able to distinguish the WT THP-1 cell stocks that were used as the parental line for knockouts and complemented knockouts in this work and our prior work (Kulsuptrakul et al., 2023), as distinct from WT THP-1 cells sourced from ATCC at 3 different loci (**Figure A1A**). Of note, unlike the THP-1 cells used here (referred to as JK THP-1), ATCC THP-1 cells elicited IL-1β secretion in the absence of HIV-1 infection in the presence of 20µg/mL DEAE-dextran (**Figure A1B**). Unless otherwise specified, any mention of "THP-1 cells" are referring to our JK THP-1 cells, not ATCC THP-1 cells. Nonetheless, given the sensitivity of some THP-1 sublines to elicit an inflammasome response in the presence of DEAE-dextran, we assessed whether or not we could establish systems to measure HIV-1 induced CARD8-dependent inflammasome activation in the absence of DEAE-dextran. Thus, we infected either wildtype (WT) or *CARD8* knockout (KO) THP-1 cells with wildtype HIV-1$_{LAI}$ in either the presence or absence of DEAE-dextran and measured cell death and IL-1β secretion 24 hours post-infection as readouts of inflammasome activation. We found that despite achieving similar levels of infection (20-30%) as measured by intracellular p24^{gag} after spinoculation with and without DEAE-dextran (**Figure A1C,** left), we detected robust CARD8-dependent inflammasome activation in WT THP-1 cells infected only in the presence of DEAE-dextran (**Figure A1C,** middle and right). These data suggest that cationic polymer is necessary to observe HIV-dependent CARD8 inflammasome activation in our cell-free system.

A

Cell Line:	JK THP-1		ATCC THP-1	
Expasy Best Match Cell Line No:	CVCL_5115		CVCL_E2QQ	
Marker	Allele 1	Allele 2	Allele 1	Allele 2
AMEL	X	Y	X	Y
D3S1358	15	17	15	17
D1S1656	16	17	16	17
D2S441	10	11	10	11
D10S1248	14		14	
D13S317	8	13	13	
Penta E	11	15	11	15
D16S539	11	12	11	12
D18S51	13	14	13	14
D2S1338	17	18	17	18
CSF1PO	11	13	11	13
Penta D	10	12	10	12
TH01	8	9.3	8	9.3
vWA	16	17	16	
D21S11	30	31.2	30	31.2
D7S820	10		10	
D5S818	11	12	11	12
TPOX	8	11	8	11
DYS391	10		10	
D8S1179	10	14	10	14
D12S391	18	19	19	
D19S433	12.2	13	12.2	13
FGA	24	25	24	25
D22S1045	11	15	11	15

B

Inflammasome activation
(JK THP-1 vs ATCC THP-1)

C

Infectivity — Inflammasome activation — Cell Death

845
Figure A1. Characterization of THP-1 cells. (A) Promega GenePrint® 24 system STR analysis
summary of our JK THP-1 cells versus ATCC THP-1 cells. Cell line authentication was done by
TransnetYX, Inc. by following the protocol described in ANSI/ATCC ASN-0002-2011. The STR alleles
were searched on the ATCC Database and the Expasy best match cell numbers for each cell line had a
100% database match. Distinguishing loci are highlighted in yellow and distinguishing alleles are in red.
(B) JK and ATCC THP-1 cells were primed with Pam3CSK4 overnight then treated with increasing doses
of DEAE-dextran for 24 hours before probing for IL-1β secretion. (**C**) Wildtype (WT) or *CARD8* knockout
(KO) THP-1 cells were infected with wildtype HIV-1$_{LAI}$ at the same MOI in the presence or absence of
DEAE-dextran (10µg/mL) then harvested after 24 hours and assayed for: left) percent infection via
intracellular p24^{gag}, middle) inflammasome activation by IL-1β secretion via IL-1R reporter assay, and
right) cell death via propidium iodide (PI) dye uptake using flow cytometry. %PI positive and IL-1 levels
are normalized to mock control. Dotted line indicates limit of detection (LoD). Datasets represent mean ±
SD (n=2 biological replicates). Two-way ANOVA with (B) Sidak's or (C) Tukey's test using GraphPad
Prism 10. ns = not significant, *p<0.05,**p<0.01, ***p<0.001, ****p<0.0001.
860

861 **Supplemental Note References**
862

Kasai F, Hirayama N, Fukushima M, Kohara A, Nakamura Y. 2022. THP-1 reference data:
Proposal of an in vitro branched evolution model for cancer cell lines. *Int J Cancer*
151:463–472. doi:10.1002/ijc.34019
Kulsuptrakul J, Turcotte EA, Emerman M, Mitchell PS. 2023. A human-specific motif facilitates
CARD8 inflammasome activation after HIV-1 infection. *eLife* **12**:e84108.
doi:10.7554/eLife.84108

869