Cellular and circuit features distinguish dentate gyrus semilunar granule cells and granule cells activated during contextual memory formation.

Laura Dovek[1,2], Krista Marrero[2], Edward Zagha[1,3,4] and Vijayalakshmi Santhakumar[1,2,3]

[1]Biomedical Sciences Graduate Program [2]Department of Molecular, Cell and Systems Biology, [3]Neuroscience Graduate Program [4]Department of Psychology, University of California Riverside, Riverside, California 92521.

Short Title: Circuit mechanisms of dentate gyrus neurons active during memory formation

Word count for Abstract: 233 words

Number of pages: 36

Number of figures: 5　　　　Supplementary figures: 4　　　　Color figures: 5

Number of tables: 0.　　Number of supplementary tables: 1

Acknowledgements: This work is supported by National Institutes of Health (NIH) NINDS R01NS069861, R01NS097750 to V.S., NIH/NINDS F31NS124290 to L.D.

Author contributions: L.D performed experiments; L.D, K.M and E.Z analyzed data; L.D, K.M. E.Z and V.S. interpreted results of experiments; L.D. and K.M. prepared figures; L.D, and V.S. conception and design of research; L.D., K.M. and V.S. drafted manuscript; L.D., K.M. E.Z and V.S. finalized and approved manuscript.

Correspondence:

Vijayalakshmi Santhakumar, PhD

Department of Molecular, Cell and Systems Biology

University of California, Riverside

3401 Watkins Drive (Rm 1308 Spieth Hall) Riverside, CA 92521

E-mail: vijayas@ucr.edu

Abstract

The dentate gyrus is critical for spatial memory formation and shows task related activation of

cellular ensembles considered as memory engrams. Semilunar granule cells (SGCs), a sparse

dentate projection neuron subtype distinct from granule cells (GCs), were recently reported to

be enriched among behaviorally activated neurons. However, the mechanisms governing SGC

recruitment during memory formation and their role in engram refinement remains unresolved.

By examining neurons labeled during contextual memory formation in TRAP2 mice, we

empirically tested competing hypotheses for GC and SGC recruitment into memory ensembles.

In support of the proposal that more excitable neurons are preferentially recruited into memory

ensembles, SGCs showed greater sustained firing than GCs. Additionally, SGCs labeled during

memory formation showed less adapting firing than unlabeled SGCs. Our recordings did not

reveal glutamatergic connections between behaviorally labeled SGCs and GCs, providing

evidence against SGCs driving local circuit feedforward excitation in ensemble recruitment.

Contrary to a leading hypothesis, there was little evidence for individual SGCs or labeled

neuronal ensembles supporting lateral inhibition of unlabeled neurons. Instead, pairs of GCs

and SGCs within labeled neuronal cohorts received more temporally correlated spontaneous

excitatory synaptic inputs than labeled-unlabeled neuronal pairs, validating a role for correlated

afferent inputs in neuronal ensemble selection. These findings challenge the proposal that

SGCs drive dentate GC ensemble refinement, while supporting a role for intrinsic active

properties and correlated inputs in preferential SGC recruitment to contextual memory engrams.

Impact Statement

Evaluation of semilunar granule cell involvement in dentate gyrus contextual memory

processing supports recruitment based on intrinsic and input characteristics while revealing

limited contribution to ensemble refinement.

29
Major subject area, keywords and organisms
31
Semilunar granule cell, inhibition, memory, engram, circuit, hippocampus

Introduction

The ability of neural circuits to represent unique experiences and events as distinct neuronal representations that can be recalled and updated is fundamental to memory formation. The hippocampal dentate gyrus (DG) is a circuit considered central for both novelty detection and to the formation of episodic memories (Hunsaker et al., 2008; Liu et al., 2012; Hainmueller and Bartos, 2020; Danieli et al., 2023). The DG receives dense information from diverse cortical regions through the perforant path projections from the entorhinal cortex (van Groen et al., 2003). Yet relatively few of the numerous closely packed dentate projection neurons, granule cells (GCs), are activated and engage downstream hippocampal circuits. This sparsening of activity is proposed as critical for pattern separation, a process by which the dentate gyrus helps disambiguate similar memories (McHugh et al., 2007; Hainmueller and Bartos, 2018). Still, the mechanisms that govern how select subsets of neurons are activated during memory formation are not fully understood.

The cellular representations of memories, known as *engrams,* refer to distinct groups of neurons activated during memory acquisition (Semon, 1909; Josselyn et al., 2015; Josselyn and Tonegawa, 2020). Recently, a sparse subset of DG projection neurons, known as semilunar granule cells (SGCs), have been found to be overrepresented among neurons labeled by the expression of the activity dependent immediate early gene (IEG) c-Fos during hippocampus dependent behaviors in TRAP2 reporter mice (Erwin et al., 2020). SGCs, like GCs, have molecular layer dendrites and project axons to CA3 (Williams et al., 2007; Afrasiabi et al., 2022). However, unbiased cluster analyses of morphometric data have revealed that structural features can reliably distinguish SGCs from GCs based on their wider dendritic arbor, greater soma width to length ratio, and more numerous primary dendrites (Williams et al., 2007; Gupta et al., 2020; Afrasiabi et al., 2022). Despite SGCs being estimated to make up only ~3% of the total GC population (Save et al., 2019), their preferential activation in memory tasks suggests that

SGCs may possess unique physiology or connectivity to support recruitment to engrams.

However, why SGCs may be preferentially recruited, and whether they shape DG ensemble

refinement is unresolved.

There are complementary theories for why certain neurons are selectively activated

during memory formation and for how the active cell ensembles may be refined by circuit

processes. One hypothesis is that neurons are recruited to memory ensembles based on

greater excitability (Yiu et al., 2014; Gouty-Colomer et al., 2016). According to this hypothesis,

distinct cohorts of neurons may have higher intrinsic excitability during certain periods; this

propensity biases them to fire preferentially in response to inputs and to be recruited into

behaviorally activated ensembles (Yiu et al., 2014). Relatedly, it has been suggested that

newborn GCs are preferentially recruited to engrams because of their higher excitability (Kee et

al., 2007). However, it is not known whether intrinsic physiological features of SGCs, which

show sustained afferent driven firing (Larimer and Strowbridge, 2010; Afrasiabi et al., 2022)

support their disproportionate representation among behaviorally activated DG ensembles.

In addition to intrinsic properties, neuronal recruitment can be refined by local circuit

feedforward or recurrent excitation. One possibility is that glutamatergic interconnectivity aids in

engram refinement. Indeed, reports of higher connection probability and strengthening of

excitatory synapses between GCs and CA3 pyramidal cells labeled based on IEG expression

following fear conditioning (Ryan et al., 2015) support this possibility. Although GCs typically do

not innervate other GCs, SGCs have axon collaterals in the molecular layer (Williams et al.,

2007; Save et al., 2019), which positions them to potentially form synaptic contacts with GCs.

However, whether SGCs directly activate GCs and whether SGCs refine their recruitment to

behaviorally active neuronal ensembles remain to be tested. Evaluation of synaptic connectivity

between neuronal pairs in ensembles labeled based on IEG expression during memory

formation would allow us to test whether recurrent glutamatergic connections support DG

ensemble recruitment. Simultaneously, since connectivity between SGCs and GCs is likely to

be sparse, this experimental paradigm allows us to address the open question of whether SGCs

synaptically activate GCs.

A leading hypothesis for DG circuit refinement of behaviorally active neuronal

ensembles, particularly in the context of pattern separation, is through lateral feedback inhibition

of surrounding GCs (Walker et al., 2010; Cayco-Gajic and Silver, 2019; Guzman et al., 2021;

Borzello et al., 2023). The characteristic robust feedback inhibition in the DG holds promise as a

mechanism by which activated engram neurons recruit interneurons to selectively inhibit

surrounding neurons(Espinoza et al., 2018). SGCs are ideally poised to mediate this effect as

their axon collaterals have been shown to form perisomatic synapses on parvalbumin

expressing fast spiking basket cells known for their feedback inhibition of GCs (Rovira-Esteban

et al., 2020). Consistent with a role for SGCs in supporting feedback inhibition, afferent evoked

persistent firing in SGCs is correlated with sustained basket cell and hilar interneuron firing and

prolonged inhibitory synaptic barrages in GCs and SGCs (Larimer and Strowbridge, 2010;

Afrasiabi et al., 2022). Activation of a large cohort of optogenetically driven GC ensembles

elicits inhibition in surrounding GCs (Stefanelli et al., 2016). However, whether the sparse

neuronal populations activated during behaviorally driven encoding mediate lateral inhibition of

surrounding neurons remains to be tested.

Finally, it is reasonable to posit that precise connectivity of afferent inputs determine

downstream activation of a sparse population of DG neurons. Indeed, there is evidence for

input-dependent recruitment of neuronal cohorts in the amygdala during fear conditioning

(Gouty-Colomer et al., 2016). However, whether shared inputs constrain coactivation of neurons

and whether input specificity acts in concert with intrinsic and circuit features to determine which

GCs and SGCs are activated is currently unknown. While several studies have focused on DG

engram formation (Liu et al., 2012; Ryan et al., 2015), only recently has there been an attempt

to explicitly distinguished GCs from SGCs (Erwin et al., 2020). Thus, the specific circuit

mechanisms underlying behaviorally relevant DG ensemble refinement during memory

encoding and roles of SGCs remain to be determined.

Here we used TRAP2 transgenic mice for c-Fos driven labeling of DG ensembles during

behavioral tasks (Guenthner et al., 2013; DeNardo et al., 2019) to label active DG ensembles

and undertook ex vivo dual patch clamp and optogenetic recordings in morphologically

characterized GCs and SGCs. We use these data to evaluate competing proposals for

refinement of cellular ensemble representations in the DG. We specifically focused on

differential recruitment of GCs and SGCs in DG ensembles and the potential roles for SGCs in

shaping DG circuit processing and refining neuronal ensembles.

118
119

Results:

Semilunar granule cells are reliably recruited during contextual memory formation.

The DG is a primary relay for memory processing (Amaral et al., 2007). However, the mechanisms by which memory-related cellular ensembles are selectively activated during memory encoding are not fully understood. For a memory encoding context suitable for physiological analyses, we compared the Barnes Maze (BM) and an enriched environment (EE) exposure. We were particularly interested in identifying a context that activated large cohorts of DG neurons, thereby enabling microcircuit analyses via physiological recordings. Behaviorally activated "*engram*" neurons, referred to henceforth as "labeled neurons", are neurons in TRAP2 mice induced to express the reporter (tdT or Chr2-YFP) downstream of the activity-dependent IEG c-Fos during a behavioral task. "Unlabeled neurons" lack reporter expression. Littermate pairs of TRAP2-tdT mice (4 pairs) were either trained in the BM spatial learning task (Supplementary Fig. 1) or exposed to an EE. Both cohorts were induced with tamoxifen during respective behavioral paradigms, on day 6 of BM acquisition or halfway through the 1-day EE exposure, to label active neurons (Fig. 1A-B). Comparison of the number of DG c-Fos expressing (tdT positive) neurons in hippocampal sections from mice one week after tamoxifen induction revealed significantly more tdT labeled neurons following EE exposure than after BM acquisition (Fig. 1C-E; # of tdT labeled cells per slice: EE: 33.90 ± 2.13, BM 13.43 ± 0.90, n=40 slices from 4 animals per group, p=0.0409 by nested t-test). Consistent with previous reports (Erwin et al., 2020), the suprapyramidal (upper) blade of the DG showed more neurons labeled than the infrapyramidal (lower) blade following both BM training and EE exposure (Fig. 1F). To determine whether tagged neurons show task specific reactivation one week after induction, mice were exposed to EE prior to perfusion and sections were immunostained for c-Fos. The distribution of neurons immunolabeled for c-Fos following EE exposure showed no apparent difference between mice previously exposed to BM followed by EE and those exposed to EE twice (Fig. 1Cii, Fig. 1Dii). However, consistent with memory related neuronal tagging, mice with

prior exposure to EE showed greater co-labeling of tdT positive neurons with c-Fos

immunostaining than mice that were initially trained in the BM task (Fig. 1G; % of co-

labeled/total labeled: BM: 2.28 ± 0.46%, EE: 6.8 ± 0.97% p=0.0003 by nested t-test). The

results suggest that a cohort of neurons, tagged following EE, reactivate when reintroduced to

the same environment, thereby demonstrating memory-specific activation. Therefore, in

subsequent experiments, we presumed that cells labeled by task related c-Fos driven reporter

expression represent engram cells. Since EE resulted in greater overall DG neuron labeling and

stable reactivation of a subset of neurons after one week, we adopted EE as the preferred

paradigm to label task-related neuronal ensembles for circuit level analysis.

We examined tagged neurons in sections from mice that underwent BM navigation and

EE exposure to determine the proportional recruitment of SGCs. SGCs were distinguished from

GCs based on (1) presence of multiple primary dendrites, (2) greater soma width than height,

(3) wide dendritic arbor and/or (4) location in or close to the inner molecular layer (Fig. 1H).

These criteria were based on our prior unbiased cluster analysis of GC and SGC morphology

(Gupta et al., 2020; Afrasiabi et al., 2022). The morphology-based classification revealed that

33.86 ± 2.18% of neurons labeled during BM acquisition and 27.83 ± 1.33% during EE exposure

were SGCs (Fig. 1I; p=0.1143 by nested t-test, based on 40 sections from 4 mice). Since SGCs

represent less than 5% of DG projection neurons (Save et al., 2019), these data suggest

preferential activation of SGCs during dentate dependent contextual memory formation.

Notably, the proportional recruitment of SGCs labeled following behavior was not different

between the BM navigation and EE exposure (Fig. 1I). These findings make a compelling case

for leveraging EE exposure to study SGC involvement in dentate-dependent microcircuits.

Contribution of intrinsic physiology to activity dependent neuronal labeling.

To test if the intrinsic physiology of GCs and SGCs labeled during EE differ from their

unlabeled counterparts, we performed whole-cell recordings from labeled and unlabeled GCs

and SGCs in slices from TRAP2ChR2/eYFP mice one week after tamoxifen induction during EE

exposure. Labeled and unlabeled neurons in the GCL and IML were visualized under

epifluorescence (λ=505 nm) and IR/DIC respectively. Recorded neurons were classified as GC

or SGC based on post hoc morphological analysis of biocytin fills (Fig. 2A-B) (Williams et al.,

2007; Gupta et al., 2020; Afrasiabi et al., 2022). Depolarizing response to blue light activation

(0.9mW, λ=470 nm, 10ms) of ChR2 was used to functionally validate cell labeling (Fig. 3E, Fig.

4E). Consistent with earlier studies (Afrasiabi et al., 2022), there was no cell-type specific

difference in resting membrane potential (RMP) between GCs and SGCs (Fig. 2C). RMP was

also not different between labeled and unlabeled cells within each cell type. Similarly, while

SGC input resistance (R_{in}) was lower than in GCs, as reported previously (Williams et al., 2007;

Afrasiabi et al., 2022), R_{in} of labeled and unlabeled neurons was not different in either cell type

(Fig. 2D). Examination of responses to a graded current injection revealed divergence of the

firing frequency between GCs and SGCs at current injections >400pA, with GCs showing

progressive reduction in frequency with increasing current injection (Fig. 2E-G) due to an

apparent depolarization block. Consistently, the firing frequency in response to +520pA current

was greater in SGCs than in GCs (Fig. 2H). Again, these cell-type specific differences were

maintained in both labeled and unlabeled neurons. The action potential parameters, including

threshold,   amplitude,   half-width,   fast   afterhyperpolarization   (fAHP),   medium

afterhyperpolarization (mAHP), and latency to first action potential were not different between

cell types or labeling of neurons (Supplemental Fig. 2B-G, Supplemental Table 1). Interestingly,

GCs showed greater amplitude attenuation during continuous firing, which was not observed in

SGCs (Fig. 2I). Once again, these cell-type specific differences were retained in both labeled

and unlabeled neurons. Finally, SGCs show higher adaptation ratios (ratio of duration between

first two and last two spikes in response to 120 pA current injection), indicating less spike

frequency adaptation than in GCs, consistent with previous findings (Williams et al., 2007).

Notably, labeled SGCs showed significantly lower adaptation in firing rate than unlabeled SGCs

(Fig. 2J; $GC_{Labeled}$: 0.33± 0.075; $GC_{Unlabeled}$: 0.28± 0.056; $SGC_{Labeled}$: 0.78 ±0.076; $SGC_{Unlabeled}$:

0.47 ±0.06; 2-Way RM ANOVA main effect of cell type, p=0.006). In contrast, labeled and

unlabeled GCs did not differ in adaptation ratio indicating that the ability to sustain firing may

distinguish labeled SGCs. These data support a role for SGC intrinsic physiology, specifically

non-attenuating, less adapting, and persistent firing in their preferential labeling during activity

dependent neuronal tagging.

Lack of evidence of local feedforward or recurrent excitation between activity driven

neuronal ensembles.

Unlike GCs, SGCs have axon collaterals in the inner molecular layer and granule cell

layer (Fig. 2A,B) (Williams et al., 2007; Save et al., 2019) raising the possibility that they could

activate GC dendrites in the IML. To test the local feedforward/recurrent excitation hypothesis

we conducted dual patch recordings from labeled neuron pairs to identify potential glutamateric

interconnections (Fig. 3A,C). Care was taken to ensure that neurons at a depth of 50 μm or

more from the surface with visible axons were targeted in order to maximize probability of

connections. As noted in Fig. 3B, a majority of the 32 pairwise connections examined between

labeled neurons were between SGCs (Fig. 3B). However, all other possibilities including

$GC_{Labeled}$ to $GC_{Labeled}$, $SGC_{Labeled}$ to $GC_{Labeled}$, and $GC_{Labeled}$ to $SGC_{Labeled}$ were also evaluated.

The presence of spontaneous EPSCs in the neurons recorded under voltage clamp served as

confirmation of overall circuit and slice health (Fig. 3D). Additionally, optically evoked (0.9 mW,

λ=470 nm, 10Hz, 10ms pulses) inward currents provided functional validation of reporter

expression (Fig. 3E). Labeled neuronal pairs were tested for glutamatergic synaptic connections

by depolarizing one of the neurons in current clamp (400pA, 10 pulses for 10ms at 50Hz) and

recording evoked current responses in the other neuron held at -70 mV (Fig. 3F). Recording

configuration was reversed to check for connections in both directions (Fig. 3G). Despite the

presence of spontaneous EPSCs, none of the 32 labeled neuronal pairs tested, including

$SGC_{Labeled}$ to $GC_{Labeled}$ (n=7) and $SGC_{Labeled}$ to $SGC_{Labeled}$ (n=16), showed functional

glutamatergic synaptic connections (Fig. 3F-G). Although our data do not eliminate the

possibility of direct excitatory connections between labeled neurons, they indicate that

glutamatergic interconnections are not critical for activation of DG neuronal ensembles.

230
Limited evidence for neurons in activity driven ensembles supporting lateral inhibition.

The role of surround inhibition and winner-take-all activation has been proposed as a

promising mechanism for establishing memory engrams and mediating dentate processing

(Espinoza et al., 2018; Guzman et al., 2021). SGCs, with preferential recruitment in DG

engrams, sustained firing, and hilar axon collaterals, are ideally suited to drive robust feedback

inhibition (Larimer and Strowbridge, 2010; Walker et al., 2010; Afrasiabi et al., 2022). We

examined the possibility that labeled neurons, particularly labeled SGCs, refine GC activity by

mediating feedback surround inhibition of unlabeled GCs. We tested this by performing dual

recordings from labeled-unlabeled neuronal pairs (Fig. 4A-C). While the majority of our

recordings focused on $SGC_{Labeled}$ to $GC_{Unlabeled}$ (49/63 pairs), we also tested for connections

between $SGC_{Labeled}$ to $SGC_{Unlabeled}$, $GC_{Labeled}$ to $SGC_{Unlabeled}$ and $GC_{Labeled}$ to $GC_{Unlabeled}$ (Fig. 4D).

The ability of wide-field optogenetic activation (0.9 mW, λ=470 nm, 10 pulses, for 10 ms at 10Hz

train) to evoke inward currents validated ChR2 expression in labeled neurons. As expected,

wide-field blue light stimulation failed to evoke inward currents in unlabeled neurons, confirming

the lack of ChR2 expression (Fig. 4E). The recordings also allowed us to test the unlikely

possibility that activation of ChR2 expressing labeled neurons evoked synaptic excitation in

unlabeled neurons. As expected, wide-field light activation, which likely activates multiple

labeled neurons throughout the slice preparation, did not evoke putative polysynaptic EPSCs in

unlabeled neurons (n= 63 pairs tested). Consistently, in paired recordings between labeled and

unlabeled neurons, current-evoked firing in labeled cells failed to evoke EPSCs in unlabeled

cells, (Fig. 4F), underscoring the lack of glutamatergic connectivity between SGCs and GCs.

In paired recordings from labeled SGCs and unlabeled GCs (n=49), depolarization

evoked firing in SGCs (400pA, 10ms, 10 pulses, 50Hz) failed to evoke polysynaptic IPSCs in

unlabeled GCs (Fig. 4G). Notably, despite the lack of evoked IPSCs, the unlabeled GCs

received spontaneous IPSCs indicating that cell and circuit health were not compromised (Fig.

4G). Interestingly, in 1/49 recordings from $SGC_{Labeled}$ to $GC_{Unlabeled}$ pairs, wide-field optogenetic

activation at a light intensity that evoked firing in the recorded labeled SGC, evoked IPSCs in

the unlabeled GC in the absence of direct synaptic connection between the pairs (Fig. 4H-I).

However, in a majority of trials, wide-field optogenetic activation of both labeled GCs and

labeled SGCs failed to evoke IPSCs in unlabeled GCs. Activating labeled neurons did not lead

to IPSCs in unlabeled neurons in any of the $GC_{Labeled}$ to $GC_{Unlabeled}$ or $SGC_{Labeled}$ to $SGC_{Unlabeled}$

pairs tested. Unexpectedly, we identified one pair (out of 63) in which current induced firing in a

labeled GC resulted in robust feedback IPSCs in an unlabeled SGC (Fig. 4J-K). These data

identify that labeled GCs can support feedback inhibition of SGCs. Collectively, these data find

limited evidence for robust lateral inhibition by neurons labeled during EE exposure onto

surrounding unlabeled neurons.

Labeled neuron pairs receive more correlated spontaneous excitatory inputs.

Since microcircuit connectivity and intrinsic physiology could not fully account for task

related coactivation of neurons, we tested the hypothesis that correlated inputs contribute to

ensemble activation. To evaluate whether temporally correlated inputs contribute to ensemble

labeling during EE exposure, we analyzed sEPSC in dual recordings from labeled-labeled (L-L)

and labeled-unlabeled (L-U) pairs for temporal correlation of synaptic event times (Fig. 5A-C).

The sEPSCs were recorded at a holding potential of -70 mV in low chloride internal solution and

the synaptic event times were extracted for analysis. To assess sEPSC temporal correlation, we

defined *peri-occurrence* as the maximum cross correlation of sEPSCs in the two recorded

neurons within a *detection window* and *cooccurrence* as the cross correlation of sEPSCs within

a more restricted time *bin duration* centered around 0 ms (Fig. 5D).

Initially, cross correlation profiles (CCPs) were developed using multiple *detection*

windows (±10ms to ±1s) with corresponding *bin durations* (21 bins within window). Temporal

correlation across full session timelines was not calculated to avoid spuriously high correlation

values from simultaneous absence of events in cell pairs (Cutts and Eglen, 2014). As an

additional measure to avoid specious correlations, sessions with too few events, low event

frequency, and short recording durations were not analyzed. For the data presented below, the

recording durations, event count, and activity rates were not different between L-L and L-U pairs

(Supplemental Fig. 3, number of sEPSCs in counts, L-L: 439±52.83, L-U: 412±48.77, p=0.71;

recording duration in s, L-L: 498±43.76, L-U: 397±48.0, p=0.15; spike rate in Hz, L-L: 0.89±0.09,

L-U: 1.06±0.11, p=0.23).

Temporal correlations of sEPSC event times in a large ±1 s detection window confirmed

an expected high cross correlation of events within the 100 ms central bin (not shown). A small

±10 ms detection window with 1 ms bin width resulted in too few cooccurring events (not

shown). These extremely large and narrow detection windows were not examined further. We

selected detection windows of ±100 ms and ±50 ms with 10 ms and 5 ms bin width,

respectively, to develop CCPs of L-L and L-U sEPSC event times (Supplemental Fig. 4,

example recording sessions with CCP for cooccurrence in L-L pairs, no coincidence in L-U

pairs, and peri-occurrence in L-L pairs, analyzed using a ±100 ms detection window). A

predetermined threshold of 2 standard deviations (2SD) above the mean correlation was

adopted to assess potential differences in temporal correlation between L-L and L-U pairs.

Peri-occurrence, quantified as the maximum cross correlation in the detection window

(R_{max}), was significantly higher in L-L than in L-U pairs for the ±100 ms detection window (R_{max}

in ±100 ms window, L-L: 0.184±0.014, L-U: 0.126±0.016, p=0.018 by t-test in 7 L-L and 8 L-U

pairs). While peri-occurrence in the ±50 ms detection window trended higher in L-L pairs, this

was not significant (R_{max} in ±50 ms window, L-L: 0.124±0.013, L-U: 0.093±0.014, p=0.12 by t-

test in 7 L-L and 8 L-U pairs). Notably, the R_{max} in sEPSC event times from 6/7 L-L pairs

crossed the threshold (2SD above the mean correlation) within each detection window and

failed to do so in 7/8 recordings in L-U pairs. These findings were consistent regardless of

whether we adopted ±100 ms or ±50 ms detection windows (Fig. 5E-F insets).

To determine if event correlations in neuronal pairs deviated from random, event time

correlations in the recorded (*temporally aligned*) data were compared with the correlations

developed from corresponding temporally *jittered* data sets. Temporally jittered data sets

included 100 iterations in which event times were jittered at random with a ±500 ms bound and

appended to one of the paired cell event timelines (Fig. 5E-F). Event time center bin correlations

in the recorded (*temporally aligned*) data sets were significantly higher than in the *jittered* data

sets for both detection windows (Fig. 5G main effect of data alignment, 10 ms bin:

$F(1,211)=63.16$, p<0.0001 by TW-ANOVA, 5 ms bin, $F(1,211)=71.60$, p<0.0001 by TW-

ANOVA). Cooccurrence, quantified as the correlation in the center bin, was higher in L-L pairs

both within the 10 ms bin (Fig. 5E,G, 10 ms center bin correlation, L-L: 0.117±0.025, L-U:

0.087±0.014, p=0.0075 by TW- ANOVA in 7 L-L and 8 L-U pairs) and within the 5 ms bin (Fig.

5F, G, 5 ms center bin correlation, L-L: 0.081±0.019, L-U: 0.042±0.009, p<0.0001 by TW-

ANOVA in 7 L-L and 8 L-U pairs

Finally, we evaluated the ability of cooccurring events to predict an L-L versus L-U

recording session. The receiver operating characteristic (ROC) curve of true versus false

positive rates defined the area under the ROC curve (AUROC) for the center 10 ms and 5 ms

bin across the ±100 ms and ±50 ms detection windows, respectively (Fig. 5H). Center bin

classification performed better than chance at predicting whether a recorded session was L-L

versus L-U ($AUROC_{Chance} = 50\%$; $AUROC_{10msCenter} = 66.96\%$; $AUROC_{5msCenter} = 71.43\%$). Thus,

the 10 ms and 5 ms center event correlations were predictive of whether a pair of recorded

neurons were likely to be a pair of labeled neurons or a labeled-unlabeled pair. Together these

results support a role for correlated inputs in driving shared neuronal activation during

contextual memory formation.

Discussion:

The recent characterization of semilunar granule cells as a unique dentate projection neuron subtype overrepresented amongst behaviorally recruited DG neurons has raised the intriguing possibility that SGCs may play a distinct role in shaping DG ensemble activity (Walker et al., 2010; Erwin et al., 2020; Afrasiabi et al., 2022). Here we evaluated competing hypotheses involving mechanisms governing recruitment of GC and SGC populations during a behavioral experience. Our data identify that *intrinsic properties of SGCs,* specifically their *less adapting firing* characteristics, likely enable preferential recruitment of SGCs among neurons labeled based on IEG expression. At the circuit level, neurons in a shared DG ensemble receive *more correlated spontaneous excitatory inputs* than neurons without shared activation, suggesting a role for common afferent inputs in behaviorally driven ensemble recruitment. Whereas GCs and downstream CA3 neurons with shared behavioral activation develop preferential glutamatergic connectivity (Ryan et al., 2015)), we found no evidence for local feedforward or recurrent excitation in DG neurons labeled as part of a memory trace. Unexpectedly, although lateral inhibition has been proposed as a leading mechanism for dentate engram refinement (Stefanelli et al., 2016), our experiments revealed that activation of labeled DG neurons rarely drove inhibitory synaptic currents in unlabeled neurons. Interestingly, we find approximately a third of the projection neurons activated as a part of a dentate dependent spatial navigation task and during exposure to enriched environment had morphology consistent with SGCs. It is possible that the reduced action potential accommodation and attenuation in SGCs contributes to enhanced SGC firing and c-Fos expression during afferent activation. This greater activity dependent c-Fos expression in SGCs may result in their preferential labeling as part of neuronal ensembles labeled during behavioral tasks. Together, these results identify that behaviorally relevant activation of SGCs and GCs in DG neuronal ensembles are determined by a combination of sustained firing characteristics of SGCs and shared afferent drive rather than by selective circuit level refinement by recurrent excitation or by lateral inhibition.

Studies evaluating mechanisms of ensemble recruitment during fear and aversive memory

encoding, by experimentally enhancing excitability or CREB expression in a sparse population

of amygdala neurons, have proposed that neurons with higher excitability outcompete

neighboring cells for allocation to behaviorally activated neuronal ensembles (Han et al., 2007;

Zhou et al., 2009; Sano et al., 2014; Yiu et al., 2014; Gouty-Colomer et al., 2016). However, our

data revealed no difference between intrinsic physiology and active properties between labeled

and unlabeled GCs. This is consistent with a previous report that firing threshold and input

resistance of GCs labeled during fear conditioning and recorded 24 hours later were not

different from unlabeled GCs (Ryan et al., 2015), although GCs and SGCs were not explicitly

distinguished. We find that unlike GCs, SGCs show less adapting firing and reduced spike

amplitude attenuation resulting in more sustained firing, particularly in response to large current

injections. Moreover, labeled SGCs had lower spike frequency adaptation than unlabeled

SGCs, indicating that sustained firing may predispose SGCs to activation during behavioral

encoding. Indeed, the sustained firing during depolarizing current injections larger than 400 pA

(this study) and in response to afferent input (Larimer and Strowbridge, 2010; Afrasiabi et al.,

2022)are quintessential functional differences between GCs and SGCs. This sustained SGC

firing is ideally suited to induce more immediate early gene (c-Fos or ARC) expression and

could contribute to higher than expected labeling of SGCs during memory engram encoding.

This is also supported by enrichment of activity dependent markers including PENK in

behaviorally activated DG neurons, including SGCs labeled in TRAP2 mice (Erwin et al., 2020)).

Our timeline for recording, one week after c-Fos-dependent ensemble labeling, does not allow a

comparison of neuronal excitability at or before task performance. Nevertheless, our data

demonstrating more sustained firing in SGCs and selectively reduced adaptation in labeled

SGCs supports a role for greater neuronal activity in preferential recruitment of SGCs to task

related dentate engrams.

It is possible that the sustained firing in SGCs, as well as higher NMDAR-mediated

currents (Larimer and Strowbridge, 2010), contribute to their increased representation (~30%)

among behaviorally tagged DG neuronal ensembles compared to their relative population (~3-

5%) (Save et al., 2019). Furthermore, the wider dendritic arbors of SGCs are ideally positioned

to receive distributed inputs and could support their preferential recruitment during behaviors. In

this regard, whether SGCs and GCs differ in the inputs they receive from the entorhinal cortex is

currently unknown. Although we find higher than expected SGCs among labeled neuronal

ensembles, the SGC labeling quantified in our study is considerably lower than the approximate

80% representation in DG engrams reported previously (Erwin et al., 2020). It is possible that

the estimation based on dendritic span and input resistance of engram neurons filled following

physiological studies contributed to a greater overrepresentation of SGCs among recorded

engram neurons in prior studies (Erwin et al., 2020) than our estimates based on morphometric

analysis of labeled neurons. This is not surprising because SGCs, especially those in the

sparsely populated molecular layer are more readily visualized and accessed for patch

physiology than labeled GCs in the densely packed cell layer. Indeed, SGCs represent greater

than 70% of the labeled neurons in our recordings as well (Supplemental Fig. 2). Thus, while

conservative, our analysis based on a combination of morphometric features previously

validated by unbiased analyses (Gupta et al., 2020; Afrasiabi et al., 2022) is likely to reflect the

proportional labeling of SGCs.

Neurons in shared ensembles, including the DG to CA3 projections, show preferential

connectivity and selective synaptic strengthening (Ryan et al., 2015; Rao-Ruiz et al., 2021).

While connectivity between GCs is rarely observed in the healthy DG, whether SGCs with axon

collaterals in the molecular layer make functional synaptic contacts on GC had not been

examined. We leveraged findings that DG ensembles stably reactivate and engage downstream

circuits up to 12 days after encoding (Kitamura et al., 2017) to evaluate local connectivity among

SGCs and GCs in behaviorally recruited DG ensembles one week after encoding. Our paired
recordings did not find evidence for glutamatergic connections between labeled SGCs and GCs,
indicating that local DG engram refinement is not supported by mutual synaptic strengthening.
Moreover, we observed no evidence of direct synaptic connectivity between GCs and SGCs
regardless of whether the neurons were labeled or unlabeled. While it is possible that slice
preparation could sever axon collaterals, we routinely record from cells over 50 μm below the
surface and recovered extensive axon collaterals from SGC-GC pairs. Moreover SGCs have
compact axonal distribution (Gupta et al., 2020), minimizing the possibility that lack of
connections was a consequence of severed axons. Furthermore, even wide-field illumination to
activate ChR2 positive terminals failed to evoke EPSCs in unlabeled GCs or SGCs, consistent
with lack of connectivity between labeled and unlabeled SGC-GC pairs. Taken together with the
evidence for increased overlap between DG ensembles labeled during encoding and recall, the
limited glutamatergic interconnectivity among GCs and SGCs supports an instructive role for
afferent inputs in DG ensemble recruitment. Consistent with this proposal, we found that the
event timing of spontaneous glutamatergic inputs to pairs of labeled DG neurons were more
correlated than inputs to labeled-unlabeled pairs suggesting that labeled neuronal pairs may
receive correlated input streams. Moreover, input correlation was effective in discriminating
between labeled-labeled versus labeled-unlabeled neuronal pairs, further supporting the role for
input dependent recruitment of neuronal ensembles.

Lateral inhibition has been long considered a promising mechanism for discriminating
amongst behaviorally relevant DG neuronal ensembles (Espinoza et al., 2018; Cayco-Gajic and
Silver, 2019; Guzman et al., 2021; Borzello et al., 2023). In support of this proposal, activation of
a large cohort of GCs, experimentally driven to fire during a behavior, was shown to mediate
robust surround inhibition of GCs (Stefanelli et al., 2016). Moreover, SGCs, with their ability to
robustly activate feedback inhibition (Larimer and Strowbridge, 2010; Afrasiabi et al., 2022),

have been proposed as an ideal cell type to drive surround inhibition (Walker et al., 2010).

However, our analysis of behaviorally tagged SGCs and GCs found limited evidence for lateral

inhibition of GCs. Even wide-field optical stimulation of labeled neurons, which would be

expected to activate labeled terminals on interneurons due to the recruitment of intact and

severed axons, rarely elicited lateral inhibition (1 of 55 trials). The apparent contrast between

prior reports of robust lateral inhibition in influencing DG engram size (Stefanelli et al., 2016)

and our results of sparse surround inhibition from behaviorally tagged ensembles could arise

from the difference between the high GC numbers in experimentally generated engrams

compared to the sparse labeling of behaviorally recruited neurons. Indeed, although existing

evidence strongly supports the role for robust interneuron activation in constraining engram size

by reducing network excitability (Morrison et al., 2016; Stefanelli et al., 2016), results of our

microcircuit analysis in hippocampal slices suggest that SGCs have limited contribution to

ensemble refinement by lateral inhibition. Curiously, we identified inhibition from a labeled GC to

an unlabeled SGC and optically induced inhibition of an unlabeled GC, which does indicate

presence of sparse lateral inhibition in the circuit. Overall, while there is considerable evidence

for robust lateral inhibition in regulating DG activity, our data do not support the hypothesis that

ensemble refinement is determined by surround inhibition mediated by a sparse population of

behaviorally active SGCs and GCs.

The contextual behavioral paradigms adopted in this study are known to engage the DG.

However, they resulted in considerably sparser labeling than reported in contextual fear

conditioning paradigms adopted in prior studies (Liu et al., 2012; Liu et al., 2014; Ryan et al.,

2015; Roy et al., 2017)). In order to specifically target neuronal cohorts activated during DG

dependent spatial learning, we initially examined neuronal activation during the BM spatial

navigation task involving spatial learning over multiple trials (Gawel et al., 2019). Tamoxifen

treatment on day 6 labeled a sparse cohort of neurons, which was not compatible with circuit

level analysis. Since the DG is preferentially activated by novelty (Hainmueller and Bartos,

2020; Mazurkiewicz et al., 2022; Borzello et al., 2023), we reasoned that learning related

decrease in novelty may have contributed to sparse DG labeling during day 6 of BM spatial

navigation. Consistent with a role for novelty in DG ensemble activation (Mazurkiewicz et al.,

2022), tamoxifen induction during a single episode of enriched environment exposure reliably

labeled a larger cohort of DG neurons, thereby enabling circuit analysis. Another caveat is the

possibility that use of the TRAP2 system may miss active neurons expressing other IEGs

(Heroux et al., 2018). Nevertheless, c-Fos driven labeling in TRAP2 mice remains the current

best approach for activity dependent labeling, especially of DG neurons (Kawashima et al.,

2014).

In summary, we find that SGCs represent about a third of dentate projection neurons

labeled based on c-Fos expression during the contextual memory encoding, which is a

considerable overrepresentation relative to their known population density. We propose that

their unique sustained firing characteristics and temporal precision of afferent inputs may

support their preferential labeling during activity dependent labeling of memory ensembles.

Taken together, these data support a role for correlated inputs, the ability to sustain action

potential firing, and sparse surround inhibition rather than glutamatergic interconnectivity as key

determinants for recruitment of neurons to dentate memory ensembles.

Materials and Methods:

Animals

All experiments were conducted under IACUC protocols approved by the University of

California at Riverside and conformed with ARRIVE guidelines. c-Fos mice (TRAP2:

Fostm2.1[(icre/ERT2)Luo/J]; Jackson Laboratories #030323) were back-crossed with C57BL6/N and

were either bred with reporter line tdT-Ai14 mice (B6;129S6-Gt(ROSA)[26Sortm14(CAG-tdTomato)Hze/J];

Jackson Laboratories # 007908) to create TRAP2-tdT mice or reporter line Chr2-YFP (B6;129S-

Gt(ROSA)[26Sortm32(CAG-COP4*H134R/EYFP)Hze/J]; Jackson Laboratories #12 569) to create TRAP2-

ChR2/eYFP mice. Male and female TRAP2-tdT and TRAP2-ChR2/eYFP mice four to eight

weeks old were used in experiments. Mice were housed with littermates (up to 5 mice per cage)

in a 12/12 h light/dark cycle. Food and water were provided ad libitum.

Behavioral training and engram labeling

Male and female experimental mice were trained in a spatial learning Barnes maze (BM)

task or placed in an enriched environment (EE) for 3 hours followed by tamoxifen induction to

induce Cre recombinase as detailed below. Since we observed TRAP2 mice exhibiting

considerable litter to litter variability in tdT labeling following identical treatments in preliminary

studies (not shown), we used littermate pairs for the following studies: ***Barnes maze (BM)***: 4-6

week old male and female TRAP2-tdT mice were trained in a spatial memory task on a Barnes

maze table (Maze Engineers, https://conductscience.com/maze/), 92cm in diameter with 20

holes (5 cm diameter each). One hole was equipped with a false floor installed with a removable

escape box that could be traded out for an additional false floor piece. The maze was set up in

the middle of 4 curtain walls with two bright lights and a camera for recording above the maze.

Different sets of visual cues (various shapes cut from felt) were attached to the curtain for

spatial orientation. The escape hole was positioned in between two visual cues. Animals were

held in their home cage outside of the curtain in a dark room until their turn to run the trial. We

observed that these mice were hyperactive, therefore mice were housed individually on the day

before training (Day 0). Mice were habituated to the behavior room in their home cages for at

least 1 hour before training on Day 1, habituated to the arena by placing them in the starter cup

on the table for 1 minute, and guided by gently moving the starter cup to the escape box (in a

temporary location different from the experimental location). <u>During task acquisition training on</u>

days 1-6, mice performed three 180-second trials during which the mouse explored the maze to

find the escape box. The three trials were separated by a minimum of 15 minute inter-trial-

intervals. If mice failed to locate the escape box at the end of the 180 seconds, the experimenter

guided them to the escape box and then placed them back into their home cage. On Day 6 of

BM acquisition, mice were brought to the room 5 hours before testing and received 4-hydroxy

tamoxifen (4-OHT, 50mg/kg i.p.) 15 minutes prior to the first acquisition session. 4-OHT was

prepared as described previously (DeNardo et al., 2019). Briefly, 4-OHT was dissolved in 100%

ethanol at a concentration of 20mg/mL by sonicating solution at 37°C for 30 minutes or until

dissolved, aliquoted and stored at -20°C. On the day of injection, 4-OHT was redissolved by

sonicating solution at 37°C for 10 minutes. A 1:4 mixture of castor oil and sunflower seed oil,

respectively, was added for a final concentration of 10mg/mL. The remaining ethanol in solution

was evaporated by speed vacuuming in a centrifuge (DeNardo et al., 2019). Behavior in the

Barnes maze paradigm was analyzed using Anymaze software, a blinded experimenter, and

BUNS analysis software (Illouz et al., 2016).

Enriched environment (EE): Experimental TRAP2-tdT and TRAP2-ChR2/eYFP mice

were housed in an enriched environment consisting of an oversized cage filled with multiple

tunnels, extra nestlets, a metal swing, and a few huts for the animals to interact with for 3 hours.

Mice received 4-OHT (50mg/kg i.p) 90 minutes into their 3 hours of enrichment. Animals were

left in the room for an additional 5 hours to limit neuronal activity labeling not related to the

behavioral paradigm. In a subset of experiments (Fig. 1), 7 days following 4-OHT induction,

littermate cohorts of TRAP2-tdT mice that underwent BM acquisition or EE exposure were

placed with their respective pair into the EE for two hours and then immediately sacrificed by

perfusion with 4% paraformaldehyde (PFA) upon removal from the EE. TRAP2-ChR2/eYFP

mice induced after EE exposure were sacrificed a week later for electrophysiology (Fig. 2-5;

Supplemental Fig. 2-4).

Immunohistochemistry and Cell Morphology

TRAP2-tdT mice, 90 minutes following EE exposure, were transcardially perfused with

PBS followed by a 4% PFA while under euthasol anesthesia. The brains were held in the 4%

PFA at 4°C for 3 hours before being transferred to PBS. Coronal brain sections (50 μm) were

obtained using a Leica vt100s vibratome and 5 sequential sections, each 250 μm apart across

the septotemporal axis, were immuno-stained for c-Fos and analyzed for quantification. Free

floating sections were blocked in 10% goat serum in PBS with 0.3% Triton X-100 for 1 hour.

Sections were incubated in 4°C overnight in primary antibody for c-Fos (1:750, Rabbit mAb Cell

Signaling Technology, cat #2250). The following day, sections were incubated in goat anti-rabbit

Alexa Fluor® 488 secondary antibody (1:500 Abcam, cat #150077) for 1 hour.

Slices from TRAP2-ChR2/eYFP mice that were used in electrophysiological studies were fixed

in 0.1mM phosphate buffer containing 4% PFA at 4°C overnight. Slices were washed with PBS

and then incubated in 10% goat serum with 0.3% Triton X-100 for 1 hour at room temperature.

Sections were incubated in 4°C overnight in primary antibody for GFP (1:500 Anti-Green

Fluorescent Protein Antibody Aves Labs, AB_2307313). The following day, sections were

incubated in goat anti-chicken Alexa Fluor® 488 secondary antibody (1:500 Abcam cat#

150169) and Alexa Fluor® 594 conjugated streptavidin (1:1000 Thermo Fisher, S11227) in PBS

with 0.3% Triton X-100 for 2 hours at room temperature.

Slices were mounted on a glass slide using Vectashield®. Sections were imaged using a

Zeiss Axioscope-5 with stereo investigator (MBF Bioscience) for analysis. Cell counts, cell type

classification, and evaluation of double labeling were conducted by an experimenter blinded to

treatments. Cells with compact dendritic arbors and somata with greater length than width were

classified as GCs and those with wide dendritic angle, 2 or more primary dendrites, and greater

somatic width than height were classified as SGCs (Gupta et al., 2020; Afrasiabi et al., 2022).

Slice Physiology

Seven to nine days after tamoxifen induction following EE exposure, TRAP2-

ChR2/eYFP mice were euthanized under isoflurane anesthesia for preparation of horizontal

brain slices (350µm) using a Leica VT1200S Vibratome in ice cold sucrose artificial

cerebrospinal fluid (sucrose-aCSF) containing (in mM): 85 NaCl, 75 sucrose, 24 $NaHCO_3$, 25

glucose, 4 $MgCl_2$, 2.5 KCl, 1.25 NaH_2PO_4, and 0.5 CaCl. Slices were bisected and incubated at

32°C for 30 min in a holding chamber containing an equal volume of sucrose-aCSF and

recording aCSF and were subsequently held at room temperature for an additional 30 min

before use. The recording aCSF contained (in mM): 126 NaCl, 2.5 KCl, 2 $CaCl_2$, 2 $MgCl_2$, 1.25

NaH_2PO4, 26 $NaHCO_3$, and 10 D-glucose. All solutions were saturated with 95% O_2 and 5%

CO_2 and maintained at a pH of 7.4 for 2-6 hours (Gupta et al., 2012; Yu et al., 2015; Afrasiabi et

al., 2022). Slices were transferred to a submerged recording chamber and perfused with

oxygenated aCSF at 33°C. Whole-cell voltage-clamp and current-clamp recordings from GCs in

the granule cell layer and presumed SGCs in the inner molecular layer or edge of the granule

cell layer were performed under IR-DIC visualization with Nikon Eclipse FN-1 (Nikon

Corporation) using 40x water immersion objective. Recordings were obtained using axon

instruments MultiClamp 700B amplifier (Molecular Devices). Data were low pass filtered at

2kHz, digitized using Axon DigiData 1400A (Molecular Devices), and acquired using pClamp11

at 10kHz sampling frequency. Recordings were obtained using borosilicate glass

microelectrodes (3-7MΩ), pulled using Narishige PC-10 puller (Narishige Japan). Recordings

were performed using K-gluconate based internal solution (K-gluc) containing 126 mM K-

gluconate, 4 mM KCl, 10 mM HEPES, 4 mM Mg-ATP, 0.3 mM Na-GTP, and10 mM PO-

creatinine or cesium methane sulfonate (CsMeSO$_4$) internal solution containing 140 mM cesium

methane sulfonate, 10 mM HEPES, 5 mM NaCl, 0.2 mM EGTA, 2 mM Mg-ATP, and 0.2 mM

Na-GTP. (pH 7.25; 270–290 mOsm). Biocytin (0.2%) was included in the internal solution for

post hoc cell identification (Yu et al., 2015; Afrasiabi et al., 2022; Gupta et al., 2022). Cells

labeled with eYFP were visualized under epifluorescence and patched under IR-DIC using

pipettes filled with K-gluc internal and held at -70mV in current clamp. 10 ms, 10Hz pulses of

blue light (λ=470 nm 0.9 mW) was used to optically evoke firing or inward currents were used to

confirm ChR2/eYFP labeling. Responses to 1s positive and negative current injections,

beginning at -200 pA with 40 pA steps up to 20 sweeps, were examined to determine active and

passive characteristics. Dual patch clamp recordings were obtained from pairs of labeled and

unlabeled neurons. Unlabeled neurons were recorded using microelectrodes with CsMeSO$_4$

internal and held at 0mV (glutamate reversal potential) to isolate inhibitory postsynaptic currents

(IPSCs) and -70mV (close to GABA reversal potential) to record excitatory postsynaptic currents

(EPSCs). Labeled neurons, held in current clamp, were depolarized by 10ms 500pA pulses at

50Hz to elicit action potentials to measure evoked responses in labeled or unlabeled cells held

in voltage clamp. Labeled neuron pairs were tested for connectivity in both directions.

Spontaneous EPSCs were (sEPSCs) were recorded in both labeled and unlabeled cells were

held at -70mV in voltage clamp for 5-10 minutes. Recordings were discontinued if series

resistance increased by > 20% or if access resistance surpassed 25MΩ. Post hoc biocytin

immunostaining and morphologic analysis was used to definitively identify SGCs and GCs

included in this study.

Data Analysis

Active and passive properties were analyzed using EasyElectrophysiology v2.6.3 (Easy

Electrophysiology Ltd). Action potential, threshold, amplitude, halfwidth, and first spike latency

were acquired from the first sweep in which the cell fired. Fast and medium

afterhyperpolarization (AHP) and spike frequency adaptation were determined based on voltage

response and firing in response to a 120 pA current injection. Action potential (AP) threshold
was calculated using the first derivative method. Amplitude was calculated by the AP peak value
minus baseline. Spike frequency accommodation was calculated using the divisor method, in
which the inter spike interval (ISI) of the first two APs is divided by the ISI of the last two APs.
First spike latency is the time from the start of a current pulse to the first AP.

Action potential and synaptic potential analysis were conducted using
EasyElectrophysiology v2.6.3 (Easy Electrophysiology Ltd). Action potential kinetics were
analyzed with a 200kHz interpolation for rise-time, decay-time, and halfwidth. Decay was
measured using a biexponential decay curve fit with a cut off of 10-90% of the AP amplitude.
Rise time is calculated between 10-90% of the AP amplitude. Half width was calculated as the
time between the two half-amplitude samples. Afterhyperpolarization values are calculated as
baseline minus fAHP or mAHP. The value is the minimum point within a search region specified
as 0-3ms for fAHP and 10-50ms for mAHP. Spontaneous EPSCs were detected and analyzed
using EasyElectrophysiology threshold search algorithm and events were confirmed by the
experimenter. Any "noise" that spuriously met trigger specifications was rejected.

Temporal Correlation of sEPSCs: Temporal correlation of sEPSCs in dual recording
sessions from cell pairs (labeled to labeled (L-L) and labeled to unlabeled (L-U)) was defined by
a session-wise cross correlation profile (CCP) of temporally binned data for select detection
windows (MATLAB *xcorr*, Wiegand & Cowen). Temporal correlations were tested using
detection windows and bin sizes were always divided into 21 equally sized bins in the window: a
±100 ms detection window with 10 ms bins (200 ms window, 21 bins aligned to sEPSC) and a
±50 ms detection window with 5 ms bins (100 ms window, 21 bins aligned to sEPSC). Temporal
correlation was determined if the CCP exceeded a 2 standard deviation (SD) threshold above
the total mean correlation (0.15 for the 100 ms detection window and 0.10 for the 50 ms
detection window). Within each detection window, "peri-occurrence" was defined by the CCP
maximum outside the center bin while "cooccurrence" was defined by the correlation in the

center bin. CCP of sEPSC event times in L-L and L-U pairs were compared with randomly

jittered event times from the same data set to identify intrinsic correlations within the data.

Session jitter was pseudo-randomly selected from an event timeline matrix (assigned to one cell

from each paired cell recording session) bound by ± 0.5 s across 100 iterations. The temporally

jittered correlation data were then compared to the temporally aligned CCP and correlation data

in the center bin (Wiegand et al., 2016). The ability of the sEPSC event cooccurrence to predict

L-L versus L-U pairs was computed by plotting the receiver operating characteristic (ROC)

curve and calculating the area under the curve (AUC) in both groups. Correlation values in the

center bin of the CCPs were used to generate histograms (correlation bins from 0 to 0.5 with

correlation bin widths of 0.001), which were reverse integrated to evaluate the cumulative sum

between categorized L-L sessions (true, n=7) and L-U sessions (false, n=8) rates. Cumulative

sums were used to find the total AUROC as the classification performance measure using

MATLAB (*hist counts*, *cumsum*, *flip*, and *trapz* functions), where 50% AUROC performance

would classify L-L versus L-U by random chance.

Sample sizes were not predetermined and conformed with those employed in the field.

Significance was set to $p < 0.05$, subject to appropriate Bonferroni correction. Statistical analysis

was performed using GraphPad Prism 10 and MATLAB. Unpaired Kolmogorov-Smirnov (KS)

Test, unpaired Mann-Whitney, one-way ANOVA, two-way ANOVA (TW-ANOVA), two-way

repeated measures ANOVA (TW-RM-ANOVA), or Kruskal-Wallis followed by post-hoc pairwise

multiple comparisons using Holm-Sidak method or Dunn's method used as appropriate.

Statistical data is reported as mean ± SEM.

654
655

References

Afrasiabi M, Gupta A, Xu H, Swietek B, Santhakumar V (2022) Differential Activity-Dependent Increase in Synaptic Inhibition and Parvalbumin Interneuron Recruitment in Dentate Granule Cells and Semilunar Granule Cells. Journal of Neuroscience 42:1090-1103.

Amaral DG, Scharfman HE, Lavenex P (2007) The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). Progress in Brain Research 163:3-22.

Borzello M, Ramirez S, Treves A, Lee I, Scharfman H, Stark C, Knierim JJ, Rangel LM (2023) Assessments of dentate gyrus function: discoveries and debates. Nature Reviews Neuroscience 24:502-517.

Cayco-Gajic NA, Silver RA (2019) Re-evaluating Circuit Mechanisms Underlying Pattern Separation. Neuron 101:584-602.

Cutts CS, Eglen XSJ (2014) Detecting pairwise correlations in spike trains: An objective comparison of methods and application to the study of retinal waves. Journal of Neuroscience 34:14288-14303.

Danieli K, Guyon A, Bethus I (2023) Episodic Memory formation: A review of complex Hippocampus input pathways. Progress in Neuro-Psychopharmacology and Biological Psychiatry 126:110757-110757.

DeNardo LA, Liu CD, Allen WE, Adams EL, Friedmann D, Fu L, Guenthner CJ, Tessier-Lavigne M, Luo L (2019) Temporal evolution of cortical ensembles promoting remote memory retrieval. Nature Neuroscience 22:460-469.

Erwin SR, Sun W, Copeland M, Lindo S, Spruston N, Cembrowski MS (2020) A Sparse, Spatially Biased Subtype of Mature Granule Cell Dominates Recruitment in Hippocampal-Associated Behaviors. Cell Reports 31:107551-107551.

Espinoza C, Guzman SJ, Zhang X, Jonas P (2018) Parvalbumin + interneurons obey unique connectivity rules and establish a powerful lateral-inhibition microcircuit in dentate gyrus. Nature Communications 9:1-10.

Gawel K, Gibula E, Marszalek-Grabska M, Filarowska J, Kotlinska JH (2019) Assessment of spatial learning and memory in the Barnes maze task in rodents—methodological consideration. Naunyn-Schmiedeberg's Archives of Pharmacology 392:1-18.

Gouty-Colomer LA, Hosseini B, Marcelo IM, Schreiber J, Slump DE, Yamaguchi S, Houweling AR, Jaarsma D, Elgersma Y, Kushner SA (2016) Arc expression identifies the lateral amygdala fear memory trace. Molecular Psychiatry 21:364-375.

Guenthner CJ, Miyamichi K, Yang HH, Heller HC, Luo L (2013) Permanent genetic access to transiently active neurons via TRAP: Targeted recombination in active populations. Neuron 78:773-784.

Gupta A, Elgammal FS, Proddutur A, Shah S, Santhakumar V (2012) Decrease in tonic inhibition contributes to increase in dentate semilunar granule cell excitability after brain injury. Journal of Neuroscience 32:2523-2537.

Gupta A, Dovek L, Proddutur A, Elgammal FS, Santhakumar V (2022) Long-Term Effects of Moderate Concussive Brain Injury During Adolescence on Synaptic and Tonic GABA Currents in Dentate Granule Cells and Semilunar Granule Cells. Frontiers in Neuroscience 16:1-10.

Gupta A, Proddutur A, Chang YJ, Raturi V, Guevarra J, Shah Y, Elgammal FS, Santhakumar V (2020) Dendritic morphology and inhibitory regulation distinguish dentate semilunar granule cells from granule cells through distinct stages of postnatal development. Brain Structure and Function.

Guzman SJ, Schlögl A, Espinoza C, Zhang X, Suter BA, Jonas P (2021) How connectivity rules and synaptic properties shape the efficacy of pattern separation in the entorhinal cortex–dentate gyrus–CA3 network. Nature Computational Science 1:830-842.

Hainmueller T, Bartos M (2018) Parallel emergence of stable and dynamic memory engrams in the hippocampus. Nature 558:292-296.

Hainmueller T, Bartos M (2020) Dentate gyrus circuits for encoding, retrieval and discrimination of episodic memories. Nature Reviews Neuroscience 21:153-168.

Han JH, Kushner SA, Yiu AP, Cole CJ, Matynia A, Brown RA, Neve RL, Guzowski JF, Silva AJ, Josselyn SA (2007) Neuronal competition and selection during memory formation. Science 316:457-460.

Heroux NA, Osborne BF, Miller LA, Kawan M, Buban KN, Rosen JB, Stanton ME (2018) Differential expression of the immediate early genes c-Fos, Arc, Egr-1, and Npas4 during long-term memory formation in the context preexposure facilitation effect (CPFE). Neurobiology of Learning and Memory 147:128-138.

Hunsaker MR, Rosenberg JS, Kesner RP (2008) The role of the dentate gyrus, CA3a,b, and CA3c for detecting spatial and environmental novelty. Hippocampus 18:1064-1073.

Illouz T, Madar R, Clague C, Griffioen KJ, Louzoun Y, Okun E (2016) Unbiased classification of spatial strategies in the Barnes maze. Bioinformatics 32:3314-3320.

Josselyn SA, Tonegawa S (2020) Memory engrams: Recalling the past and imagining the future. Science 367.

Josselyn SA, Köhler S, Frankland PW (2015) Finding the engram. Nature Reviews Neuroscience 16:521-534.

Kawashima T, Okuno H, Bito H (2014) A new era for functional labeling of neurons: Activity-dependent promoters have come of age. Frontiers in Neural Circuits 8:1-9.

Kee N, Teixeira CM, Wang AH, Frankland PW (2007) Preferential incorporation of adult-generated granule cells into spatial memory networks in the dentate gyrus. Nature Neuroscience 10:355-362.

Kitamura T, Ogawa SK, Roy DS, Okuyama T, Morrissey MD, Smith LM, Redondo RL, Tonegawa S (2017) Engrams and circuits crucial for systems consolidation of a memory. Science 356:73-78.

Larimer P, Strowbridge B (2010) Representing Information in neuronal cell assemblies: Persistent activity in the dentate gyrus mediated by semilunar granule cells. Physiology & behavior 176:139-148.

Liu X, Ramirez S, Tonegawa S (2014) Inception of a false memory by optogenetic manipulation of a hippocampal memory engram. Philosophical Transactions of the Royal Society B: Biological Sciences 369.

Liu X, Ramirez S, Pang PT, Puryear CB, Govindarajan A, Deisseroth K, Tonegawa S (2012) Optogenetic stimulation of a hippocampal engram activates fear memory recall. Nature 484:381-385.

Mazurkiewicz M, Kambham A, Pace B, Skwarzynska D, Wagley P, Burnsed J (2022) Neuronal activity mapping during exploration of a novel environment. Brain Research 1776:147748-147748.

McHugh TJ, Jones MW, Quinn JJ, Balthasar N, Coppari R, Elmquist JK, Lowell BB, Fanselow MS, Wilson MA, Tonegawa S (2007) Dentate gyrus NMDA receptors mediate rapid pattern separation in the hippocampal network. Science 317:94-99.

Morrison DJ, Rashid AJ, Yiu AP, Yan C, Frankland PW, Josselyn SA (2016) Parvalbumin interneurons constrain the size of the lateral amygdala engram. Neurobiology of Learning and Memory 135:91-99.

Rao-Ruiz P, Visser E, Mitrić M, Smit AB, van den Oever MC (2021) A Synaptic Framework for the Persistence of Memory Engrams. Frontiers in Synaptic Neuroscience 13:1-15.

Rovira-Esteban L, Hájos N, Nagy GA, Crespo C, Nacher J, Varea E, Blasco-Ibáñez JM (2020) Semilunar granule cells are the primary source of the perisomatic excitatory innervation onto parvalbumin-expressing interneurons in the dentate gyrus. eNeuro 7:1-17.

757   Roy DS, Muralidhar S, Smith LM, Tonegawa S (2017) Silent memory engrams as the basis for
758   retrograde amnesia. Proceedings of the National Academy of Sciences of the United
759   States of America 114:E9972-E9979.
760   Ryan TJ, Roy DS, Pignatelli M, Arons A, Tonegawa S (2015) Engram cells retain memory under
761   retrograde amnesia. Science 348:1007-1013.
762   Sano Y, Shobe JL, Zhou M, Huang S, Shuman T, Cai DJ, Golshani P, Kamata M, Silva AJ
763   (2014) CREB Regulates Memory Allocation in the Insular Cortex. Current Biology
764   24:2833-2837.
765   Save L, Baude A, Cossart R (2019) Temporal Embryonic Origin Critically Determines Cellular
766   Physiology in the Dentate Gyrus. Cerebral Cortex 29:2639-2652.
767   Semon RW (1909) Die nmemischen empfindungen. [Mnemic psychology]. New York, NY:
768   Leipzig: Wilhelm Engelmann.
769   Stefanelli T, Bertollini C, Lüscher C, Muller D, Mendez P (2016) Hippocampal Somatostatin
770   Interneurons Control the Size of Neuronal Memory Ensembles. Neuron 89:1074-1085.
771   van Groen T, Miettinen P, Kadish I (2003) The entorhinal cortex of the mouse: Organization of
772   the projection to the hippocampal formation. Hippocampus 13:133-149.
773   Walker MC, Pavlov I, Kullmann DM (2010) A 'sustain pedal' in the hippocampus? Nature
774   Neuroscience 13:146-148.
775   Wiegand JPL, Gray DT, Schimanski LA, Lipa P, Barnes CA, Cowen SL (2016) Age is
776   associated with reduced sharp-wave ripple frequency and altered patterns of neuronal
777   variability. Journal of Neuroscience 36:5650-5660.
778   Williams PA, Larimer P, Gao Y, Strowbridge BW (2007) Semilunar granule cells: Glutamatergic
779   neurons in the rat dentate gyrus with axon collaterals in the inner molecular layer.
780   Journal of Neuroscience 27:13756-13761.
781   Yiu AP, Mercaldo V, Yan C, Richards B, Rashid AJ, Hsiang HLL, Pressey J, Mahadevan V,
782   Tran MM, Kushner SA, Woodin MA, Frankland PW, Josselyn SA (2014) Neurons Are
783   Recruited to a Memory Trace Based on Relative Neuronal Excitability Immediately
784   before Training. Neuron 83:722-735.
785   Yu J, Swietek B, Proddutur A, Santhakumar V (2015) Dentate total molecular layer interneurons
786   mediate cannabinoid-sensitive inhibition. Hippocampus 25:884-889.
787   Zhou Y, Won J, Karlsson MG, Zhou M, Rogerson T, Balaji J, Neve R, Poirazi P, Silva AJ (2009)
788   CREB regulates excitability and the allocation of memory to subsets of neurons in the
789   amygdala. Nature Neuroscience 12:1438-1443.
790
791

    Figure Legends

Figure 1: Task associated DG labeled neurons show consistent activation of SGCs and

paradigm specific reactivation A-B) Schematic of experimental timeline for animals trained in

the Barnes maze (BM) task followed by exposure to enriched environment (EE), the BM-EE

cohort (BM) group (A) and mice housed in EE followed by reintroduction of EE, the EE-EE

cohort (EE) group (B). C-D) Representative epifluorescence image of a section from mice one

week after induction of tdT labeling (Ci, Di) following BM testing (C) or EE testing (D) and c-Fos

immunostaining (Cii, Dii) following subsequent EE exposure. E-F) Quantification of number of

tdT labeled cells per slice (E) and summary of proportion of tdT labeled cells in the upper blade

of the DG per slice (F). G)Summary of proportion of tdT cells co-labeled with c-Fos (green). H)

Representative TRAP-tdT section showing distinct SGC morphology (white arrowhead). I) Plot

of % of tdT cells that had morphology consistent with SGCs. * indicates $p<0.05$, *** indicates

$p=0.0003$ by Nested t-test, n=4 subjects/treatment. Schematics were generated using

BioRender under license.

Figure 2: Intrinsic differences in frequency adaptation distinguish labeled SGCs. A-B)

Representative images of a biocytin filled GC (A) with a narrow dendritic arbor and a smaller

somatic width and a SGC (B) with wide dendritic span, greater somatic width than height, and

axonal projections throughout the molecular and granule cell layer (arrowheads). Maximum

intensity projections of confocal image stacks are presented as gray scale, inverted images. C-

D) Summary plots of resting membrane potential (RMP in C) and input resistance (R_{in} in D)

between labeled and unlabeled GCs and SGCs. # indicates $p<0.05$ for main factor cell type by

TW-ANOVA and * indicates $p<0.05$ for labeled versus unlabeled within cell type by Šídák's

multiple comparisons post hoc test in n=11-19 cells/group. E-F) Representative cell membrane

voltage traces in response to +120 and -200pA current injections (E) and +400pA current

injection (F) in a GC (top) and SGC (bottom). G) Summary plot of firing frequency in response to

increasing current injections in labeled and unlabeled SGCs and GCs. #### indicates $p < 0.0001$

for main factor cell type by 3Way ANOVA n=9-22 cells/group. H-J) Summary plots of firing

frequency at 520pA compared to max frequency (H), spike amplitude attenuation calculated as

ratio between the amplitude of the 15th spike and 1st spike at a current injection of 400pA (I)

and spike frequency adaptation (J). # indicates $p < 0.05$, ## $p < 0.01$ for main factor cell type by

TW-ANOVA and ** i ndicates $p < 0.01$ for labeled versus unlabeled within cell type by Šídák's

multiple comparisons post hoc test in n=8-19 cells/group.

Figure 3: Tagged DG neurons do not support mutual excitatory drive. A) Schematic

showing dual patch clamp recording from labeled (green) GC-SGC pair. B) Summary

breakdown of cell-type specific connections tested in dual recordings from labeled neurons. C)

Representative maximum intensity projection of a confocal image stack of a pair of biocytin filled

SGC (left) and GC (right). Images are gray scale and inverted and are overexposed to

emphasize the intact axonal arbors in the recorded pair. D) Presence of spontaneous EPSCs in

the SGC-GC pair in E-G to verify the presence of excitatory inputs and a healthy circuit. E) Light

evoked inward currents validate expression of ChR2 in labeled cell pair. F) Representative

traces from a labeled SGC and labeled GC show that depolarization induced firing in SGC (top)

failed to evoke EPSCs in a GC (bottom) recorded in voltage clamp. Individual traces are in gray

with average trace overlaid in black. G) Depolarization induced firing in GC (bottom) fails to

evoke EPSCs in a SGC recorded in voltage clamp (top).

Figure 4: Evidence for DG engram neurons supporting sparse feedback inhibition onto

non-engram neurons. A-C) Representative confocal image of eYFP labeled neurons in a

TRAP-ChR2-eYFP mouse (A), shows biocytin staining (B) in a pair of recorded Labeled-SGC

and Unlabeled-GC. Note co-labeling for eYFP and biocytin in the SGC, while the GC does not

colocalize eYFP (C) D) Summary of cell-type specific connections tested in dual recordings from

labeled and unlabeled neurons. Inset depicts a schematic showing dual patch clamp recording

from a labeled (green) SGC and an unlabeled (blue) GC pair. E) Light evoked currents validate

the expression of ChR2 in the Labeled-SGC and lack of response in the Unlabeled-GC. F-G)

Representative traces from a Labeled-SGC and an Unlabeled GC show that depolarization

induced firing in the Labeled-SGC (top) failed to evoke EPSCs (F) and IPSCs (G) in the

Unlabeled-GC. H) Schematic of recording configuration illustrated wide field optical illumination

with labeled neurons (green), unlabeled neurons (blue), and local circuit interneuron (yellow). I)

Example traces from a recording in which wide field optical stimulation evoked inhibitory

responses in the Unlabeled-GC and firing in the Labeled-SGC. Note that the SGC firing by

depolarization in the absence of light failed to elicit IPSCs in the same GC. J-K) Schematic with

Labeled-GC (green), Unlabeled-SGC (blue), and local circuit interneuron (yellow) (J) and traces

from a recorded pair where depolarization of a Labeled-GC elicited inhibitory responses in an

Unlabeled-SGC (K).

Figure 5: Correlated Spontaneous Excitatory Inputs to Labeled Pairs. A) Representative

confocal image of eYFP labeled and biocytin stained neurons in a TRAP-ChR2-eYFP mouse. B)

Schematic for Labeled-Labeled (L-L) dual recordings with representative example of

spontaneous EPSCs (sEPSCs) in an L-L pair below. C) Schematic for Labeled-Unlabeled (L-U)

dual recordings with representative example of sEPSCs in an L-U pair below. D) Schematic for

session-wise cross correlation profiles (CCPs) defined by correlations exceeding a 2 standard

deviation (SD) threshold above the total mean correlation: EPSC peri-occurrence was tested as

event time CCP exceeding threshold within full detection window; cooccurrence was defined as

event time CCP exceeding threshold within center bin of detection window. E) CCP from

recordings from L-L pairs analyzed with ±100 ms detection window (bright blue, n=7). Overlaid

jittered data (black) was developed by appending the event timing of one cell with a randomized

lead/lag of +/- 0.5 s for 100 iterations (Top panel). Inset: Plot of maximum correlations (Rmax)

in relation to the dashed line representing 2xSD = 0.15. CCP in recordings from L-U pairs

analyzed with ±100 ms detection window (dark blue, n=8). Corresponding jittered data,

developed as detailed above, is overlaid in black (Bottom panel). Inset: Plot of Rmax in relation

to the dashed line representing 2xSD = 0.15. F) CCP from sessions with recordings from L-L

pairs analyzed with ±50 ms detection window from L-L pairs (bright purple, n=7) with jittered

data developed as detailed above is overlaid in black (Top panel). Inset: Plot of Rmax in relation

to the dashed line representing 2xSD = 0.10). CCP from recordings in L-U pairs analyzed with

±50 ms detection window (dark purple, n=8) with corresponding jittered data overlaid in black

(Bottom panel). Inset: Rmax in relation to the dashed line representing 2xSD = 0.10. G)

Comparison of center bin correlation between L-L versus L-U pairs in aligned (Align-recorded)

versus jittered (Jitter-simulated) data, analyzed using ±100 ms detection window (left, colors as

in E) and using ±50 ms detection window (right, colors as in F). H) Center bin classifier

performance (solid line) compared to chance performance (dashed line, colors as in E and F

respectively) plotted as area under the receiver-operating characteristic (ROC) curve (AUROC)

between L-L (true positive rate) and L-U (false positive rate) for analysis using ±100 ms

detection window (left panel) and for analysis using ±50 ms detection window (right panel). Data

presented as mean±SEM (dual recording sessions), * indicate $p < 0.05$; ** indicates $p < 0.01$ ***

indicates $p < 0.001$, **** indicates $p < 0.0001$; TW-ANOVA with Šídák's multiple comparisons post

hoc tests.

A)

B) Pre to Post Connections Tested in Labeled-Labeled Pairs

- Labeled GC to Labeled GC n=2
- Labeled SGC to Labeled GC n=7
- Labeled GC to Labeled SGC n=7
- Labeled SGC to Labeled SGC n=16

C) 100 μm

D) Labeled-SGC

Labeled-GC

20 pA
200 ms

E) Labeled-SGC

100 pA
200 ms

Labeled-GC

100pA
200 ms

F) Labeled-SGC

50 mV
100 ms

Labeled-GC

Holding at -70

50 pA
100 ms

G) Labeled-SGC

Holding at -70

50 pA
100 ms

Labeled-GC

50 mV
100 ms

A) TRAP::ChR2-eYFP
B) Biocytin
C) 100 μm

D) Pre to Post Connections Tested in Labeled-Unlabeled Pairs
- Labeled SGC to Unlabeled GC n=49
- Labeled GC to Unlabeled GC n=6
- Labeled GC to Unlabeled SGC n=4
- Labeled SGC to Unlabeled SGC n=4

E) Labeled-SGC
100pA
200 ms

Unlabeled-GC
100pA
200 ms

F) Labeled-SGC
50 mV
100 ms

Unlabeled-GC
Holding at -70
50pA
100 ms

G) Labeled-SGC
50 mV
100 ms

Unlabeled-GC
Holding at 0
50pA
100 ms

H) Molecular Layer / SGC / SGC / GC / GC / Hilus / IN

I) Labeled-SGC
100 ms

Unlabeled-GC
100 ms

J) Molecular Layer / Granule Cell Layer / Hilus / GC / SGC / IN

K) Labeled-GC
50 mV
100 ms

Unlabeled-SGC
50pA
100 ms