MLL1 directs gut-associated antibody responses to helminth and bacterial infections

Yan Zhang[a,b,1], Clarissa Chakma[a,b,1], Alana Kirn[a,b], Aidil Zaini[a,b,c], Rohia Farighi[a,b,c], Liang Xie[b,d,†],

Diana López-Ureña[d,e], Steven Mileto[d,e], Dena Lyras[d,e], Colby Zaph[a,b], Joanna R Groom[f,g], and Kim

L Good-Jacobson[a,b,*].

[a]Department of Biochemistry and Molecular Biology, Monash University, Clayton, Victoria, Australia 3800.

[b]Immunity Program, Biomedicine Discovery Institute, Monash University, Clayton, Victoria, Australia 3800.

[c]Department of Immunology, Alfred Medical Research and Education Precinct, Monash University, Melbourne, VIC, Australia.

[d]Department of Microbiology, Monash University, Clayton, Victoria, Australia 3800.

[e]Infection Program, Biomedicine Discovery Institute, Monash University, Clayton, Victoria, Australia 3800.

[f]Division of Immunology, Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria, Australia.

[g]Department of Medical Biology, University of Melbourne, Parkville, Victoria, Australia.

[†]Present address: Precision Medicine Translational Research Programme, Department of Obstetrics & Gynaecology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore.

[1]Equal contribution

*Correspondence: Kim Good-Jacobson

Email: kim.jacobson@monash.edu

Author contributions: KLG-J conceived the study; KLG-J and YZ designed research; YZ, CC, AK, AZ and RF performed research; KLG-J, CC, AK and YZ analyzed data; CZ, DL and JRG provided intellectual input; CZ, LX, DLU, SM and JRG provided reagents and technical expertise; and KLG-J, YZ and CC wrote the manuscript.

Competing Interest Statement: The authors declare no competing interests.

Keywords: B cells, IgA, helminth, trichuris muris, epigenetics, gut-associated lymphoid tissue, MLL1.

Abstract

Soil-transmitted helminths are one of the most common infections across the world, yet how to promote an effective humoral response to helminths is not well understood. To ensure the most effective response, molecular programs underpin the specialization of antibody effector subtype to microenvironmental signals. Here, we identify the methyltransferase mixed-lineage leukemia 1 (MLL1) as a key target to promote IgA-driven responses. *Mll1* was increased in germinal center (GC) B cells in gut-associated lymphoid tissues, and the absence of MLL1 in mature B cells led to significant reductions in GCs, IgG1 and IgE in response to the helminth *Trichuris muris*. Yet, worm expulsion occurred more rapidly in $Mll1^{f/f}Cd23^{cre/+}$ mice compared to control mice in the first two weeks of infection. Accelerated worm clearance correlated with significantly elevated IgA^{+} plasma cells (PC), serum IgA and fecal IgA in in $Mll1^{f/f}Cd23^{cre/+}$ mice. Stimulation of B cells *in vitro* confirmed that *Mll1*-deficiency accelerated the formation of IgA^{+} PC and that this effect was B cell-intrinsic. CCR9 was identified as a key MLL1-regulated molecule by RNA-sequencing and flow cytometry. Correspondingly, $Mll1^{f/f}Cd23^{cre/+}$ mice infected with either *T. muris* or the bacterium *C. rodentium* significant increased IgA^{+}CCR9^{+} PC localized in the large intestine. This tailoring of the B cell response to promote IgA-secreting PC correlated to a healthier microbiome in $Mll1^{f/f}Cd23^{cre/+}$ mice, compared to control mice. Thus, this study reveals MLL1 as a key target to tailor B cell responses to helminth infections, which may aid the development of effective mucosal vaccines or new targeted treatments.

Significance statement

Diseases caused by infections, such as soil-transmitted helminth infections or bacterial infections, continue to pose a major human health problem worldwide. One key reason why soil-transmitted

helminths continue to cause long-term health and economic burdens is the lack of an effective

vaccine for human use. Antibody responses underpin most successful vaccine designs. Therefore,

a comprehensive understanding of how B cell responses are regulated to produce tailored antibody

to different pathogens is critical for the development of effective therapeutics and vaccines. This

study uncovered MLL1 as a novel therapeutic target to promote IgA-producing plasma cells,

opening new avenues of investigation into how effective and tailored antibody responses may be

produced in response to bacterial or helminth infection.

Main Text

Introduction

Soil-transmitted helminth infections infect approximately one-quarter of the world's population,

causing severe morbidity and death in high-incidence countries (1-5). Roundworm (*Ascaris*

lumbricoides), whipworm (*Trichuris trichiura*) and hookworms (*Necator americanus and*

Ancylostoma duodenale) are the most common soil-transmitted helminths that infect humans (6,

7). Currently, the most effective treatment for soil-transmitted helminth infections is to use mass

anti-helminthic drug administrations, such as albendazole (8). However, long-term mass anti-

helminthic drug therapy induces the development of drug-resistant parasites, and the treatment of

reinfection causes a significant economic burden (9-11). Therefore, successful long-term public

health interventions such as effective vaccines are urgently needed (12). B cell-mediated antibody

responses are the basis of most successful vaccines, yet there is a lack of understanding of how to

promote a potent humoral response to intestinal helminth infections.

Helminth-induced immune responses to whipworms such as *T. trichiura* mainly occur in the gut-

associated lymphoid tissues (GALT) and are largely dominated by Th2-biased responses (13).

Humoral responses in the GALT are highly specialized to maintain gut homeostasis by regulating

commensal microbiota and responses to food antigens, pathogenic bacteria or helminths (14). The

isotype of antibody produced during infection is critical for the effectiveness of the response (15).

While IgG, IgE and IgA are all produced during mucosal responses (8), IgA is particularly crucial

for the protection of the human intestinal mucosa (16-18). Increased IgA has been correlated to

reduced helminthiasis, increased microbiota diversity and improved responses to pathogenic

bacteria such as *Salmonella* Typhimurium (19-25).

Pinpointing key drivers of effective B cell responses in the GALT may aid development of effective mucosal vaccines and new treatments for helminth infections. MLL1 (also known as KMT2A) is a histone methyltransferase which promotes B cell development (26). It also can induce *Gata3*, the master transcriptional factor for Th2 differentiation (27), and regulates Muc2 production by intestinal stem cells, the major intestinal mucin that can provide protection against worms in the colonic epithelium (28). Recent work has revealed that during immune responses, histone modifiers have unique roles in regulating B cell subset formation, B cell migration, the quality and quantity of antibody, and immunopathology (29-35). Given the role of MLL1 in promoting Th2 cells through cooperation with c-Myb (36), a transcription factor we have previously found to be important in tailoring effective antibodies in response to a Th2-induced microenvironment (15), we hypothesized that MLL1 could be a key regulator of B cell responses in the GALT. Here, we find that conditional deletion of MLL1 in mature B cells promotes IgA responses to the helminth infection *Trichuris muris*. *Mll1*[f/f]*Cd23*[Cre/+] mice demonstrated accelerated worm clearance and anti-inflammatory microbiota. Compared with control mice, *Mll1*[f/f]*Cd23*[Cre/+] mice infected with either *T. muris* or the bacteria *Citrobacter rodentium* had significantly increased IgA production and IgA[+]CCR9[+] plasma cells (PC). These results identify a role for IgA in the protection against *T. muris* infection and identify MLL1 as a therapeutic target to promote the production of IgA[+] PC.

Results

MLL1 is required for optimal GC B cell responses

To investigate whether MLL1 may tailor mucosal B cell responses, we first examined the expression of *Mll1* in mature B cell subsets in GALT. B cell subsets were sort-purified from naïve wild-type mice and assessed for *Mll1* expression by RT-qPCR in isolated mesenteric lymph nodes (mLN) and Peyer's patches. At steady-state, there is an active immune response to microbiota in the gut-draining secondary lymphoid organs, in which activated B cells can form extrafollicular antibody-secreting PC, or germinal centers (GC) (2, 37). GCs provide a site for affinity maturation of B cells, enabling the production of long-lived memory B cells and PC, two populations that can persist for long periods of time to provide immunity (38). In comparison to naïve B cells and PC, GC B cells had significantly increased expression of *Mll1* (Fig. 1A-B). To determine whether this increase in *Mll1* in GC B cells was observed systemically, we immunized wild-type mice with (4-hydroxy-3-nitrophenyl)acetyl conjugated to keyhole limpet hemocyanin (NP-KLH), a haptenated protein precipitated in the adjuvant alum, and assessed *Mll1* in sort-purified subsets post-immunization. Commensurate with GALT GCs, *Mll1* was significantly upregulated in splenic GC B cells compared to follicular B cells (Fig. 1C). A small but non-significant increase in *Mll1* expression was also detected in memory B cells, compared to follicular B cells (Fig. 1C).

Given the increased expression of *Mll1* in GC B cells, we assessed whether MLL1 was required for GC formation. To conditionally delete *Mll1* in B cells, we crossed *Mll1* floxed mice to mice in which *Cre* is under the control of the *Cd23* promoter. Thus, upon Cre expression, *Mll1* would be expressed normally during B cell development, but conditionally deleted within mature B cells. To assess Th2 responses, we used the *T. muris* model of human *T. trichuria* whipworm infection.

T. muris is antigenically similar to *T. trichuria*, and both elicit a similar type of immune response,

making *T. muris* an effective model to study the B cell response to intestinal helminth infection

(13, 39, 40). *Mll1*$^{\text{f/f}}$*Cd23*$^{\text{cre/+}}$ and *Cd23*$^{\text{cre/+}}$ controls were infected with *T. muris* and B cell responses

assessed over time (Fig. 1D). The frequency of GC B cells was significantly decreased in

Mll1$^{\text{f/f}}$*Cd23*$^{\text{cre/+}}$ mice compared to control mice at each time point post-infection. A 1.5-fold, 4.5-

fold and 1.7-fold reduction in the number of GC B cells was observed in B cell-specific MLL1-

deleted mice compared to control mice on days (d)14, d28 and d35, respectively (Fig. 1E-G).

Histological analyses confirmed that GC structures were less frequent in mLN in the absence of

Mll1 (Fig. 1H).

To understand whether the role for MLL1 in GC biology was limited to GALT, or whether it was

generalizable to other types of B cell responses, *Mll1*$^{\text{f/f}}$*Cd23*$^{\text{cre/+}}$ and *Cd23*$^{\text{cre/+}}$ mice were

immunized with alum-precipitated NP-KLH and GC B cell formation assessed in the spleen (Fig.

I-K). Use of this model antigen allows the assessment of T-dependent B cell differentiation in the

absence of the pathogen-influenced changes to the tissue microenvironment that occurs upon

infection. In the absence of *Mll1*, GC B cells were virtually absent on d14 and d28 (Fig. I-K).

Therefore, MLL1 was required for effective GC B cell responses in response to infection or

immunization.

Mll1 deletion induces preferential production of IgA over IgG1 and IgE

B cell responses are important for reducing worm burden (37, 40). Thus, given the reduction of

GC B cells, we hypothesized that there would be a reduced ability of MLL1-deficient mice to

expel worms. To our surprise, worm clearance occurred more rapidly in *Mll1*$^{\text{f/f}}$*Cd23*$^{\text{cre/+}}$ mice,

compared to $Cd23^{cre/+}$ controls, with a >3-fold reduction on d14 post-*T. muris* infection (Fig. 2A).

Thus, conditional deletion of *Mll1* within mature B cells led to a more efficient and accelerated

worm expulsion during the early stages of *T. muris* infection.

B cells and helminth-specific antibodies can contribute to relieving worm burden (40). IgG1 and

IgE isotypes were assessed on 14 days post-infection to investigate whether accelerated worm

expulsion in $Mll1^{f/f}Cd23^{Cre/+}$ mice correlated to a change in the subclass dominating the response.

T. muris-specific IgG1 was reduced significantly in the absence of functional MLL1 (Fig. 2B), as

was the frequency of IgG1$^+$ or IgE$^+$ GC B cells and IgG1$^+$ or IgE$^+$ PC (Fig. 2C-G). Thus, these

results implied that IgG1 and IgE were not responsible for the accelerated worm clearance

observed in *Mll1*-deficient mice.

IgA is also key in the protection of human and murine intestinal mucosa (16, 18). IgA antibodies

have been linked to *T. muris* worm expulsion from the intestine (25). Thus, we investigated

whether MLL1 regulates the production of IgA. As mLN and Peyer's patches are both IgA-

inductive sites in the intestinal system (41), IgA$^+$ cells were studied in both sites on d14 post-

infection. Both mLN and Peyer's patches showed increased IgA$^+$ GC after B cell-intrinsic deletion

of *Mll1* (Fig. 2H-J). Accordingly, serum total IgA (Fig. 2K) and fecal total IgA (Fig. 2L) were both

significantly increased in *T. muris*-infected $Mll1^{f/f}Cd23^{Cre/+}$ mice compared to control mice. It was

possible that this increase in IgA may already have been present in $Mll1^{f/f}Cd23^{Cre/+}$ mice prior to

infection, and not directly driven by the B cell response to *T. muris* infection. We therefore assessed

IgA levels in uninfected mice. Fecal total IgA (Fig. S1A) and serum total IgA (Fig. S1B) were

comparable between uninfected $Cd23^{cre/+}$ and $Mll1^{f/f}Cd23^{Cre/+}$ mice, indicating that the elevated

IgA detected in the absence of *Mll1* was specifically induced by *T. muris* infection. Thus, while

IgG1$^+$ and IgE$^+$ cells were decreased in the absence of MLL1 *in vivo*, IgA was significantly

increased, correlating with accelerated helminth expulsion.

To examine whether the increased production of IgA observed *in vivo* was due to B cell-intrinsic

or B cell-extrinsic mechanisms, we assessed the ability of MLL1-deficient B cells to differentiate

into IgA$^+$ PC *in vitro*. Naïve B cells from *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ and *Cd23*$^{cre/+}$ mice were stimulated *in*

vitro with IgA-promoting stimuli (CD40L, IL-4, IL-5, TGFβ and retinoic acid). After 4 days,

plasmablasts were assessed. MLL1-deficient cells had a significantly increased propensity to form

IgA$^+$ plasmablasts, compared to control B cells (Fig. 2M-O). This was also reflected in the increase

in IgA secreted in the cultures (Fig. 2P). Therefore, in the absence of MLL1, B cells were

intrinsically prone to produce IgA-producing PC.

MLL1 regulates intestinal localization of IgA$^+$CCR9$^+$ PCs

We next investigated the transcriptional programs altered in the absence of MLL1 that may tailor

B cell responses in GALT. GC B cells (CD19$^+$IgD^{lo}CD95^{hi}CD38^{lo}) were sort-purified from mLN

of *T. muris*-infected *Cd23*$^{cre/+}$ and *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ mice and RNA-sequencing undertaken. 33 genes

with > 2-fold expression change were observed in GC B cells after B cell-intrinsic deletion,

including 7 down-regulated genes and 26 up-regulated genes (Fig. 3A). Differential gene KEGG

enrichment analysis was performed to get further insight into the gene pathway in GC B cells

mediated by conditional deletion of *Mll1* in B cells. Consistent with the increased IgA in MLL1-

deficient mice, KEGG analysis identified a gene signature related to 'an intestinal immune network

for IgA production', along with several disease and stress pathways (Fig. 3B). Specifically, this

analysis identified *Ccr9* as significantly upregulated in *Mll1*^{f/f}*Cd23*^{cre/+} mice than *Cd23*^{cre/+} mice

(Fig. 3C).

Upregulation of CCR9 induces the migration of IgA$^+$ PCs from mLN to the intestinal lamina

propria in response to CCL25 (42). Given that *Ccr9* was identified as differentially expressed in

the absence of *Mll1*, we examined whether MLL1 regulated CCR9 expression on IgA$^+$ PCs in

GALT. In contrast to GC B cells (Fig. 3A-C), the frequency of CCR9$^+$ IgA$^+$ PCs and CCR9

expression on IgA$^+$ PCs in mLN were decreased in MLL1-deficient mice (Fig. 3D-E), whereas

there was no change detected in the small intestine (data not shown). As *T. muris* infection mainly

occurs in the large intestine, including colon (43), we hypothesized that CCR9 may be regulating

IgA$^+$ PC migration to the colon lamina propria (Fig. 3F). Indeed, the frequency of CCR9$^+$ IgA$^+$

PCs was significantly increased in colon lamina propria of *Mll1*^{f/f}*Cd23*^{cre/+} mice compared to

Cd23^{cre/+} mice, respectively (Fig. 3G). Thus, these results indicated that B cell-specific *Mll1*

deletion increased CCR9$^+$IgA$^+$ PCs to the colon.

To confirm that the observed regulation of IgA$^+$ PC by MLL1 was not unique to the *T. muris*-

induced GALT immune response, we used the bacteria *Citrobacter rodentium*, which also induces

IgA production (44). *Mll1*^{f/f}*Cd23*^{cre/+} and *Cd23*^{cre/+} mice were infected with *C. rodentium* and

responses assessed at d14 post-infection (Fig. 3H). In accordance with the humoral response to *T.*

muris, *Mll1*-deficiency resulted in a significant decrease of CCR9$^+$IgA$^+$ PC in mLN, coupled with

an increase of CCR9$^+$IgA$^+$ cells in the colon lamina propria, and a concordant increase in serum

IgA (Fig. 3I-K). Taken together, the absence of *Mll1* in B cells leads to increased production of

IgA$^+$ PC *in vitro*, as well as increased IgA and localization of CCR9$^+$ IgA$^+$ PCs in the colon lamina

propria in response to *T. muris* or *C. rodentium in vivo*.

B cell-intrinsic deletion of MLL1 induces changes to cecum morphology and microbiota

Another key role for B cells and IgA in the GALT is the regulation of the microbiome (45, 46).

Helminth infection also has a significant effect on microbiota diversity, and in turn, the microbiota

can either support or help expel the helminth (47, 48). Given this, we hypothesized that the changes

to the humoral response and worm clearance in the absence of B cell expression of MLL1 may

lead to significant changes in the gut microenvironment and microbiota. To investigate, we tested

whether there were detectable changes to cecum morphology and microbiota diversity in

Mll1$^{f/f}$*Cd23*$^{cre/+}$ mice. Goblet cells secrete mucin, which can induce worm expulsion (49). A

significantly increased number of goblet cells was detected in the cecum of *T. muris*-infected

Mll1$^{f/f}$*Cd23*$^{cre/+}$ mice compared to *Cd23*$^{cre/+}$ mice on d14 post-infection (Fig. 4A, B). To investigate

microbiota diversity, 16S rRNA microbiome sequencing was performed on the cecal feces

collected 14 days post-*T.muris* infection (Fig. 4C; Fig. S1C-D). Although β-diversity did not show

a significant change (*p=0.0577*) (Fig. S1C), Linear discriminant analysis Effect Size (LEfSe)

identified significant changes in composition at genus level, with 5 bacterial groups positively

associated with *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ mice compared to *Cd23*$^{cre/+}$ mice, and 4 genera found only in

Cd23$^{cre/+}$ mice (Fig. 4D, E). Functionally, bacteria with increased abundance have been noted as

beneficial, while decreased bacteria can be pathogenic (50-53). For example, the anti-

inflammatory bacteria *Prevotellaceae* and *Lachnospiraceae* (54, 55) are increased in

Mll1$^{f/f}$*Cd23*$^{cre/+}$ mice, with *Lachnospiraceae* negatively associated with *T. trichiura* egg burden

(51, 56). In contrast, microbiota that have been positively associated with helminth infection –

Rikenellaceae (48) and *Alloprevotella* (57) – were absent in in *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ mice, compared to

Cd23^{cre/+} mice (Fig. 4E). Taken together, B cell-intrinsic deletion of MLL1 altered the cecal

microbiota, correlating to a shift to a healthier microbiome during an IgA-dominated response to

helminth infection.

Discussion

Discussion

Uncovering key regulators of mucosal immune responses is important for designing vaccines that

can induce effective protective responses harnessing the GALT. Here, we revealed the regulatory

role of MLL1 in worm clearance and the microbiota during helminth infection. MLL1 deletion in

mature B cells led to decreased GC B cells but increased IgA, correlating with accelerated worm

clearance and a decrease in pathogenic microbiota. Our study thus reveals a novel role for MLL1

in tailoring the antibody response, increases understanding of molecular regulators of humoral

response within the GALT, and provides a fundamental groundwork for developing therapeutics

to prevent or treat helminth infection.

During an immune response, B cell differentiation is orchestrated by both extrinsic and intrinsic

factors. The infecting pathogen or immunizing agent induces the activation of immune cells that

will secrete cytokines and other factors into the microenvironment, creating a milieu that will

influence the fate of B cells and type of antibody produced (38). These signals can then induce the

function of transcription factors and epigenetic regulators within the nucleus, to promote or repress

gene expression programs and ultimately, regulate B cell differentiation and function (58, 59).

While some key inducers of IgA isotype switching have been discovered, such as RUNX3 and

p300, TGFβ and retinoic acid (60-62), we reveal here that inhibiting MLL1 can significantly

increase production of IgA$^+$ PC, both *in vitro* and *in vivo*, and promotes the localisation of

CCR9$^+$IgA$^+$ cells in the colon lamina propria. There appeared to be a humoral response trade-off,

however, as GC B cells were significantly reduced and showed diminished frequencies of IgG1$^+$

and IgE$^+$ cells. These findings also point to non-overlapping roles of MLL1 and KMT2D (MLL2)

in B cell biology, as *Kmt2d* deletion promotes GC B cell formation (31). Unique roles for these

methyltransferases have also been noted in other systems, and is likely due to differing histone-

modifying activity and gene targets (63-65). In contrast to KMT2D, we did not observe global

H3K4me3 loss (data not shown). MLL1 has a wider array of histone-modifying activity and

molecular partners, and thus future studies should delineate the exact histone modifications and

target loci that are modified by MLL1 in GALT B cells.

Transmissible helminth infection affects approximately one quarter of the world's population (1-

5). Long-term drug therapy induces the development of drug-resistant parasites, and the treatment

of reinfection causes a huge economic burden (9). Therefore, successful long-term public health

interventions such as preventative vaccines are needed, but generation of an effective vaccine has

remained elusive. Furthermore, the need for promoting protective IgA through new-generation

mucosal vaccines extends beyond helminth-specific vaccines (66), and is an area of intense current

research. One major hurdle to the development of helminth-specific vaccines has been vaccine-

induced formation of antigen-specific IgE, leading to an allergic response (8, 12). In this context,

targeting MLL1 in B cells could provide benefit by repressing IgE while promoting IgA. Small

molecule inhibitors of MLL1 are in development for treatment of MLL-rearranged acute

leukemias (67). It is important to note that the delivery of a MLL1 inhibitor would need to be B

cell-targeted, given the roles of MLL1 in intestinal stem cells (28) and Th2 cells (27).

In summary, our study has uncovered MLL1 as a novel regulator of GALT B cell responses.

These results provide further understanding of the roles of histone modifiers in tailoring humoral

immune responses. More broadly, improving our understanding on how to promote effective

IgA-mediated responses in GALT will offer new avenues for oral and mucosal vaccine design

and in the treatment of intestinal helminth infection.

Materials and Methods:

Mice

Cd23[cre/+] mice (68) and *Mll1*[f/f] mice (69) were generously provided by Meinrad Busslinger

(Institute of Molecular Pathology, Austria) and Patricia Ernst (University of Colorado, USA),

respectively. *Mll1*[f/f]*Cd23*[cre/+] on C57Bl/6 background were generated by crossing *Cd23*[cre/+] with

Mll1[f/f] mice. Mice were bred at the Monash Animal Research Platform under specific-pathogen-

free conditions. All animal experimentation was performed following the Australian National

Health and Medical Research Council (NHMRC) Code for the Care and Use of Animals for

Scientific Purposes. Procedures were approved by Monash and WEHI Animal Ethics Committees.

Both males and females, over 6 weeks old, were used in this study.

Immunization: Mice were intraperitoneally immunized with 100µg NP_{13}-KLH precipitated in the

adjuvant alum. *T. muris infection:* mice were infected with 200 *T. muris* eggs (70) by oral gavage.

C. rodentium infection: mice infected with $2x10^9$ colony-forming units of *C. rodentium* by oral

gavage.

Flow cytometry, cell sorting and antibodies

Single-cell suspensions were stained with fluorochrome-labelled antibodies (Table 1) in PBS

2%FCS for flow cytometric analysis or cell sorting as previously described (35). Rat IgG2b kappa

(2.4G2) was used to block the binding of antibodies by cell-surface Fc-receptors and normal rat

serum was used to block non-specific binding. For IgA staining, a transcription factor buffer set

(BD Pharmingen) was used following the manufacturer's recommendations. Cells were analyzed

using a BD-LSRFortessa X-20 flow cytometer and FACSDiva software (BD Biosciences). Data

was analyzed using FlowJo™ software (BD Life Sciences). For cell sorting, cells were sort-

purified using a BD Influx cell sorter (BD Biosciences).

RT-qPCR

Either RNeasy Plus Micro Kit (QIAGEN) or Direct-zol RNA Microprep kit (Zymo) were used to

extract RNA from sort-purified cells following the manufacturer's recommendations, and TRIzol

was used to extract RNA from homogenized proximal colon tissue. Reverse Transcription was

performed using High-Capacity cDNA Reverse Transcription kit (Thermo Fisher Scientific).

SYBR Green PCR kit (QIAGEN) was used to perform qPCR. *Rpl32* was used as an internal control

for normalization and calculating the $\Delta\Delta$Ct value (71). Primers used were: *Mll1* 5′-

GCAGATTGTAAGACGGCGAG-3′,   5′-GCAGATTGTAAGACGGCGAG-3′;   *Rpl32*   5′-

ATCAGGCACCAGTCAGACCGAT-3′, 5′- GTTGCTCCCATAACCGATGTTGG-3′.

Worm counting

To determine worm clearance, ceca were collected from *T. muris*-infected mice, worms extracted

and counted.

***T. muris*-specific ELISA**

High-binding 96-well ELISA-plates (Sarstedt) were coated with *T. muris* antigen (70) diluted in

carbonate buffer, and incubated overnight at 4°C. Following wash and blocking steps, samples that

were serially diluted and incubated for 1 hour at room temperature. After washing, anti-IgG1-HRP

(Southern Biotech) was added and incubated for 45 minutes at room temperature. The wells were

washed and TMB substrate (eBioscience) added, and after 1 minute, stopped with 1N HCl. Plates

were read at 450nm using a BMG OPTIMA plate reader.

Serum IgA ELISA and fecal IgA ELISA

Sera: blood was collected by cardiac puncture immediately post-mortem and sera collected post-

centrifugation. *Fecal extraction:* performed as previously described (72, 73) by adding 0.1g of

fecal pellets to 1ml of PBS containing complete mini protease inhibitor cocktail (Roche). Solids

were separated by centrifugation and supernatants (fecal extract) stored at -20°C until use. *IgA*

ELISAs: high-binding 96-well ELISA plates were coated using unlabelled IgA coating antibodies

(Southern Biotech). Following wash and blocking steps, samples that were serially diluted and

incubated for 37°C for 3 hours. Following washing, anti-IgA-HRP (Southern Biotech) was added

and incubated for 1 hour at 37°C. Plates were washed and developed with OPD-substrate solution

(Sigma-Aldrich). The plates were read at 450nm using a BMG OPTIMA plate reader.

Immunohistochemistry

mLN from *T. muris*-infected mice were first fixed in periodate-lysine-paraformaldehyde fixative,

immersed in 30% sucrose and then subsequently embedded and frozen in Optimal Cutting

Temperature compound (Tissue-Tek). 7μm tissue sections were cut using a microtome (Leica) and

mounted on Superfrost Plus slides (Thermo Fisher). Sections were fixed with cold acetone (Sigma)

at -20°C for 10 minutes. Sections were stained with PNA (Vector Labs) and B220 (BD Bioscience).

Slides were observed and images were captured under 4X and 10X magnification using a Nikon

microscope.

Goblet cell immunohistochemistry: formalin-fixed (10% neutral buffered), Swiss-rolled ceca were

stained with periodic acid Schiff and Alcian blue. Sections were scanned using an Aperio

Scanscope AT Turbo (Leica), at 20X magnification. Goblet cell number per crypt was counted for

20 crypts/mouse across the entire cecum using Aperio ImageScope (Leica) and ImageJ (NIH).

***In vitro* B cell culture**

B cells from the spleen were isolated using the EasySep™ Mouse B Cell Isolation Kit

(STEMCELL Technologies) and labeled with Cell Trace Violet (ThermoFisher). 5×10^4 labeled

cells were resuspended in B cell media (RPMI, 5%FCS, 50μM 2-mercaptoethanol and 2mM

glutamine) and stimulated with 50ng/ml CD40L (R&D Systems), 50ng/ml IL-4 (STEMCELL

Technologies), 5ng/ml IL-5 (R&D Systems), 10ng/ml TGFβ (R&D Systems) and 10nM retinoic

acid.

RNA-sequencing

GC B cells (B220$^+$IgD^{lo}CD95^{hi}CD38^{lo}) were sort-purified from mLN of *T.muris*-infected *Cd23*$^{cre/+}$

and *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ mice. RNA was extracted using the RNeasy Plus Micro Kit according to the

manufacturer's instructions. Total mRNA was quantified with the Qubit Fluorometer (Invitrogen)

and RNA integrity number values were assessed using the Agilent Bioanalyzer. mRNA libraries

were sequenced on an Illumina Novaseq using a 2x150 paired-end (PE) configuration according

to the manufacturer's instructions. Data was processed and analysed as previously described (33,

35). The following values were used to detect differentially expressed genes: fold change >2; q-

value (FDR-adjusted p-value) < 0.05. KEGG (Kyoto Encyclopedia of Genes and Genomes)

enrichment analysis was performed using KEGG databases. Data has been deposited at GEO and

publicly available. Accession number is GSE237391.

16S rRNA microbiome sequencing

Feces from individual mice were collected from the cecum. DNA was extracted using ISOLATE II Fecal DNA Kit (Bioline) according to the manufacturer's instructions. Amplicons were generated from the V3 and V4 regions of 16S rRNA using 341F-targeted primers (5′-CCTAYGGGRBGCASCAG -3′ and 5′-GGACTACNNGGGTATCTAAT-3′). Diversity profiling analysis were performed with QIIME 2 2019.7 (74). Demultiplexed raw reads were primer trimmed and quality filtered using the cutadapt plugin followed by denoising with DADA2 (75) (via q2-dada2). Taxonomy was assigned to ASVs using the q2-feature-classifier (76) classify-sklearn naïve Bayes taxonomy classifier. Linear Discriminant Analysis Effect Size (LefSe) was performed using MicrobiomeAnalyst (77).

Statistical analysis

Experimental data are presented as mean ± standard error of the mean (SEM). Statistical analyses performed with Mann-Whitney non-parametric test using GraphPad Prism 8 (GraphPad Software). $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$. Mice assessed to be non-responders were excluded from all analyses.

Table 1. Antibodies

Antibody	Fluorochrome	Company
CD19	APC-Cy7	Biolegend #B207588
CD95	PE-Cy7	BD Bioscience #7018663
CD38	Pacific Blue	Biolegend #B240225

IgD	A488	Biolegend #B206502
IgG1	APC	BD Bioscience #7128657
B220	PacBlue	Biolegend #B207505
IgM	PE	Biolegend #B221864
IgD	PerCp-Cy5.5	Biolegend #B186570
CD23	AF488	Biolegend #B156559
CD21/CD35	APC	Biolegend #B195498
CD5	Biotin	BD Bioscience #550874
CD3e	APC-Cy7	BD Bioscience #558117
CD19	PE	BioLegend #B207501
CD138	PE	BioLegend #B203320
IgE	FITC	BD Bioscience #5163873

Acknowledgements

We thank Moshe Olshansky, Alana Kirn and members of the Good-Jacobson lab for technical assistance; Patricia Ernst for *Mll* floxed mice, and staff of the Monash University FlowCore, Monash Animal Research Platform, Monash Histology Platform, and the Monash Genomics and Bioinformatics Platform. **Funding:** This work was supported by a Bellberry-Viertel Senior Medical Research Fellowship (KLG-J); National Health and Medical Research Council (NHMRC) Career Development Fellowship 1108066 (KLG-J); Australian Research Council (ARC) DP220102867 (KLG-J); Monash University-China Scholarship Council Joint PhD Scholarship (YZ); and Monash University Scholarships (CC, AZ, AK, LX).

References

1.   P. M. Jourdan, P. H. L. Lamberton, A. Fenwick, D. G. Addiss, Soil-transmitted helminth infections. *Lancet* **391**, 252-265 (2018).

2.   A. Zaini, K. L. Good-Jacobson, C. Zaph, Context-dependent roles of B cells during intestinal helminth infection. *PLoS Negl Trop Dis* **15**, e0009340 (2021).

3.   WHO, Soil-transmitted helminth infections. *[online]* https://www.who.int/news-room/fact-sheets/detail/soil-transmitted-helminth-infections (Accessed 2 March 2020).

4.   D. W. Crompton, M. C. Nesheim, Nutritional impact of intestinal helminthiasis during the human life cycle. *Annu Rev Nutr* **22**, 35-59 (2002).

5.   L. Chelkeba, Z. Mekonnen, Y. Alemu, D. Emana, Epidemiology of intestinal parasitic infections in preschool and school-aged Ethiopian children: a systematic review and meta-analysis. *BMC Public Health* **20**, 117 (2020).

6.   R. L. Pullan, J. L. Smith, R. Jasrasaria, S. J. Brooker, Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasit Vectors* **7**, 37 (2014).

7.   J. C. Dunn, H. C. Turner, A. Tun, R. M. Anderson, Epidemiological surveys of, and research on, soil-transmitted helminths in Southeast Asia: a systematic review. *Parasit Vectors* **9**, 31 (2016).

8.   A. Zawawi, K. J. Else, Soil-Transmitted Helminth Vaccines: Are We Getting Closer? *Frontiers in immunology* **11** (2020).

9.   C. E. James, A. L. Hudson, M. W. Davey, Drug resistance mechanisms in helminths: is it survival of the fittest? *Trends Parasitol* **25**, 328-335 (2009).

411   10.   J. C. Dunn *et al.*, Soil-transmitted helminth reinfection four and six months after mass

412   drug administration: results from the delta region of Myanmar. *PLoS Negl Trop Dis* **13**,

413   e0006591 (2019).

414   11.   H. Dixon, C. E. Johnston, K. J. Else, Antigen selection for future anti-Trichuris vaccines:

415   a comparison of cytokine and antibody responses to larval and adult antigen in a primary

416   infection. *Parasite Immunol* **30**, 454-461 (2008).

417   12.   D. J. Diemert, M. E. Bottazzi, J. Plieskatt, P. J. Hotez, J. M. Bethony, Lessons along the

418   Critical Path: Developing Vaccines against Human Helminths. *Trends Parasitol* **34**, 747-

419   758 (2018).

420   13.   R. K. Grencis, Cytokine-mediated regulation of intestinal helminth infections: the

421   Trichuris muris model. *Ann Trop Med Parasitol* **87**, 643-647 (1993).

422   14.   M. Bemark, M. J. Pitcher, C. Dionisi, J. Spencer, Gut-associated lymphoid tissue: a

423   microbiota-driven hub of B cell immunity. *Trends Immunol* **45**, 211-223 (2024).

424   15.   D. Piovesan *et al.*, c-Myb Regulates the T-Bet-Dependent Differentiation Program in B

425   Cells to Coordinate Antibody Responses. *Cell Rep* **19**, 461-470 (2017).

426   16.   J. Spencer, L. M. Sollid, The human intestinal B-cell response. *Mucosal Immunology* **9**,

427   1113-1124 (2016).

428   17.   P. E. Conrey *et al.*, IgA deficiency destabilizes homeostasis toward intestinal microbes

429   and increases systemic immune dysregulation. *Sci Immunol* **8**, eade2335 (2023).

430   18.   O. Pabst, New concepts in the generation and functions of IgA. *Nat Rev Immunol* **12**,

431   821-832 (2012).

432   19.   A. C. S. Ramos *et al.*, The role of IgA in gastrointestinal helminthiasis: A systematic

433   review. *Immunol Lett* **249**, 12-22 (2022).

434   20.   F. Fransen *et al.*, BALB/c and C57BL/6 Mice Differ in Polyreactive IgA Abundance,

435   which Impacts the Generation of Antigen-Specific IgA and Microbiota Diversity.

436   *Immunity* **43**, 527-540 (2015).

437   21.   K. Koyama, H. Tamauchi, M. Tomita, T. Kitajima, Y. Ito, B-cell activation in the

438   mesenteric lymph nodes of resistant BALB/c mice infected with the murine nematode

439   parasite Trichuris muris. *Parasitol Res* **85**, 194-199 (1999).

440   22.   M. Clerc, G. Devevey, A. Fenton, A. B. Pedersen, Antibodies and coinfection drive

441   variation in nematode burdens in wild mice. *Int J Parasitol* **48**, 785-792 (2018).

442   23.   C. McGuire, W. C. Chan, D. Wakelin, Nasal immunization with homogenate and peptide

443   antigens induces protective immunity against Trichinella spiralis. *Infect Immun* **70**, 7149-

444   7152 (2002).

445   24.   Y. Yang *et al.*, Oral vaccination with Ts87 DNA vaccine delivered by attenuated

446   Salmonella typhimurium elicits a protective immune response against Trichinella spiralis

447   larval challenge. *Vaccine* **28**, 2735-2742 (2010).

448   25.   T. I. Roach, K. J. Else, D. Wakelin, D. J. McLaren, R. K. Grencis, Trichuris muris:

449   antigen recognition and transfer of immunity in mice by IgA monoclonal antibodies.

450   *Parasite Immunol* **13**, 1-12 (1991).

451   26.   T. Gan, B. E. Li, B. P. Mishra, K. L. Jones, P. Ernst, MLL1 Promotes IL-7

452   Responsiveness and Survival during B Cell Differentiation. *J Immunol* **200**, 1682-1691

453   (2018).

454   27.   M. Yamashita *et al.*, Crucial Role of MLL for the Maintenance of Memory T Helper Type

455   2 Cell Responses. *Immunity* **24**, 611-622 (2006).

456   28.   N. Goveas *et al.*, MLL1 is required for maintenance of intestinal stem cells. *PLoS Genet*
457   **17**, e1009250 (2021).

458   29.   M. Caganova *et al.*, Germinal center dysregulation by histone methyltransferase EZH2
459   promotes lymphomagenesis. *J Clin Invest* **123**, 5009-5022 (2013).

460   30.   W. Béguelin *et al.*, EZH2 Is Required for Germinal Center Formation and Somatic EZH2
461   Mutations Promote Lymphoid Transformation. *Cancer cell* **23**, 677-692 (2013).

462   31.   J. Zhang *et al.*, Disruption of KMT2D perturbs germinal center B cell development and
463   promotes lymphomagenesis. *Nat Med* **21**, 1190-1198 (2015).

464   32.   K. Hatzi *et al.*, Histone demethylase LSD1 is required for germinal center formation and
465   BCL6-driven lymphomagenesis. *Nat Immunol* **20**, 86-96 (2019).

466   33.   A. Di Pietro *et al.*, Targeting BMI-1 in B cells restores effective humoral immune
467   responses and controls chronic viral infection. *Nat Immunol* **23**, 86-98 (2022).

468   34.   K. L. Good-Jacobson *et al.*, Regulation of germinal center responses and B-cell memory
469   by the chromatin modifier MOZ. *Proc Natl Acad Sci U S A* **111**, 9585-9590 (2014).

470   35.   L. Kealy *et al.*, The Histone Methyltransferase DOT1L Is Essential for Humoral Immune
471   Responses. *Cell Rep* **33**, 108504 (2020).

472   36.   Y. Nakata *et al.*, c-Myb, Menin, GATA-3, and MLL form a dynamic transcription
473   complex that plays a pivotal role in human T helper type 2 cell development. *Blood* **116**,
474   1280-1290 (2010).

475   37.   A. Zaini *et al.*, Heterogeneous Tfh cell populations that develop during enteric helminth
476   infection predict the quality of type 2 protective response. *Mucosal Immunol* **16**, 642-657
477   (2023).

478   38.   K. L. Good-Jacobson, Strength in diversity: Phenotypic, functional, and molecular

479   heterogeneity within the memory B cell repertoire. *Immunol Rev* **284**, 67-78 (2018).

480   39.   K. J. Montaño, C. Cuéllar, J. Sotillo, Rodent Models for the Study of Soil-Transmitted

481   Helminths: A Proteomics Approach. *Front Cell Infect Microbiol* **11**, 639573 (2021).

482   40.   N. M. Blackwell, K. J. Else, B cells and antibodies are required for resistance to the

483   parasitic gastrointestinal nematode Trichuris muris. *Infect Immun* **69**, 3860-3868 (2001).

484   41.   A. Reboldi, J. G. Cyster, Peyer's patches: organizing B-cell responses at the intestinal

485   frontier. *Immunol Rev* **271**, 230-245 (2016).

486   42.   O. Pabst *et al.*, Chemokine receptor CCR9 contributes to the localization of plasma cells

487   to the small intestine. *J Exp Med* **199**, 411-416 (2004).

488   43.   K. S. Hayes, R. K. Grencis, Trichuris muris and comorbidities - within a mouse model

489   context. *Parasitology* **148**, 1-9 (2021).

490   44.   N. Kamada *et al.*, Humoral Immunity in the Gut Selectively Targets Phenotypically

491   Virulent Attaching-and-Effacing Bacteria for Intraluminal Elimination. *Cell Host*

492   *Microbe* **17**, 617-627 (2015).

493   45.   O. Pabst, E. Slack, IgA and the intestinal microbiota: the importance of being specific.

494   *Mucosal Immunol* **13**, 12-21 (2020).

495   46.   C. Gutzeit, G. Magri, A. Cerutti, Intestinal IgA production and its role in host-microbe

496   interaction. *Immunol Rev* **260**, 76-85 (2014).

497   47.   K. Llinas-Caballero, L. Caraballo, Helminths and Bacterial Microbiota: The Interactions

498   of Two of Humans' "Old Friends". *Int J Mol Sci* **23** (2022).

499   48.   J. Kupritz, A. Angelova, T. B. Nutman, P. H. Gazzinelli-Guimaraes, Helminth-Induced

500   Human Gastrointestinal Dysbiosis: a Systematic Review and Meta-Analysis Reveals

Insights into Altered Taxon Diversity and Microbial Gradient Collapse. *mBio* **12**,

e0289021 (2021).

49. S. Z. Hasnain *et al.*, Mucin gene deficiency in mice impairs host resistance to an enteric

parasitic infection. *Gastroenterology* **138**, 1763-1771 (2010).

50. X. Liu *et al.*, Blautia-a new functional genus with potential probiotic properties? *Gut*

Microbes **13**, 1-21 (2021).

51. B. A. Rosa *et al.*, Whipworm-Associated Intestinal Microbiome Members Consistent

Across Both Human and Mouse Hosts. *Front Cell Infect Microbiol* **11**, 637570 (2021).

52. F. Zhang *et al.*, Curcumin Alleviates DSS-Induced Anxiety-Like Behaviors via the

Microbial-Brain-Gut Axis. *Oxid Med Cell Longev* **2022**, 6244757 (2022).

53. H. Wang *et al.*, Aberrant Gut Microbiome Contributes to Intestinal Oxidative Stress,

Barrier Dysfunction, Inflammation and Systemic Autoimmune Responses in MRL/lpr

Mice. *Front Immunol* **12**, 651191 (2021).

54. W. Y. Wu, P. L. Chou, J. C. Yang, C. T. Chien, Silicon-containing water intake confers

antioxidant effect, gastrointestinal protection, and gut microbiota modulation in the

rodents. *PloS one* **16**, e0248508 (2021).

55. L. Ma *et al.*, Spermidine improves gut barrier integrity and gut microbiota function in

diet-induced obese mice. *Gut Microbes* **12**, 1-19 (2020).

56. S. C. Lee *et al.*, Linking the effects of helminth infection, diet and the gut microbiota

with human whole-blood signatures. *PLoS Pathog* **15**, e1008066 (2019).

57. M. A. Toro-Londono, K. Bedoya-Urrego, G. M. Garcia-Montoya, A. L. Galvan-Diaz, J.

F. Alzate, Intestinal parasitic infection alters bacterial gut microbiota in children. *PeerJ* **7**,

e6200 (2019).

58.   Y. Zhang, K. L. Good-Jacobson, Epigenetic regulation of B cell fate and function during

an immune response. *Immunol Rev* **288**, 75-84 (2019).

59.   L. Cooper *et al.*, Type I interferons induce an epigenetically distinct memory B cell

subset in chronic viral infection. *Immunity* **57**, 1037-1055 (2024).

60.   S. R. Park, E. K. Lee, B. C. Kim, P. H. Kim, p300 cooperates with Smad3/4 and Runx3 in

TGFbeta1-induced IgA isotype expression. *Eur J Immunol* **33**, 3386-3392 (2003).

61.   A. Cerutti, The regulation of IgA class switching. *Nat Rev Immunol* **8**, 421-434 (2008).

62.   A. Reboldi *et al.*, IgA production requires B cell interaction with subepithelial dendritic

cells in Peyer's patches. *Science* **352**, aaf4822 (2016).

63.   P. Wang *et al.*, Global analysis of H3K4 methylation defines MLL family member targets

and points to a role for MLL1-mediated H3K4 methylation in the regulation of

transcriptional initiation by RNA polymerase II. *Mol Cell Biol* **29**, 6074-6085 (2009).

64.   B. P. Mishra *et al.*, The histone methyltransferase activity of MLL1 is dispensable for

hematopoiesis and leukemogenesis. *Cell Rep* **7**, 1239-1247 (2014).

65.   X. Huang *et al.*, Targeting Epigenetic Crosstalk as a Therapeutic Strategy for EZH2-

Aberrant Solid Tumors. *Cell* **175**, 186-199 e119 (2018).

66.   J. E. Oh *et al.*, Intranasal priming induces local lung-resident B cell populations that

secrete protective mucosal antiviral IgA. *Sci Immunol* **6**, eabj5129 (2021).

67.   T. R. Chern *et al.*, Discovery of Potent Small-Molecule Inhibitors of MLL

Methyltransferase. *ACS Med Chem Lett* **11**, 1348-1352 (2020).

68.   K. Kwon *et al.*, Instructive role of the transcription factor E2A in early B lymphopoiesis

and germinal center B cell development. *Immunity* **28**, 751-762 (2008).

69. C. D. Jude *et al.*, Unique and independent roles for MLL in adult hematopoietic stem

cells and progenitors. *Cell Stem Cell* **1**, 324-337 (2007).

70. F. Antignano, S. C. Mullaly, K. Burrows, C. Zaph, Trichuris muris infection: a model of

type 2 immunity and inflammation in the gut. *J Vis Exp* 10.3791/2774 (2011).

71. K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408 (2001).

72. K. A. Watt, D. H. Nussey, R. Maclellan, J. G. Pilkington, T. N. McNeilly, Fecal antibody

levels as a noninvasive method for measuring immunity to gastrointestinal nematodes in

ecological studies. *Ecol Evol* **6**, 56-67 (2016).

73. J. M. Smeekens *et al.*, Fecal IgA, Antigen Absorption, and Gut Microbiome Composition

Are Associated With Food Antigen Sensitization in Genetically Susceptible Mice. *Front*

Immunol **11**, 599637 (2020).

74. E. Bolyen *et al.*, Reproducible, interactive, scalable and extensible microbiome data

science using QIIME 2. *Nat Biotechnol* **37**, 852-857 (2019).

75. B. J. Callahan *et al.*, DADA2: High-resolution sample inference from Illumina amplicon

data. *Nat Methods* **13**, 581-583 (2016).

76. N. A. Bokulich *et al.*, q2-longitudinal: Longitudinal and Paired-Sample Analyses of

Microbiome Data. *mSystems* **3** (2018).

77. A. Dhariwal *et al.*, MicrobiomeAnalyst: a web-based tool for comprehensive statistical,

visual and meta-analysis of microbiome data. *Nucleic Acids Res* **45**, W180-W188 (2017).

Figure Legends

Figure 1. MLL1 is required for GC B cell differentiation in Th2 cell-biased responses.

(A and B) The expression of *Mll1* was assessed by RT-qPCR in B cell subsets sort-purified from

(A) the mesenteric lymph node (mLN) and (B) Peyer's patches of C57Bl/6 mice. (C) Follicular

(FO) B cells, GC B cells, PC and memory B cells were sort-purified from spleen and *Mll1*

expression assessed. FO B cells were sort-purified from C57Bl/6 mice, GC B cells were sort-

purified from C57Bl/6 mice immunized with NP-KLH in alum at d7 post-immunization, and

memory B cells and PC were sort-purified from C57Bl/6 mice immunized with NP-KLH in alum

at d28 post-immunization. Expression of *Mll1* was normalized to *Rpl32* and then normalized to

naïve (A, B) or follicular (C) B cells. (D) Schematic depiction of *T. muris* experimental setup. (E)

Flow cytometric analyses of $B220^+IgD^{lo}CD95^{hi}CD38^{lo}$ (GC B cells) in mLN *T. muris*-infected

$Cd23^{cre/+}$ and $Mll1^{f/f}Cd23^{cre/+}$ mice at d14 and d28 post-infection. The number in plots shows the

frequency of GC B cells within the $B220^+IgD^{lo}$ population. (F and G) Frequency (F) and number

(G) of GC B cells post-infection. Data are combined from three independent experiments per time

point; n=9 (d14), n=7-9 (d28) and n=10-11 (d35) per genotype. (H) Representative histological

images of mLN from d14 *T. muris*-infected $Cd23^{cre/+}$ and $Mll1^{f/f}Cd23^{cre/+}$ mice: B220 (red) and

PNA (blue). Scale bar = 100μM (I) Flow cytometric analysis of NP^+ GC B cells

$(B220^+IgD^{lo}CD95^{hi}NP^+)$ from d14 post-immunized $Cd23^{cre/+}$ and $Mll1^{f/f}Cd23^{cre/+}$ mice. (J and K)

Frequency (J) and number (K) of GC B cells post-immunization at d14 and d28. Data are combined

from two independent experiments per time point; n=7 (d14) and n=3-6 (d28) per genotype. Error

bars indicate mean ± SEM, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, Mann-Whitney test.

Figure 2. B cell-intrinsic MLL1 deletion leads to rapid worm clearance and increased IgA.

(A) Number of worms in the caecum from *T. muris*-infected $Cd23^{cre/+}$ and $Mll1^{f/f}Cd23^{cre/+}$ mice at

d14, d28 and d35 post-infection. Data are combined from 2-3 experiments per time point; n=10

(d14); n=7-9 (d28); n=7-8 (d35) per genotype. (B) *T. muris*-specific serum IgG1 at d14 post-

infection. (C) Representative flow cytometric plots of IgG1$^+$GC B cells (IgG1$^+$CD95^{hi}CD38^{lo}) and

IgE$^+$ GC B cells (IgE$^+$CD95^{hi}CD38^{lo}) in mLN at d14. The number in plots shows the frequency of

IgG1$^+$ or IgE$^+$ within the GC B cell population. (D-E) Frequency of (D) IgG1$^+$ GC B cells and (E)

IgG1$^+$ PC (IgG1$^+$B220^{lo}CD138^{hi}). Data are combined from three independent experiments, n=9

per genotype. (F-G) Frequency of (F) IgE$^+$ GC B cells and (G) IgE$^+$ PC (IgE$^+$B220^{lo}CD138^{hi}). Data

are combined from two independent experiments, n=6 per genotype. (H) Representative plots

shown of IgA$^+$ GC B cells (IgA$^+$CD95^{hi}CD38^{lo}) in mLN or Peyer's patches at d14. The number in

plots shows the frequency of IgA$^+$ within the GC B cell population. (I and J) Frequency of IgA$^+$

GC B cells in the (I) mLN and (J) Peyer's Patches (PP). Data are combined from two independent

experiments. n=8 or 9 per genotype. (K) Serum IgA and (L) fecal IgA from d14 *T. muris*-infected

$Cd23^{cre/+}$ and $Mll1^{f/f}Cd23^{cre/+}$ mice. Data are combined from 3-4 independent experiments, n=12

per genotype. Error bars indicate mean $\pm$ SEM. *p < 0.05, **p < 0.01, Mann-Whitney test. (M-O)

B cells were stimulated in culture with CD40L, IL-4, IL-5, TGFβ and retinoic acid for 4 days. (M)

Frequency of B cells, (N) frequency of PC, (O) total number of IgA$^+$ PC, and (P) IgA in supernatant

at d4 post-stimulation. Data are combined from two individual experiments, n=5 per genotype.

Error bars indicate mean +- SEM, *p < 0.05, **p < 0.01, Mann-Whitney test.

Figure 3. MLL1 regulates the expression of genes related to IgA production.

(A) Heatmap of RNA-sequencing data with fold change >2 and q-value (FDR-adjusted p-value) < 0.05; each column is an independent sample obtained from sort-purified GC B cells (B220$^+$IgD^{lo}CD95^{hi}CD38^{lo}) from mLN of d13 *T.muris*-infected *Cd23*$^{cre/+}$ and *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ mice, n=2 per genotype. (B) KEGG enrichment pathway analysis of differential gene expression. The size of the dot is positively correlated with the number of differential genes in the pathway. (C) Plots showing FPKM (Fragments per kilo base of transcript per million mapped fragments) for *Ccr9*. (D) Representative flow cytometric plots of CCR9$^+$IgA$^+$PC (CCR9$^+$IgA$^+$B220^{lo}CD138^{hi}) in mLN from d14 *T. muris*-infected *Cd23*$^{cre/+}$ and *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ mice. (E) CCR9 geometric mean fluorescence intensity (MFI) and CCR9$^+$ frequency within IgA$^+$ PC in mLN; n=4 per genotype. (F) Representative plots of CCR9$^+$IgA$^+$ PC (CCR9$^+$IgA$^+$CD3$^-$B220^{lo}Blimp1$^+$) in colon lamina propria from d14 *T. muris*-infected *Cd23*$^{cre/+}$ and *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ mice. (G) Frequency of CCR9$^+$ cells in IgA$^+$ PC population in colon lamina propria. Data shown are from 2 experiments, n=6 per genotype. Error bars indicate mean ± SEM. *p < 0.05, Mann-Whitney test. (H) *Mll1*$^{f/f}$ and *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ mice were infected with *C. rodentium* and assessed at d14 post-infection. (I) Frequency of CCR9$^+$IgA$^+$ PC in mLN. (J) Frequency of CCR9$^+$ PC and CCR9$^+$IgA$^+$ B cells in colon lamina propria. *p < 0.05, unpaired t-test. (K) Serum IgA, combined from two independent experiments. *p < 0.05, Mann-Whitney test.

Figure 4. B cell-specific MLL1 deletion induces a shift in microbiota.

(A) Representative images from histological analyses of swiss-rolled cecum tissue from d14 *T. muris*-infected *Cd23*$^{cre/+}$ and *Mll1*$^{f/f}$*Cd23*$^{cre/+}$ mice. (B) The number of goblet cells per crypt was measured for 20 crypts per mouse across the entire length of the cecum. Combined data from two

independent experiments, n=8 per genotype. Error bars indicate mean ± SEM. *p < 0.05, Mann-

Whitney test. (C) Schematic depiction of *T. muris* experimental setup to assess microbiome. (D)

Microbiota at genus level in cecum feces from d14 *T. muris*-infected *Cd23*^{cre/+} and *Mll1*^{f/f}*Cd23*^{cre/+}

mice was assessed by 16S rRNA gene sequencing. Linear Discriminant Analysis Effect Size

(LefSe) analysis with Log LDA score above 2.0 and P-value cutoff 0.05; n= 5 per genotype. (E)

The log transformed counts (abundance) of significantly changed bacteria identified by

MetagenomeSeq analysis (adjusted p-value cutoff of 0.05).

Supplemental Figure Legend

Supplemental Figure S1.

(A) Fecal IgA and (B) serum IgA from uninfected *Cd23*^{cre/+} and *Mll1*^{f/f}*Cd23*^{cre/+} mice. n= 6 per

genotype. (C) Microbiota at genus level in cecum feces from d14 *T. muris*-infected *Cd23*^{cre/+} and

Mll1^{f/f}*Cd23*^{cre/+} mice was assessed by 16S rRNA gene sequencing. β-diversity profiling of 16s

sequencing. (D) The relative abundance of microbiota at genus level in caecum feces from *T.*

muris-infected *Cd23*^{cre/+} and *Mll1*^{f/f}*Cd23*^{cre/+} mice on d14 post-infection by 16S rRNA gene

sequencing. n= 5 per genotype.

FIGURE 1

FIGURE 3

FIGURE 4

Uninfected mice

A Fecal IgA ELISA

B Serum IgA ELISA

Cd23^{Cre/+}

Mll1^{f/f}Cd23^{Cre/+}

C *T. muris-infected mice*

p-value: 0.057674

D