**Unveiling the potato cultivars with microbiome interactive traits**

**for sustainable agricultural production**

**Tianci Zhao[a], Stefanie N. Vink[a,b], Xiu Jia[a,c], Alexander Erban[d], Stephanie**

**Schaarschmidt[d], Joachim Kopka[d], Ellen Zuther[d,h], Krzysztof Treder[e], Dorota**

**Michałowska[f], Rémy Guyoneaud[g], J. Theo M. Elzenga[a], Eléonore Attard[g], Jo-**

**ana Falcão Salles[a*]**

[a]Genomics Research in Ecology and Evolution in Nature (Green), Groningen Institute for

Evolutionary Life Sciences (GELIFES), University of Groningen, Groningen, 9747AG, the

Netherlands

[b]GreenFinch Research, Emmastraat 15, Groningen, 9722 EW, the Netherlands

[c]Viral Ecology and Omics (VEO) group, Cluster of Excellence Balance of the Microverse,

Friedrich Schiller University Jena, 07745 Jena, Germany

[d]Max-Planck-Institute of Molecular Plant Physiology, 14476 Potsdam, Germany

[e]Plant Breeding and Acclimatization Institute – National Research Institute in Radzików,

Bonin Division, Department of Potato Protection and Seed Science at Bonin, 76-009 Bo-

nin, Bonin Str 3, Poland

[f]EUROPLANT Polska Sp. z o. o., Laski Koszalińskie Biesiekierz 3A, 76-039, Poland

[g]Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM, Pau, France

[h]Robert Koch Institute, Center for Artificial Intelligence in Public Health Research, Nord-

ufer 20,13353 Berlin, Germany

Keywords: leaf metabolites, root exudates, root traits, rhizosphere microbial communities,

plant-microbiome interactions, plant holobiont

## 24  ABSTRACT

Root traits significantly shape rhizosphere microbiomes, yet their impact on microbes is

often overlooked in plant breeding programs. Here, we propose that selecting modern

cultivars based on microbiome interactive traits (MITs), such as root biomass and exudate

patterns, can enhance agricultural sustainability by interacting effectively with soil

microbiomes, which in turn, promotes plant growth and resistance to stress, thereby

reducing reliance on synthetic crop protectants. Through a stepwise selection process, we

chose 51 potato cultivars based on their plant growth matrices with distinct root exudate

patterns. In a greenhouse experiment, we verified their capacity to interact with the soil

microbiome. These were further clustered into functional groups based on their MITs and

leaf metabolites and further characterised by their rhizosphere microbiome (bacterial and

fungal communities). Our results indicate a positive correlation between microbial

community diversity and root biomass. A potential influence of leaf metabolites on the

composition of rhizosphere bacteria was observed, supporting the plant holobiont

framework. We highlight the significance of incorporating positive effects on

microbiomes into the breeding targets to advance the objectives of sustainable agriculture.

Finally, a set of 11 potato cultivars was selected to verify whether those with microbiome-

interactive traits could contribute to sustainable agricultural production under field

conditions.

**1 INTRODUCTION**

Potato (*Solanum tuberosum* L.) cultivations rely heavily on conventional agricultural

practices as a staple crop. Conventional management, which includes the widespread use

of synthetic fertilisers and pesticides, is frequently used to enhance yields and protect

crops from pests and diseases (Timsina, 2018). However, this dependency on synthetic

compounds comes with substantial environmental issues. Overusing chemical inputs in

conventional agriculture leads to water pollution, soil degradation, and soil microbial

biomass and activity reduction, which are crucial in maintaining soil fertility and nutrient

cycling (AL-Ani et al., 2019; Tripathi et al., 2020). Balancing the imperative for high yields

with care for our environment is a critical challenge in the breeding of crops.

In conventional breeding, plants are bred for traits such as high yield, disease resistance,

tolerance to environmental stresses, and improved agronomic traits (Breseghello and

Coelho, 2013). While pivotal in crop development, conventional breeding has

inadvertently led to the dissociation between plant and soil microbiomes, negatively

affecting beneficial plant-microbiome interactions (Spor et al., 2020). Comparisons

between domesticated cultivars and their ancestral plants revealed that the rhizosphere

microbiome of the latter exhibited higher complexity and connectivity (da Silva et al.,

2023; Rossmann et al., 2020). However, relying solely on ancestral or wild plants may

not effectively address the challenge of optimising plant-soil microbiome interactions in

modern agricultural systems. Identifying specific genes associated with beneficial microbiomes in ancestral plants, with the aim of a subsequent transfer to modern crops (Clouse and Wagner, 2021), is complex due to environmental effects on gene expression (Raaijmakers and Kiers, 2022). Despite the reduced focus on plant-microbiome interactions during domestication (Wei and Jousset, 2017), multiple studies have demonstrated that inoculating modern crop cultivars with beneficial microbes can significantly enhance plant growth and stress resistance (Diagne et al., 2020; Fröhlich et al., 2012; Rodriguez et al., 2019; Tiwari et al., 2017). This suggests that modern cultivars retain the capacity for beneficial interactions with soil microbes, which can be leveraged or strengthened. Therefore, it is critical to identify gene markers associated with beneficial microbiomes in modern crops and apply this knowledge in real-world agricultural systems, particularly in the context of climate change.

The connection between root traits and plant productivity in crops has long been acknowledged (Lynch, 1995), and it is increasingly evident that our understanding of this connection should extend to how root traits shape the composition and functionality of the rhizosphere microbiome. While this aspect has received somewhat limited attention within plant breeding (Herms et al., 2022), many studies highlight the role of root traits in the rhizosphere microbiome. For instance, a study on maise revealed that wild maise had a significant impact on the structure of the microbial community, which was attributed to a high root-to-shoot biomass ratio (Szoboszlay et al., 2015). Plant root traits influence the soil microbial community by modulating the fungi-to-bacteria ratio, contributing to nutrient cycling (Wan et al., 2021). Moreover, a study involving beans (Pérez-Jaramillo et al., 2017) indicated that root architecture is linked to increased

specific bacterial taxa, which could be connected to plant health. In addition to the root

system, root exudates substantially impact the rhizosphere microbiome. As a survival

strategy, plants secrete an abundance of compounds in their root exudates that can

influence the diversity of the rhizosphere microbial community, promote the

development of a more complex microbial network and enhance microbial carbon cycling,

thereby facilitating the growth and survival of the host (Wang et al., 2022). A study on

wheat has demonstrated that cultivars with a strong capacity for root exudates contribute

to increased soil microbial diversity (Iannucci et al., 2021). These specific cultivars can be

further explored for their resource utilisation potential in agricultural systems.

Here, we approach the challenge of identifying plant characteristics associated with soil

microbes by exploring the microbiome interactive traits (MITs) of existing potato

cultivars in the European potato germplasm bank. The root system architecture and

exudate patterns, identified as pivotal MITs, guide our stepwise selection process for

cultivars exhibiting high to low MITs. We hypothesise that cultivars with high MITs will

select a more diverse rhizosphere microbiome, exhibiting superior growth to those with

low MITs. A greenhouse experiment was conducted to investigate the relationship

between the plant traits of 51 potato cultivars and the rhizosphere microbial community.

We propose that MITs capable of recruiting beneficial microbes to the rhizosphere should

be considered in the breeding and developing of new crop cultivars. By prioritising these

interactive traits, we aim to contribute not only to the enhancement of crop performance

but also to the interaction between plants and their microbiomes, unravelling valuable

insights for innovative and sustainable agricultural practices.

## 2 MATERIALS AND METHODS

In this project, we initially narrowed down a selection of 148 late-season potato cultivars, including a commercial cultivar Desiree as a reference, selected from a pool of a thousand potato cultivars based on their resistance to pathogens, especially viruses (Table S1). The method for determining the resistance scale is detailed in the study by Michalak and Chrzanowska (2017), which offers a comprehensive overview of the procedure used to assess resistance to potato virus Y. Detailed information is available in The European Cultivated Potato Database (www.europotato.org). These 148 cultivars were then further selected based on the plant growth matrices (e.g., shoot and root length, dry weight, and root-to-shoot dry weight ratio) under sterile conditions using 6-week-old *in vitro* plants (cf. section 2.1). From those, 51 cultivars (including Desiree) were tested for their ability to interact with soil microbiome in a greenhouse experiment (cf. section 2.2). For the selected 51 cultivars, root exudate profiles from the *in vitro* experiment were also analysed.

### 2.1 *In vitro* experiment

Two-week-old potato plantlets grown from vegetative propagation in tissue culture (Bonin, Poland) were transplanted from agar tubes into 15 ml Eppendorf tubes, each containing 12 ml of sterile 0.5 x Hoagland solution. To ensure aseptic conditions, sterile cotton wool was wrapped around the above-ground part of the plantlets, which were then gently placed into the 15 ml tubes. The roots were submerged in the Hoagland medium, while the above-ground part remained above it. The plantlets were allowed to acclimate to their new environment during a one-week growth period in a controlled climate

chamber. The climate chamber maintained a temperature of 22 °C during the day and

18 °C at night, with a photoperiod of 16 hours of light and 8 hours of darkness.

Upon completion of the initial week, the plantlets were transferred to new sterile 15 mL

tubes, maintaining the 12 ml sterile 0.5 x Hoagland solution, as described in the previous

step. To compensate for the liquid lost during the experiment, sterile water was added to

maintain the volume at 12 ml.

After the third week, the plants were harvested for length and dry weight measurements.

Additionally, 1 ml aliquots of the growth medium were collected for root exudate

metabolite analysis. The samples were immediately stored at -80°C until further analysis.

**2.2 Greenhouse experiment**

Fifty-one potato cultivars (including Desiree) were selected for a greenhouse experiment

based on plant growth traits from the *in vitro* experiment, with the cultivars spread across

a range of growth parameters (Figure S1 and Table S2). The soil in this study originated

from a sugar beet field and was sieved with a 2 mm sieve. Before starting the experiment,

six bulk soil samples were kept for physicochemical analyses and DNA extraction.

Potato plantlets obtained from tissue culture from the Institute of Plant Breeding and

Acclimation in Bonin (Bonin, Poland) were transplanted from agar tubes to small pots

filled with homogenised soil. Three replicates were established for each cultivar.

Additionally, six control pots without plants were prepared. The greenhouse temperature

was 22 °C during the day and 18 °C at night, with a photoperiod of 14 hours of light and

10 hours of darkness.

After two weeks, the potato plants were transplanted into larger pots (1.45 L). Plant leaves,

specifically the second or third leaf from the shoot top, were harvested for metabolomics

analysis in the sixth week. The harvested leaves were snap-frozen in liquid nitrogen and

then transferred to a freezer at -80 °C. The rhizosphere soil was obtained by gently

brushing off adhering soil from the roots with a disposable toothbrush. Additionally, six

bulk soil samples were collected from pots without plants. These soil samples were frozen

at -20 °C until DNA extraction the following day. Shoot height and root length were

measured on the day of harvesting, and shoot and root dry weights were measured after

drying at 65 °C for 72 hours.

**2.3 Metabolite profiling of root exudates and leaf tissue**

Metabolite fractions enriched for primary metabolites were profiled by gas

chromatography and electron impact ionisation-time of flight mass spectrometry (GC/EI-

TOF-MS) (Lisec et al., 2006; Erban et al., 2020). Root exudate samples, i.e. equal debris-

free 1 mL volumes, of the *in vitro* experiment, cf. section 2.1, were freeze-dried directly

without further sample preparation. A polar metabolite fraction was prepared from 50

mg snap-frozen leaf samples of the greenhouse experiment, cf. section 2.2. The frozen

samples were extracted by a water-methanol-chloroform solvent mixture; a polar fraction

was prepared from the extracts by water-induced liquid-liquid phase separation and dried

in a speed-vacuum-concentrator, as described earlier (Erban et al., 2020). The dried

fractions were subjected to methoxyamination and trimethylsilylation prior to GC/EI-

TOF-MS analysis. $^{13}C_6$ Sorbitol was added to all leaf samples before metabolite extraction

(Erban et al., 2020). N-alkanes were added to each sample upon chemical derivatisation

for subsequent retention index (RI) calibration (Erban et al., 2020). The GC/EI-TOF-MS

chromatograms were obtained and baseline adjusted by ChromaTOF software (LECO

Instrumente GmbH, Mönchengladbach, Germany) and background corrected using non-

sample controls. Metabolite annotation was performed using TagFinder software

(Luedemann et al., 2008), the NIST17 mass spectral database (U.S. Department of

Commerce, Gaithersburg, USA), and the RI and mass spectral reference data of the Golm

Metabolome Database, http://gmd.mpimp-golm.mpg.de/ (Hummel et al., 2010; Kopka

et al., 2005). Compounds representing known contaminants and added internal

standards were removed from further analysis. Metabolites absent from more than 75%

of all analysed exudates or leaf tissue samples were excluded, resulting in 84 characterised

leaf metabolites (Data S1) and 49 metabolites (Data S2) that were robustly present in root

exudates.

Relative concentrations were normalised for further analyses to record the sample's fresh

weight and internal standard or exudate volume (Schaarschmidt et al., 2020). An ANOVA

tool was used to perform a batch correction of the metabolite data sets for different

measurement batches and the measurement sequence within batches (Lisec et al., 2011).

All presented metabolite data are relative metabolite abundances. Three replicate

samples per cultivar were analysed. The missing values were substituted by zero.

**2.4 Soil sample sequencing and processing**

DNA was extracted from 12 bulk soil samples and 153 rhizosphere samples with DNeasy

PowerSoil Kit (Qiagen, Hilden, Germany) on 0.25 g of soil. DNA extraction followed the

kit's instructions except for the initial stage of bead beating, which was conducted with a

FastPrep-24TM 5G Instrument at 6000 rpm/s for 40 s (MP Biomedicals, Santa Ana,

USA). For sequencing of the bacterial community, the V4 region of the 16S rRNA gene

was targeted using the primer sets 515F (5′-GTGCCAGCMGCCGCGGTAA-3′) and 806R

(5′-GGACTACHVGGGTWTCTAAT-3′) (Caporaso et al., 2012, 2011). For the fungal

community, the ITS2 region was sequenced with the primer sets 5.8SR (5'-

TCGATGAAGAACGCAGCG-3')      and      reverse      primer      ITS4      (5'-

TCCTCCGCTTATTGATATGC-3') (White et al., 1990). The Illumina MiSeq platform was

used for paired-end sequencing (2 × 250bp for 16S, 2 × 300bp for ITS) at the PGTB

(Genome Transcriptome Platform of Bordeaux, Cestas, France).

The 16S rRNA gene sequencing data were processed using a QIIME2 (version 2020.8)

pipeline (Bolyen et al., 2019). Sequences were filtered, denoised, and dereplicated using

the default setting of the Divisive Amplicon Denoising Algorithm (DADA2) plugin (Calla-

han et al., 2016). 16S rRNA taxonomic classification was performed using the q2-feature-

classifier plugin (Bokulich et al., 2018) against the SILVA database (version 138) (Yilmaz

et al., 2014). ITS sequencing data was processed using the PIPITS / 2.4 pipeline (Gweon

et al., 2015). In brief, the PEAR plugin was used to join read pairs (Zhang et al., 2014).

The FASTX-Toolkit was utilised for quality filtering (Gordon and Hannon, 2010). The

fungal-specific ITS2 region was extracted via ITSx (version 1.1b) (Bengtsson-Palme et al.,

2013). The VSEARCH 2.13.3 plugin (Rognes et al., 2016) was used to dereplicate unique

sequences, clustering to 97% sequence identity, and the UNITE Uchime reference dataset

was used for chimera detection (Nilsson et al., 2015). Ultimately, the taxonomy was

assigned with the RDP Classifier against the UNITE database (version 8.0) (Kõljalg et al.,

2013).

**2.5 Statistical Analysis**

A one-way analysis of variance (ANOVA) was used to test variation in plant metabolite

diversity, plant performance (shoot and root growth), and rhizosphere microbial alpha

diversity (Data S1-S4). Post hoc comparisons were performed through Tukey's honest

significant differences or Duncan's multiple range tests.

Analysis of soil microbial community structure (bacteria and fungi) was performed in R

(version 4.2.0). Feature tables were rarefied at 14400 reads for bacterial 16S rRNA gene

and 6361 for fungal ITS sequences, resulting in 8970 amplicon sequence variants (ASVs)

for the bacterial community and 2006 operational taxonomic units (OTUs) for the fungal

community after excluding non-microbial OTUs (Data S4). The rarefactions of feature

tables were generated via 'rarefy' function in the package 'vegan' (Dixon, 2003). The same

package was used to calculate the alpha diversity metrics for microbial communities and

plant metabolites. Specifically, species richness indicates the number of unique species or

metabolites observed, evenness describes how evenly the abundances of different species

or metabolites are distributed, and the Shannon Diversity Index accounts for both the

number of species/metabolites and their relative abundances (Jost, 2006; Wagner et al.,

2018; Young and Schmidt, 2008).

The soil microbial community, root exudate and leaf metabolite compositions (Data S1,

S2 and S4) were visualised through Principal Coordinate Analysis (PCoA) based on Bray-

Curtis distance. This analysis used the 'ape' package in R (Paradis et al., 2004). PCoA

scores per axis were averaged for each sample to illustrate the distribution of distinct

cultivars in PCoA plots clearly. Permutational multivariate analysis of variance

(PERMANOVA) was conducted using the 'adonis' function within the 'vegan' package to

assess the impact of cultivars on soil microbial communities and metabolite profiles

(Dixon, 2003).

Principal Component Analysis (PCA) based on the covariance matrix was performed with

the R package 'FactoMineR' to reduce the dimensionality of the data and to visualise the

distribution of plant cultivars (Lê et al., 2008). The variables included in the analysis were

shoot and root length, shoot and root dry weight, metabolite diversity and composition of

plant leaf tissue and root exudates (Data S3). Due to the significant variation in root

exudate relative abundance observed in the *in vitro* experiment, median values were used

for PCA plotting. Normalisation of plant data and metabolite richness was performed

before analysis by calculating Z-scores. Subsequently, a PERMANOVA was conducted to

assess the influence of variables on the distribution of 51 cultivars. The cultivars were

classified into four functional groups based on their distribution across the first and

second principal components (PC1 and PC2). The classification was done by dividing the

PCA plot into four quadrants, each representing a distinct functional group of cultivars,

reflecting differences in growth characteristics and metabolite profiles. The summarised

group details can be found in Table S3.

To investigate the relationship between plant cultivars and rhizosphere microbiome

composition, we categorised the 51 potato cultivars into four distinct functional groups.

The category was determined by the distribution of cultivars and variables along the first

two axes in the PCA plot.

The correlation between plant growth and related omics datasets (Data S3) was calculated

using the Spearman correlation coefficient via the 'corr.test()' function from the 'psych'

package in R (Revelle, 2024). The resulting p-values were adjusted using the False

Discovery Rate (FDR) method to control for multiple comparisons and minimise false

positives. We used the Mantel test with Spearman's rank correlation to assess the

correlation between microbial community compositions and metabolite profiles, utilising

Bray-Curtis dissimilarity matrices. This analysis aimed to investigate the relationship

between plant leaf metabolites and rhizosphere soil community.

Using the 'ggsankey' R package (Sjoberg, 2023), a Sankey plot was generated based on

functional groups to summarise the selection of the 11 cultivars with the most potential

for future research. The distribution of rhizosphere bacterial and fungal community

compositions was illustrated for the 11 selected cultivars using Spearman correlation

analysis. The first axis of the PCoA plot of bacterial and fungal communities served as the

indicator of community composition (Data S3).

**3 RESULTS**

**3.1 Plant growth**

Plant growth varied across the 51 potato cultivars in the greenhouse experiment. Analysis

of variance (ANOVA) revealed that most of these cultivars' growth patterns differed

significantly according to their identity. Among them, the cultivars showed a more

pronounced effect on root biomass than root length (root biomass: $p = 0.001$, F = 2.26;

root length: $p = 0.003$, F = 1.91; Figure 1). Specifically, the cultivars Inwestor, Salto and

Szyper demonstrated higher root biomass than others. For shoot growth, the cultivars

had a more pronounced effect on shoot length than on shoot biomass (shoot length: $p$ =

0.001, F = 6.80; shoot biomass: $p$ = 0.009, F = 1.75; Figure S2). Astrid, Inwestor, and

Tewadi cultivars were the top three in shoot biomass. Regarding the root-to-shoot ratio,

the length ratio exhibited a significant response to distinct cultivars ($p$ = 0.001; Figure

S3). Conversely, the ratio of root-to-shoot dry weight did not show a substantial response

to different cultivars ($p$ = 0.36; Figure S3).

**Figure 1.** Root growth analysis of 51 potato cultivars. The upper panel displays root length, while the lower

panel illustrates root dry weight. Each colour represents a distinct potato cultivar in alphabetical order. The

upper right corner of each plot displays one-way ANOVA results, where the F-value explains the variation

among different cultivars, and the $p$-value indicates the statistical relationship among cultivars.

Significance levels are denoted as ** ($p$ = 0.01) and *** ($p$ = 0.001).

**3.2 Plant leaf and root exudate metabolites**

To understand the variation in metabolites produced and released by the 51 potato

cultivars, we conducted a Principal Coordinates Analysis (PCoA) and visualised the

distribution of metabolites among different cultivars (Figure 2). The cultivars differed

significantly in metabolite composition from both root exudates and leaves ($p_{\text{root exudate}}$ =

0.001 and $p_{\text{leaf}}$ = 0.001; Figure 2).

Notably, leaf metabolite structure responded more strongly to plant cultivar identity than

root exudate metabolites ($R^2_{\text{leaf metabolite}}$ = 0.66, and $R^2_{\text{root exudates}}$ = 0.55; Figure 2). The first

two axes of the PCoA plot explained more than 30% of the variation for both metabolites.

Although most cultivars tended to cluster together, several cultivars stood out, suggesting

a unique and pronounced influence on the metabolite profiles. Furthermore, we found

that the alpha diversity of plant metabolites was significantly affected by cultivars (Table

S4). These results indicate that different potato cultivars exhibit distinct metabolites,

contributing to variations in metabolite composition among cultivars.

**Figure 2.** Distribution of metabolites in root exudates and leaves across different cultivars. Principal Coordinates Analysis (PCoA) based on mean Bray-Curtis dissimilarity of metabolite composition of root exudates (a) and leaves (b). Distinct potato cultivars are represented by different colours, with the error bars for each cultivar displayed in grey. PERMANOVA results are shown in the upper right corner of each panel, indicating variations in metabolite composition among different cultivars. $R^2$ quantifies the explained variation, and p-values are derived from 9999 permutations; *** denotes statistically significant p-values ($p$ = 0.001). Four outlier cultivars (Kama, Czapla, Fianna, Orlik Cs) were excluded to better illustrate the distribution of distinct cultivars. The distribution of the complete set of 51 potato cultivars is presented in Figure S4.

## 3.3 Plant cultivar functional groups

The analysis of plant cultivar growth characteristics, including shoot and root growth, leaf metabolites and root exudates, revealed that the distribution of these traits differed significantly among different potato cultivars, as evidenced by the results of the PERMANOVA analysis ($p$ = 0.001; Figure 3). Specifically, the first axis of the PCA biplot prominently reflected the impact of root exudate metabolite diversity and composition

(Figure 3). In contrast, plant performance metrics and leaf metabolite composition

mainly drove the second axis.

Based on the distribution of cultivars and variables across the first two axes, the 51 potato

cultivars were systematically categorised into four distinct functional groups (Figure 3).

Group A featured cultivars exhibiting high leaf metabolite diversity and plant biomass,

suggesting strong performance in leaf metabolic diversity and plant development. Group

B included cultivars with high root exudate diversity and plant length, indicating a focus

on below-ground metabolic processes and vertical growth of plants. Group C consisted of

cultivars characterised by complex leaf metabolite composition. These cultivars may have

a distinct composition of leaf metabolites compared to other groups, indicating specific

leaf metabolic profiles that differentiate them from other cultivars. Group D comprised

cultivars with low root exudate metabolite diversity and different metabolite

compositions. The summarised group details and cultivars can be found in Table S3.

**Figure 3.** Distribution of 51 potato cultivars based on plant performance, root exudates and leaf metabolite profiles. Principal Component Analysis (PCA) based on the covariance matrix, highlighting variables impacting the distribution of potato cultivars. Distinct potato cultivars are represented by different colours, with the error bars for each cultivar displayed in grey. The result of PERMANOVA is shown in the upper right, elucidating the cultivar's influence on the distribution. $R^2$ quantifies the explained variation, and p-values, derived from 9999 permutations, are denoted with *** for statistically significant results (p = 0.001). Letters A, B, C and D represent the four distinct functional groups.

## 3.4 Soil microbial communities

The analysis of microbial communities in bulk and rhizosphere soil revealed a significant impact of plant presence on microbial communities. At the diversity level, there was a significant difference in bacterial evenness between different soil compartments ($p$ =

0.001; Figure S5), whereas no significant effect on fungal diversity was observed.

Regarding the composition level, bacterial and fungal communities responded to the

presence of plants (Figure S6). Specifically, all bulk soil samples were found to cluster

together in bacterial communities, separate from the rhizosphere samples (Figure S6a).

Some potato cultivars' rhizosphere fungal communities were grouped distinctly apart

from the bulk soil (Figure S6b).

Comparison across the 51 potato cultivars revealed considerable variation and no

significant effect of cultivar on bacterial and fungal alpha diversities, except for the

cultivar effect on fungal evenness ($p$ = 0.03; Figure S7). Plant cultivars showed no

significant impact at the composition level but a trend towards influencing bacterial

community distribution ($p$ = 0.093 and $R^2$ = 0.34; Figure 4a). However, cultivars

exhibited significant variation in fungal community composition ($p$ = 0.001 and $R^2$ = 0.37;

Figure 4b), indicating that cultivars can significantly influence the fungal community

composition within their rhizosphere.

**Figure 4.** Rhizosphere microbial community composition. Principal Coordinates Analysis (PCoA) based on mean Bray-Curtis dissimilarity of community composition of bacteria (a) and fungi (b). Distinct potato cultivars are represented by different colours, with the error bars for each cultivar displayed in grey. PERMANOVA results in the upper left corner of each panel elucidate the influence of cultivars on community composition. $R^2$ quantifies the explained variation, and $p$-values are derived from 9999 permutations. The symbol *** denotes statistically significant $p$-values ($p$ = 0.001).

Grouping the cultivars in the functional groups determined by MIT revealed no significant differences in the alpha diversity of bacterial and fungal communities between the different functional groups ($p$ > 0.05; Figure S8). Similarly, the composition of rhizosphere microbial communities did not vary among the functional groups ($p_{bacteria}$ = 0.49 and $p_{fungi}$ = 0.14; Figure S9).

## 3.5 Correlation between plant traits, metabolites and rhizosphere microbiome

To evaluate how different plant traits relate to root-associated microbes and metabolite release, we correlated plant shoot growth, MITs (i.e., root biomass and root exudate

metabolites) with leaf metabolites and rhizosphere microbial communities of the 51

potato cultivars. Positive correlations were observed between plant biomass and leaf

metabolite richness. In contrast, negative correlations were identified between plant

shoot growth and leaf metabolite composition (Figure 5). Regarding the microbiome, we

observed positive correlations between rhizosphere bacterial species richness, root

biomass, and root-to-shoot biomass ratio (Figure 5). Similarly, rhizosphere fungal species

richness positively correlated with the root-to-shoot biomass ratio. However, the

composition of the rhizosphere microbial community showed no significant correlations

with root growth. Notably, the bacteria-to-fungi species richness ratio (B/F) was

positively associated with root and root-to-shoot biomass ratios. Additionally, we did not

observe a pronounced correlation between root exudate metabolites and the rhizosphere

microbial community (Figure 5).

We performed Mantel tests based on Bray-Curtis dissimilarity matrices to further

investigate the relationship between plant leaf metabolites and soil microbial

communities. Our analysis revealed a subtle yet statistically significant correlation

between bacterial community compositions and leaf metabolite profiles ($r = 0.10$, $p = $

$0.02$). In contrast, the Mantel test for fungi showed no significant correlation between

fungal community compositions and leaf metabolite profiles ($r = 0.04$, $p = 0.22$).

**Figure 5.** Correlation analysis between shoot biomass, MITs, leaf metabolite, and rhizosphere microbial community of the 51 potato cultivars. Correlations are based on the Spearman's rank correlation coefficient. Dark-coloured boxes indicate a significant correlation ($p < 0.05$), with the colour intensity reflecting the strength of the correlation coefficient. Red represents a positive correlation, while blue represents a negative correlation. The numbers displayed within the boxes represent the correlation coefficient. Significance levels are denoted as follows: * ($p < 0.05$), ** ($p < 0.01$), and *** ($p < 0.001$).

When examining the correlations between plant root biomass and rhizosphere bacterial community diversity across different functional groups, functional groups A and D showed significant positive correlations ($p < 0.05$; Figure S10). Functional groups B and C show no significant correlation between these variables. A positive correlation between leaf metabolite and bacterial Shannon index was found in functional group B ($p < 0.05$; Figure S11). No significant correlations were observed between leaf metabolite and fungal Shannon index regardless of functional groups ($p > 0.05$; Figure S11).

**3.6 MIT-selected cultivars and their characterisation**

The stepwise selection workflow and the chosen cultivars are visualised in Figure 6a. We

initially selected 148 potato cultivars, including the commercial cultivar Desiree as a

reference, from a pool of a thousand based on their resistance to pathogens *in silico*. These

148 cultivars were then further grown *in vitro*, from which 51 were selected based on plant

growth matrices for subsequent greenhouse experiments. For these 51 cultivars, plant

growth, leaf metabolites and root exudate metabolites were used to classify them into four

functional groups with distinct growth traits. This classification allowed us to identify ten

representative cultivars across four groups, covering a range of MIT values, including root

growth and exudate metabolites. This diverse range of MIT values allows us to better

understand how different combinations of these traits influence plant growth and

interactions in subsequent trials. Furthermore, we illustrated the distribution of the

rhizosphere community composition of the 51 cultivars in Figure 6b, which shows a

substantial separation in bacterial and fungal compositions among the MIT-selected

cultivars. These selected cultivars can undergo further exploration in real-world

conditions to evaluate their potential for interacting with beneficial soil microbiomes.

a Workflow of potato cultivar selection

b Distribution of bacterial and fungal compositions

**Figure 6.** Workflow of the potato cultivar selection by Sankey plot (a) and distribution of rhizosphere

bacteria and fungi community composition through Spearman correlation analysis (b). The first axis of

principal coordinate analysis (PCoA1) of microbial community beta-diversity (bacteria and fungi) serves as

the indicator of community composition. The error bars for each cultivar are displayed in grey. The colour

of selected cultivars corresponds to the functional groups. The cultivars that were not selected are depicted

in grey (b).

We conducted a comprehensive characterisation of the MIT-selected cultivars (Figure 7).

The one-way ANOVA results revealed variations among the selected 11 cultivars in terms

of plant metabolites, root development, and rhizosphere microbial community richness.

Specifically, Desiree and Pasja Pomorska from group A demonstrated higher leaf

metabolite richness and microbial diversity (Figure 7a, c, d, g, h). In contrast, in group B,

Krab and Salto showed superior microbial diversity performance (Figure 7c, d, g, h). In

comparison, Danuta from group D exhibited lower plant metabolite richness, root growth,

and bacterial diversity (Figure 7a-c, e-g). At the compositional level, there were no

significant cultivar effects on metabolite or microbial profiles, except for root exudate

metabolite composition (Figure 7i-l).

**Figure 7.** Characterisation of selected cultivars. Selected potato cultivars from different functional groups are represented by different colours, with the error bars for each cultivar displayed in grey. Letters in the upper two panels indicate significant differences across cultivars (Duncan post hoc test). In the lower right corner of the last panel plots, PERMANOVA results elucidate the influence of cultivars on community composition. $R^2$ quantifies the explained variation, and p-values are derived from 9999 permutations. A significance level is denoted as *** ($p < 0.001$).

## 4 DISCUSSION

Conventional breeding primarily harnesses genetic variation to achieve the desired traits of plants. However, the impact of a cultivar's genetic information on associated microbial organisms has yet to receive much consideration, particularly those associated with the rhizosphere. Here, we emphasise that modern breeding should consider the soil microbiome as a strategy to reduce its environmental footprint. Understanding the

importance of microbiome interactive traits (MITs), such as root traits and exudates, is

vital to comprehending how plants and microbiomes interact. In the future, this

understanding will contribute to informing the breeding process, providing valuable

insights for developing new microbiome-based cultivars.

**4.1 Correlation between plant traits, metabolites and rhizosphere**

**microbiome**

Our findings underscore the significant role of plant genetics in shaping both root exudate

and leaf metabolite profiles. We observed substantial cultivar-dependent variation in root

exudate metabolite composition consistent with previous studies. This genetic influence

also extends to leaf metabolites, aligning with earlier research on rice (Schaarschmidt et

al., 2020). These results collectively demonstrate that the genetic makeup of different

plant cultivars significantly influences the metabolite composition of different plant

tissues.

We observed a positive correlation between leaf metabolite richness and plant biomass, a

relationship that can be explained through considerations of resource availability and

plant growth strategies. Environmental conditions and functional traits influence

biomass allocation among plant organs, such as leaves and roots, influencing many

growth processes (Mensah et al., 2016; Poorter et al., 2012). In optimal environmental

conditions, biomass allocation (primarily above-ground to compete for light) can enhance

the diversity of leaf metabolites instead of root metabolites (Chapin et al., 2005). The

crucial role of plant secondary metabolites in host defence may also explain this

correlation. These compounds protect hosts against herbivores, pathogens, and (biotic)

stresses (Anjali et al., 2023; Divekar et al., 2022; Yadav et al., 2021; Zaynab et al., 2018).

Plants with higher biomass may invest more in defence mechanisms, producing a wider

range of leaf metabolites for protection. This establishes positive feedback loops, wherein

increased biomass boosts photosynthetic activity and energy production, consequently

supporting the synthesis of more diverse metabolites.

Previous studies have demonstrated that plant root traits are crucial in regulating

rhizosphere microbial communities (Szoboszlay et al., 2015; Wan et al., 2021). This

relationship is further supported by Eisenhauer et al. (2017), who showed that microbial

diversity increases with increasing root biomass and exudate amount. Our results align

with these findings, revealing positive correlations between plant root biomass and root-

to-shoot biomass ratio, rhizosphere bacterial diversity, and bacteria-to-fungi species

richness ratio (B/F). A more diverse microbial community, in turn, can enhance soil

nutrient cycling and availability (Jiao et al., 2021), potentially promoting plant growth.

Previous studies have demonstrated that root exudates significantly influence bacterial

and fungal communities in the rhizosphere (Hartmann et al., 2009). In contrast, our

study did not find a significant correlation between root exudate metabolites and the

rhizosphere microbial community. This unexpected result may be because we collected

root exudate metabolites from 51 potato cultivars under *in vitro* conditions, while the

microbial community data were obtained from a greenhouse experiment. The differences

in substrates between these environments may have influenced the root exudate profiles,

leading to variances that could explain the lack of correlation.

Given the complexity of collecting root exudates from soil plants, we collected leaf
metabolites in our current experiment to explore the relationship between plant
metabolites and the soil microbial community. The marginal but positive correlation
observed between leaf metabolites and the rhizosphere bacterial community suggests a
potential influence of leaf metabolites on the composition of rhizosphere bacteria. This
finding indicates a possible link between above-ground plant tissues and below-ground
microbial communities, aligning with the holobiont concept, in which plants and their
associated microorganisms are viewed as a holistic ecological unit (Vandenkoornhuyse et
al., 2015).

Recent studies further support this interconnection. Korenblum et al. (2020)
demonstrated that the composition of the rhizosphere microbial community affects the
metabolomes and transcriptomes of tomatoes' leaves and roots. Earlier research showed
that rhizobacterium Bacillus strains can influence photosynthesis, leaf growth, and
overall plant phenotypes by producing phytohormones or volatile organic compounds
(Pang et al., 2021), potentially impacting leaf metabolomes. These findings collectively
emphasise the concept of metabolites as primary mediators regulating plant-microbiome
interactions within the holobiont framework (Carper et al., 2022). As described, the
transport of leaf-produced metabolites to the roots via the phloem (Broussard et al., 2023)
suggests a potential mechanism for how above-ground metabolites might influence root
exudate patterns and, consequently, the rhizosphere microbiome. However, it's
important to note that our study only demonstrates correlation, not causation. The
complex interactions between leaf metabolites, root exudates, and microbial communities
require further investigation to elucidate the holobiont concept fully.

**4.2 Enhancing plant-microbiome interactions for sustainable agriculture**

The soil microbiome promotes plant growth by promoting carbon, nutrient, and phosphorus cycling (Hartmann and Six, 2022). It also contributes to plant resistance by producing hormones that protect against abiotic and biotic stress (Eichmann et al., 2021). Despite its critical functions, the soil microbiome has received limited attention in conventional breeding (Mitter et al., 2019; Wei and Jousset, 2017). Compounding this issue is the fact that conventional agricultural management not only negatively affects the environment but also substantially impacts the soil microbiome (Longepierre et al., 2021). This dual impact increases the decoupling between plants and soil microbiomes (Huang et al., 2019; Spor et al., 2020).

Studies addressing the role of plants in regulating their associated microbiome remain relatively limited (Wei and Jousset, 2017). More precisely, plant roots, serving as the primary interface for interaction with soil microbes, are underexplored in plant breeding (Herms et al., 2022; Reinhold-Hurek et al., 2015). Here, we consider morphological root characteristics and root exudate metabolites as microbiome interactive traits (MITs) to explore their interaction with the rhizosphere microbiome. We aim to supply a strategy for breeding that considers plant-associated microbiota.

Indeed, the positive correlation between root growth and rhizosphere microbial diversity indicates that MITs can aid in identifying plant cultivars with the potential to interact effectively with root-associated microbiomes. Cultivars exhibiting high MITs are likely to select for a more diverse rhizosphere microbiome, which in turn can lead to enhanced plant growth through the support of beneficial microorganisms. Identifying the genes

associated with beneficial microbiomes in modern cultivars and using them in selective

breeding efforts to achieve microbial-assisted cultivars can serve as a new plant breeding

strategy. This approach represents a promising avenue for sustainable agriculture, as it

harnesses the power of beneficial microorganisms to improve crop performance while

reducing the need for chemical inputs.

**4.3 Integrating additional root traits and phyllosphere microbiome in future**

**studies**

We suggest expanding future research beyond the MITs examined in this study to include

a broader range of root phenotypic traits. While the current study focuses on root biomass

and length, future investigations should include root diameter, surface area, and root type.

Although less studied, evidence suggests that fine roots with smaller diameters have

larger surface areas, potentially recruiting a greater diversity and abundance of microbes

through enhanced nutrient and metabolite exchange (Saleem et al., 2018; Wan et al.,

2021). Pérez-Jaramillo et al. (2017) linked root types (thin or thick) to specific bacterial

phyla, highlighting the importance of root morphology in shaping microbial communities.

Additionally, the spatial distribution of microbial communities along the root should be

considered. Kawasaki et al. (2016) observed that the functional genes detected in

microorganisms near the root tip were distinct from those isolated near the root base.

Collectively, these various root phenotypic traits should be considered to improve plant-

microbiome interactions.

In addition to expanding our focus on root phenotypic traits, we propose incorporating

the phyllosphere microbiome into future studies. The phyllosphere microbiome, which

includes microorganisms inhabiting the above-ground parts of plants, plays a crucial role

in plant health and function (Thapa and Prasanna, 2018; Vorholt, 2012). Previous studies

have demonstrated that host genotypes significantly influence the composition of

phyllosphere microbial communities (Bodenhausen et al., 2014; Thapa et al., 2017). The

phyllosphere microbiome is involved in nitrogen fixation (Abadi et al., 2021), enhancing

stress tolerance (Etemadi et al., 2018; Stone et al., 2018), and suppressing plant diseases

(Fan et al., 2019; Das et al., 2023). Additionally, they can regulate plant growth through

the production of plant hormones (Stone et al., 2018). These diverse functions highlight

the importance of the phyllosphere microbiome in plant health and productivity.

By considering the phyllosphere microbiome together with the rhizosphere microbiome

and plant metabolites, we can establish a more comprehensive understanding of the plant

holobiont. This approach will allow us to bridge the gap between the plant's above- and

below-ground components. By harnessing the functions of phyllosphere and rhizosphere

microbiomes, we may enhance crop yields, improve plant resilience, and reduce reliance

on chemical inputs, ultimately contributing to more sustainable agricultural systems.

**5 CONCLUSION**

This study underscores the significant impact of plant cultivars on leaf metabolites and

root exudate metabolites. We also observe a positive correlation between leaf metabolites

and rhizosphere bacterial community; further studies are needed to verify the causation

and to involve root exudates to expand our knowledge of the holobiont framework. We

systematically selected potato cultivars to identify those with diverse microbiome

interactive traits (MITs). We lay the foundation for further studies to evaluate the
performance of MIT-selected cultivars in field conditions. This is needed to provide a
promising strategy for future breeding programs, including identifying gene markers
associated with a beneficial microbiome and utilising these genes to increase plant-
microbiome interactions. This breeding strategy could promote host growth while
reducing the reliance on synthetic chemicals in conventional agriculture. Finally, we
suggest integrating additional root phenotypic traits and the phyllosphere microbiome in
future studies to establish a more comprehensive understanding of the plant holobiont,
which can benefit plant-microbiome interactions.

## 610 Availability of Data and Materials

611 The metadata and datasets used for the bioinformatic analyses are available at the
612 following link: https://github.com/tianci-zhao/potatoMETAbiome-Greenhouse-
613 Experiment.

## 615 ACKNOWLEDGEMENTS

This study was funded by ERA-NET Cofund SusCrop project potatoMETAbiome, which
is supported by the European Union's Horizon 2020 research and innovation program
(grant agreement No 771134; French National Agency, grant number -ANR-18-SUSC-
0001) and part of the Joint Programming Initiative on Agriculture, Food Security and
Climate Change (FACCE-JPI). We thank Jolanda K. Brons, Jan Veldsink, Xipeng Liu,

Pina Brinker, Panji Cahya Mawarda, Edisa Garcia Hernandez, Georgia Voulgari, and

Florian Fischer for their help in the lab. We thank Ines Fehrle and Jessica Alpers for their

excellent assistance with the GC-MS measurements. We want to thank Marie Simonin

and Ulisses Nunes da Rocha for their insightful discussion on data analysis. T.Z. was

supported by a scholarship from the China Scholarship Council (CSC) and the University

of Groningen scholarship program. We thank PGTB for sequencing (Genome

Transcriptome Platform of Bordeaux). We thank the Center for Information Technology

at the University of Groningen for providing access to the Hábrók high-performance

computing cluster.

**631    Author contributions**

J.F.S., S.N.V., J.T.M.E., A.E., and E.A. designed the experiment; T.Z. and S.N.V.

performed the experiment, laboratory work and data analysis; X.J. participated in data

analysis; E.A. and R.G. contributed to plant sampling; D.M. and K.T. provided the potato

materials; A.E., S.S., J.K., and E.Z. measured and analysed plant metabolites. T.Z. drafted

the manuscript, with all authors contributing to its modification and approval of the final

version.

**639    Declaration of competing interest**

640    The authors declare that they have no known competing financial interests or personal

641    relationships that could have appeared to influence the work reported in this paper.

## Supporting information

**Figure S1.** Root dry weight and root-to-shoot ratio of 51 potato cultivars *in vitro* experiment. Each colour represents a distinct potato cultivar. The upper left corner of each plot displays one-way ANOVA results, where the F-value explains the variation among different cultivars, and the *p*-value indicates the statistical relationship among cultivars.

**Figure S2.** Shoot growth analysis of 51 potato cultivars. The upper panel displays the shoot length, while the lower panel illustrates the shoot dry weight. Each colour represents a distinct potato cultivar. The upper right corner of each plot displays one-way ANOVA results, where the F-value explains the variation among different cultivars, and the *p*-value indicates the statistical relationship among cultivars.

**Figure S3.** Root-to-shoot ratio analysis of 51 potato cultivars. The upper panel displays the root-to-shoot length ratio, while the lower panel illustrates the root-to-shoot dry weight ratio. Each colour represents a distinct potato cultivar. The upper right corner of each plot displays one-way ANOVA results, where the F-value explains the variation among different cultivars, and the *p*-value indicates the statistical relationship among cultivars.

**Figure S4.** Impact of 51 potato cultivars on the composition of plant root exudates (a) and leaf (b) metabolites. A Principal Coordinates Analysis (PCoA) based on Bray-Curtis dissimilarity was performed to visualise the composition. The metabolite dissimilarities of root exudates and leaf are depicted separately on the left and right. Distinct potato cultivars are represented by different colours, with the error bars for each cultivar displayed in grey. PERMANOVA (Adonis) results in the upper right corner of each panel elucidate the influence of cultivars on metabolite composition. $R^2$ quantifies the explained variation, and *p*-values are derived from 9999 permutations.

**Figure S5.** The alpha-diversity of the bacterial community in bulk and rhizosphere soil, displayed by species richness and evenness. Different colours represent bulk samples from the beginning of the

experiment (Bulk_D0), at harvest (Bulk_D36), and rhizosphere samples of 51 cultivars. The lower right

corner of each plot displays one-way ANOVA results, where the F-value explains the variation among

different soil compartments (bulk and rhizosphere), and the $p$-value indicates the statistical relationship.

**Figure S6.** The microbial community composition in bulk and rhizosphere soil. Principal Coordinates

Analysis (PCoA) based on Bray-Curtis dissimilarity was performed to visualise the community composition

of bacteria (a) and fungi (b). Different shapes represent different soil compartments (bulk and rhizosphere).

Different colours represent bulk samples from the beginning of the experiment (Bulk_D0), at harvest

(Bulk_D36), and different potato cultivars, with the error bars for each cultivar displayed in grey.

PERMANOVA results in the lower left corner of each panel elucidate the influence of soil compartments on

community composition. $R^2$ quantifies the explained variation, and $p$-values are derived from 9999

permutations.

**Figure S7.** Rhizosphere microbial community evenness of 51 potato cultivars. The upper panel is bacterial

evenness, and the lower panel is fungal evenness. Different colours indicate different cultivars. The lower

right corner of each plot displays one-way ANOVA results, where the F-value explains the variation among

different cultivars, and the $p$-value indicates the statistical relationship among cultivars.

**Figure S8.** Rhizosphere microbial community species richness. The upper panel is bacterial richness, and

the lower panel is fungal richness. Different colours indicate different cultivar functional groups. The "ns"

indicates no significant influence of groups on microbial alpha diversity (one-way ANOVA).

**Figure S9.** Rhizosphere microbial community composition. Principal Coordinates Analysis (PCoA) based

on Bray-Curtis dissimilarity was performed to visualise the community dissimilarities of bacteria (a) and

fungi (b). Distinct potato cultivar functional groups are represented by different colours, with the error bars

for each cultivar displayed in grey. PERMANOVA results in the upper right corner of each panel elucidate

the influence of groups on community composition. $R^2$ quantifies the explained variation, and $p$-values are

derived from 9999 permutations.

**Figure S10.** Correlation between plant root biomass and rhizosphere bacterial alpha diversity (species

richness and Shannon diversity) across different functional groups. Displayed by functional groups A, B, C

and D. The Spearman correlation assessed the relationship, with $R^2$ indicating the strength of the

correlation. Y is the regression equation, and a $p$-value < 0.05 represents a significant correlation between

the variables.

**Figure S11.** Correlation of leaf metabolite and microbial community Shannon index across different

functional groups. Displayed by functional groups A, B, C and D. The Spearman correlation assessed the

relationship, with $R^2$ indicating the strength of the correlation. Y is the regression equation, and a $p$-value

< 0.05 represents a significant correlation between the variables.

**Table S1.** Background information on 148 selected cultivars for the *in vitro* experiment.

**Table S2.** Plant growth data of 148 potato cultivars from *in vitro* experiment.

**Table S3.** Functional groups of potato cultivars categorised based on plant growth and metabolite profiles.

**Table S4.** The one-way analysis of variance (ANOVA) shows the influence of cultivar on the plant

metabolite alpha diversity.

**Data S1.** Leaf tissue metabolites data in greenhouse experiment

**Data S2.** Root exudate metabolites data *in vitro* experiment

**Data S3.** Plant performance in greenhouse experiment

**Data S4.** Rhizosphere microbial feature tables

## REFERENCES

Abadi, V.A.J.M., Sepehri, M., Rahmani, H.A., Dolatabad, H.K., Shamshiripour, M., Khatabi, B., 2021. Diversity and abundance of culturable nitrogen-fixing bacteria in the phyllosphere of maize. Journal of Applied Microbiology 131, 898–912. https://doi.org/10.1111/jam.14975

AL-Ani, M.A.M., Hmoshi, R.M., Kanaan, I.A., Thanoon, A.A., 2019. Effect of pesticides on soil microorganisms. J. Phys.: Conf. Ser. 1294, 072007. https://doi.org/10.1088/1742-6596/1294/7/072007

Anjali, Kumar, S., Korra, T., Thakur, R., Arutselvan, R., Kashyap, A.S., Nehela, Y., Chaplygin, V., Minkina, T., Keswani, C., 2023. Role of plant secondary metabolites in defence and transcriptional regulation in response to biotic stress. Plant Stress 8, 100154. https://doi.org/10.1016/j.stress.2023.100154

Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A., De Wit, P., Sánchez-García, M., Ebersberger, I., de Sousa, F., Amend, A., Jumpponen, A., Unterseher, M., Kristiansson, E., Abarenkov, K., Bertrand, Y.J.K., Sanli, K., Eriksson, K.M., Vik, U., Veldre, V., Nilsson, R.H., 2013. Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. Methods in Ecology and Evolution 4, 914–919. https://doi.org/10.1111/2041-210X.12073

Bodenhausen, N., Bortfeld-Miller, M., Ackermann, M., Vorholt, J.A., 2014. A Synthetic Community Approach Reveals Plant Genotypes Affecting the Phyllosphere Microbiota. PLOS Genetics 10, e1004283. https://doi.org/10.1371/journal.pgen.1004283

Bokulich, N.A., Kaehler, B.D., Rideout, J.R., Dillon, M., Bolyen, E., Knight, R., Huttley, G.A., Gregory Caporaso, J., 2018. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. Microbiome 6, 1–17. https://doi.org/10.1186/s40168-018-0470-z

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A., Brislawn, C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodríguez, A.M., Chase, J., Cope, E.K., Da Silva, R., Diener, C., Dorrestein, P.C., Douglas, G.M., Durall, D.M., Duvallet, C., Edwardson, C.F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J.M., Gibbons, S.M., Gibson, D.L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H., Huttenhower, C., Huttley, G.A., Janssen, S., Jarmusch, A.K., Jiang, L., Kaehler, B.D., Kang, K.B., Keefe, C.R., Keim, P., Kelley, S.T., Knights, D., Koester, I., Kosciolek, T., Kreps, J., Langille, M.G.I., Lee, J., Ley, R., Liu, Y.-X., Loftfield, E., Lozupone, C., Maher, M., Marotz, C., Martin, B.D., McDonald, D., McIver, L.J., Melnik, A.V., Metcalf, J.L., Morgan, S.C., Morton, J.T., Naimey, A.T., Navas-Molina, J.A., Nothias, L.F., Orchanian, S.B., Pearson, T., Peoples, S.L., Petras, D., Preuss, M.L., Pruesse, E., Rasmussen, L.B., Rivers, A., Robeson, M.S., Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A., Sinha, R., Song, S.J., Spear, J.R., Swafford, A.D., Thompson, L.R., Torres, P.J., Trinh, P., Tripathi, A., Turnbaugh, P.J., Ul-Hasan, S., van der Hooft, J.J.J., Vargas, F., Vázquez-Baeza, Y., Vogtmann, E., von Hippel, M., Walters, W., Wan, Y., Wang, M., Warren, J., Weber, K.C., Williamson, C.H.D., Willis, A.D., Xu, Z.Z., Zaneveld, J.R.,

Zhang, Y., Zhu, Q., Knight, R., Caporaso, J.G., 2019. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol 37, 852–857. https://doi.org/10.1038/s41587-019-0209-9

Breseghello, F., Coelho, A.S.G., 2013. Traditional and Modern Plant Breeding Methods with Examples in Rice (Oryza sativa L.). J. Agric. Food Chem. 61, 8277–8286. https://doi.org/10.1021/jf305531j

Broussard, L., Abadie, C., Lalande, J., Limami, A.M., Lothier, J., Tcherkez, G., 2023. Phloem Sap Composition: What Have We Learnt from Metabolomics? Int J Mol Sci 24, 6917. https://doi.org/10.3390/ijms24086917

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: High resolution sample inference from Illumina amplicon data. Nat Methods 13, 581–583. https://doi.org/10.1038/nmeth.3869

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S.M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J.A., Smith, G., Knight, R., 2012. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J 6, 1621–1624. https://doi.org/10.1038/ismej.2012.8

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J., Fierer, N., Knight, R., 2011. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proceedings of the National Academy of Sciences 108, 4516–4522. https://doi.org/10.1073/pnas.1000080107

Carper, D.L., Appidi, M.R., Mudbhari, S., Shrestha, H.K., Hettich, R.L., Abraham, P.E., 2022. The Promises, Challenges, and Opportunities of Omics for Studying the Plant Holobiont. Microorganisms 10, 2013. https://doi.org/10.3390/microorganisms10102013

Chapin, F.S., Matson, P.A., Mooney, H.A., Chapin, M.C., 2005. Principles of terrestrial ecosystem ecology, Nachdr. ed. Springer, New York, NY.

Clouse, K.M., Wagner, M.R., 2021. Plant Genetics as a Tool for Manipulating Crop Microbiomes: Opportunities and Challenges. Front Bioeng Biotechnol 9, 567548. https://doi.org/10.3389/fbioe.2021.567548

da Silva, J.L., Mendes, L.W., Rocha, S.M.B., Antunes, J.E.L., Oliveira, L.M. de S., Melo, V.M.M., Oliveira, F.A.S., Pereira, A.P. de A., Costa, G. do N., da Silva, V.B., Gomes, R.L.F., de Alcantara Neto, F., Lopes, A.C. de A., Araujo, A.S.F., 2023. Domestication of Lima Bean (Phaseolus lunatus) Changes the Microbial Communities in the Rhizosphere. Microb Ecol 85, 1423–1433. https://doi.org/10.1007/s00248-022-02028-2

Das, T., Bhattacharyya, A., Bhar, A., 2023. Linking Phyllosphere and Rhizosphere Microbiome to the Plant–Insect Interplay: The New Dimension of Tripartite Interaction. Physiologia 3, 129–144. https://doi.org/10.3390/physiologia3010011

Diagne, N., Ngom, M., Djighaly, P.I., Fall, D., Hocher, V., Svistoonoff, S., 2020. Roles of Arbuscular Mycorrhizal Fungi on Plant Growth and Performance: Importance in Biotic and Abiotic Stressed Regulation. Diversity 12, 370. https://doi.org/10.3390/d12100370

Divekar, P.A., Narayana, S., Divekar, B.A., Kumar, R., Gadratagi, B.G., Ray, A., Singh, Achuit Kumar, Rani, V., Singh, V., Singh, Akhilesh Kumar, Kumar, A., Singh, R.P., Meena, R.S., Behera, T.K., 2022. Plant Secondary Metabolites as Defense Tools against Herbivores for Sustainable Crop Protection. Int J Mol Sci 23, 2690. https://doi.org/10.3390/ijms23052690

Dixon, P., 2003. VEGAN, a package of R functions for community ecology. Journal of Vegetation Science 14, 927–930. https://doi.org/10.1111/j.1654-1103.2003.tb02228.x

Eichmann, R., Richards, L., Schäfer, P., 2021. Hormones as go-betweens in plant microbiome assembly. The Plant Journal 105, 518–541. https://doi.org/10.1111/tpj.15135

Eisenhauer, N., Lanoue, A., Strecker, T., Scheu, S., Steinauer, K., Thakur, M.P., Mommer, L., 2017. Root biomass and exudates link plant diversity with soil bacterial and fungal biomass. Sci Rep 7, 44641. https://doi.org/10.1038/srep44641

Erban, A., Martinez-Seidel, F., Rajarathinam, Y., Dethloff, F., Orf, I., Fehrle, I., Alpers, J., Beine-Golovchuk, O., Kopka, J., 2020. Multiplexed Profiling and Data Processing Methods to Identify Temperature-Regulated Primary Metabolites Using Gas Chromatography Coupled to Mass Spectrometry, in: Hincha, D.K., Zuther, E. (Eds.), Plant Cold Acclimation, Methods in Molecular Biology. Springer US, New York, NY, pp. 203–239. https://doi.org/10.1007/978-1-0716-0660-5_15

Etemadi, M., Zuther, E., Müller, H., Hincha, D.K., Berg, G., 2018. Ecotype-Dependent Response of Bacterial Communities Associated with Arabidopsis to Cold Acclimation. Phytobiomes Journal 2, 3–13. https://doi.org/10.1094/PBIOMES-04-17-0015-R

Fan, X., Matsumoto, H., Wang, Y., Hu, Y., Liu, Y., Fang, Hongda, Nitkiewicz, B., Lau, S.Y.L., Wang, Q., Fang, Hua, Wang, M., 2019. Microenvironmental Interplay Predominated by Beneficial Aspergillus Abates Fungal Pathogen Incidence in Paddy Environment. Environ. Sci. Technol. 53, 13042–13052. https://doi.org/10.1021/acs.est.9b04616

Fröhlich, A., Buddrus-Schiemann†, K., Durner, J., Hartmann, A., von Rad, U., 2012. Response of barley to root colonization by Pseudomonas sp. DSMZ 13134 under laboratory, greenhouse, and field conditions. Journal of Plant Interactions 7, 1–9. https://doi.org/10.1080/17429145.2011.597002

Gordon, A., Hannon, G., 2010. FASTX-Toolkit [WWW Document]. URL http://hannonlab.cshl.edu/fastx_toolkit/ (accessed 3.8.24).

Gweon, H.S., Oliver, A., Taylor, J., Booth, T., Gibbs, M., Read, D.S., Griffiths, R.I., Schonrogge, K., 2015. PIPITS : an automated pipeline for analyses of fungal internal transcribed spacer sequences from the I llumina sequencing platform. Methods Ecol Evol 6, 973–980. https://doi.org/10.1111/2041-210X.12399

Hartmann, A., Schmid, M., Tuinen, D. van, Berg, G., 2009. Plant-driven selection of microbes. Plant Soil 321, 235–257. https://doi.org/10.1007/s11104-008-9814-y

Hartmann, M., Six, J., 2022. Soil structure and microbiome functions in agroecosystems. Nat Rev Earth Environ. https://doi.org/10.1038/s43017-022-00366-w

Herms, C.H., Hennessy, R.C., Bak, F., Dresbøll, D.B., Nicolaisen, M.H., 2022. Back to our roots: exploring the role of root morphology as a mediator of beneficial plant–microbe interactions. Environmental Microbiology 24, 3264–3272. https://doi.org/10.1111/1462-2920.15926

Huang, R., McGrath, S.P., Hirsch, P.R., Clark, I.M., Storkey, J., Wu, L., Zhou, J., Liang, Y., 2019. Plant-microbe networks in soil are weakened by century-long use of inorganic fertilizers. Microb Biotechnol 12, 1464–1475. https://doi.org/10.1111/1751-7915.13487

Hummel, J., Strehmel, N., Selbig, J., Walther, D., Kopka, J., 2010. Decision tree supported substructure prediction of metabolites from GC-MS profiles. Metabolomics 6, 322–333. https://doi.org/10.1007/s11306-010-0198-7

Iannucci, A., Canfora, L., Nigro, F., De Vita, P., Beleggia, R., 2021. Relationships between root
morphology, root exudate compounds and rhizosphere microbial community in durum
wheat. Applied Soil Ecology 158, 103781. https://doi.org/10.1016/j.apsoil.2020.103781
Jiao, S., Peng, Z., Qi, J., Gao, J., Wei, G., 2021. Linking Bacterial-Fungal Relationships to Microbial
Diversity and Soil Nutrient Cycling. mSystems 6, 10.1128/msystems.01052-20.
https://doi.org/10.1128/msystems.01052-20
Jost, L., 2006. Entropy and diversity. Oikos 113, 363–375. https://doi.org/10.1111/j.2006.0030-
1299.14714.x
Kawasaki, A., Donn, S., Ryan, P.R., Mathesius, U., Devilla, R., Jones, A., Watt, M., 2016. Microbi-
ome and Exudates of the Root and Rhizosphere of Brachypodium distachyon, a Model for
Wheat. PLOS ONE 11, e0164533. https://doi.org/10.1371/journal.pone.0164533
Kõljalg, U., Nilsson, R.H., Abarenkov, K., Tedersoo, L., Taylor, A.F.S., Bahram, M., Bates, S.T., Bruns,
T.D., Bengtsson-Palme, J., Callaghan, T.M., Douglas, B., Drenkhan, T., Eberhardt, U., Due-
ñas, M., Grebenc, T., Griffith, G.W., Hartmann, M., Kirk, P.M., Kohout, P., Larsson, E., Lin-
dahl, B.D., Lücking, R., Martín, M.P., Matheny, P.B., Nguyen, N.H., Niskanen, T., Oja, J.,
Peay, K.G., Peintner, U., Peterson, M., Põldmaa, K., Saag, L., Saar, I., Schüßler, A., Scott,
J.A., Senés, C., Smith, M.E., Suija, A., Taylor, D.L., Telleria, M.T., Weiss, M., Larsson, K.-H.,
2013. Towards a unified paradigm for sequence-based identification of fungi. Molecular
Ecology 22, 5271–5277. https://doi.org/10.1111/mec.12481
Kopka, J., Schauer, N., Krueger, S., Birkemeyer, C., Usadel, B., Bergmüller, E., Dörmann, P., Weck-
werth, W., Gibon, Y., Stitt, M., Willmitzer, L., Fernie, A.R., Steinhauser, D., 2005.
GMD@CSB.DB: the Golm Metabolome Database. Bioinformatics 21, 1635–1638.
https://doi.org/10.1093/bioinformatics/bti236
Korenblum, E., Dong, Y., Szymanski, J., Panda, S., Jozwiak, A., Massalha, H., Meir, S., Rogachev, I.,
Aharoni, A., 2020. Rhizosphere microbiome mediates systemic root metabolite exudation
by root-to-root signaling. Proceedings of the National Academy of Sciences 117, 3874–
3883. https://doi.org/10.1073/pnas.1912130117
Lê, S., Josse, J., Husson, F., 2008. FactoMineR: An R Package for Multivariate Analysis. Journal of
Statistical Software 25, 1–18. https://doi.org/10.18637/jss.v025.i01
Lisec, J., Römisch-Margl, L., Nikoloski, Z., Piepho, H.-P., Giavalisco, P., Selbig, J., Gierl, A., Will-
mitzer, L., 2011. Corn hybrids display lower metabolite variability and complex metabolite
inheritance patterns. The Plant Journal 68, 326–336. https://doi.org/10.1111/j.1365-
313X.2011.04689.x
Lisec, J., Schauer, N., Kopka, J., Willmitzer, L., Fernie, A.R., 2006. Gas chromatography mass spec-
trometry–based metabolite profiling in plants. Nature Protocols 1, 387–396.
https://doi.org/10.1038/nprot.2006.59
Longepierre, M., Widmer, F., Keller, T., Weisskopf, P., Colombi, T., Six, J., Hartmann, M., 2021.
Limited resilience of the soil microbiome to mechanical compaction within four growing
seasons of agricultural management. ISME COMMUN. 1, 1–13.
https://doi.org/10.1038/s43705-021-00046-8
Luedemann, A., Strassburg, K., Erban, A., Kopka, J., 2008. TagFinder for the quantitative analysis
of gas chromatography--mass spectrometry (GC-MS)-based metabolite profiling experi-
ments. Bioinformatics 24, 732–737. https://doi.org/10.1093/bioinformatics/btn023
Lynch, J., 1995. Root Architecture and Plant Productivity. Plant Physiol 109, 7–13.

Mensah, S., Kakaï, R.G., Seifert, T., 2016. Patterns of biomass allocation between foliage and
woody structure: the effects of tree size and specific functional traits. Annals of Forest
Research 59, 49–60. https://doi.org/10.15287/afr.2016.458

Michalak, K., Chrzanowska, M., 2017. Evaluation of Resistance to Potato Virus Y of Cultivars in
Pre-Registration (Statutory) Trials. Plant Breeding and Seed Science 75, 63–70.
https://doi.org/10.1515/plass-2017-0009

Mitter, B., Brader, G., Pfaffenbichler, N., Sessitsch, A., 2019. Next generation microbiome appli-
cations for crop production — limitations and the need of knowledge-based solutions.
Current Opinion in Microbiology 49, 59–65. https://doi.org/10.1016/j.mib.2019.10.006

Nilsson, R.H., Tedersoo, L., Ryberg, M., Kristiansson, E., Hartmann, M., Unterseher, M., Porter,
T.M., Bengtsson-Palme, J., Walker, D.M., De Sousa, F., Gamper, H.A., Larsson, E., Larsson,
K.-H., Kõljalg, U., Edgar, R.C., Abarenkov, K., 2015. A Comprehensive, Automatically Up-
dated Fungal ITS Sequence Dataset for Reference-Based Chimera Control in Environmen-
tal Sequencing Efforts. Microbes and environments 30, 145–150.
https://doi.org/10.1264/jsme2.ME14121

Pang, Z., Chen, J., Wang, T., Gao, C., Li, Z., Guo, L., Xu, J., Cheng, Y., 2021. Linking Plant Secondary
Metabolites and Plant Microbiomes: A Review. Frontiers in Plant Science 12.

Paradis, E., Claude, J., Strimmer, K., 2004. APE: Analyses of Phylogenetics and Evolution in R lan-
guage. Bioinformatics 20, 289–290. https://doi.org/10.1093/bioinformatics/btg412

Pérez-Jaramillo, J.E., Carrión, V.J., Bosse, M., Ferrão, L.F.V., De Hollander, M., Garcia, A.A.F., Ramí-
rez, C.A., Mendes, R., Raaijmakers, J.M., 2017. Linking rhizosphere microbiome composi-
tion of wild and domesticated Phaseolus vulgaris to genotypic and root phenotypic traits.
ISME J 11, 2244–2257. https://doi.org/10.1038/ismej.2017.85

Poorter, H., Niklas, K.J., Reich, P.B., Oleksyn, J., Poot, P., Mommer, L., 2012. Biomass allocation to
leaves, stems and roots: meta-analyses of interspecific variation and environmental con-
trol. New Phytologist 193, 30–50. https://doi.org/10.1111/j.1469-8137.2011.03952.x

Raaijmakers, J.M., Kiers, E.T., 2022. Rewilding plant microbiomes. Science 378, 599–600.
https://doi.org/10.1126/science.abn6350

Reinhold-Hurek, B., Bünger, W., Burbano, C.S., Sabale, M., Hurek, T., 2015. Roots Shaping Their
Microbiome: Global Hotspots for Microbial Activity. Annu. Rev. Phytopathol. 53, 403–424.
https://doi.org/10.1146/annurev-phyto-082712-102342

Revelle, W., 2024. Procedures for Psychological, Psychometric, and Personality Research.

Rodriguez, P.A., Rothballer, M., Chowdhury, S.P., Nussbaumer, T., Gutjahr, C., Falter-Braun, P.,
2019. Systems Biology of Plant-Microbiome Interactions. Molecular Plant 12, 804–821.
https://doi.org/10.1016/j.molp.2019.05.006

Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open source
tool for metagenomics. PeerJ 4, e2584. https://doi.org/10.7717/peerj.2584

Rossmann, M., Pérez-Jaramillo, J.E., Kavamura, V.N., Chiaramonte, J.B., Dumack, K., Fiore-Donno,
A.M., Mendes, L.W., Ferreira, M.M.C., Bonkowski, M., Raaijmakers, J.M., Mauchline, T.H.,
Mendes, R., 2020. Multitrophic interactions in the rhizosphere microbiome of wheat:
from bacteria and fungi to protists. FEMS Microbiology Ecology 96, fiaa032.
https://doi.org/10.1093/femsec/fiaa032

Saleem, M., Law, A.D., Sahib, M.R., Pervaiz, Z.H., Zhang, Q., 2018. Impact of root system architecture on rhizosphere and root microbiome. Rhizosphere 6, 47–51. https://doi.org/10.1016/j.rhisph.2018.02.003

Schaarschmidt, S., Lawas, L.M.F., Glaubitz, U., Li, X., Erban, A., Kopka, J., Jagadish, S.V.K., Hincha, D.K., Zuther, E., 2020. Season Affects Yield and Metabolic Profiles of Rice (Oryza sativa) under High Night Temperature Stress in the Field. IJMS 21, 3187. https://doi.org/10.3390/ijms21093187

Sjoberg, D., 2023. ggsankey.

Spor, A., Roucou, A., Mounier, A., Bru, D., Breuil, M.-C., Fort, F., Vile, D., Roumet, P., Philippot, L., Violle, C., 2020. Domestication-driven changes in plant traits associated with changes in the assembly of the rhizosphere microbiota in tetraploid wheat. Sci Rep 10, 12234. https://doi.org/10.1038/s41598-020-69175-9

Stone, B., Weingarten, E., Jackson, C., 2018. The role of the phyllosphere microbiome in plant health and function 1, 1–24.

Szoboszlay, M., Lambers, J., Chappell, J., Kupper, J.V., Moe, L.A., McNear, D.H., 2015. Comparison of root system architecture and rhizosphere microbial communities of Balsas teosinte and domesticated corn cultivars. Soil Biology and Biochemistry 80, 34–44. https://doi.org/10.1016/j.soilbio.2014.09.001

Thapa, S., Prasanna, R., 2018. Prospecting the characteristics and significance of the phyllosphere microbiome. Ann Microbiol 68, 229–245. https://doi.org/10.1007/s13213-018-1331-5

Thapa, S., Prasanna, R., Ranjan, K., Velmourougane, K., Ramakrishnan, B., 2017. Nutrients and host attributes modulate the abundance and functional traits of phyllosphere microbiome in rice. Microbiological Research 204, 55–64. https://doi.org/10.1016/j.micres.2017.07.007

Timsina, J., 2018. Can Organic Sources of Nutrients Increase Crop Yields to Meet Global Food Demand? Agronomy 8, 214. https://doi.org/10.3390/agronomy8100214

Tiwari, S., Prasad, V., Chauhan, P.S., Lata, C., 2017. Bacillus amyloliquefaciens Confers Tolerance to Various Abiotic Stresses and Modulates Plant Response to Phytohormones through Osmoprotection and Gene Expression Regulation in Rice. Front. Plant Sci. 8, 1510. https://doi.org/10.3389/fpls.2017.01510

Tripathi, S., Srivastava, P., Devi, R.S., Bhadouria, R., 2020. Influence of synthetic fertilizers and pesticides on soil health and soil microbiology, in: Prasad, M.N.V. (Ed.), Agrochemicals Detection, Treatment and Remediation. Butterworth-Heinemann, pp. 25–54. https://doi.org/10.1016/B978-0-08-103017-2.00002-7

Vandenkoornhuyse, P., Quaiser, A., Duhamel, M., Le Van, A., Dufresne, A., 2015. The importance of the microbiome of the plant holobiont. New Phytologist 206, 1196–1206. https://doi.org/10.1111/nph.13312

Vorholt, J.A., 2012. Microbial life in the phyllosphere. Nat Rev Microbiol 10, 828–840. https://doi.org/10.1038/nrmicro2910

Wagner, B.D., Grunwald, G.K., Zerbe, G.O., Mikulich-Gilbertson, S.K., Robertson, C.E., Zemanick, E.T., Harris, J.K., 2018. On the Use of Diversity Measures in Longitudinal Sequencing Studies of Microbial Communities. Front. Microbiol. 9. https://doi.org/10.3389/fmicb.2018.01037

Wan, X., Chen, X., Huang, Z., Chen, H.Y.H., 2021. Contribution of root traits to variations in soil
microbial biomass and community composition. Plant Soil 460, 483–495.
https://doi.org/10.1007/s11104-020-04788-7
Wang, W., Jia, T., Qi, T., Li, S., Degen, A.A., Han, J., Bai, Y., Zhang, T., Qi, S., Huang, M., Li, Z., Jiao,
J., Shang, Z., 2022. Root exudates enhanced rhizobacteria complexity and microbial car-
bon metabolism of toxic plants. iScience 25, 105243.
https://doi.org/10.1016/j.isci.2022.105243
Wei, Z., Jousset, A., 2017. Plant Breeding Goes Microbial. Trends in Plant Science 22, 555–558.
https://doi.org/10.1016/j.tplants.2017.05.009
White, Bruns, T., Lee, S., Taylor, J., 1990. Amplification and direct sequencing of fungal ribosomal
RNA Genes for phylogenetics. pp. 315–322.
Yadav, B., Jogawat, A., Rahman, M., Narayan, O., 2021. Secondary metabolites in the drought
stress tolerance of crop plants: A review. Gene Reports 23, 101040.
https://doi.org/10.1016/j.genrep.2021.101040
Yilmaz, P., Parfrey, L.W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Lud-
wig, W., Glöckner, F.O., 2014. The SILVA and "All-species Living Tree Project (LTP)" taxo-
nomic frameworks. Nucleic Acids Research 42, D643–D648.
https://doi.org/10.1093/nar/gkt1209
Young, V.B., Schmidt, T.M., 2008. Overview of the Gastrointestinal Microbiota. Adv Exp Med Biol
635, 29–40. https://doi.org/10.1007/978-0-387-09550-9_3
Zaynab, M., Fatima, M., Abbas, S., Sharif, Y., Umair, M., Zafar, M.H., Bahadar, K., 2018. Role of
secondary metabolites in plant defense against pathogens. Microb Pathog 124, 198–202.
https://doi.org/10.1016/j.micpath.2018.08.034
Zhang, J., Kobert, K., Flouri, T., Stamatakis, A., 2014. PEAR: a fast and accurate Illumina Paired-
End reAd mergeR. Bioinformatics 30, 614–620. https://doi.org/10.1093/bioinformat-
ics/btt593

993