# *chrna3* modulates alcohol response

## Authors

Joshua Raine[1], Caroline Kibat [1, 2, #],  Tirtha Das Banerjee [3], Antónia Monteiro[3], Ajay S. Mathuru [1, 2, 4, 5, 6 *]

## Affiliations

1.  Department of Physiology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore, 117593

2.  Yale-NUS College, National University of Singapore, Singapore, Singapore, 138527

3.  Department of Biological Sciences, National University of Singapore, Singapore, Singapore, 117558

4.  Institute for Digital Medicine (WisDM), Yong Loo Lin School of Medicine, National University of Singapore, Singapore, 117456

5.  Institute of Molecular and Cell Biology, A*STAR, Singapore, Singapore, 138673

6.  Healthy Longevity Translational Research Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore, 117596

**\*Correspondence:** ajay.mathuru@nus.edu.sg # At the time the work was performed

## Abstract

Alcohol use disorders (AUDs) are complex phenomena governed by genetics, neurophysiology, environment, and societal structures. New methods to understand the underlying neurogenetics are valuable for designing personalised interventional strategies. Here, we used a two-choice self-administration zebrafish assay (SAZA) to isolate the function of nicotinic acetylcholine receptor (nAChR) subunit alpha3 (*chrna3*) in alcohol response. Juvenile zebrafish, prior to sex differentiation, were examined in this study. They exhibited a biphasic response when self-administering alcohol that transitioned from attraction to aversion within minutes, suggesting they can regulate exposure to alcohol. This inverted U-shaped self-administration mirrored the effect alcohol has on shoaling behaviour. Exposure to low concentration of alcohol reduced anxiety-like behaviours, while sedative effects became prominent at higher concentrations resulting in reduced locomotion and uncoordinated swimming. In contrast, these responses are blunted in *chrna3* mutants. They exhibited prolonged alcohol self-administration, and increased gregariousness. Transcriptomic analyses suggest that glutamatergic and GABAergic neurotransmission alongside cholinergic signalling is impacted in the mutant brains. Our results thus suggest that *chrna3* dysfunction has a systemic change with an increase in alcohol tolerance being one effect. These findings also highlight the use of non-rodent alternatives to understand the neurogenetics of development of AUD.

## Keywords:

Zebrafish, Preclinical model, Alcohol Use Disorders, SUD, Hormesis, *CHRNA3*

## Introduction

Substance use disorders (SUDs), encompassing nicotine, alcohol, and opioid abuse, represent a significant global health burden, accounting for approximately 162.5 million disability-adjusted life years lost worldwide (Rehm and Shield, 2019a; Moeller and Terplan, 2020). According to the WHO, harmful alcohol use alone is estimated to have resulted in 5.3% of deaths across the world in 2022 (Rehm and Shield, 2019b; Witkiewitz et al., 2019; Aslam and Kwo, 2023). Efforts to understand the interplay among social, environmental, and biological factors to effectively counter alcohol use disorders (AUDs) continue to be a focus area, as the response to alcohol in vertebrates itself is a complex multifactorial phenomenon. Alcohol interacts with a range of neuromodulatory systems, engages reward processing circuits, and affects multiple neurotransmitter systems, including glutamate and GABA neurotransmission (Chvilicek et al., 2020). Both genetic and neurobiological factors contribute significantly to AUD development (Egervari et al., 2021; Koob and Vendruscolo, 2023). Thus, a comprehensive understanding of these factors, while accounting for individual variability, is crucial for developing effective AUD prevention and interventional strategies (Witkiewitz et al., 2019; Yang et al., 2022; Aslam and Kwo, 2023).

Animal models are indispensable for dissociating genetic contributors to the development of substance dependence in humans (Smith, 2020). While such studies are typically conducted in mammalian models, high operating costs can hinder studies on genetic variability and small, single gene effects (Cox, 2015; de Magalhães, 2015). Zebrafish present a cost-effective alternative, combining high throughput genetic manipulation with whole-brain live neural activity imaging (Cully, 2019). Additionally, anatomical and functional conservation with mammalian sub-cortical circuitry is now well established (Kily et al., 2008; Gerlai, 2020).

Zebrafish exhibit behavioural and physiological responses to psychoactive substances that are consistent with mammals, providing construct validity, however, studies have primarily employed indirect

measures, like place preference or avoidance assays (Echevarria et al., 2011; Tsang et al., 2019). These

are useful, but are considered inadequate to model SUD (Smith, 2020). Despite the difficulty in designing

contingent, self-administration assays for aquatic animals, they are essential to gain predictive validity

(Smith, 2020). Few studies have explored active choice (Schneider et al., 2023) or self-administration

(Sterling et al., 2015; Bossé and Peterson, 2017; Nathan et al., 2022). Among these, studies with

juveniles (~4-5 weeks), as reported here, offer an advantage over both larvae or adults as behavioural

responses become more consistent by this age, and a relatively large number of individuals can be

examined to detect small effects.

Here, we expanded on the Self-Administration Zebrafish Assay (SAZA) (Nathan et al., 2022) to study

*chrna3*'s role in acute, dynamic alcohol responses. Human genetic studies consistently associate single

nucleotide polymorphisms (SNPs) in the *CHRNA5-A3-B4* cluster, encoding nicotinic acetylcholine

receptor (nAChR) subunits α5, α3, and β4, with nicotine addiction by GWAS (Thorgeirsson et al., 2008;

Amos et al., 2010; Bierut, 2011). Rare variants in *CHRNA3* have been linked to alcohol dependence in

humans (Haller et al., 2014; Kamens et al., 2023). While investigations into contributions of *CHRNA5* and

*CHRNB4* to SUD have progressed rapidly using rodent models (Fowler et al., 2011; Frahm et al., 2011),

such models for *CHRNA3* are fewer due to neonatal lethality of knock-outs (Xu et al., 1999; Elayouby et

al., 2021). This leaves the role of *CHRNA3* in alcohol dependence unclear and correlational. Our study

fills this gap, validating *chrna3*'s contribution to the acute alcohol response.

Using SAZA for juvenile zebrafish, we report an acute biphasic response to alcohol, transitioning rapidly

from attraction to avoidance. This correlates with the proposed role of alcohol as both stimulant and

sedative. *chrna3* mutants showed alcohol tolerance, blunting the avoidance response and resisting

changes to shoaling behaviour. In the context of the Genetically Informed Neurobiology of Addiction

(Bogdan et al., 2023), our study provides new evidence that variations in *CHRNA3* may directly influence

predisposition to alcohol addiction in humans.

# Materials & Methods

## Fish husbandry

Zebrafish (*D. rerio*, ABWT) were housed, bred, and reared as described by Nathan *et al.*, 2022(Nathan et al., 2022) at the ZebraFish Facility (Institute of Molecular and Cell Biology, A*STAR) in groups of 20–30 in 3-L tanks under standard facility conditions. All experimental protocols involving zebrafish were approved by the Institutional Animal Care and Use Committee (IACUC) of the Biological Resource Center at A*STAR. Approved experimental protocols (IACUC #201529, # 231808) were followed.

## Generation of *chrna3* mutant zebrafish

The full details of zebrafish gene editing by CRISPR/Cas9 have been described by Hwang *et al. (Hwang et al., 2013)*. CRISPR targets were determined using the web tool 'chop-chop' for the exon 5 region to disrupt key functional domains. Oligonucleotides containing the identified CRISPR target sequence ($GGN_{20}$), along with a universal reverse primer encoding the sgRNA scaffold, were used to generate a transcription template, following the method described by Bassett et al. (Bassett et al., 2013). The sequences of these oligonucleotides are listed in Table 1. CRISPR target sites are underlined, and homology arms are shown in lowercase letters. Zebrafish embryos (nacre background) were injected at the 1-cell stage with 1 nL of a mixture containing 1 µL sgRNA (≈ 4-5 µg) and 1 µL Cas9 mRNA (≈1 µg). Fish were genotyped by fin-clipping and sequencing of genomic DNA at 10 weeks post-injection to identify founder fish (F0). The identified founder fish were outcrossed to AB background WT fish to obtain F1 embryos. F1 embryos from each outcrossed family were collected and some of the embryos were verified by sequencing. Once germline transmission was confirmed by sanger sequencing, the remaining embryos were grown to adulthood. Homozygous *chrna3*$^{C246X/C246X}$ mutants (AB x nacre mixed

background) from the F3 generation onwards were used for all subsequent experiments, alongside WT

equivalents, and are henceforth referred to as *chrna3*$^{C246X}$.

Self-administration for zebrafish assay (SAZA) apparatus setup

The SAZA assay apparatus was set up as described in (Nathan et al., 2022), outlined briefly as follows.

The assay tank was constructed of opaque, 3mm thick acrylic to be 35 x 75 x 30mm (width × length ×

height). One end of the zone contained a dividing sheet of the same material, 30mm in length, to split

the zone into two halves while still permitting access (Figure 1A). The zone was placed on an LED

lightbox for illumination, providing 5000 lux at maximum intensity (LightPad LX Series, Artograph, USA).

Video footage was captured at 30 frames per second using an acA2040-90µM USB3.0 Basler camera.

Solutions were dispensed from vertically mounted 10ml syringes through silicon tubing (outer diameter:

1/16", inner diameter 1/32") into the corner of their designated zone, mediated via solenoid pinch

valves (Automate Scientific, USA, SKU: 02-pp-04i) (Figure 1A). System water was consistently supplied to

the tank through additional silicon tubing at the opposite end of the tank to the divided zones and was

removed at a matching rate (~2ml per minute) by gravitational siphoning from the end of the dividing

sheet near the centre of the tank (Figure 1A). This created a consistent flow and stopped dispensed

solutions from escaping their designated zones while permitting access to the fish being assayed

(Nathan et al., 2022). CRITTA (http://www.critta.org), a custom LABVIEW software, was used to track

fish movements as described by Krishnan *et al.* (2014) (Krishnan et al., 2014). The tracking system was

configured so that only the entry of the fish to a designated region of interest would trigger the pinch

valve to release and dispense the corresponding solution.

## Alcohol concentration measurement

Alcohol concentrations in the SAZA stimulus zone were measured using a DensitoPro (Mettler-Toledo), returning the measured solution to the zone with minimal disturbance. As attempting to measure the concentration of alcohol during a live fish assay would disturb the fish and prevent a natural response, an artificial dispensing scheme of ten continuous seconds per minute followed immediately by measurement was used as a substitute (Figure 1C). This was based on the average time fish spent in the stimulus zone under assorted alcohol dispensing solutions over 18 minutes of exposure (Table 2). After 12 minutes, the dispensing cycle was halted, but measurements continued to estimate the effect the outflow rate had on concentration in the zone if no stimulus was dispensed for a long period. The assay was repeated three times per concentration of alcohol (20, 70%).

## SAZA

SAZA was performed as described in (Nathan et al., 2022), outlined briefly as follows. First, 40 ml of system water was added to the tank, the in and outflow tubes were inserted into their respective conditions, and the flow of system water started. One zebrafish of 30-35 dpf was taken directly from its husbandry tank and added gently to each SAZA tank, then given ~5 minutes to adjust and recover post-transfer until consistent swimming behaviour had resumed and the fish had explored all zones of the SAZA tank. The recording and tracking scheme was then started for 24 minutes, consisting of three consecutive periods; three minutes of pre-exposure, 18 minutes of self-administration stimulus delivery, and three minutes post-exposure. Only during the 18-minute stimulus delivery period would entry to a designated zone trigger dispensing of the corresponding solution (Figure 1A). The valve opening time was set to a minimum of five seconds per entry. The stimuli delivered were 0/20/70% absolute alcohol diluted in the facility system water. Following each assay, the zone was emptied of all liquid, washed,

and refilled before adding the next fish. The volume of each solution dispensed was also recorded. To

reduce bias, the zone designated to deliver the stimulus solution was randomly assigned to be either the

left or right and changed periodically throughout the experiment such that ~50% of the fish were tested

in each configuration. Around 30 fish naive to SAZA were assayed per concentration of alcohol.

Following each assay, the fish were euthanized. Juvenile fish, ~5 weeks old, were used for all

behavioural experiments as  this developmental stage offers the advantages of more complex locomotor

behaviour and reduced individual variability over younger larvae, while possessing comparable

structural brain development to adult fish, yet still facilitating higher throughput and practicality of assay

design.

SAZA Data analysis

SAZA data outputs from CRITTA were processed by custom Python scripts. These provided data on time

spent in each zone and mean velocity in each zone used in these analyses. The data were separated into

three major time groups; pre-exposure, total stimulus delivery period, and post-exposure. The stimulus

delivery period was then further subdivided into either start and end (0-9m/9-18m), or three minute

segments (0-3m/3-6m/6-9m, etc…) for further analyses.

The preference index (PI) was calculated as the ratio of the difference between the times spent in the

stimulus and control zones relative to the total time spent in the two zones. The PI for a given duration

($PI_X$) was calculated by the following formula; $PI_X$ = (TinS - TinC) / (TinS + TinC), where TinS = time in

stimulus zone (s) and TinC = time in control zone (s). This gives the value for PI maximum bounds of + or

- 1 indicating more time spent relatively in the stimulus or control, respectively. For example, a PI of +1

would indicate 100% of relative time spent in the stimulus zone during one time period. Delta PI

between two time periods ($\Delta PI_{X-Y}$) was calculated as $\Delta PI_{XY}$ = $PI_Y$ - $PI_X$ , where $PI_X$ = the earlier period, and

$PI_Y$ = the later period. This gives the value for $\Delta PI$ maximum bounds of + or - 2, with the values indicating

the direction and magnitude of the change in preference from the first time point to the second. For

example, a ΔPI of +2 would indicate a change from 100% relative time spent in the control zone in the

first period, to 100% relative time spent in the stimulus zone in the second. Sparklines were also

generated to display the number of entries to each zone, and the duration of each (Figure 1A).

Shoaling cohesion assay

The shoaling cohesion assay apparatus was set up as described for SAZA, with the following alterations.

The tank had no dividing sheet, or inflow, outflow, and stimulus dispensing tubing (Figure 4A).

Additionally, the tracking of fish movements was not performed. To perform the assay, the tank was

half-filled with 20 ml of system water, then four zebrafish of 30-35 dpf were taken directly from their

husbandry tank and added gently to each shoaling cohesion tank. The fish were then given ~5 minutes

to adjust and recover post transfer until consistent swimming behaviour had resumed. Following this

acclimation period, 20 ml of 2x concentration alcohol solution was added to the tank to achieve the 1x

desired concentration for the assay. Recording began immediately and continued for 30 minutes. Six

repeats were performed per alcohol concentration. Following each assay, the fish were euthanized.

Shoaling cohesion data analysis

Video analysis was performed in imageJ. First, videos were either trimmed to the first 15 minutes, then

frames were extracted at 2fps to create an image stack (900 per assay), or one frame per 10 seconds of

footage was extracted from the full 30-minute duration (180 per assay). The image stacks for each assay

were cleaned using median background subtraction, then converted to 8-bit and auto-thresholded

(Figure 4A). From the thresholded images, particles were detected (size = 100-infinity), and the

centerpoint of each was determined. Using the centerpoints, Delaunay triangulation was performed on

each image to draw triangles covering the space between all four fish. The area of these triangles was

then combined to obtain the total area between the fish, as a measure of shoaling cohesion. These area

values were grouped into either total time for the full 30 minute duration (n = 1080 per treatment), or

one minute intervals for the first 15 minute data (n = 720 per time point per treatment). Values over

2000 mm$^2$ were identified as products of processing errors and removed (~2.5%).

As a measure of shoal kinesis, the distance travelled by the shoal centerpoint was calculated using the X

and Y coordinates for the centerpoint of the triangulation area, as calculated by imageJ, with the

equation; distance = $\sqrt{((X_2 - X_1)^2 + (Y_2 - Y_1)^2)}$. $X_2$ and $Y_2$ were the subsequent coordinates, 0.5s after $X_1$

and $Y_1$, respectively. This provided the distance the shoal centerpoint had travelled between each

selected frame. Values over 10 mm per 0.5s were discarded as products of processing errors, as the

highest velocities recorded by individual fish were ~20mm/second (Figure 3B). This amounted to < 5% of

the data. These values were grouped into one minute intervals for the first 15 minute data (n = 720 per

time point per treatment).

Quantitative PCR (qPCR)

Twelve-week old adult zebrafish were euthanized and the brains were dissected in PBS pH7.0. Brain

tissue was homogenised using a micro tube homogenizer (Thomas Scientific) in a lysis buffer made up of

Trizol (ThermoFisher #15596026). RNA was extracted using PureLink® Micro Kit (ThermoFisher

#K310250) according to the manufacturer's protocol. The concentration and purity of the extracted RNA

were determined by NanoDrop™2000 Spectrophotometer (ThermoFisher). Each brain sample yielded

around 100 ng/μL RNA with a ratio of absorbance reading 1.9-2.0 at 260/280 nm. The purified RNA was

reverse transcribed by SuperScript II First strand Synthesis System (Invitrogen #18091050) using 100

ng/μL of purified RNA to obtain approximately 1000 ng/μL of first strand cDNA. To check for genomic

DNA contamination, a negative RT (no reverse transcriptase) was set up for each sample.

The cDNA was diluted with nuclease-free water to 100 ng/μL. The qRT-PCR amplification mixtures (20

μL) contained 100 ng of cDNA, 10 μL 2x GoTaq®qPCR Master Mix (Promega #TM318), and 300 nM

forward and reverse primer. The primers against *chrn* targets used were designed using Primer3

(v.0.4.0) and are detailed in Table 3.

Reactions were performed using a 7500 Real-Time PCR system (Applied Biosystems) in 96-well plate

format. The cycling conditions were as follows: 95°C for 10m to activate the polymerase, then 40 cycles

of 95°C for 15s, 60°C for 30s. After completion, a melt curve was run at 65-95°C for 5 seconds per step.

All PCR efficiencies were above 95%. Primer specificity was validated by a single peak on the post-PCR

melt curve and single-band after electrophoresis.

The relative gene expression levels were analysed using the adjusted delta-delta CT method described

by Hellemans and Mortier et al. (2007). All data were analysed by one-way ANOVA, with genotype as

the independent variable and target gene expression, relative to reference genes, as the dependent

variable. For mRNA relative expression data, adjusted *p* values were estimated using Dunnett post-hoc

tests. Effect sizes for all differences in expression were also calculated using Cliff's Delta statistics as

described below.

Statistical analyses

Data was analysed and presented following the principles outlined by Ho *et al.* (2019), to which a p value

as a measure of significance is improved upon by the supplementary reporting of effect sizes, means,

and 95% confidence intervals (Ho et al., 2019). As the assumptions of normality and homoscedasticity

recommended for parametric testing could not be consistently met for the SAZA data, most unpaired

data were compared by non-parametric two-sided permutation T-tests, (5000 reshuffles) with a

significance level of 0.05, and unpaired Cliff's delta, while paired or repeated measures data were

compared using by paired permutation tests under the same parameters. Due to the more complex

design and higher replicate availability, the shoaling assay longitudinal data were analysed using a linear

mixed effects model, with shoal area or centerpoint travel distance as the response variable, alcohol

treatment and time as fixed effects, and assay ID as a random effect to account for repeated measures.

The residuals indicated the data were slightly skewed but appeared normally distributed and

homoscedastic enough to support a parametric test. Analysis was performed in R using the 'dabestr'

(v2024.12.24) and 'lmerTest' (v3.1-3) packages. In the case of multiple comparisons, the Holm-

Bonferroni step down family wise error correction was applied to the $p$ value. All statistical analysis

outputs can be found in supplementary tables. This data was presented in the form of Gardener-Altman

and Cummings estimation plots. Results are reported as Cliff's delta, or Paired mean difference = X, 95%

CI [lower, upper], $p$ = Y. For Cliff's delta, effect sizes greater than ± 0.4 and a $p$-value of < 0.01 was

considered meaningful and practically relevant in this study. Smaller Cliff's delta effect sizes of ± 0.2 to

0.4 with accompanying $p$-value of < 0.05 were also considered potentially meaningful, but provisional

pending experimental reproduction (Cliff, 1993; Goodman et al., 2019).

Hybridisation Chain Reaction, RNA-fluorescent in-situ hybridisation (HCR[TM] RNA-FISH)

Hybridisation Chain Reaction RNA-FISH 3.0 (HCR) (Choi et al. 2018) probes against *chrna3*, *chata*, and

*nrp1a* were prepared using an in-house Excel-based program and synthesized by IDT (please see Table 4

for details). The in-situ hybridization experiments were carried out in a controlled incubation chamber

(40°C temperature and > 90% humidity). Briefly, 14 dpf zebrafish embryos were euthanized and fixed in

4% formaldehyde diluted in 1X PBST at room temperature for 60 minutes with shaking. The larvae were

washed with 1X PBST for 3 minutes x 3 times. Embryos were permeabilized using a detergent solution

(1% SDS, 0.5% Tween 20, 150mM NaCl, 1mM EDTA, 0.05M Tris-HCL, pH 7.5) at 37°C for 30 minutes.

They were washed for 3 minutes x 3 times each with 1X PBST and then 5X SSCT.  Larvae were

transferred to 30% probe hybridization buffer (30% formamide, 9mM citric acid, 0.1% Tween 20,

50µg/mL heparin, 1x Denhardt's solution, 5X SSC) and incubated at 37°C for 30 minutes. Simultaneously,

probes were prepared in a solution containing 30ul (10uM) of probe set against each gene in 1000 µl of

30% probe hybridization buffer. The larvae were either transferred to the robotic chamber or a 1.5ml

centrifuge tube for automated or manual experiments. Larvae were incubated with the probes at 37°C

for 8-12 hrs. For both methods, on day 2, larvae were washed 5 times with 30% probe wash buffer (30%

formamide, 9mM citric acid, 0.1% Tween 20, 50µg/mL heparin, 5x SSC) at 37°C and then 5X SSCT twice.

For the secondary reaction, HCR hairpins (Molecular Instruments) were diluted and prepared in an

amplification buffer (5X SSC, 0.1% Tween 20, 2.5% dextran sulphate) and incubated overnight (for the

manual methodology) or 3 hours (for the automated) in the dark. The larvae were then washed with 5X

SSCT, 4 X 10 mins each in the dark. They were mounted with the cover slip on the dorsal side in 70%

glycerol-PBS mounting media and imaged on an Olympus FV3000 confocal microscope with the settings

consistent between fish. Images were post-processed first on Olympus viewer FLUOVIEW FV31S-SW

software with linear intensity adjustments of 150 - 4095 for all genes. Images presented for *nrp1a*, *chata*

and *chrna3* had no further processing beyond cropping to the ROI and addition of scale bars. The *elavl3*

and *chrna3* false colour images were processed in Imagej. The *elavl3* images had additional linear

brightness/contrast adjustment (15 - 110). False colour images were converted to 8bit and had the 'fire'

LUT applied, followed by linear brightness/contrast adjustment of 0 - 175, background subtraction with

25px rolling ball, and 1px gaussian blur. Fluorescence quantification (S1, S2) was performed in ImageJ on

the raw images without post-processing. The regions of interest were marked using the boundaries

observed in *chata* for the facial motor nucleus, or *nrp1a* for the dorsal habenula, and mean grey pixel

value within the region was measured. Two week old (14 dpf) larvae were used for HCR imaging as this

age represented the upper limits of transparency without additional optical clearing steps, while being

closer to behavioural testing age than the traditional imaging of 5-7 dpf larvae. Single cell analysis shows

that 97% of the cell types in a 15 dpf fish are present in 25 dpf fish (Raj et al., 2020), so we deemed it an

acceptable compromise for the benefits throughput and visualisation over imaging older animals.

RNAseq

RNA was extracted from twelve-week old adult zebrafish $chrna3^{C246X}$ and WT brains in triplicate as

described previously. The quality of each sample was verified by gel electrophoresis and bio-analyser.

Samples that passed QC were sent for sequencing by Novogene AIT. An mRNA cDNA library made up of

Paired-end sequencing reads of 150 bp in length was generated on the Illumina HiSeq instrument. Raw

reads were first processed by Novogene-AIT to remove adapter, poly-N sequences, and reads with low

quality from raw data. All downstream analyses were based on the clean data with a Phred score of 39

indicating a 99.9% base call accuracy. Further analysis was performed in Partek™ Flow™ Explore Spatial

Multiomics Data using Partek™ Flow™ software, v11.0. Reads were aligned to GRCz11 genome assembly

by STAR aligner, default parameters, and subsequently filtered to a minimum mapping quality score of >

30. Gene counts were normalised by median ratio, followed by differential expression analysis in

DeSeq2. Significance thresholds were set as follows: up/down-regulated genes = $p < 0.05$ & fold change

> ±2. Provisional up/down-regulated genes = $p < 0.05$ & fold change < ±2. Non-significant genes = $p >$

0.05. Custom gene lists and sets for analysis can be found in Table 5. Gene set enrichment analysis

(GSEA) was performed using the GRCz11-GO database, 100 permutations. Significance thresholds were

set at $p < 0.05$ and FDR < 0.2. Given the loss of spatial and structural resolution when using bulk RNA

sequencing, this investigation was primarily targeted at identifying brain-wide, compensatory

mechanisms that could potentially underlie phenotypic changes, or seed predisposition to multi-

substance dependence and comorbid disorders, we opted for the 'fully developed' brain of the sexually

mature adult. These larger adult tissues provided larger quantities of high quality RNA more consistently

than younger animals.

Results

Zebrafish preference for alcohol self-administration in SAZA is biphasic.

We refined the Self-Administration Zebrafish Assay (SAZA) first reported in (Nathan et al., 2022) to

improve the temporal resolution for quantifying the dynamic nature of the acute response. SAZA

apparatus created three chemically distinct, but physically continuous zones, or regions within the assay

chamber. The inflow and outflow were adjusted by gravity feed to maintain the integrity of the two

delivery zones throughout the assay period (Nathan et al., 2022). Subject fish behaviour was monitored

by a closed-loop, video acquisition program developed in-house that allowed for self-administration of a

stimulus or a control when subject fish entered a delivery zone from a neutral zone (Figure 1A; Movie 1).

To avoid intrinsically preferred swimming direction bias, the stimulus and control delivery zones were

pseudo-randomly assigned in each trial such that half of the fish in each group received the stimulus

(alcohol) in the left and the other half in the right delivery zone (Figure 1A). Sparkline plots in pilot

experiments showed that fish administered 20%, or 70% alcohol (at source) differentially, by changing

the frequency of entry, and duration of time spent in the stimulus area (Figure 1B; Movie 1). This

suggested that fish could quantifiably self-regulate alcohol administration, and their behaviour in the

closed-loop setup was dependent on stimulus concentration.

Next, we estimated the stimulus concentration that fish experienced during the assay. To do so, we used

a densitometer to measure the change in water density caused by the alcohol-water mixture. Alcohol

was dispensed mimicking the behaviour of fish in the pilot experiment (Figure 1B, ~10 seconds/min

discontinuously, table 2). This analysis revealed that, due to improved outflow and the lower density of

the alcohol-water mixture, the alcohol concentration in the chamber was only a fraction of that at the

source. This concentration was lower than previous estimates obtained from colorimetric analysis of

food dye-containing solution (Figure 1C). The maximum concentration plateaued at ~40-fold lower than

the source alcohol concentration, which was at least four-fold lower than previous studies (Nathan et

al., 2022). Therefore, on an average, self-administering source alcohol at 20% and 70% resulted in the

fish experiencing alcohol at ~0.5% and ~1.5% respectively in the assay (Figure 1C; for 20%, x̄ = 0.52 %, SD

= 0.13%; 70%, x̄ = 1.52%, SD = 0.44%). We refer to these two conditions as x̄0.5% condition and x̄1.5%

respectively in the rest of this study. The effective concentration reduced by half within a minute of

stopping self-administration (Figure 1C). Thus, both the pre, and post-administration periods served as

controls to compare the behaviour of animals in the naïve pre, and post exposure periods. This range of

alcohol exposure is comparable to the concentration of alcohol used in previous zebrafish studies (Kurta

and Palestis, 2010; Echevarria et al., 2011; Mathur et al., 2011; Tsang et al., 2019).

The dynamic change in alcohol concentration over time during SAZA is a key advantage for untangling

the small effects single genes may have on complex behaviours like alcohol preference. Common

variants in *CHRNA3*, for example, are risk factors for developing nicotine dependence, while several rare

variants are proposed to influence cocaine and alcohol dependence (Haller et al., 2014; Kamens et al.,

2023). *Chrna3* is proposed to mediate alcohol stimulant effects in rodents (Kamens et al., 2009).

However, this gene's role in development of alcohol dependence hasn't been tested directly. We

hypothesised that the SAZA will be effective in examining such a role, if present, in nicotinic

acetylcholine receptor alpha3 (*chrna3)* mutant generated recently (Kibat, 2020).

The *chrna3* mutant harbours an insertion that causes a frameshift and introduces a premature stop

codon within transmembrane domain 1 of the gene, resulting in a truncated protein at amino acid

position 246 (*chrna3$^{C246X}$*; Figure 1D). The mutant protein is predicted to have a large truncation of the

transmembrane and the intracellular domains (Figure 1E) and is thus expected to be damaging to

protein function. Quantitative RT-PCR of adult whole brains showed that the mutation resulted in a

significant reduction in the expression of *chrna3* transcripts in the mutants (Figure 1F, *WT* vs.

*chrna3*$^{C246X}$, Cliff's delta = -0.848, 95CI [-0.567, -0.962], *p* = < 0.001). In rodents, *Chrna3* expression is

restricted. It is expressed in a few neurons in the hippocampus, medial habenula, interpeduncular

nucleus (IPN), the ventral tegmental area (VTA), cerebellum, locus coeruleus, and spinal cord (Wada et

al., 1989; Gotti et al., 2006). We investigated the expression of *chrna3* in 14 days post fertilization (dpf)

zebrafish larvae using an in-house modified Hybridisation Chain Reaction™ RNA fluorescence in-situ

hybridisation (HCR) method (Banerjee et al., 2024). As cholinergic function in the hindbrain motor

neurons and expression in facial motor nucleus (Fmn) have also been previously reported (Rima et al.,

2020), we quantified *chrna3* expression in the mutants. While expression of *chrna3* was high in the wild

types, it was starkly and consistently reduced in the mutants (Figure 1G, S1). We also examined

expression of *chrna3* in brain regions such as the medial habenula (dorsal habenula in zebrafish),

reported to have co-expression of *CHRNA5-A3-B4* in mammals (Xu et al., 2006; Fowler et al., 2011). Only

a few cells in the left dorsal habenula expressed *chrna3* at a low level. The expression levels in the

mutant and WT were similarly low (Figure 1H, S2).

Having established a robust system for quantifying self-administration and generated *chrna3*$^{C246X}$

mutants, we examined the behaviour of juvenile fish in the SAZA at an age when they show consistency

across the population (~5 weeks old). We first examined the response of wild-type animals (WT). The

pilot (Figure 1B) suggested the preference changed dynamically over the course of the assay. To capture

this, we collapsed the preference behaviour of animals over 3-minute windows that balanced temporal

granularity in relation to the changing alcohol concentration with quantifiable observation periods. The

preference was unchanged over time when alcohol was absent (0%; Figure S3). The same analysis for

the x̄0.5% alcohol condition revealed that fish initially spent significantly more time in the stimulus zone

compared to the pre-exposure period (Figure 2A, 3-6m, Paired mean difference = 7.62, 95CI [13.9,

0.331], $p_{adj}$ = 0.028). However, after about 10 minutes, fish reversed their preference and spent

significantly less time in the stimulus zone (Figure 2A, 12-15m, Paired mean difference = -6.47, 95CI [-

2.2, -10.1], $p_{adj}$ = 0.031). This avoidance continued for the remainder of the assay, even after the self-

administration period ended. However, the early attractive portion of this biphasic response was absent

in the x̄1.5% condition (Figure 2B, 3-6m, Paired mean difference = -1.1, 95CI [4.26, -6.27], $p_{adj}$ = 1). Fish

spent less time in the stimulus zone from six minutes onwards (Figure 2B, 6-9m, Paired mean difference

= –8.57, 95CI [-3.22, -13.00], $p$ = 0.012). Mapping the behavioural changes to the alcohol concentration

expected in the stimulus zone revealed that juvenile zebrafish preferred alcohol when the concentration

was under 0.5%, while they avoided ~0.7% or higher alcohol (Figure 2C). Thus, WT zebrafish self-

regulated alcohol exposure within a narrow range.

*chrna3* mutant fish show higher tolerance to alcohol

We next examined the behaviour of *chrna3*[C246X] mutants in the SAZA. Mutants, similar to WT

counterparts showed no preference in the absence of alcohol (0%; Figure S3, A-B). In the x̄0.5%

condition, *chrna3*[C246X] fish initially behaved like their WT spending significantly more time in the stimulus

zone (Figure 2D, 3-6m, Paired mean difference = 13.9, 95CI [20.3, 8.18], $p_{adj}$ = 0.005). However, unlike

the WT, this attraction persisted throughout the assay period (Figure 2D, 9-12m, Paired mean difference

= 5.96, 95CI [10.1, 1.94], $p_{adj}$ = 0.044). No biphasic switch to avoidance was observed. Surprisingly,

mutants did not show a strong avoidance response during the assay even in the x̄1.5% condition (Figure

2E, 12-15m, Paired mean difference = -7.15, 95CI [-1.13, -12.4], $p_{adj}$ = 0.134).

The total time spent in the control delivery zone varied among conditions and genotype, but did not

show a consistent trend (Figure S3, C-H). The frequency of entry into the stimulus zone however, was

correlated with the time spent in the stimulus zone. This was similar across both genotypes in the x̄0.5%

condition and was indicative of the preference behaviour seen (Figure S4, C-D). Interestingly, the

frequency of entry into the stimulus zone in the x̄1.5% condition showed a brief increase initially across

the two genotypes (0-3 mins; Figure S4, E-F), before it plateaued off to pre-stimulus frequency rapidly,

suggesting that a reduction in total time spent in the stimulus zone observed in Figure 2B and 2E was

driven by shorter visits, rather than reduced frequency of visits.

Despite the early attraction (Figure 2A, 2D), in general alcohol appeared to be aversive to WT zebrafish

albeit less so for the mutants. This could also be inferred from the volumes delivered in alcohol

conditions compared to the volumes delivered in the 0% alcohol condition (Figure 3A, 3B). WT fish self-

administered significantly smaller volumes in the x̄0.5% condition (Figure 3A, Cliff's delta = -0.554, 95CI

[-0.266, -0.759], $p_{adj}$ = <0.001), and in the x̄1.5% condition (Figure 3A, Cliff's delta = -0.728, 95CI [-0.441,

-0.895] $p_{adj}$ = <0.001). This supports previous observations that alcohol can be aversive to zebrafish at

certain concentrations (Nathan et al., 2022). The $chrna3^{C246X}$ mutant fish however, reduced volume of

administration only in the x̄1.5% condition (Figure 3B, 0% vs x̄1.5%, Cliff's delta = -0.583, 95CI [-0.304, -

0.791], $p_{adj}$ = < 0.001), and the two genotypes differed only during the x̄0.5% condition (Figure 3C, x̄0.5%,

Cliff's delta = -0.328, 95CI [0.584, 0.009], $p_{adj}$ = 0.027).

To quantify the magnitude of the difference between genotypes we examined a Preference Index (PI;

see methods). PI was calculated for each subject as the relative preference for the stimulus compared to

control. Positive PI values (0 to +1) therefore indicated a preference for stimulus, while negative values

(0 to -1) indicated an aversion. We also quantified a delta, or the change in the PI itself in subsequent

phases: pre-exposure phase, the early phase (0-9 minutes), and the late phase (9-18 minutes). The

duration of the phases coincides with the time required (~10 minutes) for equilibration of blood alcohol

concentration in larvae immersed in alcohol (Lockwood et al., 2004). We used the "delta-delta", or the

difference of this measure between the two genotypes to quantify what aspect *chrna3* contributes to

most (Figure 3).

As anticipated, the PI range was broad, and population mean was centred around zero in the 0% alcohol

condition (Figure 3D). No significant change was observed between pre and early phases for either

genotypes. Consequently, the delta-delta between genotypes was also insignificant (Figure 3D).

However, both WT and mutants showed a positive preference shift in the early phase in the x̄0.5%

condition (Figure 3E, WT, Paired mean difference = 0.181, 95CI [0.321, 0.017], $p_{adj}$ = 0.036, Figure 3E,

*chrna3$^{C246X}$* , Paired mean difference = 0.34, 95CI [0.488, 0.232], $p_{adj}$ = < 0.001). The delta-delta between

genotypes changed only marginally (Figure 3E). The genotypic however, was significantly different in the

x̄1.5% SAZA (Figure 3F, Delta-Delta = 0.305, 95CI [0.478, 0.114], $p_{adj}$ = 0.009). In the second half of the

SAZA self-administration period, once again the 0% condition had no directional change between the

phases, or the genotypes (Figure 3G). Although both genotypes showed similar trend in the x̄0.5%

condition (Figure 3H), it was notable that the individual variance in the PI was small for mutants, while it

was considerably broader for WT, distributed across preference and aversion. Finally, in the x̄1.5%

condition, the PI shifted towards aversion, or negative values substantially for WT (Figure 3I, WT, Paired

mean difference = -0.3, 95CI [-0.144, -0.462], $p_{adj}$ = 0.001). One the other hand, the PI in the mutants

showed no directionality (Figure 3I). In addition to this increased alcohol tolerance in the x̄1.5%

condition the mutants also exhibited heightened alcohol-induced locomotor stimulation over the full

stimulus-dispensing period (x̄0.5% condition; Figure S5, A-F) that impacted other behavioural

characteristics of the mutants (Figure S6). Altogether, this suggested that manipulation of *chrna3*

function exerted larger effects on the alcohol tolerance at higher alcohol concentrations than impact the

attractive response at lower concentrations.

 *Shoaling behaviour shows biphasic effects of alcohol exposure.*

We next explored the potential reasons for this rapid and biphasic response to alcohol self-

administration. Alcohol is well-documented to act as an anxiolytic, reducing a variety of anxiety-like

behaviours in fish (Gebauer et al., 2011; Araujo-Silva et al., 2020). However, it is also known to function

as a sedative, causing reduced swimming velocity and coordination, and increased immobility

(Echevarria et al., 2011; Tsang et al., 2019). We hypothesised that the biphasic response in SAZA is

correlated to the concentrations at which these two contrasting effects of alcohol come into play. To

test this, we examined the shoaling behaviour of juvenile fish. Shoal members are proposed to swim

near others when anxious and are expected to be more exploratory when the anxiety-like state is

reduced (Miller et al., 2013), for example under an anxiolytic alcohol dose. This will be reflected by an

increase in the total area occupied by the shoal, or decreased shoal cohesion. On the other hand, shoal

members experiencing the sedative effects of alcohol are expected to swim with decreased speed and

coordination. This impacts the distance travelled by the shoal, or the shoal kinesis, in addition to

reducing the shoal cohesion. Hence, the impact of both the anxiolytic and sedative effects of alcohol can

be measured by examining shoal cohesion and kinesis in tandem (Figure 4A). Shoal cohesion was

quantified by defining each fish as a point, then triangulating the lines between them to give the total

area between fish. Shoal kinesis was calculated by the distance between the shoal centre points, as

defined by the crossing point of the triangulation between fish, between frames at 2fps (Figure 4A).

Taking into account the concentrations associated with attraction and aversion (Figure 2C), we

examined the shoaling behaviour of four fish over 30 minutes at two fixed concentrations of 0.5%, or 1%

alcohol and compared their behaviour when alcohol was absent (0%; Figure 4A, Movie 2). The total area

between WT shoal members increased at both concentrations, with the 1% alcohol having a

proportionally larger effect on the shoal area (Figure 4B, 0 vs 0.5%, Cliff's delta = 0.409, 95CI [0.453,

0.364], $p_{adj}$ = < 0.001, 0 vs 1%, Cliff's delta = 0.518, 95CI [0.559, 0.475] $p_{adj}$ = < 0.001). The same

phenomenon was also observable in the mutants (Figure 4C, 0 vs 0.5%, Cliff's delta = 0.365, 95CI [0.412,

0.320], $p_{adj}$ = < 0.001, 1%, Cliff's delta = 0.611, 95CI [0.649, 0.573] $p_{adj}$ = < 0.001).

Differences between the two genotypes emerged when their behaviours were examined at a higher

temporal resolution. Loss of cohesion in WT fish was observable within five minutes of exposure to

either concentration (Figure 4D, 0 vs 0.5% 3-6m, Cliff's delta = 0.287, 95CI [0.320, 0.252], $p_{adj}$ = 0.032;

Figure 4D 0 vs 1% 3-6m, Cliff's delta = 0.338, 95CI [0.370, 0.305], $p_{adj}$ = 0.004). The magnitude of the

difference increased over time (Figure 4D, 0 vs 0.5% 12-15m, Cliff's delta = 0.552, 95CI [0.579, 0.522],

$p_{adj}$ = < 0.001; 0 vs 1% 12-15m, Cliff's delta = 0.728, 95CI [0.749, 0.707], $p_{adj}$ = < 0.001). Shoal kinesis

changed negligibly over time with 0.5% alcohol treatment, while it was reduced almost immediately

with 1% alcohol treatment. The effect became larger over time (Figure 4E, 0 vs 1%, 6-9m, Cliff's delta = -

0.419, 95CI [-0.387, -0.449], $p_{adj}$ = 0.001). Thus, immersion in 0.5% alcohol reduced anxiety-like

behaviours, while sedative-like effects were more evident when immersed in 1% alcohol (Figure 4D, 4E).

The *chrna3*$^{C246}$ mutants seemed to experience the effects of alcohol exposure differently. Behaviour of

unexposed (0% alcohol) was similar between the two genotypes with the *chrna3*$^{C246X}$ exhibiting only

slight differences in shoal cohesion (Figure S7, A), and shoal kinesis (Figure S7B). However, unlike the WT

shoals, exposure to 0.5% alcohol reduced shoal cohesion only marginally (Figure 4F) and it did not

increase over time as it did in the WTs (Figure 4D). This difference is clearer when genotypes are

compared directly. Mutant fish retained significantly tighter shoaling cohesion compared to the wild

type (Figure 4H, 12-15m, Cliff's delta = -0.44, 95CI [-0.409, -0.47], $p_{adj}$ = < 0.001). Meanwhile, shoal

kinesis was initially heightened in *chrna3*$^{C246X}$ fish compared to the WTs (Figure 4I, 0-3m, Cliff's delta =

0.253, 95CI [0.288, 0.219], $p_{adj}$ = 0.004), which paralleled the *chrna3*$^{C246X}$ alcohol-induced motility

stimulation in the SAZA (Figure S5, A-F). The alcohol-induced kinesis difference between genotypes

gradually declined as the assay progressed (Figure 4I). This divergence in behaviours between genotypes

suggests that the anxiolysis from the alcohol exposure was much lower in *chrna3*$^{C246X}$ than that

experienced by their WT counterparts.

The behaviour differences between the mutant and wild-type shoals were minimal in 1% alcohol for the

majority of the assay duration (Figure S7, C-D). However, in the latter part of the assay mutants seemed

to resist the sedative effects of alcohol, retaining higher shoal kinesis than the wild type (Figure S7D, 15-

18m, Cliff's delta = 0.372, 95CI [0.405, 0.34], $p_{adj}$ = 0.008). The effects of alcohol on shoal cohesion and

kinesis in the first five minutes of exposure were similar between genotypes (Figure 4, S7). Overall, these

results from the shoaling assay suggest that *chrna3*$^{C246X}$ had reduced effect of alcohol exposure on

anxiolysis as well as sedative effects seen in WT fish.

Whole brain transcriptomic analysis shows major neurotransmitter signalling pathways are affected in

mutants

To gain an understanding of these behavioural differences we performed bulk RNA sequencing of WT

and mutant whole brains. Differential gene expression (DEG) analysis suggested that the two genotypes

had large differences, with PC1 accounting for ~44% of the variation in a PCA (Figure 5A). Gene Set

Enrichment Analysis (GSEA) revealed notable enrichment in transcriptional changes in mutants (Figure

5B). The significantly enriched biological functions impacted in the mutant included neuronal

development, ion channel function, and secondary messenger-associated processes (Figure 5B).

Additionally, genes associated with feeding behaviour were also enriched. In total, the expression of

~1600 genes was significantly changed (2-fold or more), and a further ~4600 genes were identified with

provisional alterations (less than 2 fold; Figure 5C). Of these, there was an approximately even split

between up- and down-regulated gene numbers (Figure 5C). Together, this perhaps indicates a wide

cascade of compensatory changes induced by the reduced expression and function of *chrna3*.

Major neurotransmitter signalling including glutamate, and GABA receptors in addition to nAChRs,

which are activated by exposure to psychoactive substances like alcohol, also showed modest changes

in expression patterns (Figure 5D-F). Both glutamate and GABA receptors appeared to show a small but

consistent upregulation amongst many receptor subunits (Figure 5D-E). The expression of most nAChR

genes however was unchanged, though a few were positively or negatively regulated with no consistent

trend (Figure 5F). Closer inspection of the nAChR subunit gene counts revealed that expression was very

low for some subunits, as was highlighted for *chrna3* in the HCR (Figure 1, 5G). For genes with low

expression counts it is a challenge to determine significant changes through RNA-seq analysis. As

*CHRNA3* expression is co-regulated with *CHRNA5* and *CHRNB4* in humans, both of which have also been

associated with substance use disorders (Wen et al., 2016), we examined whether the expression of

these genes had also been altered in the mutant. Both *chrna5* and *chrnb4* displayed significant

downregulation by qPCR, with the effect size being much larger for the latter (Figure 5H).

We also examined gene sets curated from meta-analyses of genes consistently associated with alcohol

and nicotine dependence in humans (Hu et al., 2018), and with mammalian anxiety phenotypes (Howe

et al., 2016; Koskinen and Hovatta, 2023) to address whether *chrna3* dysfunction also alters expression

of these genes (Figure 5I-K). The alcohol dependence-associated genes exhibited few changes (Figure

5I), however, ~50% of the nicotine-dependence genes were provisionally upregulated in the mutant

(Figure 6J). Finally, genes associated with anxiety phenotypes also showed a number of expression

changes, albeit some in positive and some in negative direction of expression (Figure 5K). In all cases,

this analysis revealed that though the expression of a large number of these genes changed they were

modest.

Discussion

In this study, a refinement of self-administration zebrafish assay (SAZA) revealed an acute biphasic

response to alcohol, in juvenile zebrafish, with preference transitioning to avoidance when exposure

increased to ~0.7% alcohol (Figure 2). This biphasic response to alcohol aligns zebrafish with other

animals, suggesting that hormesis to alcohol is a fundamental property across vertebrates (Figures 1, 2;

(Calabrese and Baldwin, 2003; Calabrese, 2008; Sterling et al., 2015)). The closed-loop design of SAZA,

similar to a two-bottle choice assay used for rodents, facilitated the analysis of uninhibited choice and

isolated *chrna3*'s role in modulating this biphasic response.

Isolating contributions of individual genetic factors to a phenomena such as substance dependence is

challenging. It is further compounded for multi-subunit receptor genes that cluster at a genomic locus,

such as the *CHRNA5-A3-B4* gene cluster. This genomic locus has been consistently associated with

several substance use disorder GWASes (Chen et al., 2009; Maes et al., 2011; Haller et al., 2014; Wen et

al., 2016). Although, male transgenic mice overexpressing *Chrna5-a3-b4* consume less alcohol than wild-

type mice (Gallego et al., 2012), a causal relationship can't be ascribed to *Chrna3* alone from this study

(Kamens et al., 2023). *Chrna3* knockout mice die prematurely as neonates (Xu et al., 1999) and the

conditional knockout (Elayouby et al., 2021) only examined nicotine response leaving the role of α3 in

alcohol stimulation unclear (Kamens et al., 2009). Zebrafish *chrna3* mutants, however, survive to

adulthood, offering a tractable system to investigate gene-specific effects here. *chrna3* mutants

displayed a blunted aversive switch in the biphasic response, while maintaining early attraction (Figure

2, 3). WT fish exhibited aversive behaviour to 0.7% or higher alcohol stimulus, which only became

observable in mutants at higher concentrations. This aligns with rodent studies, where a broad spectrum

nicotinic receptor antagonist 18-MC in the medial habenula (MHb) - interpeduncular nucleus (IPN)

circuit reduced transmission and increased nicotine or alcohol self-administration, respectively (Glick et

al., 2011; Miller et al., 2019). As this circuit co-expresses all the three nAChR subunits α3, α5, and β4,

they are thus proposed to mediate aversion to psychoactive substances, including alcohol (Marks et al.,

1992; Xu et al., 2006; Frahm et al., 2011; Fowler and Kenny, 2012; Mathuru, 2018; Elayouby et al., 2021).

However, as shown in the gene expression profile here (Figure 1), and in a recent report (Hua et al.,

2025), *chrna3* is expressed widely in the central and peripheral nervous system. Consequently, altered

alcohol tolerance reported here may stem from a combined effect across multiple circuits and systems.

Further to this limitation, although most physiological features and brain regions described in adult

zebrafish are present in juveniles by 25 dpf, late-stage developmental changes may influence adult

phenotypes (Schmidt et al., 2013; Singleman and Holtzman, 2014; Raj et al., 2020), warranting further

studies.

A shifted tolerance window exhibited by *chrna3* mutants (Figure 3C) suggests a threshold based

engagement of different brain circuits, similar to the proposed mammalian twin threshold model

(Grasing, 2016). This model posits that cholinergic signalling, particularly in the VTA and nucleus

accumbens (NAc), modulates drug reinforcement behaviours. In this model, lower threshold increases

positive reward association probability, stimulating behavioural repetition, while a second, higher

threshold does the opposite as different circuits come into play (Grasing, 2016). This may also be

applicable to other cholinergic brain regions, such as the MHb. Additionally, higher agonist requirements

to activate α5, α3, and β4 containing nAChRs (Coe et al., 2005), could elevate the cholinergic threshold

to engage this aversive circuit, providing a potential explanation for why *chrna3* mutation

predominantly affected the avoidance response.

Beyond preference, *chrna3* mutation also blunted the psychoactive effects of alcohol during immersion.

Immersion is a rapid, non-invasive route for administration of water-soluble pharmaceutics, uniquely

available for aquatic animals. Alcohol immersion time correlates with blood alcohol concentration (BAC)

until equilibrium is reached, and is a key measure affecting physiological responses (Lockwood et al.,

2004). As in humans, BAC > 0.05%, or the equivalent of ~2-3 Australian "standard drinks", results in

significant behavioural impairment in zebrafish, which can be reached within 1-2 minutes of immersion

in 1.5% alcohol. This manifests in humans as slower reaction times (Grant et al., 2000), and zebrafish as

loss of swimming coordination or impaired locomotion (Tran and Gerlai, 2013; Tsang et al., 2019).

However, lower concentrations are stimulatory, activating dopaminergic reward pathways and

increasing locomotion, while reducing anxiety-like behaviours and aggression (Echevarria et al., 2010;

Araujo-Silva et al., 2020).

Shoaling behaviour, that becomes prominent by 14 dpf (Miller and Gerlai, 2007), can effectively model

these contrasting effects (Oscar-Berman and Marinković, 2007; Kurta and Palestis, 2010; Miller et al.,

2013; Rueger and King, 2013) at SAZA-relevant concentrations. In wild types, both 0.5% and 1% alcohol

immersion gradually reduced shoal cohesion, but exhibited differing effects on shoal kinesis (Figure 4).

One interpretation of the coupled reduction in cohesion and kinesis in 1% alcohol is as a sedative effect,

evident also in the video recordings (Movie 2). Meanwhile, reduction of cohesion alone could indicate

an anxiolytic effect, that dominated in a narrow alcohol concentration range (0.5%). These

concentrations approximately correlated with the attractive and aversive doses during SAZA, suggesting

a potential link. However, the $chrna3^{C246X}$ mutants displayed no reduction in shoal cohesion at 0.5%

(Figure 4H). Whether this reduction in anxiolysis phenotype in the mutants was directly due to reduced

sensitivity to alcohol or independent of it needs further investigation. Irrespective, it is evident that the

$chrna3$ mutation blunted drug-induced behavioural effects, including the reduced aversion in SAZA.

Alcohol-induced kinesis also differed between genotypes, with $chrna3^{C246X}$ mutants exhibiting stronger

alcohol-induced locomotion in SAZA and shoaling assays (Figure 4, S5). This may be correlated to the

reduced expression of $chrna3$ in the caudal and rostral regions of the mutant facial motor nucleus

(Fmn), also termed the octavolateral efferent nucleus (OEN) (Figure 1, S1), a hindbrain region connected

to the lateral line.  Cholinergic signalling here is hypothesised to act as a 'brake' on locomotor auto-

sensory stimulation generated during swimming (Pichler and Lagnado, 2020; Manuel et al., 2021;

Odstrcil et al., 2022). While typically this function is assigned to α9 receptors, if α3 plays a role, as

suggested recently (Rima et al., 2020), it may account for the heightened alcohol-induced locomotor

response in chrna3$^{C246X}$ mutants. However, chrna3's influence on locomotor control requires further

investigation, as rodent studies offer inconsistent evidence (Kamens et al., 2009).

Whole-body mutation in our chrna3 mutant precluded implicating specific brain regions like the dorsal

habenula (MHb in mammals), yet, it does inform on potential consequences of CHRNA3 variants that

reduce α3 function in humans. Whole brain RNA sequencing identified broad transcriptional profile

change in the mutants. Small, but, consistent upregulation of nicotine dependence associated genes was

indicative of a bi-directional relationship (Figure 5J; (Hu et al., 2018)), aligning with GWAS studies linking

CHRNA3 polymorphisms to increased nicotine dependence (Chen et al., 2009; Maes et al., 2011).

Downregulation of chrna5 and chrnb4 (Figure 5H), also associated with substance dependence (Fowler

et al., 2011; Frahm et al., 2011) and co-expressed with chrna3 in other vertebrates, potentially

contributed to the mutant's alcohol tolerance. Generation and characterization of an independent

chrna3, chrna5, and chrnb4 mutant lines along with their combinations (currently in progress), will be

invaluable to dissociate these factors further.

Beyond nAChRs, alcohol impacts both glutamate and GABA receptors, modulating excitatory and

inhibitory tones, respectively (Davies, 2003; Möykkynen and Korpi, 2012; Jones, 2019). Consequently,

changes in expression of these receptor genes (Figure 5D, 5E) in mutants was also interesting to

observe, as many homologs also feature in nicotine dependence and anxiety phenotypes in humans

(Figure 5J, 5K). At a functional level, neuronal receptor related GO terms like synaptic signalling, ion

channel function, and cation homeostasis were significantly enriched (Figure 5B). However, neural

development terms including neuron projection, dendrite development, and axon guidance were also

enriched despite the use of adult brains (Figure 5B), perhaps indicative of compensatory plasticity.

Finally, enrichment of feeding behaviour associated genes was notable (Figure 5B), as cholinergic

signalling and CHRNA5-CHRNA3-CHRNB4 cluster SNPs have been tentatively linked by GWAS studies to

smoking/obesity interactions (Varga et al., 2013; Herman et al., 2016; Ortiz-Guzman et al., 2024). This

transcriptomic change warrants further examination to determine the simultaneous influence of *chrna3*

on substance dependence and other comorbid disorders.

Overall, these findings suggest the type of brain-wide compensatory changes that may occur

corresponding to loss of *chrna3* function, eventually resulting in heightened alcohol tolerance and

potential predisposition to multi-substance use and comorbid disorders.

Understanding alcohol's biphasic properties is imperative to address Alcohol Use Disorder (AUD). Our

study fills a gap, providing novel evidence on the role of *chrna3* in modulating alcohol preference, and

tethering it to a specific facet of the dose-response curve. Given the paucity of animal model data,

despite GWAS associations (Haller et al., 2014), the evidence of isolated gene function from a whole

body mutation in zebrafish presented here represents a valuable step forward in the context of

phenotypes leading to alcohol dependence development in humans harbouring *CHRNA3* variants.

Acknowledgements:

We would like to thank the zebrafish fish facility (ZFF) staff at the IMCB, A*STAR for their assistance with

fish husbandry. We also thank Li Xinrui for generating customized Python analysis scripts. Additionally,

during the preparation of this work the author(s) used Grammarly, and Gemini in order to assist with

grammar and consistency of phrasing. After using this tool/service, the author(s) reviewed and edited

the content as needed and take(s) full responsibility for the content of the publication.

Supplementary Materials:

Supplementary Material Figures S1 to S7, Tables S1 to S3.

Number of supplementary figures and tables = 10

Author Contributions:

Conceptualization, A.S.M, and J.R; methodology, A.S.M., T.D.B., C.K., and J.R.; formal analysis, J.R., T.D.B

and A.S.M.; investigation, J.R., C.K., and A.S.M; resources, A.S.M; writing—original draft preparation, J.R,

and A.S.M.; writing—review and editing, All.; visualisation,   J.R.; supervision, A.S.M.; project

administration, A.S.M., A.M.; funding acquisition, A.S.M.

Funding:

This research was funded by the Ministry of Education (MOE), Singapore (through grant number

T2EP30220-0020), and Yale-NUS College (through grant numbers IG16-LR003, IG18-SG103, IG19-BG106,

and SUG) to ASM.

Data Availability Statement:

The datasets used to generate the figures are included within the article and its additional file(s).

Additional data are available from the corresponding author upon request.

Conflicts of Interest:

The authors declare no conflict of interest. The sponsors had no role in the design, execution,

interpretation, or writing of the study.

# References

References

Amos CI, Spitz MR, Cinciripini P (2010) Chipping away at the genetics of smoking behavior. Nat Genet

42:366–368.

Araujo-Silva H, Leite-Ferreira ME, Luchiari AC (2020) Behavioral Screening of Alcohol Effects and

Individual Differences in Zebrafish (Danio rerio). Alcohol Alcohol 55:591–597.

Aslam A, Kwo PY (2023) Epidemiology and Disease Burden of Alcohol Associated Liver Disease. J Clin Exp

Hepatol 13:88–102.

Banerjee TD, Raine J, Mathuru AS, Chen KH, Monteiro A (2024) Spatial mRNA profiling using Rapid

Amplified Multiplexed-FISH (RAM-FISH). bioRxiv:2024.12.06.627193 Available at:

https://www.biorxiv.org/content/10.1101/2024.12.06.627193v1.abstract [Accessed May 5, 2025].

Bassett AR, Tibbit C, Ponting CP, Liu J-L (2013) Highly efficient targeted mutagenesis of Drosophila with

the CRISPR/Cas9 system. Cell Rep 4:220–228.

Bierut LJ (2011) Genetic vulnerability and susceptibility to substance dependence. Neuron 69:618–627.

Bogdan R, Hatoum AS, Johnson EC, Agrawal A (2023) The Genetically Informed Neurobiology of

Addiction (GINA) model. Nat Rev Neurosci 24:40–57.

Bossé GD, Peterson RT (2017) Development of an opioid self-administration assay to study drug seeking

in zebrafish. Behav Brain Res 335:158–166.

Calabrese EJ (2008) U-shaped dose response in behavioral pharmacology: historical foundations. Crit

Rev Toxicol 38:591–598.

Calabrese EJ, Baldwin LA (2003) Ethanol and hormesis. Crit Rev Toxicol 33:407–424.

Chen X, Chen J, Williamson VS, An S-S, Hettema JM, Aggen SH, Neale MC, Kendler KS (2009) Variants in

nicotinic acetylcholine receptors alpha5 and alpha3 increase risks to nicotine dependence. Am J

Med Genet B Neuropsychiatr Genet 150B:926–933.

Chvilicek MM, Titos I, Rothenfluh A (2020) The Neurotransmitters Involved in Alcohol-Induced

Behaviors. Front Behav Neurosci 14:607700.

Cliff N (1993) Dominance statistics: Ordinal analyses to answer ordinal questions. Psychol Bull 114:494–

509.

Coe JW et al. (2005) Varenicline: an alpha4beta2 nicotinic receptor partial agonist for smoking cessation.

J Med Chem 48:3474–3477.

Cox TC (2015) Utility and limitations of animal models for the functional validation of human sequence

variants. Mol Genet Genomic Med 3:375–382.

Cully M (2019) Zebrafish earn their drug discovery stripes. Nat Rev Drug Discov 18:811–813.

Davies M (2003) The role of GABAA receptors in mediating the effects of alcohol in the central nervous

system. J Psychiatry Neurosci 28:263–274.

de Magalhães JP (2015) The big, the bad and the ugly: Extreme animals as inspiration for biomedical

research. EMBO Rep 16:771–776.

Echevarria DJ, Hammack CM, Jouandot DJ, Toms CN (2010) Does Acute Alcohol Exposure Modulate

Aggressive Behaviors in the Zebrafish (Danio rerio), or is the Bark Worse than the Bite? Int J Comp

Psychol 23 Available at: https://escholarship.org/content/qt2px3t3jc/qt2px3t3jc.pdf?t=mwj0ph

[Accessed October 20, 2023].

Echevarria DJ, Toms CN, Jouandot DJ (2011) Alcohol-induced behavior change in zebrafish models. Rev

Neurosci 22:85–93.

Egervari G, Siciliano CA, Whiteley EL, Ron D (2021) Alcohol and the brain: from genes to circuits. Trends

Neurosci 44:1004–1015.

Elayouby KS, Ishikawa M, Dukes AJ, Smith ACW, Lu Q, Fowler CD, Kenny PJ (2021) α3* Nicotinic

Acetylcholine Receptors in the Habenula-Interpeduncular Nucleus Circuit Regulate Nicotine Intake.

J Neurosci 41:1779–1787.

Fowler CD, Kenny PJ (2012) Habenular signaling in nicotine reinforcement. Neuropsychopharmacology

37:306–307.

Fowler CD, Lu Q, Johnson PM, Marks MJ, Kenny PJ (2011) Habenular α5 nicotinic receptor subunit

signalling controls nicotine intake. Nature 471:597–601.

Frahm S, Slimak MA, Ferrarese L, Santos-Torres J, Antolin-Fontes B, Auer S, Filkin S, Pons S, Fontaine J-F,

Tsetlin V, Maskos U, Ibañez-Tallon I (2011) Aversion to nicotine is regulated by the balanced activity

of β4 and α5 nicotinic receptor subunits in the medial habenula. Neuron 70:522–535.

Gallego X, Ruiz-Medina J, Valverde O, Molas S, Robles N, Sabrià J, Crabbe JC, Dierssen M (2012)

Transgenic over expression of nicotinic receptor alpha 5, alpha 3, and beta 4 subunit genes reduces

ethanol intake in mice. Alcohol 46:205–215.

Gebauer DL, Pagnussat N, Piato AL, Schaefer IC, Bonan CD, Lara DR (2011) Effects of anxiolytics in

zebrafish: similarities and differences between benzodiazepines, buspirone and ethanol. Pharmacol

Biochem Behav 99:480–486.

Gerlai R (2020) Evolutionary conservation, translational relevance and cognitive function: The future of

zebrafish in behavioral neuroscience. Neurosci Biobehav Rev 116:426–435.

Glick SD, Sell EM, McCallum SE, Maisonneuve IM (2011) Brain regions mediating α3β4 nicotinic

antagonist effects of 18-MC on nicotine self-administration. Eur J Pharmacol 669:71–75.

Goodman WM, Spruill SE, Komaroff E (2019) A proposed hybrid effect size plus $p$-value criterion:

Empirical evidence supporting its use. Am Stat 73:168–185.

Gotti C, Zoli M, Clementi F (2006) Brain nicotinic acetylcholine receptors: native subtypes and their

relevance. Trends Pharmacol Sci 27:482–491.

Grant SA, Millar K, Kenny GN (2000) Blood alcohol concentration and psychomotor effects. Br J Anaesth

85:401–406.

Grasing K (2016) A threshold model for opposing actions of acetylcholine on reward behavior: Molecular

mechanisms and implications for treatment of substance abuse disorders. Behav Brain Res

312:148–162.

Haller G, Kapoor M, Budde J, Xuei X, Edenberg H, Nurnberger J, Kramer J, Brooks A, Tischfield J, Almasy

L, Agrawal A, Bucholz K, Rice J, Saccone N, Bierut L, Goate A (2014) Rare missense variants in

CHRNB3 and CHRNA3 are associated with risk of alcohol and cocaine dependence. Hum Mol Genet

23:810–819.

Herman AM, Ortiz-Guzman J, Kochukov M, Herman I, Quast KB, Patel JM, Tepe B, Carlson JC, Ung K,

Selever J, Tong Q, Arenkiel BR (2016) A cholinergic basal forebrain feeding circuit modulates

appetite suppression. Nature 538:253–256.

Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A (2019) Moving beyond P values: data analysis with

estimation graphics. Nat Methods 16:565–566.

Howe AS et al. (2016) Candidate genes in panic disorder: meta-analyses of 23 common variants in major

anxiogenic pathways. Mol Psychiatry 21:665–679.

Hua Y, Habicher J, Carl M, Manuel R, Boije H (2025) Novel Transgenic Zebrafish Lines to Study the

CHRNA3-B4-A5 Gene Cluster. Developmental Neurobiology 85:e22956.

Hu R, Dai Y, Jia P, Zhao Z (2018) ANCO-GeneDB: annotations and comprehensive analysis of candidate

genes for alcohol, nicotine, cocaine and opioid dependence. Database 2018 Available at:

http://dx.doi.org/10.1093/database/bay121.

Hwang WY, Fu Y, Reyon D, Maeder ML, Tsai SQ, Sander JD, Peterson RT, Yeh J-RJ, Joung JK (2013)

Efficient genome editing in zebrafish using a CRISPR-Cas system. Nat Biotechnol 31:227–229.

Jones AW (2019) Alcohol, its absorption, distribution, metabolism, and excretion in the body and

pharmacokinetic calculations. WIREs Forensic Science 1:e1340.

Kamens HM, Flarend G, Horton WJ (2023) The role of nicotinic receptors in alcohol consumption.

Pharmacol Res 190:106705.

Kamens HM, McKinnon CS, Li N, Helms ML, Belknap JK, Phillips TJ (2009) The alpha 3 subunit gene of the

nicotinic acetylcholine receptor is a candidate gene for ethanol stimulation. Genes Brain Behav

8:600–609.

Kibat C (2020) Functional Characterization of Nicotinic Acetylcholine Receptor Subunit ALPHA 3 in

Regulation of Behaviour.

Kily LJM, Cowe YCM, Hussain O, Patel S, McElwaine S, Cotter FE, Brennan CH (2008) Gene expression

changes in a zebrafish model of drug dependency suggest conservation of neuro-adaptation

pathways. J Exp Biol 211:1623–1634.

Koob GF, Vendruscolo L (2023) Theoretical Frameworks and Mechanistic Aspects of Alcohol Addiction:

Alcohol Addiction as a Reward Deficit/Stress Surfeit Disorder. Curr Top Behav Neurosci Available at:

http://dx.doi.org/10.1007/7854_2023_424.

Koskinen M-K, Hovatta I (2023) Genetic insights into the neurobiology of anxiety. Trends Neurosci

46:318–331.

Krishnan S, Mathuru AS, Kibat C, Rahman M, Lupton CE, Stewart J, Claridge-Chang A, Yen S-C, Jesuthasan

S (2014) The Right Dorsal Habenula Limits Attraction to an Odor in Zebrafish. Curr Biol 24:1167–

1175.

Kunst M, Laurell E, Mokayes N, Kramer A, Kubo F, Fernandes AM, Förster D, Dal Maschio M, Baier H

(2019) A Cellular-Resolution Atlas of the Larval Zebrafish Brain. Neuron 103:21–38.e5.

Kurta A, Palestis BG (2010) Effects of Ethanol on the Shoaling Behavior of Zebrafish (Danio Rerio). Dose

Response Available at: https://journals.sagepub.com/doi/10.2203/dose-response.10-008.Palestis

[Accessed October 17, 2023].

Lockwood B, Bjerke S, Kobayashi K, Guo S (2004) Acute effects of alcohol on larval zebrafish: a genetic

system for large-scale screening. Pharmacol Biochem Behav 77:647–654.

Maes HH, Neale MC, Chen X, Chen J, Prescott CA, Kendler KS (2011) A twin association study of nicotine

dependence with markers in the CHRNA3 and CHRNA5 genes. Behav Genet 41:680–690.

Manuel R, Iglesias Gonzalez AB, Habicher J, Koning HK, Boije H (2021) Characterization of Individual

Projections Reveal That Neuromasts of the Zebrafish Lateral Line are Innervated by Multiple

Inhibitory Efferent Cells. Front Neuroanat 15:666109.

Marks MJ, Pauly JR, Gross SD, Deneris ES, Hermans-Borgmeyer I, Heinemann SF, Collins AC (1992)

Nicotine binding and nicotinic receptor subunit RNA after chronic nicotine treatment. J Neurosci

12:2765–2784.

Mathur P, Berberoglu MA, Guo S (2011) Preference for ethanol in zebrafish following a single exposure.

Behav Brain Res 217:128–133.

Mathuru AS (2018) A little rein on addiction. Semin Cell Dev Biol 78:120–129.

Miller CN, Ruggery C, Kamens HM (2019) The α3β4 nicotinic acetylcholine receptor antagonist 18-

Methoxycoronaridine decreases binge-like ethanol consumption in adult C57BL/6J mice. Alcohol

79:1–6.

Miller N, Gerlai R (2007) Quantification of shoaling behaviour in zebrafish (Danio rerio). Behav Brain Res

184:157–166.

Miller N, Greene K, Dydinski A, Gerlai R (2013) Effects of nicotine and alcohol on zebrafish (Danio rerio)

shoaling. Behav Brain Res 240:192–196.

Moeller G, Terplan M (2020) Substance Use Disorders. Oxford University Press, USA.

Möykkynen T, Korpi ER (2012) Acute effects of ethanol on glutamate receptors. Basic Clin Pharmacol

Toxicol 111:4–13.

Nathan FM, Kibat C, Goel T, Stewart J, Claridge-Chang A, Mathuru AS (2022) Contingent stimulus

delivery assay for zebrafish reveals a role for CCSER1 in alcohol preference. Addict Biol 27:e13126.

Odstrcil I, Petkova MD, Haesemeyer M, Boulanger-Weill J, Nikitchenko M, Gagnon JA, Oteiza P, Schalek

R, Peleg A, Portugues R, Lichtman JW, Engert F (2022) Functional and ultrastructural analysis of

reafferent mechanosensation in larval zebrafish. Curr Biol 32:176–189.e5.

Ortiz-Guzman J, Swanson JL, Tantry EK, Kochukov M, Ung K, Addison AP, Srivastava S, Belfort BD, Ji E,

Dooling SW, Chen SA, Tong Q, Arenkiel BR (2024) Cholinergic Basal Forebrain Connectivity to the

Basolateral Amygdala Modulates Food Intake. eNeuro 11 Available at:

http://dx.doi.org/10.1523/ENEURO.0369-23.2024.

Oscar-Berman M, Marinković K (2007) Alcohol: Effects on Neurobehavioral Functions and the Brain.

Neuropsychol Rev 17:239–257.

Pichler P, Lagnado L (2020) Motor Behavior Selectively Inhibits Hair Cells Activated by Forward Motion in

the Lateral Line of Zebrafish. Curr Biol 30:150–157.e3.

Raj B, Farrell JA, Liu J, El Kholtei J, Carte AN, Navajas Acedo J, Du LY, McKenna A, Relić Đ, Leslie JM, Schier

AF (2020) Emergence of Neuronal Diversity during Vertebrate Brain Development. Neuron

108:1058–1074.e6.

Rehm J, Shield KD (2019a) Global Burden of Disease and the Impact of Mental and Addictive Disorders.

Curr Psychiatry Rep 21:10.

Rehm J, Shield KD (2019b) Global Burden of Alcohol Use Disorders and Alcohol Liver Disease.

Biomedicines 7:99.

Rima M, Lattouf Y, Abi Younes M, Bullier E, Legendre P, Mangin J-M, Hong E (2020) Dynamic regulation

of the cholinergic system in the spinal central nervous system. Sci Rep 10:15338.

Rueger SY, King AC (2013) Validation of the brief Biphasic Alcohol Effects Scale (B-BAES). Alcohol Clin Exp

Res 37:470–476.

Schmidt R, Strähle U, Scholpp S (2013) Neurogenesis in zebrafish - from embryo to adult. Neural Dev 8:3.

Schneider H, Pearson A, Harris D, Krause S, Tucker A, Gardner K, Chinyanya K (2023) Identification of

nicotine-seeking and avoiding larval zebrafish using a new three-choice behavioral assay. Front Mol

Neurosci 16:1112927.

Singleman C, Holtzman NG (2014) Growth and maturation in the zebrafish, Danio rerio: a staging tool for

teaching and research. Zebrafish 11:396–406.

Smith MA (2020) Nonhuman animal models of substance use disorders: Translational value and utility to

basic science. Drug Alcohol Depend 206:107733.

Sterling ME, Karatayev O, Chang G-Q, Algava DB, Leibowitz SF (2015) Model of voluntary ethanol intake

in zebrafish: effect on behavior and hypothalamic orexigenic peptides. Behav Brain Res 278:29–39.

Thorgeirsson TE et al. (2008) A variant associated with nicotine dependence, lung cancer and peripheral

arterial disease. Nature 452:638–642.

Tran S, Gerlai R (2013) Time-course of behavioural changes induced by ethanol in zebrafish (Danio rerio).

Behav Brain Res 252:204–213.

Tsang B, Ansari R, Gerlai R (2019) Dose dependent behavioral effects of acute alcohol administration in

zebrafish fry. Pharmacol Biochem Behav 179:124–133.

Varga TV, Hallmans G, Hu FB, Renström F, Franks PW (2013) Smoking status, snus use, and variation at

the CHRNA5-CHRNA3-CHRNB4 locus in relation to obesity: the GLACIER study. Am J Epidemiol

178:31–37.

Wada E, Wada K, Boulter J, Deneris E, Heinemann S, Patrick J, Swanson LW (1989) Distribution of alpha

2, alpha 3, alpha 4, and beta 2 neuronal nicotinic receptor subunit mRNAs in the central nervous

system: a hybridization histochemical study in the rat. J Comp Neurol 284:314–335.

Wen L, Jiang K, Yuan W, Cui W, Li MD (2016) Contribution of Variants in CHRNA5/A3/B4 Gene Cluster on

Chromosome 15 to Tobacco Smoking: From Genetic Association to Mechanism. Mol Neurobiol

53:472–484.

Witkiewitz K, Litten RZ, Leggio L (2019) Advances in the science and treatment of alcohol use disorder.

Sci Adv 5:eaax4043.

Xu W, Gelber S, Orr-Urtreger A, Armstrong D, Lewis RA, Ou CN, Patrick J, Role L, De Biasi M, Beaudet AL

(1999) Megacystis, mydriasis, and ion channel defect in mice lacking the alpha3 neuronal nicotinic

acetylcholine receptor. Proc Natl Acad Sci U S A 96:5746–5751.

Xu X, Scott MM, Deneris ES (2006) Shared long-range regulatory elements coordinate expression of a

gene cluster encoding nicotinic receptor heteromeric subtypes. Mol Cell Biol 26:5636–5649.

Yang W, Singla R, Maheshwari O, Fontaine CJ, Gil-Mohapel J (2022) Alcohol Use Disorder: Neurobiology

and Therapeutics. Biomedicines 10:1192.

Zhou X, Zheng W, Li Y, Pearce R, Zhang C, Bell EW, Zhang G, Zhang Y (2022) I-TASSER-MTD: a deep-

learning-based platform for multi-domain protein structure and function prediction. Nat Protoc

17:2326–2353.

Figure & Table Legends

**Table 1.** Primers for single-guide RNA template. CRISPR targets are underlined. Homology arms in lower

case.

**Table 2.** Mean time spent by zebrafish in the stimulus chamber during SAZA over an 18 minute stimulus

dispensing period (n = 30-32)

**Table 3.** Primers used for qPCR of *chrna3*, *chrna5*, *chrnb4*, and housekeeping genes.

**Table 4.** Probe sequences for FISH-HCR. B1 Probes were visualised with Alexa fluorophore 546, B2

probes were visualised with Alexa fluorophore 647, and B3 probes were visualised with Alexa

fluorophore 488. P1 and P2 probes of the same gene are paired according to number 1-8.

**Table 5.** Genes included in the custom sets 'Nicotine dependence ' and 'Alcohol dependence'. These

genes have been associated with nicotine and alcohol dependence, described by Hu *et al*. (Hu et al.

2018)

**Figure 1**. **Self-Administration Zebrafish Assay (SAZA) chamber functional design and characterisation**

**of the *chrna3*$^{C246X}$ mutant zebrafish.**

**A.** Schematic indicating locations of inflow of system water (In, top) and outflow (Out, centre). Control

and stimulus delivery inlets (stimulus shown in purple) are activated when a fish enters the

corresponding zone. Continuous inflow and outflow constrain the stimulus/control delivered to their

respective zones (Nathan et al., 2022) . 'Measurement area' denotes where alcohol concentration was

measured in panel C. **B.** Schematic shows the experimental design with three minute pre- and post-

exposure periods before and after an 18 minute self-administration period. Representative sparklines

plots show the entries and duration (thickness) of entry of the subject fish either into the stimulus or the

control zones when 20% or 70% alcohol is used as stimuli. **C.** Alcohol concentrations measured when

20% or 70% alcohol were dispensed for the duration mimicking subject fish behaviour (± standard

error). The average concentration (solid line) fish experience is 0.5% and 1.5% alcohol respectively. See

Table 2 for additional details. **D-H.** Characterisation of *chrna3*[C246X] mutant fish**. D**. *chrna3*[C246X] mutants

have a premature stop codon in the transmembrane region 1, amino acid 246 (M1 - red star), due to the

indel of 5bp at position 786bp. **E.** Predicted protein structure of chrna3 in WT and *chrna3*[C246X] mutants,

generated by iTasser (Zhou et al., 2022), colour coded by domain. **F.** Fold change of *chrna3* in 12 week

old WT (n = 21) vs. *chrna3*[C246X] (n = 20) by qRT-PCR. **G-H.** *chrna3* expression in 14 day old zebrafish larvae

visualised by Hybridisation Chain Reaction[TM] RNA-FISH (HCR[TM]) in the **G.** Predicted facial motor nucleus

(Fmn) marked by *chata* expression and, **H.** dorsal habenula (dHb) marked by *nrp1a* expression, in WT

and *chrna3*[C246X] respectively. Images are representative, see figures S1 and S2 for replicates

(WT/*chrna3*[C246X] Fmn n = 3/4, dHb n = 4/3). Schematic images with regional markers (v2.0, MECE, 2024)

were generated in mapZeBrain Atlas (mapzebrain.org, May 2025) (Kunst et al., 2019). Hb = habenula,

vHb = ventral habenula, Omn = oculomotor nucleus, Trm = trochlear motor nucleus, Ni = nucleus isthmi,

Vmn = vagus motor nucleus, srFmn = supra-rostral Fmn, rFmn = rostral Fmn, cFmn = caudal Fmn. Genes

presented left to right: all genes merged, **G.** *chata* or **H.** *nrp1a*, *chrna3*, *elavl3*, and *chrna3* with false

colour post-processing. False colour adjusted (fc) and elavl3 images have been brightened for enhanced

visibility.

**Figure 2. The preference for alcohol is biphasic.**

Gardener-Altman and Cumming estimation plots show alcohol response in three-minute intervals. No

stimulus was dispensed in the 'Pre' and 'Post' periods. Time spent in the stimulus zone during x̄0.5%

SAZA condition for **A.** WT (n = 31) and **D.** *chrna3*[C246X] mutants (n = 32). Time spent in the stimulus zone in

the x̄1.5% SAZA condition for **B.** WT (n = 30) and for **E.** *chrna3*[C246X] mutants (n = 30). **C.** Ethanol

concentrations in the stimulus zone in the x̄0.5% and x̄1.5% alcohol conditions and estimated for

extended periods. The concentration at which attraction switches to aversion is highlighted. Asterisks

indicate a significant paired mean difference between that time portion and the pre-stimulus period: * =

$p_{adj} < 0.05$ (provisional difference), ** = $p_{adj} < 0.01$ (meaningful difference). See Table S1 for the precise

effect sizes and p-values, corrected for multiple comparisons.

**Figure 3. *chrna3* mutation dynamically alters relative preference for alcohol.**

**A-C.** Gardener-Altman and Cumming estimation plots comparing the volume of alcohol dispensed over

the entire self-administration period during SAZA for **A.** WT (0%/x̄0.5%/x̄1.5%, n = 30/31/30), **B.**

*chrna3$^{C246X}$* (0%/x̄0.5%/x̄1.5%, n = 29/32/30) and **C.** intergenotype comparisons. Asterisks indicate the

following significant differences from 0% alcohol treatments (**A, B**), or the WT (**C**):  * = $p_{adj} < 0.05$, and

effect size reported by Cliff's delta between ± 0.2 and ± 0.4 (provisional difference), ** = $p_{adj} < 0.01$ and

effect size bigger than ± 0.4 (meaningful difference). **D-I.** Paired mean difference of preference index (PI)

between **D-F.** Pre and 0-9m or **G-I.** 0-9 and 9-18m time periods within genotypes, and delta delta

comparisons between genotypes, during **D, G.** 0%, **E, H.** x̄0.5%, and **F, I.** x̄1.5% SAZA. Asterisks indicate a

significant paired mean difference, or delta delta, between time periods: * = p < 0.05 (provisional

difference), ** = p < 0.01 (meaningful difference). See Table S2 for the exact effect size and p-values.

**Figure 4. *chrna3* mutants experience reduced psychoactive effects from alcohol exposure. A.**

Schematic outline of the shoaling assay metric calculation and processing steps for cohesion (shoal area)

and kinesis (shoal movement). **B, C.** Gardener-Altman, or **D, F, H.** mean (± 95% CI) shoal cohesion, and **E,**

**G, I.** kinesis, with Cumming estimation plots for the shoaling cohesion assay showing **B, D, E.** WT, **C, F, G.**

*chrna3$^{C246X}$*, and **H, I.** intergenotype comparisons.  Metrics recorded per frame at 2fps (n = 6 assays per

genotype, per treatment). Asterisks indicate a significant difference in the estimated marginal mean

from **B-G.** 0% alcohol, or **H, I.** WT within that time period:  * = $p_{adj} < 0.05$, and effect size reported by

Cliff's delta between ± 0.2 and ± 0.4 (provisional difference), ** = $p_{adj}$ < 0.01 and effect size greater than

± 0.4 (meaningful difference). See Table S3 for exact effect size and p-values.

**Figure 5. *chrna3*[C246X] mutants exhibit transcriptional changes in dependence, neuronal development,**

**and ion channel process associated genes.** RNAseq comparison of *chrna3*[C246X] (n = 3) vs *chrna3*[C246X] (n =

3) 12 week old zebrafish whole brain tissues. **A.** PCA clustering of samples by genotype. **B.** All

significantly enriched GO terms between genotypes. Positive enrichment values indicate enrichment in

the *chrna3*[C246X] mutant zebrafish. Significance thresholds are defined at $p$ = < 0.05 and FDR < 0.2. Yellow

= secondary messenger associated, Blue = ion channel associated, pink = neuronal differentiation and

development, green = behaviours. **C.** Volcano plot of gene expression changes between genotypes.

Up/down-regulated genes = $p$ < 0.05 & log2 fold change > ±2. Provisional up/down-regulated genes = $p$ <

0.05 & log2 fold change < ±2. Not significant genes = $p$ > 0.05. The colour key applies to D-F and I-K.

Positive fold change values indicate up-regulation in the *chrna3*[C246X] mutant zebrafish. **D-F.** Subsets of

volcano plot **C**, highlighting **D.** glutamate, **E.** GABA, and **F.** nicotinic acetylcholine (nAChR) receptor gene

families. **G.** Mean counts (± standard deviation) of nAChRs (gene prefix 'chrn'). Stars indicate significant

difference between genotypes, ** = p < 0.01, * = p < 0.05. **H.** Fold change in WT vs *chrna3*[C246X] of *chrna5*

(n = 21/20) and *chrnb4* (n = 22/22) by qRT-PCR. Stars indicate significant difference between genotypes,

** = p < 0.01, * = p < 0.05. **I-K.** Subsets of volcano plot **C**, highlighting genes associated with **I.** nicotine or

**J.** alcohol dependence (Hu et al., 2018), and **K.** anxiety phenotypes (Howe et al., 2016; Koskinen and

Hovatta, 2023). The full gene lists can be found in Table 4.

Movie Legends

**Movie 1:** A representative segment of the zebrafish self-administration assay (SAZA) with x̄1.5% alcohol

shown in Figure 1B.

**Movie 2:** A representative segment of the shoaling assay with 1% alcohol shown in figure 4D.

Supplementary Figure & Table Legends

**Figure S1.** Replicates of *chrna3* expression from figure 1, in *WT* (**A**) and *chrna3*$^{C246X}$ (**B**) 14dpf zebrafish

larvae facial motor nucleus (Fmn) marked by *chata* expression, visualised by *in-situ* hybridisation chain

reaction (HCR). Scale bars = 10 μm. **C.** Comparison of the mean fluorescent intensity (± standard

deviation) of *chrna3* in the demarcated zone between genotypes (Mean difference = -24.0 [95CI -26.3,-

20.9], p = 0.012).

**Figure S2.** Replicates of *chrna3* expression from figure 1, in *WT* (**A**) and *chrna3*$^{C246X}$ (**B**) 14dpf zebrafish

larvae right dorsal habenula (Hb) marked by *nrp1a* expression, visualised by *in-situ* hybridisation chain

reaction (HCR). Scale bars = 20 μm. **C.** Comparison of the mean fluorescent intensity (± standard

deviation) of *chrna3* in the demarcated zone between genotypes (Mean difference = 0.168 [95CI -2.84,

1.73], p = 0.944).

**Figure S3.** Gardener-Altman and Cumming estimation plots show alcohol response in three-minute

intervals. No stimulus was dispensed in the 'Pre' and 'Post' periods. Time spent in the stimulus zone

during 0% SAZA condition for **A.** WT (n = 31) and **B.** *chrna3*$^{C246X}$ mutants (n = 32). Time spent in the

control zone for **C, E, G.** WT (n = 30/31 /30) and **D, F, H.** *chrna3*$^{C246X}$ mutants (n = 29/32/30) in **C, D.** 0%,

**E, F.** x̄0.5% and **G, H.** x̄1.5% SAZA. Asterisks indicate a significant difference between that time portion

and the pre-stimulus period: * = $p_{adj}$ < 0.05 ** = $p_{adj}$ < 0.01. See Table S1 for the precise effect sizes and

p-values, corrected for multiple comparisons.

**Figure S4.** Gardener-Altman and Cumming estimation plots show alcohol response in three-minute

intervals. No stimulus was dispensed in the 'Pre' and 'Post' periods. Number of entries into the stimulus

zone for **A-C.** WT (n = 30/31 /30) and **D-F.** *chrna3*$^{C246X}$ mutants (n = 29/32/30) in **A, D.** 0%, **B, E.** x̄0.5% and

**C, F.** x̄1.5% SAZA. Asterisks indicate a significant difference between that time portion and the pre-

stimulus period: * = $p_{adj}$ < 0.05 ** = $p_{adj}$ < 0.01. See Table S1 for the precise effect sizes and p-values,

corrected for multiple comparisons.

**Table S1**. Statistical analysis output table for Figure 2 A-E, and Figures S3 and S4, intragenotype paired

mean difference of % time in stimulus, time in control, or number of entries, for repeated measures

against the 'Pre' period for 0%, x̄0.5% and x̄1.5% alcohol SAZA.

1002

**Figure S5.** Gardener-Altman and Cumming estimation plots for total SAZA self-administration period for

WT (**A, D**), *chrna3*$^{C246X}$ (**B, E**) and comparisons (**C, F**). Plots display the comparison of velocity inside (**A- C**)

or outside (**D-F**) the stimulus zone at different concentrations (0/x̄0.5/x̄1.5%) for WT (n = 30/31/30) and

*chrna3*$^{C246X}$ zebrafish (n = 29/32/30). **G-I.** Gardener-Altman and Cumming estimation plots for **G.** 0%, **H.**

x̄0.5%, and **I.** x̄1.5% SAZA. Plots compare preference index (PI) between WT (0/x̄0.5/x̄1.5% n = 30/ 31/30)

and *chrna3*$^{C246X}$ (0/x̄0.5/x̄1.5% n = 29/32/30) within each time period. No stimulus was dispensed in the

'Pre' and 'Post' periods. Asterisks indicate the following significant differences from 0% alcohol

treatments (**A, B, D, E**) or WT (**C, F-I**):  * = p < 0.05, and effect size reported by Cliff's delta between ± 0.2

and ± 0.4 are considered provisional, ** = p < 0.01 and effect size bigger than ± 0.4 is considered a

meaningful difference. See Table S2 for exact effect size and p-values.

1013

1014    **Figure S6**. Cliff's delta (± 95% CI) forest plots of all calculated metrics from 0/x̄0.5/x̄1.5% alcohol SAZA,

1015    comparing the response of wild type to *chrna3$^{C246X}$* mutant zebrafish. Positive values indicate a greater

1016    response in the mutant. See Table S2 for exact effect size and p-values.

1017

1018    **Table S2.** Statistical analysis output table for figure 3A, B, intragenotype, and figure 3C, intergenotype,

1019    comparisons of stimulus volume dispensed over the full self administration period for 0, x̄0.5 and x̄1.5%

1020    alcohol SAZA.  Statistical analysis output table for figure 3 D-I, paired mean difference and delta-delta in

1021    preference index (PI) between time periods, Figure S5 A-C, comparing preference index (PI), velocity,

1022    and Figure S6 other metrics for 0, x̄0.5 and x̄1.5% alcohol SAZA.

1023

**Figure S7**. ***WT* and *chrna3$^{C246X}$* mutants share shoaling cohesion and kinesis phenotypes during 0% and**

**1% alcohol immersion.** Mean (± 95% CI) shoal cohesion (**A, C**) and kinesis (**B, D**) with Cumming

estimation plots comparing 0% (**A, B**), or 1% alcohol exposure (**C, D**) between genotypes. Metrics

compared per frame at 2fps (n = 6 assays per genotype, per treatment). Asterisks indicate the following

significant differences from WT within that time period:  * = p < 0.05, and effect size reported by Cliff's

delta between ± 0.2 and ± 0.4 are considered provisional, ** = p < 0.01 and effect size bigger than ± 0.4

is considered a meaningful difference. See Table S3 for exact effect size and p-values.

1031

1032    **Table S3.** Statistical analysis output table for figure 4A-G, comparing shoaling behaviours within

1033    genotype, figure 4H-I and S7, between genotypes, displaying pairwise effect size comparisons, and

1034    repeated measures estimated marginal mean difference adjusted p values within time period.

# Figure 4

Raine, Kibat, *et. al* 2025

**Figure 3**

Raine, Kibat, *et. al* 2025