The neural mechanisms of fast versus slow decision-making

Mostafa Nashaat[1†*], Hatem Oraby[1†], Flutra Krasniqi[1], Sek Teng Goh-Sauerbier[1], Marion Bosc[2,3], Sandra Koerner[1], Sedef Karayel[4], Adam Kepecs[5], Matthew E. Larkum[1*]

[1]Institute for Biology, Humboldt University of Berlin, D-10117 Berlin, Germany
[2]Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA
[3]Neural circuits and immunity in psychosis laboratory, The Francis Crick Institute, London NW1 1AT, UK
[4]Charité – Universitätsmedizin Berlin, Einstein Center for Neurosciences Berlin, 10117, Berlin, Germany.
[5]Department of Neuroscience and Department of Psychiatry, Washington University in St. Louis, St. Louis, MO 63110, USA

[†]These authors contributed equally
[*]Corresponding authors: Matthew Larkum (matthew.larkum@hu-berlin.de), Mostafa Nashaat (mostafa.abdelhamid@hu-berlin.de)

Abstract

Is more haste less speed? Decision time variability has been attributed to speed/accuracy trade-offs(1), internal mental architecture(2), and noisy evidence accumulation(3-5). However, exploring these possibilities is difficult in rodents that consistently behave impulsively. Here, we demonstrate a novel floating-platform system that allows head-fixed mice to voluntarily vary decision times, akin to observed human behavior, in combination with sensitive neuroimaging approaches. We track the activity flow from medial to lateral frontal cortex (MFC to LFC) and record sequences of single-neuron activity. Choice-selective neurons displayed divergent temporal codes between MFC and LFC, with remarkable MFC susceptibility to optogenetic inhibition. These results suggest that LFC acts as an integrative motor threshold, while MFC plays a broader cognitive role in strategy and choice-selection.

Results

During natural decision-making paradigms in humans, there are many variables at play such as task history (priors), task difficulty (evidence), stakes (cost/reward) and attention(3, 5, 6). To disentangle the causal neural mechanisms underlying these factors, controlled experimental conditions are necessary and best studied in head-fixed mice. In this study, we developed a head-fixed mouse preparation combined with a floating nose-poke platform technology ("AirTrack"(7)), enabling the use of advanced neural acquisition and perturbation techniques, while simultaneously facilitating more natural, quasi-freely-moving behavior(8, 9). Head-fixed mice voluntarily navigated a nose-poke behavioral environment on a floating platform (Fig. 1A, B; fig. S1A), to perform a two-alternative forced-choice visual discrimination task (Video S1). To support self-paced evidence accumulation, we utilized the classical Random Dots Kinematogram (RDK) with free sampling adapted for rodents(10–12)(fig. S1B; "free sampling time", see methods). Mice were rewarded by moving in the direction that best matches the movement of the dots. By varying the dots' coherence, we could finely control stimulus salience and trial difficulty.

Unlike humans, freely moving mice performing simple detection tasks often display impulsive behavior under commonly used approaches even when sampling time is unrestricted (Fig. 1C, red histogram; fig. S1C)(13, 14). To address mice impulsivity, we adapted the RDK task to a floating platform, encouraging more voluntary control over sampling time and thus revealing richer decision-making patterns (Video S2). We have also conducted a comparable experiments in human subjects that involved speed-accuracy trade-off adjustment, adding a competitive time constraint to mirror the task design used in mice(15–17) (fig. S2A, see methods). In humans, this manipulation produced a clear shift from slow and accurate to fast and impulsive behavior (Fig. 1C, Speed = turquoise, n=20; Accuracy = violet, n=22; fig. S1C; fig. S2B). The combined reciprocal adjustments allowed us to align the human and mouse tasks, making their behavioral outputs directly comparable (fig. S2C).

Having established task comparability, we next asked what could be learned from the way sampling times vary across trials and conditions (Video S3). We began by dividing

the sampling time distribution according to trial difficulty ('easy', 'medium' & 'difficult'; see
methods). In mice, examining the trial-time distributions across all trials grouped by
difficulty, revealed that task difficulty had surprisingly little influence on sampling time (Fig.
1D). We then divided sampling times within each difficulty into thirds, labeling trials in
each segment as fast ('impulsive', red), typical ('typical', orange), or slow ('deliberate',
yellow) (Fig. 1D; n=20, session=262, trials=63,702 trials; example mouse, fig. S2D&E).
Performance scaled with both task difficulty and sampling duration, increasing with both
easy and slow trials (All mice, Fig. 1E; example mouse, fig. S2F). Next, we compared
psychometric curves across fast and slow trials (Fig. 1E). While both followed the
expected dependence on stimulus difficulty, slow decisions yielded significantly higher
accuracy than fast ones (All mice, Fig. 1E; example mouse, fig. S2F). The same pattern
was observed in humans performing the task in the speed context (Human speed: Fig.
1F, fig S2G&H; human accuracy: fig S2I-K), with fast trials showing significantly higher
lapse rate than slow ones (fig S2L).

Because fast and slow trials differed in performance, particularly on easier tasks, we
restricted our analysis to correct trials to examine whether sampling time reflected an
underlying decision strategy (Fig. 1G). In doing so, we observed a shift in policy for both
mice and humans (speed paradigm): subjects voluntarily increased sampling time for
difficult trials only during slow (deliberate) trials, but not during fast (impulsive) trials (mice,
Fig. 1G, left: fast, slope= 0.02, θ=1.08°; slow, slope=0.14, θ=7.83°; human speed, Fig.
1G, right: fast, slope= 0.06, θ=3.70°; slow, slope=0.74, θ=36.52°; human accuracy, fig
S3A). Voluntarily adjusting the time to consider the evidence during slow trials but not fast
trials suggests that factors other than difficulty can sometimes influence behavioral
strategy(*18–20*). We hypothesized that subjects update their decision-making strategy to
maximize an objective function, e.g. perceived reward outcome(*21–23*). Here, we found
that the average reward rate, i.e., performance over the last 5 trials, significantly
influenced sampling time in both mouse and human tasks (Fig. 1H, fig. S3B–D). Thus,
there was a tendency in mice to slow down if the previous trial was incorrect (Fig. 1I, left;
fig. S3E, F)(*24*). Furthermore, during fast decisions, mice were significantly biased to
switch sides if the previous trial was incorrect and a corresponding bias to stay if the
previous trial was correct (Fig. 1I, right, fig. S3G-I). This indicates that fast and slow
behavioral strategies arise from distinctively different behavioral outcomes in preceding
trials. We found that previous reward history had the highest explanatory power
(34.36%±9.94SEM), followed by evidence (27.15%±6.87SEM), trial history update
(21.02%±5.82SEM) and motor bias (19.32%±5.46SEM; fig. S3J; see methods).

We next explored what underlying computations might account for the variation in
sampling time and decision strategy. Drift-diffusion models (DDMs) are commonly used
to explain sampling time, but on their own, we found that they fell short of capturing key
aspects of strategic variability, particularly those linked to reward history(*4, 25*) (Fig. 1J;
fig. S4A). This led us to ask whether reinforcement learning components, which
incorporate trial history, might improve model predictions when combined with DDM(*25,
26*) (Fig. 1J, K). We tested two reinforcement learning models: Q-learning, where reward
sites are represented as states with associated action values, and R-learning, where the
average reward rate modulates integration noise (fig. S4A&B; see methods). Only when
both Q- and R-learning were combined with the classical DDM could we best capture the

behavioral phenotypes observed in mice, including the effects of reward rate on sampling
time and the fit of the psychometric function (Fig. 1J-L; "DDM+Q+R").

Spatiotemporal recruitment of frontal cortex mediates behavioral strategy.

To investigate the cellular mechanisms of fast versus slow decisions, we exploited the
floating platform approach in combination with widefield imaging and optogenetic
perturbation of specific cortical regions. For widefield calcium imaging, we used a
transgenic mouse line (Thy1 GP4.3) to express a genetically encoded calcium indicator
(GCaMP6s) in cortical excitatory neurons (Fig. 2A; and fig. S5A; see methods)(27), while
mice performed the RDK task using the voluntary-sampling paradigm (free sampling; Fig.
1A). Given the variable nature and timing of behavioral strategies (Fig. 1D), we first
focused on typical behavioral trials (sampling times of 1.0-1.2s; fig. S5B; see methods)
while using widefield imaging to characterize activity across the cortex (Fig. 2B&C; fig.
S5C&D)(28). We observed that all areas around the cortex were active at some point
during the sampling with a distinct spatio-temporal pattern originating from the medial axis
of the dorsal neocortex at the start of sampling (Fig. 2B, left), and traversing radially until
the animal made a decision (Fig. 2B and fig. S5E, decision made after last panel).

To identify the contribution of the different cortical areas, we used a selective optogenetic
perturbation approach in behaving head-fixed mice expressing genetically encoded ChR2
in all inhibitory neurons of mouse neocortex (VGAT-cre/ChR2; Fig. 2D , left; see
methods)(29, 30). Since the time of the animal's movement (i.e. taking the decision) was
unpredictable in the free-sampling paradigm, it was necessary to use a fixed-time
paradigm to confine the optogenetic inhibition to the entire sampling period without
overlapping with movement initiation (fig. S6A). We then applied bilateral optogenetic
inhibition to distinct cortical regions during a 1-second fixed sampling period (Fig. 2D, left
panel; primary visual cortex [V1], n=16 sessions from 7 mice; posterior parietal cortex
[PPC], n=12 sessions from 6 mice; retrosplenial cortex [RSP], n=9 sessions from 3 mice;
medial frontal cortex [MFC], n=17 sessions from 7 mice; lateral frontal cortex [LFC], n=21
sessions from 7 mice). Optogenetic silencing led to significant drops in performance,
primarily in frontal cortical regions, with the largest effects observed in MFC (28.90%±4.5
SEM) and LFC (21.53% ±6.27% SEM) (Fig. 2D, right panel; fig. S6B-F; two-tailed
permutation t-test; see methods). These findings are consistent with many previous
studies showing that proper recruitment of MFC and LFC over time is necessary for
decision-making(31, 32).

To investigate this further, we focused on the flow of calcium activity in the medial-lateral
axis of the frontal cortex (MFC→LFC) using the free-sampling paradigm (Fig. 2E). We
examined the spatiotemporal dynamics of the frontal calcium activity during the different
decision-making strategies, i.e. fast, typical and slow (Fig. 2E, n=6 mice, 18 sessions).
Across the spectrum from fast to slow strategies, frontal dynamics showed increasing
divergence between medial and lateral regions toward the end of the sampling period
(Fig. 2E&F; end). During typical and slow strategies, fluorescence activity in MFC was
antipodal (mirror-image) to activity in LFC, despite both contributing equally to
performance (Fig. 2D). Notably, LFC was most active just before the decision, when MFC

activity reached a minimum. This minimum was most pronounced in slow trials whereas
MFC activity at the end of fast trials resembled aborted decisions. Overall, the degree of
MFC-LFC antipodal activation at the end of sampling reflected behavioral strategy (Fig.
2F; fast end=0.39 ±0.06, typical end=0.66 ±0.05, slow end=0.73 ±0.04).

Consistent neural sequence activation during fast and slow behavior

These data imply that activity in MFC at the beginning of the sampling period contributes
more to the eventual decision than LFC, for which the crucial activity occurs at the end of
the sampling period(*33*). To test this hypothesis, we alternately inhibited both MFC and
LFC bilaterally either at the beginning or the end of the sampling period(*34, 35*) (Fig. 3A).
Note that since the timing of inhibition was crucial, it was necessary to use fixed sampling
to selectively block either the first (0-0.35s) or last third (0.65-1.0s) of the sampling phase
(Fig. 3B). We also repeated the widefield imaging experiments to confirm that the
spatiotemporal activity pattern across frontal cortex (MFC→LFC) was consistent under
fixed (1s) sampling (n=4 mice, sessions=12; Fig. 3B and fig. S7A&B). In accordance with
our hypothesis, optogenetic inhibition in the first third of sampling had a more potent effect
when applied to MFC (n=7, sessions=19,16.31% ±2.38% SEM) than LFC (n=10,
sessions=19, 8.79% ±2.56% SEM) (Fig. 3C; fig. S7C&D; Early inhibition; see methods).
However surprisingly, optogenetic inhibition at the end of sampling impaired performance
to a similar degree in both MFC (n=7, sessions=20, 15.18% ±3.20% SEM) and LFC (n=9,
sessions=22, 14.37% ±3.37% SEM; Fig. 3C; fig. S7E&F; Late inhibition; see methods),
showing that the decay in MFC activity does not result in a decline in its influence on the
decision.

The unexpectedly strong influence of MFC at the end of sampling, despite declining
widefield activity, raised the possibility that averaging signals across many neurons
obscured critical dynamics at the single-cell level. To address this, we used two-photon
microscopy to measure the activity of individual neurons directly in MFC and LFC during
decision-making (Figure 3D; n=6 mice, sessions=23; 609 active out of 1536 neurons in
LFC and 837 active out of 2340 neurons in MFC; see methods). Using this approach, we
found that the combined activity of cells recorded using two-photon imaging (Fig. 3E, top)
closely reflected the widefield activity (Fig. 3E; fig S8A&B). In particular, the observed
decrease in 2-photon activity at the end of sampling correlated with consistently less
active neurons in MFC (fig. S8C). This confirmed that the decline seen in the widefield
imaging signal was due to fewer active cells which were nevertheless surprisingly critical
for decision-making (Fig. 3C, right).Since sampling time differed across trials, we
normalized all neuronal activity traces to a common time base separately for fast, typical,
and slow strategies, enabling consistent comparison (fig S9A; Fig. 3F).

To characterize neural responses associated with different behavioral strategies, we
examined whether fast and slow trials recruit distinct neuronal populations. We found that
although the slow strategy engaged a slightly higher proportion of neurons, most neurons
were active during both fast and slow trials (Fig. 3G; fig. S9B), allowing for direct
comparison of their temporal activation patterns. To do this, we chronologically aligned
individual neurons according to their peak activity times to compare activation sequences
between fast and slow decision strategies (Fig. 3H; fig S9C). We next tested whether

these temporal sequences of neuronal peak activity were statistically consistent across trials, despite substantial differences in sampling duration between fast and slow strategies (Fig. 3H, I; fig. S9C&D)(*36–38*). By comparing to randomly generated permutations, we found similar activation sequences regardless of behavioral strategy (Fig. 3I; fig. S9D). To understand how neurons preserved their sequence rank order, we examined their calcium responses and found that peak activity was stretched or compressed relative to sampling duration(*39–41*) (area under the curve, AUC, and peak time; Fig. 3J bottom; fig. S9E–G; see methods). Neurons with early peak activity (predominantly located in MFC) typically maintained rigid firing patterns (Fig. 3K, top), whereas neurons peaking later often exhibited temporal stretching, a pattern particularly prevalent in LFC (Fig. 3K, bottom; fig. S9G).

These results remained consistent at the population level; averaging calcium activity across all layer 2/3 neurons yielded comparable dynamics at the single cell level to those observed under widefield imaging(*42*) (Fig. S8). Consistently, we also found that the amplitude of population activity at the start of sampling in MFC and not LFC was significantly correlated with sampling duration and easy trial's performance(*43*) (fig. S10A; fig. S8D). Additionally, MFC neurons showed a tendency to fire sparsely but reliably immediately prior to the decision while LFC exhibited an increasing neuronal engagement with consistent activity levels toward the end of sampling (fig. S10B). In conclusion, optogenetic inhibition and two-photon imaging revealed that the neural mechanisms underlying voluntary decision-making involve both decreasing neural activity in MFC (sparsity) and increasing neuronal engagement in LFC at the time of decision.

Two modes of operation in MFC versus LFC

To investigate this further, we examined the 2-photon single-neuron responses based on the animal's choice (Fig. 4A). Interestingly, the pattern of activity during the sampling was very similar for both left and right choices(*37*) (fig S11A). Nevertheless, we identified a subset of neurons as choice-selective, when the peak mean activity during sampling was significantly higher for one choice direction (Fig. 4A, green, "choice-matched") versus the other (Fig. 4A, brown, "choice-mismatched"). Using this definition, neurons fell into two categories; tuned (either left or right) or non-selective. The peak activity of choice-selective neurons occurred at different times across the whole period of sampling(*44, 45*) (Fig. 4B; fig S11B). We observed a higher percentage of choice-encoding neurons in LFC versus MFC across all behavioral strategies (Fig, 4C; 33.07 ±3.82%SEM versus 18 ±-3%SEM). However, interestingly, we found that selectivity could vary as a function of behavioral strategy even within a single neuron (Fig. 4D; fig S11C). Next, we asked whether the evolution of choice-selectivity across the sequence of neuronal firing reflected the mode of operation of MFC versus LFC. To quantify this, we binned neurons according to their temporal peak activity and compared matched versus mismatched responses for each bin (Fig. 4E, top panel, slow and bottom panel, fast; MFC 10 session, 182 neurons; LFC 10 sessions, 197 neurons; see methods). For comparison, we also overlaid the average population 2-photon activity from all L2/3 neurons (dotted lines; cf. fig. S8) on the choice-selectivity histograms.

We conclude that slow behavior commenced from a more balanced state of choice
representation (matched vs mismatched) at the start of sampling, a trend consistent in
both MFC and LFC. In contrast, fast behavior was characterized by an initial state of
imbalance in matched versus mismatched activity. Additionally, a consistent observation
across all conditions was the emergence of an imbalance in choice representation by the
end of the sampling period (fig S11D&E).

In general, these results can also be interpreted using a metric we term "coding
efficiency," defined as (matched – mismatched) / matched, reflecting the propensity of
neural activity to favor the animal's chosen direction at any stage in the sequence (Fig.
4F; fig. S11F). This metric allows for quantification of how neuronal activity evolves to
influence choice selection. We observed that coding efficiency substantially increased
during the sampling period for slow behavioral trials but decreased for fast trials. To
further elucidate the neural mechanisms underlying these differential strategies within the
DDM+Q+R computational framework (Fig. 1I-K), we found that a substantial fraction of
neurons multiplexed signals encoding previous trial outcomes together with
representations of previous or current choices (fig. S12A&B). Notably, reward history
significantly sharpened neuronal selectivity for both previous and current choice signals
during the early phase of sampling (Fig. S12C). Given the limited availability of sensory
evidence at the early stage of sampling, we interpret this early imbalance representation
as a deviation in the starting offset or noise signal during evidence accumulation. This
reward-history-driven bias provides a mechanistic link to the fast trial strategy, marked by
strong priors and elevated coding efficiency at the start of sampling (Fig. 1I; Fig. 4F; fig
S13).

Our data so far suggest that MFC and LFC operate in distinct but complementary modes
during decision-making. Both regions exhibit improved choice coding efficiency during
slow trials and reduced efficiency during fast trials (Fig. 4F). Yet they appear to achieve
this via different mechanisms. In MFC, this improvement is driven by a refinement process
in which the number of neurons representing choice decreases more steeply for
mismatched activity than for choice-aligned signals (Fig. 4E, left, fig S14). In contrast, the
number of active LFC neurons appears to gradually increase over sampling to a
consistent threshold value for triggering action (Fig. 4E, right, fig S14). The distinct
temporal trajectories in MFC and LFC, despite converging coding efficiency, suggest they
may contribute differently to how information is sampled and integrated over time(5).

To test this idea, we asked whether interfering with the sequence of firing during the
sampling period would differentially affect behavior. Specifically, we inhibited MFC or LFC
midway through free sampling (0.3–0.9s), interrupting the sequence and decision-making
process (Fig. 4G, blue shaded area). Inhibiting LFC briefly disrupted the decision-making
process, reflected in sampling duration and performance, which quickly recovered after
the end of inhibition (Fig. 4G-H, magenta; fig. S15). In contrast, inhibiting MFC caused
much longer delays in sampling and a sustained drop in performance which did not
recover even with long sampling times (Fig. 4G–H, blue; fig. S15). This indicates that
decision-making is more sensitive to the sequence of activation of neurons in MFC than
LFC.

These findings lead us to revisit the conceptual framework of serial sampling, highlighting
the distinct contributions of MFC and LFC to decision-making (Fig. 4I). In this framework,
MFC generates a sequential refinement of choice representations, culminating in a small
population of neurons encoding the eventual choice (Fig. 3C & Fig. 4G&H). Disrupting
this sequence at any point impairs performance in a sustained manner, suggesting that
MFC actively tracks the evolving state of evidence. In contrast, LFC shows a rapid
rebound following transient inhibition, implying that it does not accumulate information
locally but rather monitors upstream signals (possibly including MFC) to determine when
to act. This interpretation aligns with recent findings that LFC activity reflects the decision
state but may not integrate evidence directly(*46, 47*).

Taken together, these results suggest that MFC plays a central role in computing what
decision to make, while LFC determines when to act. Crucially, our data also imply that
the MFC may exert an inhibitory influence over downstream motor initiation, releasing this
brake only once its internal refinement process converges. In this framework, disrupting
MFC not only degrades choice quality but also delays commitment, whereas disrupting
LFC mainly affects the timing of action execution. This division of labor between
refinement and commitment offers a mechanistic account of how serial sampling is
coordinated across frontal regions and may generalize to other forms of voluntary
decision-making.

References

1. D. Kahneman, *Thinking, Fast and Slow* (Farrar, Straus and Giroux, New York, NY, US, 2011)*Thinking, fast and slow*.

2. D. Luce, "Response Times: Their Role in Inferring Elementary Mental Organization" (1986; https://www.semanticscholar.org/paper/Response-Times%3A-Their-Role-in-Inferring-Elementary-Luce/4f714b26b7ceb23ccad0e47d19a6fd94ec10bf4f).

3. R. Bogacz, E. Brown, J. Moehlis, P. Holmes, J. D. Cohen, The physics of optimal decision making: A formal analysis of models of performance in two-alternative forced-choice tasks. *Psychological Review* **113**, 700–765 (2006).

4. R. Ratcliff, G. McKoon, The Diffusion Decision Model: Theory and Data for Two-Choice Decision Tasks. *Neural Comput* **20**, 873–922 (2008).

5. J. I. Gold, M. N. Shadlen, The Neural Basis of Decision Making. *Annual Review of Neuroscience* **30**, 535–574 (2007).

6. P. W. Glimcher, The Neurobiology of Visual-Saccadic Decision Making. *Annual Review of Neuroscience* **26**, 133–179 (2003).

7. M. A. Nashaat, H. Oraby, R. N. S. Sachdev, Y. Winter, M. E. Larkum, Air-Track: a real-world floating environment for active sensing in head-fixed mice. *Journal of Neurophysiology* **116**, 1542–1553 (2016).

8. S. E. Dominiak, M. A. Nashaat, K. Sehara, H. Oraby, M. E. Larkum, R. N. S. Sachdev, Whisking Asymmetry Signals Motor Preparation and the Behavioral State of Mice. *J. Neurosci.* **39**, 9818–9830 (2019).

9. J. Voigts, M. T. Harnett, Somatic and Dendritic Encoding of Spatial Variables in Retrosplenial Cortex Differs during 2D Navigation. *Neuron* **105**, 237-245.e4 (2020).

10. T. Marques, M. T. Summers, G. Fioreze, M. Fridman, R. F. Dias, M. B. Feller, L. Petreanu, A Role for Mouse Primary Visual Cortex in Motion Perception. *Current Biology* **28**, 1703-1713.e6 (2018).

11. M. N. Shadlen, R. Kiani, Decision Making as a Window on Cognition. *Neuron* **80**, 791–806 (2013).

12. L. Hernández-Navarro, A. Hermoso-Mendizabal, D. Duque, J. de la Rocha, A. Hyafil, Proactive and reactive accumulation-to-bound processes compete during perceptual decisions. *Nat Commun* **12**, 7148 (2021).

13. C. Schwarz, H. Hentschke, S. Butovas, F. Haiss, M. C. Stüttgen, T. V. Gerdjikov, C. G. Bergner, C. Waiblinger, The head-fixed behaving rat—Procedures and pitfalls. *Somatosensory & Motor Research* **27**, 131–148 (2010).

14. J. I. Sanders, A. Kepecs, Choice ball: a response interface for two-choice psychometric
discrimination in head-fixed mice. *Journal of Neurophysiology* **108**, 3416–3423 (2012).

15. B. a. J. Reddi, R. H. S. Carpenter, The influence of urgency on decision time. *Nat Neurosci*
3, 827–830 (2000).

16. J. Palmer, A. C. Huk, M. N. Shadlen, The effect of stimulus strength on the speed and
accuracy of a perceptual decision. *Journal of Vision* **5**, 1 (2005).

17. J. Drugowitsch, R. Moreno-Bote, A. K. Churchland, M. N. Shadlen, A. Pouget, The Cost of
Accumulating Evidence in Perceptual Decision Making. *J. Neurosci.* **32**, 3612–3628
(2012).

18. N. A. Steinemann, G. M. Stine, E. M. Trautmann, A. Zylberberg, D. M. Wolpert, M. N.
Shadlen, Direct observation of the neural computations underlying a single decision. *eLife*
12 (2023).

19. Z. C. Ashwood, N. A. Roy, I. R. Stone, The International Brain Laboratory, A. E. Urai, A. K.
Churchland, A. Pouget, J. W. Pillow, Mice alternate between discrete strategies during
perceptual decision-making. *Nat Neurosci* **25**, 201–212 (2022).

20. S. Pisupati, L. Chartarifsky-Lynn, A. Khanal, A. K. Churchland, Lapses in perceptual
decisions reflect exploration. *eLife* **10**, e55490 (2021).

21. J. Drugowitsch, G. C. DeAngelis, D. E. Angelaki, A. Pouget, Tuning the speed-accuracy
trade-off to maximize reward rate in multisensory decision-making. *eLife* **4**, e06678 (2015).

22. P. Simen, D. Contreras, C. Buck, P. Hu, P. Holmes, J. Cohen, Reward rate optimization in
two-alternative decision making: empirical tests of theoretical predictions. *J Exp Psychol*
Hum Percept Perform **35**, 1865–1897 (2009).

23. Acquisition of decision making criteria: reward rate ultimately beats accuracy | Attention,
Perception, & Psychophysics. https://link.springer.com/article/10.3758/s13414-010-0049-7.

24. N. S. Narayanan, M. Laubach, Neuronal Correlates of Post-Error Slowing in the Rat
Dorsomedial Prefrontal Cortex. *Journal of Neurophysiology* **100**, 520–525 (2008).

25. A. E. Urai, J. W. De Gee, K. Tsetsos, T. H. Donner, Choice history biases subsequent
evidence accumulation. *eLife* **8**, e46331 (2019).

26. S. Shuvaev, S. Starosta, D. Kvitsiani, A. Kepecs, A. Koulakov, "R-learning in actor-critic
model offers a biologically relevant mechanism for sequential decision-making" in
Advances in Neural Information Processing Systems (Curran Associates, Inc., 2020;
https://papers.nips.cc/paper/2020/hash/da97f65bd113e490a5fab20c4a69f586-
Abstract.html)vol. 33, pp. 18872–18882.

27. H. Dana, T.-W. Chen, A. Hu, B. C. Shields, C. Guo, L. L. Looger, D. S. Kim, K. Svoboda,
Thy1-GCaMP6 Transgenic Mice for Neuronal Population Imaging In Vivo. *PLOS ONE* **9**,
e108697 (2014).

28. J. Couto, S. Musall, X. R. Sun, A. Khanal, S. Gluf, S. Saxena, I. Kinsella, T. Abe, J. P.
Cunningham, L. Paninski, A. K. Churchland, Chronic, cortex-wide imaging of specific cell
populations during behavior. *Nat Protoc* **16**, 3241–3263 (2021).

29. Z. V. Guo, N. Li, D. Huber, E. Ophir, D. Gutnisky, J. T. Ting, G. Feng, K. Svoboda, Flow of
Cortical Activity Underlying a Tactile Decision in Mice. *Neuron* **81**, 179–194 (2014).

30. L. Pinto, K. Rajan, B. DePasquale, S. Y. Thiberge, D. W. Tank, C. D. Brody, Task-
Dependent Changes in the Large-Scale Dynamics and Necessity of Cortical Regions.
Neuron **104**, 810-824.e9 (2019).

31. M. Laubach, L. M. Amarante, K. Swanson, S. R. White, What, If Anything, Is Rodent
Prefrontal Cortex? *eNeuro* **5**, ENEURO.0315-18.2018 (2018).

32. F. Barthas, A. C. Kwan, Secondary Motor Cortex: Where 'Sensory' Meets 'Motor' in the
Rodent Frontal Cortex. *Trends in Neurosciences* **40**, 181–193 (2017).

33. T.-W. Chen, N. Li, K. Daie, K. Svoboda, A Map of Anticipatory Activity in Mouse Motor
Cortex. *Neuron* **94**, 866-879.e4 (2017).

34. T. D. Hanks, C. D. Kopec, B. W. Brunton, C. A. Duan, J. C. Erlich, C. D. Brody, Distinct
relationships of parietal and prefrontal cortices to evidence accumulation. *Nature* **520**, 220–
223 (2015).

35. L. Pinto, D. W. Tank, C. D. Brody, Multiple timescales of sensory-evidence accumulation
across the dorsal cortex. *eLife* **11**, e70263 (2022).

36. M. Laubach, J. Wessberg, M. A. L. Nicolelis, Cortical ensemble activity increasingly
predicts behaviour outcomes during learning of a motor task. *Nature* **405**, 567–571 (2000).

37. C. D. Harvey, P. Coen, D. W. Tank, Choice-specific sequences in parietal cortex during a
virtual-navigation decision task. *Nature* **484**, 62–68 (2012).

38. A. J. Peters, S. X. Chen, T. Komiyama, Emergence of reproducible spatiotemporal activity
during motor learning. *Nature* **510**, 263–267 (2014).

39. J. Wang, D. Narain, E. A. Hosseini, M. Jazayeri, Flexible timing by temporal scaling of
cortical responses. *Nat Neurosci* **21**, 102–110 (2018).

40. H. Sohn, D. Narain, N. Meirhaeghe, M. Jazayeri, Bayesian Computation through Cortical
Latent Dynamics. *Neuron* **103**, 934-947.e5 (2019).

41. M. I. Leon, M. N. Shadlen, Representation of time by neurons in the posterior parietal cortex
of the macaque. *Neuron* **38**, 317–327 (2003).

42. Task-Dependent Changes in the Large-Scale Dynamics and Necessity of Cortical Regions: Neuron. https://www.cell.com/neuron/fulltext/S0896-6273(19)30731-7.

43. K. G. Thompson, K. L. Biscoe, T. R. Sato, Neuronal Basis of Covert Spatial Attention in the Frontal Eye Field. *J. Neurosci.* **25**, 9479–9487 (2005).

44. B. B. Scott, C. M. Constantinople, A. Akrami, T. D. Hanks, C. D. Brody, D. W. Tank, Fronto-parietal Cortical Circuits Encode Accumulated Evidence with a Diversity of Timescales. *Neuron* **95**, 385-398.e5 (2017).

45. S. A. Koay, A. S. Charles, S. Y. Thiberge, C. D. Brody, D. W. Tank, Sequential and efficient neural-population coding of complex task information. *Neuron* **110**, 328-349.e11 (2022).

46. Z. Yang, M. Inagaki, C. R. Gerfen, L. Fontolan, H. K. Inagaki, Integrator dynamics in the cortico-basal ganglia loop underlie flexible motor timing. bioRxiv [Preprint] (2024). https://doi.org/10.1101/2024.06.29.601348.

47. H. K. Inagaki, S. Chen, M. C. Ridder, P. Sah, N. Li, Z. Yang, H. Hasanbegovic, Z. Gao, C. R. Gerfen, K. Svoboda, A midbrain-thalamus-cortex circuit reorganizes cortical dynamics to initiate movement. *Cell* **185**, 1065-1081.e23 (2022).

48. J. R. de Leeuw, R. A. Gilbert, B. Luchterhandt, jsPsych: Enabling an Open-Source Collaborative Ecosystem of Behavioral Experiments. *Journal of Open Source Software* **8**, 5351 (2023).

49. S. Rajananda, H. Lau, B. Odegaard, A Random-Dot Kinematogram for Web-Based Vision Research. *Journal of Open Research Software* **6** (2018).

50. S. Musall, M. T. Kaufman, A. L. Juavinett, S. Gluf, A. K. Churchland, Single-trial neural dynamics are dominated by richly varied movements. *Nat Neurosci* **22**, 1677–1686 (2019).

51. D. H. Brainard, The Psychophysics Toolbox. *Spat Vis* **10**, 433–436 (1997).

52. J. Peirce, J. R. Gray, S. Simpson, M. MacAskill, R. Höchenberger, H. Sogo, E. Kastman, J. K. Lindeløv, PsychoPy2: Experiments in behavior made easy. *Behav Res* **51**, 195–203 (2019).

53. T. D. Hanks, C. Summerfield, Perceptual Decision Making in Rodents, Monkeys, and Humans. *Neuron* **93**, 15–31 (2017).

54. R. Ratcliff, F. Tuerlinckx, Estimating parameters of the diffusion model: Approaches to dealing with contaminant reaction times and parameter variability. *Psychonomic Bulletin & Review* **9**, 438–481 (2002).

55. M. Pachitariu, C. Stringer, M. Dipoppa, S. Schröder, L. F. Rossi, H. Dalgleish, M. Carandini, K. D. Harris, Suite2p: beyond 10,000 neurons with standard two-photon microscopy. bioRxiv [Preprint] (2017). https://doi.org/10.1101/061507.

Acknowledgement: We thank Moritz Drüke and Arina Lobova from the Larkum Lab for insightful discussions. We are also grateful to all members of the Larkum Lab for their valuable input and to the supporting staff for assistance with organizational tasks. Special thanks to Alexander Schill and Jan Ode from the Research Workshop at Charité – Universitätsmedizin Berlin for developing and manufacturing the experimental devices. We deeply appreciate the support of Dr. Claudia Abramuik and the Charité Mitte animal facility for providing access to facility space and equipment, as well as for their assistance in breeding various mouse lines. Special thanks to Joao Couto for advice on using the widefield preprocessing library.

Funding: Deutsche Forschungsgemeinschaft (SFB1315, Grant No. LA 3442/6-1). European Union Horizon 2020 Research and Innovation Programme (670118/ERC ActiveCortex, 101055340/ERC Cortical Coupling to M.E.L.). Einstein Foundation Berlin (Projekt A-2016-311; EVF-2019-508).

Competing interests: The authors declare no competing interests.

Data and materials availability: All data required to evaluate the conclusions of this study are included in the main text and/or the Supplementary Materials. Source data and analysis code will be deposited in a public repository and made available prior to publication.

Figures

 Fig. 1. A novel behavioral paradigm demonstrates similar adaptive decision-
 making strategies in mice and humans. (**A**) Schematic of the Random Dots
 Kinematogram (RDK) task for head-fixed mice using a nose-poke floating platform. (**B**)
 3D illustration of the floating-platform apparatus with three nose-pokes designed for
 voluntary decision-making tasks in head-fixed mice. (**C**) Comparison of sampling time

distributions in mice and human highlighting flexible decision-making strategies across
different contexts, including freely moving mice performing a visual detection task (red),
head-fixed mice on the floating platform performing the RDK task (orange), and humans
performing the RDK task under accuracy condition (violet) or speed condition (turquoise)
instructions. These distributions show a continuum from impulsive ("fast") to deliberate
("slow") decision-making. (**D**) Histograms of sampling times categorized into fast (red),
typical (orange), and slow (yellow) trials across stimulus difficulties (easy, moderate,
hard). (**E**) Performance accuracy (% correct choices) in head-fixed mice as a function of
task difficulty (motion coherence %) for fast (red) versus slow (yellow) trials. (**F**) Human
performance accuracy during speed-instructed RDK trials, formatted as in panel E. (**G**)
Mean sampling time (z-scored) of correct trials across stimulus difficulties during fast (red)
and slow (yellow) trials for both head-fixed mice and human subjects (speed condition).
(**H**) Mean sampling time (z-scored) as a function of recent reward history (reward rate
calculated from the preceding 5 trials) in both head-fixed mice and humans (speed
paradigm), illustrating behavioral sensitivity to reward feedback. (**I**) Left: Probability of
adopting fast versus slow strategies based on previous trial outcomes (correct/incorrect)
in mice. Right: Win-stay/Lose-switch behavioral update probability during fast versus slow
decisions in mice. (**J**) Schematic representation of reinforcement learning (RL)-enhanced
drift diffusion models (DDM) tested to capture mouse behavior. (**K**) Left: Reward rate
plotted against z-scored sampling time from an example mouse (avgat_1 = 2501 trials;
dashed gray line) compared to predictions from different RL-enhanced DDM variants:
basic DDM (black), Q-learning combined DDM (DDM+Q, red), R-learning combined DDM
(DDM+R, blue), and Q-learning & R-learning combined DDM (DDM+Q+R, gold). Right:
Psychometric curve of the same example mouse for fast (red dots) and slow (yellow dots)
trials, along with fits from the Q&R-DDM model (lines). (**L**) Pearson correlation (R^2) values
for model fits (DDM, DDM+Q, DDM+R, DDM+Q+R) and mice behavioral measures:
psychometric performance for fast and slow conditions ("Psy"), and sampling time/reward
rate sensitivity ("R").

480
Fig. 2. Spatiotemporal activation of frontal cortex regions causally relates to fast
and slow decision-making strategies. (**A**) Schematic illustrating the widefield calcium
imaging setup used in head-fixed mice. (**B**) Pixel-wise heatmaps depicting average
calcium fluorescence (dF/F) in an example Thy1-GCaMP6s mouse across three different
time windows during a typical sampling interval (1.0–1.2 s) in the RDK decision-making
task. Cortical area boundaries are outlined for reference according to Allen brain atlas.
(**C**) Mean normalized calcium responses across different cortical areas indicated in panel
B (primary visual cortex, V1; posterior parietal cortex, PPC; retrosplenial cortex, RSC;
primary motor cortex, M1; secondary motor cortex, M2), averaged across 6 sessions in 4
mice for trials between 1 to 1.2 seconds. Shaded regions represent SEM. (**D**) Behavioral
impact of bilateral optogenetic silencing (using VGAT-ChR2 mice) targeting distinct
cortical regions during a fixed, 1-s sampling RDK task. The cortical map illustrates
targeted areas: V1 (visual), PPC (posterior parietal), RSC (retrosplenial), MFC (medial
frontal cortex), and LFC (lateral frontal cortex). Boxplot quantifies the percentage drop in
task performance following cortical silencing (*p < 0.05, **p < 0.01, ***p < 0.001). (**E**)
Functional segmentation of frontal cortex activity into medial frontal cortex (MFC; blue)
and lateral frontal cortex (LFC; magenta), based on distinct spatiotemporal activation
patterns during the sampling period. Representative heatmaps illustrate average calcium
activity patterns for fast (impulsive), typical, and slow (deliberate) decision-making
strategies. Corresponding normalized calcium fluorescence signals for MFC and LFC
(n=6 mice) are plotted below each heatmap. The double-headed arrow labeled "δ"

indicates the magnitude of the difference in fluorescence between MFC and LFC at the
end of the sampling period. (**F**) Quantification of the average difference (δ) in normalized
calcium activity between MFC and LFC at the end of sampling across fast (impulsive),
typical, and slow (deliberate) strategies. Error bars represent SEM. This difference
systematically increases from fast to slow strategies.

507
Fig. 3. Optogenetic inhibition and two-photon imaging reveal distinct temporal
dynamics and neuronal activation sequences in MFC and LFC during fast and slow
decisions. (**A**) Schematic depicting bilateral optogenetic inhibition of medial frontal cortex
(MFC, blue) and lateral frontal cortex (LFC, magenta). (**B**) Top: Experimental design for
targeted optogenetic inhibition in two distinct time windows (early: 0–350 ms; late: 650–
1000 ms) during a fixed 1-s sampling time. Bottom: Average frontal widefield calcium
fluorescence during 1-s fixed sampling time (n=4 mice; MFC in blue, LFC in magenta).
Shaded intervals highlight early and late inhibition windows. (**C**) Behavioral effect (%
performance drop) of early (left) and late (right) optogenetic inhibition of MFC (blue) and
LFC (magenta). Significant effects indicated (*p < 0.05; **p < 0.01; ***p < 0.001; n.s., not
significant). (**D**) Left: Schematic of two-photon imaging setup targeting calcium activity of
layer 2/3 neurons in MFC and LFC. Middle: Example micrograph showing neurons in the
imaged cortical area. Right: Example calcium activity traces (dF/F) from three selected
neurons (color-coded) highlighting fast (red dotted frame) and slow (yellow dotted frame)
trials. (**E**) Mean normalized population calcium activity from layer 2/3 neurons (solid lines)
and corresponding widefield signals overlayed (dashed lines) in MFC (blue) and LFC
(magenta) during fast (left) and slow (right) behavioral strategies. Shaded areas represent
SEM. (**F**) Single-trial neuronal calcium activity traces illustrating active (green) and
inactive (gray) trials during fast (top) and slow (bottom) trials. Mini-heatmap reflects

neuron's temporal peak of average activity. (**G**) Venn diagrams showing proportions of
active (≥10% active trials) neurons selective for fast (red), slow (yellow), or both
behavioral strategies (red/yellow dashed area), in MFC (top) and LFC (bottom). (**H**)
Heatmap displaying neuronal population activity during fast (left) and slow (right)
strategies from an example MFC session, normalized and ordered by neuronal activation
peak during all trials. (**I**) Schematic illustrating the method used to quantify deviations in
the neuronal activation sequence (temporal rank) between fast and slow strategies. (**J**)
Left: Permuted behavioral strategies rank deviation null distribution (histogram) versus
observed rank deviation (vertical dashed line) in an example LFC session. Right: Scatter
plot shows rank deviation (z-scored) for neuronal sequences in individual imaging
sessions of MFC (blue dots, n = 11 sessions) and LFC (magenta dots, n = 13 sessions).
Significant deviations from chance are indicated (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$). (**K**)
Left: Examples of neuron classification based on temporal dynamics. An example of a
"rigid" neuron (top) shows no significant correlation ($r^2 < 0.3$) of activity peak timing or
area under the curve (AUC) with sampling duration. An example of a "stretching" neuron
(bottom) with significantly positive correlation ($r^2 > 0.3$) between sampling duration and
both peak timing and AUC (see methods). Right: Pie charts summarizing the proportion
of rigid (grey) vs. stretching neurons in MFC (top) and LFC (bottom). Stretching neurons
are defined as those with a positive correlation ($r^2 > 0.3$) between sampling duration and
either neural activity peak timing (brown), AUC (orange), or both (gold) (see Methods).

Fig. 4. Choice-selective neural sequences in MFC and LFC exhibit distinct modes of operation, with differential contributions to voluntary decision-making. (**A**) Schematic illustrating two-photon calcium imaging of layer 2/3 neurons in MFC and LFC. Neurons are classified as choice-selective based on differential activity patterns during trials in which neuronal preference matches ("choice-match") or mismatches ("choice-mismatch") the mouse's actual choice direction (left or right). (**B**) Left: Heatmap from an

example session (MFC) showing the temporal sequence of neuronal activation throughout the sampling period. Right: Three example neurons illustrating activity profiles—selective preference for leftward choice (#9, top), non-selective (#38, middle), and selective preference for rightward choice (#46, bottom). Green/brown traces denote matched/mismatched choices respectively; shading indicates SEM. (**C**) Pie charts quantifying the proportion of choice-selective neurons in MFC (right) and LFC (left). (**D**) Example neuron (#9 from panel B) demonstrating differential responses (choice selectivity) between choice-matched (green) and choice-mismatched (brown) trials during slow (top) and fast (bottom) decision-making strategies. The blue frame indicates the temporal bin of this neuron's activity in panel E. (**E**) Population responses of sequentially activated choice-selective neurons (ordered by temporal peak, indicated by arrows) for matched (green) and mismatched (brown) trials during correct trials, shown separately for slow (top) and fast (bottom) strategies in MFC (left) and LFC (right). Dotted curves illustrate the corresponding average sum of population activity overlayed. (**F**) Coding efficiency, quantifying the separation between matched and mismatched responses, at sampling start and decision points in MFC (blue) and LFC (magenta) during fast and slow behavioral strategies. (**G**) Top: Experimental design schematic showing timing of bilateral optogenetic inhibition of MFC (blue) or LFC (magenta) during a free-sampling task. Bottom: Example mouse showing probability distributions of sampling durations for control (gray), MFC-inhibited (blue), and LFC-inhibited (magenta) conditions. Shaded region indicates inhibition period (600 ms). (**H**) Left: Cumulative distribution of sampling durations across all mice for control (gray), MFC inhibition (blue), and LFC inhibition (magenta). Right: Behavioral performance (percentage correct) following the offset of inhibition across the sampling time, comparing control, MFC-inhibited, and LFC-inhibited trials. (**I**) Summary schematic illustrating how different parameters (e.g., reward rate, sensory evidence, previous choice) shape decision-making via distinct modes of frontal cortex operation. MFC primarily refines decisions by integrating these parameters gradually, while LFC facilitates commitment to action once evidence crosses a motor threshold, particularly differentiating fast from slow strategies.

Supplementary Materials for

The neural mechanisms of fast versus slow decision-making

Mostafa Nashaat[1†*], Hatem Oraby[1†], Flutra Krasniqi[1], Sek Teng Goh-Sauerbier[1], Marion
Bosc[2,3], Sandra Koerner[1], Sedef Karayel[4], Adam Kepecs[5], Matthew E. Larkum[1*]

[1]Institute for Biology, Humboldt University of Berlin, D-10117 Berlin, Germany
[2]Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA
[3]Neural circuits and immunity in psychosis laboratory, The Francis Crick Institute,
London NW1 1AT, UK
[4]Charité – Universitätsmedizin Berlin, Einstein Center for Neurosciences Berlin, 10117,
Berlin, Germany.
[5]Department of Neuroscience and Department of Psychiatry, Washington University in
St. Louis, St. Louis, MO 63110, USA

[†]These authors contributed equally
*Corresponding authors: Matthew Larkum (matthew.larkum@hu-berlin.de), Mostafa
596   Nashaat (mostafa.abdelhamid@hu-berlin.de)

597   **This PDF file includes:**
598   Materials and Methods
599   Figs. S1 to S15

Materials and Methods

Human psychophysical Experiment

The study was conducted at Humboldt university and approved by the local ethics committee of the Humboldt University Berlin. 56 students constituted the pool of participating subjects. As students participated on a voluntary basis but varying motivation, only the data of the top 30% best performing subjects of each group cohort were used. Each subject final score was based on their performance in both accuracy and maximum correct outcome experiments (n=18 Subjects). The experiment was conducted on 6 group cohorts, each involving 9-13 subjects. Subjects were assigned a computer rig (Dell Optiplex 7400 AIO Series), were made aware of the scoring criteria and were encouraged to compete for the highest score. Most users ran both accuracy and maximum correct outcome experiments, with each experiment duration set to 30 minutes including a 5-minute break at the 15-minute mark, as well as a mandatory 20-second break every 100 trials. Subjects performed an initial 10 test trials. For the accuracy competition, a subject's score is calculated as the percentage of correct trials over the total trials, with a minimum requirement of 50 trials. For the maximum correct outcome competition, the subject's score is the total number of correct trials. A central server controlled by the experimenter activated a countdown timer to synchronize the start of the session on all subjects' computers, thus ensuring all subjects are participating concurrently. Subjects received correct/incorrect feedback after each trial and could track their ranking among other players, viewing the leaderboard during individual 20-second breaks and the 15-minute group break. The experiments were implemented using jsPsych: and utilizing the jspsych-rdk plugin(*48, 49*). The stimulus comprised of 1,000 white dots on a black background, with a dotliftime of 1 second and a dot size of 0.5 pixels viewed by subjects at≈50cm distance resulting in a dot size≈0.02°.

Mice Experiments

All experiments were conducted in accordance with the guideline given by Landesamt für Gesundheit und Soziales Berlin and were approved by this authority.

Surgical Preparation

We used adult (>P40) C57BL/6 wild-type mice, GP4.3 thy1, (#:024275, The Jackson laboratory), and VGAT-CHR2-EYE transgenic mice (#:014548, The Jackson laboratory). Mice were anaesthetized with mit Ketamin/Xylazin (Ketamin 12,5 mg/ml, Xylazin 1 mg/ml (10 µl/g) intraperitoneal) and kept on a thermal blanket. After surgery,mice received Caprofen (5 mg/kg) und Buprenorphin (0,05-0,1 mg/kg) for 2 days. During head-fixation skin and periosteum were removed, a thin layer of light-curing adhesives was applied to the skull. A light-weight head-post was fixed on the skull with a dental cement. For widefield imaging, intact skull cleared using a layer of cyanoacrylate (Zap-A-Gap CA+, Pacer technology) to clear the bone(*50*). Green fluorescent was check under widefield microscope after surgery and double checked later with two-photon imaging calcium imaging. For optogenetics, over intact skull stereotaxic marks were identified using permanent marker, before skull clearing, to label different targets of optogenetic inhibtion: primary visual cortex, V1 (1.0 mm anterior to lambda and 3 mm lateral from the midline), retrosplenial cortex, RSC (1.0 mm anterior to lambda and 0.7 mm lateral from the midline),

posterior parietal cortex, PPC (2.0 mm anterior to lambda and 2 mm lateral from the
midline), Medial Frontal Cortex, MFC(0.8 mm anterior to bregma and 0.7 mm lateral from
the midline), LFC, LFC (2 mm anterior to lambda and 1.8 mm lateral from the midline).
For two photon imaging, a 3-mm craniotomy was made 0.3mm radially anterior from
bregma and sealed with a 3-mm glass coverslip with cyanoacrylate glue and dental
cement.

Behavior
Behavioral Setup
Our behavioural studies employed a modification for the setup developed by Nashaat et
al. 2016, that allows integrating natural behaviors with advanced neuroimaging in a head-
fixed context. The setup is composed of a Plexiglas air-table, perforated with evenly
spaced holes, rerdirecting compressed air to generate an upward current flow, levviating
a PLA platform (15cm diameter) on top. The platform features 3 nose-pokes allowing the
mice to navigate and seek rewards. A head-fixation mechanism ensures the mouse's
head remains centered. The nose pokes, equipped with LEDs, infrared sensors, and a
reward delivery tube, controlled by a solenoid valve, are extension of the Bpod
(Sanworks) behavioral control system. To display RDK stimuli, a Retina LCD screen was
positioned in front of the platform, approximately 10-15 cm away. RDK stimulus was
generated with custom script utilizing either Psychtoolbox or Psychopy(51, 52). White
dots were rendered against a black background. A single dot is set to 2.5° size with a
lifetime of 0.5s and travelling speed of 20°/s-40°/s, with wide-field sessions and 2-Photons
sessions always using 20°/s. Dots were rendered to fill the whole screen with dots filling
20% of viewing apreature. Dots traveled in 8 directions (cardinal- and inter-cardinal
directions) with a ratio of dots, determined by the trial's coherence level, formed a majority
moving either left or right, toward the direction of the reward port.

Behavioral Training
The initial phase introduced freely moving mice to the behavioral system using a simple
light-chasing task conducted over three sessions, each approximately 30 minutes. This
phase allowed the mice to acclimate to the environment and task structure. In this task,
trials began with the mice fixing their nose inside the central port, illuminated first, followed
by a light cue on either the left or right port signaling the correct choice. A successful
selection resulted in a water-based reward, sweetened with 0.05% saccharine and
sucralose, while incorrect choices triggered a high-pitched beep (4000-5000 Hz) as
negative feedback. Subsequently, mice were head-fixed to adapt to moving the platform
themselves, guided by experimenter-assisted platform movements. As mice perform
above 75% accuracy in light-chasing task, a 4-seconds timeout penalty for incorrect
choices is introduced. Trials where mice exited the central port before minimum sampling
resulted in aborted trial with 1-2 second timeout. Expert-level was marked by sustained
>75% accuracy across sessions, after which RDK stimuli were gradually integrated
alongside light cues. Initially paired, the reliance on light cues was reduced until the task
relied solely on RDK stimuli, with varying coherence levels (100%, 50%, 40%, 10%, 0%)
introduced as the training advanced. Motion direction and coherence varied pseudo-
randomly to prevent biases.

Fixed sampling Paradigm

For the fixed time (FT) Paradigm, the approach mirrored tasks previously conducted(*10*,
11, *53*). In FT trials, Mice must remain poked-in for the entire 1s stimulus duration. Early-
Withdrawals counted as aborted trials. For the chronometry experiment (Extended Fig.
1c), trials between 0.3s to 1.2s, in increments of 0.3s, were presented in pseudo-random
order with 20% probability in-between 1s trials.

Free sampling Paradigm
For free sampling paradigm, the approach mirrored tasks previously conducted with
humans and monkeys. In this paradigm, the stimulus is continously presented and only
terminated on exiting the center port, with a maximum of 5s cap. The mice had the
autonomy to end the sampling period anytime between a minimum of 0.3s and a
maximum of 5s, facilitating a balance between impulsivity control and decision-making
flexibility. Trials with less than 0.3s sampling duration are counted as aborted trials.

Optogenetics
The targeting coordinates for optogenetic inactivation were determined using the Allen
Brain Map API for Mouse Brain data(*29*, *35*). 473-nm laser (OEM Laser Systems) was
administered through bifurcated fiber bundle (Thorlabs BFYL2LF01), each emitting light
covering approximately 250 µm in diameter at a 40 Hz with an 80% duty cycle and an
intensity of 6 mW/mm² to ensure that photoinhibition was confined to the targeted cortical
areas(*29*). Additionally, blue masking light was positioned above the viewing screen.
Training data indicated that mice exposed only to the masking light maintained an
average performance exceeding 75%. In each experimental session, mice experienced
perturbation in one brain area per session, with optogenetic trials pseudorandom
introduced among control trials. To sustain high performance levels (70% or higher) and
avoid demotivation, the probability of light illumination was set at 15-20%. During
optogenetic inactivtion under fixed-time paradigm, inactivation duration was either
throughout the entire delivery period of sampling (1s, Full), during early sampling (0-
0.35s, Early), or the late sampling (0.65-1s, Late) phases of stimulus presentation (Figure
2)(*34*).

Widefield imaging
We used an adaptation of widefield imaging setup that utilizes two light source for
hemodynamic responses correction. We used the following variation: for the inverted
tandem lens macroscope, we utilized a top lense with focal length, 105 mm (DC-Nikkor,
Nikon) and a bottom one with focal length 85 mm (85M-S, Rokinon)(*28*, *50*).

Two-Photon Calcium Imaging
Imaging utilized a resonant-scanning two-photon microscope (ThorLabs B-scope) at a 5
kHz scanning rate. GCaMP6s excitation occurred at 940 nm via a Ti: Deepsee laser (90-
110 mw power). Data collection was through a 16x water immersion objective (Nikon, 0.8
NA), capturing 512 x 512 pixel full-frame images of layer II/III neurons within a 1 mm²
area, at 150-300 µm depths. Selection criteria for imaging fields included cell density and
minimal vascular interference. Imaging operated at 30 Hz frame rate, employing
ThorImage software.

Data Analysis

Data analysis was performed using Python utilizing Numpy, Scipy, Pandas Scikit-Learn,
Matplotlib, Seaborn and statsmodels libraries.

Human and Mice Behavioral Analysis
Each session trials were grouped by difficulty. Within each difficulty, the data were
grouped into 3 thirds (1st Quantile: Fast, 2nd Quantile: Typical, 3rd Quantile: Slow). In mice,
early-withdrawal trials, i.e. trials where mouse spent <0.3s sampling (88 trials), and trials
where mice stayed in the center-poke after the end of the maximum 5 seconds stimulus
presentation were excluded (131 trials). Additionally, mice with few trials (<300 trials) or
that followed a non-typical strategy, dissimilar to Fig. S2 E, where correct trials are slower
than incorrect trials, were excluded (21,628 out of 90,973 Trials were excluded). For
human subjects, stimulus was set to be indefinitely presented. Here, trials with more than
5 seconds were excluded (0.2% of maximum correct outcome competition, 7.84% for
accuracy competition).

In analyses where reaction-time Z-score normalization is applied (Fig 1D, G-H, Sup. Fig.
3. A-B,E-G) each mouse's data is pooled across all its sessions. If only a subset of data
is selected, e.g. only correct trials in Fig 1G, then Z-score normalization is first applied
beforehand. For human data, for analyses that characterize each experiment type (Fig.
1F-H. Sup. Fig. S2 G-H,J-K), data was normalized only for the given experiment type. For
analyses characterizing change in behavioral signature across experiment types (Fig. 1C,
Sup. Fig. S2 B-C) data was normalized across both experiment types.

Psychometric curves (Fig. 1 E-F, J Right, Sup. Fig. 1 B Middle, Sup. Fig. 2F,I, Sup. Fig.
4 A, Sup. Fig. 6 B-F, Sup. Fig. 7. C-F, Sup. Fig. 15 E-F) were fit using the Psychofit library
(https://github.com/cortex-lab/psychofit) using error function configuration. Single-sided
psychometric functions, characterizing performance and perception threshold, were fit
using a single-lapse-rate configuration, while two sided psychometric functions,
additionally characterizing choice bias, were fit using two-lapse-rate configuration. As the
region of greatest rate of change within the psychometrics differ between humans and
mice subjects, human data were binned in [0, 1, 5, 10, 20, 50, 100] bin intervals. Mice
data were binned in power of two increments, yet when excluding the empty bins the
remaining yield [0, 16, 32, 64, 100] bin intervals. For Fig 1I, Sup. Fig. 2L and Sup. Fig 3I,
a paired t-test was performed across each subject.

For Sup. Fig. 2B-C, mice were limited to a maximum sampling time of 3 seconds to match
the maximum stimulus time used in human experiments. To compare the 3 groups, and
after testing each group for normality (Shapiro-Wilk test), we deployed one-way ANOVA
followed by Tukey's test with Bonferroni correction for post-hoc pairwise comparisons.

In Sup. Fig. 3F, groups were compared using the Kruskal-Wallis test, with post-hoc
comparisons performed using Dunn's test with Holm–Bonferroni correction. In Sup. Fig.
3G, trials were pooled acris all mice and were classified as Win (previous trial correct) or
Lose (previous trial incorrect). Each classification was split into bins of 800 trials each,
where the stay/switch update value within each bin was calculated relative to the system-
generated baseline stay/switch distribution within the given bin. A Gaussian kernel was
applied to smooth changes between neighboring bins. Two mice were excluded due to

abnormal low z-score distribution (mean percent of trials < -1 z-score=3.28%, excluded
mice=18.62% & 13.26%). The stay/switch update per sub-panel in Sup. Fig. 3H was
calculated similarly to a single bin calculation in Sup. Fig. 3G but without Gaussian kernel
smoothing.

Average Reward-Rate figures (Fig 1H, Fig. S3 B-D) are calculated per subject then
subjects' mean of means is calculated. A subject average reward rate (including Fig 1K,
Fig. S4 Reward Rate column) is calculated as the subject's sessions' mean of means
sampling for each reward values over the last 5 trials. A session's average reward rate
entry with 5 trials or less is excluded. As Avg. reward rate value=0, i.e. no reward over
the last 5 trials, is the least occurring event with few data points, it is excluded from the
plots.

To quantify the contribution of individual predictors to sampling time (Fig. S3 J), we used
ordinary least squares (OLS) regression from the statsmodels library, with the model log-
likelihood serving as the measure of fit. For each animal, we log-transformed the sampling
times and fit both a null model (intercept-only) and a full model including all predictors.
The difference in log-likelihood between the full and null models was used to define the
total log-likelihood improvement — a measure of the overall explanatory power of the
predictors. To assess the unique contribution of each predictor, we fit reduced models
omitting one predictor at a time and computed the percentage loss in log-likelihood
improvement relative to the full model. This procedure was repeated for each subject,
and group-level statistics were computed as the mean and standard error of the mean
(SEM) across animals. (Mention here the parameters used, and they were used to avoid
redundancy information between variables)

Model
To model the effects of incorporating Q-learning and R-learning reinforcement learning
influence on the Drift Diffusion Model (DDM) (Fig. 1J-L, Fig S4), we fitted each mouse's
behavioral data against four models:
DDM
We used the standard DDM as the baseline comparison model:
Let offset starting point:

$$Z = 0 \tag{1}$$

807 Let the initial evidence state be:

$$x_0 = Z \cdot \frac{a}{2} \tag{2}$$

809 The decision variable evolves according to:

$$x_{t+\Delta t} = x_t + k \cdot Cohr \cdot \Delta t + s \cdot \epsilon_t \cdot \Delta t \tag{3}$$

811 A decision is made when x_k reaches either bound ($\pm \frac{a}{2}$). The time to decision is:

$$T_{decision} = i \cdot \Delta t \tag{4}$$

Trial Total decision time is calculated as follows:

$$T_{st} = \begin{cases} T_0 + T_{decision} & \to T_0 + T_{decision} \leq T_{max} \\ T_{max} & \to T_0 + T_{decision} > Tmax \end{cases} \tag{5}$$

Where a is the distance between decision bounds at $\pm\frac{a}{2}$, ($a > 0$), Z is normalized

starting point ($Z \in [-1,1]$). x_t is accumulated evidence at time step t. Δt is a time step

increment ($\Delta t > 0$). $Cohr$ is the dots coherence and direction on the trial ($Cohr \in$

$[-1,1]$). ϵ_t is random standard normal noise at time step t ($\epsilon_t \sim \mathcal{N}(0,1)$). $T_{decision}$ is

time until a boundary is crossed. i is number of time steps until boundary crossing. T_0 is

non-decision time, e.g., motor and sensory latency. T_{max} is maximum simulation time

per trial. T_{st} is trial total time including non-decision time.

Model free parameters are: k: drift scaling factor ($k > 0$), s: noise scaling factor ($s > 0$).

Q-learning + DDM

To capture action preferences, we modeled a simple Q-learning process starting from a

center nose-poke state S_0, with left and right nose-pokes as terminal states each

associated with an action value. We allowed an additional parameter to capture motor

bias:

$$Q_L^{n=1} = Q_R^{n=1} = 0.5 \tag{6}$$

$$Q_{L|R}^n = Q_{L|R}^{n-1} + \begin{cases} \alpha \left(Reward^{n-1} - Q_{L|R}^{n-1}\right) & \to ChoiceDir^{n-1} = (L|R) \\ 0 & \to ChoiceDir^{n-1} \neq (L|R) \end{cases} \tag{7}$$

$$Z^n = clip\left(\delta \cdot \left[log\left(\frac{Q_L^n}{Q_R^n}\right) + offset\right], -1,1\right) \tag{8}$$

Where n is the trial number within a session. $Reward^n$ flags whether a reward is received

on trial n ($Reward^n \in \{0,1\}$). $ChoiceDir^n$ is choice direction on trial n ($ChoiceDir^n \in$

$\{0,1,nil\}$). Q_L^n, Q_R^n are action values for left/right at trial n ($Q_{L|R}^n \in [0,1]$).

Model free parameters are: $offset$: motor bias term ($offset \in [-1,1]$), α: Q-learning

update rate ($\alpha \in [0,1]$), δ: starting offset scaling factor ($\delta \in [0,1]$).

R-Learning + DDM

To model adaptive strategy changes in response to reward rate, we applied R-learning

as modulating of noise gain rather than a change in decision bound. This choice was

guided by the latter findings of the constant motor threshold value of LFC.

$$RR^n = RR^{n-1} + \beta(Reward^{n-1} - R^{n-1}) \tag{9}$$

$$x_{t+1}^n = x_t^n + k \cdot Cohr^n \cdot \Delta t + RR^n \cdot s \cdot \epsilon_t \cdot \Delta t \tag{10}$$

Where RR^n: reward rate at trial n ($RR^n \in [0,1]$).

Model free parameters: β: R-learning update rate ($\beta \in [0,1]$).

Q-Learning + R-Learning + DDM

We combined both Q- and R-learning models to evaluate their joint influence on

behavioral strategies.

In all models, to reduce interaction effects when fitting both the decision boundary and

noise parameters simultaneously, we fixed the decision boundary to $a = 2$, setting

bounds at $(-1, 1)$. Using differential evolution optimization, models' loss functions were
minimized using Chi-Square fitting method to lessen the difference between the subjects'
sampling times and those generated by the model(*54*). Loss was first computed
separately for correct and incorrect choices, and the resulting loss values were then
summed. For each choice condition, the Chi-square statistic was calculated using the
following quantile intervals from the subject's data: $[0, 0.1, 0.3, 0.5, 0.7, 0.9, 1]$. The
integration step was set to $\Delta t = 0.005s$, with a maximum allowable sampling time of
$T_{\max} = 3s$. Trials that did not reach a decision threshold within this time were categorized
as no-choice and no-reward trials.

The coefficient of determination R^2 (Fig 1L) is computed for both the 2-sided
psychometric, thus capturing performance and choice bias, and the average reward rate.
For psychometric R^2, the difference in each difficulty-direction bin between slow and fast
psychometric fits of the subject's real data is compared against the corresponding data in
the model. For reward-rate R^2, sessions' average reward-rate for each reward-rate value
is compared in real-data vs model data. A session's reward-rate value entry with less than
5 trials is dropped.

Optogenetics Data Analysis
Optogenetics data was collected for one brain-region at a time and its control trials span
between the first and last brain-region optogenetics trials. To quantify the optogenetics
performance effect, we computed the drop in performance across each session as
follows:

871
$$Drop = -100 \times \left(\frac{\overline{Opto\ Performance}}{\overline{Control\ Perfomance}} - 1 \right) \tag{11}$$

We adapted the approach in to test whether optogenetics inhibition effect is significant
(Fig. 2D)(*29, 30*), data were pooled across mice, and for each of 10,000 iterations,
subjects, sessions, and trials were resampled with replacement in that sequence. For
each iteration, change in performance was computed. The p-value was determined by
calculating the proportion of iterations where the performance changed sign, e.g. drop in
performance changed to performance improvement, was detected. As a two-tailed test
was used, the significance threshold was set at 0.025 and adjusted for multiple
comparisons using the Holm-Bonferroni correction.
To compare whether the performance effect of optogenetics inhibition significantly differs
between two brain regions (Fig. 3C), we first perform Shapiro-Wilk test for normality on
both groups. If data is not normally distributed for any of the groups (Fig 3C left),
Wilcoxon–Mann–Whitney two-sided test is used, when data is normal for both groups (Fig
3C right), two-sided t-test is used.
To regress the relation between sampling time and performance under control and
optogenetics inhibition conditions (Fig. S15C), we performed linear regression on the z-
score normalization on the sampling time of each animal control session. The
optogenetics trials were z-score transformed using the corresponding control data
distribution. To acquire mean performance, trials were grouped in bins of 50 trials. A linear
fit for each of the control and optogenetics sessions as well as the overall data were
plotted using a fixed common origin value at -1.5 z-score and 50% performance.
892

Wide-Field Imaging Data Analysis
We adapted a variation of the analysis in (https://github.com/jcouto/wfield) with the
following changes: 1) using 800 singular value decomposition (SVD) components instead
of default value (200 components) to reduce dimensionality-reduction loss. 2) Following
image registration step, pixels outside the brain were set to black to minimize their
assigned SVD weights. 3) we used separate left and right hemispheres for a better fit that
accounts for unequal sizes due to zooming artifacts caused by small angle tilt in the
acquisition camera. To characterize the wide-field activity of a typical trial (Fig. 2B, left),
we constructed a reference set by pooling median-time trials across sessions (Fig. S5B).
For each session, we identified trials falling between the 0.45 and 0.55 quantiles of the
sampling duration distribution (Fig. S5B, vertical colored bars). We then computed the
global median of these session-level medians (Fig. S5B, red horizontal line). The
reference set included all trials whose sampling durations were within ±0.1 seconds of
this global median (i.e., vertical colored bars falling between the horizontal dotted lines).

Two-Photon Data Analysis
Preprocessing
Raw two-photon imaging data were preprocessed using Suite2p for image registration
and neurons extraction followed by manual inspection(*55*). To identify active neurons and
neurons active trials, we adapted a variation of the steps used in previous 2 photon
imaging studies(*44*). We extracted raw traces 0.1 seconds before start of sampling until
0.1 seconds after end of sampling. For each trace, a z-score normalization is temporarily
applied across a concatenation of all the sampling trials. Each trial is re-extracted, and
the standard deviation is computed after applying a 1-frame gaussian smoothing. A
neuron is marked to have an active trial if the standard deviation of the trace within the
trial exceeds a threshold value of 3 x 5^{th} percentile of all the trials standard deviation.
Neurons with less than 5 percent active trials are discarded. 837 out of 2,340 neurons in
MFC (36.44% ±6.9%SD per session), and 609 out of 1,0536 neurons were marked as
active in LFC (38.88% ±7.51%SD per session).Neurons $\Delta F/F_0$ was calculated using the
whole session trace after applying a gaussian smoothing with a standard deviation=1. F_0
is computed by computing the mode value of all the frames baseline. A frame's baseline
was computed as the maximum of the minimum values in the surrounding 60 seconds'
window.

Heatmaps
Fast and Slow heatmaps (Fig 3H) were sorted in reference to all trials' heatmap or in
reference to all Slow trials (Fig S11 B). The activity is normalized across the minimum
and maximum values found in both heatmaps.

Tuning selectivity
To test neuronal selectivity for specific criteria, e.g left/right choices (Fig. 4C) and previous
left/right choices (Fig. S12 A, we extracted the maximum trial activity for each condition
and compared them using a two-sided Wilcoxon–Mann–Whitney test.

To quantify the percentage of neurons that are modulated by previous trial reward
outcome (Fig. S12 C Pie-Charts), we computed a modulation index for each neuron,
defined as the difference in activity between matched and mismatched stimulus

conditions, separately for trials following a correct and incorrect outcome. We then
calculated the change in this modulation index based on the previous trial's outcome. To
assess statistical significance, we performed a permutation test of 1,000 shuffles in which
trial labels for previous outcome were shuffled, generating a null distribution of selectivity
differences.

To determine the percentage of neurons sensitive to reward history (Fig. S12B), we
binned each neuron's trials based on reward history, capping the history at a maximum
of three consecutive correct or incorrect outcomes. If the neuron was determined to be
signficantly selectivy tuned to a criteria (e.g. previous-choice left/right), the analysis is
then restricted to trials matching neuron's selectivity (e.g preferred direction), otherwise
all the trials are used. We computed the mean activity for each bin and ordered bins in
descending sequence of incorrect choices followed by ascending correct choices (aligned
with the reward history timeline in Fig. S12C, top). Neurons whose activity increased
monotonically across this sequence were classified as reward-history sensitive. These
results were compared to a null distribution generated by 1,000 shuffles of reward-history
labels across trials.

Sequence characterization
To determine each neuron's rank within a sequence, we used the median peak firing
position across its active trials within time normalized trials.
To assess the consistency of sequential neuronal firing across trials (Fig. 3I–J), we
compared the rank deviations between fast and slow trials. For each session, a neuron's
rank was defined as its median peak firing position in fast and slow trials separately. Tied
ranks were assigned the same value, with subsequent indices skipped (e.g., 1, 2, 2, 2,
5). The rank deviation between fast and slow trials was calculated as:

$$960 \quad \text{deviation} = \frac{\sum |\text{NeuronRank}_{Fast} - \text{NeuronRank}_{slow}|}{Number\ of\ neurons} \quad (12)$$

We then performed a permutation test with 100,000 iterations, shuffling the ranks to
generate a null distribution. The original deviation's p-value was projected onto the
resulting z-score distribution. A variation of this analysis (Fig. S9D–E) was also
conducted, comparing fast and slow trial ranks to those from typical trials (defined as the
middle third quantile). Permutation tests were run by reshuffling the typical trial ranks.
This approach allowed us to assess whether the typical sequence of activation resembled
those observed in both fast and slow trials, using an unbiased, independently derived
reference set for evaluating rank consistency.

To investigate how single neurons modulate their responses with respect to sampling
duration (Fig. 3K, Fig. S9F–H), we analyzed the correlation between each neuron's firing
position and area under the curve (AUC) of calcium activity as a function of increasing
sampling time. AUC was computed from calcium traces while accounting for two key
caveats. First, strong motion artifacts in the two-photon imaging field, particularly along
the Z-axis, can induce sharp frame-to-frame deflections in the calcium trace that are
inconsistent with expected GCaMP6s dynamics. While such artifacts are typically
mitigated in trial-averaged data, single-trial traces remain more sensitive. To reduce these
effects, we applied Gaussian smoothing with a standard deviation of 2. The resulting
traces (Fig. 3K left; Fig. S9F,G left panels) were visually inspected to ensure removal of

high-frequency artifacts. Second, the slow decay kinetics of GCaMP6s fluorescence can
cause residual signal from prior events to appear in subsequent time points. To ensure
AUC reflects ongoing or sustained activity, we restricted calculations to positive or
sustained deflections. A trace may contain zero or more such events. For each, AUC was
defined as the sum of trace values above 60% of the peak amplitude within the event
window (i.e., ≥ 0.6 × [peak – value at deflection onset]). Using neurons' active trials (see
"2-Photon Preprocessing" for the definition), we performed linear regression of both AUC
and peak firing position against sampling time, and computed Pearson correlation
coefficients (Fig. S9F,G middle and right panels). To avoid skewing results, trials with
sampling durations ≥ 3 standard deviations from the mean (Z-score) were excluded.
Neurons with an absolute Pearson correlation ≥ 0.3 were considered non-rigid (Fig. 3K
right; Fig. S9H).

To further assess the significance of these correlations, we plotted histograms of
observed values against distributions obtained via 1,000 reshuffled controls, in which
sampling times were randomly permuted within each neuron (Fig. S9H). Notably, neurons
with early peak firing positions can exhibit gain modulation (i.e., changes in AUC), but
their firing time remains fixed (Fig. S9H top). In contrast, neurons peaking later in the
sampling epoch can exhibit both gain and timing modulation (Fig. S9H middle and
bottom).

In Fig.4E and Fig. S11C, each pair of bins contain the set of significantly direction-
selective neurons, pooled across all brain-region sessions, that their peak firing position
IQR fall within the bin-pair boundary. Thus, each pair of green and brown bins contain the
same set of neurons under different trial conditions: when the trial matches the neuron
selectivity (green) and when trial mismatches it (brown). The bin height and error bar
reflect the average and SEM of the neurons mean z-score activity under the bin condition.
We split time-normalized sampling duration into 7 bins pairs: boundaries pair before and
after sampling start/end and 5 equal-time pairs during sampling. We excluded neurons
that their median peak firing position falls within the first and last bins pairs, to avoid
including predominantly motor tuned neurons.

To further confirm the observed effects, a logistic-regression choice classifier (Fig S11D)
was run on significant choice neurons. After applying z-score normalization to each
significant neuron traces during the sampling epoch, the features input to the classifier is
each significant neuron maximum firing at each trial at either the start of sampling (0.1
before and 0.2 after start of sampling) or end of sampling (0.2 before and 0.1 after end of
sampling). The classifier input labels are the animal direction choice. For each test
combination, a 1,000 of random training and test splitting is performed, with 30% of the
data used for testing on each run.
1016
1017 We defined Coding-efficiency (Fig. 4F), a measure of the mean separation between
1018 match and mismatched selectivity, as follows:

1019
$$Eff = \frac{\overline{\text{Match}} - \overline{\text{Mismatch}}}{\overline{\text{Match}}} \tag{13}$$

1020 Coding-efficiency error bars are computed using the propagation of uncertainty
1021 approximation method:

1022
$$\Delta Eff = \sqrt{\left(\frac{100 \cdot \overline{\text{Match}} - 100 \cdot \left(\overline{\text{Match}} - \overline{\text{Mismatch}}\right)}{\overline{\text{Match}}^2}\, \sigma_{\text{Match}}\right)^2 + \left(\frac{-100}{\overline{\text{Match}}}\, \sigma_{\text{Mismatch}}\right)^2} \quad (14)$$

To describe a two-photon population activity, either the mean activity of all neurons (Fig.
3E solid lines, Fig. 4E background dotted lines, Fig. S8), or the mean activity of only
significant neurons (Fig S10 A left, B left, Fig S11 C bottom), the sum of z-scored activity
is computed across each time point.

To investigate whether early neural activity within a brain region influences fast versus
slow behavioral strategies (Fig. S10A, middle), we quantified the percentage of active
neurons during the early sampling window (from 0.1 seconds before to 0.3 seconds after
sampling start), relative to the total number of traces in the session (see "2-Photon
Preprocessing" for the definition of active trials). The 0.3-second cutoff corresponds to
the minimum enforced sampling duration in the behavioral task. Trials were pooled across
animals and binned into 0.25-second intervals based on their sampling durations. A linear
regression and Pearson's correlation coefficient were then computed to quantify the
relationship between early activity and sampling duration.

To examine how the density, or sparseness, of population activity within a brain region
evolves with sampling duration (Fig. S10B, right), we repeated the early activity analysis
described above (Fig. S10A, middle), but applied it to the final phase of the sampling
period. Specifically, we analyzed the window spanning the last 30% of the sampling
duration up to 0.1 seconds after the end of sampling (i.e., movement start).

To investigate whether early neural activity within a brain region influences performance
(Fig. S10A, right), we repeated the early activity analysis described above (Fig. S10A,
middle) with the following modifications: only easy trials were included to control for task
difficulty, and trials were binned by early activity levels, ranging from 0% to 30% in 3%
intervals. For each bin within a session, performance was calculated as the difference
between the mean performance of trials in the bin and the mean performance of all easy
trials. A linear regression and Pearson's correlation coefficient were then computed to
assess the relationship between early activity and performance.

Supplementary Figures

Fig. S1. A novel floating-platform nose-poke system to study neural mechanisms of decision-making in head-fixed mice. (**A**) 3D illustrations showing side (left), top (middle), and schematic bottom (right) views of the nose-poke floating-platform setup designed for head-fixed mice performing a Random Dots Kinematogram (RDK) task. (**B**) Free-sampling RDK task structure (left), psychometric curves showing choice performance (% choice-left) across individual sessions for an example mouse (middle; thin gray lines: single sessions, bold black line: average), and average sampling time as a function of stimulus coherence levels for rightward (R) and leftward (L) choices (right). (**C**) Left: Schematics illustrating experimental setups comparing freely-moving (nose-poking) and head-fixed (nose-poking floating-platform) paradigms. Right: Kernel density estimates (KDE) of sampling-time distributions comparing an example mouse performing

1061   freely-moving light-chasing (red), head-fixed light-chasing (orange), and head-fixed RDK
1062   (yellow) tasks. Mean reaction times ± SEM indicated.

Fig. S2. Switching between fast and slow behavior is a common strategy during voluntary behavior in both humans and mice. (**A**) Schematic of the human RDK task. Participants performed the task in two contexts: Context I (Accuracy)—emphasizing correct performance, and Context II (Speed)—emphasizing maximizing correct

responses. Subjects responses, reported by pressing left or right arrow keys, were followed by outcome feedback. (**B**) Distribution of normalized sampling times (z-scored) across contexts. Violet: human accuracy context; turquoise: human speed context; orange: head-fixed mice on the floating platform. Dashed lines indicate means for each distribution. (**C**) Sampling times were significantly longer in the human accuracy context compared to both the human speed and mouse conditions. Sampling time distributions in human speed and mouse contexts were not significantly different (ns). (**D**) Sampling time histograms (seconds) across RDK stimulus difficulties (easy, medium, hard) for an example mouse (M#85, 3,519 trials), subdivided into fast, typical, and slow trials. (**E**) Mean sampling time (seconds) for correct (green) and incorrect (red) trials across difficulty levels (hard, medium, easy) for the same example mouse. (**F**) Psychometric performance (choice-left %) as a function of motion coherence for fast (red) and slow (yellow) trials. (**G–H**) Data from human participants in the speed context (n=22). (**G**) Sampling time distributions (z-scored) across difficulties for fast, typical, and slow trials. **H**) Mean sampling time (z-scored) for correct (green) vs. incorrect (red) trials across difficulty levels. (**I**) Psychometric curve showing choice accuracy across motion coherences for fast (red) vs. slow (yellow) trials during human accuracy context (n = 20). (**J–K**) Same as panels G–H, but for human participants in the accuracy context (n = 20). (**J**) Sampling time histograms by difficulty level. (**K**) Sampling time comparison between correct and incorrect trials across difficulty. (**L**) Accuracy during easy trials for fast (red) and slow (yellow) decisions across three groups: human accuracy context, human speed context, and head-fixed mice. Grey lines represent individual subjects or mice; bars indicate group means.

Fig. S3. Trial history and decision outcome and sensory evidence modulate sampling time. (**A**) Mean sampling time (z-scored) across stimulus difficulties during fast (red) and slow (yellow) behavioral strategies humans under the accuracy context. (**B–D**) Sampling time (z-scored) as a function of recent reward rate (averaged over the preceding 5 trials) in: (**B**) Head-fixed mice (orange), (**C**) Human subjects under the speed context (turquoise), and (**D**) Human subjects under the accuracy context (violet). Thin gray lines represent individual animals or participants; bold lines show group means. (**E**) Heatmap showing average sampling time as a function of previous trial outcome (correct/incorrect) and current trial difficulty. Left: example mouse (M#46); Right: mean across all mice. Warmer colors indicate longer sampling times. (**F**) Violin plots illustrating distributions of sampling time (z-scored) across combinations of previous trial outcome (correct/incorrect) and current trial difficulty. Trial history and trial difficulty significantly affect sampling time. (**G**) Probability of win (green) and lose (red) behavioral updates as

a function of sampling time (z-scored). Fast decisions are more likely to be driven by
outcome-insensitive strategies. (**H**) Impact of previous trial outcome (correct/incorrect) on
win /lose behavior update across current trial difficulties (easy, medium, hard).
Percentages indicate update probabilities. (**I**) Choice bias (preference for one direction)
is higher during fast strategies compared to slow strategies across mice. Paired
comparisons across animals reveal significant reduction in bias during slow decisions (p
= 0.016). (**J**) Variance in sampling time explained (OLS model) by different task variables,
including decision evidence (stimulus strength), previous outcome, previous choice, and
stay/switch behavior. (**K**) Linear regression of sampling time (seconds) over trials for an
example mouse (M#85), showing reduced sampling over the course of a session. Each
gray line represents a single session; the bold line shows the fit across sessions. (**L**)
Same as panel K, averaged across all mice.

Fig. S4. Reinforcement learning (RL)-combined drift diffusion models (DDMs) capture distinct behavioral strategies during fast and slow decisions. (**A**) Four computational models were fit to mouse correct and incorrect trial distribution, each panel (left to right) reflecting sampling time distributions, psychometric performance, reward rate sensitivity, and trial-history dependencies: (Top to bottom) 1. Classic Drift Diffusion Model (DDM). 2. Q-DDM: DDM with ΔQ-learning action values incorporated as a starting point bias. 3. R-DDM: DDM with R-learning value (reward rate) modulating noise to affect drift rate. 4. Q&R-DDM: Combined model with ΔQ-learning action values setting the starting point and R-learning value modulating the drift rate. Each model (black) is fit to mouse

data (grey) and the resulting distributions are shown**:** Sampling time distributions (seconds) for correct and incorrect trials, psychometric performance (choice left%) across motion coherence levels during fast (red) and slow (yellow) trials, reward rate as a function of sampling time, and previous trial outcome vs. current behavioral strategy (fast, typical, slow), showing model vs. mouse performance dependence on history. (**B**) Schematic illustrating how RL variables are integrated into a classical DDM framework to explain behavioral strategy. Evidence (coherence %) is accumulated toward a decision threshold (θ). ΔQ-learning action values biases the starting point (z), affecting early decision tendencies. R-learning value modulates noise amplitude, influencing the drift rate (slope), with lower reward rates yielding more stable accumulation in slow trials and more variable trajectories in fast trials.

Fig S5. Widefield calcium imaging reveals cortical dynamics during typical trials
across mice. (A) Schematic of the widefield imaging setup for monitoring neural activity
across the dorsal cortex (left). Top right: clear skull under white light illumination. Bottom
right: same view under blue light for calcium imaging in Thy1-GCaMP-expressing mice.
(B) Overview of selected imaging sessions across mice focused on typical trials with
sampling durations between 1000–1200 ms. Each point indicates a session, colored by
mouse identity; dashed lines indicate inclusion thresholds. **(C)** Pixel-wise heatmaps from
an example mouse show spatiotemporal dynamics of calcium activity during the sampling

period in typical-duration trials, averaged over successive 200-ms windows. **(D)** Trial-averaged calcium dynamics from selected cortical areas in the example mouse, aligned to normalized sampling time. **(E)** Cross mice average of calcium dynamics from multiple cortical regions (n=4 mice, 7 sessions), showing consistent spatiotemporal frontal dynamics frontal during typical decisions. **(F)** Cross mice (n=5 mice, 14 sessions), exhibit similar spatiotemporal frontal dynamics similar (E). However, most cortical areas exhibit a noticeable decay in activity before the end of sampling.

Fig. S6. Optogenetic inactivation of distinct cortical areas during fixed sampling paradigm reveals region-specific contributions to decision-making. (**A**) Schematic showing bilateral optogenetic inactivation during a fixed 1-second sampling window in VGAT-ChR2-YFP mice. Inactivation was delivered at 6 mW and 40 Hz using chronically implanted optical fibers targeting specific cortical areas under a clear skull preparation. (**B–F**) Psychometric performance curves comparing control trials (solid black lines) and optogenetic inactivation trials (dashed lines) across motion coherence levels for each targeted brain region: (**B**) Primary visual cortex (V1, green), (**C**) Retrosplenial cortex

1159   (RSP, orange), (**D**) Posterior parietal cortex (PPC, violet), (**E**) Medial frontal cortex (MFC,
1160   blue), (**F**) Lateral frontal cortex (LFC, magenta). Data show a region-specific effect of
1161   inactivation on perceptual performance, with frontal areas (particularly MFC and LFC)
1162   showing more pronounced impairments of behavior compared to early sensory areas.

Fig. S7. Frontal cortex spatiotemporal dynamics and its causal contributions to decision-making during sampling. (**A**) Frontal cortical calcium dynamics during a fixed 1-second sampling task, showing normalized widefield fluorescence from medial frontal cortex (MFC, blue) and lateral frontal cortex (LFC, magenta) in an example mouse. MFC shows an earlier, more transient peak, while LFC exhibits delayed and sustained activity. (**B**) Same as (A), but during free sampling trials with durations between 1 and 1.2 seconds

for the same mouse. The spatiotemporal activity pattern across MFC and LFC is
preserved and reflects voluntary sampling dynamics. (**C–F**) Psychometric performance
across different difficulties during optogenetic inhibition at early (0–350 ms) and late (650–
1000 ms) phases of the sampling period. (**C, D**) Early inhibition in MFC (C, blue) and LFC
(D, magenta). (**E, F**) Late inhibition in MFC (E) and LFC (F). Solid lines: control trials;
dashed lines: optogenetic inhibition trials. Data reveal timing- and region-specific effects
on perceptual decisions, with MFC inhibition having the most pronounced impact across
all time.

Fig. S8. Consistent spatiotemporal calcium dynamics in layer 2/3 frontal neurons across behavioral strategies. (**A**) Schematic illustrating calculating population-averaged calcium responses from individual neurons. Left: trial-aligned calcium traces from example L2/3 neurons. Right: resulting normalized population activity over time. (**B**) Normalized population calcium responses of L2/3 population from medial frontal cortex (MFC, blue) and lateral frontal cortex (LFC, magenta) during fast (left) and slow (right) trials in an example mouse. While MFC activity peaks early in fast trials, LFC shows delayed and more sustained activity, particularly during slow trials. (**C**) Comparison of frontal cortex calcium dynamics aligned to motor initiation, using both widefield imaging (left) and two-photon imaging (right). In both methods, LFC shows a ramping activity pattern consistent with motor-related tuning, especially in slow and typical trials. MFC activity peaks earlier and declines before movement onset.

Fig. S9. Temporal organization of L2/3 frontal neuron sequences during fast and slow behavioral strategies. (**A**) Top: Single-neuron calcium traces showing fast and slow decision epochs. Bottom left: Decision probability distribution for fast (red) and slow (yellow) trials in an example session. Bottom right: Average z-scored dF/F trace for fast and slow trials aligned to sampling time. (**B**) Cumulative distribution function (CDF) of trial-by-trial firing reliability for L2/3 neurons in MFC and LFC during fast and slow trials. (**C**) Heatmap of calcium activity from LFC neurons in an example session, aligned by average peak response across trials during fast (left) and slow (right) strategies. Bottom: Rank penalty scores across all MFC and LFC sessions showing consistent differences between strategies. (**D**) Schematic of the permutation-based approach for computing rank mismatch penalties across fast, typical, and slow trials. Neuronal sequences from each

strategy are compared to permuted typical-trial sequences as a reference. (**E**) Left:
Histogram showing rank mismatch penalties for typical (orange), fast (red), and slow
(yellow) trials compared to permuted typical-trial sequences from a single example
session. Right: Summary across all sessions showing consistent sequence deviation in
all three strategies (see methods). (**F**) Example L2/3 neuron responses across trials
(color-coded from red (fast) to yellow (slow) by sampling duration), aligned to sampling
onset. Left: single trials dF/F traces. Middle: Correlation between sampling duration and
peak time. Right: Correlation between sampling duration and response amplitude (AUC).
(**G**) Same example neuron as in F, but responses aligned to movement onset. Left: single
trials dF/F traces. Middle: Sampling duration vs. peak time. Right: Sampling duration vs.
AUC. Both timing and amplitude are significantly modulated by sampling duration. (**H**)
Population-level analysis showing distributions of correlation coefficients between
sampling time and neuronal responses: Top: AUC during early sampling (<0.2 seconds
of sampling time). Middle: AUC during late sampling (>0.2 seconds of sampling time).
Bottom: Peak response timing during late sampling. Blue: MFC neurons; magenta: LFC
neurons; grayed blue and magenta are shuffled controls.

Fig. S10. Frontal L2/3 population activity is differentially tuned to sampling time and performance in MFC but not LFC. A) Left: Mean z-scored population activity in MFC (blue) and LFC (magenta) aligned to the onset of sampling, separated by fast (faint lines) and slow (dark lines) behavioral strategies. Middle: Percentage of active L2/3 neurons during the early phase of sampling as a function of sampling time across mice and sessions. Dashed lines show linear fits with Pearson correlation coefficients and p-values**.** Right**:** Correlation between early sampling activity and behavioral performance (mean % correct) during easy trials across mice and sessions (see methods). **B**) Left**:** Mean z-scored population activity in MFC and LFC across mice and sessions aligned to the end of sampling (just before movement initiation), separated by fast and slow trials. Right**:** Percentage of neurons active at the end of sampling as a function of sampling time across mice and sessions (see methods). MFC activity shows a stronger negative correlation with trial duration than LFC, indicating tighter tuning to behavioral timing in MFC.

Fig. S11. Choice-selective neural population dynamics during fast and slow
decisions. (**A**) Heatmaps of normalized calcium activity from MFC neurons in an example
session for rightward choices (left panel) and leftward choices (right panel), illustrating
population sequence alignment is consistent for left and right choices. (**B**) Proportion of
choice-selective neurons over normalized sampling time in MFC (blue) and LFC
(magenta). LFC neurons show delayed encoding compared to MFC, which encode choice

homogeneously across sampling time. (**C**) Percent of choice encoding neurons at the
start of sampling during fast and slow trials (see methods). (**D**) Top: Population activity of
choice-preferring (green) and non-preferring (brown) neural subgroups at the start
(dashed box) and end (dashed box) of sampling. Bottom: Subpopulation mean activity for
matched and unmatched subgroups in MFC (top) and LFC (bottom), showing divergence
in dynamics between fast and slow strategies at the start and end of sampling. (**E**)
Difference in mean population activity (z-scored) between the last neural bin before the
decision and the first neural bin at the onset of sampling. Values are shown separately
for MFC and LFC, and across matched vs. non-matched choice-neurons during fast and
slow trials. This metric captures the net buildup or declines in activity from initial sensory
sampling to the moment of choice commitment. (**F)** Choice decoder accuracy (coding
efficiency) computed at the start and end of sampling, separately for fast and slow trials.
Both MFC (blue) and LFC (magenta) show improved decoding performance toward the
end of the sampling time.

Fig. S12. Previous trial outcomes and reward history modulate frontal cortex encoding of current and past choices. (**A**) **Left:** Proportion of outcome-sensitive neurons in MFC and LFC. Right: The influence of previous outcome on neurons selective

for previous choice direction (left) and current choice direction (right) at the start of
sampling. (**B**) Left: Example neuron activity is modulated by recent reward history (reward
rate over the last 3 trials), aligned to normalized sampling time. Right: Pie charts showing
the proportion of reward-rate-sensitive neurons in MFC and LFC. (**C**) Top: Example
neurons encoding the current choice, showing modulation by recent reward rate. Bottom:
Example neurons encoding the previous choice, also modulated by recent reward rate.
Right: Pie charts showing the proportion of current choice (green) and previous choice
(yellow) encoding neurons modulated by previous outcome. (**D**) Schematic illustrating
how outcome and choice history jointly influence decision strategy. (**E**) Schematic
representation of how sampling time variability reflects history-dependent strategy
shifting.

Fig. S13. Neural representation of trial history and evidence during fast and slow behavior. (A) Population activity (z-scored) in MFC (blue) and LFC (magenta) during fast (impulsive, light lines) and slow (deliberate, dark lines) trials. Left: Aligned to the start of sampling, showing stronger early activation in MFC during slow trials. Right: Aligned to motor initiation (end of sampling), showing greater LFC activation prior to movement. Center: schematic of the two-photon imaging setup used for single-neuron resolution. **(B)** Neural tuning at the start of sampling for two decision-related variables: Left column: Previous trial's choice direction (left vs. right). Right column: Current trial's stimulus strength (easy vs. difficult motion coherence). Top: Example neuron responses for fast and slow strategies. Bottom: Percentage of significantly tuned neurons in MFC and LFC during fast and slow (see methods). **(C)** Same analysis as panel B, but for the end of sampling, aligned to movement initiation.

Fig. S14. Choice coding efficiency reflects sensitivity to sensory evidence across behavioral strategies. (**A**) **Top:** Sequence-aligned choice-selective neuron responses in MFC (left) and LFC (right) during slow trials, plotted as mean population activity for matched-choice (green) vs mismatch (brown) conditions, aligned to different temporal bins along sampling time. Bottom: Breakdown of neural activity by stimulus difficulty (easy, medium, difficult) for both matched and mismatch trials at the start and end of sampling. During slow decisions, matched-choice neurons show stronger evidence-related tuning, particularly in MFC. (**B**) Same analysis as in (**A**), but for fast trials. Matched-choice neuron activity is more similar across difficulties..

Fig. S15. Mid-sampling inhibition of MFC and LFC disrupts decision probability, accuracy during free sampling paradigm. (**A**) Decision probability during the optogenetic inhibition window (0.3–0.9 s after sampling onset) significantly decreased with **MFC** (blue) and **LFC** (magenta) inhibition. (**B**) Task performance during optogenetic inhibition showed a significant drop under both MFC and LFC inhibition compared to control. (**C**) Linear fit of task performance as a function of z-scored sampling time, separated by condition: control (black), MFC inhibition (blue), and LFC inhibition (magenta). MFC inhibition flattens the performance-sampling time relationship. (**D**) Cumulative trial distribution across z-scored sampling time. MFC inhibition delays the accumulation of decisions compared to control and LFC inhibition. (**E–F**) Psychometric curves show behavioral performance across RDK motion coherence in control slow

1296   (yellow), control fast (red), and MFC-inhibited (E, blue) or LFC-inhibited (F, magenta)

1297   trials.

1298

Video S1. Head-fixed mice navigate a nose-poking floating platform. Side view of a head-fixed mouse controlling a circular floating platform equipped with three nose-pokes. Each trial begins when the mouse initiates a nose-poke in the center port, triggering the presentation of a random-dot kinematogram (RDK) stimulus. The mouse then voluntarily withdraws from the center port to terminate stimulus sampling and rotates the platform to reach either the left or right choice port to report its decision and collect a reward.

Video S2. Head-fixed mice display both fast and slow sampling strategies under identical sensory conditions. Top-view of a head-fixed mouse performing on a nose-poking platform during a decision-making task. The video shows two representative trials at intermediate stimulus difficulty (50% coherence), illustrating that mice can adopt either fast or slow sampling strategies for the same sensory evidence.

Video S3. Head-fixed mice exhibit variable sampling durations while maintaining robust performance during RDK task in nose-poking floating platform. Top left: Psychometric function from an example behavioral session, showing performance across stimulus difficulties. Top right: Histogram of sampling time distribution from the same session, illustrating within-session variability in sampling behavior. Bottom: Top-down video of a head-fixed mouse engaged in the decision-making task on the nose-poking platform.