**Gut microbiota–mediated lipid accumulation as a driver of**

**evolutionary adaptation to blue light toxicity in *Drosophila***

Yuta Takada[1], Toshiharu Ichinose[2,3], Naoyuki Fuse[4], Kokoro Saito[3], Wakako Ikeda-Ohtsubo[1],

Hiromu Tanimoto[3], Masatoshi Hori[1]

[1]Graduate School of Agricultural Sciences, Tohoku University, Japan

[2]Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Japan

[3]Graduate School of Life Sciences, Tohoku University, Japan

[4]Graduate School of Pharmaceutical Sciences, Tohoku University, Japan

Correspondence to Yuta Takada or Masatoshi Hori

yutatakada38@gmail.com

masatoshi.hori.a3@tohoku.ac.jp

## Abstract

Human agriculture has always raced against insect adaptation, requiring updated pest control methods and knowledge of evolutionary processes. Excessive exposure to blue light (BL) kills a wide range of insect species and has attracted attention as an alternative to chemical pesticides. In this study, to understand the evolutionary adaptation of insects under the pressure of BL toxicity, we revealed evolutionary responses to BL toxicity over 70 generations using laboratory selection experiments in *Drosophila melanogaster*. The selected line exhibited an obese phenotype accompanied by midgut elongation, with BL tolerance dependent on gut microbiota–mediated lipid accumulation. Whole-genome and transcriptome analyses consistently highlighted interactions between the microbiota and lipogenesis genes. Remarkably, manipulation of lipogenesis via these genes conferred BL tolerance even in the absence of selection. We suggest that the acquisition of BL tolerance occurs through 'adaptive obesity'. Our study introduces a mechanism of evolutionary adaptation of insects against BL-based selective pressure by maximising the benefits from the gut microbiota via midgut elongation.

## Introduction

One of the most challenging issues in pest control is the rapid evolution of insects. Traditional chemical pesticides have led to the development of resistance, environmental contamination and unintended effects on non-target organisms, highlighting the need for alternative approaches[1,2].

Blue light (BL) is within the 400–500 nm range, is abundant in sunlight, and is crucial for insect physiology and ecology because it influences processes such as visual perception, circadian rhythms, and photoreactivation (Fig. S1)[3–6]. However, excessive exposure to BL is lethal in a variety of insect species[7]. We propose the use of BL as a pest control method and have demonstrated that exposure to BL at intensities below 40% of the BL content of sunlight for several days can be lethal to fruit fly *Drosophila melanogaster* (hereinafter referred to as *Drosophila*), mosquito *Culex pipiens* form *molestus,* flour beetle *Tribolium confusum,* and leaf beetle *Galerucella grisescens*[7–9]. Notably, in *Drosophila*, *C. pipiens* f. *molestus*, and *G. grisescens*, BL around 420, 440, and 465 nm has been found to be more lethal than shorter wavelength 375–405 nm[7–9]. In *Drosophila*, BL around 465 nm is particularly toxic to both pupae and adults, with males being more sensitive than females[9]. These results suggest that the sensitivity of BL toxicity cannot be explained by photon energy alone. The BL toxicity is thought to be associated with reactive oxygen species (ROS) generated by mitochondria and other cellular components, leading to oxidative damage[9–14]. This damage can impair fitness and act as a selective pressure in insects, yet tolerance and evolutionary adaptation to this stressor remain largely unexplored. Achieving the societal implementation of chemical-free, BL-based pest control requires the gathering of evolutionary insights.

Laboratory selection has contributed to understanding the evolutionary adaptations of organisms to environmental stressors[15,16]. *Drosophila* is well-suited for laboratory selection experiments due to its short life cycle and the extensive genetic and physiological knowledge available[17]. Importantly, these experiments often reveal both evolutionary responses and trade-offs that were not anticipated at the outset[18]. Observing evolutionary processes in the

laboratory and identifying the phenotypic, genomic and epigenomic factors that lead to tolerance in *Drosophila* are crucial to understanding the mechanisms of evolution in insects. Here, by selecting flies that are tolerant to excessive BL exposure, we established *Drosophila* strain with enhanced BL tolerance. Characterisation of these selected line (SL) flies revealed that obesity is a key adaptive trait for enhanced BL tolerance. Moreover, we found that parental BL exposure and selection under BL toxicity induce midgut elongation and that increased gut bacterial load is critical for acquired BL tolerance. This study highlights that maximising the benefits of the gut microbiota has been key to evolutionary adaptation in the host.

## Results

## *Drosophila* evolved BL tolerance through laboratory selection

Reproductive capacity is one of the most important determinants of fitness. We investigated the effects on ovarian development to determine whether damage caused by BL toxicity acts as a selective pressure. Older flies are known to be more sensitive to BL toxicity[13], but our results showed that even very young flies, such as those on the first day after eclosion with underdeveloped ovaries, were particularly susceptible (Fig. S2a and Data S1)[19]. BL toxicity consistently suppressed ovarian development at all stages between days 1 and 10 post eclosion (Fig. S2a). Even after a recovery period of up to 7 days post-BL exposure, ovarian recovery remained minimal (Fig. S2b and Data S1). These results highlight that BL toxicity has a dramatic effect on fitness.

First, we generated a *Drosophila* strain with BL tolerance through laboratory selection. In brief, we derived two experimental strains from the same population: one maintained without selective pressure and the other reared under BL toxicity-driven selection (Fig. 1 a). This procedure was replicated in three independent lines for each strain and each line was repeated for over 70 generations (Fig. 1 a). For laboratory selection, a Canton Special (Canton-S) strain, which is a long-inbred strain of the common wild type, was used as the

ancestral population (for details, see "Laboratory selection for BL tolerance -Artificial

selection-" in Methods)[20]. Over the generations, 1–5 day-old flies were exposed for 3 days

to BL at around 465 nm, a wavelength highly toxic to *Drosophila* adults, at a photon flux

density at which approximately 50% of individuals exhibited the "low impact" phenotype,

which was defined as being capable of climbing after a 2-day recovery (Fig. 2b and Fig. S1

and Fig. S3a)[9]. We then selected the low impact flies and let them breed to develop the SL

lines (Fig. 1b). The unselected control line (UCL), maintained without BL irradiation, served

as a negative control for comparative analyses with the SL. In the negative control group, 3

days of BL irradiation were replaced by darkness and UCL parents were randomly chosen

(Fig. 1b and Fig. S3a). Although not selected, a subset of UCL flies in each generation was

exposed to BL under the same conditions as the SL to serve as a positive control group (Fig.

S3a). Across generations, no strain-specific differences in climbing ability were observed

without BL irradiation (Fig. S3b and Data S1). In SL, both male and female flies showed

greater climbing ability after BL irradiation following a single selection event (Fig S3a). In

contrast, fecundity in SL remained suppressed by BL toxicity even after more than 10

generations of selection (Fig. 2a). After around 20 generations, SL flies began to show

improved adaptability, with fecundity maintained at similar levels to flies in the non-irradiated

UCL (Fig. 2a–c, Data S2 and Data S3). Ovarian development was affected little by BL

exposure in the SL flies but severely in the UCL flies (Fig. 2b and Data S3). Moreover,

fecundity in SL flies was slightly increased in response to BL irradiation, whereas that in UCL

flies was reduced (Fig. 2c and Data S3). Additionally, SL flies survived longer than UCL flies

under continuous BL irradiation (Fig. 2d and Data S3). After 73 generations of selection, the

lifespan was further extended, with disparity from the UCL increasing as light intensity

decreased (Fig. 2d, e, Fig. S4 and Data S3). These results collectively suggest that we have

successfully established a new *Drosophila* strain with enhanced BL tolerance.

**SL flies exhibited obese phenotype, lipid accumulation, and**

**oxidative stress resistance**

Next, we examined the phenotypic characteristics of the SL flies in non-irradiated conditions.

We noticed that they exhibited an obese phenotype characterised by abdomen bloating,

increased body weight, and well-developed fat bodies with markedly increased lipid droplets

(Fig. 3a, b and Fig. 4a–b' and Fig. S5a and Data S4). They also showed an elongated midgut

and shortened wings (Fig. 4c and Fig. S5b and Data S4). Additionally, we observed a slightly

low number of oocytes in the SL flies (Fig. S5c and Data S4) and a strong inter-individual

correlation between oocyte number and body weight in the UCL flies (Fig. S5d and Data S4).

SL flies maintained higher body weight than UCL flies despite similar food consumption levels

(Fig. 4d and Data S4). Taken together, these results show that the increased body weight of

SL flies was not due to larger ovaries or higher food intake but may instead be associated

with increased lipid reserves and elongated midguts. SL flies were rich in major lipids,

including fatty acids (fatty acid methyl esters, FAMEs) and triacylglycerol (TAG), and were

starvation resistant, providing evidence of lipid accumulation (Fig. 4e–g and Data S4)[21,22].

Consistent with previous reports that lipid droplets and their constituent TAG may protect

against oxidative damage[23–25], resistance to oxidative stress was higher in SL flies than in

UCL flies (Fig. 4h and Data S4). We did not find any physical traits associated with light

reflection by cellular melanin, despite previous reports suggesting these as adaptive traits to

phototoxicity (Fig. S6)[26]. These results suggest that BL tolerance in SL flies is acquired

physiologically through obesity, rather than through physical traits.

**Increased bacterial abundance in SL flies**

Gut microbiota–derived acetate serves as a potential substrate for de novo lipogenesis in the

host via acetyl-CoA[27,28]. Considering the elongated midgut and accumulated lipids in SL flies,

we compared the gut microbiota profiles and amounts in SL and UCL flies. In both strains,

16S rRNA gene sequencing of whole-body DNA revealed that a single species of

*Acetobacter* (*A. persici)*, a common gut bacteria in *Drosophila*, dominated (>99%) the

microbiota (Fig. S7a and Data S5)[29–31]. *Wolbachia* was not detected in our microbiota

analyses (Data S5). *A. persici* has also been detected in wild flies, and several studies have

reported that in some cases more than 90% of the microbiota consisted of this or other

*Acetobacter* species[30–34]. Relatedly, several studies have reported that *Acetobacter* can

dominate the gut microbiota in natural environments[35,36]. Notably, bacterial counts were

higher in SL flies than in UCL flies, and a mild positive relationship between body weight and

bacterial abundance was observed, with no change in microbiota profiles (Fig. 5a–a' and Fig.

S7b, b' and Fig. S8 and Data S5 and Data S6). All randomly sequenced colonies were

identified as *A. persici* (Data S6). These results consistently show that while both strains

colonise the same species, SL has the higher bacterial abundance.

## Gut microbiota promotes BL tolerance and lipid accumulation in SL flies

Acetic acid bacteria, including *Acetobacter*, are known to influence fecundity and TAG levels

in *Drosophila*[37–45]. In addition, both bacterial abundance and body weight were drastically

reduced in SL flies following BL irradiation, suggesting that the sensitivity of these flies to BL

toxicity may be related to the gut microbiota (Fig. S8). To test the hypothesis that gut

microbiota play a role in shaping the phenotype of SL flies, we used an antibiotic mixture

(Abx) that depletes the bacterial abundance (Fig. S9a). Interestingly, we found that BL

tolerance in the SL flies, as measured by fecundity, was totally abolished by Abx treatment,

suggesting the critical role of gut microbiota (Fig. 5b and Data S6). Feeding the Abx-treated

SL flies with the gut contents of SL flies rescued these phenotypes, whereas feeding with

those of UCL flies barely did (Fig. S9b–d, Fig. 5b, Data S6 and Data S7). Consistently, Abx

treatment significantly reduced the body weight and the TAG level selectively in the SL flies

(Fig. 5c–e and Data S6). This result highlights the dependence of SL flies' body weight and

lipid accumulation on their gut microbiota.

The midgut of Abx-treated SL flies was longer than that of UCL flies regardless of treatment,

and it was even longer in non-treated SL flies (Fig. 5f and Data S7). A positive correlation

between body weight and midgut length in non-treated SL flies (Fig. S10 and Data S7)

suggested that gut microbiota may contribute to midgut elongation. Furthermore, enhanced

oxidative stress resistance was observed only in non-treated SL flies (Fig. S11a, b and Data

S7). These observations align with previous reports that gut microbiota is linked to gut

morphology and oxidative stress resistance[29,46,47]. Overall, these results suggest that the

characteristic phenotype in SL flies is shaped by their gut microbiota.

## Parental exposure to BL affects gut microbiota–associated traits in progeny

We investigated why only the gut contents from SL flies could rescue BL tolerance in infection

experiments, despite the dominance of the same gut bacterial species across strains (Fig.

S7a and Fig. 5b). We noticed that the flies used for gut contents had different parental

experiences depending on whether they belonged to the UCL or SL. Additionally, parental

exposure to BL has been reported to alter the transcriptome of the progeny[48]. Thus, we

hypothesised that parental exposure to BL irradiation might influence gut microbiota–

associated traits in the progeny, leading to quantitative or qualitative changes in gut

microbiota. We exposed adult males and females from both strains to the same BL toxicity

used in the selection experiment for 3 days, followed by a 3-day recovery period. We then

randomly picked flies to be the parent, producing progeny that were influenced by the

parental experience (Fig. 6a). First, we examined the body weight and BL tolerance in Abx-

treated SL flies that were fed with gut contents from flies whose parents had or had not

received BL irradiation. Abx-treated SL flies fed with gut contents from SL flies whose parents

had experienced BL irradiation showed the greatest increase in body weight (Fig. S12a–b

and Data S8). However, flies fed with gut contents from *A. persici* in different cocktails, each

standardized in concentration, exhibited similar body weight increases (Fig. S12c and Data S8). This suggests that quantitative rather than qualitative changes in gut microbiota due to parental BL irradiation are responsible for the increases in host body weight. Furthermore, gut contents from UCL flies whose parents had experienced BL irradiation also rescued the phenotypes of Abx-treated SL flies, restoring fecundity to the level conferred by gut contents from SL flies (Fig. S13a–b and Data S8). These results indicate that parental BL irradiation affects the progeny's bacterial abundance. Consistent with this idea, parental BL irradiation increased the progeny's body weight, bacterial abundance, and midgut length in both UCL and SL flies (Fig. S14, Fig. 6b, c and Data S8). Notably, even without parental BL irradiation, the midgut of SL flies was longer than that of non-irradiated UCL flies, suggesting that this inherited trait had been conserved during the laboratory selection of SL flies (Fig. 6c). In both strains, parental BL irradiation induced midgut elongation even after antibiotic treatment, suggesting that in *Drosophila* this is a common response dependent on parental experience. The increased abundance of beneficial bacteria under BL toxicity may improve fitness, highlighting the dependence of BL tolerance in SL flies on gut microbiota.

## Host genome and transcriptomic profiles explain the characteristic phenotype of SL flies

To investigate how the SL flies develop their phenotypes through interactions with the gut microbiota, we first performed whole-genome resequencing aimed at detecting genetic characteristics unique to SL. The first approach was to explore genes that show strain-specific variation to understand the genetic variation of SL. We identified 1,917 genes with SL-specific variation above 5% for the full sequence length of each gene (Fig. 7a and Data S9 and Data S10). Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis of these variations revealed their association with detoxification and antioxidant response such as *Glutathione S transferase* (e.g. *GstD4*, *GstD5, GstD6, GstD7, GstD10, GstE1, GstE2,*

*GstE5, GstE6, GstE8, GstE11, GstE14, GstO2, GstT3*) (Fig. 7b and Data S10)[49]. Given the

role of oxidative stress in BL toxicity[9,11,13], the variation in these detoxification enzymes

suggests functional changes and the effect of selective pressures. The original of Canton-S

strain is isogenic in the 1920-1930s, although it has already accumulated variants due to long

laboratory breeding (note that the sequencing data is from the Canton-S strain obtained from

the Bloomington *Drosophila* Stock Center (Fig. S15a))[20,50]. Even in UCL, the detection of

694,867 total sites with heterozygous and homozygous variants across the three replicate

lines supports the possibility that multiple standing variants were present (Fig S15b–d).  We

analysed strain-specific homozygous variants to further elucidate the genetic basis of SL.

Homozygous variants shared among the three replicate lines in SL and absent in UCL and

in the Canton-S strain were identified (Fig. S16). SL-specific SNPs and indels were detected

in 658 and 2,625 genes, respectively, giving a total of 2,970 genes, and the majority of

alterations were located outside the coding sequences (CDS) (Fig. 7c and Data S11). KEGG

enrichment analysis identified the Notch signaling and the Hippo signaling pathway as

enriched pathways (Fig. 7d), consistent with their known function in organ size regulation[51–

53]. In particular, the presence of CDS variants in genes that regulate or interact with the Hippo

signaling pathway (*Par-1* (missense mutation), *misshapen* (frameshift mutation)), together

with the elongated midgut observed in Abx-treated SL flies, suggests that midgut elongation

was potentially influenced by these variants (Data S11, Fig. 5f and Fig. 6c)[52,54,55].

Furthermore, the Toll and Imd pathways of the immune system were identified as enriched

pathways of 4,612 genes with SL genetic characteristics (Fig. 7e and Data S12)[56,57]. The

genetic features of SL suggest that the acquisition of gut microbiota-dependent phenotypes

may have been facilitated by midgut elongation and immune system modifications.

Because variations located outside the CDS are generally thought to influence gene

expression and because gut microbiota may alter host phenotypes through changes in the

host transcriptome[37,58,59], we performed transcriptome analysis on the whole body of non-

irradiated UCL and SL flies, with or without the Abx treatment. We identified 1,242

differentially expressed genes (DEGs), of which 1,192 (96%) were specific to non-treated SL

flies (Fig. S17a, b, Fig. 8a and Data S13). KEGG enrichment analysis of the 1192 DEGs

revealed gene enrichment in oxidative phosphorylation, metabolic pathways and the

tricarboxylic acid (TCA) cycle (Fig. 8b and Data S13). Suppressed mitochondrial metabolism

is known to lead to lipid accumulation in various animal species[60,61]. Consistent with this fact,

non-treated SL flies showed downregulation of most of the genes involved in oxidative

phosphorylation and the TCA cycle, which are associated with mitochondrial metabolism (Fig.

8c). In addition, several genes whose downregulation is known to induce lipid accumulation

were also specifically downregulated in non-treated SL flies (Fig. 8c)[21,22,62], including

*Tachykinin* (*Tk*), which also harbored mutation in the SL flies (Fig. 8c and Data S13). Actually,

*Tk* expression is known to be modulated in response to acetate produced by gut microbiota,

thereby regulating lipid utilisation in enterocytes[37,63]. As expected, approximately 30% of

DEGs (387 genes) reflected the distinctive genomic profile of SL, with the Toll and Imd

signaling pathways appearing most enriched (Fig. S18a–d). Of these, 276 genes contained

homozygous variants, including 6 genes associated with the Toll and Imd signaling pathways,

such as *Toll* (Fig. S18b and Data S14).  Our results suggest that the phenotypes of SL flies

are influenced by the Hippo signaling pathway, *Tk*, and mitochondrial genes, and by their

interactions with gut microbiota.

## Lipid accumulation via genetic manipulation confers BL

## tolerance

267  Our finding suggest that SL has acquired a gut microbiota-driven lipid accumulation (Fig.4,

268  Fig. 5 and Fig. 8c). To prove the causal relationship between lipid accumulation and BL

269  tolerance, we examined BL tolerance in genetically manipulated flies that are reported to

270  have increased TAG levels[21,22]. We used a range of mutants and GAL4/UAS-mediated

271  manipulations, including knockdowns of *Tk*, *AkhR*, and *Lipid storage droplet-1 (Lsd-1)*, which

were downregulated in the SL flies (Fig. 8c, Data S13 and Data S15). Interestingly, knocking

down these genes enhanced BL tolerance to a level comparable to that in the SL flies (Fig.

9a). Furthermore, BL tolerance was also enhanced by other genetic manipulations that are

known to increase lipid levels but were not detected as DEGs in our transcriptome analysis

(Fig. 8c), such as knockdown of *brummer* (*bmm*), *Phosphatase and tensin homolog* (*Pten*),

and *Stromal interaction molecule* (*Stim*), and overexpression of *Insulin-like receptor* (*InR*),

*Lk6 kinase* (*Lk6*), and *Lipid storage droplet-2* (*Lsd-2*) (Fig. 9a). Indeed, we found a significant

positive relationship between fecundity and TAG levels among flies with all of these genetic

modifications (Fig. 9b), which strongly suggests that lipid accumulation is a trait that confers

BL tolerance (Fig. 9c).

## The transcriptome of *A. persici* in the host and the growth patterns vary between the host strains

Finally, we tried to investigate the possible mechanisms underlying the contribution of *Acetobacter*. To our knowledge, seventeen *Acetobacter* species have been identified as gut bacteria in *Drosophila* and are known to affect host metabolism and fitness[31,33,35,38,41–46,64,65]. Phylogenetic analysis based on reported full-length 16S rRNA gene sequences suggesting that *A. persici* is closely related to *A. malorum* and *A. cerevisiae* (Fig. 10a). In both UCL and SL, the dominant OTU ID (397 bp) representing 89-98% of the microbiota was identified as *A. persici* Dm-48 strain with 99.75% sequence identity (Fig. 10a, Fig. S19). Analysis of the complete genome of *A. persici* using publicly available data showed that its chromosome is 3,230,507 bp, while its plasmid is 526,169 bp, containing 2,898 and 492 CDS, respectively (Fig. 10b). To characterise the transcriptomic profile of *A. persici* within the host, we first extracted reads that were not mapped to the host genome in the RNA-seq data and subsequently mapped them to the complete genome of *A. persici*. As this mRNA was obtained by poly(A) selection of the host, there is a possibility of bias in the results. However, the sequencing depth was approximately 10X, ensuring a minimum number of reads for

comparative analysis (Fig. 10c, Fig. S20a and Data S16). This approach primarily captured

extreme variation and may have missed low-expressed genes. Nevertheless, a total of 225

DEGs were detected, including 124 downregulated and 101 upregulated genes (Fig. S20b

and Fig. 10d). Gene Ontology (GO biological process) enrichment analysis revealed a

frequent enrichment of metabolic pathways, suggesting that the colonisation state of the

*A.persici* within the SL differs from that in the UCL (Fig. S20c and Data S16). *A. persici*

harbours 15 genes involved in acetic acid production[66]. Interestingly, key genes such as

*Alcohol dehydrogenase* (*ADH*) and *Acetyl-CoA hydrolase* (EC 3.1.2.1) were found to be

specifically upregulated in the SL (Fig. 10d and Data S16). These upregulated genes may

contribute to the microbiota-dependent lipid accumulation of SL. The top 5 genes with the

lowest *p*-values were all upregulated and included potentially oxidative stress-related genes

such as *NADPH-dependent FMN reductase* and *Paraquat-inducible protein* (Fig.10d). The

ability of *A. tropicalis*, a close relative of *A. persici*, to restore host survival via microbiota-

mediated detoxification was previously documented (Fig. 10a)[64]. Given that antioxidant

mechanisms are thought to contribute to resilience to BL toxicity, it is noteworthy that genes

associated with the oxidation-reduction process were detected as enriched DEGs (Fig. 10f

and Data S16). The upregulation of oxidoreductases and translocases indicates their

potential antioxidant role by neutralising ROS and transporting key metabolites that support

cellular redox balance (Fig. 10f). These results suggest that *A. persici* in SL has a different

transcriptomic profile, especially in metabolism. However, it remains uncertain whether this

difference is due to bacterial genomic variation, differences in bacterial colonisation state or

host-mediated effects. We then investigated whether *A. persici* showed phenotypic

differences by analysing the growth patterns of isolates from each host strain. Fifty colonies

were cultured from each host strain. Consistent with the transcriptomic variation, *A. persici*

isolates from the UCL and SL strains showed divergent growth patterns (Fig. 10g–g''' and

Data S16). In the SL, 26% (13 colonies) failed to grow, whereas only 8% (4 colonies) failed

to grow in the UCL. Additionally, UCL showed a higher variance, whereas SL showed a lower

variance (Fig. 10g''). This phenotypic difference indicated a potential bias towards SL-

dominant bacterial strains. Overall, these results suggest that BL toxicity may have acted as

a selective pressure shaping bacterial strain dynamics or driving host-microbiota co-evolution.

## Discussion

In this study, we established BL-tolerant *Drosophila* strains through laboratory selection,

demonstrating that excessive BL exposure drives the 'adaptive obesity', particularly lipid

accumulation mediated by gut microbiota. This supports the increasingly recognized role of

microbiota in host adaptation to environmental stress, in which specific bacterial populations

contribute to adaptive host phenotypes[33,43,67–69]. Our findings further reveal that elongation

of the host midgut in response to parental BL toxicity facilitates an increase in beneficial

bacteria, ultimately enhancing host fitness via lipid accumulation under excessive BL

exposure (Fig. S21).

**Lipid storage is one of the factors that determines BL tolerance**

SL flies exhibited increased lipid accumulation (Fig. 4), and genetic manipulation that

increased lipid contents conferred BL tolerance (Fig. 9). This acquisition of BL tolerance can

be explained by the role of lipid accumulation in protecting against ROS, which are produced

upon BL irradiation[12,13]. Intracellular lipid droplets play a critical role in shielding cells from

oxidative stress, as reduced TAG levels are associated with increased sensitivity to such

stress[23–25,70]. Indeed, the accumulation of lipid droplets in response to intracellular ROS has

been observed in many species[25,70]. Additionally, lipid droplets are closely linked to fecundity,

as higher TAG levels are associated with improved fecundity, whereas reduced TAG levels

can lead to oocyte degeneration[71,72]. Thus, lipid accumulation not only helps defend against

BL toxicity but also supports the maintenance of fecundity. Furthermore, the transcriptome of

SL flies revealed a broad downregulation of genes for mitochondrial metabolism (Fig. 8c).

Mitochondria, while crucial for energy production, also contribute to ROS generation[73]. It has

been reported that BL toxicity causes a decrease in ATP production[12,13]. Therefore, inhibition

of both mitochondrial metabolism and lipid utilisation may be an adaptive strategy to

accumulate lipids while minimising oxidative stress, and obesity may serve to compensate

for insufficient ATP levels.

**358 The potential of the gut microbiota for benefit enhancement in *Drosophila***

*Acetobacter* is generally the dominant bacteria in *Drosophila*[34–36,41,42]. The bacterial

composition and relative abundance within the microbiota varied between laboratories, with

*Acetobacter* sometimes accounting for over 90% of the community[30,32–34]. Generally, the

microbiota of laboratory flies is influenced by factors such as the host genome and

environmental conditions, including diet[34,36,42]. We identified *A. persici* as the predominant

bacterial species and showed the critical role of its increased abundance in the acquisition

of BL tolerance. *Acetobacter* species, which are well-known symbionts of *Drosophila* and

other insects, are recognized for their acetate production[74]. Acetyl-CoA, derived from acetate,

is essential for lipid utilisation[28,75] and also plays a crucial role in the acetylation of both

histone and non-histone proteins, facilitating adaptive responses through tissue

plasticity[37,76,77]. The elevated TAG levels observed in SL flies may be explained by the

increased acetyl-CoA pool resulting from gut microbiota–derived acetate. The increased

expression of acetic acid production genes in *A. persici* from the SL supports this hypothesis

(Fig. 10d)[66]. Additionally, gut microbiota might act as a barrier against physiological stress

induced by BL toxicity. The significant reduction in gut microbiota following BL irradiation in

SL flies suggests the evolution of mechanisms that utilise bacteria as a direct nutritional

resource or as enzymes (Fig. S8 and Fig. 10f). These findings align with previous research

showing that *Acetobacter* enhances the fitness of the host fly[29,35,43–45]. Notably, *Acetobacter*

infection enhances oxidative stress tolerance in *Drosophila*, whereas oxidative stress

reduces *Acetobacter* abundance, consistent with the findings of Brown *et al.* 2021 (Fig. S8

and Fig. S11)[64]. Several studies in *Pseudomonadota*, including *Acetobacter*, have shown

that gut bacterial enzymes in nematodes and insects adaptively contribute to host stress

tolerance[33,64,78,79]. Consistently, our transcriptome analysis revealed that antioxidant

enzymes are part of the upregulated cluster in *A. persici* from SL (Fig. 10d, f and Data S16),

suggesting their potential role as bacterial metabolism benefiting the host. In addition, the

infection experiments primarily suggested that bacterial abundance strongly influenced the

outcome; yet infection with gut contents did not fully rescue the fecundity of SL (Fig. 5b and

Fig. S13b). This implies that in addition to bacterial abundance, qualitative differences in *A.*

*persici* (such as colonisation status, within the host response, metabolism and bacterial

strains) may also play a critical role in host BL tolerance. Given that bacterial evolution is

likely to be faster than host genetic change, it is plausible that some genetic differences have

arisen in SL-derived *A. persici*. This is suggested by the slightly higher weight gain rate

observed for SL-derived *A. persici* in infection experiments with standardised bacterial

concentrations, and by differences in the growth patterns of isolated bacteria, with SL strains

showing less variance (Fig. S12c and Fig.10g, g''). However, the rate of microbiome evolution

during the experimental evolution process is unknown. The difference in bacterial

phenotypes highlights the importance of investigating bacterial strain dynamics in response

to BL toxicity and host-microbiota co-evolution in future studies.

**Acquisition of BL tolerance in SL via adaptive obesity**

When specific bacteria are implicated in host fitness-related phenotypes, their relative

enrichment often leads to reduced microbiota diversity[33,68]. Given that the microbiota

composition did not differ between strains (Fig. S7a), the initial diversity of the gut microbiota

in our study was probably low. Despite low microbiota diversity, increased abundance of *A.*

*persici*, possibly together with reduced strain diversity (dominance of specific strains),

appears to confer an adaptive phenotype in SL. Characteristic features such as shortened

wings and an elongated midgut were also observed in the SL phenotype. We propose that

gut elongation plays a key role in evolutionary adaptation driven by the gut microbiota. Both

scenarios could contribute to the formation of the SL phenotype: changes in organ size may

be influenced by independent pathways or result from a trade-off in resource allocation. The

increase in bacterial abundance appears to be associated with midgut elongation and genetic

changes, particularly those in the Hippo signaling pathway and the Toll and Imd signaling

pathways (Fig. 4c, Fig. 5f, Fig. 6c, Fig. 7d, e and Fig. S18b). Because our bacterial

quantification was based on CFU measurements from whole-body homogenates, the direct

relationship between midgut elongation and bacterial abundance remains uncertain.

Nevertheless, the observation that SL flies retain an elongated midgut after Abx treatment,

but experience a significant reduction in body weight, suggests that gut microbiota are

necessary for the acquisition of obesity (Fig. S10). The resulting phenotype from laboratory

selection may have been influenced by a combination of epigenetics, de novo mutations,

genetic drift, and variant selection. We are not able to separate these contributions, and

further analysis will be needed to clarify the underlying mechanisms. The *Drosophila* total

sequence length is approximately 140 Mbp[80], with a de novo single nucleotide mutation

(SNM) rate of $3.3 \times 10^{-9}$ per site per generation[81]. This corresponds to an estimated 0.462 de

novo SNMs per generation across the genome. Over 44 generations (as in the WGS

performed), this amounts to approximately 26 de novo SNMs. However, we identified SL-

specific homozygous variants at 1,985 sites (corresponding to 658 genes) shared by the

three replicate lines, a level that cannot be explained by the accumulation of naturally

occurring variants alone. Although de novo mutations may have occurred, we predict that

most of the SL-specific variants, including both SNPs and indels, are derived from standing

variations already present in the ancestral population. The possibility that BL toxicity

increases the SNM rate cannot be ruled out and requires further investigation of the

molecular evolutionary process. We also found that parental irradiation with BL, when

combined with accumulated mutations in later generations of SL flies, is critical for the midgut

elongation and bacterial abundance (Fig. 6b, c). This is consistent with previous studies

showing that parental exposure to environmental stressors, such as starvation and BL toxicity,

induces epigenetic inheritance or alternations in the transcriptome[48,82]. In *Drosophila*, the

midgut undergoes plastic elongation or shortening in response to environmental

conditions[83,84]. This physical expansion of the gut, which increases spatial capacity, offers

several advantages, including more efficient nutrient absorption, increased bacterial

abundance, and maintenance of gut homeostasis[71,85–89]. However, midgut elongation alone,

even after one generation with parental exposure to BL toxicity, is insufficient for BL tolerance

(Fig. 2a and Fig. 6c). Full BL tolerance requires both midgut enlargement and specific genetic

changes. Among these genetic factors, the genes for the neuropeptide *Tk* and its receptor

*TkR99D* are particularly important (Fig. 8c, Fig. 9 and Data S11), as they may regulate lipid

utilisation in response to acetate produced by the gut microbiota[37,62]. Laboratory evolution

studies typically use genetically diverse populations, such as wild populations or multiple

lines from the *Drosophila* Genetics Reference Panel[90]. As we used Canton-S (low genetic

diversity), it is possible that microbiota-mediated lipid accumulation was more optimal than

the development of dedicated lipid synthesis systems driven by host genetic changes.

Microbiota-mediated evolutionary adaptation and co-evolution under low host genetic

diversity remain interesting topics for future research.

**Implications of our findings for pest control**

BL toxicity indicates wavelength-specific effects across insect species and developmental

stages[9]. Reports also show no changes in sensitivity between mutant or overexpression

models of photoreceptors[10], suggesting a complex underlying mechanism. Our study

identifies lipid storage and gut microbiota as critical factors in BL tolerance. In addition, we

have shown that BL toxicity strongly inhibits ovarian development and exerts persistent

suppressive effects even in the absence of lethality (Fig. S2). In the context of pest control,

the combination of BL irradiation with treatments that disrupt insect lipid storage and gut

microbiota could potentially improve efficacy. In parallel with the development of these

methods, research into whether similar tolerance mechanisms exist in other insect species

could lead to more effective chemical-free BL pest control.

**Conclusion**

Our findings highlight the importance of gut microbiota–mediated lipid accumulation as an

adaptive trait for BL tolerance. The elongation of the midgut, associated with an increased

abundance of beneficial bacteria, turns out to be a key factor in this evolutionary

adaptation. Such phenotypes may be driven by genetic and epigenetic changes from

selective pressures and parental BL exposure. These evolutionary responses to excessive

BL exposure demonstrate how host traits evolve under a toxic BL environment by

maximising the benefits provided by gut microbiota. Our study reveals an insect

evolutionary process via the microbiome and presents a concept of evolutionary adaptation

initiated by midgut elongation.

## Methods

### Fly cultures and diet

Flies were maintained at 23–26°C on a 16:8 hour light:dark cycle using cold cathode fluorescent lamps as the light source. Detailed light conditions are provided in Fig. S1. The fly diet consisted of 5 g glucose (Fujifilm Wako Pure Chemicals Co., Osaka, Japan), 6 g dry brewer's yeast (Asahi Group Holdings Ltd., Tokyo, Japan), 1 g agar (Fujifilm Wako Pure Chemicals Co.), 0.4 mL propionic acid (Fujifilm Wako Pure Chemicals Co.), 0.4 mL 70% ethanol (Fujifilm Wako Pure Chemicals Co.), 0.01 g butyl $p$-hydroxybenzoate (Fujifilm Wako Pure Chemicals Co.), and 100 mL distilled water.

### BL irradiation

Virgin flies aged 1–5 days were used. Unless otherwise specified, flies were exposed to 462 nm BL for 3 days. BL irradiation conditions varied across experiments:

Fig. 2b–c, Fig. 5b, Fig. 6b, c, Fig. S2a, b, Fig. S3a, Fig. S8, Fig. S9c, d, Fig. S12b, c and Fig. S13b: Photon flux density $10.3 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$ for females.

Fig. 2a, c, Fig. S3a: $6.5 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$ for males.

Fig. 2d: Continuous BL irradiation for 5 days, $8 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$ for males and $10 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$ for females.

Fig. S4: Continuous BL irradiation for 10 and 4 days, $5 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$ and $15 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$ for females.

Fig. 9: 465 nm BL, $9.5 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$ for females.

Irradiation was performed in a multi-room incubator (LH-30CCFL-8CT; Nippon Medical & Chemical Instruments Co., Ltd, Osaka, Japan) at 24–26°C, with LED panels (IS-mini®, ISL-150 × 150 Series; CCS Inc., Kyoto, Japan) installed on the ceiling. Photon flux density was measured using a high-resolution spectrometer HSU-100S (Asahi Spectra Co., Ltd., Tokyo, Japan) with ND 0050 filter (Asahi Spectra Co., Ltd) and spectroradiometer MS-730 (EKO

500　Instruments Co., Ltd., Tokyo, Japan). For details see Fig. S1. All light condition

501　measurements used in the experiment are included in the source data (Light condition).

## Laboratory selection for BL tolerance

**Artificial selection**

We used the Canton-S strain of *Drosophila* as the ancestral population for laboratory

selection. The Canton S strain used in this study was obtained from Tokyo Metropolitan

University in 2016. It has been maintained under laboratory conditions for at least 30 years

and has been bred in our laboratory in over 1,000 populations over 6 years. Two strains

were established, SL and UCL; the former was subjected to BL toxicity every generation

and the latter was used for controls. Each strain included three replicate lines. For each

strain, 60–100 virgin flies aged 1–5 days were placed in two 60 mm × 90 mm glass Petri

dishes for BL irradiation. SL and UCL flies (positive control group) were exposed to 462 nm

BL for 3 days. Instead of BL irradiation, the UCL, used as a negative control group, was

kept in darkness for 3 days. We followed this standard approach as dark conditions are

commonly used as a control in BL toxicity studies[7,10,11,48].

**Climbing assay**

Irradiated flies were placed in a breeding environment (for details, see "Fly cultures and diet"

in Methods) for a 2-day recovery period. For the data shown in Fig. S3a and Fig. S9c, the

impact of BL toxicity was assessed in a 4-minute climbing assay and the flies were

categorized into four states:

"Low impact":

1. Intact: capable of climbing

"High impact" (unable to climb):

2. Phase 1: alive but unable to climb

3. Phase 2: alive but non-ambulatory

4. Phase 3: dead

**Artificial selective breeding and evaluation of fecundity**

"Low impact" SL flies were allowed 5 days for mating and egg-laying. In the early stages of selection, there was no clear difference in climbing ability, and fecundity was low; thus, we used as many intact individuals as possible as parents. From the 17th generation, when tolerance began to emerge, we prioritised intact individuals that climbed above the 2 cm threshold and selected preferentially for higher tolerance. Each vial containing 3 mL of diet medium 3-mL contained 9–13 pairs, with a total of 4–8 vials (for details, see Source data (Fig. 2a)). After 5 days, the diet medium containing eggs and larvae was transferred to a larger plastic box (40 mL diet medium, 72 × 72 × 100 mm; Insect Breeding Square Dish, SPL Life Sciences, Gyeonggi-do, Korea) to ensure sufficient food supply. One box was used for every 4 or 5 vials. UCL parents were randomly chosen from the negative control group (for details, see "Laboratory selection for BL tolerance -Artificial selection-" in Methods) and reared under the same conditions as the SL flies. The total number of pupae was counted and divided by the number of parents to determine the fecundity per female for each generation.

**Negative geotaxis climbing assay in non-irradiated conditions**

Young adult flies, which typically climb 4 cm in 4 seconds, were tested for negative geotaxis[91]. In Fig. S3b, ten flies aged 5–10 days were placed in an empty vial, gently tapped three times to ensure they reached the bottom and given 10 seconds to climb. Climbing rate was defined as the flies reaching a 6 cm reference line within this time. Each vial was tested three times with a 1-minute interval between trials and the mean climbing rate was calculated.

**Survival assays**

Virgin flies aged 5–10 days were used. In the continuous BL irradiation assay, 50 or 42 flies were placed in each glass Petri dish containing standard diet medium and exposed to 462

nm BL (Fig. 2d and Fig. S4). In the starvation assay (Fig. 4g), 10 flies were placed in each

vial with 1% (w/v) agar (Fujifilm Wako Pure Chemicals Co.) in distilled water. In the

oxidative stress assay, 10 flies were placed in each vial with a medium prepared with either

10 mM paraquat dichloride standard (Fujifilm Wako Pure Chemicals Co.) (Fig. 4h) or 10

mM 1,1'-dimethyl-4,4'-bipyridinium dichloride (Tokyo Chemical Industry Co., Ltd., Tokyo,

Japan) (Fig. S11a, b), 5% (w/v) glucose (Fujifilm Wako Pure Chemicals Co.), and 1% (w/v)

agar (Fujifilm Wako Pure Chemicals Co.) in distilled water. The surviving flies were

manually counted every 2 h. Survival curves were visualised using the KaplanMeierfitter

from the lifelines (Python version 3.8.3 version 0.25.9).

**Measurement of fecundity**

To measure fecundity (Fig. 2a, Fig. 5b, Fig. 9, and Fig. S13b), after 3 days of darkness

(non-irradiation) or BL irradiation followed by a 3-day recovery period, 5 surviving females

were randomly chosen and mated with 5 non-irradiated virgin males (aged 1–10 days) in

individual pairs within vials containing 5 mL of diet medium for 4 days of egg-laying. The

total numbers of pupae and emerged adults were counted to evaluate fecundity and

emergence rate (in Fig. 2a and Fig. 2c, only the total number of pupae was measured).

Fig. 2c, Fig. 5b and Fig. S13b used non-irradiated Abx-treated males from the UCL or SL

and Fig. 9 used non-irradiated Canton-S males.

**Measurement of body weight, organ size and oocyte number**

Phenotypic measurements were performed on non-irradiated flies in Fig. 3a, b, Fig. 4c, Fig.

5a, d, f, Fig. 6c, Fig. S8 and Fig. S14 and on BL-irradiated flies in Fig. 2b, Fig. S2a, b and

Fig. S8. As ovarian development can progress with age, we ensured that phenotypic

comparisons between strains were made at the same chronological age[19]. Virgin flies aged

4–12 days were anesthetized by chilling on ice. Body weight was measured using an

AUW220D balance (Shimadzu, Kyoto, Japan). Wings or ovaries were dissected and

photographed under a SteREO Discovery.V12 stereomicroscope and an Axiocam 305

color camera (Carl Zeiss, Oberkochen, Germany). Wing length was measured as the

distance between the alula opening and the distal edge of the 3rd longitudinal vein, using

Fiji ImageJ2 (version 2.3/1.54f), following a previous method[92]. Stage 14 oocytes were

counted under the stereomicroscope. Midgut lengths of regions R1 to R5 were measured

manually from the midgut images using Fiji ImageJ2[89].

**Lipid droplet staining and imaging**

In Fig. 4b and b', we performed dissection and fluorescence microscopy to visualise lipid

droplets in the fat body of the abdominal epithelium of virgin female flies aged 6–12 days

kept in non-irradiated conditions. After removing other tissues from the abdomen, we fixed

the abdominal epithelium in PBS containing 4% paraformaldehyde (Fujifilm Wako Pure

Chemicals Co.) for 20 minutes. After fixation, we washed the samples twice with PBS

(Gibco, Carlsbad, CA, USA) for 15 minutes each time. The fixed abdominal epithelium was

stained with Nile Red (Fujifilm Wako Pure Chemicals Co.) at a concentration of 10 µg/mL

overnight according to a previous method[93]. Following this, we washed the samples twice

with PBS for 10 minutes each time and then stained them with NucBlue (Thermo Fisher

Scientific, Waltham, MA, USA) for 3 minutes, followed by a 5-minute wash with PBS. All

staining procedures were conducted in a light-shielded environment at room temperature.

Imaging was performed under an Eclipse Ti2 microscope equipped with a DS-Ri2 camera

(Nikon, Tokyo, Japan). The images were analysed using the Python package OpenCV

(Python version 3.8.3 version 4.5.3.56) to obtain the mean fluorescent intensity of Nile Red.

**Lipid extraction and derivatization of fatty acids**

Lipids were extracted using the Bligh and Dyer method[94,95]. Two virgin female flies aged 6–

12 days and kept in non-irradiated conditions were homogenised in 100 µL of PBS (Gibco),

added 600 µL of methanol (Fujifilm Wako Pure Chemicals Co.) and chloroform (Fujifilm

Wako Pure Chemicals Co.) in a 2:1 (v/v) ratio, vortexed for 1 minute, and shaken at 1000

rpm for 2 h at 4°C. Next, we added 200 µL of chloroform and 250 µL of distilled water,

vortexed for 1 min, centrifuged at 9000 rpm for 2 min, and collected the organic lower

phase. We then added 400 µL of chloroform to the remaining mixture, vortexed for 1

minute, and centrifuged as above. The two organic extracts were pooled and dried under

vacuum. The dried lipids were derivatized using the Fatty Acid Methylation Kit (Nacalai

Tesque, Inc., Tokyo, Japan) and the Fatty Acid Methyl Ester Purification Kit (Nacalai

Tesque, Inc.). FAMEs were collected.

### 617     Gas chromatography mass spectrometry (GC-MS) analysis

The FAMEs were analysed using a GC-MS instrument (GCMS-QP2010 Ultra (Shimadzu))

equipped with a DB-5 ms column (30 m × 0.25 mm ID, 0.25 µm film thickness) (J&W

Scientific, Folsom, CA, USA). Helium was used as the carrier gas at a pressure of 100 kPa.

The column oven was set at 40°C for 5 minutes. The temperature was increased at a rate

of 10°C/min to 280°C and maintained at 280°C for 10 minutes. The injection port

temperature was 220°C, and the analysis was performed in split mode. Fatty acids were

identified using a similarity search against the standards Supelco F.A.M.E. Mix C4-C24

(Supelco, Bellafonte, PA, USA) and relevant references[96,97]. The total ion chromatogram

area for each compound was measured. On the basis of the obtained data, we compared

the total amount of FAMEs. The log-transformed average values of each compound in each

replicate line were visualized using a cluster map. Clusters were classified by hierarchical

clustering using the Ward method based on Euclidean distances between compound

amounts[98].

### 632     Measurement of TAG

Relative TAG levels were measured by homogenising 10 virgin female flies aged 6–12

days and kept in non-irradiated conditions, in 1 mL of PBS (Fujifilm Wako Pure Chemicals

Co.) containing 0.1% Triton X-100 (Nacalai Tesque Inc.). The homogenate was heated at

70°C for 10 minutes and then centrifuged at 20,000 × $g$ for 15 minutes at 4°C. A 10-μL

aliquot of the supernatant was used for TAG measurement using the LabAssay Triglyceride

kit (Fujifilm Wako Pure Chemicals Co.). For protein normalization, protein concentration

was measured using the Bradford method with the Protein Assay CBB Clean Up Kit and

Protein Assay CBB Solution (both from Nacalai Tesque Inc.), with the standard calibration

curve prepared using bovine serum albumin solution (Nacalai Tesque Inc.). Absorbance

was measured using the BioTek Synergy H1 Multimode Reader (Agilent Technologies Inc.,

Santa Clara, CA, USA). Relative TAG levels were normalised to protein content, adjusted

on the basis of the average values from UCL (Fig. 4f and Fig. 9) or non-treated UCL (Fig.

5e), and the $Log_2$ fold change was computed accordingly.

**Measurement of reflectance and transmittance**

Virgin female flies, aged 6–12 days and kept in non-irradiated conditions, were

anesthetized on ice. The abdominal reflectance and transmittance were measured by

capturing hyperspectral images under a BX51WI microscope (Olympus, Tokyo, Japan)

fitted with an SC-108 hyperspectral imaging system (EBA Japan Co., Ltd., Tokyo, Japan).

**DNA extraction**

Virgin female flies, aged 5–8 days and kept in non-irradiated conditions, were used. For

Fig. 7 and Fig. S7a, the same sample of 30 flies was used, with 10 flies used specifically for

Fig. S7b'. The flies were rinsed in a 1.5-mL tube with 99% ethanol (Fujifilm Wako Pure

Chemicals Co.) for 30 seconds. Then, 180 μL of PBS and beads were added, and the flies

were homogenised at 3500 rpm for 30 seconds using the Micro Smash MS100R (Tomy

Seiko Co., Ltd., Tokyo, Japan) while keeping the mixture cold. DNA was extracted using

the DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) according to the kit protocols.

## 16S rRNA amplicon sequencing (V3–V4 hypervariable region)

Virgin female adult flies were used after surface sterilisation under non-irradiated conditions. For details of DNA samples, see "DNA extraction" in Methods. Quality checks were performed using the 2100 BioAnalyzer (Agilent Techologies Santa Clara, CA, USA), followed by library preparation with the NEBNext® Ultra™ II DNA Library Prep Kit (Illumina, San Diego, CA, USA). DNA amplification was performed by PCR using the V3-V4 region primers 341F (5'-CCTAYGGGRBGCASCAG-3') and 806R (5'-GGACTACNNGGGTATCTAAT-3'). PCR cycling conditions were as follows: 95°C for 3 minutes, followed by 25 cycles of 95°C for 30 seconds, 55°C for 30 seconds, 72°C for 30 seconds, with a final extension at 72°C for 5 minutes and holding at 4°C. Amplicon sequencing was conducted on an Illumina NovaSeq 6000 with the following parameters: paired-end reads of 250 bp × 2 (PE250) and an output of 50,000 reads per sample. In silico analyses were performed on a Mac OS (Catalina version 10.15) terminal using Jupyter Notebook (Python version 3.9 version 6.4.11). Quality was assessed with FastQC (version 0.11.8). Data were analysed using QIIME 2 (version 2023.7) with the Silva 138 99% OTUs full-length database (MD5: b8609f23e9b17bd4a1321a8971303310). The QIIME dada2 denoise-paired parameters were set as follows: --p-trim-left-f 30, --p-trim-left-r 30, --p-trunc-len-f 250, and --p-trunc-len-r 250. To improve classification accuracy, taxonomic filtering was performed by removing all unassigned taxa using the qiime taxa filter-table command, resulting in a refined feature table. Sequences of operational taxonomic units were classified to determine gut microbiota composition. Relative abundance calculations were performed using the qiime feature-table relative-frequency function.

## Bacterial abundance and identification

Body weights of 4 virgin adult females were measured after 3 days of non-irradiation or BL irradiation followed by a 3-day recovery period (Fig. 5a–a', Fig. 6b, Fig. S8). 4 surviving females aged 6–12 days were randomly chosen, immersed in 70% ethanol for 1 minute,

washed three times with PBS homogenised in 500 μL of PBS in a 2-mL tube (stock

solution), and diluted 1:100 with PBS. A 10-μL aliquot was plated onto an MRS agar plate

(5.5 g MRS broth (Becton, Dickinson and Company, Franklin Lakes, NJ, USA), 1 g agar

(Fujifilm Wako Pure Chemicals Co.), and 100 mL distilled water). The plates were

incubated for 72 hours at 30–31°C, and colony-forming units (CFU) were counted. Single

bacterial colonies were isolated, and their 16S rRNA genes were amplified by PCR using

universal bacterial primers 27F (5' AGAGTTTGATCCTGGCTCAG 3') and 1492R (5'

GGTTACCTTGTTACGACTT 3'). PCR cycling conditions were as follows: 95°C for 3

minutes, followed by 30 cycles of 94°C for 30 seconds, 55°C for 45 seconds, 72°C for 1

minute, with a final extension at 72°C for 7 minutes and holding at 4°C. Sequencing of the

amplified 16S RNA gene fragments was performed on an ABI3500 Sanger sequencing

platform (Thermo Fisher Scientific) using the BigDye™ Terminator v3.1 Cycle Sequencing

Kit (Thermo Fisher Scientific)[99,100]. Bacterial species were identified using Microbial

Nucleotide BLAST.

**Abx treatment**

705    The Abx was prepared according to established protocols[101]; the diet was supplemented

706    with 100 μg/mL streptomycin sulfate (Fujifilm Wako Pure Chemicals Co.), 50 μg/mL

707    tetracycline hydrochloride (Fujifilm Wako Pure Chemicals Co.), and 200 μg/mL rifampicin

708    (Fujifilm Wako Pure Chemicals Co.). Flies were reared on a diet containing Abx from the

709    embryo stage to adult.

**Capillary feeding (CAFE) assay**

712    The assay was adapted from a previous study with minor modifications[102]. Briefly, a group

713    of four male flies, aged 5-8 days, was kept in a plastic vial with two glass capillaries

714    inserted (BF100-50-15, Sutter Instrument, CA, USA). The gender was selected because

715    female flies often lay eggs inside the capillaries, which prevents precise measurement.

Flies were fed for 24h with liquid food (5% sucrose, 2% yeast extract, 0.25% nipagin in

ethanol, and 0.005% Sulforhodamine B sodium salt (red dye) in water), and images of the

capillaries were taken before and after the experiments. Reduction in liquid volume was

quantified to estimate consumption. To account for evaporation, control capillaries without

flies were included and their loss of fluid was used as a correction.

**Parental experience of BL irradiation**

Virgin flies aged 1–5 days were BL-irradiated for 3 days (462 nm BL, $6.5 \times 10^{18}$

photons·m$^{-2}$·s$^{-1}$ for males and $10.3 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$ for females) and allowed to

recover for 3 days. Non-irradiated parents were used as controls to ensure that the

offspring had no parental experience of irradiation. Flies exposed to BL irradiation were

assumed to experience BL toxicity. Parental choice was made without assessing

differences in survival status, such as climbing ability, to produce offspring with different

parental experiences. Randomly chosen males and females were allowed to mate and lay

eggs for 4 days.

**Infection with gut contents or *A. persici* cocktails**

In Fig. 5b, Fig. S12a and Fig. S13, 30 virgin female flies aged 5–8 days were immersed in

70% ethanol for 1 minute, washes three times with PBS and homogenised in 500 µL of

PBS in a 2-mL tube (stock solution). The latter was diluted 1:10 with PBS. To infect with the

gut microbiota, 20 µL of the diluted stock solution was added to the diet, and virgin females

aged 1–4 days were maintained on this diet for 4 days. PBS was used as a control. Body

weight was measured before and after infection. Climbing ability and fecundity were

evaluated after 3 days of non-irradiation or BL irradiation followed by a 3-day recovery

period. In Fig. S12b, *A. persici* was cultured and isolated on MRS plates for 48 hours.

These isolates, along with the standard strain of *A. persici* (JCM 25330), were cultured in

200 µL of MRS broth (5.5 g MRS broth (Becton, Dickinson and Company) and 100 mL

distilled water) in 96-well plates for 72 hours. Bacterial suspensions were centrifuged to

obtain pellets, and the 10-µL bacterial culture pellet was diluted 1:30 with PBS and

inoculated into 20 µL of diet. Autoclaved MRS was also diluted 1:30 with PBS, and 20 µL of

this mixture was used as the control.

**Whole-genome resequencing, variation estimation, variant identification,**

**and enrichment analysis**

Thirty virgin adult females were used after surface sterilisation under non-irradiated

conditions. For details of DNA samples, see "DNA extraction" in Methods. Quality checks

were performed using a BioAnalyzer, followed by library preparation with the NEBNext®

Ultra DNA Library Prep Kit (Illumina). Sequencing was conducted on an Illumina NovaSeq

6000 with the following parameters: paired-end reads of 150 bp (PE150), a data output of

10 G bases per sample, and approximately 66.7 million reads per sample. In silico

analyses were performed using a macOS (Catalina version 10.15.7) terminal and Jupyter

Notebook (Python version 3.9 version 6.4.11). The quality of the fastq files was assessed

using seqkit stat (version 0.16.1) and then fastqc (version 0.11.8). Reads were trimmed

using Trimmomatic (version 0.39) with the following parameters: LEADING:20,

TRAILING:20, SLIDINGWINDOW:4:15, and MINLEN:20, based on Illumina adapter

sequences (CTGTCTCTTATACACATCT,

AGATCGGAAGAGCACACGTCTGAACTCCAGTCA,

AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT). Trimmed reads were re-evaluated

with seqkit stat and fastqc to confirm their quality. Mapping was performed using the

reference genome BDGP6.46, INSDC Assembly GCA_000001215.4 (details provided in

Data S9). Indexing and mapping were conducted using bwa (version 0.7.17), followed by

conversion from SAM to BAM format using samtools (version 1.10). PCR duplicates were

marked with GATK (version 4.2.6.1), and alignment statistics were calculated from the BAM

file. Paired-end reads were merged using samtools, and the merged BAM file was sorted.

Consensus sequences for UCL, SL, and Canton-S strains were generated using the

reference genome and VCF files. Gene sequences were extracted based on BDGP

6.32.109.chr.gtf3 annotations and processed using BioPython's SeqIO module. Gene

sequence alignments between strains were performed using BioPython's AlignIO module,

MAFFT (version 7.0), and variation for each gene were calculated using Canton-S

(SRX8038113)[50] as a reference. Variation was calculated as (the number of mismatched

bases to the reference gene / the total sequence length of the gene) × 100. Gaps were

counted as 0.5 to account for insertions, deletions and sequencing errors. SL-specific

variation was determined by subtracting the variation of each SL replicate line from the

mean variation of UCL replicate lines 1-3. Genes with a mean variation of more than 5%

across replicate lines 1-3 were classified as SL-specific variation. Variants were called

using sambamba (version 1.0.1) and varScan (version 2.4.6) with the following parameters:

sambamba mpileup, Heterozygous: SNP; pileup2snp --min-coverage 10 --min-var-freq 0.20

--p-value 0.05 (filter; default), Indel; pileup2indel --min-coverage 10 --min-var-freq 0.20 --p-

value 0.10 (filter; --min-reads2 4 --min-var-freq 0.15 --p-value 0.05), Homozygous: SNP;

pileup2snp --min-coverage 10 --min-var-freq 0.90 --p-value 0.05 (filter; default), Indel;

pileup2indel --min-coverage 10 --min-var-freq 0.90 --p-value 0.10 (filter; --min-reads2 4 --

min-var-freq 0.15 --p-value 0.05). Homozygous variants were identified by extracting rows

with matching Cons and VarAllele values. Common chromosomes and positions across

replicates (lines 1, 2, and 3) within each strain were defined to identify strain-specific

shared homozygous variants. Shared and unique homozygous variants among UCL, SL,

and Canton-S[102] were analysed. homozygous variants were annotated using SnpEff

(version 5.0) and visualized using IGV (version 2.8.12). Variants classified as having a low

impact ("LOW") by SnpEff were excluded from the analysis list (full list available in Data

S11). Enrichment analysis was conducted with DAVID (version 2023q4) and KEGG

(version 108.1).

## RNA extraction

Ten virgin female flies, aged 5–8 days and kept in non-irradiated conditions, were frozen in liquid nitrogen and homogenised in Trizol (Molecular Research Center, Inc., Cincinnati, OH, USA). Chloroform (Fujifilm Wako Pure Chemicals Co.) was then added, and the mixture was vortexed twice and centrifuged at 20,000 × $g$ for 10 minutes at 4°C. The upper aqueous phase was transferred to a new tube, and 200 μL of isopropyl alcohol was added. The mixture was vortexed, incubated at room temperature for 10 minutes and centrifuged again at 20,000 × $g$ for 10 minutes at 4°C; the supernatant was discarded. To wash the RNA pellet, 400 μL of 75% ethanol (prepared with DEPC-treated water) was added, followed by vortexing. The sample was then centrifuged at 20,000 × $g$ for 5 minutes at 4°C. The supernatant was discarded, and the pellet was centrifuged once more for 1 minute. The pellet was air-dried and resuspended in DEPC-treated water.

## RNA-sequence and enrichment analysis

Quality assessment was initially performed using a BioAnalyzer. Library preparation was conducted with the NEBNext Ultra II Directional RNA Library Prep Kit (Illumina) and the NEBNext Poly(A) mRNA Magnetic Isolation Module (New England Biolabs, Ipswich, MA, USA). Sequencing was performed on an Illumina NovaSeq 6000, with a read length of PE150 (150 bp × 2 paired-end), generating a data output of 6 G bases per sample and approximately 40 million reads per sample. In silico analyses and read trimming followed the protocols described in the "Whole-genome resequencing, variant identification, and enrichment analysis" section in Methods. Poly(A) tails were removed using cutadapt, and mapping was conducted with hisat2 (version 2.2.1) using the BDGP6.46 reference genome (INSDC Assembly GCA_000001215.4) (for details, see Data S13). Gene-wise read counting was accomplished using HTSeq (version 2.0.4) with the dmel-all-r6.54.gtf annotation file. Read counts were normalised with edgeR (version 3.40.2) using the trimmed mean of M values method. Subsequent principal component analysis and

differential expression analysis were carried out with edgeR and scikit-learn (version 1.0.2).

DEGs were identified by comparing non-treated SL with non-treated UCL, Abx-treated SL,

or Abx-treated UCL groups, applying a false discovery rate–adjusted *p*-value threshold of

<0.05. Python libraries such as pandas (version 2.0.3), numpy (version 1.25.2), scipy

(version 1.11.2), and stats were used to filter DEGs with *p*-values below 0.05. For the

DEGs, $\log_2$-fold change values and Z scores normalised within the range of −2 to 2 were

calculated. These Z scores were then used to create a cluster map using the Ward method,

which classified genes and treatments on the basis of their similarity patterns. Enrichment

analysis was performed using DAVID and KEGG. Gene lists related to specific biological

functions were retrieved from KEGG (for TCA cycle, oxidative phosphorylation and Toll and

Imd signaling pathways) and FlyBase (for lipid metabolic processes). In Fig. 8c and Fig.

S18d, heatmaps illustrating functional profiles were generated using $\log_2$-fold change

values. These values were compared for each gene and each group and normalised

against the mean value of the non-treated UCL group.

**Genetics and evaluation of TAG levels and BL tolerance**

We collected genetically engineered flies with known elevated TAG levels and reared them

at 23–25°C under a 12-hour light/dark cycle. Using the GAL4/UAS system and null

mutants, we generated flies with RNAi knockdown or overexpression of target genes. The

following strains from the Bloomington *Drosophila* Stock Center were used (see Data S15):

Null mutants: w;*Akh* (84448), w;*Bmm* (15828), *yw,Pdgy* (19990), *yw;Chico* (14337), *w;InR*

(8267), *w;Ilp 2 3 5* (30889), *w;Lk6* (87087)

GAL4 drivers: *yw;Lsp2* (6357), *w;ppl* (58768), *w;Tk* (51973)

UAS strains: *yv;UAS-AkhR$^{RNAi}$* (51710), *yv;UAS-Pect$^{RNAi}$* (63710), *yv;UAS-Pten$^{RNAi}$*

(33643), *yv;UAS-Bmm$^{RNAi}$* (25926), *y sc v sev;UAS-Stim$^{RNAi}$* (52911), *y sc v sev;UAS-Lsd-*

*1$^{RNAi}$* (65020), *yv;UAS-Tk$^{RNAi}$* (25800), *yw;UAS-Gbb* (63059), *yw;UAS-InR* (8263), *w;UAS-*

*Lsd-2* (98116), *w;UAS-Lk6* (8709), *w,UAS-Rpr1* (5823)

We used F1 progenies for the experiments, obtained by crossing GAL4 driver males with

UAS effector females. As controls, we used wild-type Canton-S and w1118 genetically

washed by crossing with the Canton-S flies, F1 progenies from crosses of w1118 with UAS

effectors, and F1 progenies from crosses of GAL4 drivers with *y sc v;eGFP^RNAi* (41555) and

*y sc v;eGFP^RNAi* (41556) UAS effectors. Virgin female flies aged 1–5 days were irradiated

with either darkness or BL for 3 days and recovered for 3 days. Their survival rates were

assessed, and 5 flies were randomly chosen to mate with 5 Canton-S males for 4 days (for

details, see "Measurement of fecundity" in Methods). TAG storage was measured in virgin

female flies aged 7–12 days. Relative TAG levels are presented as $\log_2$-fold changes

relative to the mean values across all treatments. Fecundity was calculated as $\log_2$-fold

changes relative to the non-irradiation condition for each treatment. Data are from SL and

UCL (replicate lines 1, generation 57).

**Phylogenetic tree based on the 16S rRNA gene**

16S rRNA gene sequences were obtained from the Integrated Microbial Genomes (IMG)

database, with the addition of *A. persici* DS-1 strain[66] for *A. persici*. Based on previous

studies, *Neokomagataea thailandica* and *Swingsia samuiensis* were used as outgroups[103].

Sequence lengths ranged from 1,391 to 1,498 bp. The phylogenetic tree in Fig. S19 was

constructed by including the *A. persici* sequence (397 bp) from the dominant OTU ID,

which accounted for 89-98% of the microbiome analysis in Fig. S7a. Based on these

results, the phylogenetic position of *A. persici* in our laboratory was inferred in Fig. 10a.

Phylogenetic analyses were performed using MEGA (version 11.013), with alignments

performed using ClustalW. A maximum likelihood (ML) method with 1,000 bootstrap

replicates was used (substitution type: general time reversible model; ML heuristic method:

Nearest-Neighbour Interchange; initial tree for ML: NJ/BioNJ). The resulting phylogenetic

tree was visualised using iTOL (version 7.0).

### *A.persici* circular genome visualisation and transcriptome analysis

The complete genome of *A. persici* (GCA_002006565.1_ASM200656v1) was used for

circular genome visualisation and transcriptome analysis for reference genome. The

circular genome representation was generated using GenoVi (version 0.2.16)[104]. After

mapping to the host genome, reads that unmatched to the host were extracted and

subsequently mapped to the complete *A. persici* genome. Unmapped reads were filtered

from the SAM file using Samtools (samtools view -b -f 4), followed by conversion to FASTQ

format. Sequence quality was assessed using FastQC and overrepresented sequences,

excluding *A. persici*, were removed. Reads were trimmed using Trimmomatic with the

following parameters: LEADING:20, TRAILING:20, SLIDINGWINDOW:4:15, and

MINLEN:20. The *A. persici* genome index was generated using HISAT2, and reads were

aligned using parameters: hisat2 -x reference genome index -1 -2 -k 3 --mp 2.1 --np 1 --rdg

10.5 --rfg 10.5 --seed 42 --score-min L,0,-0.9) (for details, see Data S16). Gene expression

was quantified using featureCounts (subread-1.5.2.-MacOSX-x86_64, version 2.0.8) and

parameters: featureCounts -t CDS -g gene_id -a genomic.gft (GCA_002006565.1) -M -p.

Gene expression levels were quantified using transcripts per million (TPM). Reads per

kilobase (RPK) were calculated as expression / (length / 1000). TPM values were then

derived by normalising the RPK of each gene to the total RPK within its respective GeneID.

Sequence depth was estimated using the formula (total bases mapped to the gene) /

(target gene length). DEGs were identified by comparing the UCL and SL groups using an

FDR adjusted $p$-value threshold of <0.05. To quantify changes in gene expression, we

calculated the $log_2FC$ of the transcripts. The mean TPM was calculated for each gene in

the UCL group, and the fold change was determined as the ratio of individual sample TPM

values to this mean. The dataset was then $log_2$ transformed. Enrichment analysis was

performed using ShinyGO (version 0.82) and Gene ontology (GO) biological process

(version Ensembl 92). Gene lists related to oxidation-reduction process was retrieved from

GO biological process. The genomic.gft (GCA_002006565.1), UniProt (version 2025_01)

and KEGG databases were used to determine protein identities, enzyme classifications and

functional annotations for each gene. For details of the host transcriptome analysis and

visualisation, see "RNA-sequence and enrichment analysis" in Methods.

**Growth experiments of isolated *A. persici***

*A. persici* was isolated from 10 adult females and cultured on MRS plates for 48 hours. 50

isolates were pre-cultured in 200 μL MRS broth (5.5 g MRS broth and 100 mL distilled

water) in 96-well plates for 72 hours. Bacterial growth was then assessed by culturing in 50

mL of MRS broth with continuous shaking at 60 rpm. Cultures were maintained at 30-31°C

throughout the incubation period. Optical density (OD) was measured using a compact

rocking incubator equipped with a spectrophotometer (TVS062CA ADVANTEC Tokyo,

Japan).

**Statistical analysis**

**State space model and Bayesian estimation**

The state space model is a statistical framework designed to analyse time series data with

autocorrelation, based on state and observation equations[105–107]. It was applied to estimate

fecundity across generations, incorporating replicate lines as random effects. In Fig. 2a,

intergenerational fecundity data were analysed using a Bayesian state space model

implemented with pystan (Python 3.9, version 2.17.0.0). This model included the following

components:

Data block: Defined the number of generations (length of the time series data), a vector of

explanatory variables, and three outcome variables corresponding to each replicate line,

each following a normal distribution.

Parameters block: Estimated a vector of levels with drift components (*mu*), vectors of

random effects (*r1*, *r2*, *r3*), coefficients of the explanatory variables (*b1*, *b2*, *b3*), the

standard deviation representing the magnitude of the drift component ($s\_z$), and the

standard deviations of the random effects ($s\_r1$, $s\_r2$, $s\_r3$).

Transformed parameters block: Calculated the expected values for each outcome variable

(*lambda1*, *lambda2*, *lambda3*) as the sum of the level with drift component (mu), the

product of the explanatory variables' coefficients (*b*) and values (*ex*), and the random

effects (*r*).

Model block: Assumed that the random effects followed a normal distribution with a mean

of 0 and a given standard deviation. The state equation, *mu[i] ~ normal(2 * mu[i-1] - mu[i-2],*

*s_z*), expressed changes in the level with drift components using a second-order local

linear regression. The observation equations, *y1[i] ~ normal(lambda1[i], sigma1), y2[i] ~*

*normal(lambda2[i], sigma2)*, and *y3[i] ~ normal(lambda3[i], sigma3)*, defined the observed

values as following a normal distribution.

Generated quantities block: Sampled the predicted values for each outcome variable using

the normal distribution.

Random numbers were generated according to the posterior distribution using the Markov

Chain Monte Carlo (MCMC) method with parameters set to iter=10000, chains=4, and

control={'adapt_delta': 0.8, 'max_treedepth': 20}. Convergence was confirmed by ensuring

that the *R*hat of the predicted values was <1.10 and by examining the trace plots (for

details, see Data S2).

**Hierarchical Bayesian model**

Unless otherwise specified, hierarchical uncertainty estimation, accounting for random

effects, was performed using Hierarchical Bayesian Model (HBM)[108]. The model

specification and execution used the following tools and versions:

Software: R (version 4.0.1), rstan (version 2.32.2, Stan version 2.26.1), brms (version

2.20.4), and bayesplot (version 1.10.0).

Bayesian model comparison: The Bayes factor ($BF_{10}$) quantifies the strength of evidence

for the alternative hypothesis ($H_1$) over the null hypothesis ($H_0$)[109]. A $BF_{10} > 1$ indicates

support for $H_1$, whereas a $BF_{10} < 1$ favours $H_0$. Guidelines for interpretation are as follows:

$BF_{10} > 3$ indicates moderate evidence, $BF_{10} > 10$ indicates strong evidence, and $BF_{10} > 100$

provides conclusive support for $H_1$. Conducted using $BF_{10}$ with the loo function, computing

*exp(0.5 * (elpd1 - elpd2))*, where elpd1 denotes the model's expected log pointwise

predictive density (elpd_loo) and elpd2 denotes that of the null model.

MCMC sampling: Parameters were set to seed=1, chains=4, iter=20000, warmup=5000,

and thin=4. If convergence was not attained, adjustments were made to iter=25000 and

warmup=10000, ensuring that $R$hat values remained below 1.10.

Prior distributions: Uniformly set to Gaussian.

Model details: Comprehensive details including the response variable, predictors, random

effects, and estimation results are available in Data S1, Data S3, 4, Data S6–8, Data S16.

Results included posterior median estimates, Bayesian 95% credible intervals (CI), and

approximate $p$-values ($p$MCMC) computed from posterior sampling.

Posterior sampling and $p$MCMC calculation: To test whether the posterior distribution of the

difference in each treatment was above or below 0, we followed references[110–113]. Posterior

samplings of each parameter were stored in diff_samples, and pairwise comparisons of

treatments were made using the formula: $p$MCMC = 2 * min(mean(diff_samples > 0),

mean(diff_samples < 0)).

**Logistic function**

Lethal dose: The survival rate at each dose was calculated by assigning a value of 1 if the

survival time exceeded the median survival time of UCL at $10 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$ (88 h)

and 0 otherwise. The $LD_{50}$ was estimated using a logistic function: The logistic function,

defined as $f(x) = 100 / (1 + exp(slope \times (x - LD_{50})))$, describes the survival probability as a

sigmoidal function of dose.

Bacterial growth curves: modelled using a logistic function defined as $f(t) = K / (1 + exp(-r \times$

$(t - t_0)))$, where $K$ is the maximum optical density (OD), $r$ is the growth rate, and $t_0$ is the

midpoint of the growth phase. Parameter estimation was performed by fitting this function

to the OD measurements over time.

**Correlation and Regression analysis**

The Pearson correlation coefficient or linear regression and 95% confidence intervals were

calculated using the pearsonr function or linregress function from the scipy.stats library.

# Data availability

991 16S rRNA amplicon sequencing, whole-genome resequencing, and RNA-sequence data

992 have been deposited in the DNA Data Bank of Japan (DDBJ). All raw read data have been

993 uploaded to DDBJ under BioProject PRJDB19030, PRJDB19027, and PRJDB19028. The

994 source data for all figures are available in the supplementary files. Detailed experimental

995 methods are available on reasonable request from the corresponding author, Y.T.

# Code availability

997 The code for all analyses and visualisations is available on reasonable request from the

998 corresponding author Y.T.

# References

1. Mallet, J. The evolution of insecticide resistance: Have the insects won? *Trends Ecol. Evol.* **4**, 336-340 (1989). doi: 10.1016/0169-5347(89)90088-8

2. Barathi, S., Sabapathi, N., Kandasamy, S. & Lee, J. Present status of insecticide impacts and eco-friendly approaches for remediation-a review. *Environmental Research* **240**, 117432 (2024). doi: 10.1016/j.envres.2023.117432

3. Spitschan, M., Aguirre, G. K., Brainard, D. H. & Sweeney, A. M. Variation of outdoor illumination as a function of solar elevation and light pollution. *Sci. Rep.* **6**, 26756 (2016). doi: 10.1038/srep26756

4. Song, B. M. & Lee, C. H. Toward a mechanistic understanding of color vision in insects. *Front. Neural Circuits.* **12**, 16 (2018). doi: 10.3389/fncir.2018.00016

5. Numata, H., Miyazaki, Y. & Ikeno, T. Common features in diverse insect clocks. *Zool. Lett.* **1**, 10 (2015). doi: 10.1186/s40851-014-0003-y

6. Kavakli, I. H., Ozturk, N. & Gul, S. DNA repair by photolyases. *Adv. Protein. Chem. Struct. Biol.* **115**, 1–19 (2019). doi: 10.1016/bs.apcsb.2018.10.003

7. Hori, M., Shibuya, K., Sato, M. & Saito, Y. Lethal effects of short-wavelength visible light on insects. *Sci. Rep.* **4**, 7383 (2014). doi: 10.1038/srep07383

8. Hori, M. & Suzuki, A. Lethal effect of blue light on strawberry leaf beetle, *Galerucella grisescens* (Coleoptera: Chrysomelidae). *Sci. Rep.* **7**, 2694 (2017). doi: 10.1038/s41598-017-03017-z

9. Shibuya, K., Onodera, S. & Hori, M. Toxic wavelength of blue light changes as insects grow. *PLoS One* **13**, e0199266 (2018). doi: 10.1371/journal.pone.0199266

10. Nash, T. R. *et al.* Daily blue-light exposure shortens lifespan and causes brain neurodegeneration in *Drosophila*. *npj Aging Mech. Dis.* **5**, 8 (2019). doi: 10.1038/s41514-019-0038-6

11. Hall, H., Ma, J., Shekhar, S., Leon-Salas, W. D. & Weake, V. M. Blue light induces a neuroprotective gene expression program in *Drosophila* photoreceptors. *BMC Neurosci.* **19**, 43 (2018). doi: 10.1186/s12868-018-0443-y

12. Yang, J. *et al.* Chronic blue light leads to accelerated aging in *Drosophila* by impairing energy metabolism and neurotransmitter levels. *Front. Aging* **3**, 983373 (2022). doi: 10.3389/fragi.2022.983373

13. Song, Y. *et al.* Age-dependent effects of blue light exposure on lifespan, neurodegeneration, and mitochondria physiology in *Drosophila melanogaster*. *npj Aging* **8**, 11 (2022). doi: 10.1038/s41514-022-00092-z

14. Stanhope, S. C. *et al.* Proteome-wide quantitative analysis of redox cysteine availability in the *Drosophila melanogaster* eye reveals oxidation of phototransduction machinery during blue light exposure and age. *Redox Biol.* **63**, 102723 (2023). doi: 10.1016/j.redox.2023.102723

1052
1053   15.   Scheiner, S. M. Selection experiments and the study of phenotypic plasticity. *J Evol.*
1054 *Biol.* **15**, 889–898 (2002). doi: 10.1046/j.1420-9101.2002.00468.x
1055
1056   16.   Martínez, A. A. & Lang, G. I. Identifying targets of selection in laboratory evolution
1057 experiments. *J. Mol. Evol.* **91**, 345–355 (2023). doi: 10.1007/s00239-023-10096-2
1058
1059   17.   Harshman, L. G. & Hoffmann, A. A. Laboratory selection experiments using
1060 *Drosophila*: What do they really tell us? *Trends Ecol. Evol.* **15**, 32-36 (2000). doi:
1061 10.1016/S0169-5347(99)01756-5
1062
1063   18.   Watson, M. J. O. & Hoffmann, A. A. Acclimation, cross-generation effects, and the
1064 response to selection for increased cold resistance in *Drosophila*. *Evolution* **50**, 1182–
1065 1192 (1996). doi: 10.1111/j.1558-5646.1996.tb02359.x
1066
1067   19.   Zhao, R., Xuan, Y., Li, X. & Xi, R. Age-related changes of germline stem cell activity,
1068 niche signaling activity and egg production in Drosophila. *Aging Cell* **7**, 344–354
1069 (2008). doi: 10.1111/j.1474-9726.2008.00379.x
1070
1071   20.   Lindsley, D. L. & Grell, E. H. Genetic Variations of *Drosophila melanogaster*. 2nd edn
1072 431 *Carnegie Institution of Washington Publication* (1968).
1073
1074   21.   Chatterjee, N. & Perrimon, N. What fuels the fly: Energy metabolism in *Drosophila* and
1075 its application to the study of obesity and diabetes. *Sci. Adv.* **7**, eabg4336 (2021). doi:
1076 10.1126/sciadv.abg433
1077
1078   22.   Heier, C. & Kühnlein, R. P. Triacylglycerol metabolism in *Drosophila melanogaster*.
1079 *Genetics* **210**, 1163–1184 (2018). doi: 10.1534/genetics.118.301583
1080
1081   23.   Akhmetova, K., Balasov, M. & Chesnokov, I. *Drosophila* STING protein has a role in
1082 lipid metabolism. *eLife* **10**, e67358 (2021). doi: 10.7554/eLife.67358
1083
1084   24.   Bailey, A. P. *et al.* Antioxidant role for lipid droplets in a stem cell niche of *Drosophila*.
1085 *Cell* **163**, 340–353 (2015). doi: 10.1016/j.cell.2015.09.020
1086
1087   25.   Moulton, M. J. *et al.* Neuronal ROS-induced glial lipid droplet formation is altered by
1088 loss of Alzheimer's disease-associated genes. *Proc. Natl. Acad. Sci. U.A.S.* **118**,
1089 e2112095118 (2021). doi: 10.1073/pnas.2112095118
1090
1091   26.   Bastide, H., Yassin, A., Johanning, E. J. & Pool, J. E. Pigmentation in *Drosophila*
1092 *melanogaster* reaches its maximum in Ethiopia and correlates most strongly with ultra-
1093 violet radiation in sub-Saharan Africa. *BMC Evol. Biol.* **14**, 179 (2014). doi:
1094 10.1186/s12862-014-0179-y
1095
1096   27.   Paiva, P. *et al.* Animal fatty acid synthase: a chemical nanofactory. *Chem. Rev.* **121**,
1097 9502–9553 (2021). doi: 10.1021/acs.chemrev.1c00147
1098
1099   28.   Miao, T. *et al.* Acetyl-CoA-mediated autoacetylation of fatty acid synthase as a
1100 metabolic switch of de novo lipogenesis in *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.*
1101 **119**, e2212220119 (2022). doi: 10.1073/pnas.2212220119

1102
1103 29. Onuma, T. et al. Recognition of commensal bacterial peptidoglycans defines
1104 *Drosophila* gut homeostasis and lifespan. *PLoS Genet.* **19**, e1010709 (2023). doi:
1105 10.1371/journal.pgen.1010709
1106
1107 30. Sekihara, S., Shibata, T., Hyakkendani, M. & Kawabata, S. I. RNA interference
1108 directed against the Transglutaminase gene triggers dysbiosis of gut microbiota in
1109 Drosophila. *J. Biol. Chem.* **291**, 25077–25087 (2016). doi: 10.1074/jbc.M116.761791
1110
1111 31. McMullen, J. G., Bueno, E., Blow, F. & Douglas, A. E. Genome-inferred
1112 correspondence between phylogeny and metabolic traits in the wild *Drosophila* gut
1113 microbiome. *Genome Biol. Evol.* **13**, (2021). doi: 10.1093/gbe/evab127
1114
1115 32. Suyama, R., Cetraro, N., Yew, J. Y. & Kai, T. Microbes control *Drosophila* germline
1116 stem cell increase and egg maturation through hormonal pathways. *Commun. Biol.* **6**,
1117 (2023). doi: 10.1038/s42003-023-05660-x
1118
1119 33. Henry, L. P., Fernandez, M., Webb, A. & Ayroles, J. F. Host evolutionary history and
1120 ecological context modulate the adaptive potential of the microbiome. Preprint
1121 https://doi.org/10.1101/2020.09.21.306779 (2020).
1122
1123 34. Wong, A. C. N. *et al.* The host as the driver of the microbiota in the gut and external
1124 environment of *Drosophila melanogaster*. *Appl. Environ. Microbiol.* **81**, 6232–6240
1125 (2015). doi: 10.1128/AEM.01442-15
1126
1127 35. Pais, I. S., Valente, R. S., Sporniak, M. & Teixeira, L. *Drosophila melanogaster*
1128 establishes a species-specific mutualistic interaction with stable gut-colonizing
1129 bacteria. *PLoS Biol.* **16**, e2005710 (2018). doi: 10.1371/journal.pbio.2005710
1130
1131 36. Staubach, F., Baines, J. F., Künzel, S., Bik, E. M. & Petrov, D. A. Host species and
1132 environmental effects on bacterial communities associated with *Drosophila* in the
1133 laboratory and in the natural environment. *PLoS One* **8**, e70749 (2013). doi:
1134 10.1371/journal.pone.0070749
1135
1136 37. Jugder, B. E., Kamareddine, L. & Watnick, P. I. Microbiota-derived acetate activates
1137 intestinal innate immunity via the Tip60 histone acetyltransferase complex. *Immunity*
1138 **54**, 1683-1697.e3 (2021). doi: 10.1016/j.immuni.2021.05.017
1139
1140 38. Chaston, J. M., Newell, P. D. & Douglas, A. E. Metagenome-wide association of
1141 microbial determinants of host phenotype in *Drosophila melanogaster*. *mBio* **5**,
1142 e01631-14 (2014). doi: 10.1128/mbio.01631-14
1143
1144 39. Newell, P. D. & Douglas, A. E. Interspecies interactions determine the impact of the
1145 gut microbiota on nutrient allocation in *Drosophila melanogaster*. *Appl. Environ.*
1146 *Microbiol.* **80**, 788–796 (2014). doi: 10.1128/AEM.02742-13
1147
1148 40. He, Y. *et al.* Oxidative fermentation of acetic acid bacteria and its products. *Front.*
1149 *Microbiol.* **13**, 879246 (2022). doi: 10.3389/fmicb.2022.879246
1150
1151 41. Shin, S. C. *et al. Drosophila* microbiome modulates host developmental and metabolic

homeostasis via insulin signaling. *Science* **334**, 670–674 (2011). doi: 10.1126/science.1212782

42. Chaston, J. M., Dobson, A. J., Newell, P. D. & Douglas, A. E. Host genetic control of the microbiota mediates the *Drosophila* nutritional phenotype. *Appl. Environ. Microbiol.* **82**, 671–679 (2016). doi: 10.1128/AEM.03301-15

43. Gould, A. L. *et al*. Microbiome interactions shape host fitness. *Proc. Natl. Acad. Sci. U.S.A.* **115**, E11951–E11960 (2018). doi: 10.1073/pnas.1809349115

44. Elgart, M. *et al*. Impact of gut microbiota on the fly's germ line. *Nat. Commun.* **7**, 11280 (2016). doi: 10.1038/ncomms11280

45. Henriques, S. F. *et al*. Metabolic cross-feeding in imbalanced diets allows gut microbes to improve reproduction and alter host behaviour. *Nat. Commun.* **11**, 4236 (2020). doi: 10.1038/s41467-020-18049-9

46. Dodge, R. *et al*. A symbiotic physical niche in *Drosophila melanogaster* regulates stable association of a multi-species gut microbiota. *Nat. Commun.* **14**, 1557 (2023). doi: 10.1038/s41467-023-36942-x

47. Broderick, N. A., Buchon, N. & Lemaitre, B. Microbiota-induced changes in *Drosophila melanogaster* host gene expression and gut morphology. *mBio* **5**, e01117-14 (2014). doi: 10.1128/mBio.01117-14

48. Huang, J. *et al.* Systematic assessment of transcriptomic and metabolic reprogramming by blue light exposure coupled with aging. *PNAS Nexus* **2**, 1–15 (2023). doi: 10.1093/pnasnexus/pgad390

49. Tu, C. P. D. & Akgül, B. *Drosophila* glutathione S-transferases. *Methods Enzymol.* **401**, 204–226 (2005). doi: 10.1016/S0076-6879(05)01013-X

50. Wierzbicki, F.*, Schwarz, F.*, Cannalonga, O. & Kofler, R. Novel quality metrics allow identifying and generating high-quality assemblies of piRNA clusters. *Mol. Ecol. Resour.* **22**, 102–121 (2022). doi: 10.1111/1755-0998.13455

51. Yu, F. X., Zhao, B. & Guan, K. L. Hippo pathway in organ size control, tissue homeostasis, and cancer. *Cell* **163**, 811–828 (2015). doi: 10.1016/j.cell.2015.10.044

52. Li, Q. et al. Ingestion of food particles regulates the mechanosensing Misshapen-Yorkie pathway in *Drosophila* intestinal growth. *Dev. Cell* **45**, 433-449.e6 (2018). doi: 10.1016/j.devcel.2018.04.014

53. Yu, F. X., Meng, Z., Plouffe, S. W. & Guan, K. L. Hippo pathway regulation of gastrointestinal tissues. *Annu. Rev. Physiol.* **77**, 201–227 (2015). doi: 10.1146/annurev-physiol-021014-071733

54. Huang, H. L. *et al. Par-1* regulates tissue growth by influencing Hippo phosphorylation status and Hippo-Salvador association. *PLoS Biol.* **11**, e1001620 (2013). doi:

10.1371/journal.pbio.1001620

55.   Li, Q. et al. The conserved Misshapen-Warts-Yorkie pathway acts in enteroblasts to
regulate intestinal stem cells in *Drosophila*. *Dev. Cell* **31**, 291–304 (2014). doi:
10.1016/j.devcel.2014.09.012

56.   Valanne, S., Wang, JH. & Rämet, M. The *Drosophila* Toll Signaling Pathway. *J.*
*Immunol.* **186**, 649–656 (2011). doi: 10.4049/jimmunol.1002302

57.   Myllymäki, H., Valanne, S. & Rämet, M. The *Drosophila* Imd Signaling Pathway. *J.*
*Immunol.* **192**, 3455–3462 (2014). doi: 10.4049/ jimmunol.1303309

58.   Albert, F. W. & Kruglyak, L. The role of regulatory variation in complex traits and
disease. *Nat. Rev. Genet.* **16**, 197–212 (2015) doi: 10.1038/nrg3891

59.   Massouras, A. *et al.* Genomic variation and its impact on gene expression in
*Drosophila    melanogaster*.    *PLoS    Genet*    **8**,    e1003055    (2012)    doi:
10.1371/journal.pgen.1003055

60.   Moschandrea, C. *et al*. Mitochondrial dysfunction abrogates dietary lipid processing
in enterocytes. *Nature* **625**, 385–392 (2024). doi: 10.1038/s41586-023-06857-0

61.   Byrns, C. N. et al. Senescent glia link mitochondrial dysfunction and lipid accumulation.
*Nature* **630**, 475–483 (2024). doi: 10.1038/s41586-024-07516-8

62.   Song, W., Veenstra, J. A. & Perrimon, N. Control of lipid metabolism by *Tachykinin* in
*Drosophila*. *Cell Rep.* **9**, 40–47 (2014). doi: 10.1016/j.celrep.2014.08.060

63.   Kamareddine, L., Robins, W. P., Berkey, C. D., Mekalanos, J. J. & Watnick, P. I. The
*Drosophila* immune deficiency pathway modulates enteroendocrine function and host
metabolism. *Cell Metab.* **28**, 449-462.e5 (2018). doi: 10.1016/j.cmet.2018.05.026

64.   Brown, J. B. *et al.* An integrated host-microbiome response to atrazine exposure
mediates toxicity in *Drosophila*. *Commun. Biol.* **4**, 1324 (2021). doi: 10.1038/s42003-
021-02847-y

65.   Chongtham, A. *et al.* Gut bacteria regulate the pathogenesis of Huntington's disease
in    *Drosophila*    model.    *Front.    Neurosci.*    **16**,    902205    (2022).    doi:
10.3389/fnins.2022.902205

66.   Tripathi, N. M. *et al.* Draft genome sequence of *Acetobacter persici* DS1, a strain for
high-acidity vinegar fermentation. *Microbiol. Resour. Announc.* **14**, e01119-24 (2025).
doi: 10.1128/mra.01119-24

67.   Henry, L. P., Bruijning, M., Forsberg, S. K. G. & Ayroles, J. F. The microbiome extends
host evolutionary potential. *Nat. Commun.* **12**, 5141 (2021). doi: 10.1038/s41467-021-
25315-x

68.   Henry, L. P. & Ayroles, J. F. Meta-analysis suggests the microbiome responds to

evolve and resequence experiments in *Drosophila melanogaster*. *BMC Microbiol.* **21**,
108 (2021). doi: 10.1186/s12866-021-02168-4
1252

1253    69.    Bruijning, M., Henry, L. P., Forsberg, S. K. G., Metcalf, C. J. E. & Ayroles, J. F. Natural
1254      selection for imprecise vertical transmission in host–microbiota systems. *Nat. Ecol.*
1255      *Evol.* **6**, 77–87 (2022). doi: 10.1038/s41559-021-01593-y
1256

1257    70.    Liu, L. *et al.* Glial lipid droplets and ROS induced by mitochondrial defects promote
1258      neurodegeneration. *Cell* **160**, 177–190 (2015). doi: 10.1016/j.cell.2014.12.019
1259

1260    71.    Reiff, T. *et al*. Endocrine remodelling of the adult intestine sustains reproduction in
1261      *Drosophila*. *eLife* **4**, e06930 (2015). doi: 10.7554/eLife.06930
1262

1263    72.    Buszczak, M., Lu, X., Segraves, W. A., Chang, T. Y. & Cooley, L. Mutations in the
1264      Midway gene disrupt a *Drosophila* Acyl Coenzyme A: diacylglycerol acyltransferase.
1265      *Genetics* **160**, 1511–1518 (2002). doi: 10.1093/genetics/160.4.1511
1266

1267    73.    Murphy, M. P. How mitochondria produce reactive oxygen species. *Biochem. J.* **417**,
1268      1–13 (2009). doi: 10.1042/BJ20081386
1269

1270    74.    Crotti, E. et al. Acetic acid bacteria, newly emerging symbionts of insects. *Appl.*
1271      *Environ. Microbiol.* **76**, 6963–6970 (2010). doi: 10.1128/AEM.01336-10
1272

1273    75.    Shi, L. & Tu, B. P. Acetyl-CoA and the regulation of metabolism: mechanisms and
1274      consequences. *Curr. Opin. Cell Biol.* **33**, 125–131 (2015). doi:
1275      10.1016/j.ceb.2015.02.003
1276

1277    76.    Li, X. & Karpac, J. A distinct Acyl-CoA binding protein (ACBP6) shapes tissue plasticity
1278      during nutrient adaptation in *Drosophila*. *Nat. Commun.* **14**, 7599 (2023). doi:
1279      10.1038/s41467-023-43362-4
1280

1281    77.    Narita, T., Weinert, B. T. & Choudhary, C. Functions and mechanisms of non-histone
1282      protein acetylation. *Nat. Rev. Mol. Cell Biol.* **20**, 156–174 (2019). doi: 10.1038/s41580-
1283      018-0081-3
1284

1285    78.    Schiffer, J. A. *et al.* Caenorhabditis elegans processes sensory information to choose
1286      between freeloading and self-defense strategies. *eLife* **9**, e56186 (2020). doi:
1287      10.7554/eLife.56186
1288

1289    79.    Sato, Y. *et al.* Insecticide resistance by a host-symbiont reciprocal detoxification. *Nat.*
1290      *Commun.* **12**, 6432 (2021). doi: 10.1038/s41467-021-26649-2
1291

1292    80.    Hoskins *et al*. The release 6 reference sequence of the *Drosophila melanogaster*
1293      genome. *Genome Res.* **25**, 445–458 (2015). doi: 10.1101/gr.185579.114
1294

1295    81.    Wang, Y. *et al.* Variation in mutation, recombination, and transposition rates in
1296      *Drosophila melanogaster* and *Drosophila simulans*. *Genome Res.* **33**, 587–598
1297      (2023). doi: 10.1101/gr.277383.122
1298

1299 82. Fitz-James, M. H. & Cavalli, G. Molecular mechanisms of transgenerational epigenetic
1300 inheritance. *Nat. Rev. Genet.* **23**, 325–341 (2022). doi: 10.1038/s41576-021-00438-5
1301

1302 83. Christensen, C. F., Laurichesse, Q., Loudhaief, R., Colombani, J. & Andersen, D. S.
1303 *Drosophila* activins adapt gut size to food intake and promote regenerative growth.
1304 *Nat. Commun.* **15**, 273 (2024). doi: 10.1038/s41467-023-44553-9
1305

1306 84. Nagai, H. *et al.* Nutrient-driven dedifferentiation of enteroendocrine cells promotes
1307 adaptive intestinal growth in *Drosophila*. *Dev. Cell* **58**, 1764–1781 (2023). e10. doi:
1308 10.1016/j.devcel.2023.08.022
1309

1310 85. Karasov, W. H., Martínez, del Rio, C. & Caviedes-Vidal, E. Ecological physiology of
1311 diet and digestive systems. *Annu. Rev. Physiol.* **73**, 69–93 (2011). doi:
1312 10.1146/annurev-physiol-012110-142152
1313

1314 86. Stojanović, O., Miguel-Aliaga, I. & Trajkovski, M. Intestinal plasticity and metabolism
1315 as regulators of organismal energy homeostasis. *Nat. Metab.* **4**, 1444–1458 (2022).
1316 doi: 10.1038/s42255-022-00679-6
1317

1318 87. Miguel-Aliaga, I., Jasper, H. & Lemaitre, B. Anatomy and physiology of the digestive
1319 tract of *Drosophila melanogaster*. *Genetics* **210**, 357–396 (2018). doi:
1320 10.1534/genetics.118.300224
1321

1322 88. Chevalier, C. *et al.* Gut microbiota orchestrates energy homeostasis during cold. *Cell*
1323 **163**, 1360–1374 (2015). doi: 10.1016/j.cell.2015.11.004
1324

1325 89. Buchon, N. *et al.* Morphological and molecular characterization of adult midgut
1326 compartmentalization in *Drosophila*. *Cell Rep.* **3**, 1725–1738 (2013). doi:
1327 10.1016/j.celrep.2013.04.001
1328

1329 90. MacKay, T. F. C. *et al.* The Drosophila melanogaster Genetic Reference Panel. *Nature*
1330 **482**, 173–178 (2012). doi: 10.1038/nature10811
1331

1332 91. Gargano, J. W., Martin, I., Bhandari, P. & Grotewiel, M. S. Rapid iterative negative
1333 geotaxis (RING): A new method for assessing age-related locomotor decline in
1334 *Drosophila*. *Exp. Gerontol.* **40**, 386395 (2005). doi: 10.1016/j.exger.2005.02.005
1335

1336 92. Torquato, L. S., Mattos, D., Matta, B. P. & Bitner-Mathé, B. C. Cellular basis of
1337 morphological variation and temperature-related plasticity in *Drosophila melanogaster*
1338 strains with divergent wing shapes. *Genetica* **142**, 495–505 (2014). doi:
1339 10.1007/s10709-014-9795-0
1340

1341 93. Lee, J. H., Bassel-Duby, R. & Olson, E. N. Heart- and muscle-derived signaling system
1342 dependent on MED13 and Wingless controls obesity in *Drosophila*. *Proc. Natl. Acad.*
1343 *Sci. U.S.A.* **111**, 9491–9496 (2014). doi: 10.1073/pnas.1409427111
1344

1345 94. Bligh, E. G. & Dyer. W. J. A rapid method of total lipid extraction and purification. *Can.*
1346 *J. Biochem. Physiol.* **37**, 911–7 (1959). doi: 10.1139/o59-099
1347

1348   95.   Guan, X. L. *et al.* Biochemical membrane lipidomics during *Drosophila* development.
1349          *Dev. Cell.* **24**, 98–111 (2013). doi: 10.1016/j.devcel.2012.11.012
1350

1351   96.   Shen, L. R. *et al. Drosophila* lacks C20 and C22 PUFAs. *J. Lipid Res.* **51**, 2985–2992
1352          (2010). doi: 10.1194/jlr.M008524
1353

1354   97.   Wood, J. G. *et al. Sirt4* is a mitochondrial regulator of metabolism and lifespan in
1355          *Drosophila melanogaster. Proc. Natl. Acad. Sci. U.S.A.* **115**, 1564–1569 (2018). doi:
1356          10.1073/pnas.172067311
1357

1358   98.   Ward, J. H. Hierarchical grouping to optimize an objective function. *J. Am. Stat. Assoc.*
1359          **58**, 236–244 (1963). doi: 10.1080/01621459.1963.10500845
1360

1361   99.   Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J. 16S Ribosomal DNA
1362          amplification for phylogenetic study. *J. Bacteriol.* **173**, 697–703 (1991). doi:
1363          10.1128/jb.173.2.697-703.1991
1364

1365   100.  Edwards, U., Rogall, T., Blöcker, H., Emde, M. & Böttger, E. C. Isolation and direct
1366          complete nucleotide determination of entire genes. Characterization of a gene coding
1367          for 16S ribosomal RNA. *Nucleic Acids Res.* **17**, 7843–7853 (1989). doi:
1368          10.1093/nar/17.19.7843
1369

1370   101.  Leftwich, P. T., Clarke, N. V. E., Hutchings, M. I. & Chapman, T. Gut microbiomes and
1371          reproductive isolation in *Drosophila. Proc. Natl. Acad. Sci. U.S.A.* **114**, 12767–12772
1372          (2017). doi: 10.1073/pnas.1708345114
1373

1374   102.  Kanno, M., Hiramatsu, S., Kondo, S., Tanimoto, H. & Ichinose, T. Voluntary intake of
1375          psychoactive substances is regulated by the dopamine receptor *Dop1R1* in
1376          *Drosophila. Sci. Rep.* **11**, 3432 (2021). doi: 10.1038/s41598-021-82813-0
1377

1378   103.  Yukphan, P. *et al. Acetobacter garciniae* sp. nov., an acetic acid bacterium isolated
1379          from fermented mangosteen peel in Thailand. *Int. J. Syst. Evol. Microbiol.* **71**, 10
1380          (2021). doi: 10.1099/ijsem.0.005052
1381

1382   104.  Cumsille, A. *et al.* GenoVi, an open-source automated circular genome visualizer for
1383          bacteria and archaea. *PLoS Comput. Biol.* **19**, e1010998 (2023). doi:
1384          10.1371/journal.pcbi.1010998
1385

1386   105.  Auger-Méthé, M. *et al.* A guide to state–space modeling of ecological time series. *Ecol.*
1387          *Monogr.* **91**, e01470 (2021). doi: 10.1002/ecm.1470
1388

1389   106.  Frühwirth-Schnatter, S. & Wagner, H. Stochastic model specification search for
1390          Gaussian and partial non-Gaussian state space models. *J. Econom.* **154**, 85–100
1391          (2010). doi: 10.1016/j.jeconom.2009.07.003
1392

1393   107.  Aeberhard, W. H., Flemming, J. M. & Nielsen, A. Review of state-space models for
1394          fisheries science. *Annu. Rev. Stat. Appl.* **5**, 215–235 (2018). doi: 10.1146/annurev-
1395          statistics-031017-100427
1396

1397   108.  Lee, M. D. How cognitive modeling can benefit from hierarchical Bayesian models. *J.*

*Math. Psychol.* **55**, 1–7 (2011). doi: 10.1016/j.jmp.2010.08.013

109. Kass, R. E. & Raftery, A. E. Bayes factors. *J. Am. Stat. Assoc.* **90**, 773–795 (2012). doi: 10.1080/01621459.1995.10476572

110. Bürkner, PC. brms: An R package for Bayesian multilevel models using Stan. *J. Stat. Softw.* **80**, 1–28 (2017). doi: 10.18637/jss.v080.i01

111. Nott, D. J., Drovandi, C. C., Mengersen, K. & Evans, M. Approximation of Bayesian predictive *p*-values with regression ABC. *Bayesian Anal.* **13**, 59–83 (2018). doi: 10.1214/16-BA1033

112. Cornwallis, C. K. *et al*. Symbioses shape feeding niches and diversification across insects. *Nat. Ecol. Evol.* **7**, 1022–1044 (2023). doi: 10.1038/s41559-023-02058-0

113. Hadfield, J. D. MCMC methods for multi-response generalized linear mixed models: The MCMCglmm R package. *J. Stat. Softw.* **33**, 1–22 (2010). doi: 10.18637/jss.v033.i02

## Acknowledgements

We are grateful to Dr. Takema Fukatu, Dr. Yuto Yoshinari, Dr. Takashi Nishimura, and Dr. Koichiro Tamura for their valuable discussions regarding the study. We also thank Dr. Masaki Iwata, Dr. Shuhei Miyashita, Dr. Chisato Kobayashi, Dr. Katsuya Taniyama, Dr. Takashi Makino, Dr. Yuma Takahashi, Dr. Takahisa Ueno, Dr. Masashi Murakami, Dr. Misako Okumura, Dr. Takahiro Chihara Dr. Toshinori Hayashi, Dr. Takeshi Igawa, Dr. Ritsuko Suyama and Dr. Sota Tanaka for their important suggestions. We would like to thank Dr. Atsuhiko Nagasawa and Dr. Yuki Chiba for teaching us the GC-MS analysis, and Dr. Fu Namai for teaching us microbiome analysis. We also thank Mr. Yusaku Hayashi for instructing us on how to dissect the fat body. Ms. Ayako Abe and Mr. Kento Fukuoka provided technical assistance. We extend our gratitude to Dr. Haruki Kitazawa and Dr. Shoichiro Kurata for lending us their laboratories. We thank the Rhelixa, Inc. for conducting the next-generation sequencing and assisting with the data acquisition, ELSS, Inc for proofreading and editing the English language of this manuscript, Bloomington Drosophila Stock Center for providing the *Drosophila* stocks used in this study, and Japan Collection of Microorganisms RIKEN BioResource Research Center for providing the bacterial strains used in this research. This work was supported by JST SPRING, grant numbers JPMJSP2102 and JPMJSP2114 to Y.T.

## Author information

Authors and Affiliations

**Graduate School of Agricultural Sciences, Tohoku University, Sendai, Japan**

Yuta Takada, Wakako Ikeda-Ohtsubo, and Masatoshi Hori

**Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Sendai, Japan**

1444    Toshiharu Ichinose

1445

1446    **Graduate School of Pharmaceutical Sciences, Tohoku University, Sendai, Japan**

1447    Naoyuki Fuse

1448

1449    **Graduate School of Life Sciences, Tohoku University, Sendai, Japan**

1450    Toshiharu Ichinose, Kokoro Saito, and Hiromu Tanimoto

1451

Contributions

Y.T. and M.H. conceived and designed the study. Y.T. performed all experiments and

analysed all data, wrote the original manuscript, third manuscripts and fourth manuscript.

Y.T. and W.I. designed and performed the infection experiment with gut contents and

bacterial experiments. Y.T., T.I., and N.F. performed whole-genome and RNA sequencing.

Y.T., K.S., T.I., and H.T. designed the genetics and CAFE assays. Y.T. and K.S. performed

the CAFE assay. Y.T., W.I., N.F., and T.I. edited the first manuscript. Y.T. and T.I. edited

the second manuscript and third manuscript. H.T. and M.H. supervised the study. All

authors participated in the discussions and reviewed the manuscript.

1461

1462    **There are no conflicts of interest to declare in relation to this study.**

# Figures

1463

1464

1465

**Fig 1: Experimental design of laboratory selection using blue light (BL)**

**toxicity.**

**a**, Overview of the ancestral population and the laboratory selection experiment. We

established two strains from the Canton Special (Canton-S) strain: a selected line

subjected to BL toxicity pressure at the adult stage in each generation, and an unselected

control line without selection pressure. Each strain was maintained with three replicate

lines. **b**, Detail of the selection experiment. Flies were BL-irradiated for 3 days, recovered

for 2 days, and categorized in a climbing assay into those capable of climbing ("low

impact") and those unable to climb ("high impact"). In SL, parents selected from low impact

flies. Flies were allowed 5 days for mating and egg-laying. UCL parents were randomly

chosen from the negative control group and reared under the same conditions as the SL

flies.

1478

1479

**Fig 2: Establishment of *Drosophila* lines with BL tolerance evolved by**

**laboratory selection.**

**a**, Changes in fecundity across generations. UCL represents fecundity under non-irradiated

conditions, while SL represents fecundity after BL irradiation. Fecundity was measured as

the number of offspring per female (pupae). *n* = 3. State space model with Markov Chain

Monte Carlo (MCMC) method, posterior median and Bayesian 95% credible intervals (CI).

**b**, Ovarian development differences after 3 days of BL irradiation followed by 3 days of

recovery.   Number of stage 14 oocytes after a 3-day recovery period. *n* = 9–15. Bayes

factor (BF$_{10}$ ) = 50.74. (Replicate lines 2, generation 25). **c**, Fecundity measured as the

number of offspring (pupae). $n$ = 11–15. $BF_{10}$ > 100, generation 38. **d**, Lifespan under

continuous BL irradiation. Male $n$ = 100. $BF_{10}$ >100. Estimates: UCL 90.62 h (89.38–91.82),

SL 96.35 h (95.16–97.52), $p$MCMC < 0.001. Female $n$ = 92. $BF_{10}$ = 0.72. Estimates: UCL

92.09 h (90.38–93.78), SL 100.39 h (98.74–102.13), $p$MCMC < 0.001, replicate lines 1,

generation 25. Female $n$ = 100. $BF_{10}$ > 100. Estimates: UCL 86.16 h (84.05–88.27), SL

107.18 h (105.09–109.28), $p$MCMC < 0.001, replicate lines 1, generation 73. **e**, Dose-

response curve for females exposed to BL. The lethal dose ($LD_{50}$) was estimated by logistic

regression: $10.35 \times 10^{18}$ photons-m-$^2$-s-$^1$ for UCL and $12.50 \times 10^{18}$ photons-m-$^2$-s-$^1$ for SL.

See Fig. S4 for details of lifespan at $5 \times 10^{18}$ photons-m-$^2$-s-$^1$ and $15 \times 10^{18}$ photons-m-$^2$-s-$^1$.

Statistical analyses for panels **b**–**d** used a Hierarchical Bayesian Model (HBM) with MCMC

methods, posterior median, and Bayesian 95% CI. In **b** and **c**, different letters indicate

significant differences at $p$MCMC < 0.05. See Data S2 and Data S3 for details of all results,

including statistical models.

1502

1503

**Fig 3: Selected line (SL) flies characterised by enlarged abdomen and heavier body weight in both sexes.**

**a,** Phenotype of adult virgin male and virgin female aged 6 days. The zoomed view of the abdomen is brightened with maximum exposure. **b**, Body weight, $n$ = 90, $BF_{10}$ >100, generation 39. Statistical analysis for panel **b** was performed using a HBM with MCMC methods, posterior median, and Bayesian 95% CI. Asterisks indicate significance levels: *** $p$MCMC < 0.001. See Data S4 for details of all results, including statistical models.

1511

1512

1513

**Fig 4: SL flies exhibited obese phenotype, lipid accumulation, and oxidative**

**stress resistance.**

**a**, Phenotypic measurements of virgin adult females (aged 4-12 days) under non-irradiated

conditions (**b-h**). Age matching was performed between strains at the time of

measurement. See Fig. S5 and Fig. S6 for details of wing, ovary and abdomen spectrum.

**b**, Fluorescence imaging of the abdominal fat body and lipid droplets (lipid droplets: Nile

red; nuclei: NucBlue). **b'**, Mean fluorescence intensity from Nile red staining, $n$ = 30, $BF_{10}$

>100, generation 35. **c**, Midgut length, $n$ = 15, $BF_{10}$ >100, generation 45. **d**, Food

consumption, $n$ = 12–13, $BF_{10}$ = 1.85, replicate line 1, generation 52. **e**, Gas

chromatography mass spectrometry analyses of fatty acid methyl esters (FAMEs), total

FAMEs, $n$ = 9, $BF_{10}$ = 1.69, generation 30. **f**, Relative triacylglycerol (TAG) level, $n$ = 5, $BF_{10}$

=15.14, replicate line 1, generation 57. **g**, Starvation resistance, Kaplan–Meier survival

curve, $n$ = 40, $BF_{10}$ = 4.49. Estimates: UCL 53.04 h (49.20–56.89), SL 61.91 h (58.09–

65.43), $p$MCMC < 0.01, replicate line 2, generation 43. **h**, Oxidative stress resistance (10

mM paraquat), Kaplan–Meier survival curve, $n$ = 40, $BF_{10}$ >100. Estimates: UCL 53.07 h

(45.90–59.03), SL 77.94 h (71.19–84.08), $p$MCMC < 0.001, replicate line 3, generation 43.

Statistical analyses for panels **b** and **d**–**h** were performed using a HBM with MCMC

methods, posterior median, and Bayesian 95% CI. Asterisks indicate significance levels: *

$p$MCMC < 0.05, *** $p$MCMC < 0.001, n.s = not significant. See Data S4 for details of all

results, including statistical models. FC, fold change.

1534

1535

**Fig 5: High bacterial abundance is the key to BL tolerance and lipid accumulation in SL flies.**

**a,** Relating body weight to bacterial abundance. $n$ = 16, UCL; $y$ = 0.00 × $x$ +1.15, $p$ = 0.54, SL; $y$ = 0.00 × $x$ +1.27, $p$ = 0.06. Bacterial measurements from virgin adult females (aged 6-12 days) under non-irradiated conditions. After measuring body weight, homogenised samples from four flies were diluted 100-fold and cultured on MRS plates for 72 hours to quantify colony-forming units (CFU). The variance represents the 80% confidence interval. See Fig. S7 for microbiota analysis. **a',** Bacterial abundance. $n$ = 16, $BF_{10}$ = 11.17. Asterisks indicate significance: ** $p$MCMC < 0.01; replicate line 1, generation

48. CFU from same dataset as Fig. 5a. **b,** Impact of antibiotic mixture (Abx) treatment and gut content infection on fecundity under BL toxicity. Fecundity measured as the number of offspring (adults), $n = 5$, $BF_{10} > 100$; replicate line 1, generation 47. **c**, Phenotypic effects of Abx treatment (**d**–**f**). Abx was administered continuously from the embryo to adult, resulting in a significant reduction in the adult microbiota. The efficacy of Abx treatment was confirmed by regular plating of adults on MRS plates, where CFU counts were zero in all observations. See Fig. S9a for details. **d**, Body weight, $n = 100$, $BF_{10} > 100$; replicate line 1, generation 55. **e**, Relative TAG level, $n = 10$, $BF_{10} = 1.47$; replicate line 1, generation 55. **f,** Midgut length, $n = 16$, $BF_{10} > 100$; replicate line 1, generation 48. Statistical analyses for panels **a'**–**f** were performed using a HBM with MCMC methods, posterior median, and Bayesian 95% CI. Different letters indicate significant differences at $p$MCMC < 0.05. See Data S6 for details of all results, including statistical models.

1559

**Fig 6: Parental experience of BL irradiation promotes midgut elongation**

**and increased bacterial abundance in progeny.**

**a**, The effect of parental exposure to BL toxicity on offspring traits. Offspring with different

parental experiences were generated. **b**, Bacterial abundance. $n$ = 16. $BF_{10}$ = 8.93; replicate

line 1, generation 53. **c**, Midgut length. $n$ = 16, $BF_{10}$ >100; replicate line 1, generation 54.

Statistical analyses for panels **b** and **c** were performed using HBM with MCMC methods,

posterior median, and Bayesian 95% CI. Different letters indicate significant differences at

$p$MCMC < 0.05. For details, see Data S8 for all results, including statistical models.

1570

**Fig 7: Genetic characteristics unique to SL are enriched in detoxification, antioxidant response, Hippo signaling pathway and Toll and Imd signaling pathway.**

**a**, Manhattan plot of variation for each gene from whole genome resequencing analysis; generation 44. Variation of genes are referenced to Canton-S strain (SRX8038113). SL-specific variation is defined as the difference between the UCL mean variation and the SL variation per replicate line and mean greater than 5%. Each SL replicate line is shown in light grey, the SL mean is shown in dark grey and SL means with more than 5% variation

are highlighted in yellow. Negative values indicate higher variation in Canton-S or UCL. A

value of zero indicates no difference between UCL and SL. **b**, Enrichment analysis of SL-

specific variation (above 5%) using Kyoto Encyclopedia of Genes and Genomes (KEGG).

**c**, Identification of SL-specific common homozygous variants. CDS, coding sequence. **d**,

Enrichment analysis of SL-specific variants using KEGG. **e**, Enrichment analysis of SL-

specific variation (above 5%) and variants using KEGG. For details, see Data S9–Data S1

2.

1586

1587

**Fig 8: The characteristic phenotype of SL flies is explained by the host**

**transcriptome.**

**a**, Transcriptome analysis of the whole body of virgin females under non-irradiated

conditions. Cluster map of non-treated SL-specific differentially expressed genes (DEGs)

with a false discovery rate (FDR)-adjusted *p*-value threshold of <0.05. The total number of

DEGs across all comparisons was 1,242, of which 1,192 (96%) were specific to non-treated

SL (pie chart); replicate line 1, generation 50. **b**, Enrichment analysis of non-treated SL

DEGs using KEGG. **c**, Heatmaps illustrating functional profiles. All genes shown are DEGs

(FDR-adjusted *p*-value threshold of < 0.05). Expression levels are presented as mean log2

fold change for each group, calculated by first calculating $\log_2$ FC for individual samples

relative to the average expression in non-treated UCL and then averaging across groups.

The list of genes includes those known to induce lipid accumulation through genetic

manipulation and related mitochondrial metabolism. Genes with orange symbols represent

either SL-specific variations (above 5%) or common homozygous variants in SL. Circles

indicate mean read counts across all groups. For details, see Data S13.

1603

1604

**Fig 9: Lipid accumulation via genetic manipulation confers BL tolerance.**

**a**, Relative TAG levels, survival rates following BL irradiation, and normalised fecundity. TAG levels are presented as $Log_2$ FC values relative to mean UCL levels. Data are arranged in the order of UCL, SL, the *Tachykinin* (neuropeptide) targeting *Tk-GAL4* driver, the fat body targeting *ppl-GAL4* and *Lsp2-GAL4* drivers and mutant lines. The order is sorted by TAG levels, with higher values on the right. Survival rates after 3 days of BL exposure were measured on a minimum of 40 and a maximum of 120 flies per genotype. Genotypes with less than 10% survival were excluded from the data. The full list is available in Data S4. Fecundity was normalised by calculating the $log_2$ FC in the number of offspring after BL

exposure relative to the unexposed control for each genotype. Negative values indicate a reduction in fecundity due to BL toxicity, while zero indicates no effect. Positive values observed in SL only indicate a beneficial effect. Raw fecundity data, including the number of offspring in both non-irradiated and BL irradiated conditions, are presented in Data S7. Genotype labels and colors indicate the effector, the target gene and the type of manipulation (overexpression; red, knockdown; skyblue or null mutation; royalblue). UCL and SL TAG levels were derived from the same dataset as in Fig. 4f (replicate line 1, generation 57). **b**, Correlation analysis. $n$ = 33, Data using mean values for each genotype in Fig.9a. The variance represents the 95% confidence interval. **c**, Summary of experimental results: Lipid accumulation induced by genetic manipulation confers BL tolerance. For details, see Data S15.

1626

**Fig 10: Phylogenetic relationships, distinct transcriptome and divergent growth patterns of *A. persici* in SL.**

**a**, Phylogenetic tree based on the 16S rRNA gene of *Acetobacter* species identified in the *Drosophila* gut microbiota. *Neokomagataea thailandica* and *Swingsia samuiensi*s were used as outgroups. Several strains of *A. persici* were analysed to improve resolution. The arrows indicate the inferred phylogenetic position of the dominant *A. persici* in our laboratory. As the full-length sequence was not available, it was excluded from the phylogenetic tree. See Fig. S19 for the phylogenetic relationships of the *A. persici* strains. The tree was constructed using maximum likelihood with 1,000 bootstrap replicates. **b**,

Complete genome of *A. persici* based on ASM200656v1. The alphabetical annotations

represent functional categories based on the Database of Cluster of Orthologous Genes. **c**,

Sequencing depth calculated from unmapped *Drosophila* reads mapped to *A. persici.*

Sequencing depth estimated from the length per gene length and total bases mapped to

the gene (mean values for UCL: 13.10, SL: 10.63). Genes corresponding to the same locus

are connected by lines. The central white line represents the mean value. See Fig. S19a for

an overview of the depth distribution. See Fig. 8 for host transcriptome data. **d**, Number of

*A. persici* DEGs across host strains and corresponding volcano plot. Genes with the lowest

*p*-values in the top five are shown in lime, while those involved in acetic acid production are

highlighted in red. Genes highlighted in the main text are shown in bold. **e**, Cluster map of

*A. persici* DEGs (FDR-adjusted *p*-value <0.05) in the SL. **f**, List of genes associated with

the oxidation-reduction process (Gene Ontology: biological process). Only DEGs (FDR-

adjusted *p*-value <0.05) are shown. Bold indicates oxidoreductases or translocases. **g**,

Bacterial growth curve between isolates from host strains. $n$ = 44 (UCL), 37 (SL). The

bacterial growth curve shows the mean OD values with a 95% confidence interval,

estimated by bootstrap resampling ($n$ = 10,000, seed = 4). **g'**, Maximum OD ($K$); $BF_{10}$ =

0.52, **g''**, Midpoint time ($t_0$); $BF_{10}$ = 0.37, **g'''**, growth rate ($r$); $BF_{10}$ = 8.94, Asterisks indicate

significance: ** *p*MCMC < 0.01; *** *p*MCMC < 0.001, n.s = not significant; replicate line 1,

generation 52. **g**–**g'''** represent the same dataset. Statistical analyses for panels **g'**–

**g'''**were performed using HBM with MCMC methods, posterior median, and Bayesian 95%

CI. For details, see Data S16 for all results, including statistical models.

1657

# Supplementary information

## Gut microbiota–mediated lipid accumulation as a driver of evolutionary adaptation to blue light toxicity in *Drosophila*

Yuta Takada[1], Toshiharu Ichinose[2,3], Naoyuki Fuse[4], Kokoro Saito[3], Wakako Ikeda-Ohtsubo[1], Hiromu Tanimoto[3], Masatoshi Hori[1]

[1]Graduate School of Agricultural Sciences, Tohoku University, Japan
[2]Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Japan
[3]Graduate School of Life Sciences, Tohoku University, Japan
[4]Graduate School of Pharmaceutical Sciences, Tohoku University, Japan

Correspondence to Yuta Takada or Masatoshi Hori
yutatakada38@gmail.com
masatoshi.hori.a3@tohoku.ac.jp

1676

**Fig. S1: Sunlight contains a significant amount of blue light.**

Sunlight photon flux density: Sendai, Japan; 38°N, 140°E, clear sky; November 21, 2021;

11:30 AM (approximately 4000 µmol·m⁻²·s⁻¹, blue light (BL): approximately 470

µmol·m⁻²·s⁻¹). 460–470 nm is a highly toxic wavelength range for *Drosophila* adults.

Laboratory selection experiment (female) light condition photon flux density: approximately

175 µmol·m⁻²·s⁻¹, BL: approximately 175 µmol·m⁻²·s⁻¹ ($10.3 \times 10^{18}$ photons·m⁻²·s⁻¹)).

Laboratory selection experiment (male) light condition photon flux density: approximately

120 µmol·m⁻²·s⁻¹, BL: approximately 120 µmol·m⁻²·s⁻¹ ($6.5 \times 10^{18}$ photons·m⁻²·s⁻¹)).

Breeding light condition photon flux density: approximately 3 µmol·m⁻²·s⁻¹, BL:

approximately 0.3 µmol·m⁻²·s⁻¹. For details, see Source data (Light condition and Fig. S1).

1687

**Fig. S2: Dramatic impact on fitness due to BL toxicity**

**a**, Impact of BL toxicity on flies (Canton-S strain) of different ages subjected to a single irradiation for 3 days and a recovery period (RP) of 3 days. $n$ = 50, Bayes factor 10 ($BF_{10}$) > 100. Δ Number is percent change in same age non-irradiation vs following BL irradiation.

**b**, Impact of BL toxicity on UCL flies (generation 34) subjected to a single irradiation for 3 days: ovarian development, the number of stage 14 oocytes after recovery. $n$ = 34–42. $BF_{10}$ > 100. Statistical analyses for panels **a** and **b** were performed using Hierarchical Bayesian Model (HBM) with Markov Chain Monte Carlo (MCMC) methods, posterior median, and Bayesian 95% credible intervals (CI). In **a** and **b**, different letters indicate significant differences at $p$MCMC < 0.05. For details, see Data S1 for all results, including statistical models.

1699

1700

**Fig. S3: SL flies are BL tolerant.**

**a**, Climbing ability and generational changes in male and female flies after exposure to BL irradiation. Climbing ability is represented by a total of 60–100 flies per bar, with bars representing replicate lines 1, 2, and 3. All categories were exposed to BL except the negative control group (top). Negative control group: The UCL flies were kept in darkness for 3 days instead of BL irradiation. Positive control group: A subset of UCL flies in each generation was exposed to BL under the same conditions as SL. **b**, Climbing rate under non-irradiated conditions. The percentage of 10 flies that crossed the 6 cm reference line within 10 seconds of tapping. Each measurement was repeated three times, and the average was taken as the climbing rate. $n$ = 12 (corresponding to 120 flies). Statistical analyses for panels **b** were performed using a HBM with MCMC methods, posterior median, and Bayesian 95% CI. n.s = not significant. For details, see Data S1 for all results,

1713 including statistical models. The source data (Light condition) contains all the light condition

1714 measurements used in the experiment.

1715

1716

**Fig. S4: The 73rd generation SL flies show high survival under BL irradiation.**

Lifespan under continuous BL irradiation. Kaplan–Meier survival curve, $5 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$, Female $n$ = 100, BF$_{10}$ >100. Estimates: UCL 149.25 h (145.09–153.11), SL 198.65 h (194.71–202.77). $15 \times 10^{18}$ photons·m$^{-2}$·s$^{-1}$, Female $n$ = 100, BF$_{10}$ >100. Estimates: UCL 59.12 h (58.00–60.31), SL 75.28 h (74.01–76.57), $p$MCMC < 0.001. Replicate line 1, generation 73. For details, see Data S3 for all results, including statistical models.

1725

**Fig. S5: SL flies have heavier bodies, shorter wings and fewer oocytes.**

**a**, Body weight. $n$ = 90, $BF_{10}$ > 100; generation 28. **b**, Wing length. $n$ = 90, $BF_{10}$ = 91.1; generation 28. **c**, Number of stage 14 oocytes. $n$ = 90, $BF_{10}$ = 2.46; generation 28. **d,** Correlation between traits, based on data from panels **a**–**c**. $n$ = 30. Asterisks indicate significance: *** $p$ < 0.001. Statistical analyses for panels **a**–**c** were performed using a HBM with MCMC methods, posterior median, and Bayesian 95% CI. Asterisks indicate significance levels: *** $p$MCMC < 0.001. See Data S4 for details of all results, including statistical models.

1736

1737 **Fig. S6: No differences in light reflection or transmission due to physical**

1738 **characteristics.**

1739 Reflectance rate and transmittance rate in female abdomens, *n* = 16; replicate line 1,

1740 generation 49.

1741

1742

**Fig. S7: The gut microbiota of SL flies consists of a single *Acetobacter* species.**

**a**, Relative bacterial abundance based on 16S rRNA amplicon sequencing in SL and UCL. The DNA samples used were the same as those used for whole-genome resequencing in Fig 6 (generation 44). **b**, Overview of body weight distribution in the SL flies. replicate line 1, *n* = 240; generation 45. **b'**, Relative bacterial abundance based on 16S rRNA amplicon sequencing in SL replicate line 1, generation 45. For details, see Data S5.

1750

1751

**Fig. S8: Dramatic reduction in gut microbiota and body weight of**

***Drosophila* after BL irradiation.**

**a**, Bacterial abundance and Body weight, $n$ = 13–15, $BF_{10}$ >100; generation 38. Randomly

picked colonies were sequenced for 16S rRNA genes and all bacteria were identified as *A.*

*persici.* Statistical analyses were performed using HBM with MCMC methods, posterior

median, and Bayesian 95% CI. Different letters indicate significant differences at *p*MCMC <

0.05. See Data S6 for details of all results, including statistical models.

1759

1760

**Fig. S9: The unique phenotypes of SL flies depend on gut microbiota.**

**a**, Effect of antibiotic mixture (Abx) treatment on gut microbiota. **b**, Impact of Abx treatment

and infection with gut contents on body weight. Grey bars represent pre-infection, while

white bars indicate post-infection after 4 days. $n$ = 30, $BF_{10}$ > 100; replicate line 1,

generation 47. **c**, Climbing ability following BL irradiation. Each bar represents a total of 30

flies; replicate line 1, generation 47. **d**, Emergence rate, based on data shown in Fig 4B.

$BF_{10}$ = 1.30; replicate line 1, generation 47. Statistical analyses for panels **b** and **d** were

performed using HBM with MCMC methods. Different letters indicate significant differences

at $p$MCMC < 0.05. See Data S7 for details of all results, including statistical models.

1770

1771 **Fig. S10: *Drosophila* body weight and midgut length positively correlate in**

1772 **the presence of gut microbiota.**

1773 Correlation between body weight and midgut length, based on data from Fig. 5f. *n* = 16,

1774 replicate line 1, generation 48. For details, see Data S7.

1775

1776

**Fig. S11: Resistance to oxidative stress in SL flies depends on gut**

**microbiota.**

**a**, Oxidative stress resistance in SL flies treated with 10 mM paraquat. Kaplan–Meier

survival curve, $n$ = 120; replicate line 1, generation 52. **b**, Oxidative stress resistance in SL

and UCL flies treated with 10 mM paraquat. $n$ = 120, $BF_{10}$ = 0.01. Survival: non-treated

UCL 56.36 h (53.29–59.43), Abx-treated UCL 59.57 h (56.33–62.61), non-treated SL 62.48

h (59.42–65.64), Abx-treated SL 53.10 h (50.02–56.22); replicate line 1, generation 52.

Statistical analysis was performed using HBM with MCMC methods. Different letters indicate

significant differences at $p$MCMC < 0.05. See Data S7 for details of all results, including

statistical models.

1787

**Fig. S12: Greater bacterial abundance leads to higher body weight.**

**a**, Schematic of the infection experiment. **a'**, Overview of gut contents or isolated *Acetobacter persici* cocktail from offspring of parents exposed to BL. **b**, Impact of infection with gut contents on body weight. *n* = 30. **c**, Impact of infection with *A. persici* cocktail on body weight. *n* = 23,24. In both panels, grey bars represent pre-infection; white bars indicate post-infection after 4 days. BF$_{10}$ >100; replicate line 1, generation 53. Statistical analyses were performed using HBM with MCMC methods. Different letters indicate significant differences at *p*MCMC < 0.05. See Data S8 for details of all results, including statistical models.

1798

**Fig. S13: Gut contents from BL-exposed UCL parental flies restore fecundity in Abx-treated SL flies.**

**a**, Schematic of the infection experiment. **a'**, Overview of gut contents from offspring of parents exposed to BL. **b**, Fecundity: number of offspring (adults). $n = 5$. $BF_{10} > 100$; replicate line 1, generation 53. Statistical analysis was performed using HBM with MCMC methods. Different letters indicate significant differences at $p$MCMC < 0.05. See Data S8 for details of all results, including statistical models.

1806

1807

**Fig. S14: Parental BL experience increases body weight in progeny.**

Overview of the offspring of parents exposed to BL. Data is based on female body weight. $n$ = 72–90, $BF_{10} > 100$; replicate line 1, generation 46. Statistical analysis was performed using HBM with MCMC methods. Different letters indicate significant differences at $p$MCMC < 0.05. See Data S8 for details of all results, including statistical models.

1813

1814

**Fig. S15: Number of variants potentially present in the strains used for the selection experiment.**

Number of variants in the Canton-S strain (SRX8038113), UCL and SL. Variants in UCL and SL are the total number of variants from replicate lines 1 to 3. The bottom panel shows the number of variants for each replicate line. Colour differences indicate SNPs and indels as well as homozygous and heterozygous variants. The numbers on or within the bars represent the total number of variants.

**Homozygous variants**

1822

1823 **Fig. S16: Number of homozygous variants specific to SL and shared by all**

1824 **three replicate lines, potentially selected under BL toxicity.**

1825 Common and unique variants identified in each replicate line across different strains.

1826 Variants shared within a strain were considered strain-specific, whereas homozygous

1827 variants absent in both UCL and Canton-S were defined as SL-specific.

1828

**Fig. S17: Overview of the host transcriptome profiles.**

**a**, Principal component analysis (PCA) of transcriptome profiles using normalised read

counts. **b**, Volcano plots comparing non-treated SL with each condition. Differentially

expressed genes (DEGs) with a false discovery rate (FDR)-adjusted *p*-value threshold of

<0.05 are shown in magenta. Housekeeping genes were defined based on Ferrari and

Aitken, 2006, *BMC Genomics*. For details, see Data S13.

1835

1836

**Fig. S18: Genes shared between genetic characteristics unique to SL and non-treated SL DEGs are enriched in immune-related pathways.**

**a**, Venn diagram of genetic characteristics unique to SL and differentially expressed genes (DEGs) in non-treated SL, showing a 32.5% overlap. **b**, Enrichment analysis of 387 overlapping genes using the Kyoto Encyclopedia of Genes and Genomes (KEGG), with the top hit identified as the Toll and Imd signaling pathway. **c**, Volcano plots comparing non-treated SL with each condition, highlighting genetic characteristics unique to SL. Approximately 32% overlap was observed across all treatments. **d**, Expression levels of Toll and IMD signaling pathway genes shared between genetic characteristics unique to SL and non-treated SL DEGs. Expression levels are presented as mean $\log_2$ FC for each group, calculated by first calculating $\log_2$ FC for individual samples relative to the average

expression in non-treated UCL and then averaging across groups. Circles indicate mean read counts across all groups. Homozygous variants are indicated in bold. For details, see Data S14.

1852
**Fig. S19: Phylogenetic tree based on the 16S rRNA gene of *A. persici***

Phylogenetic relationship between the dominant OTU ID (397 bp), which comprised 89-98%

of *A. persici* in our laboratory, and other *A. persici* strains. Other bacterial species were

analysed using full-length sequences (1,391-1,498 bp). *Neokomagataea thailandica* and

*Swingsia samuiensi*s were used as outgroups. The tree was constructed using maximum

likelihood with 1,000 bootstrap replicates.

1859

1860

**Fig. S20: Transcriptome profile of the gut microbiota *A. persici***

**a**, Sequence depth based on mapping results. After extracting unmapped reads from the *Drosophila melanogaster* host (GCA_000001215.4), the dominant gut bacteria *A. persici* (ASM200656v1) was used as a reference for mapping. The same genes are connected by lines. The gene with the highest sequencing depth across all samples was *Large ribosomal subunit protein bL12*. See Fig 10 c for magnification and mean values. **b**, PCA based on transcripts per million (TPM). **c**, Enrichment analysis of 225 DEGs (Gene Ontology (GO); biological process), predominantly indicative of metabolic pathways. GO enrichment analysis was then performed on 166 genes mapped by DEGs to the STRING database.

1870

**Fig. S21: Gut microbiota–mediated obesity drives evolutionary adaptation to BL toxicity.**

Evolutionary adaptation to BL toxicity involves selective pressure and the intergenerational transmission of parental experience, which drives midgut elongation, increased bacterial abundance, and optimisation of both host genetic information and microbiota interaction. The shaping of bacterial strain dynamics or bacterial evolution may occur in parallel with host evolution. This process may be associated with increases in bacterial abundance and phenotypic changes, suggesting that both quantitative and qualitative changes in bacterial traits may alter host interactions. These changes can alter gene expression related to lipid metabolism, facilitating the formation of adaptive obesity characterised by lipid accumulation via the gut microbiota. Red arrows indicate an increase, while light blue arrows indicate a suppression.