Blue light toxicity drives the gut microbiota-mediated lipid

accumulation in *Drosophila*

Authors

Yuta Takada[1], Toshiharu Ichinose[2,3], Naoyuki Fuse[4], Kokoro Saito[3], Wakako Ikeda-Ohtsubo[1],

Hiromu Tanimoto[3], Masatoshi Hori[1]

Affiliations

[1]Graduate School of Agricultural Sciences, Tohoku University, Japan

[2]Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Japan

[3]Graduate School of Life Sciences, Tohoku University, Japan

[4]Graduate School of Pharmaceutical Sciences, Tohoku University, Japan

Corresponding to Yuta Takada or Masatoshi Hori

yutatakada38@gmail.com

masatoshi.hori.a3@tohoku.ac.jp

Abstract

Blue light (BL), abundant in sunlight, is highly toxic to many insect species when exposure is excessive. While the physiological mechanisms of BL toxicity are being revealed, the evolutionary responses remain less explored. In this study, using *Drosophila melanogaster* as a model system, we conducted laboratory selection for over 60 generations under excessive BL exposure. The selected line (SL) flies exhibited enhanced BL-tolerance, and also increased body weight and lipid accumulation. Interestingly, we found elongated midgut in the SL flies, and increased microbiota, which was dominated by *Acetobacter persici*. The BL-tolerance and the lipid accumulation were dependent on the increment of gut microbiota. Genomic analysis identified mutations in the Hippo signaling pathway linked to midgut elongation, while transcriptome analysis showed downregulation of *Tachykinin* (*Tk*), a key suppressor of intestinal lipogenesis. Genetically induced lipid accumulation through manipulation of *Tk* or related lipogenic genes was sufficient to confer BL-tolerance. Moreover, our findings indicate that parental BL irradiation, along with accumulated mutations from laboratory selection, played a crucial role in midgut elongation and increased bacterial abundance. These findings reveal evolutionary responses to excessive BL exposure that shape host traits to maximize the benefits provided by gut microbiota.

Introduction

Organisms utilize various environmental cues for survival, yet excessive exposure to these cues can act as stressors. Toxicity resulting from environmental stressors impacts the fitness and evolution of organisms.

Blue light (BL) within the 400-500 nm range, abundant in sunlight, is crucial for insect physiology and ecology, influencing processes such as visual perception, circadian rhythms, and photoreactivation (Fig. 1a)[1–4]. However, excessive BL exposure is lethal across various insect species, with toxicity varying by species and developmental stage[5,6]. Our previous studies have demonstrated that exposure to BL at intensities below 40 % of sunlight's BL content for several days can be lethal to various insects[5–7]. Notably, in some species, BL is more lethal than ultraviolet A (315-400 nm) or shorter-wavelength BL near 400 nm[5–7]. In *Drosophila melanogaster* (hereinafter referred to as *Drosophila*), BL around 465 nm is particularly toxic to both pupae and adults, with males being more sensitive than females[5].

The underlying mechanism of BL toxicity likely involves oxidative stress induced by reactive oxygen species (ROS) generated by mitochondria and other cellular components, leading to organ damage[5,8–12]. While the physiological mechanisms of BL toxicity are increasingly understood, the evolutionary responses to this stressor remain poorly characterized.

Laboratory selection has contributed to understanding the evolutionary responses of organisms to environmental stressors[13,14]. *Drosophila* is well-applied for laboratory selection experiments[15]. Importantly, these experiments often reveal evolutionary responses that were not anticipated at the outset. For example, laboratory selection has demonstrated trade-offs such as increased cold stress resistance accompanied by reduced fecundity and has identified shared proteomic evolutionary changes across various stressors[16,17]. Therefore, *Drosophila*-based laboratory selection is a valuable tool for investigating evolutionary responses to environmental stress.

In this study, we established *Drosophila* lines with BL-tolerance through laboratory selection across consecutive generations to characterize the acquired phenotypes. By characterizing the selected line (SL) flies in terms of morphological, physiological, and genetic traits, we

65 found that the gut-microbiota mediated lipid accumulation is a key adaptive trait for enhanced

66 BL-tolerance.

Result

Drosophila evolved BL-tolerance through laboratory selection

We generated a *Drosophila* line with BL-tolerance using laboratory selection. Specifically, we

selected *Drosophila* adults that exhibited low impact following BL irradiation and conducted

artificial selective breeding to develop BL-tolerant lines (Fig. 1b). After a 3-day BL irradiation

and a 2-day recovery period, we conducted a climbing assay and classified individuals

capable of climbing as low impact. BL irradiation was performed at around 470 nm, the

wavelength most toxic to *Drosophila* adults, with a photon flux density where approximately

50 % of individuals exhibited low impact (Supplementary Fig. 1a)[5]. The un-selected control

line (UCL), maintained without BL irradiation, served as a negative control for comparative

analyses with the SL (Fig. 1b). This procedure was replicated in three independent lines for

each regime.

Generally, BL irradiation negatively affected *Drosophila* fecundity, strongly inhibiting ovarian

development (Fig. 1c). However, in SL flies, improved adaptability was evident after over 20

generations of selection, where fecundity and ovarian development was maintained despite

BL irradiation (Fig. 1-f and Supplementary Fig. 1b). On the contrary, fecundity of SL flies

increased in response to BL irradiation, whereas UCL flies showed reduced fecundity (Fig.

1f). Additionally, the lifespan of SL flies under continuous BL irradiation was longer than that

of UCL flies (Fig. 1h). These results collectively suggest that we have successfully

established a new *Drosophila* line (SL) with BL-tolerance.

SL flies exhibited obese phenotype, lipid accumulation, and oxidative stress resistance

Next, we examined the phenotypic characteristics of SL flies exhibiting BL-tolerance. We

noticed that the SL flies exhibited an obese phenotype characterized by abdomen bloating

(Fig. 2a). This led us to hypothesize that these flies might be undergoing lipid accumulation.

SL flies are characterized by increased body weight and a developed fat body with

numerous lipid droplets (Fig. 2 b-d). They also showed an elongated midgut, and reduction

in wing length (Fig. 2 e and Supplementary Fig. 2 a and b). Unexpectedly, we observed low

number of oocytes in the SL flies despite increased fecundity upon BL irradiation (Fig 1g

and Supplementary Fig. 2c) and strong inter-individual correlation among oocyte number

and body weight in the UCL flies (Supplementary Fig 2 d). These results suggest that,

without BL irradiation, energy sources are preferentially allocated to the fat body and

midgut rather than to oocyte development. We did not find physical traits associated with

light reflection through cellular melanin, despite previous reports suggesting these as

adaptive traits to phototoxicity (Supplementary Fig. 3)[18].

SL flies were rich in major lipids, including fatty acids (fatty acid methyl esters) and

triacylglycerol (TAG), and showed starvation resistance, providing evidence of lipid

accumulation (Fig. 2f-h). The lipid droplets and their constituent TAG have been suggested

to protect against oxidative damage[19–21]. Consistent with these findings, SL flies displayed

higher oxidative stress resistance than UCL flies (Fig. 2i).

Increased bacterial abundance in SL flies

It has been reported that host lipid homeostasis is linked to gut microbiota-derived

acetate[22,23]. Acetate is converted to acetyl-CoA, which serves as a substrate for de novo

lipogenesis[24,25]. Given this insight and considering the elongated midgut in SL flies

accumulating lipids, we investigated gut microbiota profiles in SL and UCL flies. In both flies,

16S rRNA gene sequencing of whole body DNA revealed that a single species of *Acetobacter*

(*Acetobacter persici*), a common gut bacteria in *Drosophila*, dominated (>99 %) the

microbiota (Supplementary Fig. 4a)[26]. Notably, higher bacterial counts were obtained in SL

flies compared to UCL flies, and a positive correlation was observed between body weight

and bacterial abundance, although the microbiota profile remained unchanged (Fig. 3a and

Supplementary Fig. 4b and c). Since all randomly sequenced colonies were identified as *A.*

persici, the difference in bacterial abundance between SL and UCL flies is likely to reflect the

gut colonization level of this bacterium (Supplementary Table 11).

Gut microbiota promotes BL-tolerance and lipid accumulation in SL flies

Following BL irradiation, both bacterial abundance and body weight in SL flies drastically

decreased (Fig. 3b and c), suggesting that these flies might utilize gut microbiota for BL-

tolerance. Consistently, acetic acid bacteria, including *Acetobacter*, which metabolizes

acetate, are known to influence fecundity and TAG levels in *Drosophila*[23,27–34]. We tested the

hypothesis that gut microbiota plays a role in shaping the phenotype of SL flies using an

antibiotic mixture (Abx) (Supplementary Fig. 5a).

We investigated the impact of restoring gut microbiota on BL-tolerance in SL flies by infecting

Abx-treated flies with gut contents. We examined the effects on body weight, climbing ability,

fecundity, and emergence rate. The fecundity and emergence rate of Abx-treated SL flies

were inhibited by BL irradiation (Fig. 3d). Infection of Abx-treated SL flies with gut contents

from SL flies rescued these phenotypes (Supplementary Fig. 4b-d and Fig. 3d). In contrast,

infection with gut contents from UCL flies barely rescued the phenotypes in Abx-treated SL

flies (Supplementary Fig. 5a-c and Fig. 3e). Without irradiation, non-treated SL flies showed

lower fecundity than UCL flies, whereas Abx-treated SL flies displayed fecundity equivalent

to UCL flies (Fig. 3e).

Consistently, Abx treatment significantly impacted the body weight of SL flies, with elevated

TAG levels observed only in non-treated flies (Fig. 3e and f). Non-treated SL flies maintained

higher body weight than non-treated UCL flies despite similar food consumption levels (Fig.

3g and Supplementary Table 3). This result highlights the dependence of SL flies' body

weight and lipid accumulation on their gut microbiota.

Furthermore, the midgut of Abx-treated SL flies was smaller than that of non-treated SL flies

but longer than that of UCL flies, with a positive correlation observed between body weight

and midgut length (Fig. 3h and Supplementary Fig. 6). Enhanced oxidative stress resistance

was observed only in non-treated SL flies (Supplementary Fig. 7a and b). These observations

align with previous reports that gut microbiota is linked to gut morphology and oxidative stress

resistance[26,35,36]. Overall, these results suggest that the characteristic phenotype in SL flies

is shaped by their gut microbiota.

153 Host genome and transcriptomic profiles explain the characteristic

154 phenotype of SL flies

To investigate whether SL flies develop their phenotypes through interactions with gut

microbiota, we conducted comparative genomic analyses of UCL and SL flies. First, we

performed variant discovery using whole genome sequencing to detect common variants

selected in SL flies. Homozygous variants shared among the three replicate lines in each

regime and absent in Canton-S were identified. SL-specific SNPs and indels were detected

in 633 and 1936 genes, respectively. A total of 2224 genes were found to have SL-specific

variants, with the majority located outside the coding sequences (CDS) (Fig. 4a). Enrichment

analysis using the Kyoto Encyclopedia of Genes and Genomes (KEGG) identified the Hippo

signaling pathway as the only significantly enriched pathway (Fig. 4b).

The Hippo signaling pathway regulates intestinal elongation[37,38]. The presence of Hippo

pathway-regulated genes (*Par-1*, *Misshapen*) among CDS variants[38–40], along with the

elongated midgut observed in Abx-treated SL flies, suggests that midgut elongation in SL

flies is potentially influenced by these variants (Supplementary Table 12 and Fig. 3h).

Variants located outside of the CDS are generally thought to influence gene expression, and

gut microbiota may alter host phenotypes through changes in the host transcriptome[41]. We

performed transcriptome analysis on the whole body of non-irradiated UCL and SL flies by

RNA sequencing and compared gene expression profiles between non-treated and Abx-

treated flies. We identified 1242 differentially expressed genes (DEGs), with 1192 (96 %)

specific to non-treated SL flies (Fig. 4c and Supplementary Fig. 8 and Supplementary Table

13). KEGG analysis of the 1192 DEGs revealed gene enrichment in oxidative

phosphorylation, metabolic pathways, Toll and Imd signaling, and the TCA cycle (Fig. 4d).

Suppressed mitochondrial metabolism is known to lead to lipid accumulation in various

animal species[42,43]. Non-treated SL flies showed broad downregulation of oxidative

phosphorylation and the TCA cycle, which are associated with mitochondrial metabolism (Fig.

4e). Several genes, whose downregulation is known to induce lipid accumulation, were also

specifically downregulated in non-treated SL flies (Fig. 4e)[44–46]. Among those, *Tachykinin*

(*Tk)* was also identified in the genome sequencing as variants exhibited specific

downregulation in non-treated SL flies (Fig. 4e). In addition, *Tk* expression is known to be

modulated in response to acetate produced by gut microbiota, thereby regulating lipid

utilization in enterocytes[23,47]. These results suggest that the phenotypes of SL flies are

shaped through interactions with their gut microbiota.

Lipid accumulation via genetic manipulation confers BL-tolerance

To prove the causal-relationship between the lipid accumulation and BL-tolerance, we

examined the BL-tolerance in genetically manipulated flies that are reported to increase the

TAG level [41,42]. We employed a range of mutants and GAL4/UAS-mediated manipulations,

including knockdown of *Tk*, *Adipokinetic hormone receptor* (*AkhR*), and *Lipid storage dloplet-*

1 (*Lsd1*), which were downregulated in the SL flies (Fig. 4). Interestingly, knocking down

these genes resulted in particularly enhanced BL-tolerance, to a comparable level to the SL

flies (Fig. 5a). Furthermore, we also observed enhanced BL-tolerance by other genetic

manipulations, such as knockdown of *Brummer* (*Bmm*), *Phosphatase and tensin homolog*

(*Pten*), and *Stromal interaction molecule* (*Stim*), and overexpression of *Insulin-like receptor*

(*InR*), *Lk6 kinase* (*Lk6*), and *Lipid storage droplet-2* (*Lsd2*). Indeed, we found a significant

positive relationship between fecundity and TAG levels was observed (Fig. 5b), strongly

suggesting that lipid accumulation in general is a trait that confers BL-tolerance.

Parental BL irradiation affects progeny's gut microbiota-associated traits

Finally, we investigated why only the gut microbiota from SL flies could rescue BL-tolerance

in infection experiments, despite the dominance of the same gut bacterial species across

regimes (Supplementary Fig. 4a and Fig. 3e). We hypothesized that parental BL irradiation

might impact gut microbiota-associated traits (e.g., increased bacterial abundance) in the

progeny.

To test this hypothesis, we examined the phenotypes (body weight, BL-tolerance) in Abx-

treated SL flies that had regained gut microbiota through inoculation with gut contents from

flies with different parental experiences. Abx-treated SL flies inoculated with gut contents

from SL flies whose parents had experienced BL irradiation showed the greatest increase in

body weight (Supplementary Fig. 9a). However, flies inoculated with different *A. persici*

cocktails, each standardized in concentration, exhibited similar body weight increases

(Supplementary Fig. 9b). This suggests that quantitative rather than qualitative changes in

gut microbiota due to parental BL irradiation are responsible for the increases in host body

weights.

Furthermore, gut contents from UCL flies whose parents had experienced BL irradiation also

rescued the phenotypes of Abx-treated SL flies, restoring fecundity to the level conferred by

gut contents from SL flies (Fig. 6a). These results indicate that parental BL irradiation affects

the progeny's gut microbiota-associated traits. Consistently, parental BL irradiation increased

the progeny's body weight, bacterial abundance, and midgut length in both UCL and SL flies

(Supplementary Fig. 10 and Fig. 6b, c). Notably, even without parental BL irradiation, the

midgut of SL flies was longer than that of non-irradiated UCL flies, suggesting that this

inherited trait has been conserved during the laboratory selection of SL flies (Fig. 6c). Given

that parental BL irradiation leads to an increase in bacterial abundance in progeny, the

different outcomes observed in gut content inoculation experiments likely reflect variations in

bacterial loads.

Discussion

In this study, we established a BL-tolerant *Drosophila* line through laboratory selection,

demonstrating that excessive BL irradiation over generations drives the evolution of adaptive

traits, particularly gut microbiota-mediated lipid accumulation. This supports the increasingly

recognized role of microbiota in host adaptation to environmental stress, where specific

bacterial populations expand under laboratory-driven selection pressures, contributing to

adaptive host phenotypes[32,48–51]. Our findings further reveal that the elongation of the host

gut in response to BL toxicity facilitates an increase in beneficial bacteria, ultimately

enhancing host fitness under excessive BL exposure.

SL flies exhibited increased lipid accumulation (Fig. 2), and genetic manipulation to increase

lipid conferred BL-tolerance (Fig. 5). Acquisition of BL-tolerance through lipid accumulation

can be explained by its role in protecting against ROS, which is produced upon BL

irradiation[10,11]. Intracellular lipid droplets play a critical role in shielding cells from the oxidative

stress, as reduced TAG levels are associated with increased sensitivity to such stress[19–21,52].

The accumulation of lipid droplets in response to intracellular ROS has been observed in

many species[21,52]. Additionally, lipid droplets are closely linked to fecundity, as higher TAG

levels are associated with improved fecundity, whereas reduced TAG levels can lead to

oocyte degeneration[53,54]. Thus, lipid accumulation not only helps defend against BL toxicity

but also supports the maintenance of fecundity. Furthermore, the transcriptome of SL flies

revealed a broad downregulation of mitochondrial metabolism (Fig. 4e). Mitochondria, while

crucial for energy production, also contribute to ROS generation[55]. Therefore, the inhibition

of mitochondrial metabolism and lipid utilization could be an adaptive strategy to accumulate

lipids while minimizing oxidative stress.

We identified *Acetobacter persici* as the predominant bacterial species, and showed the

critical role of their increased abundance in the acquisition of BL-tolerance. *Acetobacter*

species are well-known symbionts of *Drosophila* and other insects[56], recognized for their

acetate production. Acetyl-CoA, derived from acetate, is essential for lipid utilization[25,57] and

also plays a crucial role in the acetylation of both histone and non-histone proteins, facilitating

adaptive responses through tissue plasticity[23,58,59]. The elevated TAG levels observed in SL

flies may be explained by the increased acetyl-CoA pool resulting from gut microbiota-

derived acetate. Additionally, gut microbiota might act as a barrier against physiological

stress induced by BL toxicity. The significant reduction in gut microbiota following BL

irradiation in SL flies (Fig. 3b) suggests the evolution of mechanisms that utilize bacteria as

a buffer against BL toxicity or as a direct nutritional resource. This finding aligns with previous

research showing that *Acetobacter* enhances the fecundity and stress tolerance of the host

fly[26,32–34].

The increase in bacterial abundance appears to be associated with midgut elongation and

genetic changes, particularly those in the Hippo signaling pathway (Fig. 3h and Fig. 4b). We

also found that the parental irradiation to BL, in addition to the accumulated mutations, is

critical for the midgut elongation and bacterial abundance (Fig. 6b and c). In *Drosophila*, the

midgut undergoes plastic elongation or shortening in response to environmental

conditions[60,61]. This physical expansion of the gut, which increases spatial capacity, offers

several advantages, including more efficient nutrient absorption, increased bacterial

abundance, and maintenance of gut homeostasis[53,62–66]. However, midgut elongation alone,

even after one generation of parental exposure to BL toxicity, is insufficient for BL-tolerance

(Fig. 1d and Fig. 6c). Full BL-tolerance requires both midgut enlargement and specific genetic

changes. Among these genetic factors, the neuropeptide *Tk* and its receptor *TkR99D* are

particularly important (Fig. 5 and Supplementary Table 12), as they may regulate lipid

utilization in response to acetate produced by the gut microbiota[23,46].

Parental exposure to environmental stressors, such as starvation, has been reported to

induce epigenetic inheritance, altering the information passed to progeny[67]. Similarly,

parental BL exposure may have led to epigenetic modifications that influence gut microbiota-

associated traits in the progeny. This proposed mechanism is consistent with evidence that

parental BL irradiation affects the progeny's transcriptome[68].

To conclude, our findings highlight that lipid accumulation mediated by gut microbiota is a

critical adaptive trait for BL-tolerance. The elongation of the midgut, associated with

increased bacterial abundance, emerges as a key factor in this adaptation, may drive by

genetic and epigenetic changes from selective pressures and parental experience of BL

toxicity. These evolutionary responses to excessive BL exposure demonstrate how host traits

evolve to maximize the benefits provided by gut microbiota.

Methods

Fly cultures and diet

Flies were maintained at 23-26 °C on a 16:8 hour light-dark cycle using cold cathode fluorescent lamps as the light source. Detailed lighting conditions are provided in the Supplementary Table 14.

The fly diet consisted of 5 g glucose (Fujifilm Wako Pure Chemicals Co., Osaka, Japan), 6 g dry brewer's yeast (Asahi group holdings Ltd., Tokyo, Japan), 1 g agar (Fujifilm Wako Pure Chemicals Co.), 0.4 mL propionic acid (Fujifilm Wako Pure Chemicals Co.), 0.4 mL 70 % ethanol (Fujifilm Wako Pure Chemicals Co.), 0.01 g butyl p-hydroxybenzoate (Fujifilm Wako Pure Chemicals Co.), and 100 mL distilled water.

Blue light (BL) irradiation

We previously reported that blue light at around 465 nm exhibits high toxicity to adult *Drosophila*, with males being more sensitive than females[5]. Unless otherwise specified, flies were exposed to 462 nm BL for 3 days. Virgin flies aged 1-5 days were collected for experiments. BL irradiation conditions varied across experiments:

Fig. 1c-g, 3d, 6a-c: Photon flux density 10.3×10^{18} photons·m^{-2}·s^{-1} (approximately 175 µmol·m^{-2}·s^{-1}, 20,000 Lux) for females.

Fig. 5: 465 nm BL, photon flux density 9.5×10^{18} photons·m^{-2}·s^{-1} (approximately 19,500 Lux) for females.

Fig. 1c-g, Fig.6b and c: Photon flux density 6.5×10^{18} photons·m^{-2}·s^{-1} (approximately 122 µmol·m^{-2}·s^{-1}, 13,500 Lux) for males.

Fig. 1h: Photon flux density 8×10^{18} photons·m^{-2}·s^{-1} for males and 10×10^{18} photons·m^{-2}·s^{-1} for females.

For details in Supplementary Table 15.

Irradiation was performed in a multi-room incubator (LH-30CCFL-8CT; Nippon Medical & Chemical Instruments Co., Ltd, Osaka, Japan) at 24-26 °C, with LED panels (IS-mini®, ISL-150 × 150 Series; CCS Inc., Kyoto, Japan) installed on the ceiling. Photon flux density was

319   measured using a high-resolution spectrometer HSU-100S (Asahi Spectra Co., Ltd., Tokyo,

320   Japan) with ND 0050 filter (Asahi Spectra Co., Ltd), lux meter LX-90 (AMTAST USA Inc.,

321   Florida, USA) and spectroradiometer MS-730 (EKO Instruments Co., Ltd., Tokyo, Japan).

Laboratory selection for BL-tolerance

Artificial selection

We used the Canton-S strain of *Drosophila*, a common wild-type strain, as the ancestral

population for laboratory selection. Two regimes were established: a selected line (SL)

subjected to BL toxicity every generation, and an un-selected control line (UCL). Each

regime included three replicate lines. The UCL, used as a negative control group, was kept

in darkness for 3 days. For each regime, 60-100 virgin flies aged 1-5 days were placed in

two 60 mm × 90 mm glass Petri dishes for BL irradiation. Because blue light at around 470

nm highly toxicity to adult, SL and UCL flies (positive control group) were exposed to 462

nm BL for 3 days[5].

Climbing assay

Following irradiation, flies were placed in a breeding environment (for details, see "Fly

cultures and diet" in Methods) for a 2-day recovery period. For the data shown in

Supplementary Fig. 1a, the impact of BL toxicity was assessed based on a 4-minute climbing

assay and categorized into five states:

"Low impact" (capable of climbing):

1. Intact; Climbing >2 cm (as defined from the 17th generation onwards)

2. Semi-intact; Capable of climbing, but less efficiently

"High impact" (unable to climb):

3. Phase1; Alive but unable to climb

4. Phase2; Non-ambulatory

5. Phase3; Dead

Artificial selective breeding and evaluation of fecundity

Following the climbing assay, SL individuals classified as "Low impact" were allowed 5 days

for mating and egg-laying. Each 3 mL diet medium vial contained 9-13 pairs, with a total of

4-8 vials (for details, see Supplementary Table 16). After 5 days, the diet medium

containing eggs and larvae was transferred to a larger plastic box (40 mL diet medium, 72

× 72 × 100 mm; Insect Breeding Square Dish, SPL Life Sciences, Gyeonggi-do, Korea) to

ensure sufficient food supply. One box was used for every 4-5 vials. UCL parents were

randomly chosen from the negative control group (for details, see "Laboratory selection for

BL-tolerance -Artificial selection-" in Methods) and reared under the same conditions as the

SL flies. The total number of pupae was counted and divided by the number of parents to

determine the fecundity per female for each generation.

Survival assays

Survival assays were conducted for three different stress conditions: continuous blue light

irradiation, starvation, and oxidative stress. For all assays, virgin female flies aged 5-10

days were used. The number of surviving individuals was manually counted every 2 hours

for all assays.

For the continuous BL irradiation assay in Fig.1h, flies were exposed to 462 nm blue light.

Fifty or forty-two flies were placed in each glass petri dish containing standard diet medium.

For the starvation and oxidative stress assays, ten flies were placed in each vial. For the

starvation assay in Fig.2h, flies were placed in vials containing a medium composed of 1%

(w/v) agar (Fujifilm Wako Pure Chemicals Co.) in distilled water.

For the oxidative stress assay, flies were placed in vials containing a medium prepared with

either 10 mM Paraquat dichloride standard (Fujifilm Wako Pure Chemicals Co.) (Fig. 2i) or

10 mM 1,1'-Dimethyl-4,4'-bipyridinium Dichloride (Tokyo Chemical Industry Co., Ltd.,

Tokyo, Japan) (Supplementary Fig. 7a and b), 5% (w/v) glucose (Fujifilm Wako Pure

Chemicals Co.), and 1% (w/v) agar (Fujifilm Wako Pure Chemicals Co.) in distilled water.

Measurement of fecundity

To measure fecundity (Fig. 1g, Fig. 3d, Fig.5 and Fig. 6a), after 3 days of dark conditions

(non-irradiation) or BL irradiation followed by a 3-day recovery period, 5 surviving females

were randomly chosen and mated with 5 non-irradiated virgin males (aged 1-10 days) in

individual pairs within 5 mL diet medium vials for 4 days of egg-laying. The total number of

pupae and emerged adults were counted to evaluate fecundity and emergence rate (Fig.

1g measured only the total number of pupae).

Fig. 1g utilized non-irradiated males corresponding to the UCL or SL regime, Fig. 5 used

non-irradiated Canton-S males, and Fig. 3d and Fig. 6a employed non-irradiated males

treated with an antibiotic mixture corresponding to the UCL or SL regime.

Measurement of body weight and organ size

Virgin flies aged 4-12 days were collected and anesthetized by chilling on ice for all

measurements. Body weight was measured using an AUW220D balance (Shimadzu,

Kyoto, Japan) for experiments in Fig. 2b, 3c, 3e, Supplementary Fig. 2a, Supplementary

Fig. 5b, Supplementary Fig. 2a, Supplementary Fig. 6, Supplementary Fig. 9a,

Supplementary Fig. 9b and Supplementary Fig. 10.

For experiments in Fig. 1c, Fig. 1e, Supplementary Fig. 2b and Supplementary Fig. 2c,

wings or ovaries were dissected and photographed using a SteREO Discovery.V12

stereomicroscope and an Axiocam 305 color camera (Carl Zeiss, Oberkochen, Germany).

Wing length was measured as the distance between the alula opening and the distal edge

of the 3rd longitudinal vein, using Fiji ImageJ2 (version 2.3/1.54f), following the previous

method[69]. The number of stage 14 oocytes was counted under the microscope. In Fig. 2e,

Fig. 3h and Fig. 6c, midgut lengths of the R1-R5 regions were manually measured from the

midgut images using Fiji ImageJ2[66].

Lipid droplet staining and imaging

We performed dissection and fluorescence microscopy to visualize lipid droplets in the fat

body of the abdominal epithelium from virgin female flies aged 6-12 days. After removing

other tissues from the abdomen, we fixed the abdominal epithelium in PBS containing 4 %

paraformaldehyde (Fujifilm Wako Pure Chemicals Co.) for 20 minutes. After fixation, we

washed the samples twice with PBS (Gibco, Carlsbad, USA) for 15 minutes each. The fixed

abdominal epithelium was stained with Nile Red (Fujifilm Wako Pure Chemicals Co.) at a

concentration of 10 µg/mL overnight according to previous study[70]. Following this, we

washed the samples twice with PBS for 10 minutes each and then stained them with

NucBlue (Thermo Fisher Scientific, Waltham, MA, USA) for 3 minutes, followed by a 5-

minute wash with PBS. All staining procedures were conducted in a light-shielded

environment at room temperature. Imaging was performed using an ECLIPSE Ti2

microscope and a DS-Ri2 camera (Nikon, Tokyo, Japan). The images were analyzed using

the Python package OpenCV (Python version 3.8.3 version 4.5.3.56) to obtain the mean

fluorescent intensity of Nile Red.

Lipid extraction and derivatization of fatty acids

We homogenized 2 virgin female flies aged 6-12 days in 100 µL of PBS (Gibco). Lipids

were extracted using the Bligh and Dyer method[71,72]. To the homogenate, we added 600 µL

of methanol (Fujifilm Wako Pure Chemicals Co.) and chloroform (Fujifilm Wako Pure

Chemicals Co.) in a 2:1 (v/v) ratio, vortexed for 1 minute, and then shook at 1,000 rpm at

4°C for 2 hours. Next, we added 200 µL of chloroform and 250 µL of distilled water to the

mixture, vortexed for 1 minute, and centrifuged at 9,000 rpm for 2 minutes to separate the

phases. The organic lower phase was collected. We then added an additional 400 µL of

chloroform to the remaining mixture, vortexed for 1 minute, and centrifuged at 9,000 rpm for

2 minutes to achieve phase separation. The two organic extracts were pooled, the

container was sealed, and the extracts were dried under vacuum. The dried lipids were

derivatized using the Fatty Acid Methylation Kit (Nacalai Tesque, Inc., Tokyo, Japan) and

the Fatty Acid Methyl Ester Purification Kit (Nacalai Tesque, Inc.) according to the kit

protocols. Fatty acid methyl esters (FAMEs) were collected.

Gas chromatography-tandem mass spectrometry (GC-MS/MS) analysis

The FAMEs were analyzed using a GCMS-QP2010 Ultra (Shimadzu) equipped with a DB-5 ms column (30 m × 0.25 mm ID, 0.25 μm film thickness) (J&W Scientific, Folsom, USA). The carrier gas used was helium at a pressure of 100 kPa. The analysis conditions were as follows:

The column oven was set to an initial temperature of 40°C and held for 5 minutes. The temperature was increased at a rate of 10°C/min to 280°C and maintained at 280°C for 10 minutes. The injection port temperature was set to 220°C, and the analysis was performed in split mode.

Fatty acids were identified using a similarity search against the standard substance Supelco F.A.M.E. Mix C4-C24 (Supelco, Bellafonte, PA, USA) and relevant references[73,74]. The total ion chromatogram area for each compound was measured.

Based on the obtained data, we compared the total amount of FAMEs. The log-transformed average values of each compound in each replicate line were visualized using a cluster map. Clusters were classified by hierarchical clustering using the Ward method, based on Euclidean distances of compound amounts[75].

Measurement of triacylglycerol (TAG)

Relative TAG levels were measured by homogenizing 10 virgin female flies (aged 6-12 days) in 1000 μL of PBS (Fujifilm Wako Pure Chemicals Co.) containing 0.1% Triton X-100 (Nacalai Tesque Inc.). The homogenate was heated at 70°C for 10 minutes and then centrifuged at 20,000 × g for 15 minutes at 4°C. A 10 μL aliquot of the supernatant was used for TAG measurement using the LabAssay Triglyceride kit (Fujifilm Wako Pure Chemicals Co.). For protein normalization, protein concentration was measured using the Bradford method with the Protein Assay CBB Clean Up Kit and Protein Assay CBB Solution (both from Nacalai Tesque Inc.), with the standard calibration curve prepared using Albumin, Bovine Serum, Solution (Nacalai Tesque Inc.). Absorbance was measured using the BioTek Synergy H1 Multimode Reader (Agilent Technologies Inc., Santa Clara, USA).

Relative TAG levels were normalized to protein content, adjusted based on the average

values from UCL (Fig. 2g and Fig. 5) or non-treated UCL (Fig. 3f), and the Log_2 fold change

was computed accordingly.

Measurement of reflectance and transmittance

Virgin female flies aged 6-12 days were collected and anesthetized on ice. The abdominal

reflectance and transmittance were measured by capturing hyperspectral images using a

BX51WI microscope (Olympus, Tokyo, Japan) fitted with an SC-108 hyperspectral imaging

system (EBA Japan Co., Ltd., Tokyo, Japan).

DNA extraction

DNA was extracted from virgin female flies aged 5-8 days using the DNeasy Blood &

Tissue Kit (Qiagen, Hilden, Germany). For both Fig. 4a and Supplementary Fig. 3a, the

same sample of 30 flies was used, with 10 flies specifically used for Supplementary Fig. 3c.

The surface of the flies was first washed in a 1.5 mL tube with 99 % ethanol (Fujifilm Wako

Pure Chemicals Co.) for 30 seconds. Then, 180 µL of PBS (Fujifilm Wako Pure Chemicals

Co.) and beads were added, and the flies were homogenized at 3500 rpm for 30 seconds

using the Micro Smash MS100R (Tomy Seiko Co., Ltd., Tokyo, Japan) while keeping the

mixture cold. Following homogenize, DNA extraction was performed according to kit

protocols.

16S/ITS rRNA amplicon sequencing (V3-V4 hypervariable region)

Quality checks were performed using the BioAnalyzer, followed by library preparation with

the NEBNext® Ultra™ II DNA Library Prep Kit (Illumina, San Diego, CA, USA). Sequencing

was conducted on an Illumina NovaSeq 6000 with the following parameters: paired-end

reads of 250 bp × 2 (PE250) and an output of 50,000 reads per sample. In silico analyses

were performed on a Mac OS (Catalina version 10.15) terminal using Jupyter Notebook

(Python version 3.9 version 6.4.11). Quality assessment was conducted with FastQC

(version 0.11.8). Data analysis was carried out using QIIME 2 (version 2023.7) with the

Silva 138 99% OTUs full-length database (MD5: b8609f23e9b17bd4a1321a8971303310).

The QIIME dada2 denoise-paired parameters were set as follows: --p-n-threads 0, --p-trim-

left-f 30, --p-trim-left-r 30, --p-trunc-len-f 250, and --p-trunc-len-r 250. Operational

Taxonomic Unit sequences were classified to determine the gut microbiota composition.

Bacterial abundance and identification

Body weight (4 virgin adult females) measurements were conducted following either non-

irradiation or BL irradiation. In Fig. 3b, after 3 days of BL irradiation followed by a 3-day

recovery period, 4 surviving females were randomly chosen and virgin female flies, aged 5-

8 days, were surface-sterilized by immersion in 70 % ethanol for 1 minute, followed by

three washes with PBS (Fujifilm Wako Pure Chemicals Co.). The flies were homogenized

in 500 µL of PBS in a 2 mL tube to produce a stock solution, which was then diluted 1:10

with PBS. A 10 µL aliquot of the diluted solution was plated onto MRS agar plates

(comprising 5.5 g MRS broth (Becton, Dickinson and Company, Franklin Lakes, NJ, USA),

1 g agar (Fujifilm Wako Pure Chemicals Co.), and 100 mL distilled water). Colony-forming

units (CFU) were counted after 72 hours of incubation at 30-31 °C. In Fig.3 a, fold change

values are calculated based on the mean of the corresponding UCL replicate line. In Fig.3

b, $\log_2$ fold change values are calculated based on the mean of the corresponding non-

treated UCL replicate line. In Fig. 6a, fold change values are calculated based on the mean

of the corresponding UCL non-experience.

Following CFU measurement, single colonies were isolated for identification. The 16S

rRNA genes of isolated bacterial strains were amplified by PCR using universal bacterial

primers 27F and 1492R. Sequencing was performed on an ABI3500 Sanger sequencing

platform (Thermo Fisher Scientific) using the BigDye™ Terminator v1.1 Cycle Sequencing

Kit (Thermo Fisher Scientific)[76,77].

Antibiotic mixture treatment

The antibiotic mixture (Abx) was prepared according to established protocols[78]. Flies were

reared on a diet containing Abx from the embryo stage to adult. The diet was supplemented

with 100 µg/mL streptomycin sulfate (Fujifilm Wako Pure Chemicals Co.), 50 µg/mL

tetracycline hydrochloride (Fujifilm Wako Pure Chemicals Co.), and 200 µg/mL rifampicin

(Fujifilm Wako Pure Chemicals Co.).

Capillary feeding (CAFE) assay

521   The CAFE assay was conducted using a method adapted from previous research[79]. The

522   feeding solution comprised 5 % sucrose, 2 % yeast extract, 0.25 % nipagin (dissolved in

523   ethanol), 0.005 % red dye, and water. To prevent egg-laying within the capillaries, 4 virgin

524   male flies aged 5-8 days were employed in the experiment. Consumption was measured by

525   allowing the flies to feed for 24 hours.

Infection with gut contents or *Acetobacter persici* cocktails

In Fig. 3d, Supplementary Fig. 4b-d, Fig. 6c, and Supplementary Fig. 9a, 30 virgin female

flies aged 5-8 days were surface-sterilized by immersion in 70 % ethanol for 1 minute,

followed by three washes with PBS (Fujifilm Wako Pure Chemicals Co.). The flies were

homogenized in 500 µL of PBS in a 2 mL tube to create a stock solution, which was then

diluted 1:10 with PBS.

To infect the gut microbiota, 20 µL of the diluted solution was added to the diet, and virgin

females aged 1-4 days were maintained on this diet for 4 days. PBS was used as a control.

Body weight was measured before and after infection. After 3 days of non-irradiation or BL

irradiation followed by a 3-day recovery period, climbing ability and fecundity were

evaluated.

In Supplementary Fig. 9b, *Acetobacter persici* was cultured and isolated on MRS plates for

48 hours. These isolates, along with the standard strain of *A. persici* (JCM 25330), were

cultured in 200 µL of MRS broth (5.5 g MRS broth (Becton, Dickinson and Company) and

100 mL distilled water) in 96-well plates for 72 hours. The bacterial culture pellet, grown for

48 hours in a 96-well plate containing 200 µL of MRS broth per well, was diluted 1:30 with

PBS and inoculated into 20 µL of diet. Autoclaved MRS was also diluted 1:30 with PBS,

and 20 µL of this mixture was used as the control.

Whole-genome resequencing, variant identification, and enrichment

analysis

Quality checks were performed using a BioAnalyzer, followed by library preparation with

the NEBNext® Ultra DNA Library Prep Kit (Illumina). Sequencing was conducted on an

Illumina NovaSeq 6000 with the following parameters: paired-end reads of 150 bp (PE150),

a data output of 10 G bases per sample, and approximately 66.7 million reads per sample.

In silico analyses were performed using macOS (Catalina version 10.15.7) terminal and

Jupyter Notebook (Python version 3.9 version 6.4.11). The quality of the fastq files was

assessed using seqkit stat (version 0.16.1). Quality assessment was further performed with

fastqc (version 0.11.8). Reads were trimmed using trimmomatic (version 0.39) with the

following parameters: LEADING:20, TRAILING:20, SLIDINGWINDOW:4:15, and

MINLEN:20, based on Illumina adapter sequences (CTGTCTCTTATACACATCT). Trimmed

data was re-evaluated with seqkit stat and fastqc to confirm the quality of the trimmed

reads.

Mapping was performed using the reference genome BDGP6.46, INSDC Assembly

GCA_000001215.4 (details provided in the Supplementary Table 12). Indexing and

mapping were conducted using bwa (version 0.7.17), followed by conversion from SAM to

BAM format using samtools (version 1.10). PCR duplicates were marked with GATK

(version 4.2.6.1), and alignment statistics were calculated from the BAM file. Paired-end

reads were merged using samtools, and the merged BAM file was sorted.

Variant calling was executed with sambamba mpileup and varScan pileup2snp, using the

parameters: SNP; --min-coverage 10 --min-var-freq 0.90 --p-value 0.05 (filter; default),

Indel; --min-coverage 10 --min-var-freq 0.90 --p-value 0.10 (filter; --min-reads2 4 --min-var-

freq 0.15 --p-value 0.05). Homozygous variants were identified by extracting rows with

matching Cons and VarAllele values. Common positions across replicates (lines 1, 2, and

3) within each regime were defined to identify regime-specific shared variants. Shared and

unique variants among UCL, SL, and Canton-S (SRX8038113)[80] were analyzed.

Variants were annotated using SnpEff (version 5.0) and visualized using IGV (version

2.8.12). Enrichment analysis was conducted with ShinyGO 0.80 and Kyoto Encyclopedia of

Genes and Genomes (KEGG), based on the unique variants identified (details in the

Supplementary Table 12). Variants classified as having a low impact by SnpEff were

excluded from the analysis list (full list available in the Supplementary Table 18).

RNA extraction

Ten virgin female flies aged 5-8 days were immediately frozen in liquid nitrogen. The frozen

flies were then homogenized in Trizol (Molecular Research Center, Inc, OH, USA).

Following homogenization, chloroform (Fujifilm Wako Pure Chemicals Co.) was added to

the mixture, which was vortexed twice. The sample was then centrifuged at 20,000 × g for

10 minutes at 4°C. The upper aqueous phase was transferred to a new tube, and 200 μL of

isopropyl alcohol was added. After vortexing, the mixture was incubated at room

temperature for 10 minutes. The sample was centrifuged again at 20,000 × g for 10

minutes at 4°C, and the supernatant was discarded. To wash the RNA pellet, 400 μL of

75% ethanol (prepared with DEPC-treated water) was added, followed by vortexing. The

sample was then centrifuged at 20,000 × g for 5 minutes at 4°C. The supernatant was

discarded, and the pellet was centrifuged once more for 1 minute. After allowing the pellet

to air-dry, it was resuspended in DEPC-treated water.

RNA-sequence and enrichment analysis

Quality assessment was initially performed using a BioAnalyzer. Library preparation was

conducted with the NEBNext Ultra II Directional RNA Library Prep Kit (Illumina) and the

NEBNext Poly(A) mRNA Magnetic Isolation Module (New England Biolabs, Ipswich, MA,

USA). Sequencing was executed on an Illumina NovaSeq 6000, with a read length of

PE150 (150 bp × 2 paired-end), generating a data output of 6 G bases per sample and

approximately 40 million reads per sample.

In silico analyses and read trimming followed the protocols described in the " Whole-

genome resequencing, variant identification, and enrichment analysis" section. Poly(A) tail

removal was performed using cutadapt, and mapping was conducted with hisat2 (version

2.2.1) using the BDGP6.46 reference genome (INSDC Assembly GCA_000001215.4) (for

details, see Supplementary Table 19). Gene-wise read counting was accomplished using

HTSeq (version 2.0.4) with the dmel-all-r6.54.gtf annotation file. Read count normalization

was executed with edgeR (version 3.40.2) using the trimmed mean of M values method.

Subsequent principal component analysis and differential expression analysis were carried

out with edgeR and scikit-learn (version 1.0.2). Differentially expressed genes (DEGs) were

identified by comparing non-treated SL with non-treated UCL, Abx-treated SL, or Abx-

treated UCL groups, applying a false discovery rate (FDR)-adjusted p-value threshold of <

0.05. Python libraries such as pandas (version 2.0.3), numpy (version 1.25.2), scipy

(version 1.11.2), and stats were used to filter DEGs with p-values below 0.05. For the

DEGs, $\log_2$ fold change values and Z-scores normalized within the range of -2 to 2 were

calculated. These Z-scores were then used to create a cluster map using the Ward method,

classifying genes and treatments based on their similarity patterns.

Enrichment analysis was performed using ShinyGO 0.80. Gene lists related to specific

biological functions were curated from KEGG (TCA cycle, oxidative phosphorylation, Toll

and Imd signaling pathways) and FlyBase (lipid metabolic processes).

In Fig. 4e, heatmaps illustrating functional profiles were generated using $\log_2$ fold change

values. These values were compared for each gene and each group, normalized against

the mean value of the non-treated UCL group.

Genetics and evaluate of TAG levels and BL tolerance

We collected genetically engineered flies with known elevated TAG levels and reared them

at 23-25°C under a 12-hour light/dark cycle. Using the GAL4/UAS system and null mutants,

we generated flies with RNAi knockdown or overexpression of target genes. The following

strains from the Bloomington Stock Center were utilized (see Supplementary Table 5):

Null mutants: w;*Akh* (BL84448), w;*Bmm* (BL15828), *yw,Pdgy* (BL19990), *yw;Chico*

(BL14337), *w;InR* (BL8267), *w;Ilp 2 3 5* (BL30889), *w;Lk6* (BL87087)

GAL4 drivers: *yw;Lsp2* (BL6357), *w;ppl* (BL58768), *w;Tk* (BL51973)

UAS strains: *yv;UAS-AkhR^{RNAi}* (BL51710), *yv;UAS-Pect^{RNAi}* (BL63710), *yv;UAS-Pten^{RNAi}*

(BL33643), *yv;UAS-Bmm^{RNAi}* (25926), *y sc v sev;UAS-Stim^{RNAi}* (BL52911), *y sc v sev;UAS-*

Lsd-1^{RNAi} (BL65020), *yv;UAS-Tk^{RNAi}* (BL25800), *yw;UAS-Gbb* (BL63059), *yw;UAS-InR*

(BL8263), *w;UAS-Lsd-2* (BL98116), *w;UAS-Lk6* (BL8709), *w,UAS-Rpr1* (BL5823)

We used F1 progenies for the experiments, obtained by crossing GAL4 driver males with

UAS effector females. As controls, we used wild-type Canton-S and w1118 genetically

washed with the Canton-S flies, F1 progenies from crosses of w1118 with UAS effectors,

and F1 progenies from crosses of GAL4 drivers with *y sc v;eGFP^{RNAi}* (BL41555) and *y sc*

v;eGFP^{RNAi} (BL41556) UAS effectors.

Virgin female flies aged 1-5 days were irradiated with either DD or BL for 3 days. Three days

after the recovery period, survival rates were assessed, and 5 female flies were randomly

chosen to mate with 5 Canton-S males for 4 days (for details, see "Measurement of fecundity"

in methods). TAG storage was measured in virgin female flies aged 7-12 days. Relative TAG

levels are presented as $\log_2$ fold changes relative to the mean values across all treatments.

Fecundity is calculated as $\log_2$ fold changes relative to the non-irradiation condition for each

treatment. Data are from SL and UCL (Replicate line 1, Generation 57).

Parental experience of BL irradiation (BL irradiation on the parents)

After 3 days of BL irradiation followed by a 3-day recovery period, flies (male and female)

were randomly chosen as parents. These flies were then allowed to mate and lay eggs for

4 days.

Statistical analysis

Bayesian state space model and estimation

In Fig. 1d, intergenerational fecundity data were analyzed using a Bayesian state space

model implemented with pystan (Python 3.9, version 2.17.0.0). The Stan model included

the following components:

Data block: Defined the number of generations (length of the time series data), a vector of

explanatory variables, and three outcome variables corresponding to each replicate line,

each following a normal distribution.

Parameters block: Estimated a vector of levels with drift components (mu), vectors of

random effects (r1, r2, r3), coefficients of the explanatory variables (b1, b2, b3), the

standard deviation representing the magnitude of the drift component (s_z), and the

standard deviations of the random effects (s_r1, s_r2, s_r3).

Transformed parameters block: Calculated the expected values for each outcome variable

(lambda1, lambda2, lambda3) as the sum of the level with drift component (mu), the

product of the explanatory variables' coefficients (b) and values (ex), and the random

effects (r).

Model block: Assumed that the random effects followed a normal distribution with a mean

of 0 and a given standard deviation. The state equation, mu[i] ~ normal(2 * mu[i-1] - mu[i-2],

s_z), expressed changes in the level with drift components using a second-order local

linear regression. The observation equations, y1[i] ~ normal(lambda1[i], sigma1), y2[i] ~

normal(lambda2[i], sigma2), and y3[i] ~ normal(lambda3[i], sigma3), defined the observed

values as following a normal distribution.

Generated quantities block: Sampled the predicted values for each outcome variable using

the normal distribution rather than the Poisson log-link function.

Random numbers were generated according to the posterior distribution using the Markov

Chain Monte Carlo (MCMC) method with parameters set to iter=10000, chains=4, and

control={'adapt_delta': 0.8, 'max_treedepth': 20}. Convergence was confirmed by ensuring

that the Rhat of the predicted values was <1.10 and by examining the trace plots (for

details, see Supplementary Table 20).

Bayesian hierarchical model (HBM)

Unless otherwise specified, hierarchical uncertainty estimation, accounting for random

effects, was performed using Hierarchical Bayesian Model (HBM). The model specification

and execution utilized the following tools and versions:

Software: R (version 4.0.1), rstan (version 2.32.2, Stan version 2.26.1), brms (version

2.20.4), and bayesplot (version 1.10.0).

Bayesian model comparison: Conducted using Bayes Factor$_{10}$ with the loo function,

computing exp(0.5 * (elpd1 - elpd2)), where elpd1 denotes the model's expected log

pointwise predictive density (elpd_loo) and elpd2 denotes that of the null model.

MCMC sampling: Parameters were set to seed=1, chains=4, iter=20000, warmup=5000,

and thin=4. If convergence was not attained, adjustments were made to iter=25000 and

warmup=10000, ensuring Rhat values remained below 1.10.

Prior Distributions: Uniformly set to Gaussian.

Model details: Comprehensive details including the response variable, predictors, random

effects, and estimation results are available in the supplementary table#. Results included

posterior median estimates, Bayesian 95% credible intervals, and approximate p-values

(pMCMC) computed from posterior sampling.

Posterior sampling and pMCMC calculation: To test whether the posterior distribution of the

difference in each treatment was above or below 0, we followed references[81–84]. Posterior

samplings of each parameter were stored in diff_samples, and pairwise comparisons of

treatments were made using the formula: pMCMC = 2 * min(mean(diff_samples > 0),

mean(diff_samples < 0)).

Correlation analysis

The Pearson correlation coefficient and confidence intervals were computed using the

pearsonr function from the scipy.stats library.

References

References

1. Spitschan, M., Aguirre, G. K., Brainard, D. H. & Sweeney, A. M. Variation of outdoor

illumination as a function of solar elevation and light pollution. *Sci. Rep.* **6**, 26756

(2016).

DOI: 10.1038/srep26756

2. Song, B. M. & Lee, C. H. Toward a mechanistic understanding of color vision in insects.

Front. Neural Circuits **12**, 16 (2018).

DOI: 10.3389/fncir.2018.00016

3. Numata, H., Miyazaki, Y. & Ikeno, T. Common features in diverse insect clocks.

Zoological Lett. **1**, 10 (2015).

DOI: 10.1186/s40851-014-0003-y

4. Kavakli, I. H., Ozturk, N. & Gul, S. DNA repair by photolyases. *Adv Protein. Chem.*

Struct. Biol. **115**, 1–19 (2019).

DOI: 10.1016/bs.apcsb.2018.10.003

5. Shibuya, K., Onodera, S. & Hori, M. Toxic wavelength of blue light changes as insects

grow. *PLoS One* **13**, e0199266 (2018).

DOI: 10.1371/journal.pone.0199266

6. Hori, M., Shibuya, K., Sato, M. & Saito, Y. Lethal effects of short-wavelength visible

light on insects. *Sci. Rep.* **4**, 7383 (2014).

DOI: 10.1038/srep07383

7. Hori, M. & Suzuki, A. Lethal effect of blue light on strawberry leaf beetle, Galerucella

grisescens (Coleoptera: Chrysomelidae). *Sci. Rep.* **7**, 2694 (2017).

DOI: 10.1038/s41598-017-03017-z

8. Nash, T. R., Chow, E. S., Law, A. D., Fu, S. D., Fuszara, E., Bilska, A., Bebas, P.,

Kretzschmar, D & Giebultowicz, J. M. Daily blue-light exposure shortens lifespan and

causes brain neurodegeneration in *Drosophila*. *npj Aging Mech. Dis.* **5**, 8 (2019).

DOI: 10.1038/s41514-019-0038-6

9. Hall, H., Ma, J., Shekhar, S., Leon-Salas, W. D. & Weake, V. M. Blue light induces a

neuroprotective gene expression program in *Drosophila* photoreceptors. *BMC*
Neurosci. **19**, 43 (2018).

DOI: 10.1186/s12868-018-0443-y

10.   Yang, J., Song, Y., Law, A. D., Rogan, C. J., Shimoda, K., Djukovic, D., Anderson, J.
C., Hendrix, D. A & Giebultowicz, J. M. Chronic blue light leads to accelerated aging
in *Drosophila* by impairing energy metabolism and neurotransmitter levels. *Front.*
Aging **3**, 983373 (2022).

DOI: 10.3389/fragi.2022.983373

11.   Song, Y., Yang, J., Law, A. D., Hendrix, D. A., Kretzschmar, D., Robinson, M &
Giebultowicz, J. M. Age-dependent effects of blue light exposure on lifespan,
neurodegeneration, and mitochondria physiology in *Drosophila melanogaster*. *npj*
Aging **8**, 11 (2022).

DOI: 10.1038/s41514-022-00092-z

12.   Stanhope, S. C., Brandwine-Shemmer, T., Blum, H. R., Doud, E. H., Jannasch, A.,
Mosley, A. L., Minke, B & Weake, V. M. Proteome-wide quantitative analysis of redox
cysteine availability in the *Drosophila melanogaster* eye reveals oxidation of
phototransduction machinery during blue light exposure and age. *Redox Biol.* **63**,
102723 (2023).

DOI: 10.1016/j.redox.2023.102723

13.   Scheiner, S. M. Selection experiments and the study of phenotypic plasticity. *J Evol.*
Biol. **15**, 889–898 (2002).

DOI: 10.1046/j.1420-9101.2002.00468.x

14.   Martínez, A. A. & Lang, G. I. Identifying targets of selection in laboratory evolution
experiments. *J. Mol. Evol.* **91**, 345–355 (2023).

DOI: 10.1007/s00239-023-10096-2

15.   Harshman, L. G. & Hoffmann, A. A. Laboratory selection experiments using
Drosophila: what do they really tell us? *Trends Ecol. Evol.* **15**, 32-36 (2000).

DOI: 10.1016/S0169-5347(99)01756-5

16. Watson, M. J. O. & Hoffmann, A. A. Acclimation, cross-generation effects, and the response to selection for increased cold resistance in *Drosophila*. *Evolution* **50**, 1182-1192 (1996).

DOI: 10.1111/j.1558-5646.1996.tb02359.x

17. Sørensen, J. G., Schou, M. F. & Loeschcke, V. Evolutionary adaptation to environmental stressors: a common response at the proteomic level. *Evolution* **71**, 1627–1642 (2017).

DOI: 10.1111/evo.13243

18. Bastide, H., Yassin, A., Johanning, E. J. & Pool, J. E. Pigmentation in *Drosophila* melanogaster reaches its maximum in Ethiopia and correlates most strongly with ultra-violet radiation in sub-Saharan Africa. *BMC Evol. Biol.* **14**, 179 (2014).

DOI: 10.1186/s12862-014-0179-y

19. Akhmetova, K., Balasov, M. & Chesnokov, I. *Drosophila* STING protein has a role in lipid metabolism. *eLife* **10**, e67358 (2021).

DOI: 10.7554/eLife.67358

20. Bailey, A. P., Koster, G., Guillermier, C., Hirst, E. M. A., MacRae, J. I., Lechene, C. P., Postle, A. D & Gould, A. P. Antioxidant role for lipid droplets in a stem cell niche of *Drosophila*. *Cell* **163**, 340–353 (2015).

DOI: 10.1016/j.cell.2015.09.020

21. Moulton, M. J., Barish, S., Ralhan, I., Chang, J., Goodman, L., Harland, J. G., Marcogliese, P. C., Johansson, J. O., Ioannou, M. S. & Bellen, H. J. Neuronal ROS-induced glial lipid droplet formation is altered by loss of Alzheimer's disease-associated genes. *Proc. Natl. Acad. Sci. UAS* **118**, e2112095118 (2021).

DOI: 10.1073/pnas.2112095118

22. Zhao, S., Jang, C., Liu, J., Uehara, K., Gilbert, M., Izzo, L., Zeng, X., Trefely, S., Fernandez, S., Carrer, A., Miller, K. D., Schug, Z. T., Snyder, N. W., Gade, T. P., Rabinowitz, J. D. & Wellen, K. E. Dietary fructose feeds hepatic lipogenesis via microbiota-derived acetate. *Nature* **579**, 586–591 (2020).

DOI: 10.1038/s41586-020-2101-7

23.   Jugder, B. E., Kamareddine, L. & Watnick, P. I. Microbiota-derived acetate activates

intestinal innate immunity via the Tip60 histone acetyltransferase complex. *Immunity*

54, 1683-1697.e3 (2021).

DOI: 10.1016/j.immuni.2021.05.017

24.   Paiva, P., Medina, F. E., Viegas, M., Ferreira, P., Neves, R. P. P., Sousa, J. P. M.,

Ramos, M. J. & Fernandes, P. A. Animal fatty acid synthase: a chemical nanofactory.

Chem. Rev. **121**, 9502–9553 (2021).

DOI: 10.1021/acs.chemrev.1c00147

25.   Miao, T., Kim, J., kang, P., Fujiwara, H., Hsu, FF & Bai, H. Acetyl-CoA-mediated

autoacetylation of fatty acid synthase as a metabolic switch of de novo lipogenesis in

Drosophila. *Proc. Natl. Acad. Sci. USA* **119**, e2212220119 (2022).

DOI: 10.1073/pnas.2212220119

26.   Onuma, T., Yamauchi, T., Kosakamoto, H., Kadoguchi, H., Kuraishi, T., Murakami, T.,

Mori, H., Miura, M. & Obata, F. Recognition of commensal bacterial peptidoglycans

defines *Drosophila* gut homeostasis and lifespan. *PLoS Genet.* **19**, e1010709 (2023).

DOI: 10.1371/journal.pgen.1010709

27.   Chaston, J. M., Newell, P. D. & Douglas, A. E. Metagenome-wide association of

microbial determinants of host phenotype in *Drosophila melanogaster*. *mBio* **5**,

e01631-14 (2014).

DOI: 10.1128/mbio.01631-14

28.   Newell, P. D. & Douglas, A. E. Interspecies interactions determine the impact of the

gut microbiota on nutrient allocation in *Drosophila melanogaster*. *Appl. Environ.*

Microbiol. **80**, 788–796 (2014).

DOI: 10.1128/AEM.02742-13

29.   He, Y., Xie, Z., Zhang, H., Liebl, W., Toyama, H. & Chen, F. Oxidative fermentation of

acetic acid bacteria and its products. *Front. Microbiol.* **13**, 879246 (2022).

DOI: 10.3389/fmicb.2022.879246

30.   Shin, S. C., Kim, SH., You, H., Kim, B., Kim, A. C., Lee, KA., Yoon, JH., Ryu, JH. &Lee,

WJ. *Drosophila* microbiome modulates host developmental and metabolic

homeostasis via insulin signaling. *Science* **334**, 670–674 (2011).

DOI: 10.1126/science.1212782

31.   Chaston, J. M., Dobson, A. J., Newell, P. D. & Douglas, A. E. Host genetic control of

the microbiota mediates the *Drosophila* nutritional phenotype. *Appl. Environ. Microbiol.*

82, 671–679 (2016).

DOI: 10.1128/AEM.03301-15

32.   Gould, A. L., Zhang, V., Lamberti, L., Jones, E. W., Obadia, B., Korasidis, N.,

Gavryushkin, A., Carlson, J. M., Beerenwinkel, N. & Ludington, W. B. Microbiome

interactions shape host fitness. *Proc. Natl. Acad. Sci. USA* **115**, E11951-E11960

(2018).

DOI: 10.1073/pnas.1809349115

33.   Elgart, M., Stern, S., Salton, O., Gnainsky, Y., Heifetz, Y. & Soen, Y. Impact of gut

microbiota on the fly's germ line. *Nat. Commun.* **7**, 11280 (2016).

DOI: https://doi.org/10.1038/ncomms11280

34.   Henriques, S. F., Dhakan, D. B., Serra, L., Francisco, A. P., Carvalho-Santos, Z.,

Baltazar, C., Elias, A. P., Anjos, M., Zhang, T., Maddocks, O. D. K. & Ribeiro, C.

Metabolic cross-feeding in imbalanced diets allows gut microbes to improve

reproduction and alter host behaviour. *Nat. Commun.* **11**, 4236 (2020).

DOI: 10.1038/s41467-020-18049-9

35.   Dodge, R., Jones, E. W., Zhu, H., Obadia, B., Martinez, D. J., Wang, C., Aranda-Díaz,

A., Aumiller, K., Liu, Z., Voltolini, M., Brodie, E. L., Huang, K. C., Carlson, J. M., Sivak,

D. A., Spradling, A. C. & Ludington, W. B. A symbiotic physical niche in *Drosophila*

melanogaster regulates stable association of a multi-species gut microbiota. *Nat.*

Commun. **14**, 1557 (2023).

DOI: 10.1038/s41467-023-36942-x

36.   Broderick, N. A., Buchon, N. & Lemaitre, B. Microbiota-induced changes in *Drosophila*

melanogaster host gene expression and gut morphology. *mBio* **5**, e01117-14 (2014).

DOI: 10.1128/mBio.01117-14

37. Yu, FX., Zhao, B. & Guan, KL. Hippo pathway in organ size control, tissue

homeostasis, and cancer. *Cell* **163**, 811–828 (2015).

DOI: 10.1016/j.cell.2015.10.044

38. Li, Q., Nirala, N. K., Nie, Y., Chen, HJ., Ostroff, G., Mao, J., Wang, Q., Xu, L. & Ip, Y.

T. Ingestion of food particles regulates the mechanosensing Misshapen-Yorkie

pathway in *Drosophila* intestinal growth. *Dev. Cell* **45**, 433-449.e6 (2018).

DOI: 10.1016/j.devcel.2018.04.014

39. Huang, HL., Wang, S., Yin, MX., Dong, L., Wang, C., Lu, Y., Feng, M., Dai, C., Guo,

X., Li, L., Zhao, B., Zhou, Z., Ji, H., Jiang, J., Zhao, Y., Liu, XY. & Zhang, L. Par-1

regulates tissue growth by influencing Hippo phosphorylation status and Hippo-

Salvador association. *PLoS Biol.* **11**, e1001620 (2013).

DOI: 10.1371/journal.pbio.1001620

40. Li, Q., Li, S., Mana-Capelli, S., Flach, R. J. R., Danai, L. V., Amcheslavsky, A., Nie, Y.,

Kaneko, S., Yao, X., Chen, X., Cotton, J. L., Mao, J., McCollum, D., Jiang, J., Czech,

M. P., Xu, L. & Ip, Y. T. The conserved Misshapen-Warts-Yorkie pathway acts in

enteroblasts to regulate intestinal stem cells in *Drosophila*. *Dev. Cell* **31**, 291–304

(2014).

DOI: 10.1016/j.devcel.2014.09.012

41. Nyholm, L., Koziol, A., Marcos, S., Botnen, A. B., Aizpurua, O., Gopalakrishnan, S.,

Limborg, M. T., Gilbert, M. T. P. & Alberdi, A. Holo-omics: integrated host-microbiota

multi-omics for basic and applied biological research. *iScience* **23**, 101414 (2020).

DOI: 10.1016/j.isci.2020.101414

42. Moschandrea, C., Kondylis, V., Evangelakos, I., Herholz, M., Schneider, F., Schmidt,

C., Yang, M., Ehret, S., Heine, M., Jaeckstein, M. Y., Szczepanowska, K., Schwarzer,

R., Baumann, L., Bock, T., Nikitopoulou, E., Brodesser, S., Krüger, M., Frezza, C.,

Heeren, J., Trifunovic, A. & Pasparakis, M. Mitochondrial dysfunction abrogates

dietary lipid processing in enterocytes. *Nature* **625**, 385–392 (2024).

DOI: 10.1038/s41586-023-06857-0

43.	Byrns, C. N., Perlegos, A. E., Miller, K. N., Jin, Z., Carranza, F. R., Manchandra, P.,

Beveridge, C. H., Randolph, C. E., Chaluvadi, V. S., Zhang, S. L., Srinivasan, A. R.,

Bennett, F. C., Sehgal, A., Adams, P. D., Chopra, G. & Bonini, N. M. Senescent glia

link mitochondrial dysfunction and lipid accumulation. *Nature* **630**, 475–483 (2024).

DOI: 10.1038/s41586-024-07516-8

44.	Chatterjee, N. & Perrimon, N. What fuels the fly: energy metabolism in *Drosophila* and

its application to the study of obesity and diabetes. *Sci. Adv.* **7**, eabg4336 (2021).

DOI: 10.1126/sciadv.abg433

45.	Heier, C. & Kühnlein, R. P. Triacylglycerol metabolism in *Drosophila melanogaster*.

Genetics **210**, 1163–1184 (2018).

DOI: 10.1534/genetics.118.301583

46.	Song, W., Veenstra, J. A. & Perrimon, N. Control of lipid metabolism by Tachykinin in

Drosophila. *Cell Rep.* **9**, 40–47 (2014).

DOI: 10.1016/j.celrep.2014.08.060

47.	Kamareddine, L., Robins, W. P., Berkey, C. D., Mekalanos, J. J. & Watnick, P. I. The

Drosophila immune deficiency pathway modulates enteroendocrine function and host

metabolism. *Cell Metab.* **28**, 449-462.e5 (2018).

DOI: 10.1016/j.cmet.2018.05.026

48.	Henry, L. P., Bruijning, M., Forsberg, S. K. G. & Ayroles, J. F. The microbiome extends

host evolutionary potential. *Nat. Commun.* **12**, 5141 (2021).

DOI: 10.1038/s41467-021-25315-x

49.	Henry, L. P. & Ayroles, J. F. Meta-analysis suggests the microbiome responds to

evolve and resequence experiments in *Drosophila melanogaster*. *BMC Microbiol.* **21**,

108 (2021).

DOI: 10.1186/s12866-021-02168-4

50.	Bruijning, M., Henry, L. P., Forsberg, S. K. G., Metcalf, C. J. E. & Ayroles, J. F. Natural

selection for imprecise vertical transmission in host–microbiota systems. *Nat. Ecol.*

Evol. **6**, 77–87 (2022).

DOI: 10.1038/s41559-021-01593-y

51.   Henry, L. P., Fernandez, M., Webb, A. & Ayroles, J. F. Host evolutionary history and

ecological context modulate the adaptive potential of the microbiome. Preprint at

https://doi.org/10.1101/2020.09.21.306779 (2020).

52.   Liu, L., Zhang, K., Sandoval, H., Yamamoto, S., Jaiswal., M., Sanz, E., Li, Z., Hul, J.,

Graham, B. H., Quintana, A. & Bellen, H. J. Glial lipid droplets and ROS induced by

mitochondrial defects promote neurodegeneration. *Cell* **160**, 177–190 (2015).

DOI: 10.1016/j.cell.2014.12.019

53.   Reiff, T., Jacobson, J., Cognigni, P., Antonello, Z., Ballesta, E., Junn, Tan, K. J., Yew,

J. Y., Dominguez, M. & Miguel-Aliaga, I. Endocrine remodelling of the adult intestine

sustains reproduction in *Drosophila. eLife* **4**, e06930 (2015).

DOI: 10.7554/eLife.06930

54.   Buszczak, M., Lu, X., Segraves, W. A., Chang, T. Y. & Cooley, L. Mutations in the

Midway gene disrupt a *Drosophila* Acyl Coenzyme A: diacylglycerol acyltransferase.

Genetics **160**, 1511-1518 (2002).

DOI: 10.1093/genetics/160.4.1511

55.   Murphy, M. P. How mitochondria produce reactive oxygen species. *Biochem. J.* **417**,

1–13 (2009).

DOI: 10.1042/BJ20081386

56.   Crotti, E., Rizzi, A., Chouaia, B., Ricci, I., Favia, G., Alma, A., Sacchi, L., Bourtzis, K.,

Mandrioli, M., Cherif, A., Bandi, C. & Daffonchio, D. Acetic acid bacteria, newly

emerging symbionts of insects. *Appl. Environ. Microbiol.* **76**, 6963–6970 (2010).

DOI: 10.1128/AEM.01336-10

57.   Shi, L. & Tu, B. P. Acetyl-CoA and the regulation of metabolism: mechanisms and

consequences. *Curr. Opin. Cell Biol.* **33**, 125–131 (2015).

DOI: 10.1016/j.ceb.2015.02.003

58.   Li, X. & Karpac, J. A distinct Acyl-CoA binding protein (ACBP6) shapes tissue plasticity

933   during nutrient adaptation in *Drosophila*. *Nat. Commun.* **14**, 7599 (2023).

934   DOI: 10.1038/s41467-023-43362-4

935   59.   Narita, T., Weinert, B. T. & Choudhary, C. Functions and mechanisms of non-histone

936   protein acetylation. *Nat. Rev. Mol. Cell Biol.* **20**, 156-174 (2019).

937   DOI: 10.1038/s41580-018-0081-3

938   60.   Christensen, C. F., Laurichesse, Q., Loudhaief, R., Colombani, J. & Andersen, D. S.

939   *Drosophila* activins adapt gut size to food intake and promote regenerative growth.

940   *Nat. Commun.* **15**, 273 (2024).

941   DOI: 10.1038/s41467-023-44553-9

942   61.   Nagai, H., Nagai, L. A. E., Tasaki, S., Nakato, R, Umetsu, D., Kuranaga, E., Miura, M.

943   & Nakajima, Y. Nutrient-driven dedifferentiation of enteroendocrine cells promotes

944   adaptive intestinal growth in *Drosophila*. *Dev. Cell* **58**, 1764-1781.e10 (2023).

945   DOI: 10.1016/j.devcel.2023.08.022

946   62.   Karasov, W. H., Martínez Del Rio, C. & Caviedes-Vidal, E. Ecological physiology of

947   diet and digestive systems. *Annu. Rev. Physiol.* **73**, 69–93 (2011).

948   DOI: 10.1146/annurev-physiol-012110-142152

949   63.   Stojanović, O., Miguel-Aliaga, I. & Trajkovski, M. Intestinal plasticity and metabolism

950   as regulators of organismal energy homeostasis. *Nat. Metab.* **4**, 1444–1458 (2022).

951   DOI: 10.1038/s42255-022-00679-6

952   64.   Miguel-Aliaga, I., Jasper, H. & Lemaitre, B. Anatomy and physiology of the digestive

953   tract of *Drosophila melanogaster*. *Genetics* **210**, 357–396 (2018).

954   DOI: 10.1534/genetics.118.300224

955   65.   Chevalier, C., Stojanović, O., Colin, D. J., Suarez-Zamorano, N., Tarallo, V., Veyrat-

956   Durebex, C., Rigo, D., Fabbiano, S., Stevanović, A., Hagemann, S., Montet, X.,

957   Seimbille, Y., Zamboni, N., Hapfelmeier, S. & Trajkovski, M. Gut microbiota

958   orchestrates energy homeostasis during cold. *Cell* **163**, 1360–1374 (2015).

959   DOI: 10.1016/j.cell.2015.11.004

66. Buchon, N., Osman, D., David, F. P. A., Fang, H. Y., Boquete, JP., Deplancke, B. &

Lemaitre, B. Morphological and molecular characterization of adult midgut

compartmentalization in *Drosophila*. *Cell Rep.* **3**, 1725–1738 (2013).

DOI: 10.1016/j.celrep.2013.04.001

67. Fitz-James, M. H. & Cavalli, G. Molecular mechanisms of transgenerational epigenetic

inheritance. *Nat. Rev. Genet.* **23**, 325–341 (2022).

DOI: 10.1038/s41576-021-00438-5

68. Huang, J., Zhou, F., Zhou, H., Zheng, X., Huo, Z., Yang, M., Xu, Z., Liu, R., Wang, L.

& Wang, X. Systematic assessment of transcriptomic and metabolic reprogramming

by blue light exposure coupled with aging. *PNAS Nexus* **2**, 1-15(2023).

DOI: 10.1093/pnasnexus/pgad390

69. Torquato, L. S., Mattos, D., Matta, B. P. & Bitner-Mathé, B. C. Cellular basis of

morphological variation and temperature-related plasticity in *Drosophila melanogaster*

strains with divergent wing shapes. *Genetica* **142**, 495–505 (2014).

DOI: 10.1007/s10709-014-9795-0

70. Lee, J. H., Bassel-Duby, R. & Olson, E. N. Heart- and muscle-derived signaling system

dependent on MED13 and Wingless controls obesity in *Drosophila*. *Proc. Natl. Acad.*

Sci. USA **111**, 9491–9496 (2014).

DOI: 10.1073/pnas.1409427111

71. Bligh. E. G & Dyer. W. J. A rapid method of total lipid extraction and purification. *Can.*

J. Biochem. Physiol. **37**, 911-7 (1959).

DOI: 10.1139/o59-099

72. Guan, X. L., Cestra, G., Shui, G., Kuhrs, A., Schittenhelm, R. B., Hafen, E., van der

Goot, F. G., Robinett, C. C., Gatti, M., Gonzalez-Gaitan, M. & Wenk, M. R. Biochemical

membrane lipidomics during *Drosophila* development. *Dev. Cell* **24**, 98–111 (2013).

DOI: 10.1016/j.devcel.2012.11.012

73. Shen, L. R., Lai, C. Q., Feng, X., Parnell, L. D., Wan, J. B., Wang, J. D., Li, D., Ordovas,

J. M. & Kang, J. X. *Drosophila* lacks C20 and C22 PUFAs. *J. Lipid Res.* **51**, 2985–

2992 (2010).

DOI: 10.1194/jlr.M008524

74. Wood, J. G., Schwer, B., Wickremesinghe, P. C., Hartnett, D. A., Garcia, M., Li, M.,

Verdin, E. & Helfand, S. L. Sirt4 is a mitochondrial regulator of metabolism and lifespan

in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* **115**, 1564–1569 (2018).

DOI: 10.1073/pnas.172067311

75. Ward, J. H. Hierarchical grouping to optimize an objective function. *J. Am. Stat. Assoc.*

58, 236-244 (1963).

DOI: 10.1080/01621459.1963.10500845

76. Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J. 16S Ribosomal DNA

amplification for phylogenetic study. *J. Bacteriol.* **173**, 697-703 (1991).

DOI: 10.1128/jb.173.2.697-703.1991

77. Edwards, U., Rogall, T., Blöcker, H., Emde, M. & Böttger, E. C. Isolation and Direct

Complete Nucleotide Determination of Entire Genes. Characterization of a Gene

Coding for 16S Ribosoma] RNA. *Nucleic Acids Res.* **17**, 7843-7853 (1989).

DOI: 10.1093/nar/17.19.7843

78. Leftwich, P. T., Clarke, N. V. E., I. Hutchings, M. & Chapman, T. Gut microbiomes and

reproductive isolation in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **114**, 12767-12772

(2017).

DOI: 10.1073/pnas.1708345114

79. Kanno, M., Hiramatsu, S., Kondo, S., Tanimoto, H. & Ichinose, T. Voluntary intake of

psychoactive substances is regulated by the dopamine receptor Dop1R1 in

Drosophila. *Sci. Rep*. **11**, 3432 (2021).

DOI: 10.1038/s41598-021-82813-0

80. Wierzbicki, F., Schwarz, F., Cannalonga, O. & Kofler, R. Novel quality metrics allow

identifying and generating high-quality assemblies of piRNA clusters. *Mol. Ecol.*

Resour. **22**, 102–121 (2022).

DOI: 10.1111/1755-0998.13455

81. Bürkner, P. C. brms: An R package for Bayesian multilevel models using Stan. *J. Stat.*

Softw. **80**, 1-28 (2017).

DOI: 10.18637/jss.v080.i01

82. Nott, D. J., Drovandi, C. C., Mengersen, K. & Evans, M. Approximation of Bayesian

predictive *p*-values with regression ABC. *Bayesian Anal.* **13**, 59–83 (2018).

DOI: 10.1214/16-BA1033

83. Cornwallis, C. K., van't Padje, A., Ellers, J., Klein, M., Jackson, R., Kiers, E. T., West,

S. A. & Henry L. M. Symbioses shape feeding niches and diversification across insects.

Nat. Ecol. Evol. **7**, 1022–1044 (2023).

DOI: 10.1038/s41559-023-02058-0

84. Hadfield, J. D. MCMC methods for multi-response generalized linear mixed models:

The MCMCglmm R package. *J. Stat. Softw.* **33**, 1-22 (2010).

DOI: 10.18637/jss.v033.i02

1029

1030

Acknowledgements

We are grateful to Dr. Takema Fukatu, Dr. Yuto Yoshinari, Dr. Takashi Nishimura, and Dr. Koichiro Tamura for their valuable discussions regarding the study. We also thank Dr. Masaki Iwata, Dr. Shuhei Miyashita, Dr. Chisato Kobayashi, Dr. Yuma Takahashi, Dr. Takahisa Ueno, Dr. Masashi Murakami and Dr. Sota Tanaka for their important suggestions. We would like to thank Dr. Atsuhiko Nagasawa and Mr Yuki Chiba for teaching us the GC/MS-MS analysis, and Dr. Fu Namai for teaching us about microbiome analysis. We also thank Mr. Yusaku Hayashi for instructing us on how to dissect the fat body. Ms. Ayako Abe and Mr. Kento Fukuoka provided technical assistance. We extend our gratitude to Dr. Haruki Kitazawa and Dr. Shoichiro Kurata for lending us their laboratories. We thank the Rhelixa, Inc, Bloomington Drosophila Stock Center and Japan Collection of Microorganisms RIKEN BioResource Research Center.

This work was supported by JST SPRING, Grant Number JPMJSP2102.

Author information

Authors and Affiliations

Graduate School of Agricultural Sciences, Tohoku University, Sendai, Japan

Yuta Takada, Wakako Ikeda-Ohtsubo and Masatoshi Hori

Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Sendai, Japan

Toshiharu Ichinose

Graduate School of Pharmaceutical Sciences, Tohoku University, Sendai, Japan

Naoyuki Fuse

Graduate School of Life Sciences, Tohoku University, Sendai, Japan

Toshiharu Ichinose, Kokoro Saito and Hiromu Tanimoto

1059

Contributions

Y.T. and M.H. conceived and designed d the study. Y.T. performed all experiments and

analyzed all data, wrote the original manuscript, edited the first manuscript, and reviewed

and edited the second manuscript. Y.T. and W.I. designed and performed the infection

experiment with gut contents and bacterial experiments. Y.T., T.I., and N.F. performed and

analyzed whole genome sequencing and RNA-sequencing. Y.T., K.S., T.I., and H.T.

designed the genetics and CAFE assay. Y.T. and K.S. performed the CAFE assay. W.I.,

N.F., and T.I. edited the first manuscript. H.T. and M.H. supervised the study. All authors

participated in the discussion and reviewed the final manuscript.

1069

Figures

1072 Fig. 1 | Establishment of *Drosophila* lines with blue light (BL)-tolerance

1073 evolved by laboratory selection

a, Sunlight conditions. Photon flux density: 4,050 µmol·m⁻²·s⁻¹ (BL (400-500 nm): 471

µmol·m⁻²·s⁻¹). 460-470 nm is the most toxic wavelength range for *Drosophila* adults.

b, Experimental design of laboratory selection using BL toxicity. After 3 days of BL

irradiation and a 2-day recovery period, a climbing assay was performed. Flies capable of

climbing were classified as "low impact", while those unable to climb were classified as

"high impact". In the selected line (SL), "low impact" flies were allowed 5 days for mating

and egg-laying. Unselected control line (UCL) parents were randomly chosen from the

negative control group and reared under the same conditions as the SL flies. (for details,

see "Laboratory selection for BL-tolerance" in Methods). **c**, Impact of BL toxicity on ovarian

development. Number of stage 14 oocytes in virgin flies after different recovery periods.

Gray: Non-irradiation; Blue: 3 days of BL irradiation. n = 34-42. BF_{10} > 100. (UCL,

Generation 34). **d**, Changes in fecundity across generations resulting from laboratory

selection. Fecundity measured as the number of offspring per female (pupae). n = 3.

Bayesian state space model with Markov Chain Monte Carlo (MCMC) method, posterior

median and Bayesian 95% Credible Intervals (CI). **e**, Ovarian development differences

after 3 days of BL irradiation followed by 3 days of recovery. **f**, Number of stage 14 oocytes

after a 3-day recovery period. n = 9-15. BF_{10} = 50.74. (Replicate line 2, Generation 25). **g**,

Fecundity measured as the number of offspring (pupae). n = 11-15. BF_{10} = 50.74. Each plot

involved the mating and egg-laying of 5 treated females and 5 untreated males for 4 days.

(Generation 38). **h**, Lifespan under continuous BL irradiation. Top: male n = 100. BF_{10}

>100. Estimates: UCL 90.62 h (89.38 to 91.82), SL 96.35 h (95.16 to 97.52), pMCMC <

0.001. Bottom: female n = 92. BF_{10} = 0.72. Estimates: UCL 92.09 h (90.38 to 93.78), SL

100.39 h (98.74 to 102.13), pMCMC < 0.001. (Replicate line 1, Generation 25).

Statistical analyses for panels **b**-**h** used a Bayesian Hierarchical Model (HBM) with MCMC

methods, posterior median, and Bayesian 95% Credible Intervals (CI). In **c**, **f** and **g**,

significant differences between different letters indicate pMCMC < 0.05 (Supplementary

Table 1).

1101

1102

Fig. 2 | Selected line flies exhibited obese phenotype, lipid accumulation, and oxidative stress resistance.

a, Phenotype of adult virgin females aged 6-8 days: left, UCL; right, SL. **b**, Body weight, $n = 90$, $BF_{10} > 100$, (Generation 39). **c**, Fluorescence imaging of abdominal fat body and lipid droplets (Lipid droplet: Nile red; Nucleus: NucBlue). **d**, Mean fluorescence intensity from Nile red staining, $n = 30$, $BF_{10} > 100$, (Generation 35). **e**, Midgut length, $n = 15$, $BF_{10} > 100$, (Generation 45). **f**, GC/MS analyses of Fatty Acid Methyl Esters (FAMEs), total FAMEs, $n = 9$, $BF_{10} = 1.69$ (Generation 30). **g**, Relative triacylglycerol (TAG) level, $n = 5$, $BF_{10} = 15.14$, (Replicate line 1, Generation 57). **h**, Starvation resistance, Kaplan-Meier survival curve, $n =$

40 , BF_{10} =4.49. Estimates: UCL 53.04 h (49.20 to 56.89), SL 61.91 h (58.09 to 65.43)

(Replicate line 2, Generation 43). **i**, Oxidative stress resistance (10 mM paraquat), Kaplan-

Meier survival curve, n = 40, BF_{10} >100. Estimates: UCL 53.07 h (45.90 to 59.03), SL 77.94

h (71.19 to 84.08), (Replicate line 3, Generation 43). Statistical analyses for panels **b**, **d**, **e**,

f, **g**, **h**, and **i** were performed using a HBM with MCMC methods, posterior median, and

Bayesian 95% CI. Asterisks indicate significance levels: * pMCMC < 0.05, ** pMCMC < 0.01,

*** pMCMC < 0.001. For details, see Supplementary Table 2 for all results, including

statistical models.

1120

1121

Fig. 3 | High bacterial abundance is the key to BL-tolerance and lipid accumulation in SL flies.

a, Bacterial abundance, n = 15, BF_{10} = 9.22. Asterisks indicate significance: ** pMCMC < 0.01. (Generation 38). **b**, Bacterial abundance, n = 13-15, BF_{10} >100. (Generation 38). **c**, Body weight, n = 16, BF_{10} >100. (Generation 38). **d,** Fecundity measured as the number of offspring (adults), n = 5, BF_{10} >100. (Replicate line 1, Generation 47) **e**, Body weight, n = 100, BF_{10} >100. (Replicate line 1, Generation 55). **f**, Relative TAG level, n = 10, BF10 = 1.47. (Replicate line 1, Generation 55). **g**, Food consumption, n = 12-13, BF_{10} >100. (Replicate line 1, Generation 52). **h,** Midgut length, n = 16, BF_{10} >100. (Replicate line 1, Generation 48).

1131   Statistical analyses for panels **b-h** were performed using HBM with MCMC methods,

1132   posterior median, and Bayesian 95% CI. Significant differences between different letters

1133   indicate pMCMC < 0.05. For details, see Supplementary Table3 for all results, including

1134   statistical models.

1135

1136

Fig. 4 | Host genome and transcriptomic profiles explain the characteristic phenotype of SL flies

a, Identification of SL-specific variants through whole genome resequencing (Generation 44).

b, Enrichment analysis of SL-specific variants using Kyoto Encyclopedia of Genes and Genomes (KEGG) with an FDR-adjusted *p* value threshold of < 0.05. **c**, Cluster map of differentially expressed genes (DEGs) (Replicate line 1, Generation 50). **d**, Enrichment analysis of non-treated SL DEGs using KEGG with an FDR-adjusted *p* value threshold of < 0.05. **e**, Heatmaps illustrating functional profiles. All genes shown are DEGs (FDR < 0.05). Genes mutated in SL are indicated in orange.

1146  For details, see Supplementary Table 4 for all results.

1147

1148

Fig. 5 | Lipid accumulation via genetic manipulation confers BL-tolerance

a, Top: Relative TAG levels. Middle: Survival rates following BL irradiation. Bottom: Normalized fecundity. **b**, Correlation analysis: Left - Survival rate with Relative TAG levels (r = 0.24, p < 0.0001). Right - Number of offspring with Relative TAG levels (r = 0.54, p < 0.0001). For details, see Supplementary Table 5 for all results, including target genes.

1154

Fig. 6 | Parental experience of BL irradiation affects progeny's gut microbiota-associated traits

a, Fecundity; number of offspring (adults). n = 5. BF_{10} >100. (Replicate line 1, Generation 53). **b**, Bacterial abundance. n = 16. BF_{10} = 63.02. (Replicate line 1, Generation 53). **c**, Midgut length. n = 16, BF_{10} >100 (Replicate line 1, Generation 54). **d**, Evolutionary response to blue light toxicity. The selective pressure and the intergenerational repetition of parental experiences can alter genetic information, driving midgut elongation and increasing bacterial abundance. This process facilitates the formation of an adaptive physiological phenotype, characterized by lipid accumulation via the gut microbiota. Statistical analyses for panels **a-**

1165　　**c** were performed using HBM with MCMC methods, posterior median, and Bayesian 95% CI.

1166　　Significant differences between different letters indicate pMCMC < 0.05. For details, see

1167　　Supplementary Table 6 for all results, including statistical models.

1168

Supplementary Figures

Supplementary Fig. 1 | SL flies exhibit BL-tolerance

a, Climbing ability following BL irradiation and generational changes resulting from

laboratory selection. Climbing ability is represented by a total of 60-100 flies per bar, with

bars representing replicate lines 1, 2, and 3. Left: males, right: females The three

categories are: Top - UCL non-irradiation (negative control group), Middle - UCL following

BL irradiation (positive control group), and Bottom - SL following BL irradiation. **b,** Changes

in fecundity across generations resulting from laboratory selection. The number of offspring

per female (pupae) is shown. Data from Fig. 1d were Log_2-transformed relative to the

average UCL value.

1182

Supplementary Fig. 2 | SL flies are characterized by heavier body weight,

shorter wing length, and a lower oocyte count

a, Body weight. $n = 90$, $BF_{10} > 100$ (Generation 28). **b**, Wing length. $n = 90$, $BF_{10} = 91.1$

Generation 28). **c**, Number of stage 14 oocytes. $n = 90$, $BF_{10} = 2.46$ (Generation 28).

d, Correlation between traits, based on data from panels **a**-**c**. Asterisks indicate

significance: *** $p < 0.001$.

Statistical analyses for panels a-c were performed using a HBM with MCMC methods,

posterior median, and Bayesian 95% CI. Asterisks indicate significance levels: *** pMCMC

< 0.001. For details, see Supplementary Table 7 for all results, including statistical models.

1192

1193

Supplementary Fig. 3 | SL flies show no physical differences in response to light

Reflectance rate and transmittance rate in female abdomen: left, reflectance rate; right, transmittance rate, n = 16 (Replicate line 1, Generation 49).

1198

1199

Supplementary Fig. 4 | The gut microbiota of SL flies consists of a single

***Acetobacter* species, with bacterial abundance positively correlated with**

body weight

a, 16S/ITS rRNA amplicon sequencing results. The DNA samples used were the same as

those employed for WGS in Fig.4 a (Generation 44). **b**, Correlation between body weight

and bacterial abundance, based on data from panel Fig. 3a. Data for body weight and

bacterial abundance were normalized to the mean UCL values from corresponding

replicate lines. **c**, 16S/ITS rRNA amplicon sequencing results (SL Replicate line 1

Generation 45).

1209

1210

Supplementary Fig. 5 | The unique phenotypes of SL flies depend on gut

microbiota

a, Effect of antibiotic mixture (Abx) treatment on gut microbiota. **b**, Impact of Abx treatment

and gut contents infection on body weight. Grey bars represent pre-infection, while white

bars indicate post-infection after 4 days. n = 30, BF_{10} > 100 (Replicate line 1, Generation

47). **c**, Climbing ability following BL irradiation. Climbing ability is represented by a total of

30 flies per bar (Replicate line 1, Generation 47). **d**, Emergence rate, based on data from

Fig. 3d. BF_{10} = 1.30 (Replicate line 1, Generation 47).

Statistical analyses for panels **b** and **d** were performed using HBM with MCMC methods.

Significant differences between different letters indicate pMCMC < 0.05. For details, see

Supplementary Table 8 for all results, including statistical models.

1222

1223

1224 **Supplementary Fig. 6 | A positive correlation between midgut length and**

1225 **body weight is observed under microbiota-present conditions**

1226 Correlation between body weight and midgut length, based on data from Fig. 3h.

1227

1228

Supplementary Fig. 7 | Oxidative stress resistance in SL flies is observed

only in the presence of microbiota

a, Oxidative stress resistance in SL flies treated with 10 mM paraquat. Kaplan-Meier

survival curve, n = 120 (Generation 52, Replicate line 1).

b, Oxidative stress resistance in SL and UCL flies treated with 10 mM paraquat. n = 120,

BF_{10} = 0.01. Survival: Estimates: non-treated UCL 56.36 h (53.29 to 59.43), Abx-treated

UCL 59.57 h (56.33 to 62.61), non-treated SL 62.48 h (59.42 to 65.64), Abx-treated SL

53.10 h (50.02 to 56.22). (Generation 52, Replicate line 1).

Statistical analysis was performed using HBM with MCMC methods. Significant differences

between different letters indicate pMCMC < 0.05. For details, see Supplementary Table 9 for

all results, including statistical models.

1240

1241

Supplementary Fig. 8 | Characteristic transcriptome profiles of non-treated

SL flies

Principal component analysis of transcriptome profiles from non-treated SL flies. Count data

were standardized using Log_{10} fold change.

1246

1247

Supplementary Fig. 9 | Greater bacterial abundance leads to higher body weight

a, Impact of gut contents infection on body weight. Grey bars represent pre-infection, while white bars indicate post-infection after 4 days. n = 30, BF_{10} > 100 (Replicate line 1 Generation 53). **b**, Impact of *Acetobacter persici* cocktail infection on body weight. Grey bars represent pre-infection, while white bars indicate post-infection after 4 days. n = 30, BF_{10} >100 (Replicate line 1 Generation 53).

Statistical analyses for panels **b** and **d** were performed using HBM with MCMC methods. Significant differences between different letters indicate pMCMC < 0.05. For details, see

1257   Supplementary Table 10 for all results, including statistical models.

1258

1259

Supplementary Fig. 10 | Parental BL experiences induce increased

progeny's body weight

Female body weight. n = 72-90, BF_{10} > 100. Statistical analysis was performed using HBM

with MCMC methods. Significant differences between groups are indicated by different letters,

with pMCMC < 0.05 (Replicate line 1, Generation 46). For details, see Supplementary Table

11 for all results, including statistical models.

1266