Title: Activation and Evasion of the FEAR Pathway by RNA Viruses

Short title: RNA viruses and the FEAR Pathway

Emily A. Rex[1], Dahee Seo[1], Aaron Embry[1], Rozanne Arulanandam[2,3], Marcus M.
Spinelli[2,3], Jean-Simon Diallo[2,3], and Don B. Gammon[1]*.

[1]Department of Microbiology, University of Texas Southwestern Medical Center, Dallas,
TX, 75390, USA
[2]Center for Innovative Cancer Research, Ottawa Hospital Research Institute, 501 Smyth
Road, Ottawa, Ontario K1H 8L6, Canada.
[3]Department of Biochemistry, Microbiology and Immunology, Faculty of Medicine,
University of Ottawa, 451 Smyth Road, Ottawa, Ontario K1H 8M5, Canada.
*Corresponding author: Don B. Gammon. Contact: don.gammon@utsouthwestern.edu

Abstract

We recently identified the FACT-ETS-1 Antiviral Response (FEAR) pathway as an

interferon-independent innate immune response that restricts DNA virus replication and

is countered by poxvirus-encoded A51R proteins (Rex *et al.*, 2024, *Nature Microbiology*).

The human FEAR pathway is mediated by the FACT complex, consisting of hSpt16 and

SSRP1 subunits, that remodels chromatin to activate expression of the antiviral

transcription factor, ETS-1. To counter this pathway, poxvirus A51R proteins tether

SUMOylated hSpt16 subunits to microtubules to prevent ETS-1 expression. While these

observations indicate a role for the FEAR pathway in DNA virus restriction, it was

unclear if RNA viruses interact with this pathway. Here, we show that RNA viruses are

also restricted by the FEAR pathway, yet encode mechanisms distinct from poxviruses

to counter this response. We show vesicular stomatitis virus (VSV), a rhabdovirus,

utilizes its matrix (M) protein to promote proteasome-dependent degradation of

SUMOylated hSpt16 and to block ETS-1 nuclear import. Strains encoding mutant M

proteins that cannot antagonize the FEAR pathway exhibit replication defects in human

cells that can be rescued by hSpt16 or ETS-1 depletion. Moreover, FACT inhibitor

treatment enhanced the replication of oncolytic VSV strains encoding defective M

proteins in restrictive cancer cells, suggesting FEAR pathway inhibition may improve

oncolytic virotherapy. Strikingly, we provide evidence that the inability of VSV M to degrade SUMOylated Spt16 in lepidopteran insect cells results in abortive infection, suggesting VSV-Spt16 interactions influence virus host range. Lastly, we show that human and murine paramyxovirus target SUMOylated Spt16 proteins for degradation in human and murine cells utilizing a conserved N-terminal motif in their accessory "C" proteins. Collectively, our study illustrates that DNA and RNA viruses have independently evolved diverse mechanisms to antagonize SUMOylated host Spt16 proteins, underscoring the physiological importance of the FEAR pathway to antiviral immunity.

Keywords: FACT complex; ETS-1; FEAR pathway; RNA virus; viral immune evasion

Abbreviations

FACT-ETS-1 Antiviral Response (FEAR); vesicular stomatitis virus (VSV); matrix (M); interferon (IFN); facilitates chromatin transcription (FACT); vaccinia virus (VV); suppressor of Ty 16 homologue (Spt16); human Spt16 (hSpt16); SUMOylated hSpt16 (hSpt16^{SUMO}); structure specific recognition protein 1 (SSRP1); E26 transformation-specific sequence-1 (ETS-1); Ran GTPase activating protein (RanGAP); ribonucleic acid export 1 (Rae1); nucleoporin 98 (Nup98); nonsegmented negative-strand (NNS); *Lymantria dispar* Spt16 (LdSpt16); SUMOylated LdSpt16 (LdSpt16^{SUMO}); firefly luciferase (LUC); Sendai virus (SeV); Human parainfluenza virus-1 (HPIV-1).

Introduction

The activation of transcription-dependent responses during infection is a fundamental aspect of eukaryotic antiviral immunity. In turn, evolutionary pressures on viruses to efficiently replicate in their eukaryotic hosts have driven the acquisition of viral countermeasures to these host responses [1-3]. Perhaps the strongest evidence for the importance of an innate immune response pathway to combating infection is the identification of pathogen-encoded antagonists of that pathway. The fact that virtually all mammalian viruses encode antagonists of the Type I interferon (IFN) response highlights the importance of this antiviral pathway [3-5]. However, relatively little is known regarding IFN-independent transcriptional response pathways that restrict virus replication and the mechanisms used by viruses to thwart such responses.

We recently reported the discovery of an IFN-independent "FACT-ETS-1 Antiviral Response (FEAR)" pathway that requires the cellular Facilitates Chromatin Transcription (FACT) complex for activation [6]. FACT is an ancient chromatin remodeling factor that is conserved from yeast to humans. In humans, FACT is comprised of human suppressor of ty 16 homologue (hSpt16) and structure specific recognition protein 1 (SSRP1) subunits that regulate cellular gene transcription by assembling and disassembling nucleosomes at specific loci [7, 8]. Using vaccinia virus (VV), a large DNA virus belonging to the *Poxviridae* family, we showed that viral infection triggers the FACT-dependent expression of E26 transformation-specific sequence-1 (ETS-1), an antiviral transcription factor, which subsequently promotes VV restriction [6]. ETS-1 expression during infection is dependent upon FACT complexes that contain a specialized, SUMOylated form of hSpt16 (hSpt16^{SUMO})[6]. However, we found the VV-

encoded A51R protein to block ETS-1 expression by outcompeting SSRP1 for direct

binding to hSpt16^{SUMO} subunits in the cytosol and by tethering hSpt16^{SUMO} to

microtubules [6]. VV mutant strains lacking A51R or encoding A51R mutants unable to

bind hSpt16^{SUMO} strongly induce ETS-1 expression and display attenuated replication in

human cell culture and in mice, suggesting the FEAR pathway is a critical component of

eukaryotic antiviral immunity and that poxvirus A51R proteins function as FEAR pathway

antagonists [6].

Prior to our identification of the FEAR pathway, we had serendipitously discovered that

poxvirus A51R proteins could rescue the abortive infection of lepidopteran (moth and

butterfly) cells by vesicular stomatitis virus (VSV), a small RNA virus belonging to the

Rhabdoviridae family [9]. Although VSV is an arbovirus and can be naturally transmitted

by blood-feeding dipteran (flies and mosquitoes) insects to mammalian hosts,

lepidopterans are unnatural hosts for this virus. Thus, we suspected that A51R proteins

may promote productive VSV replication in lepidopteran cells by suppressing an immune

response that VSV is incapable of evading in these host cells [9]. Given that A51R

proteins are only encoded by vertebrate poxviruses [9], we suspected that these viral

proteins are likely immunosuppressive in insect cells because they target antiviral

machinery conserved between invertebrate and vertebrate hosts. However, the nature of

the host response restricting VSV and suppressed by A51R has remained elusive.

Understanding innate immune responses restricting VSV replication is important for

many reasons. For one, VSV it is associated with outbreaks in livestock in the United

States and thus poses a significant economic burden to the agriculture industry [10].

Moreover, the early development of a reverse genetic system for VSV [11, 12], its ability

to infect a broad range of host cell types [13], and inability to cause significant disease in

humans [14], have made VSV an important model for understanding fundamental

aspects of nonsegmented negative-strand (NNS) RNA virus biology. VSV has also

become an important vaccine platform for a wide variety of other viral diseases [14, 15].

VSV is also being investigated as an oncolytic agent in basic and clinical studies [16,

17]. A rapid and lytic replication cycle, and lack of preexisting immunity in the human

population to VSV, are key advantages of its use in oncolytic virotherapy. Oncolytic VSV

strains encoding either a deletion or Arg substitution at Met51 in the VSV M protein

(VSV$^{\Delta M51}$ and VSV^{M51R}) have been widely studied in oncolytic virotherapy [16-23]. These

strains typically display attenuated replication due to their reduced ability to block host

gene expression, although how wild-type VSV M normally impedes cellular gene

expression is controversial [24, 25]. Prior work has shown that antiviral gene expression

by the IFN response is antagonized by wild-type, but not VSV$^{\Delta M51}$ or VSV^{M51R} strains [16-

19]. It has been suggested that the reduced ability of VSV$^{\Delta M51}$ and VSV^{M51R} strains to

replicate in untransformed cells may be due to defective IFN antagonism [18, 21, 25,

26]. However, in transformed cells, the IFN pathway is often defective because this

pathway can also activate anti-proliferative and pro-death responses that would be

detrimental to cancer cell replication and survival [27]. Thus, part of the selectivity of

VSV$^{\Delta M51}$ and VSV^{M51R} strains for replication in cancer cells may be in part due to

defective IFN signaling in these cells. However, not all transformed cells are IFN-

defective, possibly explaining why some cancer cell types are refractory to oncolytic VSV

strains [18, 21, 23]. Indeed, prior work has shown that certain cancer cell lines that are

refractory to VSV$^{\Delta M51}$ replication can become sensitized to infection by treatment with

compounds that inhibit IFN responses [18, 21]. However, it is unclear if IFN-independent

responses also contribute to VSV$^{\Delta M51}$ and VSV^{M51R} restriction in refractory cancer cell

types. Thus, identifying and overcoming innate immune responses present in cancer

cells that restrict oncolytic VSV strain replication is critical for improving virotherapy

because enhanced VSV replication is associated with increased oncolysis and improved

virotherapy outcomes [18, 20, 21, 28, 29].

Interestingly, upregulation of FACT and ETS-1 have been linked to oncogenesis and/or

tumor invasion [30-32]. For example, FACT subunit upregulation is strongly associated

with aggressive tumor types [31], suggesting that FACT can drive oncogenic gene

expression programs that favor cancer development. This led to the development of

small molecule inhibitors of FACT called "curaxins" that are currently being pursued in

clinical trials for the treatment of human malignancies [33]. ETS-1 is a proto-oncoprotein

and its upregulation promotes cellular transformation and tumor invasion [30, 34]. These

observations suggest that the FEAR pathway involves host factors that are both antiviral

and pro-tumoral. Thus, if the FEAR pathway is retained in transformed cells, it may be

relevant to the restriction of oncolytic viruses and thus blocking this pathway with FACT

inhibitors such as curaxins may be a logical strategy to improve the oncolytic virus

replication.

With our discovery of poxvirus A51R proteins as the first DNA virus-encoded FEAR

pathway antagonists [6], we wanted to use VSV as a model to investigate if RNA viruses

activate and/or antagonize this pathway. Here, we show that the attenuated replication

of VSV$^{\Delta M51}$/VSV^{M51R} strains is at least in part due their inability to suppress FEAR

pathway activation in human cells. In contrast, VSV strains encoding wild-type M

proteins effectively suppress this pathway by promoting the proteasome-dependent

degradation of hSpt16^{SUMO} to abrogate ETS-1 expression and by blocking ETS-1 nuclear

accumulation. We find that inhibition of the FEAR pathway in human cancer lines

normally resistant to VSV$^{\Delta M51}$/VSV^{M51R} infection by hSpt16 RNA interference (RNAi) or

curaxin treatment sensitizes these cancer cells to infection with these oncolytic viruses.

Moreover, we show that the host range restriction of VSV in lepidopteran cells likely

results from an inability of VSV M to degrade SUMOylated Spt16 proteins encoded by

these insect hosts. Our data suggests that VV A51R rescues this VSV host range

restriction by interacting with SUMOylated Spt16 expressed in these insect cells. Finally,

we show that murine and human paramyxovirus-encoded C proteins target SUMOylated

Spt16 proteins for proteasomal degradation in mouse and human cells using a

conserved motif in their N-termini, suggesting that other NNS RNA viruses have

independently evolved strategies to degrade these host factors. Collectively, our work

illustrates that RNA viruses can also activate and antagonize the FEAR pathway and, to

our knowledge, identifies VSV M and paramyxovirus C proteins as the first RNA virus-

encoded FEAR pathway antagonists.

Results

hSpt16 or ETS-1 depletion rescues the replication defect of a VSV^{M51R} mutant

We previously showed that hSpt16 RNAi significantly enhances the replication of VSV in

both A549 cells and primary neonatal human dermal fibroblasts, suggesting that FACT

contributes to VSV restriction [6]. However, we wanted to ask if ETS-1 was also

restrictive to VSV replication because it was possible that FACT-mediated VSV

restriction is independent of ETS-1. Moreover, because VSV$^{\Delta M51}$ or VSV^{M51R} strains are

thought to exhibit replication defects because of their inability to block host gene

expression-dependent antiviral responses, we wondered whether these replication

defects may be related to a differential ability of these mutant viruses to antagonize the

FEAR pathway.

To begin to address these questions, we assessed the replication of recombinant VSV

strains encoding eGFP and either wild-type (VSV-eGFP) [35] or M51R mutant M

(VSV^{M51R}-eGFP) proteins [35] in A549 cells transfected with small interfering RNAs

(siRNAs) targeting hSpt16 or ETS-1 [6]. We found hSpt16 or ETS-1 RNAi to significantly

enhance VSV-eGFP replication compared to scrambled (control) RNAi conditions,

suggesting the FEAR pathway is indeed restrictive to VSV. The VSV^{M51R}-eGFP strain

exhibited a replication defect compared to VSV-eGFP under control RNAi conditions but

strikingly, under either hSpt16 or ETS-1 RNAi conditions, this mutant replicated to titers

indistinguishable from VSV-eGFP (**Fig 1A and B**). These results suggest that the FEAR

pathway contributes to VSV restriction and that the VSV^{M51R}-eGFP mutant strain is more

sensitive to restriction by this pathway.

Wild-type VSV, but not VSV^{M51R}, inhibits ETS-1 expression to evade restriction

The results above were reminiscent of our VV studies where we found a VV strain

lacking A51R (ΔA51R) to exhibit a replication defect that could be rescued by FACT

subunit or ETS-1 depletion. We also found this ΔA51R strain to more strongly induce

ETS-1 expression than wild-type VV [6]. Thus, we next used infection time courses and

immunoblots (IBs) to determine if ETS-1 levels differ between VSV-eGFP and VSV^{M51R}-

eGFP infections. Interestingly, while ETS-1 protein levels were not significantly different

between mock- and VSV-eGFP-infected cells, VSV^{M51R}-eGFP-infected WCE showed

significantly elevated ETS-1 by 4 h post-infection (hpi) (**Fig 1C and D**). These results

indicate that infection with the VSV^{M51R}-eGFP mutant induces ETS-1 expression while

wild-type (VSV-eGFP) infection does not induce ETS-1.

We hypothesized that wild-type VSV suppresses ETS-1 expression to prevent activation

of antiviral gene expression programs that require elevated ETS-1 levels. Therefore, we

next asked if ETS-1 overexpression prior to infection could suppress VSV-eGFP

infection. To do this, we overexpressed an HA-tagged human ETS-1 construct (ETS-1-

HA) in cells for 48 h (**Fig 1E**) and then challenged cells with VSV-eGFP. Using

fluorescence microscopy, we quantified GFP signals as a readout for infection and

normalized these signals to total cell number by labeling cell nuclei with NucBlue.

Compared to cultures transfected with empty vector (control), ETS-1-HA-transfected

cells displayed significantly reduced GFP signal (**Fig 1F and G**). Collectively, these data

suggest ETS-1 expression results in a block to VSV infection but VSV-eGFP evades this

restriction by suppressing ETS-1 expression. In contrast, VSV^{M51R}-eGFP is unable to

block ETS-1 induction.

VSV^{M51R} infection induces ETS-1 expression independently of RIG-I and the IFN

response

We showed ΔA51R VV strains to induce ETS-1 expression in the absence of an intact

IFN response [6]. Therefore, we wanted to determine if ETS-1 induction by VSV^{M51R}-

eGFP infection was also independent of the IFN response. Since the pattern recognition

receptor RIG-I is required for the activation of IFN responses during VSV infection of

non-immune cell types [36], we first examined ETS-1 induction in RIG-I-deficient A549

cells. No significant differences in ETS-1 induction were found between wild-type and

RIG-I knockout cells during VSV^{M51R}-eGFP infection (**Fig S1A and B**). We also

quantified ETS-1 expression levels in A549 cells lacking IRF3 or STAT1, two critical

components of the IFN signaling pathway [37, 38]. ETS-1 induction by VSV^{M51R}-eGFP

infection was unaffected in these IFN pathway knockout cell lines (**Fig S1C-F**).

Collectively, these data indicate that, like VV, VSV activates ETS-1 expression

independently of the IFN response.

VSV, but not VSV^{M51R}, promotes hSpt16^{SUMO} depletion to prevent ETS-1 induction

During VV infection, FACT complexes comprised of hSpt16^{SUMO} and SSRP1 subunits

are required to induce ETS-1 expression [6]. Therefore, we hypothesized that VSV-

eGFP may suppress ETS-1 expression by altering FACT subunit levels. Using infection

time courses and IBs, we determined if hSpt16^{SUMO} or SSRP1 levels differ between

VSV-eGFP and VSV^{M51R}-eGFP infections. It is important to note that both hSpt16^{SUMO}

and SUMOless hSpt16 forms can be resolved from one another by SDS-PAGE using

low percentage acrylamide protein gels and then simultaneously detected via IB with

hSpt16 antibodies (Ab) [6]. Strikingly, our IBs revealed a clear and significant reduction

of hSpt16^{SUMO} (upper band in **Fig 2A**) levels compared to mock-infected WCE during

VSV-eGFP infection by 4 hpi and hSpt16^{SUMO} became virtually undetectable by 8 hpi. In

contrast, hSpt16^{SUMO} levels remained unchanged throughout the entire VSV^{M51R}-eGFP

infection time course (**Fig 2A and B**). Interestingly, SUMOless hSpt16 (lower band in

Fig 2A) or SSRP1 levels did not change during infection with either virus (**Fig 2C and**

D), suggesting that VSV infection leads to the specific depletion of hSpt16^{SUMO} subunits.

When comparing the effect of MOI on hSpt16^{SUMO} depletion during VSV-eGFP infection,

we noticed that complete depletion of hSpt16^{SUMO} levels occurred earlier (by 4 hpi) when

using higher MOIs (e.g. 10-30) versus during lower MOI (e.g. 3) infections. However, the

VSV^{M51R}-eGFP strain could not deplete hSpt16^{SUMO} at any MOI or time point (**Fig 2A**

and S2).

To confirm these data, and to determine if loss of hSpt16^{SUMO} levels during VSV-eGFP

infection was not due to a global down regulation of intracellular SUMOylation, we used

Ab against SUMO-1 or SUMO-2/3 to immunoprecipitate (IP) SUMOylated protein

fractions from mock-, VSV-eGFP-, or VSV^{M51R}-eGFP-infected WCE [6]. As expected,

hSpt16^{SUMO} was specifically depleted in VSV-eGFP-infected WCE while present in

VSV^{M51R}-eGFP-infected lysates. Moreover, no overt reductions in either total

SUMOylation banding patterns or levels of SUMOylated-RanGAP1 (an abundant,

SUMOylated protein [39]) were observed in VSV-eGFP versus VSV^{M51R}-eGFP infections

(**Fig 2E and F**). These data suggest that VSV-eGFP can promote depletion of hSpt16

that is SUMOylated with either SUMO-1 or SUMO-2/3 and that the specific loss of

hSpt16^{SUMO} levels during VSV-eGFP infection is not due to a generalized decrease in

intracellular SUMOylation.

Finally, because hSpt16^{SUMO} is required for ETS-1 induction during ΔA51R VV infection

[6], we wanted to determine if hSpt16 RNAi treatment would inhibit ETS-1 induction by

VSV^{M51R}-eGFP infection. Consistent with our VV studies, hSpt16 RNAi treatment

abrogated ETS-1 induction after VSV^{M51R}-eGFP infection (**Fig 2G**). Collectively, these

data indicate that the inability of VSV^{M51R}-eGFP to deplete hSpt16^{SUMO} results in robust

ETS-1 induction during infection.

VSV M is sufficient to promote proteasomal degradation of cytoplasmic hSpt16^{SUMO} and to block ETS-1 expression

Given that the only difference between VSV-eGFP and VSV^{M51R}-eGFP strains is the M51R substitution in the VSV M protein, we hypothesized that this viral factor was responsible for hSpt16^{SUMO} depletion. Consistent with this, a VSV$^{\Delta M51}$ strain expressing GFP (VSV$^{\Delta M51}$-GFP [18]) was also unable to deplete hSpt16^{SUMO} levels when compared to its wild-type VSV-GFP control strain (**Fig S3**). However, we still conducted a screen of 4 of the 5 VSV-encoded proteins to determine if any of these viral factors could reduce hSpt16^{SUMO} levels when expressed from plasmids in uninfected human cells. The VSV L protein, encoding the viral RNA polymerase, was excluded from this screen as it seemed unlikely to be involved in immune evasion. Consistent with our hypothesis, only expression of Flag-tagged VSV M (VSV M-Flag) resulted in hSpt16^{SUMO} depletion (**Fig 3A**). Moreover, expression of either VSV M$^{\Delta M51}$-Flag or VSV M^{M51R}-Flag mutant constructs failed to deplete hSpt16^{SUMO} levels (**Fig 3B**), consistent with our infection results. These data indicate that wild-type VSV M proteins, but not M51R or ΔM51 mutants, are sufficient for hSpt16^{SUMO} depletion in the absence of other VSV proteins.

VSV M can inhibit host gene expression by binding to Rae1, a component of the cellular Rae1-Nup98 export complex that promotes mRNA export from the nucleus [25]. However, the mechanism by which VSV M-Rae1 interaction inhibits host gene expression is controversial [24, 25]. In one proposed model, VSV M binds Rae1 to inhibit Rae1-Nup98-mediated mRNA export from the nucleus while an alternative model suggests VSV M may usurp Rae1-Nup98 complexes to inhibit cellular gene transcription

[24, 25]. Notably, VSV M51R/∆M51 M mutants cannot interact with Rae1 [25]. Despite

the unresolved nature of VSV M-Rae1 interaction models, we wanted to determine if

Rae1 or Nup98 was involved in hSpt16^{SUMO} level regulation. Thus, we tested if depletion

of either Rae1 or Nup98 altered hSpt16^{SUMO} levels in mock-infected cells or inhibited

hSpt16^{SUMO} depletion during VSV-eGFP infection. In either case, we observed no effect

of Rae1 or Nup98 depletion on hSpt16^{SUMO} levels (**Fig S4**). This suggests that VSV M

antagonism of the FEAR pathway cannot be explained by its interaction with Rae1-

Nup98 complexes.

We next wanted to gain additional insight into the mechanism by which VSV M depletes

hSpt16^{SUMO} levels. Given that VV A51R blocks the FEAR pathway by binding to cytosolic

hSpt16^{SUMO} subunits, we next asked if VSV M also targets cytosolic hSpt16^{SUMO}

subunits. To do this, we co-expressed VSV M-Flag constructs with either a wild-type or

nuclear localization sequence (NLS)-deleted (NLS) HA-tagged hSpt16 constructs we

previously generated [6]. Importantly, we confirmed that wild-type HA-hSpt16 displayed

both cytosolic and nuclear localization while the HA-hSpt16$^{∆NLS}$ mutant was restricted to

the cytosol (**Fig 3C**). Upon co-transfection with wild-type VSV M-Flag, the SUMOylated

forms of both wild-type and NLS-deleted HA-hSpt16 constructs were depleted (**Fig 3D**),

indicating VSV M-Flag likely targets hSpt16^{SUMO} subunits in the cytoplasm. To determine

if VSV M promotes proteasome-dependent degradation of hSpt16^{SUMO}, we expressed

VSV M-Flag in human cells in the absence or presence of proteasome inhibitors

bortezomib or MG132 and assessed hSpt16^{SUMO} levels by IB. Interestingly, hSpt16^{SUMO}

depletion was blocked in the presence of either proteasome inhibitor (**Fig 3E and F**),

suggesting VSV M promotes the proteasome-dependent degradation of hSpt16^{SUMO}.

Finally, we asked if plasmid-based expression of VSV M-Flag was sufficient to block

ETS-1 induction by either VSV^{M51R}-eGFP (**Fig 3G and H**) or ΔA51R VV (**Fig 3I and J**)

infection. In both cases prior expression of VSV M-Flag, but not VSV M^{M51R}-Flag, was

able to suppress ETS-1 induction by either of these mutant viruses (**Fig 3G-J**). The

ability of VSV M to complement the ΔA51R strains inability to block ETS-1 expression

demonstrates VSV M is a bona fide FEAR pathway antagonist that can function

independently of other VSV proteins. Collectively, our prior and current findings suggest

that VV and VSV have independently evolved unique strategies to inhibit FACT-

dependent ETS-1 expression by either tethering hSpt16^{SUMO} to microtubules (VV A51R)

or by promoting hSpt16^{SUMO} degradation (VSV M), illustrating that FEAR pathway

suppression is important for both DNA and RNA viruses.

VSV M encodes a second mechanism to inhibit ETS-1 function: blockade of

nuclear import

While the specific antiviral function that ETS-1 performs after FEAR pathway activation

is still an active area of investigation, ETS-1 is known to regulate host transcription in the

nucleus. Thus, we determined if ETS-1 localizes to the nucleus during VSV infection.

First, we used immunofluorescence microscopy to examine endogenous ETS-1

localization in mock-, VSV-eGFP-, or VSV^{M51R}-eGFP-infected human A549 cells. In

mock-infected cells, both cytosolic and nuclear ETS-1 staining was observed, consistent

with the known nucleocytoplasmic cycling of ETS-1 [40, 41]. However, during VSV-

eGFP infection, ETS-1, was largely excluded from the nucleus and restricted to the

cytosol, suggesting that VSV infection impairs ETS-1 nuclear translocation. In contrast,

ETS-1 signals were stronger and predominantly nuclear in VSV^{M51R}-eGFP-infected cells

(**Fig 4A**). To complement our microscopy experiments, we conducted cell fractionation

and IB studies to quantify the percentage of total ETS-1 protein in nuclear fractions.

Consistent with our microscopy results, ~37% of total ETS-1 was found in nuclear

fractions of mock-infected cells, while only ~10% of ETS-1 was nuclear during VSV-

eGFP infection. In contrast, ~78% of ETS-1 was found in the nucleus during VSV^{M51R}-

eGFP infection (**Fig 4B and C**). Importantly, immunofluorescence experiments in U2OS

cells also demonstrated a similar exclusion of ETS-1 from the nucleus during VSV-eGFP

infections as observed in A549 cells (**Fig 4D**), suggesting this was not cell type-specific.

Finally, to determine if VSV M was able to impede ETS-1 nuclear localization in the

absence of other VSV proteins, we co-transfected the ETS-1-HA expression vector with

empty vector, VSV M-Flag, or VSV^{M51R}-Flag constructs into U2OS cells. In the presence

of empty vector or VSV^{M51R}-Flag constructs, ETS-1-HA displayed both nuclear and

cytosolic localization. However, in the presence of wild-type VSV M-Flag, ETS-1-HA

staining was largely excluded from the nucleus and restricted to the cytoplasm (**Fig 4E**).

Collectively, these data indicate that, during infection, ETS-1 accumulates in the nucleus

in the presence of the M51R mutant M protein but not in the presence of wild-type VSV

M. Moreover, our transfection experiments suggest VSV M can prevent ETS-1 nuclear

accumulation in the absence of other VSV proteins, indicating it encodes an additional

mechanism of antagonizing the FEAR pathway- blocking ETS-1 nuclear import.

FEAR pathway inhibition enhances oncolytic VSV strain replication in refractory

cancer cells

VSV$^{\Delta M51}$ and VSV^{M51R} viruses are being pursued for use in oncolytic virotherapy. While

effective at infecting some human cancer cell types, some transformed cell types have

proven to be refractory to infection by these oncolytic VSV strains, posing a major

challenge to their wide-spread use [18, 21, 23]. Thus, we asked if the FEAR pathway

may contribute to restriction of oncolytic VSV strain replication in human cancer lines

known to be refractory to $VSV^{\Delta M51}$/VSV^{M51R} viruses. First, we measured replication of

$VSV^{\Delta M51}$-GFP in human 786-0 renal adenocarcinoma cells, which are relatively resistant

to infection with this virus [18, 21], after hSpt16 RNAi. As shown in **Fig 5A**, $VSV^{\Delta M51}$-GFP

replication was significantly increased in 786-0 cells after hSpt16 depletion by RNAi.

Importantly, we confirmed that both hSpt16 and ETS-1 were reduced in 786-0 cells

infected with $VSV^{\Delta M51}$-GFP during hSpt16 knockdown (**Fig 5B**). These data suggest the

FEAR pathway contributes to the restriction of this virus in this restrictive cell type.

Since the use of hSpt16 RNAi as a therapeutic strategy to inhibit the FEAR pathway is

likely to be technically challenging *in vivo*, we conducted a "proof of principle" series of

cell culture experiments to determine if treatment with the small molecule FACT inhibitor,

CBL0137, could similarly enhance $VSV^{\Delta M51}$-GFP replication in restrictive cancer cell

types. CBL0137 is a second-generation curaxin and has broad activity against diverse

cancer cell types in culture and in mouse models and is being pursued in clinical trials as

a therapeutic for various malignancies [33, 42-45]. Curaxins inhibit FACT by causing this

complex to become trapped on chromatin, preventing it from functioning properly in gene

expression regulation [33]. First, we used quantitative fluorescence microscopy and GFP

signal as a simple readout of $VSV^{\Delta M51}$-GFP replication in 786-0 cells in the presence of a

broad range of CBL0137 concentrations 24 hpi. Addition of CBL0137 to media 1 hpi led

to a dose-dependent increase in GFP signal, peaking at an ~34-fold increase in GFP

signal with addition of 3 μM CBL0137 compared to infected cultures without curaxin (**Fig**

5C and D, right axis). We also analyzed cell viability via fluorometric resazurin assays

of parallel 786-0 cultures treated with either CBL0137 alone or VSV$^{\Delta M51}$-GFP+CBL0137

after 48 h and found a clear synergistic effect of the combined treatment on cancer cell

killing. For example, in cultures treated with 1.3 μM CBL0137 only, cell viability was

~72% of untreated cultures. However, combining 1.3 μM curaxin with VSV$^{\Delta M51}$-GFP

treatment reduced 786-0 cell viability to ~10% of controls (**Fig 5D, left axis**). Importantly,

we verified that the increased GFP signal observed in CBL0137-treated cultures

correlated with elevated VSV$^{\Delta M51}$-GFP titers (**Fig 5E**) and with inhibition of ETS-1

expression (**Fig 5F**). We also tested the human melanoma cell line, RPMI 7951, which is

resistant to infection with VSV^{M51R} oncolytic strains [23] and observed enhanced

VSV$^{\Delta M51}$-GFP replication in these cells in the presence of CBL0137 (**Fig 5G**), suggesting

curaxin may sensitize other types of cancer cells to oncolytic VSV strain infection.

Collectively, these data suggest that hSpt16 depletion or FACT inhibition with curaxins

can enhance the replication of an oncolytic VSV strain lacking a functional FEAR

pathway antagonist in restrictive human cancer cell types.

FACT contributes to VSV host range restriction in *Lymantria dispar* insect cells

We originally became interested in understanding poxvirus A51R protein function after

discovering that these proteins rescue the post-entry abortive infection of VSV in

Lymantria dispar (gypsy moth or spongy moth)-derived LD652 cells [9]. With the

knowledge that VV A51R binds to hSpt16^{SUMO} to antagonize its antiviral function [6] and

that Spt16 proteins are conserved in all eukaryotes, we wondered if VV A51R-mediated

rescue of VSV in LD652 cells related to its ability to antagonize insect Spt16 function.

After sequencing the *L. dispar* genome [46], we discovered that hSpt16 and *L. dispar*

Spt16 (LdSpt16) proteins share an overall a.a. identity of ~60% [6]. However, alignment

of the A51R-binding domain in hSpt16 [6] with the corresponding region in LdSpt16

showed this region to be ~73% identical between these Spt16 proteins (**Fig S5**).

However, because VV A51R only binds SUMOylated forms of hSpt16, it was important

to determine if LdSpt16 was also SUMOylated. Interestingly, IBs of LD652 WCE

revealed two LdSpt16 forms as found with hSpt16 in human cells. Treatment of LD652

cultures with tannic acid, a global SUMOylation inhibitor [47], eliminated the upper

LdSpt16 form on IBs, suggesting this band represents SUMOylated LdSpt16

(LdSpt16[SUMO]) (**Fig 6A**). To determine if VV A51R interacts with LdSpt16[SUMO], we

infected LD652 cells with a recombinant VV strain encoding Flag-tagged A51R (FA51R)

under its natural *A51R* gene promoter (ΔA51R[FA51R]) and then performed co-

immunoprecipitation [6]. Interestingly, LdSpt16[SUMO], but not SUMOless LdSpt16 proteins,

specifically co-immunoprecipitated with FA51R in LD652 WCE (**Fig 6B)**, suggesting VV

A51R can interact with SUMOylated Spt16 proteins in this insect host. Next, we tested if

A51R-LdSpt16 interactions were required for A51R-mediated rescue of VSV replication

in LD652 cells. To do this, we transfected LD652 cells with a control expression vector

encoding Flag-tagged GFP (FGFP) or vectors encoding wild-type FA51R, or a FA51R

mutant that cannot bind to SUMOylated Spt16 proteins FA51R[158-162Ala] [6]. 48 h post-

transfection, we challenged cells with a recombinant VSV strain encoding firefly

luciferase (VSV-LUC) [48]. We then assessed VSV-LUC gene expression with luciferase

assays and measured viral titers in collected LD652 culture supernatants [9]. While the

wild-type FA51R construct was able to rescue both VSV gene expression and

replication, the FA51R[158-162Ala] mutant was completely defective in this regard (**Fig 6C**

and D). These results suggest that VV A51R-mediated rescue of VSV requires A51R-

LdSpt16^{SUMO} interaction.

To directly test if LdSpt16 was involved in VSV restriction, we used *in vitro*-transcribed

dsRNAs to target LdSpt16 by RNAi. Compared to control RNAi targeting an irrelevant

GFP mRNA sequence, VSV-LUC gene expression (**Fig 6E**) and replication (**Fig 6F**)

were significantly enhanced in LD652 cultures transfected with dsRNAs targeting

LdSpt16 mRNA. These results suggested that the abortive infection of wild-type VSV in

LD652 cells is at least in part dependent upon LdSpt16^{SUMO}-mediated restriction.

Therefore, we hypothesized that wild-type VSV M may be defective in its ability to target

LdSpt16^{SUMO} for degradation. Consistent with this hypothesis, VSV M-Flag transfection

into LD652 cells failed to promote LdSpt16^{SUMO} depletion at any expression level tested

(**Fig 6G**). Collectively, these results suggest that VSV host range restriction in LD652

cells may be due to an inability of VSV M to antagonize LdSpt16^{SUMO} function.

Paramyxovirus-encoded C proteins also promote proteasomal degradation of

hSpt16^{SUMO}

Finally, we asked if other NNS RNA viruses, unrelated to VSV, also interact with the

FEAR pathway. We focused on viruses belonging to the *Paramyxoviridae* family

because they include important animal and human pathogens. For example, Sendai

virus (SeV) is a natural pathogen of rodents and a common cause of viral pneumonia in

wild and laboratory rodents [49]. Although not known to be pathogenic to humans, SeV

can productively replicate in human cells and has been an important model for

understanding paramyxovirus replication and immune evasion mechanisms [50, 51].

To determine if the FEAR pathway is relevant to paramyxovirus restriction, we

challenged A549 cells with a recombinant SeV strain encoding GFP (SeV-GFP) after

hSpt16 RNAi. Compared to control RNAi conditions, hSpt16 RNAi led to a significant

increase in SeV-GFP replication (**Fig 7A**), suggesting the FEAR pathway may also

restrict paramyxoviruses. We next examined if FACT subunit levels were altered during

SeV-GFP infection. Due to the slower replication cycle of SeV compared to VSV, we

collected WCE at relatively later time points (8 and 24 hpi). Interestingly, while FACT

subunit levels were largely unchanged at 8 hpi, hSpt16^{SUMO} subunits were specifically

eliminated in SeV-GFP-infected cells by 24 hpi, suggesting SeV encodes an antagonist

of hSpt16^{SUMO} (**Fig 7B**).

We suspected that the SeV-encoded accessory "C" proteins may be involved in

mediating hSpt16^{SUMO} depletion because they function as immune evasion factors [50].

During SeV infection, C proteins are expressed as a nested set of four different proteins

(C', C, Y1, and Y2) that derive from the same SeV *P* gene-derived mRNA and contain

common C-termini but differ in their N-termini due to the use of alternative translation

start sites [52-54] (**Fig 7C**). SeV C' is the largest of these four proteins so we first

examined if Flag-tagged SeV C' (SeV C'-Flag) expression could promote hSpt16^{SUMO}

depletion. Interestingly, expression of SeV C'-Flag was sufficient to promote hSpt16^{SUMO}

depletion in 293T cells in the absence of other SeV proteins. Moreover, this SeV C'-

mediated hSpt16^{SUMO} depletion was blocked in the presence of bortezomib (**Fig 7D**).

Interestingly, SeV C'-Flag could also deplete SUMOylated forms of the HA-hSpt16$^{\Delta NLS}$

mutant that cannot enter the nucleus (**Fig S6**). Importantly, SeV C'-Flag expression also

reduced SUMOylated Spt16 levels in mouse 3T3 cells (**Fig S7**), indicating that this viral

activity is retained in a natural SeV host species. These results collectively suggest that

like VSV M, SeV C' promotes proteasomal degradation of SUMOylated host Spt16

proteins in the cytoplasm.

We next examined if any of the shorter C proteins (C, Y1, and Y2) could also deplete

hSpt16^{SUMO} levels. Interestingly, only C' and C proteins were capable of depleting

hSpt16^{SUMO} in 293T cells (**Fig 7E**), suggesting the N-terminal 23 a.a. common between

C' and C proteins, but absent in Y1 and Y2 proteins (**Fig 7C**), are critical for targeting

hSpt16^{SUMO} for degradation. To determine if this 23 a.a. region was sufficient to promote

hSpt16^{SUMO} depletion, we fused these residues to the N-terminus of a GFP-Flag

construct, creating SeV C'$^{12-34}$-GFP-Flag. Strikingly, expression of SeV C'$^{12-34}$-GFP-Flag

was sufficient to deplete hSpt16^{SUMO} levels (**Fig 7F**).

Human parainfluenza virus-1 (HPIV-1), also known as human respirovirus 1, is a

clinically important paramyxovirus that is the leading cause of laryngotracheobronchitis

(croup) in children, a disease associated with swelling of the airways and difficulty

breathing [55]. HPIV-1 is closely related to SeV, sharing similar genetic and antigenic

properties. For example, the C proteins of SeV and HPIV-1 are ~70% identical [56].

Therefore, we examined the effect of hSpt16 RNAi on the replication of a GFP-encoding

HPIV-1 strain (HPIV-1-GFP) and the ability of this virus to target hSpt16^{SUMO} for

degradation. We found hSpt16 RNAi to enhance HPIV-1-GFP replication in human cells

(**Fig 7G**). Interestingly, as with SeV infection, hSpt16^{SUMO} was specifically depleted in

A549 cells by 24 hpi with HPIV-1-GFP (**Fig 7H**). Moreover, expression of a HPIV-1 C'-

Flag construct in 293T cells was sufficient to deplete hSpt16^{SUMO} and this depletion was

blocked by bortezomib (**Fig 7I**). Collectively, these results suggest that murine and

human paramyxovirus-encoded C proteins function as FEAR pathway antagonists that

target SUMOylated host Spt16 proteins for degradation.

519   VSV M and paramyxovirus C proteins are structurally unrelated

Given that both VSV M and paramyxovirus C proteins promote hSpt16^{SUMO} degradation,

we compared these proteins for similarity at primary (a.a. sequence) and tertiary

(structural) levels. Using basic local alignment search tool (BLAST) [57] we did not

observe any significant similarity between VSV M and either SeV C' or HPIV-1 C'

sequences.

We next examined tertiary structures of these proteins. Although a full-length VSV M

structure for the Indiana serotype used in our study has not been solved, the structure of

a VSV M fragment encompassing the globular domain (a.a. 44-229) (**Fig S8A**) bound to

Rae1-Nup98 complexes has been reported (4OWR) [58] (**Fig S8B**). In 4OWR, M

contacts Rae1-Nup98 complexes using three distinct regions that project from a "fist-

like" globular domain: a finger (a.a. 49-61), a thumb (a.a. 213-223), and a "web" region

(a.a. 142-144). Importantly, the finger motif, encompassing the M51 residue, makes

critical contacts with residues in a Rae1 β-propeller [58], illustrating the importance of

this region to host interactions. Since it was our goal to compare complete VSV M and

paramyxovirus C' protein structures to one another, we generated a full-length (a.a. 1-

229) VSV M structure using Alphafold2 [59](**Fig S8C**) and then compared this with

4OWR by overlaying them (**Fig S8D**) and conducting pairwise similarity tests using

RSCB PDB [60] to assign root mean square deviation (RMSD) values and template

modeling (TM) scores (**Fig S8E**). RMSD scores of <3 Å and TM scores >0.5 were

considered to have similar overall folds while those producing RMSD values >3 Å and

TM scores <0.5 were considered to be structurally unrelated [61]. Pairwise comparisons

between 4OWR and our predicted VSV M structure produced a RMSD of 1.82 Å and a

TM score of 0.93, providing confidence in our Alphafold2-generated VSV M structure.

We then used full-length SeV C' and HPIV-1 C' proteins (**Fig S9A**) to generate

structures with Alphafold2. We aligned our Alphafold2 SeV C' structure with a solved

structure from a SeV C fragment (a.a. 99-204) in complex with host Alix proteins (6KP3)

[62] (**Fig S9B-D**). Although 6KP3 is missing the N-terminal 23 a.a. region in SeV C' (a.a.

12-34) involved in hSpt16^{SUMO} depletion (**Fig S9A**), it contains most of the structured

globular domain (a.a. 88-215), allowing us to conduct a pairwise comparison to

determine how well the Alphafold2 structure aligned with 6KP3. We obtained a RSMD

value of 1.22 Å and TM score of 0.92 (**Fig S9E**), suggesting that the regions we could

compare were similar in structure, giving us confidence in the Alpahfold2 model.

Interestingly, the N-terminal domain of SeV C' (a.a. 1-87) in the Alphafold2 model is

completely unstructured with the exception of an alpha helix (a.a. 14-34) we term "α1"

that largely encompasses the region involved in hSpt16^{SUMO} depletion (a.a. 12-34) (**Fig**

S9A and C). Unfortunately, structures for HPIV-1 C proteins are not currently available,

therefore we used an Alphafold2 model of full-length HPIV-1 C' (**Fig S9F**) to overlay with

our predicted SeV C' structure (**Fig S9G**). Importantly, pairwise comparisons of these

SeV and HPIV-1 structures generated a RSMD of 2.57 Å and TM score of 0.66 (**Fig**

S9H), consistent with the high degree of conservation of these paramyxovirus proteins

[56]. Interestingly, we again found the N-terminal region (a.a. 1-93) of HPIV-1 C' to be

entirely disordered with the exception of a short alpha helix (a.a. 18-25, α1) that

corresponds to the N-terminal half of α1 in SeV C' (**Fig S9A and G**). These data suggest

that paramyxovirus C proteins have a conserved alpha helical structure that is likely

important for mediating hSpt16^{SUMO} depletion.

Finally, we overlaid our Alphafold2-generated models of full length VSV M and

paramyxoviruses C' proteins (**Fig S10A**) and used RMSD and TM analyses to compare

VSV M to either paramyxovirus protein. In both pairwise comparisons, we found VSV M

to exhibit RMSD values >3 Å and TM scores <0.5, indicating that these proteins are

structurally unrelated (**Fig S10B**). Collectively, these data suggest that VSV M and

paramyxovirus proteins are structurally unrelated, supporting the notion that

rhabdoviruses and paramyxoviruses evolved distinct factors to target hSpt16^{SUMO} for

degradation.

Discussion

The FEAR pathway was revealed by our discovery that poxvirus A51R proteins

antagonize this pathway [6], highlighting the fact that viral immune evasion factors can

provide insights into the critical host defenses that combat viral infection. While this

antiviral pathway was clearly important for combating DNA viruses, its relevance to RNA

viruses was less clear.

Here, we used VSV as a model RNA virus to examine if these viruses actively induce

and evade the FEAR pathway. We chose this virus in part because we previously

observed a measurable increase in VSV replication after hSpt16 depletion in human

cells [6], hinting at the possibility that this virus may be restricted by the FEAR pathway.

However, our discovery here that hSpt16 or ETS-1 depletion more strongly enhanced

the replication of VSV^{M51R} versus wild-type VSV, implicated the VSV M protein as a

suppressor of the FEAR pathway. Analysis of VSV-infected cell lysates led to our

unexpected discovery that, unlike VV A51R, which tethers hSpt16^{SUMO} to microtubules

[6], VSV M specifically promotes the proteasomal degradation of hSpt16^{SUMO} (**Fig 8**).

The inability of VSV^{M51R} strains to degrade hSpt16^{SUMO} explains the strong induction of

ETS-1 expression by these viruses. Importantly, the fact that VSV M expressed from a

plasmid could not only deplete hSpt16^{SUMO}, but also block ETS-1 induction by the ΔA51R

VV strain, indicates it can antagonize ETS-1 expression independently of other VSV

proteins. Moreover, our microscopy and cell fractionation studies suggest that wild-type,

but not M51R mutant, VSV M also prevents ETS-1 nuclear accumulation as a second

mechanism to inhibit the FEAR pathway.

The molecular mechanisms by which VSV M promotes hSpt16^{SUMO} degradation and

prevents ETS-1 nuclear accumulation are active areas of investigation. Given that the

most well-characterized defect of the M51R M mutant is its inability to interact with the

host Rae1-Nup98 mRNA export complex, we were surprised that Rae1 or Nup98

depletion had no effect on VSV-mediated hSpt16^{SUMO} depletion. Given that the M51

residue resides in a "finger-like" projection that protrudes from the globular domain of

VSV M [58], it may be a docking site for not only Rae1 but also for additional host

factors. We speculate that VSV M may use this finger projection to recruit ubiquitin-

proteasome system-related machinery (e.g. E3 ubiquitin ligases) that target hSpt16^{SUMO}

for degradation (**Fig 8**). Indeed, the repurposing of host E3 ligases to degrade antiviral

host factors has been widely observed among DNA and RNA viruses [63, 64]. Prior work

has shown that VSV M can inhibit nuclear translocation mediated by specific

karyopherin/importin protein family members in xenopus oocytes and this activity is lost

in the M51R mutant [65, 66]. Thus, it is possible that VSV M interferes with import

factors required for ETS-1 nuclear translocation.

Our discovery that VSV$^{\Delta M51}$/ VSV^{M51R} strains are defective in FEAR pathway antagonism

has led to important implications for their use in oncolytic virotherapy. A major drawback

of these strains is their inability to effectively replicate in cancer cell types that are still

capable of mounting effective antiviral responses [17, 40, 67]. Prior work had shown that

786-0 cells are highly refractory to VSV$^{\Delta M51}$ but that compounds termed "viral sensitizers"

that inhibit IFN responses can enhance the replication in, and killing of, these cells by

this oncolytic strain [18, 21, 68]. Our data demonstrating that hSpt16 RNAi enhances the

replication of a VSV$^{\Delta M51}$strain in these cells suggest that IFN-independent responses (i.e.

the FEAR pathway) are also relevant to the restriction of this strain. Moreover, we found

the FACT inhibitor, CBL0137, to be a promising viral sensitizer that improved VSV$^{\Delta M51}$

replication in these and other human cancer lines known to be resistant to infection by

VSV$^{\Delta M51}$/VSV^{M51R} oncolytic strains. Importantly, CBL0137 is already being pursued in

clinical trials as a cancer chemotherapeutic [33] as FACT upregulation is often observed

during oncogenesis and is associated with more aggressive cancers [31]. Such FACT

upregulation may also lead to more potent FACT-dependent antiviral responses,

potentially posing barriers to oncolytic virus replication in tumors. Thus, it will be

important to determine in future studies if combined (VSV$^{\Delta M51}$+CBL0137) treatment can

improve oncolytic virotherapy efficacy in animal models.

This study has also provided insight into the poorly understood abortive infection of VSV

in lepidopteran insect cells and the rescue of VSV replication in these cells by poxvirus

A51R proteins [9]. We previously found Spt16 proteins from various mammalian, insect,

and nematode cell lysates to exhibit the same, two-band pattern on IBs found in human

cell extracts, suggesting SUMOylation of Spt16 proteins is highly conserved among both

vertebrate and invertebrate eukaryotes [6]. More formal evidence for this was provided

here where we demonstrated that treatment with a SUMOylation inhibitor (tannic acid)

[47] could specifically eliminate the higher molecular weight form of LdSpt16 in *L. dispar-*

derived LD652 cells. Strikingly, despite being encoded by a mammalian poxvirus, VV

A51R was able to interact with LdSpt16^{SUMO}. This is likely due to the high degree of

conservation of the A51R-binding domain [6] among human and moth Spt16 proteins.

Our findings that: 1) VSV M cannot deplete LdSpt16^{SUMO} levels; 2) A51R proteins lacking

a functional Spt16-binding domain cannot rescue VSV replication in *L. dispar* cells; and

3) LdSpt16 RNAi rescues VSV replication, all strongly implicate LdSpt16^{SUMO} in VSV

host range restriction in these cells. Importantly, these results also suggest an ancient

role for SUMOylated Spt16 proteins in eukaryotic antiviral immunity. Future studies will

be needed to determine if insect ETS transcription factors [69] are also involved in virus

restriction and whether the FEAR pathway is indeed intact in invertebrates.

Investigation of paramyxovirus-FEAR pathway interactions revealed a remarkable

example of independent evolution of a common immune evasion mechanism by NNS

RNA viruses. Despite a lack of primary a.a. sequence or structural homology between

VSV M and paramyxovirus C proteins, they could both promote proteasome-dependent

degradation of hSpt16^{SUMO} (**Fig 8**). The specific targeting of SUMOylated Spt16 proteins

by pox-, rhabdo-, and paramyxoviruses suggests these host factors activate a critical

node of antiviral immunity. Interestingly, SeV and HPIV-1 C proteins have also been

shown to antagonize STAT protein function to inhibit IFN signaling [50, 51]. Thus, SeV

and HPIV-1 are likely under similar evolutionary pressure to inhibit key immune

responses pathway components such as STAT1 and Spt16^{SUMO} in their murine and

human hosts. Future studies will be aimed at determining if paramyxovirus C and VSV M

use a common or distinct host interactions to ultimately achieve hSpt16^{SUMO} degradation,

but our current work clearly defines these proteins as the first RNA virus-encoded FEAR

pathway antagonists.

In conclusion, our work has demonstrated that RNA viruses can activate and evade the

FEAR pathway, indicating this pathway is relevant to this important class of pathogens.

Major questions remain regarding how RNA viruses activate the FEAR pathway and how

ETS-1 activation eventually leads to virus restriction. Studies are ongoing to explore

these exciting areas of FEAR pathway biology. Moreover, whether other RNA virus

families antagonize the FEAR pathway is an active area of investigation. Ultimately, a

greater understanding of the activation and evasion of the FEAR pathway by viruses will

help us better appreciate the role of this host response in the evolutionary arms race

between viruses and their hosts.

Materials and Methods

Specific details regarding the source of all key experimental reagents (primers, plasmids,

Abs, viruses, cell lines, etc.) can be found in **Table S1**.

Cell lines and primary cultures. Mammalian cell lines were maintained at 37°C in 5%

CO_2 atmosphere. A549, U2OS, 293T, 786-0, NIH 3T3, Vero, and BHK-21 cells were

cultured in DMEM supplemented with 10% FBS. RPMI 7951 cells were cultured in

EMEM supplemented with 10% FBS. BSC-40 cells were passaged in MEM containing

5% FBS. All media above additionally contained 1% non-essential amino acids, 1% L-

glutamine, and 1% antibiotic/antimycotic (Gibco). LD652 were cultured in a 1:1 mixture

of Ex-Cell 420 (Sigma) and Graces insect medium (Invitrogen) supplemented with 10%

FBS at 27°C under normal atmospheric conditions [9, 70].

Viruses and virus titrations. VSV-GFP and VSV$^{\Delta M51}$-GFP strains were amplified in

either Vero or BHK-21 cells using low MOI infections [9, 71]. All other VSV strains were

amplified using low MOI infections in BHK-21 cells. SeV-GFP and HPIV-1-GFP stocks

were obtained from Viratree. BSC-40 cells were used for VV stock preparation [9].

Unless otherwise noted, virus strains were titrated by plaque assay (VSV-LUC,

ΔA51R^{FA51R}, ΔA51R) or fluorescent foci/plaque assay (VSV-eGFP, VSV^{M51R}-eGFP,

VSV$^{\Delta M51}$-GFP, SeV-GFP, HPIV-1-GFP) on BSC-40 cell monolayers, with a 1.5% low-

melting point agarose (Invitrogen) overlay used for VSV titrations [6]. In some cases,

VSV-GFP and VSV$^{\Delta M51}$-GFP stocks were titered on Vero cells [71]. Experimental viral

infections were incubated for 1 h in serum free DMEM at 37°C before the inoculum was

replaced with complete media for the remainder of the infection with the exception of

SeV-GFP and HPIV-1 cultures, which were kept in serum free DMEM containing either

1% TrypLE Select (Invitrogen) or 5 μg/ml acetylated trypsin (Sigma). Where indicated,

replacement media after 1 h of infection contained indicated doses of CBL0137 (Selleck

Chemicals) and was kept in the media until the end of the experiment. At indicated times

post-infection, infected cell culture supernatants were collected, clarified by

centrifugation, and clarified supernatant titers were determined by serial dilution followed

by plaque assay or fluorescent foci/plaque assay [6].

Expression vectors. The human ETS-1-HA pEZ-M07 expression vector was obtained

from GeneCopoeia. VSV M-Flag and M51R/ΔM51 mutant derivatives were constructed

by gene synthesis (Gene Universal) with inclusion of *Sac*II and *Pac*I sites for cloning into

SacII/PacI-digested pcDNA3 [6]. VSV M-Flag was also cloned into the insect p166

expression vector [9] using SacII/PacI sites. SeV C'-Flag and HPIV-1 C'-Flag expression

constructs were also generated using gene synthesis and cloned into pCNDA3 using

flanking SacII/PacI sites. SeV C-, Y1-, and Y2-Flag expression plasmids were

constructed from PCR amplification from the full-length SeV C' sequence. The SeV

C'$^{M12-R34}$-GFP-Flag pcDNA3 construct was made from PCR amplifying GFP-Flag from

pcDNA3 with a forward primer encoding the M12-R34 sequence. All SeV-related

constructs were cloned into pCDNA3 using flanking SacII/PacI sites. Wild-type VV Flag-

A51R and Flag-A51R$^{158-162Ala}$ open reading frames were cloned into the SacII/PacI sites

of p166 by SacII/PacI digest of Flag-A51R and Flag-A51R$^{158-162Ala}$ pcDNA3 vectors [6].

The HA-hSpt16 and HA-hSpt16^{ΔNLS} pEZ-M06 expression vectors have been described

[6]. The Flag-GFP pcDNA3 and p166 vectors have been described [6, 9]. All genes were

sequence verified. Expression plasmids were transfected into indicated mammalian cell

lines as previously described using Lipofectamine 2000 (Invitrogen) in Opti-MEM I

(Gibco) [9] overnight and then media was replaced with complete culture media.

Plasmids transfected into LD652 cells were used FuGENE HD (Promega) (LD652 cells)

[72]. Cells were for the indicated times prior to further processing [e.g. protein extraction

(see below)].

General protein extraction and inhibitor treatments. Cells were washed with

phosphate-buffered saline (PBS) prior to scraping and transfer into a 1.5 mL

microcentrifuge tube for centrifugation at 800 x g at 4°C for 15 min. Cell pellets were

resuspended in either 1x Reporter Lysis Buffer (Promega) containing cOmplete™

EDTA-free Protease Inhibitor Cocktail (Roche) and 1 mM phenylmethylsulfonyl fluoride

(PMSF) and freeze-thawed prior to addition of 5x SDS-PAGE loading buffer (100 mM tris

HCl, pH 6.8, 4% SDS, 12% (v/v) glycerol, 4 mM DTT, 0.02% (w/v) Bromophenol Blue) or

were resuspended in Pierce RIPA buffer (containing cOmplete™ EDTA-free Protease

Inhibitor Cocktail and 1 mM PMSF) prior to addition of 5x SDS-PAGE loading buffer.

Where indicated, cells were treated with 10 µM tannic acid for 4 h prior to protein

harvest. Also, where indicated, bortezomib (Selleck Chemicals) (25 nM), MG132

(Sigma) (40 μM), or DMSO (Corning) (vehicle control) was added to cells 6 h post-

transfection and was kept in media until protein extraction.

Immunoblotting. WCE were boiled for 10 min prior to SDS-PAGE electrophoresis at 50

V for approximately 4 h. Separated proteins were transferred in Towbin Buffer (BioRad)

onto nitrocellulose membranes at 150 mA at 4°C for 90 min and blocked with Odyssey

Blocking Buffer (LI-COR) for 1 h at RT. Membranes were blotted with primary Ab

overnight at 4°C, with actin serving as a loading control. After 3 x 5 min PBS-T (PBS,

0.1% Tween) washes, membranes were incubated in secondary Ab conjugated to an

IRDye (LI-COR) for 1 h, washed 3 x 5 min in PBS-T, then a final 5 min PBS wash.

Membranes were then imaged with an Odyssey Fc Imager (LI-COR).

Immunoprecipitation. Infection-based co-immunoprecipitation experiments were

performed as described with minor modifications [6]. Briefly, 10^6 LD652 cells were

seeded overnight, then infected with ΔA51R^{FA51R} (MOI=25) for 24 h. Prior to cell lysis,

cells were washed with equal volumes of PBS twice. Cells were lysed in IP lysis buffer

(cOmplete™ EDTA-free Protease Inhibitor Cocktail, 1 mM PMSF, 25 mM Tris-HCl at pH

7.4, 150 mM NaCl, 0.5% NP-40) and subjected to shearing and sonication (two-15

second sonications with a 30 second interval). Samples were benzonase (Sigma)

treated (250 units/mL) for 1 h at RT. 10% of "input" was collected, and remaining lysate

was end-over-end incubated with 5 µg of primary Ab overnight at 4°C (rabbit-anti-Flag or

isotype control Ab). Then, lysates were incubated with PureProteome protein A/G

magnetic beads (Sigma) for 1-2 h, extensively washed, and immunoprecipitants eluted

in 60 µl 2x SDS-PAGE loading buffer. IPs of total SUMOylated protein fractions in mock-

, VSV-eGFP, or VSV^{M51R}-eGFP-infected (MOI=10) A549 cells for 12 h used either

SUMO-1 or SUMO-2/3 Ab included with the Signal Seeker SUMOylation 1 or 2/3

Detection Kit (Cytoskeleton) and IPs were carried out according to the manufacturer

instructions [6].

Cell fractionation. Cell fractionation and subsequent densitometric analyses were

conducted as previously described [6]. Briefly, cells were infected at the indicated MOI

and times for the indicated times prior to fractionation using the Subcellular Protein

Fractionation Kit (Thermo Scientific) into either cytosolic or nuclear fractions.

Densitometry-based band quantifications were performed using ImageJ software (NIH,

version 1.51n). It is important to note that because the lysis buffer volume used in the

fractionation procedure is dependent upon the initial cell pellet size (which varied

between infection conditions), and the fractionation buffers were not compatible with

protein quantification assays, we only compared cytosolic versus nuclear distribution of a

particular protein within (and not between) each infection condition [6]. Quantification of

the percentage of total cellular ETS-1 protein present in the nuclear fraction within each

indicated infection condition was calculated by dividing the nuclear fraction ETS-1 band

densitometric values by the sum of total (cytosolic +nuclear) ETS- 1 densitometric

values.

RNAi in cell culture. All siRNAs were obtained from Sigma's pre-designed siRNA

library (see **Table S1)**. The hSpt16 and ETS-1-targeted siRNAs have been previously

validated [6]. Transient siRNA-mediated knockdown was achieved by reverse

transfecting mammalian cells with Lipofectamine 2000 according to manufacturer's

protocol for 48-72 h prior to viral infection [6]. RNAi knockdowns in LD652 cells were

performed as previously described using *in vitro*-transcribed dsRNAs transfected with

Cellfectin II (Invitrogen) in Sf-900 II serum free medium [9]. Transfection medium was

replaced with complete culture medium after overnight culture. Sequences of primers

used to generate dsRNA targeting GFP (control) or *LdSpt16* gene sequences are

in **Table S1**.

Luciferase assays. Luciferase assays in LD652 WCE were performed as previously

described with minor modifications [9, 72]. Briefly, at the indicated times post-infection,

cells were washed in PBS, collected by centrifugation (2000 rpm, 10 min, 4°C) and lysed

in reporter lysis buffer (Promega). Lysates were spotted to 96-well dishes, mixed with

Luciferase Assay Reagent (Promega) and arbitrary light units (LU) were measured using

an FLUOstar Omega Plate reader (BMG Labtech).

Antibodies. Information regarding antibodies used in this study, including their source,

is available in **Table S1**.

Fluorescence microscopy and cell viability assays. At indicated times post-infection,

cells were stained with 200 μL of serum free media containing NucBlue (Thermo Fisher)

for 30 min followed by replacement with 1 mL of PBS. Cells were imaged using a 4X

objective on an EVOS-FL fluorescence microscope (Thermo Fisher) using DAPI and

GFP cubes. Each condition had three replicate wells, and at least four images/well were

collected for analysis. Image analysis was conducted using Fiji (NIH) to quantify the

normalized GFP signal/field by dividing total GFP signal by total DAPI (NucBlue) signals.

Positive signal was determined by using uninfected wells lacking NucBlue stain to set a

minimum threshold in both DAPI and GFP channels that was applied to the remaining

dataset. Fold change in GFP signals were calculated by dividing normalized GFP values

of experimental treatments with normalized GFP values in control treatments (indicated

in each figure). For GFP foci imaging and cell viability analysis of 786-0 cells infected

with VSV$^{\Delta M51}$-GFP, cells were grown to confluency in multi-well dishes, infected with

VSV$^{\Delta M51}$-GFP at the indicated MOIs and treated with 0.3-3µM CBL0137 (or media

alone) 1 hpi using the Agilent Bravo liquid handling system (Agilent, CA, USA). 24 hpi,

GFP images were captured by fluorescence microscopy using an ArrayScan Thermo

Scientific Cellomics) and GFP foci were quantified. Cell viability was assessed 48

hpi using resazurin sodium salt (Sigma-Aldrich) after a 2.5 h incubation using a BioTek

Microplate Reader (BioTek Instruments Inc.).

Immunofluorescence microscopy. For staining cells under mock, VSV-eGFP, or

VSV^{M51R}-eGFP infection conditions, cells were seeded at a density of 30,000 cells per

coverslip, cultured overnight, then infected (MOI=10) for 12 hpi. Cells were fixed with 4%

paraformaldehyde (Sigma), incubated with blocking buffer (PBS with 1% BSA and 0.1%

Triton-X) for 1 h, stained with mouse anti-ETS-1 Ab for 2 h, washed extensively with

blocking buffer, incubated with Alexa Fluor-conjugated secondary Ab for 1h, and then

again washed extensively. Coverslips were then mounted onto glass slides using

ProLong™ Diamond anti-fade with DAPI (Thermo Scientific) and imaged with an

Olympus FV10i confocal laser scanning microscope (version 2.1) equipped with

Olympus Fluoview software (version 4.2a). Images were deconvoluted with CellSens

Imaging software (version 1.18). Similar imaging methods were used for U2OS cells

transiently transfected with VSV M-Flag, VSV M^{M51R}-Flag and ETS-1-HA expression

plasmids, except rabbit anti-Flag and mouse anti-HA primary Ab were used for staining.

Protein sequence alignment and structure prediction

Sequences of VSV M (NCBI accession: 7UMK_B), SeV C' (P04862.3), and HPIV-1 C'

(NP_604434.1) were predicted using AlphaFoldv2 [59] and analyzed using PyMOL **(**The

PyMOL Molecular Graphics System version 1.2r3pre, Schrödinger, LLC**.).** To determine

structural similarity between VSV M, SeV C' and HPIV-1 C', pairwise structural

alignments were conducted using RSCB PDB [60] to assign RMSD and TM scores.

RMSD values of <3 Å and TM scores >0.5 were considered to have similar overall fold,

while pairwise comparisons producing RMSD values >3 Å and TM scores <0.5 were

considered to be structurally unrelated [61].

Statistical analyses. All statistical analyses were conducted using GraphPad Prism

(version 8.0) software and *P* values <0.05 were considered statistically significant.

Sample sizes, statistical tests used, and *P* value information are indicated in the

respective figure or figure legend for each quantitative experiment.

Data Availability

The authors declare that the main data supporting the findings of this study are available

within the article and/or Supplementary Information. Source data will be provided after

initial peer review as per the editor's instructions as Supplementary Information files due

to the size and time required to prepare.

Author Contributions

Conceptualization, E.A., R.A., J.S.D., and D.B.G.; Methodology, E.A., R.A., J.S.D., and

D.B.G.; Investigation, E.A., D.S., A.E., R.A., M.M.S., and D.B.G.; Data Curation, E.A.,

D.S, A.E., R.A., M.M.S., and D.B.G.; Writing—original draft, E.A. and D.B.G.; Writing—

Review & Editing, E.A., D.S., A.E., R.A., M.M.S., J.S.D., and D.B.G.; Visualization, E.A.,

A.E., R.A., and D.B.G. Supervision, D.B.G., R.A., and J.S.D.; Funding Acquisition, J.S.D

and D.B.G.

Competing Interest Statement

873 The authors declare no competing interests. The funders had no role in the design of the

874 study; in the collection, analyses, or interpretation of data; in the writing of the

875 manuscript; or in the decision to publish the results.

Acknowledgments

879 We thank members of the Gammon lab for helpful discussion regarding the manuscript.

880 We also thank Dr. Beatriz Fontoura (UTSW Medical Center) for advice regarding the

881 manuscript. This work was supported by NIH grants 1R35GM137978 and

882 1R21AI180551, and American Cancer Society Research Scholar grant
883
884 RSG-23-1153296-01-DMC to DBG. EAR and AE were supported by NIH Training Grant
885
886 T32 AI007520.

References

1.   Tenthorey JL, Emerman M, Malik HS. Evolutionary Landscapes of Host-Virus Arms Races. Annu Rev Immunol. 2022;40:271-94. Epub 2022/01/27. doi: 10.1146/annurev-immunol-072621-084422. PubMed PMID: 35080919.

2.   Chan YK, Gack MU. Viral evasion of intracellular DNA and RNA sensing. Nat Rev Microbiol. 2016;14(6):360-73. Epub 2016/05/14. doi: 10.1038/nrmicro.2016.45. PubMed PMID: 27174148; PubMed Central PMCID: PMCPMC5072394.

3.   Zhu J, Chiang C, Gack MU. Viral evasion of the interferon response at a glance. J Cell Sci. 2023;136(12). Epub 2023/06/21. doi: 10.1242/jcs.260682. PubMed PMID: 37341132; PubMed Central PMCID: PMCPMC10411950.

4.   Nan Y, Nan G, Zhang YJ. Interferon induction by RNA viruses and antagonism by viral pathogens. Viruses. 2014;6(12):4999-5027. Epub 2014/12/17. doi: 10.3390/v6124999. PubMed PMID: 25514371; PubMed Central PMCID: PMCPMC4276940.

5.   Beachboard DC, Horner SM. Innate immune evasion strategies of DNA and RNA viruses. Curr Opin Microbiol. 2016;32:113-9. Epub 2016/06/12. doi: 10.1016/j.mib.2016.05.015. PubMed PMID: 27288760; PubMed Central PMCID: PMCPMC4983539.

6.   Rex EA, Seo D, Chappidi S, Pinkham C, Brito Oliveira S, Embry A, et al. FEAR antiviral response pathway is independent of interferons and countered by poxvirus proteins. Nat Microbiol. 2024;9(4):988-1006. Epub 2024/03/28. doi: 10.1038/s41564-024-01646-5. PubMed PMID: 38538832.

7.   Winkler DD, Luger K. The histone chaperone FACT: structural insights and mechanisms for nucleosome reorganization. J Biol Chem. 2011;286(21):18369-74. Epub 2011/04/02. doi: 10.1074/jbc.R110.180778. PubMed PMID: 21454601; PubMed Central PMCID: PMCPMC3099653.

8.   Duina AA. Histone Chaperones Spt6 and FACT: Similarities and Differences in Modes of Action at Transcribed Genes. Genet Res Int. 2011;2011:625210. Epub 2011/01/01. doi: 10.4061/2011/625210. PubMed PMID: 22567361; PubMed Central PMCID: PMCPMC3335715.

9.   Gammon DB, Duraffour S, Rozelle DK, Hehnly H, Sharma R, Sparks ME, et al. A single vertebrate DNA virus protein disarms invertebrate immunity to RNA virus infection. Elife. 2014;3. Epub 2014/06/27. doi: 10.7554/eLife.02910. PubMed PMID: 24966209; PubMed Central PMCID: PMCPMC4112549.

10. Liu G, Cao W, Salawudeen A, Zhu W, Emeterio K, Safronetz D, et al. Vesicular Stomatitis Virus: From Agricultural Pathogen to Vaccine Vector. Pathogens. 2021;10(9). Epub 2021/09/29. doi: 10.3390/pathogens10091092. PubMed PMID: 34578125; PubMed Central PMCID: PMCPMC8470541.

11. Lawson ND, Stillman EA, Whitt MA, Rose JK. Recombinant vesicular stomatitis viruses from DNA. Proc Natl Acad Sci U S A. 1995;92(10):4477-81. Epub 1995/05/09. doi: 10.1073/pnas.92.10.4477. PubMed PMID: 7753828; PubMed Central PMCID: PMCPMC41967.

12. Whelan SP, Ball LA, Barr JN, Wertz GT. Efficient recovery of infectious vesicular stomatitis virus entirely from cDNA clones. Proc Natl Acad Sci U S A. 1995;92(18):8388-92. Epub 1995/08/29. doi: 10.1073/pnas.92.18.8388. PubMed PMID: 7667300; PubMed Central PMCID: PMCPMC41162.

13. Abdelmageed AA, Ferran MC. The Propagation, Quantification, and Storage of Vesicular Stomatitis Virus. Curr Protoc Microbiol. 2020;58(1):e110. Epub 2020/08/25. doi: 10.1002/cpmc.110. PubMed PMID: 32833351; PubMed Central PMCID: PMCPMC7449588.

14. Suder E, Furuyama W, Feldmann H, Marzi A, de Wit E. The vesicular stomatitis virus-based Ebola virus vaccine: From concept to clinical trials. Hum Vaccin Immunother. 2018;14(9):2107-13. Epub 2018/05/15. doi: 10.1080/21645515.2018.1473698. PubMed PMID: 29757706; PubMed Central PMCID: PMCPMC6183239.

15. Case JB, Rothlauf PW, Chen RE, Kafai NM, Fox JM, Smith BK, et al. Replication-Competent Vesicular Stomatitis Virus Vaccine Vector Protects against SARS-CoV-2-Mediated Pathogenesis in Mice. Cell Host Microbe. 2020;28(3):465-74 e4. Epub 2020/08/18. doi:

10.1016/j.chom.2020.07.018. PubMed PMID: 32798445; PubMed Central PMCID:
PMCPMC7391951.
16.   Holbrook MC, Goad DW, Grdzelishvili VZ. Expanding the Spectrum of Pancreatic
Cancers Responsive to Vesicular Stomatitis Virus-Based Oncolytic Virotherapy: Challenges and
Solutions. Cancers (Basel). 2021;13(5). Epub 2021/04/04. doi: 10.3390/cancers13051171.
PubMed PMID: 33803211; PubMed Central PMCID: PMCPMC7963195.
17.   Felt SA, Grdzelishvili VZ. Recent advances in vesicular stomatitis virus-based oncolytic
virotherapy: a 5-year update. J Gen Virol. 2017;98(12):2895-911. Epub 2017/11/17. doi:
10.1099/jgv.0.000980. PubMed PMID: 29143726; PubMed Central PMCID: PMCPMC5845697.
18.   Selman M, Ou P, Rousso C, Bergeron A, Krishnan R, Pikor L, et al. Dimethyl fumarate
potentiates oncolytic virotherapy through NF-kappaB inhibition. Sci Transl Med. 2018;10(425).
Epub 2018/01/26. doi: 10.1126/scitranslmed.aao1613. PubMed PMID: 29367345.
19.   Stojdl DF, Lichty BD, tenOever BR, Paterson JM, Power AT, Knowles S, et al. VSV
strains with defects in their ability to shutdown innate immunity are potent systemic anti-cancer
agents. Cancer Cell. 2003;4(4):263-75. Epub 2003/10/31. doi: 10.1016/s1535-6108(03)00241-1.
PubMed PMID: 14585354.
20.   Le Boeuf F, Diallo JS, McCart JA, Thorne S, Falls T, Stanford M, et al. Synergistic
interaction between oncolytic viruses augments tumor killing. Mol Ther. 2010;18(5):888-95. Epub
2010/03/18. doi: 10.1038/mt.2010.44. PubMed PMID: 20234341; PubMed Central PMCID:
PMCPMC2890112.
21.   Arulanandam R, Batenchuk C, Varette O, Zakaria C, Garcia V, Forbes NE, et al.
Microtubule disruption synergizes with oncolytic virotherapy by inhibiting interferon translation and
potentiating bystander killing. Nat Commun. 2015;6:6410. Epub 2015/03/31. doi:
10.1038/ncomms7410. PubMed PMID: 25817275.
22.   Alluqmani N, Jirovec A, Taha Z, Varette O, Chen A, Serrano D, et al. Vanadyl sulfate-
enhanced oncolytic virus immunotherapy mediates the antitumor immune response by
upregulating the secretion of pro-inflammatory cytokines and chemokines. Front Immunol.
2022;13:1032356. Epub 2022/12/20. doi: 10.3389/fimmu.2022.1032356. PubMed PMID:
36532027; PubMed Central PMCID: PMCPMC9749062.
23.   Blackham AU, Northrup SA, Willingham M, D'Agostino RB, Jr., Lyles DS, Stewart JHt.
Variation in susceptibility of human malignant melanomas to oncolytic vesicular stomatitis virus.
Surgery. 2013;153(3):333-43. Epub 2012/10/30. doi: 10.1016/j.surg.2012.09.003. PubMed PMID:
23102637; PubMed Central PMCID: PMCPMC3561511.
24.   Rajani KR, Pettit Kneller EL, McKenzie MO, Horita DA, Chou JW, Lyles DS. Complexes
of vesicular stomatitis virus matrix protein with host Rae1 and Nup98 involved in inhibition of host
transcription. PLoS Pathog. 2012;8(9):e1002929. Epub 2012/10/03. doi:
10.1371/journal.ppat.1002929. PubMed PMID: 23028327; PubMed Central PMCID:
PMCPMC3460625.
25.   Faria PA, Chakraborty P, Levay A, Barber GN, Ezelle HJ, Enninga J, et al. VSV disrupts
the Rae1/mrnp41 mRNA nuclear export pathway. Mol Cell. 2005;17(1):93-102. Epub 2005/01/05.
doi: 10.1016/j.molcel.2004.11.023. PubMed PMID: 15629720.
26.   Bishnoi S, Tiwari R, Gupta S, Byrareddy SN, Nayak D. Oncotargeting by Vesicular
Stomatitis Virus (VSV): Advances in Cancer Therapy. Viruses. 2018;10(2). Epub 2018/02/24. doi:
10.3390/v10020090. PubMed PMID: 29473868; PubMed Central PMCID: PMCPMC5850397.
27.   Fenton SE, Saleiro D, Platanias LC. Type I and II Interferons in the Anti-Tumor Immune
Response. Cancers (Basel). 2021;13(5). Epub 2021/04/04. doi: 10.3390/cancers13051037.
PubMed PMID: 33801234; PubMed Central PMCID: PMCPMC7957896.
28.   Kottke T, Tonne J, Evgin L, Driscoll CB, van Vloten J, Jennings VA, et al. Oncolytic
virotherapy induced CSDE1 neo-antigenesis restricts VSV replication but can be targeted by
immunotherapy. Nat Commun. 2021;12(1):1930. Epub 2021/03/28. doi: 10.1038/s41467-021-
22115-1. PubMed PMID: 33772027; PubMed Central PMCID: PMCPMC7997928.
29.   Nguyen TL, Abdelbary H, Arguello M, Breitbach C, Leveille S, Diallo JS, et al. Chemical
targeting of the innate antiviral response by histone deacetylase inhibitors renders refractory
cancers sensitive to viral oncolysis. Proc Natl Acad Sci U S A. 2008;105(39):14981-6. Epub

2008/09/26. doi: 10.1073/pnas.0803988105. PubMed PMID: 18815361; PubMed Central PMCID:
PMCPMC2567479.
30.   Oda N, Abe M, Sato Y. ETS-1 converts endothelial cells to the angiogenic phenotype by
inducing the expression of matrix metalloproteinases and integrin beta3. J Cell Physiol.
1999;178(2):121-32. Epub 1999/02/27. doi: 10.1002/(SICI)1097-4652(199902)178:2<121::AID-
JCP1>3.0.CO;2-F. PubMed PMID: 10048576.
31.   Garcia H, Miecznikowski JC, Safina A, Commane M, Ruusulehto A, Kilpinen S, et al.
Facilitates chromatin transcription complex is an "accelerator" of tumor transformation and
potential marker and target of aggressive cancers. Cell Rep. 2013;4(1):159-73. Epub 2013/07/09.
doi: 10.1016/j.celrep.2013.06.013. PubMed PMID: 23831030; PubMed Central PMCID:
PMCPMC5886782.
32.   Dittmer J. The role of the transcription factor Ets1 in carcinoma. Semin Cancer Biol.
2015;35:20-38. Epub 2015/09/24. doi: 10.1016/j.semcancer.2015.09.010. PubMed PMID:
26392377.
33.   Gasparian AV, Burkhart CA, Purmal AA, Brodsky L, Pal M, Saranadasa M, et al.
Curaxins: anticancer compounds that simultaneously suppress NF-kappaB and activate p53 by
targeting FACT. Sci Transl Med. 2011;3(95):95ra74. Epub 2011/08/13. doi:
10.1126/scitranslmed.3002530. PubMed PMID: 21832239; PubMed Central PMCID:
PMCPMC6281439.
34.   Park YH, Jung HH, Ahn JS, Im YH. Ets-1 upregulates HER2-induced MMP-1 expression
in breast cancer cells. Biochem Biophys Res Commun. 2008;377(2):389-94. Epub 2008/10/15.
doi: 10.1016/j.bbrc.2008.09.135. PubMed PMID: 18851945.
35.   Whitlow ZW, Connor JH, Lyles DS. New mRNAs are preferentially translated during
vesicular stomatitis virus infection. J Virol. 2008;82(5):2286-94. Epub 2007/12/21. doi:
10.1128/JVI.01761-07. PubMed PMID: 18094194; PubMed Central PMCID: PMCPMC2258916.
36.   Kato H, Sato S, Yoneyama M, Yamamoto M, Uematsu S, Matsui K, et al. Cell type-
specific involvement of RIG-I in antiviral response. Immunity. 2005;23(1):19-28. Epub 2005/07/26.
doi: 10.1016/j.immuni.2005.04.010. PubMed PMID: 16039576.
37.   Yu H, Bruneau RC, Brennan G, Rothenburg S. Battle Royale: Innate Recognition of
Poxviruses and Viral Immune Evasion. Biomedicines. 2021;9(7). Epub 2021/08/07. doi:
10.3390/biomedicines9070765. PubMed PMID: 34356829; PubMed Central PMCID:
PMCPMC8301327.
38.   Schoggins JW. Interferon-Stimulated Genes: What Do They All Do? Annu Rev Virol.
2019;6(1):567-84. Epub 2019/07/10. doi: 10.1146/annurev-virology-092818-015756. PubMed
PMID: 31283436.
39.   Mahajan R, Delphin C, Guan T, Gerace L, Melchior F. A small ubiquitin-related
polypeptide involved in targeting RanGAP1 to nuclear pore complex protein RanBP2. Cell.
1997;88(1):97-107. Epub 1997/01/10. doi: 10.1016/s0092-8674(00)81862-0. PubMed PMID:
9019411.
40.   Boulukos KE, Pognonec P, Rabault B, Begue A, Ghysdael J. Definition of an Ets1 protein
domain required for nuclear localization in cells and DNA-binding activity in vitro. Mol Cell Biol.
1989;9(12):5718-21. Epub 1989/12/01. doi: 10.1128/mcb.9.12.5718-5721.1989. PubMed PMID:
2555704; PubMed Central PMCID: PMCPMC363744.
41.   Lulli V, Romania P, Morsilli O, Gabbianelli M, Pagliuca A, Mazzeo S, et al.
Overexpression of Ets-1 in human hematopoietic progenitor cells blocks erythroid and promotes
megakaryocytic differentiation. Cell Death Differ. 2006;13(7):1064-74. Epub 2005/11/19. doi:
10.1038/sj.cdd.4401811. PubMed PMID: 16294212.
42.   Fleyshman D, Prendergast L, Safina A, Paszkiewicz G, Commane M, Morgan K, et al.
Level of FACT defines the transcriptional landscape and aggressive phenotype of breast cancer
cells. Oncotarget. 2017;8(13):20525-42. Epub 2017/04/21. doi: 10.18632/oncotarget.15656.
PubMed PMID: 28423528; PubMed Central PMCID: PMCPMC5400524.
43.   Kim M, Neznanov N, Wilfong CD, Fleyshman DI, Purmal AA, Haderski G, et al.
Preclinical Validation of a Single-Treatment Infusion Modality That Can Eradicate Extremity

Melanomas. Cancer Res. 2016;76(22):6620-30. Epub 2016/11/05. doi: 10.1158/0008-5472.CAN-
15-2764. PubMed PMID: 27680682; PubMed Central PMCID: PMCPMC5609527.
44.   Dermawan JK, Hitomi M, Silver DJ, Wu Q, Sandlesh P, Sloan AE, et al. Pharmacological
Targeting of the Histone Chaperone Complex FACT Preferentially Eliminates Glioblastoma Stem
Cells and Prolongs Survival in Preclinical Models. Cancer Res. 2016;76(8):2432-42. Epub
2016/02/28. doi: 10.1158/0008-5472.CAN-15-2162. PubMed PMID: 26921329; PubMed Central
PMCID: PMCPMC4873320.
45.   Burkhart C, Fleyshman D, Kohrn R, Commane M, Garrigan J, Kurbatov V, et al. Curaxin
CBL0137 eradicates drug resistant cancer stem cells and potentiates efficacy of gemcitabine in
preclinical models of pancreatic cancer. Oncotarget. 2014;5(22):11038-53. Epub 2014/11/18. doi:
10.18632/oncotarget.2701. PubMed PMID: 25402820; PubMed Central PMCID:
PMCPMC4294371.
46.   Zhang J, Cong Q, Rex EA, Hallwachs W, Janzen DH, Grishin NV, et al. Gypsy moth
genome provides insights into flight capability and virus-host interactions. Proc Natl Acad Sci U S
A. 2019;116(5):1669-78. Epub 2019/01/16. doi: 10.1073/pnas.1818283116. PubMed PMID:
30642971; PubMed Central PMCID: PMCPMC6358702.
47.   Suzawa M, Miranda DA, Ramos KA, Ang KK, Faivre EJ, Wilson CG, et al. A gene-
expression screen identifies a non-toxic sumoylation inhibitor that mimics SUMO-less human
LRH-1 in liver. Elife. 2015;4. Epub 2015/12/15. doi: 10.7554/eLife.09003. PubMed PMID:
26653140; PubMed Central PMCID: PMCPMC4749390.
48.   Cureton DK, Massol RH, Saffarian S, Kirchhausen TL, Whelan SP. Vesicular stomatitis
virus enters cells through vesicles incompletely coated with clathrin that depend upon actin for
internalization. PLoS Pathog. 2009;5(4):e1000394. Epub 2009/04/25. doi:
10.1371/journal.ppat.1000394. PubMed PMID: 19390604; PubMed Central PMCID:
PMCPMC2667253.
49.   Mohd-Qawiem F, Nawal-Amani AR, Faranieyza-Afiqah F, Yasmin AR, Arshad SS,
Norfitriah MS, et al. Paramyxoviruses in rodents: A review. Open Vet J. 2022;12(6):868-76. Epub
2023/01/19. doi: 10.5455/OVJ.2022.v12.i6.14. PubMed PMID: 36650879; PubMed Central
PMCID: PMCPMC9805762.
50.   Wang C, Wang T, Duan L, Chen H, Hu R, Wang X, et al. Evasion of Host Antiviral Innate
Immunity by Paramyxovirus Accessory Proteins. Front Microbiol. 2021;12:790191. Epub
2022/02/18. doi: 10.3389/fmicb.2021.790191. PubMed PMID: 35173691; PubMed Central
PMCID: PMCPMC8841848.
51.   Pisanelli G, Pagnini U, Iovane G, Garcia-Sastre A. Type I and Type II Interferon
Antagonism Strategies Used by Paramyxoviridae: Previous and New Discoveries, in Comparison.
Viruses. 2022;14(5). Epub 2022/05/29. doi: 10.3390/v14051107. PubMed PMID: 35632848;
PubMed Central PMCID: PMCPMC9145045.
52.   Morita N, Tanaka Y, Takeuchi K, Kitagawa Y, Sakuma R, Koide N, et al. SeV C Protein
Plays a Role in Restricting Macrophage Phagocytosis by Limiting the Generation of Intracellular
Double-Stranded RNA. Front Microbiol. 2022;13:780534. Epub 2022/03/11. doi:
10.3389/fmicb.2022.780534. PubMed PMID: 35265056; PubMed Central PMCID:
PMCPMC8899396.
53.   Garcin D, Latorre P, Kolakofsky D. Sendai virus C proteins counteract the interferon-
mediated induction of an antiviral state. J Virol. 1999;73(8):6559-65. Epub 1999/07/10. doi:
10.1128/JVI.73.8.6559-6565.1999. PubMed PMID: 10400752; PubMed Central PMCID:
PMCPMC112739.
54.   Grogan CC, Moyer SA. Sendai virus wild-type and mutant C proteins show a direct
correlation between L polymerase binding and inhibition of viral RNA synthesis. Virology.
2001;288(1):96-108. Epub 2001/09/07. doi: 10.1006/viro.2001.1068. PubMed PMID: 11543662.
55.   Marcink TC, Englund JA, Moscona A. Paramyxoviruses: Parainfluenza Viruses. In:
Kaslow RA, Stanberry LR, Powers AM, editors. Viral Infections of Humans: Epidemiology and
Control. New York, NY: Springer US; 2020. p. 1-50.
56.   Lo MK, Sogaard TM, Karlin DG. Evolution and structural organization of the C proteins of
paramyxovirinae. PLoS One. 2014;9(2):e90003. Epub 2014/03/04. doi:

10.1371/journal.pone.0090003. PubMed PMID: 24587180; PubMed Central PMCID:
PMCPMC3934983.
57.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool.
J Mol Biol. 1990;215(3):403-10. Epub 1990/10/05. doi: 10.1016/S0022-2836(05)80360-2.
PubMed PMID: 2231712.
58.   Quan B, Seo HS, Blobel G, Ren Y. Vesiculoviral matrix (M) protein occupies nucleic acid
binding site at nucleoporin pair (Rae1 * Nup98). Proc Natl Acad Sci U S A. 2014;111(25):9127-
32. Epub 2014/06/14. doi: 10.1073/pnas.1409076111. PubMed PMID: 24927547; PubMed
Central PMCID: PMCPMC4078809.
59.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly
accurate protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9. Epub
2021/07/16. doi: 10.1038/s41586-021-03819-2. PubMed PMID: 34265844; PubMed Central
PMCID: PMCPMC8371605 have filed non-provisional patent applications 16/701,070 and
PCT/EP2020/084238, and provisional patent applications 63/107,362, 63/118,917, 63/118,918,
63/118,921 and 63/118,919, each in the name of DeepMind Technologies Limited, each pending,
relating to machine learning for predicting protein structures. The other authors declare no
competing interests.
60.   Bittrich S, Segura J, Duarte JM, Burley SK, Rose Y. RCSB protein Data Bank: exploring
protein 3D similarities via comprehensive structural alignments. Bioinformatics. 2024;40(6). Epub
2024/06/13. doi: 10.1093/bioinformatics/btae370. PubMed PMID: 38870521; PubMed Central
PMCID: PMCPMC11212067.
61.   Xu J, Zhang Y. How significant is a protein structure similarity with TM-score = 0.5?
Bioinformatics. 2010;26(7):889-95. Epub 2010/02/19. doi: 10.1093/bioinformatics/btq066.
PubMed PMID: 20164152; PubMed Central PMCID: PMCPMC2913670.
62.   Oda K, Matoba Y, Sugiyama M, Sakaguchi T. Structural Insight into the Interaction of
Sendai Virus C Protein with Alix To Stimulate Viral Budding. J Virol. 2021;95(19):e0081521. Epub
2021/07/22. doi: 10.1128/JVI.00815-21. PubMed PMID: 34287046; PubMed Central PMCID:
PMCPMC8428403.
63.   Mahon C, Krogan NJ, Craik CS, Pick E. Cullin E3 ligases and their rewiring by viral
factors. Biomolecules. 2014;4(4):897-930. Epub 2014/10/15. doi: 10.3390/biom4040897. PubMed
PMID: 25314029; PubMed Central PMCID: PMCPMC4279162.
64.   Loboda AP, Soond SM, Piacentini M, Barlev NA. Lysine-specific post-translational
modifications of proteins in the life cycle of viruses. Cell Cycle. 2019;18(17):1995-2005. Epub
2019/07/12. doi: 10.1080/15384101.2019.1639305. PubMed PMID: 31291816; PubMed Central
PMCID: PMCPMC6681785.
65.   Her LS, Lund E, Dahlberg JE. Inhibition of Ran guanosine triphosphatase-dependent
nuclear transport by the matrix protein of vesicular stomatitis virus. Science.
1997;276(5320):1845-8. Epub 1997/06/20. doi: 10.1126/science.276.5320.1845. PubMed PMID:
9188527.
66.   Petersen JM, Her LS, Varvel V, Lund E, Dahlberg JE. The matrix protein of vesicular
stomatitis virus inhibits nucleocytoplasmic transport when it is in the nucleus and associated with
nuclear pore complexes. Mol Cell Biol. 2000;20(22):8590-601. Epub 2000/10/25. doi:
10.1128/MCB.20.22.8590-8601.2000. PubMed PMID: 11046154; PubMed Central PMCID:
PMCPMC102164.
67.   Dittmer J. The biology of the Ets1 proto-oncogene. Mol Cancer. 2003;2:29. Epub
2003/09/16. doi: 10.1186/1476-4598-2-29. PubMed PMID: 12971829; PubMed Central PMCID:
PMCPMC194255.
68.   Phan M, Watson MF, Alain T, Diallo JS. Oncolytic Viruses on Drugs: Achieving Higher
Therapeutic Efficacy. ACS Infect Dis. 2018;4(10):1448-67. Epub 2018/08/29. doi:
10.1021/acsinfecdis.8b00144. PubMed PMID: 30152676.
69.   Garrett-Sinha LA. Review of Ets1 structure, function, and roles in immunity. Cell Mol Life
Sci. 2013;70(18):3375-90. Epub 2013/01/05. doi: 10.1007/s00018-012-1243-7. PubMed PMID:
23288305; PubMed Central PMCID: PMCPMC3636162.

70.　Winkler DD, Muthurajan UM, Hieb AR, Luger K. Histone chaperone FACT coordinates
nucleosome interaction through multiple synergistic binding events. J Biol Chem.
2011;286(48):41883-92. Epub 2011/10/05. doi: 10.1074/jbc.M111.301465. PubMed PMID:
21969370; PubMed Central PMCID: PMCPMC3308894.
71.　Diallo JS, Vaha-Koskela M, Le Boeuf F, Bell J. Propagation, purification, and in vivo
testing of oncolytic vesicular stomatitis virus strains. Methods Mol Biol. 2012;797:127-40. Epub
2011/09/29. doi: 10.1007/978-1-61779-340-0_10. PubMed PMID: 21948474.
72.　Embry A, Baggett NS, Heisler DB, White A, de Jong MF, Kocsis BL, et al. Exploiting
bacterial effector proteins to uncover evolutionarily conserved antiviral host machinery. PLoS
Pathog. 2024;20(5):e1012010. Epub 2024/05/16. doi: 10.1371/journal.ppat.1012010. PubMed
PMID: 38753575; PubMed Central PMCID: PMCPMC11098378.
73.　Wu W, Zhang W, Duggan ES, Booth JL, Zou MH, Metcalf JP. RIG-I and TLR3 are both
required for maximum interferon induction by influenza virus in human lung alveolar epithelial
cells. Virology. 2015;482:181-8. Epub 2015/04/17. doi: 10.1016/j.virol.2015.03.048. PubMed
PMID: 25880109; PubMed Central PMCID: PMCPMC4461467.
74.　Loo YM, Fornek J, Crochet N, Bajwa G, Perwitasari O, Martinez-Sobrido L, et al. Distinct
RIG-I and MDA5 signaling by RNA viruses in innate immunity. J Virol. 2008;82(1):335-45. Epub
2007/10/19. doi: 10.1128/JVI.01080-07. PubMed PMID: 17942531; PubMed Central PMCID:
PMCPMC2224404.

1173

1174　**Main Figure Legends**

1175

Fig 1. The FEAR pathway contributes to VSV restriction in human cells and is

strongly activated by VSV^{M51R}-eGFP. (A-B) VSV-eGFP (WT) and VSV^{M51R}-eGFP

(M51R) titers 24 h post-infection (hpi) (MOI=0.001) in A549 cells transfected with

indicated RNAi treatments using two independent siRNAs for either hSpt16 (A) or ETS-1

(B) knockdown. Scram., scrambled siRNA. Data are means $\pm$ SD; n=3. Statistical

significance in A and B were determined by unpaired two-tailed Student's t-tests

between indicated treatments. * = P< 0.05; ** = P< 0.01; ns = not significant. (C)

Immunoblot (IB) of endogenous ETS-1 in A549 whole cell extract (WCE) after infection

with VSV-eGFP (WT) or VSV^{M51R}-eGFP (M51R) (MOI=10). (D) Densitometric

quantification of ETS-1 from multiple IB experiments as in C. Data are means $\pm$ SD; n=3.

Results of unpaired two-tailed Student's t-test between ETS-1 levels in mock WCE and

infected WCE are shown above each bar graph as: * = P< 0.05, ** = P< 0.01, or ns = not

significant. (E) IB of U2OS cell WCE transfected with empty vector (EV) or ETS-1-HA

constructs 48 h post-transfection. (F) Representative fluorescence microscopy images of

U2OS cells transfected as in E and then infected with VSV-eGFP for 16 h. NucBlue was

used to mark cell nuclei and to account for cell number. Scale bar = 500 μm. (G)

Quantification of total GFP signal/field normalized to NucBlue signal for experiments

from F. Data are means ± SD; n=3. Statistical significance was determined by unpaired

two-tailed Student's t-test between indicated treatments. ** = P< 0.01.

1195

Fig 2. VSV, but not VSV^{M51R}, promotes hSpt16^{SUMO} depletion to prevent ETS-1

induction. (A) IB of FACT subunit levels in WCE from A549 cells infected with VSV-

eGFP (WT) or VSV^{M51R}-eGFP (M51R) for indicated hpi (MOI=10). (B-D) Densitometric

quantification of IB experiments in A for hSpt16^{SUMO} (B) SUMOless hSpt16 (C) or SSRP1

levels (D). Data are means ± SD; n=3. In B-D, results of unpaired two-tailed Student's t-

test between protein levels in mock WCE and infected WCE are shown above each bar

graph as: * = P< 0.05, ** = P< 0.01, or ns = not significant. (E-F) IB of

immunoprecipitated SUMO-1- (E) or SUMO-2/3- (F) conjugated protein fractions in WCE

from mock-, VSV-eGFP -, or VSV^{M51R}-eGFP-infected A549 cells (MOI=3) 18 hpi. Cont.,

control IP. RanGAP1 is a known SUMOylated protein and used as a control for

enrichment of SUMOylated fractions [39]. GFP is a marker for infection. (G) IB of

endogenous ETS-1 and hSpt16 levels and in WCE of scrambled- or hSpt16-siRNA-

treated A549 cells 8 hpi with indicated strains (MOI=10).

1209

Fig 3. VSV M is sufficient to promote proteasomal degradation of cytoplasmic

hSpt16^{SUMO} and to block ETS-1 expression. (A) IB of endogenous hSpt16 in 293T

WCE transfected with indicated Flag-tagged VSV protein expression constructs for 24 h.

Empty vector, EV. Asterisks indicate probable VSV M degradation products. (B) IB of

endogenous hSpt16 in 293T WCE transfected with indicated VSV M expression

constructs 24 h post-transfection. EV and Flag-GFP constructs were negative controls.

Asterisks indicate probable VSV M degradation products. (C) Immunofluorescence (IF)

images of U2OS cells transfected with indicated HA-tagged hSpt16 constructs for 24 h.

Scale bar = 5 μm. (D) IB of 293T WCE 24 h post-transfection with indicated HA-tagged

hSpt16 constructs along with either EV, VSV M-Flag (WT), or VSV M^{M51R}-Flag (M51R)

pcDNA3 vectors. Asterisks indicate probable VSV M degradation products. (E-F) 293T

WCE transfected with indicated VSV M expression constructs in the absence or

presence of bortezomib (E) or MG132 (F) proteasome inhibitor treatment. Vehicle

(DMSO) treatments served as negative controls. Asterisks indicate probable VSV M

degradation products. (G) IB of endogenous ETS-1 in U2OS WCE after transfection with

indicated expression constructs for 24 h and then infected with VSV^{M51R}-eGFP (MOI

=10) for 8 h. (H) Densitometric quantification of IB experiments in G. Data are means ±

SD; n=3. (I) IB of endogenous ETS-1 in U2OS WCE after transfection with indicated

expression constructs for 24 h and then infected with ΔA51R VV strain (MOI =10) for 8h.

(J) Densitometric quantification of IB experiments in I. Data are means ± SD; n=4.

Statistical significance in H and J were determined by unpaired two-tailed Student's t-

tests between indicated treatments. ** = P< 0.01; ns = not significant.

1232

Fig 4. VSV M inhibits ETS-1 nuclear accumulation. (A) IF staining of ETS-1 in mock-,

VSV-eGFP (WT)-, or VSV^{M51R}-eGFP (M51R)-infected A549 cells 12 hpi (MOI=10). (B) IB

of A549 cytoplasmic (C) and nuclear (N) fractions 12 hpi with indicated strains (MOI=10).

Vimentin marks cytosolic fractions while histone H4 marks nuclear fractions [6]. (C)

Densitometric quantification of multiple IB experiments as in B expressed as percentage

of total (cytosolic+nuclear fractions) ETS-1 protein present in nuclear fractions. Data are

means ± SD; n=3. Statistical significance was determined by unpaired two-tailed

Student's t-test between indicated treatments. ** = P< 0.01. (D) IF staining of ETS-1 in

mock-, VSV-eGFP-, or VSV^{M51R}-eGFP-infected U2OS cells 12 hpi (MOI=10). (E) IF

staining of U2OS cells 24 h post co-transfection with ETS-1-HA expression vectors and

either EV, VSV M-Flag, or VSV M^{M51R}-Flag expression vectors. Scale bars in A, D, and E

= 5 μm.

1245

Fig 5. Suppression of FEAR pathway activation promotes oncolytic VSV strain

replication in refractory cancer cell lines. (A) VSV$^{\Delta M51}$-GFP titers 24 hpi (MOI=0.01) in

786-0 cells transfected with indicated RNAi treatments. Data are means ± SD; n=3. (B)

IB of 786-0 WCE from experiments in A at 24 hpi. (C) Representative fluorescence

microscopy images 24 hpi with VSV$^{\Delta M51}$-GFP (MOI=0.05) in 786-0 cells in the presence

of indicated doses of CBL0137. (D) Quantification of GFP foci 24 hpi (green line, read on

right y-axis) after VSV$^{\Delta M51}$-GFP infection of 786-0 cells in the absence or presence of

indicated doses of CBL0137. Cell viability 48 h post-treatment (read on left y-axis) of

cultures containing only CBL0137 (blue line) or CBL0137+ VSV$^{\Delta M51}$-GFP (red line) is

also shown. Data are means ± SD; n=3. (E) Titer of VSV$^{\Delta M51}$-GFP-infected 786-0 cells

24 hpi (MOI=0.01) in the presence of indicated doses of CBL0137. Data are means ±

SD; n=3. (F) IB of 786-0 WCE from experiments in E at 24 hpi. (G) Titer of VSV$^{\Delta M51}$-

GFP-infected RPMI 7951 cells 24 hpi (MOI=0.01) in the presence of indicated doses of

CBL0137. Data are means ± SD; n=3. Statistical significance in A, E, G, and H were

determined by unpaired two-tailed Student's t-test between indicated treatments. * = P<

0.05; ** = P< 0.01; ns, not significant.

1262

Fig 6. Abortive VSV replication in *L. dispar* cells can by complemented by VV

A51R expression or by Spt16 RNAi. (A) IB of *L. dispar* Spt16 (LdSpt16) in LD652

WCE in the absence or presence of tannic acid (TA) treatment. (B) Co-IP of endogenous

LdSpt16 with FA51R in ΔA51R^{FA51R}-infected (MOI=25) LD652 WCE. (C) Fold-change in

luciferase (LUC) signal in LD652 cells transfected with indicated FA51R expression

constructs for 24 h and then infected with VSV-LUC (MOI=10) for 48 h. Fold-change in

LUC is plotted as relative to LUC signal in Flag-GFP (FGFP) transfection treatments

(negative control). Data are means ± SD; n=3. (D) VSV-LUC titers from supernatants

collected from C. Data are means ± SD; n=3. (E) Fold-change in LUC signal after 72 h of

transfection with dsRNAs (for RNAi) targeting either GFP (control) or LdSpt16 (Spt16-

A/B) and infection with VSV-LUC (MOI=10) for 48 h. Data are means ± SD; n=4. (F)

VSV-LUC titers from supernatants collected from E. Data are means ± SD; n=4. (G) IB

of LdSpt16 levels in LD652 WCE 48 h post-transfection with empty vector (EV) or

increasing amounts of VSV M-Flag p166 expression vector. Statistical significance in C,

D, E, and F were determined by unpaired two-tailed Student's t-test between indicated

treatments. ** = P< 0.01; ns, not significant.

1279

Fig 7. Murine and human paramyxovirus-encoded C proteins also promote

proteasome-dependent degradation of hSpt16^{SUMO}. (A) SeV-GFP titers 48 hpi

(MOI=0.01) in A549 cells after indicated RNAi treatments. Data are means ± SD; n=3.

(B) IB of FACT subunit levels in mock- or SeV-GFP-infected (MOI=1) A549 WCE at

indicated hpi. GFP is a marker of infection. (C) Schematic illustrating the four accessory

SeV "C" proteins (C'1, C, Y1, and Y2) and their relative translation initiation sites (nts) in

the SeV *P* transcript and resulting protein sizes (a.a.). Image modified from [54]. (D) IB

of endogenous hSpt16 in 293T WCE transfected with indicated expression constructs for

24 h in the absence or presence of bortezomib treatment. Asterisks indicates SeV C'-

Flag degradation product. (E) IB of endogenous hSpt16 in 293T WCE transfected with

indicated expression constructs for 24 h. Asterisks indicates SeV C'/C-Flag degradation

products. (F) IB of endogenous hSpt16 in 293T WCE transfected with indicated

expression constructs for 24 h. (G) HPIV-1-GFP titers 48 hpi (MOI=0.01) in A549 cells

after indicated RNAi treatments. Data are means $\pm$ SD; n=3. (H) IB of FACT subunit

levels in mock- or HPIV1-GFP-infected (MOI=1) A549 WCE at indicated hpi. GFP is a

marker of infection. (I) IB of endogenous hSpt16 in 293T WCE transfected with indicated

expression constructs for 24 h in the absence or presence of bortezomib treatment.

Statistical significance in A and G were determined by unpaired two-tailed Student's t-

test between indicated treatments. * = P< 0.05; ** = P< 0.01; ns, not significant.

1299

Fig 8. Model for DNA and RNA virus-encoded mechanisms to antagonize the

FEAR pathway. Poxvirus A51R proteins tether hSpt16^{SUMO} subunits to microtubules to

prevent FACT complex formation and activation of ETS-1 expression in the nucleus. In

contrast, VSV M and paramyxovirus (SeV, HPIV-1) C proteins promote degradation of

hSpt16^{SUMO} subunits in the proteasome to block ETS-1 expression. This degradation-

based mechanism may involve recruitment of host E3 ubiquitin ligases to target

hSpt16^{SUMO} for degradation. VSV M may also inhibit FEAR pathway activation by

impeding ETS-1 nuclear accumulation (not shown). Model for FEAR pathway is adapted

from [6] and was created using Biorender.com.

1309

1310

1311   **Supplementary Figure Legends**

1312

Fig S1. VSV-induced ETS-1 expression is independent of RIG-I or IFN signaling.

(A, B) Representative IB (A) and quantification (B) of ETS-1 expression using IB of WCE from control or RIG-I knockout (KO) A549 cells infected with the indicated strains (MOI=10) for 8 h. Note: RIG-I is expressed at a low level in uninfected cells and is known to be induced by viral infection [73, 74], consistent with our results. (C, D) Representative IB (C) and quantification (D) of ETS-1 expression using IB of WCE from control or IRF3 KO A549 cells infected with the indicated strains (MOI=10) for 8 h. (E, F) Representative IB (E) and quantification (F) of ETS-1 expression using IB of WCE from control or STAT1 KO A549 cells infected with the indicated strains (MOI=10) for 8 h. Data in B, D, and F are means $\pm$ SD for n=3 experiments. Statistical significance in B, E, and F were determined by unpaired two-tailed Student's t-test between indicated treatments. For brevity, only the most relevant statistical comparisons are shown. * = P< 0.05; ** = P< 0.01; ns, not significant.

1326

Fig S2. The VSV^{M51R}-eGFP strain is unable to deplete hSpt16^{SUMO} regardless of MOI. IB of endogenous hSpt16 in A549 WCE after infection with VSV-eGFP or VSV^{M51R}-eGFP at the indicated MOI. GFP is used as a marker for infection.

1330

Fig S3. A VSV$^{\Delta M51}$ strain is unable to deplete hSpt16^{SUMO}. IB of endogenous hSpt16 in A549 WCE after infection with VSV-GFP (WT) or VSV$^{\Delta M51}$-GFP (ΔM51) (MOI=3). GFP is used as a marker for infection.

1334

Fig S4. VSV M-mediated depletion of hSpt16^{SUMO} is independent of Rae1 and Nup98. IB of endogenous hSpt16 in A549 whole cell extract (WCE) 72 h after RNAi of

1337   Rae1 (A) or Nup98 (B) under mock- or VSV-eGFP-infection (MOI = 10) conditions for 12

1338   h. VSV N and GFP are markers for infection. Scram., scrambled.

1339

1340   **Fig S5. Human and *Lymantria dispar*-encoded Spt16 proteins share a similar**

1341   **A51R-binding domain.** Alignment of the VV A51R-binding domain of hSpt16 [6]

1342   (accession: NP_009123.1) with corresponding region in LdSpt16 (accession:

1343   GCA_004115105.1). The overall amino acid identity of these regions is 72.8%.

1344   Alignment was made using NCBI Blast2p Software.

1345

1346   **Fig S6. SeV C' promotes depletion of SUMOylated Spt16 subunits present in the**

1347   **cytosol**. IB of HA-hSpt16 or HA-hSpt16$^{\Delta NLS}$ in 293T WCE 24 h after transfection with

1348   empty vector (EV) and or SeV C'-Flag expression constructs.

1349

1350   **Fig S7. SeV C' promotes depletion of SUMOylated Spt16 subunits in mouse 3T3**

1351   **cells.** IB of endogenous mouse Spt16 in 3T3 WCE 24 h after transfection with empty

1352   vector (EV) and or SeV C'-Flag expression constructs.

1353

Fig S8. VSV M protein structural modeling. (A) VSV M protein map showing region

surrounding M51 residue (red). NTE = N-terminal extension. Protein map based on

reference [58]. B) 4OWR structure of VSV M fragment (a.a. 44-229) bound to Rae1-

Nup98 complex [58]. (C) Alphafold2-generated structure of full-length (a.a. 1-229) VSV

M (strain Indiana). (D) Overlay of structures from B and C. E) Results of pairwise

comparison of structures in B and C showing RMSD (in Å) and TM values.

1360

Fig S9. Paramyxovirus C protein structural modeling. (A) Protein maps of SeV C'

and HPIV-1 C' showing region in SeV C' N-terminus sufficient for hSpt16^{SUMO}

degradation (a.a. 12-34) aligned with corresponding region in HPIV-1 C'. NTE = N-

terminal extension. Residues predicted to form alpha helical structures in the NTE are in

red and purple for SeV C' and HPIV-1 C', respectively. (B) 6KP3 structure of SeV C'

fragment (a.a. 99-204) bound to cellular Alix protein [62]. (C) Alphafold2-generated

structure of full-length (a.a. 1-215) SeV C'. (D) Overlay of structures from B and C. (E)

Results of pairwise comparison of structures in B and C showing RMSD (in Å) and TM

values. (F) Alphafold2-generated structure of full-length (a.a. 1-219) HPIV-1 C'. (G)

Overlay of structures from C and F. (H) Results of pairwise comparison of structures in B

and C showing RMSD (in Å) and TM values.

1372

1373 **Fig S10. Structural comparison of VSV M to paramyxovirus C proteins.** (A) Overlay

1374 of Alphafold2-generated VSV M (Fig S8C), SeV C' (Fig S9C), and HPIV-1 C' (Fig S9F)

1375 structures. (B) Results of pairwise comparison of VSV M to either SeV C' or HPIV-1 C'

1376 structures showing RMSD (in Å) and TM values.

1377
1378 **Supplementary Tables**
1379
1380 **Table S1. Key Experimental Reagents.**

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure S1

Figure S2

Figure S3

Figure S4

```
hSpt16  758 HDRDDLYAEQMEREMRHKLKTAFKNFIEKVEALTKEELEFEVPFRDLGFNGAPYRSTCLLQPTSSALVNATEWPPFVVTLDEVELI
LdSpt16 761 HDRDDLAAEQSERELRHKLKVAFKSFCERVETMTKQEVEFDTPFRELGFPGAPFRSTVLLQPTSGALVNLTEWPPFVIALEDVELV

hSpt16  844 HFERVQFHLKNFDMVIVYKDYSKKVTMINAIPVASLDPIKEWLNSCDLKY
LdSpt16 846 HFERVQFHLKNFDMVFVFKDYAKKVAMVNAVPMNMLDHVKEWLNSCDIRY
```

Figure S5

Figure S6

Figure S7

E

RCSB PDB Pairwise Structural Alignment

Entry		Length	RMSD	TM-Score	Identity	Aligned Residues
🟩	VSV M	1-229	0	1	100%	229/229
🟨	4OWR	42-229	1.82	0.93	97%	167/188

Figure S8

A

SeV C': NTE | α1 | | Globular Domain

1 14   34   88   215

```
(12)MPSFLKKILKLRGRRQEDESRSRMLSDSSML(42)
    MPSFL+ ILK + R  E+++ S+M SDS
(16)MPSFLRGILKPKERHHENKNHSQMSSDSLTS(46)
```
Per.
Iden.
;69%

HPIV C': NTE | α1 | | Globular Domain

B   Alix

C

D

E

RCSB PDB Pairwise Structural Alignment

Entry	Length	RMSD	TM-Score	Identity	Aligned Residues
🟩 SeV C'	1-215	0	1	100%	215/215
🟪 6KP3	85-215	1.22	0.92	92%	109/119

F

G

H

RCSB PDB Pairwise Structural Alignment

Entry	Length	RMSD	TM-Score	Identity	Aligned Residues
🟩 SeV C'	1-215	0	1	100%	215/215
🟦 HPIV-1 C'	1-219	2.57	0.66	77%	144/219

Figure S9

A

B

RCSB PDB Pairwise Structural Alignment

Entry		RMSD	TM-Score	Identity	Aligned Residues
🟨	VSV M	0	1	100%	229/229
🟩	SeV C'	5.52	0.22	7%	46/216
🟦	HPIV-1 C'	6.45	0.21	5%	43/219

Figure S10