1

Two residues reprogram immunity receptor kinases to signal in nitrogen-fixing symbiosis

Magdalini Tsitsikli[1,2]*, Bine W. Simonsen[1]*, Thi-Bich Luu[1]*, Maria M. Larsen[1], Camilla G. Andersen[1],

Kira Gysel[1], Damiano Lironi[1,3], Christina Krönauer[1,4], Henriette Rübsam[1,2], Simon B. Hansen[1,5], René

Bærentsen[1], Jesper Lundsgaard Wulff[1], Sarah Holt Johansen[1], Gülendam Sezer Kaya[1], Jens

Stougaard[1], Kasper Røjkjær Andersen[1,#], Simona Radutoiu[1,#]

[1]Department of Molecular Biology and Genetics, Aarhus University, 8000 Aarhus C, Denmark.

[2]Present address: Albert-Ludwigs University Freiburg, Institute of Biology II, 79104 Freiburg,
Germany.

[3]Present address: FMC, Genvej 2, 2970 Hørsholm, Denmark.

[4]Present address: Hopfenforschungszentrum Hüll, Hüll 5 1/3, 85283 Wolnzach, Germany.

[5]Present address: Department of Biochemistry, University of Zürich, CH-8057 Zürich, Switzerland.

*Joint first authorship.

[#]Corresponding authors.

Kasper Røjkjær Andersen: kra@mbg.au.dk

Simona Radutoiu: radutoiu@mbg.au.dk

Abstract

Receptor signalling determines cellular responses and is crucial for defining specific biological
outcomes. In legume root cells, highly similar and structurally conserved chitin and Nod factor
receptor kinases activate immune or symbiotic pathways, respectively, upon perception of chitinous
ligands. Here, we show that specific amino acid residues in the intracellular part of the Nod factor
receptor NFR1 determine signalling specificity and enable the distinction between immune and
symbiotic responses. Functional investigation of CERK6, NFR1 and receptor variants hereof revealed
a conserved motif that we term *Symbiosis Determinant 1* in the juxtamembrane region of the kinase
domain that is key for symbiotic signalling. We demonstrate that two residues in *Symbiosis*
Determinant 1 are indispensable hallmarks for NFR1-type receptors and are sufficient to convert
Lotus CERK6 and barley RLK4 kinase outputs to enable symbiotic signalling in *Lotus japonicus*.

Main

Receptor kinases initiate and regulate diverse signalling events in eukaryotic cells in response to extracellular signals[1,2]. Evolution has diversified the extracellular, ligand-binding domains of these receptors while retaining the coupling to intracellular kinases to maintain the catalytic function[3-5]. Kinases share structurally similar domains but differ greatly in their determinants of specificity mediating the activation of specific downstream signalling pathways[6,7]. Identification and prediction of determinants of signalling specificity has proven a challenge. In plants, Chitin Elicitor Receptor Kinases (CERK) recognize chitin molecules and initiate an immune response[8-10], and facilitate symbiosis with arbuscular mycorrhizal fungi[11-16]. Legume plants have evolved an additional, but highly similar NFR1 receptor to perceive decorated chitin ligands called Nod factors, produced by symbiotic bacteria to initiate nitrogen-fixing symbiosis[17,18]. Studies based on receptor mutants determined that in legumes, the chitin-triggered immunity and nitrogen-fixing symbiosis are genetically and functionally separated[8,12] even though both receptors can be expressed in the same cell[19] (Fig. 1a). This suggests that the receptors themselves encode signalling specificity. Therefore, LysM receptor kinases offer an attractive test system for identifying determinants of specificity to understand how similar kinases control divergent signalling pathways.

Previous studies of CERK6 and NFR1 from *Lotus japonicus* (*Lotus*), and LYK3 (the NFR1 homologue) from *Medicago truncatula* (*Medicago*) identified that their ligand affinity and signalling ability depend on distinct regions present in their ectodomains distinguishing chitin and Nod factors[5]. These studies also found that while the ectodomains are in place to recognize the correct extracellular signal, the nature of the cellular pathways is dependent on residues present in the intracellular regions of NFR1 and CERK6 receptors. The CERK6 kinase domain can activate symbiosis only in the presence of the transmembrane-juxtamembrane (TM-JM) region from NFR1. The NFR1 kinase, on the other hand, is not able to activate immune related ROS production after chitin exposure irrespective of the origin of the TM-JM region[5]. This indicates that root nodule symbiosis is controlled independently by specific residues present in the TM-JM region or in the kinase of NFR1, while chitin immunity is controlled by residues present in the CERK6 kinase. The regions in the intracellular part of NFR1 and CERK6 that control this divergent signal output are currently unknown. To discover the specificity determining regions, we performed a detailed functional investigation of chimeric NFR1-CERK6 receptors. Crystal structures informed the design of receptor variants that were functionally tested in *nfr1*, Nod factor insensitive, or in *cerk6*, chitin insensitive loss-of-function mutant plants by expression from their native promoters. We identified key regions and residues that control signalling specificity and demonstrated that engineering symbiotic functionalities into chitin receptors from legume and non-legume plants is possible.

Symbiosis Determinant 1 defines the signalling specificity of NFR1

Residues present in the transmembrane and juxtamembrane regions have been identified to play

distinct regulatory roles for the core kinases of single-pass receptors[20,21]. We aimed to identify

residues controlling the specificity of signalling for nitrogen-fixing symbiosis in NFR1[5]. For this, we

created a series of fifteen receptor variants where the NFR1 ectodomain was coupled to the CERK6

core kinase via different configurations of the transmembrane (TM) and juxtamembrane (JM) (Fig.

1b, c, receptors 2-16) and investigated their function *in vivo* compared to NFR1 (receptor 1). The first

results confirmed that the CERK6 kinase could initiate signalling for root nodule formation and

infection only when preceded by the TM-JM of NFR1 (Fig. 1b, receptor 2), but not by the TM-JM of

CERK6 (Fig. 1b, receptor 3)[5]. A region spanning 23 residues located in the C-terminal part of the JM

adjacent to the kinase domain was found to be critical for the symbiotic proficiency of these

receptors (Fig. 1b, c). All variants with the NFR1 version of this region (receptors 11-16) induced

formation of infected root nodules expressing the symbiotic marker *Nin*[22](Fig. 1b, c, Fig. S1). By

contrast, all receptors with the corresponding version of CERK6 (receptors 3-10) were unable to

induce nodule organogenesis or *Nin* expression in *nfr1* roots, irrespective of the origin of the

remaining regions of the JM domain (Fig. 1b, c, S1). We named this region of NFR1 the Symbiosis

Determinant 1 (SD1). To verify protein integrity, we expressed the receptor variants in tobacco

leaves or *Lotus* root protoplasts and found them to be localized at the plasma membrane, indicating

that receptors which are symbiotically inactive are expressed and correctly localized (Fig. S2, S3).

Next, we investigated the activation of the symbiotic epidermal program characterized by rhizobial

infection via root hair deformation, microcolony and infection thread formation, as well as the early

activation of *Nin* promoter in the epidermis. We inspected the root hairs of *nfr1* expressing key

receptor variants (receptors 1, 2, and 6) analysed for nodule organogenesis (Fig. 1b). Roots

expressing the full-length NFR1 (receptor 1) had typical root hair curling, bacteria entrapment, and

long, elongated infection threads (Fig. 1d, e, S4). Such elongated infection threads were also

identified in roots expressing the symbiotically functional receptor 2 (Fig. 1b, d, e), albeit in a

significantly reduced number, while none were detected for the nonfunctional receptor 6 (Fig. 1b, d,

e). Importantly, roots expressing receptors 2 and 6 responded to the presence of bacteria with root

hair deformation and microcolony formation even when intracellular root hair infection was not

enabled (Fig. S4). This shows that receptor 6 which recognizes the Nod factors via the NFR1

ectodomain initiates only root hair deformation responses in the *nfr1* mutant, but that key events

leading to symbiosis, such as the *Nin* promoter activation in the epidermis, intracellular infection,

and nodule organogenesis, fail to be activated (Fig.1e, S4). The lack of symbiotic functionality was

further confirmed in an independent experiment where the synthetic coupling of receptor 3 to NFR5

using the nanobody-mediated receptor complex formation strategy[23] showed an inability to induce

spontaneous nodule formation (Fig. 1f- receptor 3, 17). However, this was reconstituted if the SD1

was present (Figure 1f- receptor 13) demonstrating the key role of this region in activating the

symbiotic signalling from a heteromeric complex with NFR5. Together, our studies demonstrate that

SD1 controls root nodule symbiosis ensuring activation of both organogenesis and root hair infection

downstream of receptor complex formation.

113   The region containing SD1 is a surface exposed motif in the kinase domain

The region containing SD1 has an amino acid composition that distinctly separates Nod factor

receptors from their chitin-binding counterparts and other members of this protein family in legume

plants (Fig. S5 and Supporting Material 1). To understand the molecular details of SD1, we expressed

and purified the intracellular domains of CERK6 and NFR1 (Fig. S6) for structural investigation. The

purified proteins were functional and able to bind ATP (Fig. S7). We obtained well-diffracting crystals

for CERK6 when an inhibitory mutation, D460N, in the DFG motif was introduced, but this was not

the case for the tested NFR1 versions. To obtain crystals of a symbiotic kinase, we turned to the

NFR1 orthologue from *Medicago*, LYK3, which crystalized with the same inhibitory mutation, D459N,

as CERK6 (Fig. S8). The atomic structures of CERK6 and LYK3 kinases were determined to 1.85 Å and

2.5 Å resolution, respectively (Table S1). Both LYK3 and CERK6 showed a classical fold with clearly

defined N- and C-lobes and hallmark structural features of active kinases (Fig. 2a, b). LYK3 and CERK6

are both crystallized in an inactive conformation closely resembling the conformation of the

CDK2/Src inactive kinase, which is an autoinhibition state of the kinase, where the αC is in the "out"-

position, while the aspartate of the DFG motif points towards the nucleotide binding side. LYK3, but

not CERK6, was crystallized with a bound ATP analogue (Fig. S8a). However, despite this, the two

structures show an overall high similarity (RMSD=0.724 Å) (Fig. S8b). The SD1 region is clearly

defined in the density map of the LYK3 structure as a loop followed by the αB helix, which is a

conserved feature of IRAK/Pelle-type kinases. SD1 is kept in place by specific residues that are

conserved between chitin and Nod factor receptors, stabilizing the interactions to the N-lobe of the

kinase (Fig. 2c). SD1 contains only six residues differing between CERK6 and NFR1, three of these are

in the N-terminal loop, while the remaining three are on the αB helix. All six defining residues in SD1

are surface exposed making this motif accessible for interactions (Fig. 2d, e).

The core kinase domains are important for the two diverging signalling pathways

Using the available structural information for the NFR1 and CERK6 receptors, we next aimed to

identify which regions or residues in the core kinases control the specificity of the two signalling

pathways. Structural overlay revealed that the differing residues between CERK6 and NFR1 were

dispersed across both the N- and C-lobes and were primarily localized on the surface (Fig. S9), likely

to facilitate protein-protein interactions[6]. Knowing that core kinases of NFR1 and CERK6 can initiate

symbiosis or immunity independently of the TM-JM regions[5], underscores the importance of

diverging, surface-localized residues contribute to signalling specificity. To identify contributing

surfaces, we investigated which residues of the NFR1 core kinase differing from CERK6 can drive

symbiosis independent of SD1. We identified seven regions (A to G) containing multiple different

divergent residues that were structurally clustered, defining potential interaction surfaces (Fig. S9,

S10). These were exchanged individually from CERK6 to NFR1 in receptor 6, which was symbiotically

nonfunctional (Fig. 1b, Fig. S10). None of the surfaces investigated alone was able to activate root

nodule formation in *nfr1*, indicating a possible collective contribution (Fig. S10). We next tested if

their combined differences might be a limiting factor for symbiosis establishment (Fig. 3a). Receptor

20, where both the αC helix (region E) and the activation loop (region A) of CERK6 were exchanged

to those of NFR1, was indeed found to be able to activate the root hair symbiotic program (Fig. S11)

and to induce nodule formation on *nfr1* roots, albeit with very low efficiency, when compared to

functional receptors 1, and 18 containing the full-length NFR1, or the entire NFR1 kinase (Fig. 3a).

This provides further indications that multiple surfaces of the core NFR1 kinase contribute to

symbiosis signalling. Therefore, we tested two variants where the N or C-terminal regions of CERK6

were exchanged to NFR1 in receptor 6 (Fig. 3a receptors 21,22, Fig. S11) without a negative impact

on protein folding and ATP binding activities (Fig. S5). Receptor 22 which has the N-terminal region

of CERK6 and the C-terminal of NFR1 ensured the activation of both epidermal symbiotic signalling

(Fig. S11) and formation of functional nodules more efficiently than receptor 20 where only the αC

helix and the activation loop of CERK6 have been exchanged to NFR1 (Fig. 3a). The opposite variant,

receptor 21 (containing the N-terminal region of NFR1 and the C-terminal of CERK6), did not have

the same effect. The symbiotically functional receptor 22 contains 45 residues diverging from CERK6,

scattered across several surface areas (Fig. S9, 10). This shows that residues from the C-terminal

region of NFR1 create collectively a larger or a particularly well-defined surface enabling symbiosis.

The NFR1 core kinase does not have the ability to activate immunity[5], and our findings show that

residues in its C-terminal region have diverged from CERK6 to enable symbiosis (Fig. 3a, d). This

raises the question of whether the specificity for symbiotic signalling evolved in the core kinase at
the expense of immunity signalling. We investigated this possibility by exchanging the C-terminal
region of the full-length CERK6 with the symbiosis-permissive region of NFR1. This receptor version
expressed in the *cerk6* mutant from the *Cerk6* promoter was able to induce the production of
reactive oxygen species (ROS) after chitin treatment (Fig. 3b, receptor 26), indicating no negative
interference from these NFR1-type residues on this signalling pathway. By contrast, receptor 25
containing the NFR1-type residues in the N-terminus of the kinase failed to complement the mutant
phenotype (Fig. 3b, receptor 25) indicating that the N-terminal region of CERK6 contains
determinants for immune signalling. The C-terminal region of the NFR1 core kinase acts
independently of SD1 in activating symbiosis (Fig. 3a, receptor 22), and importantly, it does not
interfere in immunity. Therefore, we asked if the SD1 in NFR1 may affect the immune signalling from
the CERK6 core kinase. We exchanged the corresponding region with the NFR1 variant within the
full-length CERK6 (Fig. 3c, receptor 23) and found that this receptor variant 23 also induced
production of ROS after chitin treatment when expressed in *cerk6* mutant. Together, these results
demonstrate that neither of the two regions found here to be determinants for symbiosis have a
negative impact on the capacity of CERK6 to activate ROS and that the functionality from the
intracellular region of CERK6 or NFR1 can be independently modulated by residues located in
distinct regions (Fig.3d, e).

Specific residues in SD1 coordinate organogenesis and infection

All land plants contain CERK receptors that activate immunity, but only legumes contain NFR1-type
receptors that activate root nodule symbiosis[24-26]. The identification of two distinct regions in NFR1
extending the molecular functions of CERK6 prompted us to investigate if this can be employed for
engineering of CERKs. The C-terminus of the NFR1 core kinase contains 45 residues diverging from
CERK6 and our extensive analysis of receptor variants revealed a synergetic action of these variable
residues (Fig. 3a). On the other hand, the SD1 contains only 6 residues that differ from CERK6 (Fig.
4a, Fig. S5) and are thus more amenable to engineering. Therefore, we asked if all defining residues
of SD1 are crucial for the symbiotic program. We examined variants of the symbiotically active
receptor 2 where individual variable residues of SD1 were exchanged from NFR1 to CERK6. (Fig. 1b,
4b). All these variants retained the ability to induce nodule formation after inoculation with *M. loti*,
indicating a synergetic contribution to signalling. Closer inspection of the phenotypes, however,
revealed residue-specific variations (Fig. 4b, c, Fig. S12, S13). Receptor 27 (M306T) ensured the
formation of both infected and white nodules albeit on fewer plants (18 out of 24) and in a

significantly reduced number compared to receptor 2 (29 out of 30) (Fig. 4b, c). This indicates an

impact of M306 on the efficiency of symbiotic signalling. Receptor 28 (A308D) had the lowest

symbiotic capacity. Only 15 of the 40 *nfr1* roots expressing this receptor formed infected nodules, a

significantly reduced number when compared to those expressing receptor 2 (Fig. 4b, c, Fig. S12).

This shows that A308 in SD1 is critical for root nodule formation. Nodule organogenesis and rhizobial

infection are well coordinated processes in symbiosis and the phenotypes observed in receptors 27

and 28 imply that variable residues of SD1 may not only control the initiation of symbiotic signalling,

but they may also coordinate organogenesis and infection. To investigate this, we inspected the

transgenic roots in more detail for the expression pattern of the *Nin* promoter providing a better

insight on the early signalling events. No infection threads or epidermal *Nin* activation were detected

at two weeks post inoculation when receptors 27 or 28 were inspected for early symbiotic events

(Fig. S13). At 5 weeks, however we found that *nfr1* roots expressing receptors 28 (A308D), 30

(Q316D) and 31 (K320T) showed activation of the cortical program where a significantly large

number of nodule primordia were present inside the roots, and only a few were developed into

nodules. These primordia were only detectable after monitoring for cortical expression of the *Nin*

promoter (Fig. S12). It is well-established in root nodule symbiosis that numerous primordia are

initiated if there is an impairment of infection[27-30]. Thus, the large number of *Nin*-expressing

organogenetic events initiated by these specific receptors indicates that signalling from NFR1 via SD1

is key for the efficiency of signalling as well as for coordinating organogenesis and infection

programs.

Two residues reprogram chitin receptors to signal in symbiosis

Knowing that a synergetic action of the residues from SD1 is needed, we asked whether minimal

engineering could be achieved to reroute the signalling from the intracellular region of CERK6 for

root nodule symbiosis. As proof-of-concept we targeted the symbiotically nonfunctional receptor 6,

where the SD1 and kinase domain derived from CERK6 (Fig. 1b, c). Changing individual residues in

the SD1 of CERK6 did not reconstitute symbiosis of the *nfr1* mutant (Fig. S14), but T304M and D306A

substitutions together were sufficient to complement nodule formation and rhizobial infection (Fig.

4d, receptor 33) demonstrating that these two residues (methionine and alanine) are the main

drivers of specificity for symbiotic signalling. One additional change, T318K, significantly improved

the efficiency of the symbiotic signalling (Fig. 4d, receptor 34). Next, we attempted to engineer RLK4,

the closest receptor of NFR1 and CERK6 in barley[23]. The kinase of RLK4 is, like CERK6, able to induce

immunity by producing ROS in *cerk6* upon chitin treatment (Fig. S15), but unlike CERK6 it has a very

limited symbiotic signalling capacity when coupled to the ectodomain of NFR1. Eleven out of forty-

three *nfr1* roots expressing receptor 36 with the intracellular region of RLK4 induced a barely

detectable symbiotic activity in the roots and produced only one or two nodules (Fig. 4e, S16, S17,

receptor 36). This is different compared to CERK6 (Fig. 1b- receptor 3). Nonetheless, the residues

within the region of SD1 are conserved between CERK6 and RLK4 (Fig. 4a). Therefore, we asked if

engineering this region in the cereal receptor would have a similar impact on symbiosis signalling.

We created an engineered receptor 37 where the intracellular RLK4 contained SD1. This receptor,

like receptor 13 (SD1 engineered in CERK6), was also able to efficiently initiate root nodule

formation and rhizobial infection when expressed in the *nfr1* mutant (Fig. 4e, S16, S17 receptor 37).

Unlike CERK6, engineering SD1 in the intracellular region of RLK4 abolishes its capacity to activate

ROS production (Fig. S15). Together, these results demonstrate that SD1 provides both *Lotus* CERK6

and barley RLK4 with symbiotic signalling capacity leading to root nodule organogenesis and

intracellular bacterial infection.

Discussion

The ability to detect specific signals present on the cell surface and to respond with the appropriate

internal signalling pathway is important for all life forms. Receptor kinases play a major role in

eukaryotes in relaying the correct information in the cell, and despite high structural

conservation[2,6,31-33], little is known about how signalling specificity is ensured in plants. Here we

investigated the mechanism enabling NFR1 and CERK6 to initiate symbiosis or immune signalling

pathways. NFRs evolved as homologs of CERKs and became neofunctionalized in legumes to enable

root nodule organogenesis and intracellular infection of nitrogen-fixing bacteria[18,25,34,35] (Fig. 3d).

When Nod factor recognition was ensured by the NFR1 ectodomain, the intracellular region of

CERK6 was able to activate root hair deformation, albeit in a rather uncontrolled manner (Figure 1e-

receptors 2 and 6), a phenotype previously observed for the symbiotic *nin* mutants[22]. Thus, it is

conceivable that specific residues have diverged from CERK receptors to enable initiation of the

epidermal infection and cortical root nodule organogenesis. We demonstrate here that two regions,

the SD1 in the juxtamembrane and the C-terminus of the NFR1 core kinase, contain residues that

control the activation of *Nin* and determine symbiosis signalling. These regions can activate

symbiosis independently since the necessity for SD1 can be overcome by NFR1 residues present in

the C-terminal region of the kinase and vice versa (Fig. 1b and Fig. 3a, d). Interestingly, an

intracellular conformation where the C-terminus is of the NFR1-type can serve a dual function,

initiating both symbiosis and immunity, depending on the type of the ectodomain (Fig. 3a, b). This is

reminiscent to a certain extent of the functionality of barley RLK4 which was able, albeit with a very

low efficiency, to induce root nodule formation (Figure 4e). LYK receptors of *Parasponia andersonii*

(Parasponia)[25] and those from plants of the FaFaCuRo clade forming nitrogen fixing symbiosis with

Nod factor-producing Frankia[36] also lack the SD1 (Figure S4). The LYK1, LYK3a and LYK3b from

Parasponia were shown to function in both symbiosis and chitin immunity[25]. SD1 is thus a specific

signature of Nod factor receptors from legumes, and we show here that the motif is necessary and

sufficient (Fig. 4) for the initiation and coordination of root nodule organogenesis and root hair

infection which in legumes runs efficiently. It is thus conceivable that the evolution and maintenance

of SD1 in the legume clade have contributed to the evolution of a very efficient symbiosis in

Papilionoids[37].

The residues in SD1 are surface exposed (Fig. 2d, e), allowing this motif to be accessible for potential

interactions. Detailed inspection of CERK6 and LYK3 crystal structures revealed no evidence of intra-

molecular interactions between variable residues and known regulatory regions of the core kinase

that could explain a direct control over signalling (Fig. 2). However, kinases have a dynamic

structure[33,38,39] and both intra- and intermolecular interactions between protomers or with

interacting partners e.g. NFR5 may occur once the compatible ligand is bound by the ectodomains.

Deciphering the specific contribution of SD1 in NFR1 remains a future challenge due to the inherent

structural plasticity of the juxtamembrane region that includes SD1, the temporally dynamic

landscape of their interacting partners, and the fact that SD1, specifically regulates the signalling

output of the NFR1-NFR5 heteromeric complex (Fig. 1f). The necessity of a large surface in the C-

terminal region of NFR1 kinase corroborates with a scenario whereby multiple surface-exposed

residues contribute to symbiotic signalling possibly by creating more stable protein-protein

interaction interfaces or by enabling dynamic interactions with multiple partners.

Minimal changes were identified here that enabled chitin receptors, CERK6 from *Lotus* and RLK4

from barley, to activate the symbiotic program by inducing expression of *Nin* that controls root hair

infection and nodule organogenesis in the cortex. Changing two residues, T304M and D306A, was

sufficient for the CERK6 kinase to induce the formation of nodules (Fig. 4d). The symbiotic

proficiency was improved by changing T318K and NFR1-levels of nodulation were reached when the

entire SD1 was present (Fig. 1b and Fig. 4d). Importantly, our detailed investigations revealed that

engineering the intracellular regions of chitin receptors extends their functional capacities and that

both symbiotic and immune functionalities can be controlled from a singular version of the

intracellular domain. Together with the opportunity of minimal alterations, we provide the blueprint

for engineering native receptors to support root nodule symbiosis in non-legumes[40].

Acknowledgments and funding

We would like to thank Lene H. Madsen for expertise on gene cloning and Lotus resources, Majken K. Sørensen for plant care and seed production in our greenhouse, and Taylor G. Fitzgerald for proofreading. MT thanks Thomass Ott for support. This work is funded by the Novo Nordisk Foundation (NNF18OC0052855) and the project Enabling Nutrient Symbioses in Agriculture (ENSA), which is funded by Bill & Melinda Gates Agricultural Innovations (INV- 57461), the Bill & Melinda Gates Foundation and the Foreign, Commonwealth and Development Office (INV-55767).

Competing interests

Aarhus University has filed a provisional patent application authored by SR, KRA, MT, BWS, CS, CK, MML, DL, SBH and KG, on using these findings for engineering LysM receptor kinases. The other authors declare no competing interests.

Author contributions:

Conceptualization: SR, KRA.

Methodology: SR, KRA, KG, TBL, CK, DL, MT, BWS, SH.

Investigation: MT, BWS, TBL, CGA, MML, HR, GS, SH, JLW, SHJ.

Visualization: MT, BWS, KRA, KG.

Funding acquisition: SR, KRA.

Project administration: SR, KRA.

Supervision: SR, KRA, KG, TBL, CK, DL.

Writing – original draft: SR.

Writing – review & editing: MT, BWS, TBL, CGA, MML, KG, KRA.

References

1   Lemmon, M. A. & Schlessinger, J. Cell Signaling by Receptor Tyrosine Kinases. *Cell* **141**, 1117-1134 (2010). https://doi.org/10.1016/j.cell.2010.06.011
2   Shiu, S. H. & Bleecker, A. B. Receptor-like kinases from Arabidopsis form a monophyletic gene family related to animal receptor kinases. *Proc Natl Acad Sci U S A* **98**, 10763-10768 (2001). https://doi.org/10.1073/pnas.181141598
3   Zipfel, C. & Oldroyd, G. E. Plant signalling in symbiosis and immunity. *Nature* **543**, 328-336 (2017). https://doi.org/10.1038/nature22009
4   Ngou, B. P. M., Wyler, M., Schmid, M. W., Kadota, Y. & Shirasu, K. Evolutionary trajectory of pattern recognition receptors in plants. *Nat Commun* **15**, 308 (2024). https://doi.org/10.1038/s41467-023-44408-3

5   Bozsoki, Z. *et al.* Ligand-recognizing motifs in plant LysM receptors are major determinants
of specificity. *Science* **369**, 663-670 (2020). https://doi.org/10.1126/science.abb3377

6   Johnson, J. L. *et al.* An atlas of substrate specificities for the human serine/threonine
kinome. *Nature* **613**, 759-766 (2023). https://doi.org/10.1038/s41586-022-05575-3

7   Ma, Y. *et al.* Comparisons of two receptor-MAPK pathways in a single cell-type reveal
mechanisms of signalling specificity. *Nat Plants* **10**, 1343-1362 (2024).
https://doi.org/10.1038/s41477-024-01768-y

8   Bozsoki, Z. *et al.* Receptor-mediated chitin perception in legume roots is functionally
separable from Nod factor perception. *P Natl Acad Sci USA* **114**, E8118-E8127 (2017).
https://doi.org/10.1073/pnas.1706795114

9   Miya, A. *et al.* CERK1, a LysM receptor kinase, is essential for chitin elicitor signaling in. *P
Natl Acad Sci USA* **104**, 19613-19618 (2007). https://doi.org/10.1073/pnas.0705147104

10   Wang, G. *et al.* Release of a ubiquitin brake activates OsCERK1-triggered immunity in rice.
Nature **629**, 1158-1164 (2024). https://doi.org/10.1038/s41586-024-07418-9

11   Zhang, X. W. *et al.* The receptor kinase CERK1 has dual functions in symbiosis and immunity
signalling. *Plant J* **81**, 258-267 (2015). https://doi.org/10.1111/tpj.12723

12   Zhang, J. *et al.* A receptor required for chitin perception facilitates arbuscular mycorrhizal
associations and distinguishes root symbiosis from immunity. *Curr Biol* **34**, 1705-1717 e1706
(2024). https://doi.org/10.1016/j.cub.2024.03.015

13   Han, R. C. *et al.* Wild species ricerice OsCERK1DY-mediated arbuscular mycorrhiza symbiosis
boosts yield and nutrient use efficiency in rice breeding. *Mol Breeding* **44** (2024).
https://doi.org/ARTN 22 10.1007/s11032-024-01459-8

14   Miyata, K. *et al.* OsCERK2/OsRLK10, a homolog of OsCERK1, has a potential role for chitin-
triggered immunity and arbuscular mycorrhizal symbiosis in rice. *Plant Biotechnol-Nar* **39**,
119-+ (2022). https://doi.org/10.5511/plantbiotechnology.21.1222a

15   Feng, F. *et al.* A combination of chitooligosaccharide and lipochitooligosaccharide
recognition promotes arbuscular mycorrhizal associations in. *Nature Communications* **10**
(2019). https://doi.org/ARTN 5047 10.1038/s41467-019-12999-5

16   Liao, D. H., Sun, X., Wang, N., Song, F. M. & Liang, Y. Tomato LysM Receptor-Like Kinase
SlLYK12 Is Involved in Arbuscular Mycorrhizal Symbiosis. *Frontiers in Plant Science* **9** (2018).
https://doi.org/ARTN 1004 10.3389/fpls.2018.01004

17   Limpens, E. *et al.* LysM domain receptor kinases regulating rhizobial Nod factor-induced
infection. *Science* **302**, 630-633 (2003). https://doi.org/10.1126/science.1090074

18   Radutoiu, S. *et al.* Plant recognition of symbiotic bacteria requires two LysM receptor-like
kinases. *Nature* **425**, 585-592 (2003). https://doi.org/10.1038/nature02039

19   Frank, M. *et al.* Single-cell analysis identifies genes facilitating rhizobium infection in Lotus
japonicus. *Nature Communications* **14** (2023). https://doi.org/ARTN 7171 10.1038/s41467-
023-42911-1

20   Red Brewer, M. *et al.* The Juxtamembrane Region of the EGF Receptor Functions as an
Activation Domain. *Mol Cell* **34**, 641-651 (2009).
https://doi.org/10.1016/j.molcel.2009.04.034

21   Zhou, Q. *et al.* The juxtamembrane domains of CERK1, BAK1, and FLS2 play a conserved role
in chitin-induced signaling. *J Integr Plant Biol* **62**, 556-562 (2020).
https://doi.org/10.1111/jipb.12847

22   Schauser, L., Roussis, A., Stiller, J. & Stougaard, J. A plant regulator controlling development
of symbiotic root nodules. *Nature* **402**, 191-195 (1999). https://doi.org/Doi 10.1038/46058

23   Rübsam, H. *et al.* Nanobody-driven signaling reveals the core receptor complex in root
nodule symbiosis. *Science* **379**, 272-277 (2023). https://doi.org/10.1126/science.ade9204

24   Buendia, L., Girardin, A., Wang, T., Cottret, L. & Lefebvre, B. LysM Receptor-Like Kinase and
LysM Receptor-Like Protein Families: An Update on Phylogeny and Functional
Characterization. *Front Plant Sci* **9**, 1531 (2018). https://doi.org/10.3389/fpls.2018.01531

25 Rutten, L. *et al.* Duplication of Symbiotic Lysin Motif Receptors Predates the Evolution of
Nitrogen-Fixing Nodule Symbiosis. *Plant Physiol* **184**, 1004-1023 (2020).
https://doi.org/10.1104/pp.19.01420
26 Miyata, K. *et al.* The Bifunctional Plant Receptor, OsCERK1, Regulates Both Chitin-Triggered
Immunity and Arbuscular Mycorrhizal Symbiosis in Rice. *Plant Cell Physiol* **55**, 1864-1872
(2014). https://doi.org/10.1093/pcp/pcu129
27 Li, X. *et al.* Atypical Receptor Kinase RINRK1 Required for Rhizobial Infection But Not Nodule
Development in Lotus japonicus. *Plant Physiol* **181**, 804-816 (2019).
https://doi.org/10.1104/pp.19.00509
28 Liu, M., Soyano, T., Yano, K., Hayashi, M. & Kawaguchi, M. ERN1 and CYCLOPS coordinately
activate NIN signaling to promote infection thread formation in. *J Plant Res* **132**, 641-653
(2019). https://doi.org/10.1007/s10265-019-01122-w
29 Schiessl, K. *et al.* Recruits the Lateral Root Developmental Program for Symbiotic Nodule
Organogenesis in. *Current Biology* **29**, 3657-+ (2019).
https://doi.org/10.1016/j.cub.2019.09.005
30 Lee, T. *et al.* Light-sensitive short hypocotyl genes confer symbiotic nodule identity in the
legume Medicago truncatula. *Current Biology* **34** (2024).
https://doi.org/10.1016/j.cub.2024.01.018
31 Hohmann, U., Lau, K. & Hothorn, M. The Structural Basis of Ligand Perception and Signal
Activation by Receptor Kinases. *Annu Rev Plant Biol* **68**, 109-137 (2017).
https://doi.org/10.1146/annurev-arplant-042916-040957
32 Chiu, C. H. & Paszkowski, U. Receptor-Like Kinases Sustain Symbiotic Scrutiny. *Plant Physiol*
182, 1597-1612 (2020). https://doi.org/10.1104/pp.19.01341
33 Huse, M. & Kuriyan, J. The conformational plasticity of protein kinases. *Cell* **109**, 275-282
(2002). https://doi.org/Doi 10.1016/S0092-8674(02)00741-9
34 Gust, A. A., Willmann, R., Desaki, Y., Grabherr, H. M. & Nürnberger, T. Plant LysM proteins:
modules mediating symbiosis and immunity. *Trends Plant Sci* **17**, 495-502 (2012).
https://doi.org/10.1016/j.tplants.2012.04.003
35 Murakami, E. *et al.* Epidermal LysM receptor ensures robust symbiotic signalling in Lotus
japonicus. *Elife* **7** (2018). https://doi.org/ARTN e33506 10.7554/eLife.33506
36 Nguyen, T. V. *et al.* An assemblage of Cluster II strains from California contains the canonical
nod genes and also the sulfotransferase gene nodH. *Bmc Genomics* **17** (2016).
https://doi.org/ARTN 796 10.1186/s12864-016-3140-1
37 Griesmann, M. *et al.* Phylogenomics reveals multiple losses of nitrogen-fixing root nodule
symbiosis. *Science* **361** (2018). https://doi.org/ARTN eaat1743 10.1126/science.aat1743
38 Mudumbi, K. C. *et al.* Distinct interactions stabilize EGFR dimers and higher-order oligomers
in cell membranes. *Cell Rep* **43** (2024). https://doi.org/ARTN 113603
10.1016/j.celrep.2023.113603
39 Sheetz, J. B., Lemmon, M. A. & Tsutsui, Y. Dynamics of protein kinases and pseudokinases by
HDX-MS. *Methods Enzymol* **667**, 303-338 (2022).
https://doi.org/10.1016/bs.mie.2022.03.037
40 Jhu, M. Y. & Oldroyd, G. E. D. Dancing to a different tune, can we switch from chemical to
biological nitrogen fixation for sustainable food security? *Plos Biol* **21** (2023).
https://doi.org/ARTN e3001982 10.1371/journal.pbio.3001982
41 Malolepszy, A. *et al.* A plant chitinase controls cortical infection thread progression and
nitrogen-fixing symbiosis. *Elife* **7** (2018). https://doi.org/10.7554/eLife.38874

440   **FIGURES**

Figure 1. Symbiosis Determinant 1 controls symbiosis with rhizobia. **a** *Lotus* root cells expressing *NFR1* (red), *CERK6* (blue), or both (magenta) 10 dpi (UMAP generated by https://lotussinglecell.shinyapps.io/lotus_japonicus_single-cell/). **b** Number of pink or white

nodules[41] formed by *nfr1* roots expressing the indicated receptor variants. Nodules were counted 5

weeks post inoculation with *M. loti DsRed*. Fractions under the boxplots indicate the number of

nodulating plants out of total plants tested. **c** Representative photos of transgenic roots carrying the

indicated receptor variants. *YFP*: Yellow Fluorescent Protein (transformation marker), *M. loti DsRed*

(fluorescent bacteria), *pNIN:GUS*: Bright Field photos after GUS staining. Scale bars: 5 mm. **d** Number

of infection threads (ITs) formed per cm of transgenic root expressing the indicated receptor variant.

ITs counted 3 weeks post inoculation with *M. loti MAFF*. '*n*' indicates the number of analysed plants.

e Photos of transgenic roots expressing the indicated receptors at 3 weeks post inoculation with *M.*

loti MAFF. Red shows *M. loti MAFF* bacteria. Yellow arrowheads indicate branched and scrambled

root hair tips where microcolonies of bacteria are visible. Scale bars: 30 µM. **f** Number of

spontaneous nodules formed on *nfr1* roots expressing the indicated constructs. Fractions under the

boxplots indicate the number of nodulating plants out of total plants tested. In **b**, **d** and **f** lowercase

letters indicate significant differences based on Kruskal-Wallis analysis of variance and post-hoc

analysis (Dunn's test) $p < 0.05$.

Figure 2. Crystal structures of LYK3 and CERK6 kinase domains. Structure of the LYK3 (**a**) and CERK6 (**b**) kinase domains annotated with conserved secondary structural elements (LYK3, PDB-ID: 9GFZ, CERK6 PDB-ID: 9GB9). LYK3 kinase was crystallized with AMP-PNP in the nucleotide binding site. The identified SD1 region is indicated in gray. **c** Structural overlay of the LYK3 and CERK6 showing the region with SD1. The indicated amino acids that interact with the kinase core domain and SD1 are conserved. **d-e** Zoom-in on SD1 of LYK3 and the same region in CERK6 seen from the top showing that this region is surface exposed and accessible. In CERK6, the T304 (dotted line in **e**) is not visible in the structure. In (**d**) residues in red are from LYK3 and those in black are from NFR1.

Figure 3. The kinase N- and C-lobes contain determinants controlling independently the two signalling pathways. a The number of pink or white nodules formed on *nfr1* roots expressing the indicated receptor variants. Nodules were counted 5 weeks post inoculation with *M. loti DsRed*. Fractions under the boxplots indicate the number of nodulating plants out of total plants tested. Lowercase letters indicate significant differences based on Kruskal-Wallis analysis of variance and post-hoc analysis (Dunn's test) p < 0.05. **b-c** Reactive oxygen species (ROS) generated by *cerk6* roots expressing the indicated receptor variants within a period of 30 min after treatment with 0.1 mg/ml chitin. Top: curves show the values of RLU (Relative Luminescence Units) during the 30-min period after chitin treatment. Bottom: boxplots show the maximum value of RLU observed for each sample within the 30-min period after chitin treatment. (100% is the highest value observed in WT expressing the empty vector construct). **d** Schematic representation of the observed functional capacities for the intracellular domains (IC) of the indicated LysM receptors. These capacities are dependent on the capacity of the ectodomains (EC), to recognize and bind Nod factors or chitin[5].

489 **Figure 4. Two residues of Symbiosis Determinant 1 are essential for root nodule symbiosis. a**
490 Amino acid alignment of Symbiosis Determinant 1 (SD1) in NFR1, CERK6 and RLK4. Numbers above
491 the alignment indicate the amino acid positions in NFR1 protein. **c** Representative photos of
492 transgenic roots expressing the indicated receptor variants. YFP: Yellow Fluorescent Protein
493 (transformation marker), *M. loti DsRED* (fluorescent bacteria), *pNIN:GUS*: Bright Field photos after
494 GUS staining. Scale bars: 5 mm. **b**, **d**, and **e** Number of pink or white nodules formed on *nfr1* roots
495 expressing the indicated receptor variants. Nodules were counted 5 weeks post inoculation with *M.*
496 *loti DsRed*. Fractions under the boxplots indicate the number of nodulating plants out of total plants
497 tested. Lowercase letters indicate significant differences based on Kruskal-Wallis analysis of variance
498 and post-hoc analysis (Dunn's test) $p < 0.05$.

499