Dysfunctional mechanotransduction regulates the progression of PIK3CA-driven vascular malformations

Wen Yih Aw[1], Aanya Sawhney[1], Mitesh Rathod[1], Chloe P. Whitworth[1,2], Elizabeth L. Doherty[1], Ethan Madden[2], Jingming Lu[1], Kaden Westphal[1], Ryan Stack[1], and William J. Polacheck[1,3,4,*]

[1]Joint Department of Biomedical Engineering, University of North Carolina at Chapel Hill and North Carolina State University, Chapel Hill, NC and Raleigh, NC, USA

[2]Department of Genetics and Molecular Biology, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC, USA

[3]Department of Cell Biology and Physiology, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC, USA

[4]McAllister Heart Institute, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC, USA

*Correspondence: polacheck@unc.edu

Abstract

Somatic activating mutations in *PIK3CA* are common drivers of vascular and lymphatic malformations. Despite common biophysical signatures of tissues susceptible to lesion formation, including compliant extracellular matrix and low rates of perfusion, lesions vary in clinical presentation from localized cystic dilatation to diffuse and infiltrative vascular dysplasia. The mechanisms driving the differences in disease severity and variability in clinical presentation and the role of the biophysical microenvironment in potentiating progression are poorly understood. Here, we investigate the role of hemodynamic forces and the biophysical microenvironment in the pathophysiology of vascular malformations, and we identify hemodynamic shear stress and defective endothelial cell mechanotransduction as key regulators of lesion progression. We found that constitutive PI3K activation impaired flow-mediated endothelial cell alignment and barrier function. We show that defective shear stress sensing in *PIK3CA^{E542K}* endothelial cells is associated with reduced myosin light chain phosphorylation, junctional instability, and defective recruitment of vinculin to cell-cell junctions. Using 3D microfluidic models of the vasculature, we demonstrate that *PIK3CA^{E542K}* microvessels apply reduced traction forces and are unaffected by flow interruption. We further found that draining transmural flow resulted in increased sprouting and invasion responses in *PIK3CA^{E542K}* microvessels. Mechanistically, constitutive PI3K activation decreased cellular and nuclear elasticity resulting in defective cellular tensional homeostasis in endothelial cells which may underlie vascular dilation, tissue hyperplasia, and hypersprouting in *PIK3CA*-driven venous and lymphatic malformations. Together, these results suggest that defective nuclear mechanics, impaired cellular mechanotransduction, and maladaptive hemodynamic responses contribute to the development and progression of *PIK3CA*-driven vascular malformations.

Key words

Vascular malformations, PIK3CA, hemodynamics, mechanotransduction, transmural flow, vascular tissue engineering, organs-on-chip

Introduction

Vascular malformations (VMs) are a class of rare genetic disorders associated with localized developmental abnormalities of venous, arterial, capillary, or lymphatic vessels [1-4]. Histologically, VMs are characterized by complexes of structural lesions with enlarged, irregular lumens that are lined with vascular endothelial cells (ECs) and surrounded by disorganized extracellular matrix [3, 5]. These lesions are often congenital, progress in severity over time, are associated with vascular obstruction and impaired drainage, and can be life threatening. Treatment for VMs, especially for complex malformations that occur in multiple tissues, is difficult and rarely curative, and the standard of care relies on surgery, sclerotherapy, and a limited repertoire of drugs.

VMs are caused by somatic mutations in genes that are involved in vascular development, and these same genes are commonly implicated in malignant tumor angiogenesis [4]. VMs are classified based on blood flow dynamics and are categorized as either slow-flow or fast-flow malformations [3]. The majority of slow-flow malformations, which do not involve an arterial component, are caused by mutations that result in upregulated phosphoinositide 3-kinase (PI3K) and mammalian target of rapamycin (mTOR) signaling [4]. In particular, somatic activating mutations of *PIK3CA* have been identified as the cause of the majority (~80%) of cystic lymphatic malformations, and other pathologies, including generalized lymphatic anomalies, capillary malformations, and venous malformations [2, 6-9]. Several studies have demonstrated that overstimulation of PI3K signaling, independent of genetic mutations in *PIK3CA,* is associated with the pathological development of slow-flow VMs, including cerebral cavernous malformation [10-13] and fast-flow arteriovenous malformations in mouse models of hereditary hemorrhagic telangiectasia [14-16]. Given the implication of PI3K overactivation in both slow- and fast-flow VMs, understanding how hemodynamic forces contribute to the pathogenesis of *PIK3CA*-driven VMs could shed light on novel therapeutic approaches for various vascular anomalies.

PIK3CA encodes the p110α catalytic subunit (PI3Kα) of phosphatidylinositol-3-kinases (PI3K), a lipid kinase that is ubiquitously expressed and essential for regulating effector signaling pathways involved in cell proliferation, metabolism, migration, and survival [17]. At the cellular level, PI3K is activated by receptor tyrosine kinases, focal adhesion kinase, and G-protein coupled receptor signaling [17]. Upon activation, PI3K catalyzes the production of phosphatidylinositol 3,4,5-bisphosphate (PtdIns(3,4,5)P_3 or PIP3) at the plasma membrane, leading to membrane translocation and downstream activation of AKT and mTOR signaling cascades that promote cell growth and survival [17]. PI3K signaling is also involved in the remodeling of the actin cytoskeleton. In particular, activation of PI3Kα has been demonstrated to modulate actomyosin contractility and cell motility in endothelial cells by regulating the activity of myosin light chain phosphatase [18], Rac1- [19], and RhoA-GTPases [20]. In addition to being regulated by growth factors signaling, PI3K functions downstream of the fluid shear stress sensitive mechanosensory complex that consists of platelet endothelial cell adhesion molecule 1 (PECAM-1), vascular endothelial (VE-) cadherin, and vascular endothelial growth factor receptor 2 (VEGFR2) [21-23]. While shear-stress induced activation of PI3K signaling is associated with endothelial cell elongation, alignment, and vasodilatory response, how overactivation of PI3K signaling impacts endothelial mechanotransduction remains unknown.

Multiple studies using mouse models have revealed that PI3Kα plays a crucial role in regulating venous and lymphatic vessel growth and development [6-9, 24-26]. Both the loss of function and inhibition of PI3Kα result in embryonic lethality due to deficient angiogenesis and venous fate

specification [24, 27]. Conversely, expression of gain-of-function $PIK3CA^{H1047R}$ variant of PI3Kα in
the endothelium leads to over-proliferation of venous and lymphatic endothelial cells [25, 26].
Interestingly, expression of the same oncogenic $PIK3CA^{H1047R}$ variant resulted in distinct vascular
overgrowth and malformation defects in blood and lymphatic vessels [25]. Specifically, *PIK3CA*-
driven vascular malformations manifest as locally restricted vessel overgrowth and dilation in
veins and capillaries, but as dense hypersprouting vascular networks in the lymphatics [25].
However, the mechanisms by which one *PIK3CA* activating mutation produces distinct
pathological phenotypes in different vascular beds remain to be investigated. The lack of arterial
involvement in slow-flow VMs, together with common biophysical signatures of tissues
susceptible to lesion formation including compliant extracellular matrix, low rates of perfusion, and
drainage flow, suggest significant contributions of vascular hemodynamics and the biophysical
microenvironment to *PIK3CA*-driven lesion progression.

Previously, we established a microphysiological model of VMs and identified that *PIK3CA*
activating mutations result in elevated Rac1 activity in endothelial cells, which drives lesion
formation through dysregulated cytoskeletal dynamics [19]. In this study, we address the role of
fluid shear stress and the biophysical microenvironment in the development of *PIK3CA*-driven
slow-flow malformations, and we identify hemodynamic shear stress and defective cellular
mechanotransduction as key regulators of malformation progression. We demonstrate that
endothelial cells expressing $PIK3CA^{E542K}$ activating mutation fail to elongate, align, and establish
diffusive barrier function in response to laminar shear stress. We observe increased cellular and
nuclear compliance, and we further identified deficient recruitment of vinculin, a mechanosensitive
adaptor protein, to intercellular adherens junctions in human umbilical vein endothelial cells
(HUVECs) expressing a PIK3CA activating mutation commonly associated with VMs. Additionally,
using tissue-engineered microvessels, we demonstrate that mutant HUVECs form dilated
microvessels that are insensitive to flow interruption and are hypersensitive to basal-to-apical
transmural drainage, resulting in sprouting and invasion responses to transmural flow. These
results demonstrate that altered endothelial cellular and nuclear mechanical properties, as well
as dysfunctional mechanotransduction of hemodynamic stresses, contribute to pathological
vascular overgrowth and hypersprouting in *PIK3CA*-driven VMs.

Results

Constitutive PI3K activation impedes flow-mediated ECs remodeling and alignment to laminar
shear stress.

The development of PIK3CA-driven vascular lesions is restricted to slow-flow regions of the
vasculature [28]. To determine whether VM causal mutations in *PIK3CA* impact
mechanotransduction of hemodynamic shear stress, we generated human umbilical vein
endothelial cells (HUVECs) expressing wild-type *PIK3CA* ($PIK3CA^{WT}$) or *PIK3CA* harboring a
PI3K-activating mutation ($PIK3CA^{E542K}$), as described previously [19], and applied flow via orbital
shaking (**Figure 1**). This approach allowed us to apply a range of shear stresses, both laminar
and oscillatory, to cultured $PIK3CA^{WT}$ and $PIK3CA^{E542K}$ monolayers. At a wall shear stress
magnitude similar to that experienced by venous ECs (~11 dyn/cm^2) [29, 30] near the edge of the
well, $PIK3CA^{WT}$ HUVECs became highly elongated and aligned with the local flow direction
(**Figure 1A-C**). In response to lower magnitude wall shear stress and oscillatory flow toward the
center of the well, $PIK3CA^{WT}$ HUVECs adopted a more isometric morphology (**Figure 1A, C**). In
contrast, $PIK3CA^{E542K}$ HUVECs remained isometric at all magnitudes of wall shear stress (**Figure
1A-C**). We further observed increased cell density in $PIK3CA^{E542K}$ HUVECs across all applied

shear stress magnitudes, indicating impaired flow-mediated quiescence with excess PI3K
activation (**Figure 1D**). Consistent with orbital shaking results, *PIK3CA^E542K* HUVECs remained
misaligned and overcrowded in response to 10 dyn/cm^2 laminar shear stress applied in a Hele-
Shaw flow cell (**Figure 1F**).

To determine if the defective mechanotransduction we observed in blood endothelial cells could
play a role in lymphatic malformation progression, we generated lymphatic endothelial cells
(LECs) expressing either *PIK3CA^WT* or *PIK3CA^E542K* (**Figure S1**). Similar to *PIK3CA^E542K* HUVECs,
we observed that *PIK3CA^E542K* LECs were larger in cell size and more isometric compared to
PIK3CA^WT LECs in response to orbital shaking (**Figure S1**), suggesting that excess PI3K
activation and effector signaling perturb mechanotransduction of hemodynamic shear stress in
both the blood and the lymphatic endothelium. To determine the functional consequences of
cellular misalignment to flow, we measured the permeability of *PIK3CA^WT* and *PIK3CA^E542K*
HUVEC monolayers in response to flow. We cultured *PIK3CA^WT* and *PIK3CA^E542K* HUVEC under
orbital shaking conditions on biotinylated-fibronectin substrates, and permeability was quantified
by fluorescence intensity of Cy3-conjugated streptavidin that bound to the biotinylated surface
(**Figure 1A, E**). In agreement with previous reports [31], we observed improved barrier function
in *PIK3CA^WT* HUVECs at regions corresponding to high laminar shear stress and a gradual
increase in permeability with decreasing wall shear stress magnitude (**Figure 1A, E**).
Interestingly, *PIK3CA^E542K* HUVECs demonstrated a dichotomous relationship between shear
stress magnitude and barrier function, with increased permeability as compared to wild-type cells
at high magnitudes of shear stress and reduced permeability at lower shear stress magnitudes
(**Figure 1A, E**). Collectively, these findings demonstrated that constitutive PI3K activation hinders
flow-mediated endothelial cell alignment and barrier function.

Priming with hemodynamic shear stress prior to PIK3CA^E542K *induction restores cell alignment*
but does not impact permeability.

To establish the temporal relationship between elevated PI3K signaling, defective endothelial cell
remodeling, and loss of vascular barrier function, we generated HUVECs expressing either *tet-*
ON-PIK3CA^WT or *tet-ON-PIK3CA^E542K* in which the expression of WT or mutant variant of *PIK3CA*
can be induced by the addition of doxycycline (**Figure S2A**). Doxycycline induced similar levels
of p110α expression as measured by Western blot, indicating a comparable level of lentivirus
integration and protein overexpression in both cell lines (**Figure S2A**). Upon addition of
doxycycline, we observed increased levels of phospo-AKT1(Ser473) in *tet-ON-PIK3CA^E542K*, but
not in *tet-ON-PIK3CA^WT* endothelial cells, demonstrating activation of PI3K effector signaling in
response to induction (**Figure S2A**). To verify the functional relevance of this construct, we
generated 3D human microvascular networks with *tet-ON-PIK3CA^E542K* HUVECs (**Figure S2B-**
C). We observed that the addition of doxycycline during seeding of *tet-ON-PIK3CA^E542K* HUVECs
accelerated lumenogenesis and resulted in the formation of microvascular networks with enlarged
and irregular lumens (**Figure S2B**), in agreement with our previous study demonstrating that
constitutive activation of PIK3CA^E542K resulted in pathologic network formation [19].

To establish the temporal relationships between dysregulated PI3K activation and
mechanotransduction of hemodynamic shear stress, we induced expression of *PIK3CA^E542K*
either prior to or after the application of flow (**Figure 2A**). When *PIK3CA^E542K* expression was
induced before the application of flow, ECs remained unresponsive to shear stress, as measured
by cell orientation and elongation (**Figure 2A-C**). However, when *PIK3CA^E542K* expression was
triggered one day after flow, flow-conditioned HUVECs remained elongated and aligned parallel

to the flow direction (**Figure 2A-C**). Interestingly, although flow preconditioning rescued cell
morphology and alignment, PI3K activation significantly reduced vascular barrier function
compared with uninduced (no doxycycline) control (**Figure 2D**). Collectively, these results indicate
a mechanistic separation between flow-mediated cell alignment and barrier function, whereby
excessive PI3K signaling may have a more direct and immediate impact on EC adherens junction
stability and resultant permeability.

Defective recruitment of vinculin at adherens junctions in PIK3CA mutant endothelium.

We previously reported the presence of discontinuous VE-cadherin-containing adherens
junctions, decreased cortical actin localization, and increased stress fibers formation in HUVECs
expressing *PIK3CA* activating mutations [19]. These findings suggest that the mechanical
coupling between the cytoskeleton and adhesion complexes may be hampered with constitutive
PI3K activation. VE-cadherin junctions are mechanically connected to the actin cytoskeleton
through the interaction of VE-cadherin with β- and α-catenin [32, 33]. Under mechanical load,
tensile stress unfolds α-catenin, allowing for vinculin binding to α-catenin and subsequent force-
dependent strengthening of cell-cell junctions [34-36]. To address whether the mechanical
coupling between actin cytoskeleton and adhesion complexes could underlie the increased
permeability in mutant endothelium, we investigated the spatial localization of vinculin in flow-
conditioned $PIK3CA^{WT}$ and $PIK3CA^{E542K}$ HUVECs (**Figure 3A-B**). In $PIK3CA^{WT}$ HUVECs, vinculin
localized at the cell cortex and co-localized with VE-cadherin containing cell-cell adhesions and
along actin stress fibers (**Figure 3A-B**). In contrast, $PIK3CA^{E542K}$ HUVECs demonstrated reduced
recruitment of vinculin to intercellular adherens junctions and decreased stress fiber formation as
compared to $PIK3CA^{WT}$ HUVECs (**Figure 3A-B**). Despite significant changes in vinculin
localization, western blot analysis showed no measurable differences in the amounts of VE-
cadherin, β-catenin, and vinculin between control and mutant HUVECs (**Figure 3C-D**).
Interestingly, in probing for changes for overall actomyosin contractility, we observed reduced
level of phosphorylated myosin light chain II (pMLC2) in mutant ECs compared with control ECs
(**Figure 3C-D**). These data demonstrate defective transmission of mechanical forces and
stabilization of cell-cell and cell-matrix adhesion complexes in $PIK3CA^{E542K}$ HUVECs.

$PIK3CA^{E542K}$ HUVECs form dilated microvessels with increased sprouting and reduced traction
forces.

To relate our findings in 2D cell culture to the pathophysiological phenotype of vascular
malformations, we generated and characterized the effect of flow on $PIK3CA^{WT}$ or $PIK3CA^{E542K}$
tissue engineered human microvessels (**Figure 4**). Engineered microvessels were molded using
an acupuncture needle to generate a hollow 160-μm cylindrical channel lined with HUVECs and
fully embedded within 2.5 mg/mL type I collagen hydrogel [31, 37]. Cyclic flow of approximately
1-3 dyn/cm^2 was imparted by culturing microvessels on a laboratory rocker [37]. After two days of
culture under a slow flow environment, $PIK3CA^{WT}$ HUVECs formed uniform microvessels with a
limited number of sprouts and no apparent HUVECs invasion into the subluminal matrix (**Figure**
4 A-B). In contrast, $PIK3CA^{E542K}$ microvessels were significantly dilated and developed numerous
sprouts along the vessel length (**Figure 4 A-B**). The expanded mutant vessel diameter coincides
with an increased number of $PIK3CA^{E542K}$ HUVECs, implicating defective regulation of flow-
mediated quiescence with excessive PI3K signaling. Treatment with rapamycin, an mTOR
inhibitor, or alpelisib, a PIK3CA-specific kinase inhibitor, reduced vessel dilation (**Figure 4 A-B**).
Importantly, when compared to rapamycin, we observed a more effective reduction in number of
sprouts with alpelisib treatment (**Figure 4 A-B**). This observation is consistent with our previous

finding and the notion that in addition to contributing to uncontrolled growth through downstream
mTORC1/2 and growth signaling activation, PI3K hyperactivation alters cytoskeletal dynamics
and mechanical properties of endothelial cells [19].

Given the defective recruitment of vinculin to adherens junctions, increased vascular permeability,
loss of cortical actin organization, and prominent stress fiber formation in *PIK3CA^E542K* HUVECs,
we reasoned that the dilated microvessel phenotype may reflect reduced cell-cell adhesion
strength and altered cell-matrix traction forces in mutant microvessels. To assess changes in cell-
generated forces, we visualized the collagen matrix surrounding the microvessels through
reflectance imaging and compared the displacement of matrix before and after lysing ECs with
detergent (**Figure 4 C-D**). Notably, we observed significantly lower matrix displacement in mutant
microvessels when compared to control (**Figure 4 B-D**), indicating reduced traction forces in
PIK3CA^E542K microvessels.

PIK3CA^E542K microvessels are unresponsive to flow interruption.

To determine whether the slow-flow environment of native lesions potentiates VM progression,
we next investigated the effect of transitioning from flow to static culture of *PIK3CA^WT* and
PIK3CA^E542K microvessels by preconditioning microvessels with flow for 24 hrs then transitioning
to static conditions for the subsequent 24 hrs. After transition to static culture, *PIK3CA^WT* HUVECs
delaminated and detached into the apical lumen, resulting in gaps in the endothelium (**Figure
5A**). In contrast, *PIK3CA^E542K* HUVECs were insensitive to flow interruption, with no observable
delamination and the formation of tortuous, sprouted, and dilated microvessels that are akin to
mutant vessels cultured under flow conditions (**Figure 5B-C**). In all culture conditions,
PIK3CA^E542K microvessels contained approximately twice the number of cells and were
approximately 2-fold wider than *PIK3CA^WT* microvessels (**Figure 5C**). While *PIK3CA^E542K*
microvessels showed minimal effects from flow interruption, flow interruption in *PIK3CA^WT*
microvessels resulted in gaps covering approximately 6.5% of the vessel area (**Figure 5C**).
Collectively, these findings indicate that constitutive PI3K activation renders ECs insensitive to
vascular regression upon flow interruption and may lead to improved survival and favorable
expansion of mutant ECs in the slow flow characteristics of PIK3CA-driven VMs.

Basal-to-apical transmural flow exacerbates hypersprouting in PIK3CA mutant microvessels.

In addition to luminal shear stress, venous and lymphatic ECs are subjected to draining
transmural flow. Hydrostatic and oncotic pressure differences across the vessel wall causes fluid
to extravasate from blood capillaries, resulting in the accumulation of interstitial fluid in the
surrounding matrix which is subsequently reabsorbed and collected by venous and lymphatic
vessels [38]. Therefore, in addition to experiencing low level of wall shear stress, venous and
lymphatic ECs are also subjected to fluid stresses arising from transmural drainage. To investigate
the effect of transmural flow on *PIK3CA^WT* and *PIK3CA^E542K* microvessels, we cultured control or
mutant HUVECs microfluidic devices that contain two parallel microvessels in a shared collagen
I hydrogel [39]. This two-channel configuration allows us to establish a pressure gradient between
the two microchannels to drive transmural flow into (draining vessel, basal-to-apical flow) or out
of (source vessel, apical-to-basal flow) a microvessel (**Figure 6A, Figure S3A**). We observed
increased HUVEC invasion, reminiscent of angiogenic sprouting [40], into the subluminal collagen
I hydrogel upon application of transmural flow in the basal-to-apical direction in *PIK3CA^E542K*
microvessels as compared to *PIK3CA^WT* microvessels (**Figure 6A-C**). Consistent with previous

reports [41-43], this increase in cell invasion primarily occurred with the application of basal-to-
apical, but not apical-to-basal transmural flow (**Figure 6A-B**).

PIK3CA^{E542K} activation drives non-cell-autonomous sprouting in mosaic microvessels.

Slow-flow VMs arise from sporadic somatic mutations, resulting in vascular lesions consisting of
endothelial cells with and without the driving mutations. The mosaic nature of vascular lesions
implies a role of non-cell autonomous signaling where mutated endothelial cells influence the
behavior of neighboring non-mutated cells to amplify lesion growth and progression. To determine
how *PIK3CA^{E542K}* HUVECs influence non-mutated HUVECs, we generated mosaic microvessels
containing HUVECs expressing histone *H2B-mCherry* and inducible *tet-ON-PIK3CA^{E542K}*. We
then applied basal-to-apical transmural flow and investigated whether doxycycline induced
expression of *PIK3CA^{E542K}* influenced sprouting and invasion behavior of control *H2B-mCherry*
HUVECs (**Figure 7A**). We found that doxycycline induction increased both the number of *H2B-*
mCherry and *PIK3CA^{E542K}* HUVECs located within sprouts (**Figure 7B**), indicating a non-cell-
autonomous effect of PI3K activation on sprouting angiogenesis. We further quantified the tip cell
distribution in mosaic sprouts and observed that *tet-ON-PIK3CA^{E542K}* and *H2B-mCherry* HUVECs
contributed to 40% and 60% of tip cells, respectively in untreated mosaic microvessels (**Figure**
7C-D). Doxycycline treatment drastically changed the proportion of control and mutant tip cells,
where *PIK3CA^{E542K}* HUVECs occupied 87% of the tips and *H2B-mCherry* HUVECs contributed
mostly as stalk cells in mosaic sprouts of doxycycline treated microvessels (**Figure 7C-D**).
Together, these results suggest that *PIK3CA^{E542K}* endothelial cells are more migratory and have a
propensity to lead the sprouting and invasion of non-mutated endothelial cells.

We next determined whether the non-autonomous effect on sprouting is a result of matrix
remodeling by *PIK3CA^{E542K}* HUVECs. Proteolytic degradation of the subluminal collagen matrix
is known to be required for angiogenic sprout initiation and extension [44]. Previously, we
described an increase in protease secretion and ECM degradation by *PIK3CA-mutant* HUVECs
[19], and we thus hypothesized that ECM degradation and remodeling by *PIK3CA^{E542K}* HUVECs
may prime *PIK3CA^{WT}* HUVECs for angiogenic sprouting. To test this hypothesis, we formed
PIK3CA^{WT} and *PIK3CA^{E542K}* microvessels and lysed cells after 24 hrs, then reseeded the resulting
channels with *PIK3CA^{WT}* HUVECs (**Figure 7E**). We then applied basal-to-apical transmural flow
and quantified the number of sprouts formed in primed ECM (**Figure 7F**). We found that
PIK3CA^{WT} HUVECs cultured in microvessels primed by *PIK3CA^{E542K}* HUVECs failed to reproduce
the hypersprouting phenotype (**Figure 7F**). This finding suggests that changes in matrix
architecture and mechanics caused by *PIK3CA^{E542K}* HUVECs are not sufficient to account for the
etiopathology of VMs.

PIK3CA activation leads to increased endothelial cell and nuclear compliance:

Increased cell and nuclear compliance has been shown to promote 3D migration in various cell
types [45], and we thus hypothesized that the hypersprouting observed in *PIK3CA^{E542K}*
microvessels was driven by changes in cell and/or nuclear mechanical properties of *PIK3CA^{E542K}*
HUVECs. To distinguish changes in cell intrinsic mechanics from tissue mechanics, we quantified
cellular and nuclear deformation on a single cell basis using a microfluidic micropipette aspiration
assay [46, 47] (**Figure 8, Figure S4**). The microfluidic device was designed to have two inlets
allowing for the perfusion and loading of cells through the main channel. The main channel
contains an array of 18 pockets with narrow constriction micropipette channels and are connected
to an outlet port which is open to the atmosphere (**Figure S4**). A pressure of 7.0 kPa was applied

to inlet 1 and 1.4 kPa to inlet 2, and both outlets are drained through constricted channels to
atmospheric pressure (**Figure S4B**). Cells are initially trapped in larger pockets before deforming
into 5 μm × 5 μm constricted channels under a fixed pressure gradient, and deformation was
imaged via optical microscope for 2 minutes (**Figure 8B, Figure S4**). Interestingly, while
untransduced HUVECs maintained cellular integrity for the duration of the experiment, $PIK3CA^{WT}$
and $PIK3CA^{E542K}$ endothelial cells were prone to fracture as the cells passed through the narrow
constriction, with $PIK3CA^{E542K}$ cells having the lowest probability of survival throughout the
experiment (**Figure 8C**). The cytoplasm and nuclear protrusion length were measured as a
function of time, and while the cytoplasm of $PIK3CA^{WT}$ and $PIK3CA^{E542K}$ HUVECs deformed
similarly in response to the applied pressure gradient, both cell types demonstrated greater
cytoplasmic deformation and cytoplasm fracture than untransduced HUVECs (**Figure 8C-D**). The
nuclei of the three cell types deformed at different rates, with untransduced HUVEC nuclei
experiencing the least deformation over time and $PIK3CA^{E542K}$ expressing nuclei experiencing the
greatest deformation (**Figure 8E**). Nuclear and cytoplasm viscoelastic properties were computed
from the protrusion curves, as described previously [47]. Interestingly, $PIK3CA^{E542K}$ and $PIK3CA^{WT}$
HUVECs share similar cytoplasmic viscoelastic properties and are more compliant and less
viscous than the cytoplasm of untransduced HUVECs (**Figure 8F**). The nuclei of $PIK3CA^{E542K}$
HUVECs showed reduced elasticity as compared to the nuclei of $PIK3CA^{WT}$ and untransduced
HUVECs but exhibited the same viscous properties as $PIK3CA^{WT}$ nuclei (**Figure 8G**). Thus,
$PIK3CA^{E542K}$ HUVECs were distinguished only by their reduced nuclear elastic modulus
compared to $PIK3CA^{WT}$ HUVECs. Together, these results demonstrated increased cellular and
nuclear compliance in PIK3CA-mutant ECs which may underlie dilation and overgrowth of venous
and lymphatic vessels.

Discussion and conclusion

PIK3CA-driven VMs stereotypically present in soft tissues with low rates of perfusion [4, 48]. Here,
using a series of engineering and microfluidic tools, we investigated the role of the biophysical
microenvironment in the development of slow-flow vascular malformations to test the hypothesis
that ECM compliance and low hemodynamic shear stress potentiate lesion formation. Alignment
and elongation of the cytoskeleton in the direction of flow is a hallmark of endothelial
mechanotransduction [49], and laminar shear stress has been shown to regulate vascular barrier
function locally and acutely [31, 50]. Furthermore, a lack of alignment and barrier function in
response to hemodynamic shear stress is indicative of endothelial dysfunction [51]. Here, using
both an orbital flow system to impart a range of shear stress magnitudes and a Hele-Shaw flow
cell for unidirectional, laminar flow, we observed that HUVECs and LECs expressing a
constitutively active PIK3CA-activating mutation fail to elongate and align in response to laminar
shear stress (**Figures 1-2**). Furthermore, the permeability of PIK3CA-mutant endothelial cells was
insensitive to shear stress and remained elevated even in the presence of physiologic flow
(**Figures 1-2**). Together, these results suggest that dysfunctional mechanotransduction and
resultant insensitivity to flow could be a key mechanism that underlies the dysregulated vascular
networks characteristic of VMs.

We observed that expression of $PIK3CA^{E542K}$ reduced MLC phosphorylation and recruitment of
vinculin to adherens junction complexes, resulting in the discontinuity of junctional VE-cadherin
and increased permeability (**Figure 3**). Previous work has demonstrated that maturation of
adherens junction complexes requires a balance of MLC-mediated cell-generated forces and
Rac1 activity [52], and previously we identified that *PIK3CA* activating mutations result in excess

Rac1 activity [19]. Together, these results suggest that pathophysiologic PI3K activation results in
adherens junction instability through mechanical imbalances at cell-cell junctions. This
mechanism is supported by our observations in 3D microvessels, where $PIK3CA^{E542K}$
microvessels exhibit lower traction forces compared to $PIK3CA^{WT}$ microvessels (**Figure 4**),
consistent with published data demonstrating that PIK3CA signaling inhibits NUAK1-dependent
phosphorylation of myosin phosphatase targeting-1 in ECs, leading to reduced overall actomyosin
contractility [18]. Collectively, these results suggest that adherens junction instability and lack of
maturation in response to hemodynamic signals render PIK3CA mutant cells prone to
angiogenesis and could underlie the overconnected topologies characteristic of VMs.

Clinically, VMs are described as soft and compressible, and histologically they appear as a single
layer of endothelial cells in enlarged vein-like channels with disorganized pericellular matrix [4].
To recapitulate the compliant ECM characteristic of tissues susceptible to VMs, we developed a
3D microfluidic model of vascular malformations and observed cellular signatures consistent with
the clinical presentation of VMs, including irregular and dilated microvessels (**Figure 4**).
Previously, we have shown that reducing hemodynamic shear stress leads to rapid adherens
junction dissolution [31], and experiments conducted in developmental animal models have
demonstrated that reduction or interruption of flow induces vessel regression and network pruning
[53], which is thought to be essential for optimization of efficient delivery of oxygen and nutrients
[54, 55]. To recapitulate the hemodynamic signals associated with pruning, we exposed
microvessels to flow for 24 hrs, then interrupted flow and cultured microvessels in static conditions
for 24 hrs. Interestingly, PIK3CA mutant microvessels did not respond to flow interruption, and
mutant ECs remain attached and continue to grow under static culture conditions, whereas wild-
type ECs delaminate from the vessel wall (**Figure 5**). These results suggest that cells expressing
activating PIK3CA mutations are primed for survival in the unique hemodynamic environment of
slow-flow lesions, and that this survival results in a lack of pruning in response to reduced or
interrupted perfusion to allow VM progression in the slow-flow microenvironment.

Computational modeling and a limited number of experiments have demonstrated that changes
in vascular network topology modulate not only local wall shear stress but also luminal pressure
[56]. To determine whether gradients in fluid pressure could play a role in lesion formation, we
fabricated microfluidic devices to expose wild-type and mutant microvessels to transmural
pressure gradients. We observed that transmural pressure gradients and resultant flow in the
basal-to-apical direction results in increased sprouting and invasion of PIK3CA-mutant ECs as
compared to control microvessels (**Figure 6**). Previously, immature endothelial monolayers,
characterized by discontinuous junctional VE-cadherin have been shown to sprout in response to
lower basal-to-apical pressure drops than mature monolayers with continuous pericellular VE-
cadherin [57], suggesting that the increased sprouting in PIK3CA-mutant ECs could be driven by
immature adherens junctions. Furthermore, basal-to-apical transmural flow has been shown to
synergize with VEGF gradients to increase angiogenic sprouting [42], and a recent study has
demonstrated increased paracrine VEGF-C signaling in $PIK3CA$-driven mosaic LM mouse model
[26]. Together, these results suggest that transmural pressure gradients and flow are a pro-
angiogenic stimulus that drives lesion elaboration and growth.

Previous studies have demonstrated that nuclear stiffness and deformability play a role in 3D
cellular migration through dense collagen matrices and spatially constraining micropores [45, 58].
Using a microfluidic micropipette aspiration assay, we found that $PIK3CA^{WT}$ and $PIK3CA^{E542K}$ ECs
exhibit increased cellular compliance compared to untransduced HUVECs (**Figure 8**) and that

$PIK3CA^{E542K}$ ECs have significantly more compliant nuclei than $PIK3CA^{WT}$ ECs. The increased
cellular and nuclear compliance in PIK3CA-mutant ECs may facilitate EC migration and sprout
formation, potentially contributing to hypervascularization in VM lesions. While the mechanisms
relating PIK3CA activation to nuclear compliance remain unknown, increased invasion of
colorectal cancer cells expressing PIK3CA activating through 8 μm pores and through Matrigel
has been observed [59], while knockdown of PIK3CA in glioblastoma cells reduces invasion into
Matrigel [60], collectively suggesting that PIK3CA-mediated changes in nuclear mechanics could
underlie a wide variety of pathogenic processes.

In summary, our findings demonstrate a role of altered mechanotransduction and cell mechanics
in PIK3CA-driven vascular malformations. These findings could potentially be leveraged for
genetic diagnosis and treatment of vascular anomalies. Molecular genetic testing for VMs remains
challenging as the lesions are mosaic, with only a small population of cells expressing the driving
mutations. Therefore, these reported changes in cellular mechanics may potentially be used as a
biomarker for the sorting and enrichment of mutated ECs from patient biopsies. Further, targeting
the defective contractility and recalibrating the mechanotransduction in mutant endothelial cells
could represent an avenue to reduce or regress lesion progression.

Materials and methods

Cell culture, cloning, and lentiviral transduction

HUVECs (Lonza) and juvenile HDLECs (Promocell) were cultured in endothelial growth medium (EGM)-2 growth medium (Lonza) in a humidified incubator at 37 ºC and 5% CO_2. HUVECs were used between passages 6 and 12. HDLECs were used between passage 3 and 7. Lentivirus generation and transduction of HUVECs with pHAGE-PIK3CAWT, pHAGE-PIK3CAE542K, and pLenti6-H2B-mCherry (Addgene, plasmid #89766, gift from Torsten Wittmann) were performed as previously described [19]. PIK3CAE542K cDNA was amplified from pHAGE-PIK3CAE542K (Addgene, plasmid # 116479, gift from G. Mills and K. Scott) and was subcloned into pCW57.1, a third-generation inducible lentiviral expression vector (Addgene, plasmid # 41393, gift from David Root). All constructs were expressed through lentiviral transduction. HUVECs expressing pCW57.1-tetON-PIK3CAE542K were selected with 2 µg/mL of puromycin (Corning) and 5 µg/mL of blasticidin (Fisher) in EGM-2, respectively for at least two cell passages. Doxycycline (Sigma, 10 µg/mL in EGM-2) was used to induce the expression of tet-ON-PIK3CAE542K.

Hemodynamic shear stress via orbital shaking and 2D permeability assay

HUVECs were seeded at 0.15 x 10^6 cells per well in a 6-well plate (Corning Costar 6-well Clear TC-treated, Cat #3516) coated with 50 µg/mL of rat-tail derived type I collagen (Corning 354326) in 0.02 M acetic acid solution. Cells were allowed to adhere overnight prior to the application of orbital shaking. Fresh EGM-2 (2 mL) was added into each well prior to the application of orbital flow. Orbital flow was applied by culturing ECs for 3 days on an orbital shaker set to 200 rpm, which corresponds to a maximum wall shear stress of 15 Dynes/cm^2 [61]. For measurements of local changes in vascular permeability, HUVECs were seeded in a 6-well plate coated with 50 µg/mL of biotinylated-human fibronectin (Corning 354008) in PBS. Fibronectin biotinylation was performed with EZ-link-Sulfo-NHS-LC-Biotin (Thermo Fisher A39257) as described previously [62]. Briefly, 0.1 mg/mL of fibronectin in PBS was incubated with 0.5 mM of EZ-link-Sulfo-NHS-LC-Biotin for 30 min at room temperature. At the end of orbital flow culture, cells were incubated with 25 mg/mL Cy3-Streptavidin (Jackson Immuno Research Labs 016160084) in PBS containing calcium and magnesium (PBS++) for 2 minutes. After streptavidin labeling of exposed matrix, cells were rinsed once with PBS++ and were fixed with 4% paraformaldehyde (Electron Microscopy Sciences) in PBS++ for 15 min at room temperature. Permeability was measured by quantifying the streptavidin positive area using ImageJ.

Hydrogel formation and cell seeding in microvessel microfluidic devices

Microfluidic devices were fabricated as described in the *Supplementary Methods*. To improve hydrogel adhesion to PDMS devices prepared by photolithography, 2 mg/mL dopamine hydrochloride in 10 mM of pH 8.5 Tris HCl buffer (BioWorld) was pipetted into the hydrogel chamber, incubated for 1 hr at room temperature, rinsed twice with DI-H_2O, left dry at room temperature, and sterilized via UV for 20 min. Sterile 160 µm stainless steel acupuncture needles (Llahsa Oms) were coated with 0.01% bovine serum albumin (BSA) in PBS for at least 30 min prior to hydrogel injection. BSA-coated needles were inserted into each device, and 2.5 mg/mL of rat tail-derived collagen I hydrogel was prepared and pipetted into the hydrogel chamber as described previously [37], and allowed to polymerize at 37 ºC for 30 min prior to hydrating with EGM-2. After 4 hrs of further incubation at 37 ºC, acupuncture needles were gently removed from devices to form a cylindrical channel within the collagen I hydrogel. Needle inlets were sealed with vacuum grease (Millipore Sigma) and fresh EGM-2 was added to the media ports of the

devices. Devices were kept on a rocker at 5 cycles/min at 37 ºC and 5% CO_2 for at least 12 hrs
prior to cell seeding. To form microvessels, HUVECs were resuspended at 0.3×10^6 cells/mL in
EGM-2. Media was removed from both media ports and gel filling pipette tips were used to
aspirate and empty the hydrogel ports prior to seeding. To promote pressure driven flow into the
microchannel, 50 µL and 40 µL of cell suspension were added to the inlet and outlet media ports,
respectively. Devices were inverted every minute for 8 min at room temperature, followed by
incubating upside-down in a tissue culture incubator for 6 minutes. Devices were subsequently
inverted and incubated for an additional 6 minutes in the incubator. After 20 min of seeding,
devices were placed on a rocker for at least 2 hours before changing EGM-2 media, which was
subsequently changed daily.

Transmural flow in two-channel devices

Two-channel devices were used to apply transmural flow into or out of microvessels (**Figure S3**).
For the application of transmural flow, microvessels were first cultured on rocker (5 cycles per
minute at ± 30° angle) for 2 days. Media reservoirs were inserted into all four media ports and any
gaps between reservoirs and PDMS were sealed with vacuum grease. Pre-warmed EGM-2 was
added either to the empty or microvessel channel to apply a 20 mm H_2O hydrostatic pressure
driven transmural flow into or out of the microvessels, respectively. Microvessels were fixed 24
hours after the application of transmural flow.

Microfluidic micropipette aspiration

Microfluidic devices were fabricated as described in the *Supplementary Methods.* The
micropipette aspiration device was used to deform cells under an applied pressure gradient.
Untransduced HUVECs, *PIK3CA^{WT}*, and *PIK3CA^{E542K}* cells were cultured as described previously
before being stained with Hoechst 3342 (10 mg/mL, Thermo Scientific) and CellMask Deep Red
plasma membrane stain (5 mg/mL, 1:2000 v/v, Invitrogen) for 10 min to stain the nucleus and
plasma membrane respectively. Stained cells were washed with PBS and allowed to recover in
EGM-2 for 30 minutes. Cells were then trypsinized, counted, and resuspended at a concentration
of 5,000,000 cells/mL in BSA (20 mg/mL in PBS) before being put on ice. Constant-pressure
microfluidic flow controllers ("Flow EZ," Fluigent) were used to apply flow to the opposing inlets at
constant pressures, creating flow from one inlet to the other using a known pressure gradient.
Two flow controllers were each connected to a dry air source and a power outlet. One controller
was used to drive stained and suspended cells from a 15 mL conical tube into one inlet of the
microfluidic device at a pressure of 7.0 kPa. A second controller was used to drive cell-free PBS
from a 15 mL conical tube into the second inlet of the microfluidic device at a pressure of 1.4 kPa.
The outlet port of the device was open to the atmosphere (**Figure S4**).

Decellularization and reseeding of decellularized microvessels

Fresh sterile decellularization solution (0.5% Triton with 50 mM NH_4OH in PBS) was prepared on
the day of decellularization. Microvessels were rinsed once with sterile PBS prior to
decellularization. After PBS wash, 100 µL of decellularization solution was added to the media
ports of each device. Devices were kept on a rocker at 37 ºC and 5% CO_2 for 15 min.
Decellularized devices were washed three times with PBS and residual DNA was digested via
treatment with 10 µg/mL of DNAseI (Sigma Aldrich, D4527) in PBS++ for 45 minutes at 37 ºC.
DNAseI treated devices were washed three times with PBS and were incubated overnight on a
rocker in fresh EGM-2 media. Devices were reseeded with *PIK3CA^{WT}* cells the next day as

described in the "Hydrogel formation and cell seeding" section. Basal-to-apical pressure driven
transmural flow was applied a day after reseeding as described above.

Immunostaining

Mouse Anti–VE-cadherin (F-8) was from Santa Cruz Biotechnology. Rabbit Anti-VE-cadherin
(ab33168) was from Abcam. Mouse Anti-Vinculin (hVIN-1) was from Sigma Aldrich. 4′,6-
Diamidino-2-phenylindole (DAPI), rhodamine phalloidin (1 mg/ml), Alexa Fluor 488 phalloidin (1
mg/ml), and Alexa Fluor–conjugated secondary antibodies were from Life Technologies. For
endpoint immunostaining, fixed cells or microvessels were permeabilized with 0.3% Triton X-100
(Millipore Sigma) for 30 minutes at room temperature and non-specific antibody binding was
blocked by treatment with 2% (w/v) BSA in PBS for an hour at room temperature. Primary
antibodies against VE-cadherin (1:200, v/v, F-8, Santa Cruz Biotechnology; 1:250, v/v, ab33168,
Abcam), vinculin (1: 400, hVIN-1, Sigma Aldrich) were diluted in 2% BSA blocking solution and
applied overnight on a laboratory rocker at 4 ºC. Cells and devices were subsequently rinsed
three times over 15 min with PBS, and were incubated in secondary antibodies (1:1000, v/v, goat
anti-mouse or goat-anti rabbit immunoglobulin G conjugated to Cy3 or Alexa Fluor 647, Thermo
Fisher Scientific), Alexa Fluor 488 or rhodamine phalloidin (1:200, v/v, Thermo Fisher Scientific),
and DAPI (1:1000, v/v, Thermo Fisher Scientific) diluted in blocking solution for an hour at room
temperature. Labeled cells and devices were rinsed three times over 15 min with PBS and were
stored in 0.01% Sodium Azide solution diluted in PBS until imaging.

Western blots

Cell lysates were collected from cells cultured in complete media for 24 hours. Briefly, cells were
washed once with ice cold PBS and lysed on ice with radioimmunoprecipitation assay buffer (RIPA
buffer, Thermo Fisher Scientific) containing 1X HALT protease and phosphatase inhibitor (Thermo
Fisher Scientific). Lysates were homogenized by pulse-vortexing and clarified by centrifugation at
14,000 x g for 5 min at 4 ºC. The concentration of clarified lysates was quantified using Pierce
BCA Protein Assay (Thermo Fisher Scientific). Lysates were then reduced with NuPAGE LDS
reducing agent and dithiothreitol and denatured for 5 min at 100 ºC. Denatured proteins were
resolved in NuPAGE 4 to 12% bis-tris gradient gels (Thermo Fisher Scientific) and transferred to
iBlot2 PVDF ministack membranes. Membranes were blocked with SuperBlock Blocking Buffer
(Thermo Fisher Scientific) for an hour with rocking at room temperature and were incubated with
the following primary antibodies for overnight with rocking at 4 ºC: anti-p110α (1:1000, Cell
Signaling Technology, #9272), anti-vinculin (1:10,000, Sigma-Aldrich, V9264), anti-VE-cadherin
(1:1000, Santa Cruz, SC9989), anti-Beta-catenin (1:1000, Cell Signaling Technology, #8480),
anti-glyceraldehyde-3-phosphate dehydrogenase (1:5000, Cell Signaling Technology, #2118),
and anti-phospho-myosin light chain 2 (Thr18/Ser19) (1:1000, Cell Signaling Technology, #3674).
After overnight incubation with primary antibodies, membranes were washed three times with
TBS-T and were subsequently incubated with horseradish peroxidase conjugated antibodies
(1:10,000, v/v, diluted in 5% milk (w/v)) for 1 hour with rocking at room temperature. Membranes
were then washed three times over a period of 15 min with TBS-T and were developed via
chemiluminescence using either SuperSignal West Femto Maximum Sensitivity Substrate
reagent (Thermo Fisher Scientific) or Clarity Western ECL reagent (BioRad). Images of western
blots were quantified using ImageJ.

Imaging for Microvessels and flow-conditioned cells

Confocal z-stacks of microvessels (4 μm step size, captured from top to bottom of the
microvessels) were acquired with an Olympus FV3000 laser scanning confocal microscope with
a 10x U Plan S-Apo NA 0.4 air objective. Fluorescence images of cells after orbital flow were
captured at 10x magnification (10X U Plan Fluor NA 0.3 air objective, Olympus) using an
automated scanning stage on an Olympus wide-field microscope. Briefly, the edge of a well of a
6-well plate was positioned over the objective, and a uniform rectangular area from the edge to
the middle of the well was selected. A focus map was generated to include at least 10 focus points
along the length of the rectangular area. The autofocus feature was used to cycle through the
focus map and set the z-focus for each imaging point used throughout the area scan.
Computational image processing is described in the *Supplementary Methods*.

Imaging for Microfluidic Micropipette Aspiration Assay

Cell deformation into the micropipette channels was imaged using a timelapse acquisition on an
inverted widefield fluorescence microscope (Olympus IX83) with a white light source 20X/NA 0.45
objective. DAPI and Cy5 filters were used to image the nucleus and cytoplasm respectively, and
images were acquired every 2.935 seconds for a total acquisition time of 2.5 min. At the end of
each timelapse acquisition, deformed cells were cleared from the cell pockets by applying
pressure at the outlet port by pipette. Cleared pockets were then filled again by flowing cells during
the next acquisition period.

Statistical analysis

Graphs were generated and statistical analyses were performed in Prism 10 (GraphPad).
Unless otherwise mentioned, data were plotted as mean ± standard deviation of ≥3 experiments
and data points denote values from each cell or microvessel. Data set with normal distribution
were compared with two-tailed Student's test and one-way ANOVA followed by Tukey's post hoc
test. Kruskal-Wallis test followed by Dunn's multiple-comparison was used for comparison of
non-Gaussian distributed data. Two-way ANOVA followed by Šidák's test was used for
comparison of more than two variables.

 Figures and captions

Figure 1. **Constitutive PI3K activation impedes shear stress-induced endothelial cell alignment and vascular barrier function.** (**A**) Representative images of *PIK3CA^WT^* and *PIK3CA^E542K^* HUVECs cultured on biotinylated fibronectin for 72 hr under orbital shaking conditions. Cells were labeled with DAPI (cyan) and VE-cadherin (magenta). Monolayer permeability was assessed through streptavidin labeling (yellow). Scale bar = 10 μm. (**B**) Quantification of cellular orientation in the central, middle, or outer edge region of the well after exposure to orbital shaking (n = 3; mean ± s.d., Kruskal-Wallis test followed by Dunn's test). (**C**) Quantification of cellular eccentricity as a function of radial distance from the center of the well for

PIK3CA^{WT} and *PIK3CA^{E542K}* HUVECs (n = 3; mean ± s.d., two-way analysis of variance (ANOVA)
followed by Šidák's test). (**D**) Quantification of cell density as a function of radial distance from the
center of the well for PIK3CA^{WT} and PIK3CA^{E542K} HUVEC (n = 3; means ± s.d., two-way ANOVA
followed by Šidák's test). (**E**) Quantification of streptavidin intensity as a function of radial distance
from the center of the well for *PIK3CA^{WT}* and *PIK3CA^{E542K}* HUVECs (n = 3; mean ± s.d., two-way
ANOVA followed by Šidák's test). (**F**) Representative images of *PIK3CA^{WT}* and *PIK3CA^{E542K}*
HUVECs exposed to 10 dyn/cm^2 laminar shear stress in a Hele-Shaw flow cell. Scale bar = 10
µm. *p< 0.05, **p< 0.01, ***p< 0.001, ****p< 0.0001 for all plots.

VE-CADHERIN

Figure 2. Preconditioning with shear stress prior to the induction of _PIK3CA^{E542K}_ expression restores shear stress-mediated EC alignment but not barrier function. (**A**) Schematic of experimental workflow and representative images of VE-cadherin staining in control cells and cells where expression of _PIK3CA^{E542K}_ was induced 24 hr before flow, at the onset of flow, or 24 hr after flow. Images were captured near the center of the well or near the outer edge of the well. Scale bar = 10 μm. (**B**) Quantification of cellular eccentricity as a function of radial distance for control cells as well as for cells induced with doxycycline 24 hr before flow, at the onset of flow, or 24 hr after flow (n = 3; mean ± s.d., two-way ANOVA followed by Šidák's test).

(**C**) Quantification of cellular orientation in the central, middle, or outer region of the well for cells
induced with doxycycline 24 hr before flow, at the onset of flow, or 24 hr after flow (n = 3; mean ±
s.d., Kruskal-Wallis tests followed by Dunn's test). (**D**) Quantification of streptavidin area as a
function of radial distance from the center of the well for control cells, cells induced with
doxycycline 24 hr before flow, and cells induced with doxycycline 24 hr after flow (n = 3; mean ±
s.d. (dotted lines), ANOVA followed by Dunnett test). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p <$
0.0001 for all plots.

Figure 3. Loss of junctional vinculin recruitment in *PIK3CA^E542K* endothelium. (**A**)

Representative images of *PIK3CA^WT* and *PIK3CA^E542K* HUVECs exposed to orbital shaking and

stained for vinculin (red), F-actin (cyan), and VE-cadherin (grey). Scale bar = 10 μm. (**B**)

Quantification of percent vinculin area, mean intensity of vinculin, and mean VE-cadherin

intensity at cell-cell borders (n = 957 *PIK3A^WT* and 400 *PIK3CA^E542K* cells; mean ± s.d., two-

tailed unpaired t-test). (**C**) Western blots for β-catenin, VE-cadherin, phospho-myosin light chain

2, and vinculin. (D) Quantification of relative VE-cadherin, vinculin, β-catenin, and phospho-

myosin light chain 2 in *PIK3CA^WT* or *PIK3CA^E542K* cells (n = 3; mean ± s.d., two-tailed unpaired t-

test). *p< 0.05, **p< 0.01, ***p< 0.001, ****p< 0.0001 for all plots.

Figure 4. *PIK3CA^E542K* ECs form dilated microvessels with increased sprouting and reduced traction forces. (**A**) Representative confocal images (maximum intensity projections) of *PIK3CA^WT* and *PIK3CA^E542K* microvessels treated with DMSO, 1 μM alpelisib, or 1 μM rapamycin. Scale bar = 100 μm. (**B**) Quantification of vessel width in *PIK3CA^WT* microvessels or *PIK3CA^E542K* microvessels treated with DMSO load control, rapamycin (rap.), or alpelisib (n = 3; mean ± s.d., one-way ANOVA followed by Tukey test). (**C**) Representative phase contrast (grey) and confocal reflectance (cyan and magenta) images of *PIK3CA^WT* and *PIK3CA^E542K* microvessels before and after decellularization. (**D**) Quantification of subluminal ECM displacement before and after decellularization of *PIK3CA^WT* or *PIK3CA^E542K* microvessels (n ≥ 4 microvessels, 3 measurements per microvessel; data points from same devices are color-matched; mean ± s.d., two-tailed unpaired t-test). *p< 0.05, **p< 0.01, ***p< 0.001, ****p< 0.0001 for all plots.

Figure 5. *PIK3CA*^{E542K} **microvessels are unresponsive to flow interruption.** (**A**) Representative images of microvessels generated from *PIK3CA*^{WT} and *PIK3CA*^{E542K} HUVECs. Microvessels were exposed to continuous rocking 48 hr and stained for DAPI (cyan), VE-cadherin (white), and F-actin (magenta). Scale bars = 100 μm. (**B**) Representative images of *PIK3CA*^{WT} and *PIK3CA*^{E542K} microvessels exposed to 24 hr rocking followed by 24 hr static conditions and stained for DAPI (cyan), VE-cadherin (white), and F-actin (magenta). Scale bars = 100 μm. (**C**) Quantification of vessel width, number of cells, vessel area, and gap area for *PIK3CA*^{WT} and *PIK3CA*^{E542K} microvessels exposed to either continuous or interrupted flow (n ≥ 4; mean ± s.d., two-way ANOVA). *p< 0.05, **p< 0.01, ***p< 0.001, ****p< 0.0001 for all plots.

Figure 5. *PIK3CA^{E542K}* **microvessels are unresponsive to flow interruption.** (**A**) Representative images of microvessels generated from *PIK3CA^{WT}* and *PIK3CA^{E542K}* HUVECs. Microvessels were exposed to continuous rocking 48 hr and stained for DAPI (cyan), VE-cadherin (white), and F-actin (magenta). Scale bars = 100 μm. (**B**) Representative images of *PIK3CA^{WT}* and *PIK3CA^{E542K}* microvessels exposed to 24 hr rocking followed by 24 hr static conditions and stained for DAPI (cyan), VE-cadherin (white), and F-actin (magenta). Scale bars = 100 μm. (**C**) Quantification of vessel width, number of cells, vessel area, and gap area for *PIK3CA^{WT}* and *PIK3CA^{E542K}* microvessels exposed to either continuous or interrupted flow (n ≥ 4; mean ± s.d., two-way ANOVA). *p< 0.05, **p< 0.01, ***p< 0.001, ****p< 0.0001 for all plots.

Figure 6. Basal-to-apical transmural flow exacerbates hypersprouting in PIK3CA mutant microvessels. (**A**) Representative images of *PIK3CA^WT^* and *PIK3CA^E542K^* microvessels exposed to continuous rocking for 48 hr and subsequently subjected to apical-to-basal (source) or basal-to-apical (sink) transmural flow for 24 hr. Scale bar = 100 μm. (**B**) Quantification of number of spouts in microvessels exposed to either luminal, apical-to-basal, or basal-to-apical flow (n ≥ 4; mean ± s.d., two-way ANOVA followed by Tukey test). (**C**) Fluorescence intensity profiles of actin channel were used to quantify number of sprouts in *PIK3CA^WT^* or *PIK3CA^E542K^* microvessels. Dashed white boxes indicate where intensity profiles were extracted. *p< 0.05, **p< 0.01, ***p< 0.001, ****p< 0.0001 for all plots.

Figure 7. Mosaic co-culture of HUVECs with PIK3CA^E542K ECs resulted in increased sprouting of non-mutated HUVECs. (**A**) Representative images of mosaic microvessels

generated using HUVECs expressing H2BmCherry and inducible tet-ON-PIK3CA^{E542K}.
Microvessels were first exposed to continuous rocking 48 hr and then were subjected to basal-to-
apical transmural flow for the next 24 hrs. PIK3CA^{E542K} expression was induced upon the
application of transmural flow. Scale bar = 100 µm. (**B**) Quantification of the distribution of
H2BmCherry and tet-ON-PIK3CA^{E542K} cells in mosaic vessels that were subjected to basal to
apical transmural flow (n ≥ 3; chi-square test). (**C**) Quantification of percent H2BmCherry and tet-
ON-PIK3CA^{E542K} tip cells in mosaic sprouts (n = 5 control and 15 doxycycline treated mosaic
sprouts; chi-square test). (**D**) Images of mosaic H2BmCherry (yellow) and tet-ON-PIK3CA^{E542K}
sprouts stained for DAPI (magenta) and actin (cyan). Scale bar = 10 µm. (**E**) Decellularized
PIK3CA^{WT} and *PIK3CA^{E542K}* microvessels were reseeded with *PIK3CA^{WT}* HUVECs. Reseeded
microvessels were subjected to basal-to-apical transmural flow for 24 hr. Scale bar = 100 µm. (**F**)
Representative confocal images and number of sprout quantifications of transmural flow
conditioned *PIK3CA^{WT}* microvessels cultured on extracellular matrix modified by control or mutant
cells. Scale bar = 10 µm. (n = 3; mean ± s.d., unpaired two-tailed t-test).

Figure 8. Altered cytoplasmic and nuclear mechanical properties of single cells expressing *PIK3CA^{WT}* or *PIK3CA^{E542K}* activating mutations. (**A**) An image of the microfluidic micropipette aspiration device labeled with an outlet and two inlets at differential pressures causing a flow vector between the two inlets. A magnified image depicts the dashed area, showing the array of pockets and constriction channels which trap and deform flowing cells under a constant applied pressure gradient. (**B**) Time-lapse images of nucleus (Hoechst) and cytoplasm (CellMask) deformation of untransduced HUVECs and endothelial cells expressing *PIK3CA^{WT}* or *PIK3CA^{E542K}* activating mutations. (**C**) Percentage of HUVECs that withstood deformation, as measured by the number of endothelial cells with intact cytoplasm after passage through the constriction channel relative to the total number of cells that entered the channels. (**D**) Protrusion length over time of endothelial cell cytoplasm into the constriction channels (mean ± s.e.m.). (**E**) Quantification of nuclear deformation over time of endothelial cells into constriction channels (mean ± s.e.m.). (**F**) Quantification of viscoelastic properties of the cytoplasm of endothelial cells (mean ± s.e.m., one-way ANOVA). (**G**) Elastic modulus, short-term viscosity, and long-term viscosity of the nuclei of HUVECs and endothelial cells expressing *PIK3CA^{WT}* or *PIK3CA^{E542K}* activating mutations (mean ± s.e.m., one-way ANOVA).

Acknowledgements

This work was funded by the National Institute of Health (R35GM142944), by the American Heart Association (CDA857738), and by research grants from the CLOVES Syndrome Community and the Lymphatic Malformation Institute. W.Y.A. is supported by grant from the CLOVES Syndrome Community. C.P.W. and E.L.D. are supported by a training fellowship from the Integrative Vascular Biology Training Program (T32HL69768), C.P.W. is supported by an American Heart Association predoctoral fellowship (24PRE1192185), and E.L.D. is supported by a Ruth L. Kirchstein predoctoral individual fellowship (F31HL162462). E.M. acknowledges support from an NIH training fellowship (T32GM135128). We thank Bob Geil for advice and help with microfabrication. The fabrication of microfluidic devices was performed in the Chapel Hill Analytical and Nanofabrication Laboratory (CHANL), a member of the North Carolina Research Triangle Nanotechnology Network (RTNN), which is supported by the National Science Foundation (ECCS-2025064), as part of the National Nanotechnology Coordinated Infrastructure (NNCI).

Author Declarations

Conflict of Interest

The authors have no conflicts to disclose.

Ethics Approval

Ethics approval is not required.

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

[1] K. Zenner, C.V. Cheng, D.M. Jensen, A.E. Timms, G. Shivaram, R. Bly, S. Ganti, K.B. Whitlock, W.B. Dobyns, J. Perkins, J.T. Bennett, Genotype correlates with clinical severity in PIK3CA-associated lymphatic malformations, JCI Insight 4(21) (2019).

[2] K.C. Kurek, V.L. Luks, U.M. Ayturk, A.I. Alomari, S.J. Fishman, S.A. Spencer, J.B. Mulliken, M.E. Bowen, G.L. Yamamoto, H.P. Kozakewich, M.L. Warman, Somatic mosaic activating mutations in PIK3CA cause CLOVES syndrome, Am J Hum Genet 90(6) (2012) 1108-15.

[3] M. Sadick, R. Muller-Wille, M. Wildgruber, W.A. Wohlgemuth, Vascular Anomalies (Part I): Classification and Diagnostics of Vascular Anomalies, Rofo 190(9) (2018) 825-835.

[4] A. Queisser, E. Seront, L.M. Boon, M. Vikkula, Genetic Basis and Therapies for Vascular Anomalies, Circ Res 129(1) (2021) 155-173.

[5] J.A. Cox, E. Bartlett, E.I. Lee, Vascular malformations: a review, Semin Plast Surg 28(2) (2014) 58-63.

[6] N. Limaye, J. Kangas, A. Mendola, C. Godfraind, M.J. Schlogel, R. Helaers, L. Eklund, L.M. Boon, M. Vikkula, Somatic Activating PIK3CA Mutations Cause Venous Malformation, Am J Hum Genet 97(6) (2015) 914-21.

[7] P. Castel, F.J. Carmona, J. Grego-Bessa, M.F. Berger, A. Viale, K.V. Anderson, S. Bague, M. Scaltriti, C.R. Antonescu, E. Baselga, J. Baselga, Somatic PIK3CA mutations as a driver of sporadic venous malformations, Sci Transl Med 8(332) (2016) 332ra42.

[8] S.D. Castillo, E. Tzouanacou, M. Zaw-Thin, I.M. Berenjeno, V.E. Parker, I. Chivite, M. Mila-Guasch, W. Pearce, I. Solomon, A. Angulo-Urarte, A.M. Figueiredo, R.E. Dewhurst, R.G. Knox, G.R. Clark, C.L. Scudamore, A. Badar, T.L. Kalber, J. Foster, D.J. Stuckey, A.L. David, W.A. Phillips, M.F. Lythgoe, V. Wilson, R.K. Semple, N.J. Sebire, V.A. Kinsler, M. Graupera, B. Vanhaesebroeck, Somatic activating mutations in Pik3ca cause sporadic venous malformations in mice and humans, Sci Transl Med 8(332) (2016) 332ra43.

[9] L. Rodriguez-Laguna, N. Agra, K. Ibanez, G. Oliva-Molina, G. Gordo, N. Khurana, D. Hominick, M. Beato, I. Colmenero, G. Herranz, J.M. Torres Canizalez, R. Rodriguez Pena, E. Vallespin, R. Martin-Arenas, A. Del Pozo, C. Villaverde, A. Bustamante, C. Ayuso, P. Lapunzina, J.C. Lopez-Gutierrez, M.T. Dellinger, V. Martinez-Glez, Somatic activating mutations in PIK3CA cause generalized lymphatic anomaly, J Exp Med 216(2) (2019) 407-418.

[10] T. Hong, X. Xiao, J. Ren, B. Cui, Y. Zong, J. Zou, Z. Kou, N. Jiang, G. Meng, G. Zeng, Y. Shan, H. Wu, Z. Chen, J. Liang, X. Xiao, J. Tang, Y. Wei, M. Ye, L. Sun, G. Li, P. Hu, R. Hui, H. Zhang, Y. Wang, Somatic MAP3K3 and PIK3CA mutations in sporadic cerebral and spinal cord cavernous malformations, Brain 144(9) (2021) 2648-2658.

[11] M. Peyre, D. Miyagishima, F. Bielle, F. Chapon, M. Sierant, Q. Venot, J. Lerond, P. Marijon, S. Abi-Jaoude, T. Le Van, K. Labreche, R. Houlston, M. Faisant, S. Clemenceau, A.L. Boch, A. Nouet, A. Carpentier, J. Boetto, A. Louvi, M. Kalamarides, Somatic PIK3CA Mutations in Sporadic Cerebral Cavernous Malformations, N Engl J Med 385(11) (2021) 996-1004.

[12] A.A. Ren, D.A. Snellings, Y.S. Su, C.C. Hong, M. Castro, A.T. Tang, M.R. Detter, N. Hobson, R. Girard, S. Romanos, R. Lightle, T. Moore, R. Shenkar, C. Benavides, M.M. Beaman, H. Muller-Fielitz, M. Chen, P. Mericko, J. Yang, D.C. Sung, M.T. Lawton, J.M. Ruppert, M. Schwaninger, J. Korbelin, M. Potente, I.A. Awad, D.A. Marchuk, M.L. Kahn, PIK3CA and CCM mutations fuel cavernomas through a cancer-like mechanism, Nature 594(7862) (2021) 271-276.

[13] R. Huo, Y. Yang, Y. Sun, Q. Zhou, S. Zhao, Z. Mo, H. Xu, J. Wang, J. Weng, Y. Jiao, J. Zhang, Q. He, S. Wang, J. Zhao, J. Wang, Y. Cao, Endothelial hyperactivation of mutant MAP3K3 induces cerebral cavernous malformation enhanced by PIK3CA GOF mutation, Angiogenesis 26(2) (2023) 295-312.

[14] K. Banerjee, Y. Lin, J. Gahn, J. Cordero, P. Gupta, I. Mohamed, M. Graupera, G. Dobreva, M.A.
Schwartz, R. Ola, SMAD4 maintains the fluid shear stress set point to protect against arterial-
venous malformations, J Clin Invest 133(18) (2023).
[15] R. Ola, A. Dubrac, J. Han, F. Zhang, J.S. Fang, B. Larrivee, M. Lee, A.A. Urarte, J.R. Kraehling, G.
Genet, K.K. Hirschi, W.C. Sessa, F.V. Canals, M. Graupera, M. Yan, L.H. Young, P.S. Oh, A. Eichmann,
PI3 kinase inhibition improves vascular malformations in mouse models of hereditary
haemorrhagic telangiectasia, Nat Commun 7 (2016) 13650.
[16] E. Alsina-Sanchis, Y. Garcia-Ibanez, A.M. Figueiredo, C. Riera-Domingo, A. Figueras, X. Matias-
Guiu, O. Casanovas, L.M. Botella, M.A. Pujana, A. Riera-Mestre, M. Graupera, F. Vinals, ALK1 Loss
Results in Vascular Hyperplasia in Mice and Humans Through PI3K Activation, Arterioscler Thromb
Vasc Biol 38(5) (2018) 1216-1229.
[17] D.A. Fruman, H. Chiu, B.D. Hopkins, S. Bagrodia, L.C. Cantley, R.T. Abraham, The PI3K Pathway
in Human Disease, Cell 170(4) (2017) 605-635.
[18] A. Angulo-Urarte, P. Casado, S.D. Castillo, P. Kobialka, M.P. Kotini, A.M. Figueiredo, P. Castel, V.
Rajeeve, M. Mila-Guasch, J. Millan, C. Wiesner, H. Serra, L. Muixi, O. Casanovas, F. Vinals, M.
Affolter, H. Gerhardt, S. Huveneers, H.G. Belting, P.R. Cutillas, M. Graupera, Endothelial cell
rearrangements during vascular patterning require PI3-kinase-mediated inhibition of actomyosin
contractility, Nat Commun 9(1) (2018) 4826.
[19] W.Y. Aw, C. Cho, H. Wang, A.H. Cooper, E.L. Doherty, D. Rocco, S.A. Huang, S. Kubik, C.P.
Whitworth, R. Armstrong, A.J. Hickey, B. Griffith, M.L. Kutys, J. Blatt, W.J. Polacheck,
Microphysiological model of PIK3CA-driven vascular malformations reveals a role of dysregulated
Rac1 and mTORC1/2 in lesion formation, Sci Adv 9(7) (2023) eade8939.
[20] L. Gambardella, M. Hemberger, B. Hughes, E. Zudaire, S. Andrews, S. Vermeren, PI3K signaling
through the dual GTPase-activating protein ARAP3 is essential for developmental angiogenesis, Sci
Signal 3(145) (2010) ra76.
[21] S. Dimmeler, B. Assmus, C. Hermann, J. Haendeler, A.M. Zeiher, Fluid shear stress stimulates
phosphorylation of Akt in human endothelial cells: involvement in suppression of apoptosis, Circ
Res 83(3) (1998) 334-41.
[22] B.G. Coon, N. Baeyens, J. Han, M. Budatha, T.D. Ross, J.S. Fang, S. Yun, J.L. Thomas, M.A.
Schwartz, Intramembrane binding of VE-cadherin to VEGFR2 and VEGFR3 assembles the
endothelial mechanosensory complex, J Cell Biol 208(7) (2015) 975-86.
[23] E. Tzima, M. Irani-Tehrani, W.B. Kiosses, E. Dejana, D.A. Schultz, B. Engelhardt, G. Cao, H.
DeLisser, M.A. Schwartz, A mechanosensory complex that mediates the endothelial cell response
to fluid shear stress, Nature 437(7057) (2005) 426-31.
[24] L. Bi, I. Okabe, D.J. Bernard, A. Wynshaw-Boris, R.L. Nussbaum, Proliferative defect and
embryonic lethality in mice homozygous for a deletion in the p110alpha subunit of
phosphoinositide 3-kinase, J Biol Chem 274(16) (1999) 10963-8.
[25] M. Petkova, M. Kraft, S. Stritt, I. Martinez-Corral, H. Ortsater, M. Vanlandewijck, B. Jakic, E.
Baselga, S.D. Castillo, M. Graupera, C. Betsholtz, T. Makinen, Immune-interacting lymphatic
endothelial subtype at capillary terminals drives lymphatic malformation, J Exp Med 220(4) (2023).
[26] I. Martinez-Corral, Y. Zhang, M. Petkova, H. Ortsater, S. Sjoberg, S.D. Castillo, P. Brouillard, L.
Libbrecht, D. Saur, M. Graupera, K. Alitalo, L. Boon, M. Vikkula, T. Makinen, Blockade of VEGF-C
signaling inhibits lymphatic malformations driven by oncogenic PIK3CA mutation, Nat Commun
11(1) (2020) 2869.
[27] S.P. Herbert, J. Huisken, T.N. Kim, M.E. Feldman, B.T. Houseman, R.A. Wang, K.M. Shokat, D.Y.
Stainier, Arterial-venous segregation by selective cell sprouting: an alternative mode of blood vessel
formation, Science 326(5950) (2009) 294-8.

[28] K. Kunimoto, Y. Yamamoto, M. Jinnin, ISSVA Classification of Vascular Anomalies and
Molecular Biology, Int J Mol Sci 23(4) (2022).
[29] T.G. Papaioannou, E.N. Karatzis, M. Vavuranakis, J.P. Lekakis, C. Stefanadis, Assessment of
vascular wall shear stress and implications for atherosclerotic disease, Int J Cardiol 113(1) (2006)
12-8.
[30] W.W. Nichols, M.F. O'Rourke, E.R. Edelman, C. Vlachopoulos, McDonald's blood flow in
arteries : theoretical, experimental and clinical principles, CRC Press,, Boca Raton, FL, 2022, p. 1
online resource.
[31] W.J. Polacheck, M.L. Kutys, J. Yang, J. Eyckmans, Y. Wu, H. Vasavada, K.K. Hirschi, C.S. Chen, A
non-canonical Notch complex regulates adherens junctions and vascular barrier function, Nature
552(7684) (2017) 258-262.
[32] G. Charras, A.S. Yap, Tensile Forces and Mechanotransduction at Cell-Cell Junctions, Curr Biol
28(8) (2018) R445-R457.
[33] C. Bertocchi, Y. Wang, A. Ravasio, Y. Hara, Y. Wu, T. Sailov, M.A. Baird, M.W. Davidson, R. Zaidel-
Bar, Y. Toyama, B. Ladoux, R.M. Mege, P. Kanchanawong, Nanoscale architecture of cadherin-based
cell adhesions, Nat Cell Biol 19(1) (2017) 28-37.
[34] T.J. Kim, S. Zheng, J. Sun, I. Muhamed, J. Wu, L. Lei, X. Kong, D.E. Leckband, Y. Wang, Dynamic
visualization of alpha-catenin reveals rapid, reversible conformation switching between tension
states, Curr Biol 25(2) (2015) 218-224.
[35] M. Yao, W. Qiu, R. Liu, A.K. Efremov, P. Cong, R. Seddiki, M. Payre, C.T. Lim, B. Ladoux, R.M.
Mege, J. Yan, Force-dependent conformational switch of alpha-catenin controls vinculin binding,
Nat Commun 5 (2014) 4525.
[36] D.L. Huang, N.A. Bax, C.D. Buckley, W.I. Weis, A.R. Dunn, Vinculin forms a directionally
asymmetric catch bond with F-actin, Science 357(6352) (2017) 703-706.
[37] W.J. Polacheck, M.L. Kutys, J.B. Tefft, C.S. Chen, Microfabricated blood vessels for modeling the
vascular transport barrier, Nat Protoc 14(5) (2019) 1425-1454.
[38] H. Wiig, M.A. Swartz, Interstitial fluid and lymph formation and transport: physiological
regulation and roles in inflammation and cancer, Physiol Rev 92(3) (2012) 1005-60.
[39] G.H. Lee, S.A. Huang, W.Y. Aw, M.L. Rathod, C. Cho, F.S. Ligler, W.J. Polacheck, Multilayer
microfluidic platform for the study of luminal, transmural, and interstitial flow, Biofabrication 14(2)
(2022).
[40] D.-H.T. Nguyen, S.C. Stapleton, M.T. Yang, S.S. Cha, C.K. Choi, P.A. Galie, C.S. Chen,
Biomimetic model to reconstitute angiogenic sprouting morphogenesis in vitro, Proceedings of the
National Academy of Sciences 110(17) (2013) 6712-6717.
[41] V. Vickerman, R.D. Kamm, Mechanism of a flow-gated angiogenesis switch: early signaling
events at cell-matrix and cell-cell junctions, Integr Biol (Camb) 4(8) (2012) 863-74.
[42] J.W. Song, L.L. Munn, Fluid forces control endothelial sprouting, Proc Natl Acad Sci U S A
108(37) (2011) 15342-7.
[43] Y. Abe, M. Watanabe, S. Chung, R.D. Kamm, K. Tanishita, R. Sudo, Balance of interstitial flow
magnitude and vascular endothelial growth factor concentration modulates three-dimensional
microvascular network formation, APL Bioeng 3(3) (2019) 036102.
[44] S. Quintero-Fabian, R. Arreola, E. Becerril-Villanueva, J.C. Torres-Romero, V. Arana-Argaez, J.
Lara-Riegos, M.A. Ramirez-Camacho, M.E. Alvarez-Sanchez, Role of Matrix Metalloproteinases in
Angiogenesis and Cancer, Front Oncol 9 (2019) 1370.
[45] T. Harada, J. Swift, J. Irianto, J.W. Shin, K.R. Spinler, A. Athirasala, R. Diegmiller, P.C. Dingal, I.L.
Ivanovska, D.E. Discher, Nuclear lamin stiffness is a barrier to 3D migration, but softness can limit
survival, J Cell Biol 204(5) (2014) 669-82.

[46] C.P. Whitworth, W.Y. Aw, E.L. Doherty, C. Handler, Y. Ambekar, A. Sawhney, G. Scarcelli, W.J.
Polacheck, P300 Modulates Endothelial Mechanotransduction of Fluid Shear Stress, Cellular and
Molecular Bioengineering (2024).
[47] P.M. Davidson, G.R. Fedorchak, S. Mondesert-Deveraux, E.S. Bell, P. Isermann, D. Aubry, R.
Allena, J. Lammerding, High-throughput microfluidic micropipette aspiration device to probe time-
scale dependent nuclear mechanics in intact cells, Lab Chip 19(21) (2019) 3652-3663.
[48] A. Queisser, L.M. Boon, M. Vikkula, Etiology and Genetics of Congenital Vascular Lesions,
Otolaryngol Clin North Am 51(1) (2018) 41-53.
[49] C. Aitken, V. Mehta, M.A. Schwartz, E. Tzima, Mechanisms of endothelial flow sensing, Nature
Cardiovascular Research 2(6) (2023) 517-529.
[50] Y.S. Li, J.H. Haga, S. Chien, Molecular basis of the effects of shear stress on vascular
endothelial cells, J Biomech 38(10) (2005) 1949-71.
[51] M.A. Gimbrone, Jr., J.N. Topper, T. Nagel, K.R. Anderson, G. Garcia-Cardena, Endothelial
dysfunction, hemodynamic forces, and atherogenesis, Ann N Y Acad Sci 902 (2000) 230-9;
discussion 239-40.
[52] Z. Liu, J.L. Tan, D.M. Cohen, M.T. Yang, N.J. Sniadecki, S.A. Ruiz, C.M. Nelson, C.S. Chen,
Mechanical tugging force regulates the size of cell-cell junctions, Proc Natl Acad Sci U S A 107(22)
(2010) 9944-9.
[53] Q. Chen, L. Jiang, C. Li, D. Hu, J.W. Bu, D. Cai, J.L. Du, Haemodynamics-driven developmental
pruning of brain vasculature in zebrafish, PLoS Biol 10(8) (2012) e1001374.
[54] T.W. Secomb, J.P. Alberding, R. Hsu, M.W. Dewhirst, A.R. Pries, Angiogenesis: an adaptive
dynamic biological patterning problem, PLoS Comput Biol 9(3) (2013) e1002983.
[55] A.R. Pries, T.W. Secomb, Making microvascular networks work: angiogenesis, remodeling, and
pruning, Physiology (Bethesda) 29(6) (2014) 446-55.
[56] A.R. Pries, B. Reglin, T.W. Secomb, Structural adaptation of vascular networks: role of the
pressure response, Hypertension 38(6) (2001) 1476-9.
[57] P.A. Galie, D.-H.T. Nguyen, C.K. Choi, D.M. Cohen, P.A. Janmey, C.S. Chen, Fluid shear stress
threshold regulates angiogenic sprouting, Proceedings of the National Academy of Sciences
111(22) (2014) 7968-7973.
[58] K. Wolf, M. Te Lindert, M. Krause, S. Alexander, J. Te Riet, A.L. Willis, R.M. Hoffman, C.G. Figdor,
S.J. Weiss, P. Friedl, Physical limits of cell migration: control by ECM space and nuclear deformation
and tuning by proteolysis and traction force, J Cell Biol 201(7) (2013) 1069-84.
[59] Y. Samuels, L.A. Diaz, Jr., O. Schmidt-Kittler, J.M. Cummins, L. Delong, I. Cheong, C. Rago, D.L.
Huso, C. Lengauer, K.W. Kinzler, B. Vogelstein, V.E. Velculescu, Mutant PIK3CA promotes cell
growth and invasion of human cancer cells, Cancer Cell 7(6) (2005) 561-73.
[60] G.L. Weber, M.O. Parat, Z.A. Binder, G.L. Gallia, G.J. Riggins, Abrogation of PIK3CA or PIK3R1
reduces proliferation, migration, and invasion in glioblastoma multiforme cells, Oncotarget 2(11)
(2011) 833-49.
[61] M.M. Salek, P. Sattari, R.J. Martinuzzi, Analysis of fluid flow and wall shear stress patterns
inside partially filled agitated culture well plates, Ann Biomed Eng 40(3) (2012) 707-28.
[62] M.L. Rathod, W.Y. Aw, S. Huang, J. Lu, E.L. Doherty, C.P. Whithworth, G. Xi, P. Roy-Chaudhury,
W.J. Polacheck, Donor-Derived Engineered Microvessels for Cardiovascular Risk Stratification of
Patients with Kidney Failure, Small (2024) e2307901.

924