Title: Tracheal terminal cells of *Drosophila* are immune privileged to maintain their Foxo-dependent structural plasticity

Authors: Judith Bossen [1,2*], Larissa Fritz[1], Reshmi Raveendran[1], Leizhi Shi[1,2], Jingjing He[1] & Thomas Roeder[1,2*]

Affiliations:

[1] Kiel University, Zoology, Dept. Molecular Physiology; Kiel, Germany.

[2] Airway Research Center North (ARCN), German Center for Lung Research (DZL).

*Corresponding authors: Judith Bossen, email: jbossen@zoologie.uni-kiel.de, Thomas Roeder, email: troeder@zoologie.uni-kiel.de.

Abstract:

Respiratory organs must balance their primary function of gas exchange with the constant threat of inhaled pathogens. In the *Drosophila* tracheal system, gas exchange occurs at the tracheal terminal cells (TTCs), the functional equivalents of mammalian alveoli. While bacterial infection triggers a robust innate immune response throughout the broader airway epithelium, we reveal that TTCs are uniquely exempt from this reaction. Mechanistically, TTCs lack expression of the membrane-associated peptidoglycan recognition receptor PGRP-LC. This absence protects these highly susceptible cells from Immune deficiency (Imd) pathway activation and subsequent JNK-mediated cell death, establishing TTCs as a distinct, immune-privileged niche. Ectopic immune activation via targeted *PGRP-LCx* overexpression in TTCs caused a severe reduction in branching, cellular damage, and ultimately cell death, phenotypes that were fully rescued by the depletion of AP-1 or *foxo*. Because both structural plasticity (in response to nutritional cues and hypoxia) and innate immune responses strictly require the transcription factor FoxO, we demonstrate that potent immune signaling is fundamentally incompatible with dynamic TTC remodeling. Ultimately, the immune-privileged status of TTCs represents an essential evolutionary trade-off, restricting local inflammation to preserve *foxo*-dependent structural plasticity and vital respiratory function.

Short title: Tracheal terminal cells of *Drosophila* are immune privileged to maintain their Foxo-dependent structural plasticity.

INTRODUCTION

The tracheal system of *Drosophila* comprises three different parts: the main dorsal trunks, the smaller tracheal branches, and the terminal cells (Whitten 1957; Uv et al. 2003). The tracheal terminal cells (TTCs) form the most distal part of the respiratory system. They are the site at which gas exchange takes place. TTCs fulfill an essential role by supplying all body organs with oxygen. To fulfill their task, even under ever-changing conditions, these cells possess a high degree of structural plasticity. For example, TTCs can sprout in response to local hypoxia (Jarecki et al. 1999), or in response to differences in nutrient availability (Centanin et al. 2010); this process is analogous to angiogenesis in mammals. Sprouting is dependent on growth factors, as well as the hypoxia-inducible factor (HIF)-α homolog sima, again analogous to endothelial cells during angiogenesis in the mammalian vascular system (Jarecki et al. 1999; Centanin et al. 2008; Horowitz and Simons 2008).

The primary role of the proximal tracheal tubes is to conduct air to more distal parts. The epithelial cells lining these tubes can mount an effective immune response to infections (Wagner et al. 2008; Faisal et al. 2014). This immune response is driven mainly by expression of several antimicrobial peptides (AMP) (Tzou et al. 2000; Akhouayri et al. 2011). One out of the two major immune pathways operative in *Drosophila*, the immune deficiency (Imd) pathway, is solely responsible for responses in the larval airway epithelium; this is because the Toll pathway is not functional in these cells (Wagner et al. 2008) Klicken oder tippen Sie hier, um Text einzugeben.. The Imd signaling pathway is homologous to the human tumor necrosis factor-α (TNF-α) pathway and converges on activation of NF-κB factors. Moreover, it is connected to c-Jun N-terminal kinase (JNK) signaling via transforming growth factor-β (TGF-β) activated kinase 1 (dTak1) (Silverman et al. 2003). Infection of the airway system induces an immune response by the tracheal epithelium, which includes expression of canonical Imd and JNK target genes (Tzou et al. 2000; Wagner et al. 2009). Chronic activation of tracheal immune signaling leads to marked structural changes in the epithelium

(Wagner et al. 2021). This tissue remodeling is mediated by JNK and its downstream

transcription factor, forkhead box sub-group O (foxo), which is a as a terminal target of the

epithelial immune system. FoxOs are also central proteins of the insulin signaling pathway,

which also plays an important role in the structural plasticity of TTCs (Linneweber et al.

2014; Wong et al. 2014; Burguete et al. 2019).

The present study shows that TTCs differ fundamentally from the rest of the tracheal

epithelium. In the case of a natural infection of the tracheal system, these cells show only a

negligible immune response. The very few TTCs that do show AMP expression are

structurally impaired. We found that TTCs are immune privileged. The Imd pathway in TTCs

is deactivated since the cells do not express the transmembrane peptidoglycan recognition

protein (PGRP)-LCx. Chronic activation of the Imd signaling pathway in TTCs leads to JNK-

dependent apoptosis. Therefore, we hypothesize that the immune privileged status of TTCs

maintains the normal function of foxo, which acts as a regulator of structural plasticity.

RESULTS

Tracheal infection revealed that AMP expression in terminal cells is rare

Larvae carrying reporters for antimicrobial peptide genes were studied following infection with *Pectobacterium carotovorum* (formerly *Erwinia carotovora*) or under control conditions. These larvae express GFP under the control of promoters for different antimicrobial peptide (AMP) genes, which are known to be activated by a natural infection in different tissues (Tzou et al. 2000). GFP fluorescence was analyzed after 24 h, focusing on the TTCs (Fig. 1). Without infection AMP expression was rarely or never observed in the tracheal system (Fig. 1A-E). Expression in the TTCs was never observed. In infected animals, AMP expression was observed frequently in the tracheal system, but again, never in the TTCs (Fig. 1A'-E'). The emergence of AMP expression was quantified by counting the number of animals with fluorescence in the 3^{rd} tracheal metamere with focus on the dorsal part, especially in the dorsal branch/fusion cells (Fig. 1F; GFP+, light green) and fluorescence in the TTCs (GFP TTCs, dark green). In 5-33% of the larvae, fluorescence was detected in the dorsal branch (DB)/fusion cell (FC) region; this number increased after infection to 24-70 % (Fig. 1G). But infection could not trigger an antimicrobial peptide response in the TTCs. In addition to a response in the trachea, fluorescence was observed in various tissues, such as hemocytes and fat body, especially in the anterior part, indicating the animals' general responsiveness to infection (Fig. S1A-D).

For a more detailed analysis, we focused on the AMP Drosomycin, (*Drs*) which showed the strongest response to infection in the trachea. A total of 169 infected larvae were examined over all tracheal metameres. About 34% showed no GFP signal in the focus area, a finding consistent with previous observations (Akhouayri et al. 2011). The remaining larvae showed a GFP signal in the DB/FCs (57.4%, Fig. 2A, A'). The absence of a response in the TTCs was not limited to the dorsal TTCs but was also observed in TTCs supplying other body parts

(Fig. 2B). However, 8.3% showed *Drs* expression in TTCs (Fig. 2C-G). Most of the few

fluorescent TTCs showed clear structural differences from TTCs without a GFP signal

(Fig. 2D, D'). Although some cells had a normal structure (Fig. 2E, E'), most had shortened

branches (Fig. 2D, D', F, F'), showed signs of melanization (Fig. 2G, arrow), or were no

longer air-filled (Fig. 2G, arrowhead). These observations suggest an immune triggered

response in the TTC itself that act negatively on these cells.

Tracheal terminal cells do not express the Imd-pathway receptor PGRP-LCx

We wondered whether constitutive immune activation within the tracheal system would

exclude TTCs, as expected from the results of the infection experiments. To ensure that all

parts of the tracheal epithelium, including TTCs, were exposed to this activating stimulus, we

ectopically expressed the secreted (and intracellular) Imd pathway receptor PGRP-LE using

the *ppk4-Gal4* driver. AMP expression (a readout of the immune response) was visualized by

simultaneous expression of GFP-tagged *Drs* (Fig. 3). A strong immune response was

observed in the tracheal tissue, from the dorsal trunks down to the smaller branches (Fig. 3A–

C). While expression of *Drs* was consistent throughout most parts of the tracheal system, it

was absent from the most distal parts (Fig. 3B–E). Closer observation of the TTCs revealed a

distinct breakpoint in the GFP signal at the proximal end of the TTCs. This was true for TTCs

attached to the outer cuticle (Fig. 3D) as well as for cells attached to the intestine (Fig. 3E).

Activation of the Imd pathway may not be fully functional in these cells. We infected the

animals and quantified the corresponding response in the 3^{rd} tracheal metamere. We observed

an increased response upon infection and a few fluorescence events in the TTCs in 3% of the

animals (Fig. 3F, G).

To activate the Imd pathway successfully, secreted PGRP-LE has to bind to transmembrane

PGRP-LC, meaning that only cells expressing PGRP-LC can be activated (Takehana et al.

2002; Takehana et al. 2004). Therefore, to investigate PGRP-LC expression in the tracheal

system, specifically in TTCs, we expressed *Gal4* controlled by the *PGRP-LC* promoter to

drive concurrent GFP expression (Fig. 3H, I). While GFP was visible throughout the entire

tracheal system, it was absent from all TTCs associated with the cuticle (Fig. 3H, H') and

intestine (Fig. 3I, I'). To demonstrate that GFP expression can be visualized in TTCs, and that

the lack of a signal was not an artifact, expression of GFP in TTCs was driven by the tracheal

driver *btl-Gal4* (*btl-Gal4; UAS-GFP*) (Fig. 3J, K). TTCs on the dissected cuticle (Fig. 3J) and

on the intestine (Fig. 3K) expressed GFP. Similarly, as before, we infected the *PGRP-LCx-P-*

Gal4 larvae and observed an increased number of animals with fluorescence in the 3rd

metamere DB/FCs but no animals with fluorescence in the TTCs (Fig. 3L).

Thus, the data indicate that lack of expression of the Imd-pathway receptor PGRP-LC by

TTCs is the reason for the difference between TTCs and cells within the rest of the tracheal

system. The observation that only 3 % of the PGRP-LCx expressing animals show *Drs*

expression in the TTCs after infection (Fig. 3F, G) opens up alternative modes of induction

including potential stress responses mediated through e.g., Foxo activation.

Ectopic activation of PGRP-LCx in the tracheal system leads to the death of TTCs

To further investigate the effects of PGRP-LCx-mediated immune responses on the tracheal

system, we expressed *PGRP-LCx* exclusively in TTCs (*PGRP-LCx^{OE}*; Fig. 4). We found that

Imd-mediated immune activation was restricted to these cells, as indicated by concurrent

expression of GFP. Compared with the widely branched and diversified TTCs in control

samples (Fig. 3A), *PGRP-LCx*-expressing TTCs showed a striking phenotype characterized

by smaller TTCs with shortened terminal branches (Fig. 4A, B). Measurement of TTC

branches in *PGRP-LCx*-expressing larvae revealed a significant reduction in the number and

length of branches compared with the controls (Fig. 4C, D). While cells in the control had

about 14 branches, with a total length of about 1550 µm, TTCs in PGRP-LCx^{OE} (III) insects

had seven branches at the most, and the length did not exceed 514 µm. This means that the

branched surface of TTCs expressing *PGRP-LCx* was reduced to less than one-third of that in

controls. Additionally, we tested another PGRP-LCx expressing line (PGRP-LCx[OE] (II), also

used later for mechanistic evaluation). We could see a less severe but still prominent TTC-

phenotype (Fig. 4C, D). To prove the less severe phenotype, we drove PGRP-LCx[OE] (III) and

(II) in the dorsal trunks of the trachea and compared the tracheal thickness (Fig. S2A-D). The

PGRP-LC-dependent epithelial thickness phenotype was shown in a previous study (Wagner

et al. 2021). PGRP-LCx[OE] (III) produced a significantly increased thickening phenotype

compared to PGRP-LCx[OE] (II) (Fig. S2A-D).

To investigate whether this cellular phenotype has an impact on physiological oxygen supply,

we exposed larvae to low oxygen levels (2–3 %). Leaving the lawn under hypoxic conditions

is the normal response of larvae, and this response can be monitored over time. A

significantly higher percentage of larvae with TTCs overexpressing *PGRP-LCx* showed the

escape phenotype after 15–25 minutes of exposure (Fig. 4E, S4). Activation of the Imd

pathway by ectopic expression of *PGRP-LCx* clearly impairs TTCs, not only with respect to

size reduction, but also to the functionality of these cells (i.e., the enhanced response to

hypoxic conditions). To elucidate whether this size reduction is part of an apoptotic cell

program, we analyzed expression of two apoptotic factors: head involution defective (hid) and

reaper (rpr) (White et al. 1994; Grether et al. 1995). We targeted expression of these factors to

TTCs using the *DSRF*-Gal4 driver line. Larvae expressing *hid; rpr* showed a strong response

to hypoxia by leaving the lawn at the very early larval stages; these larvae did not survive

(Fig. 4F). Since PGRP LCx-expressing larvae survived and did not show this extreme

phenotype, we examined Tak1 (a downstream component of the Imd pathway). Expression of

Tak1 in TTCs also resulted in early larval death, similar to expression of *hid; rpr* (Fig. 4F).

Dissection of *hid; rpr*-expressing insects revealed a complete absence of TTCs, with only

rudimentary cell remnants connected to the intestine (Fig. 3G, H). By contrast, overexpression

of *PGRP-LCx* in TTCs resulted in similar cellular alterations, although the cells remained

alive. Nevertheless, the overall phenotype was strikingly similar to that of apoptotic terminal

cells (Fig. 4I).

Drosophila death caspase-1 (Dcp-1) is orthologous to mammalian Caspase-3, and is an

established marker of apoptosis (*21*). Immunohistochemical analysis using an anti-Dcp-1

antibody (Fig. 4J, K) revealed a lack of Dcp-1 in the control; however, strong GFP staining of

TTCs was visible (Fig. 4J-J"). TTCs expressing *PGRP-LCx* were positive for cleaved Dcp-1,

whereas counterstaining for GFP revealed reduced expression (Fig. 4K-K"). These data could

indicate that TTCs are undergoing apoptosis in consequence of PGRP-LCx overexpression.

To complete our results, we stained trachea with PGRP-LCx expression in the tracheal

epithelium and saw an increase of cleaved Dcp-1 in these tracheae as well (Fig. S3).

We expressed PGRP-LE in the trachea for only 24 h to induce a mild activation of the

pathway in TTCs and epithelial cells (using *btl-Gal4; tubGal80ts*) and compared hypoxia

response under control and infected conditions (Fig. S4). While we saw the same increase in

the escape response of PGRP-LE larvae under control conditions, the response was even more

enhanced for the infected larvae.

PGRP-LCx-induced terminal cell death is mediated by JNK signaling

The *Drosophila* Imd signaling pathway, which classically leads to activation of the Nf-κB

factor Relish, is homologous to the TNF-α signaling pathway in mammals. Like the TNF-α

pathway, the Imd pathway branches out to the JNK signaling pathway; the branch point is

Tak1 (Fig. 5A). To test whether the Nf-κB or the JNK branch is necessary to translate PGRP-

LCx activation in TTCs into triggering of apoptosis, we expressed these downstream

components in flies, which were then phenotyped. The number and length of the terminal

branches were measured and compared with those in control TTCs, and with the phenotype

produced by *PGRP-LCx* expression (Fig. 5B–F). Ectopic expression of an activated *Relish*

allele had no effect on the TTC branching phenotype (Fig. 5B, E, F). Expression of a

constitutively active form of dJNKK, *hemipterous* (hep^{CA}), resulted in a phenotype like that

caused by expression of the upstream receptor PGRP-LCx (Fig. 5C, E, F). Branching of TTCs

was reduced significantly and did not differ significantly from that of TTCs with *PGRP-LCx*

expression. Overexpression of the wild-type JNK, *basket* (bsk^{OE}), also resulted in reduced

branching when compared with the control (Fig. 5D–F). These results suggest that JNK

activation is responsible for the TTC phenotype. To further support this hypothesis, the cells

were stained for the Nf-κB factor Relish (Rel) and for the phosphorylated JNK (pJNK) basket

(Fig. 5G, H).

Staining of the transcription factor Relish was observed throughout intestinal tissue, as well as

in TTCs, from controls, and in TTCs expressing *PGRP-LCx*; however, we observed no

translocation of Relish to the nucleus (Fig. 5G, arrow). In contrast to controls, *PGRP-LCx-*

expressing cells showed distinct nuclear staining by an anti-pJNK antibody (Fig. 5H, arrow).

Moreover, nuclear pJNK was also observed in the nuclei of subjacent intestinal cells. This

suggests that intestinal cells experience some additional JNK-mediated stress.

JNK-mediated TTC damage can be rescued by AP-1 and foxo depletion

The *Drosophila* JNK signaling pathway induces apoptosis (Adachi-Yamada et al. 1999;

Takatsu et al. 2000). The canonical transcription factors (TFs) AP-1, foxo and Ets21C act

downstream of Tak1 and JNK to initiate transcription of target genes (Fig. 6A). Activation of

each of these factors can lead to expression of apoptotic genes (Hafezi et al. 1997; Lehmann

et al. 2002; Luo et al. 2007; Mundorf et al. 2019). Klicken oder tippen Sie hier, um Text

einzugeben.To evaluate whether one of these TFs is responsible for the *PGRP-LCx^{OE}*-induced

phenotype, we silenced each of them. This experimental approach required the use of

different controls with the *PGRP-LCx^{OE}* cassette located on different chromosomes. Beside

the PGRP-LCx^{OE} (III) on the 3rd chromosome, we used PGRP-LCx^{OE} (II) which produces a

less severe phenotype and is located on the 2nd chromosome (Fig. 6F, G; red and blue). First,

we used a dominant-negative form of *Tak1* (*Tak1^{DN}*) and *bsk* (*bsk^{DN}*) together with *PGRP-*

LCx^{OE} to confirm that PGRL-LC-induced apoptosis of TTCs is mediated by JNK. Concurrent

expression of *PGRP-LCx^{OE}* with either *Tak1^{DN}* or *bsk^{DN}* rescued the apoptotic phenotype (Fig.

6B, F, G). Fos and Jun proteins form homo- or heterodimers that act as AP-1 transcription

factors, also a canonical target of the JNK pathway. To target AP-1, we focused on the

Drosophila Fos ortholog kayak (kay). We co-expressed a dominant-negative *kay* allele

(*kay^{DN}*) together with *PGRP-LCx^{OE}* in TTCs. Although *kay^{DN}* rescued the *PGRP-LCx^{OE}*

phenotype with respect to the number and length of branches, it was not complete (Fig. 6C, F,

G; red). Co-expression of *foxo^{RNAi}* rescued the *PGRP-LCx^{OE}* phenotype to the same level as

kay^{DN} (Fig. 6D, F, G; red). RNAi of *Ets21C* failed to rescue the *PGRP-LCx^{OE}* phenotype (Fig.

6E, F, G; blue); rather, it worsened the phenotype with respect to the length of branches.

Because depletion of the TFs foxo and kayak rescued the TTC phenotype induced by *PGRP-*

LCx overexpression, we next tested whether upregulation of either induces a similar TTC

phenotype (Fig. 6H–K). Overexpression of *foxo* led to a slightly reduction in the number and

length of the branches (Fig. 6H–J), while overexpression of a combination of *kay* and *Jra*,

which drives concurrent overexpression of both AP-1 components, led to a severe reduction

in number and length of the branches, to levels below that induced by *PGRP-LCx*

overexpression (Fig. 6H–J, K). Therefore, we asked whether foxo and AP-1 are present and

active in TTCs. To do this, we used an AP-1 responsive TRE-RFP reporter line to detect AP-1

activity in TTCs (Chatterjee and Bohmann 2012). We observed AP-1 activity in wild-type

TTCs, and upon overexpression of *PGRP-LCx* (Fig. 6L, M). When we used a *Gal4* line with a

foxo promoter, we observed strong *foxo* promoter activity in all TTCs (Fig. 6N). Thus, both

TFs appear to be expressed and functional in TTCs.

Foxo controls TTC branching under normal conditions and under conditions of oxygen

deprivation

A previous study reported a role for foxo in maintaining homeostasis of tracheal epithelial

cells (Wagner et al. 2021). One of the main features that ensures full functionality of TTCs

under changing conditions is the ability to adapt to local hypoxia to maintain the supply of

oxygen to target tissues; therefore, we subjected wild-type larvae and larvae with foxo-RNAi

in TTCs to mild hypoxia (5% O_2) and measured their branching ability (Fig. 7). As expected,

wild-type TTCs responded with increased branching (Fig. 7A, B, E). The $foxo^{RNAi}$ TTCs

already had an increased number of branches under control conditions (21% O_2; Fig. 7C–E),

and the number of branches did not increase further after oxygen deprivation; this suggest that

foxo is required for appropriate responses to hypoxic conditions (Fig. 7D, E).

In summary, the data suggest that TTCs differ from the other parts of the tracheal system in

terms of immune pathway activation (Imd/JNK); this is a mechanism that circumvents cell

death and prevents impairment of functionality (Fig. 8A). The Imd pathway receptor *PGRP-*

LCx does not appear to be expressed in TTCs, indicating that no downstream signaling takes

place (Fig. 8B). Ectopic expression of *PGRP-LCx* exclusively in these cells leads to JNK-

mediated cell death; however, all downstream components must be present and functional.

We have shown that depletion of the JNK-associated TFs AP-1 and foxo can rescue the

PGRP-LCx-mediated phenotype. Both are present under physiological conditions, and their

overexpression can induce a severe TTC phenotype. In addition, the universal transcription

factor foxo, which is not only associated with JNK signaling, plays a role in branching of

TTCs under normal and hypoxic conditions, indicating its importance for TTC homeostasis.

DISCUSSION

Activation of the immune system is a double-edged sword in that although it fights infection,

it can also damage the organism's tissues (Ruff et al. 2020); therefore, appropriate regulation

and restriction of the immune system is of prime importance. Tight regulation of the immune

system is essential when delicate, sensitive tissues are involved, or when immune-relevant

signaling pathways are also needed for other cellular functions. Tracheal TTCs appear to fall

into both categories; they are susceptible due to their specific structure but require foxo

signaling to maintain structural plasticity. Here, we found that TTCs, in contrast to the other

epithelial cells in the trachea, barely respond to bacterial infection. Immune-privileged status

might protect TTCs from damage caused by a strong immune response, a strategy that also

protects other organ systems (Galea et al. 2007; Forrester et al. 2008; Hill et al. 2021). Here,

we found that the entire tracheal epithelium expressed the transmembrane receptor PGRP-LC,

but not TTCs. We propose that the lack of *PGRP-LCx* expression by TTCs is the reason for

their lack of an immune response. This hypothesis is supported by the observation that

targeted overexpression of *PGRP-LC* in TTCs induced an immune response, suggesting that

all other pathway components are active in these cells. Since targeted overexpression of

PGRP-LC leads to degeneration and, ultimately, apoptosis of these cells, this finding is

particularly interesting. Targeted induction of apoptosis in TTCs by *hid* and *rpr* induced a

similar, but more severe, phenotype; indeed, these larvae die after complete ablation of the

cells. The weaker phenotype we observe in *PGRP-LCx* expressing TTCs could be also

explained by a non-apoptotic function of Dcp-1. Dcp-1 encodes for an effector caspase during

the apoptotic process but does also have non-apoptotic functions. Dcp-1 is involved in

autophagy regulation and mitochondrial morphology (Hou et al. 2009; DeVorkin et al. 2014).

In our case Dcp-1 is not only cleaved upon *PGRP-LCx* expression in the TTCs, but also in the

tracheal epithelial cells. As described, *PGRP-LCx* expression in these cells does mediate

tissue hyperplasia and renewal (Wagner et al. 2021). Furthermore, a role in inhibition of

innate Toll signaling by cleavage of NF-κBs was reported (Wu et al. 2022).

Herein, we show that A) TTCs respond to a strong immune response by undergoing

apoptosis, which leads to the death of the animal; and B) that immune activation is prevented

by a lack of expression of the central receptor of the Imd pathway, PGRP-LCx. To better

understand the first finding, we needed to elucidate the mechanism of apoptosis induction on

the one hand and to identify mechanisms that still protect against infection on the other.

Activation of the Imd pathway by ectopic expression of *PGRP-LCx* throughout the trachea

leads to JNK-dependent meta- and hyperplasia of the trunk cells (Wagner et al. 2021). The

JNK pathway, as well as activation of its canonical TFs, is associated with both proliferation

and apoptosis. The intertwining of the Imd pathway with the JNK pathway occurs at the level

of Tak1 and is a general organizational principle within these signaling pathways. This

architecture means that strong activation of the Imd pathway in these cells also triggers the

multifaceted JNK pathway. Downstream of JNK (*bsk* in *Drosophila*), we found that foxo and

AP-1 are both necessary and sufficient for the apoptotic phenotype in TTCs. AP-1 is a

classical TF that is activated by JNK signaling. The reason underlying the incomplete rescue

is unknown but might be due to a specific role for foxo in this context.

We also demonstrated that depleting *foxo* rescues the apoptotic phenotype, at least in part.

Although the effects triggered by *foxo* depletion or *foxo* overexpression were statistically

significant, they were certainly not as strong as those triggered by AP-1. This difference in

efficiency is because the tools used to manipulate foxo (RNAi and overexpression) are much

weaker than those used to manipulate AP-1 signaling. For example, in the latter case, a

dominant-negative form of *kay* and a fully active heterodimer of *Jra* and *kay* were available;

in the case of foxo, the overexpression allele does not give rise to an activated version of foxo,

and RNAi is generally less effective than dominant-negative alleles. In the tracheal cells of

the dorsal trunk, and the primary and secondary branches, foxo plays a role in tracheal

remodeling downstream of Imd pathway activation (Wagner et al. 2021). It seems reasonable

then that foxo would also be activated in TTCs downstream of Imd signaling, which would in

turn disrupt branching control and thus TTC functionality, as well as the ability to respond to

environmental changes. Our central hypothesis derived from these observations is as follows:

since foxo is essential for structural plasticity, and this property is critical for the survival of

the organism, foxo should not be activated by other signals such as those from the Imd

pathway. Foxo is activated not only by JNK signaling but also is a canonical member of the

insulin signaling pathway. Linneweber and colleagues showed that insulin signaling, and

especially the insulin receptor, is necessary for TTCs to respond (i.e., with reduced branching)

to changing nutritional conditions such as low protein concentrations in food (Linneweber et

al. 2014). Moreover, different insulin receptor alleles affect the size of TTCs (Burguete et al.

2019). Here, we showed that foxo is also part of this signaling pathway and that it is

(presumably) required to translate different insulin signaling activities into structural changes

in TTC branching. A very similar scenario for FoxO signaling is operative in endothelial cells

(ECs). Insulin and VEGF signaling in human ECs targets FoxO to control its transcriptional

activity directly via PI3K/Akt signaling (Abid et al. 2004; Abid et al. 2006).

An alternative explanation for the observed lack of an immune response in TTCs could be

their maximal distance from the spiracles. In this scenario, a gradient of bacterial inducers

along the tracheal system might be expected, resulting in a gradual decrease in immune

activation from the spiracles toward the TTCs. However, this is not what we observed. In

tracheae that displayed an immune response, the response was largely homogeneous along the

entire length of the tracheal system, from the spiracles to the TTCs. Only at the transition to

the TTCs did the immune response drop abruptly. This observation argues against the

gradient hypothesis and suggests that TTCs are specifically excluded from the immune

response.

Following our discussion of why it is sensible and necessary to omit this vital part of

epithelial immunity from TTCs, we will focus on alternative strategies that operate to ensure

that the cells are protected against infection. The strong immune response of the tracheal

epithelium may be sufficient to protect the trachea, making an immune response in distant

TTCs unnecessary. This strong and efficient immune response in the airway epithelia includes

contact structures that connect the tracheal system with the outside world: the dorsal trunks

and the primary and secondary branches (Wagner et al. 2008; Davis and Engstrom 2012;

League and Hillyer 2016; Bossen et al. 2023). TTCs reside at the very ends of these systems,

so potential pathogens would have to travel a long distance and escape powerful immune

responses before reaching the TTCs. In addition, the diameters of 0.1–1 μm are so small that

most bacteria will not gain access (Jarecki et al. 1999).

The mechanisms underlying immune privilege vary. The same is true for the mechanisms that

dampen the immune response. Here, we found that the dampening of the immune response

was almost complete (based on non-expression of the proximal receptor within the canonical

signaling pathway controlling epithelial immunity). A similar strategy was reported in a

subpopulation of intestinal epithelial cells, called enterocytes that do not express the Imd

receptor *PGRP-LCx* (Fink et al. 2016). It was hypothesized that this prevents chronic MAMP-

mediated activation of innate epithelial immune responses in these cells, which are constantly

exposed to gut bacteria. Organs such as the liver, kidney, lymphoid organs, brain, and lung

are immunotolerant (Amersfoort et al. 2022). Relevant immune pathways that induce

apoptosis are dampened to different extents. In *Drosophila* epithelia, this dampening applies

primarily to the Imd pathways because Toll signaling is not functional in these structures

(Ferrandon et al. 1998; Wagner et al. 2008) Interestingly, the mammalian homolog of the Imd

pathway, the TNF-α pathway, also invokes these protective strategies. This can be seen in

ECs, which are the first cells to meet pathogens in the circulation and share some

characteristics with TTCs. For example, they express several PRRs, as well as secrete pro-

inflammatory cytokines to initiate immune responses (Mai et al. 2013) To prevent EC death

and apoptosis, protective genes and negative feedback loops become active when immune

pathways are triggered (Bach et al. 1997). For example, VEGF/VEGFR signaling is

immunosuppressive (Yang et al. 2018) because it inhibits TNF-α induced apoptosis of ECs;

VEGF inhibits secretion of TNF in a concentration-dependent manner (Spyridopoulos et al.

1997). For example, during angiogenesis and VEGFR activation, ECs express the anti-

apoptotic gene *surviving*, which reduces caspase-3 activity and inhibits TNF-α-mediated

apoptosis (O'Connor et al. 2000). The anti-apoptotic gene A20, when expressed in ECs,

protects them from TNF-α-induced cell death, and inhibits inflammatory responses triggered

by NF-κB activation (Daniel et al. 2004). Trabid, the equivalent of A20 in flies, is a negative

regulator of the Imd signaling pathway (Fernando et al. 2014) TNF-α signaling, which

induces death of various cell types, is a classical apoptotic pathway. TNF-α-induced apoptosis

is highly regulated in ECs, which resist TNF-α-induced and physiological inflammatory

apoptosis via TAK1 (Naito et al. 2019), which also regulates necroptosis and metastasis of

ECs (Yang et al. 2019).

A recent study from our lab shows an inability of TTCs to respond to low oxygen level upon

Trabid knockout, which supports our hypothesis that this signaling pathway converging onto

foxo activation is necessary for this response (Bossen et al. 2025). The branching process in

TTCs is comparable with that observed during angiogenesis, in which the outgrowth of new

capillaries is regulated by VEGF/VEGFR signaling. Angiogenesis is influenced by events that

induce survival or apoptosis in ECs; it can only be maintained by EC survival or inhibition of

apoptosis mediated; for example, by growth factors such as VEGF/VEGFR and PI3K/Akt

(Chavakis and Dimmeler 2002). Expression of the Ig-family member CD31 by ECs prevents

cell death and renders them immune privileged (Cheung et al. 2015). When the TNF-α

pathway is activated in ECs, CD31 is also activated; this in turn activates the Erk/Akt

pathway and subsequent exclusion of FoxO3 from the nucleus, leading ultimately to

inhibition of apoptosis *in vitro*. FoxO factors play a central role in controlling cell fate (i.e.,

apoptosis or proliferation). Constitutively active FOXO3a promotes EC apoptosis by

downregulating protective factors, whereas dominant-negative FoxO protects ECs from

apoptosis (Skurk et al. 2004). In line with this observation, loss of *mFoxO1* from murine ECs

leads to uncontrolled overgrowth and hyperplasia. Moreover, FoxO controls endothelial

quiescence by reducing glycolysis and mitochondrial respiration (Wilhelm et al. 2016). Active

FoxO inhibits EC migration, whereas its silencing has the opposite effect (Potente et al.

2005). By coincidence, we discovered a similar role for foxo in *Drosophila* larval TTCs. We

showed that foxo controls the branching of TTCs. Overexpression of *foxo* led to reduced

branching, whereas *foxo* depletion increased TTC branching. We hypothesize that TTCs need

to avoid activation of the Imd pathway because they depend on foxo to maintain full

functionality in response to changing environmental conditions. The exact role of foxo in

TTCs, and the signaling pathways involved, remain to be elucidated. Our study highlights the

importance of *Drosophila* TTCs as a model for human ECs and thus may provide information

that is useful for angiogenesis-related research.

The data presented herein demonstrate how immune privilege is necessary to maintain the

functionality of specific cell types. In the case of TTCs, which should mount potent immune

responses, the architecture and interconnectivity of the Imd (TNF-α) and JNK pathways mean

that cell fate is inevitably linked to foxo. This is, however, incompatible with the role of foxo

in controlling the structure and functionality of these cells under changing conditions. For

these reasons, switching off the Imd pathway is the only solution to this dilemma.

MATERIAL AND METHODS

Fly lines and husbandry

Flies were raised on corn meal medium at 25°C. Tracheal humoral immune responses were

tracked using *Drs-GFP* (BDSC_55707), which allows measurement of the GFP-tagged

promotor activity of the antimicrobial peptide gene *Drosomycin*. Similar to this, we employed

GFP reporters for Defensin, Attacin, Metchnikowin and Diptericin (Tzou et al. 2000). The

expression pattern of PGRP-LCx was analyzed by crossing *PGRP-LCx-Gal4* (BDSC_77776)

flies with *UAS-GFP* (BDSC_52262) flies. Tracheal driver lines *ppk4-Gal4* (Liu et al. 2003)

and *btl-Gal4; UAS-GFP* (*tub-Gal80ts*) (Leptin Group, EMBL Heidelberg), or TTC driver line

UAS-GFP; DSRF-Gal4 (Gervais and Casanova 2011)), were crossed to the corresponding

responder lines *UAS-PGRP-LCx(III)* (Kathryn Anderson, New York), *UAS-PGRP-LC(II)*

(BDSC_30918), *UAS-PGRP-LE* (*Dipt.-lacZ,UAS-Flag-PGRP-LE/CyO*, Shoichiro Kurata,

Sendai), *UAS-Rel68* (BDSC_55777), *UAS-Tak1* (BDSC_58810), *UAS-hid;rpr* (*UAS-hid; rpr;*

Christian Wegener, Würzburg), *UAS-bsk^{OE}* (BDSC_9310), TRE-RFP attP40 (Chatterjee &

Bohmann 2012), *UAS-bsk^{DN}* (BDSC_44801), *UAS-kay^{DN}* (BDSC_7214), *UAS-Ets21C^{RNAi}*

(BDSC_39069), *UAS-foxo^{RNAi}* (BDSC_27656), *UAS-kay^{OE}* (BDSC_7213), and *UAS-Jra^{OE}*

(BDSC_7216), or *w^{1118}* (for control crossing, BDSC_5905).

Infection experiments

Natural infection of larvae was performed as described (Tzou et al. 2000; Akhouayri et al.

2011). The gram-negative bacterium *Pectobacterium carotovorum* (*Erwinia carotovora*, Ecc-

15, 2141) was cultured overnight at 30°C in LB broth. The bacteria were pelleted by

centrifugation for 20 min at 3200 × g, resuspended in PBS, and absorbance at OD_{600}

measured. Next, 200 µl of the bacteria solution (OD_{600} = 160) was dropped into a vial

containing developing 2nd or early 3rd instar larvae (3 days after egg laying) in standard

cornmeal medium. After 24 h, the 3rd instar larvae were used for microscopy. The larvae were

heat killed in a drop of glycerol at 70°C for 20–25 s and arranged on a microscopy slide with

the dorsal side facing up. GFP expression in the tracheal DB and TTC between the dorsal

trunks was analyzed.

Branching analysis

The 3rd instar larvae were washed in PBS and deposited on a slide in a drop of glycerol. The

larvae were heat killed at 70°C for 20–25 s and arranged dorsal side up before microscopy.

Images of the TTCs were taken using the GFP channel (at 10× or 20× magnification). The

right TTC from the 3rd dorsal segment was chosen for analysis. Images were taken by an Axio

Imager.Z1 (Zeiss, Munich, Germany) with ApoTome in Z-stack mode to capture all TTC

branches. Measurement of the number and length of TTC branches was undertaken using the

ImageJ plugin NeuronJ (Meijering et al. 2004). The branches were measured, as described

previously (Jones and Metzstein 2011).

Immunohistochemistry and microscopy

For microscopic and immunohistochemical analyses of intestinal TTCs, the gut of 3rd instar

larvae (along with the connected TTCs) was dissected. As an alternative, a fillet dissection for

the cuticle-attached TTCs was used for microscopy. The tissue was dissected in PBS, fixed

with 4% PFA, and blocked with 5% normal goat serum (Sigma-Aldrich, Munich, Germany)

before overnight incubation at 4°C with an anti-GFP antibody (1:200, DSHB, Iowa City,

USA). Cells were also stained with anti-Dcp-1 (1:200, Cell Signaling Technology, Danvers,

USA), anti-pJNK (Promega, V7931), and anti-Relish-C (DSHB, 21F3). Images were taken

with an Axio Imager.Z1 (Zeiss, Munich, Germany) with ApoTome in the Z-stack mode.

Hypoxia sensitivity assay

The 3rd instar larvae were washed with PBS and placed into a new food vial with a scratched

surface; 20 larvae were used per replicate. The vials were incubated for at least 10 min until

all larvae were buried in the food and then deposited in a sealed desiccator. Nitrogen gas was

introduced until the O_2 concentration reached 2–5%. This O_2 level was maintained for 25 min.

The number of larvae outside of the food was counted every 5 min.

Measurement of epithelial thickness of the dorsal trunks

For temporal control using the TARGET system, crosses (with *btl-Gal4; UAS-GFP tub-*

Gal80ts) were maintained at 18 °C (restrictive temperature) until larvae reached third instar

stage. Larvae were then shifted to 29 °C (permissive temperature) for 24 h to induce transgene

expression. Late L3 wandering larvae were washed in PBS to remove residual food and

dissected in PBS. Z-stack DIC and GFP images were acquired using an Axio Imager Z1

microscope equipped with ApoTome (Zeiss, Munich, Germany). Epithelial thickness was

quantified using the AxioVision SE64 Rel. 4.9 software (Zeiss, Munich, Germany).

Measurements were performed in the 9th tracheal metamere.

Acknowledgments

We would like to thank Britta Laubenstein and Christiane Sandberg for their excellent

technical assistance.

Funding

This work was supported by Christian-Albrechts University, Kiel, as part of the Leibniz

Campus EvoLung (TR, JB), by the International Max-Planck Research School for

Evolutionary Biology (RR), and by the Deutsche Forschungsgemeinschaft (DFG) (as part of

the CRC 1182 (project C2))(TR).

Author contributions

Conceptualization: JB, TR

Methodology: RR, JB

Investigation: JB, LZ, RR, LS, JH

Visualization: JB, LZ, RR, LS, JH

Funding acquisition: TR

Project administration: TR

Supervision: TR, JB

Writing – original draft: JB, TR

Writing – review & editing: TR, JB, RR

Competing interests

The authors declare no conflict of interest.

  Figures and figure legends:

Fig. 1: Terminal tracheal immune reaction to natural infection measured via AMP

reporter response. GFP reporter larvae were infected with *P. carotovorum* for 24 h,

and GFP fluorescence in the terminal structures of the tracheal system was monitored.

The Images were taken in the DIC and GFP channels. Scale, 50 µm. (**A-E**) Dorsal

tracheal structures of non-infected larvae. (**A'-E'**) Dorsal tracheal structures of infected

larvae. (**F**) Dorsal view of the tracheal system, showing the dorsal trunks branching

into the dorsal branch (DB), fusion cells (FC) and the dorsal TTCs. The quantification

explanation is shown in magnification emphasized with colors. Presence of

fluorescence in DB/FC counts as a GFP positive larva (light green, GFP+). Presence

of fluorescence in the TTCs in GFP+ animals is indicated by dark green color (GFP

TTCs). (**G**) Proportion of animals with tracheal fluorescence following the mentioned

quantification procedure. Scale bar, 50 µm. Drs = Drosomycin, Def = Defensin, Att =

Attacin, Mtk = Metchnikowin, Dpt = Diptericin.

Fig. 2: Tracheal terminal cells (TTCs) show very rare *Drs* expression upon natural infection. *Drs-GFP* larvae were infected with *P. carotovorum* for 24 h, and GFP fluorescence in the TTCs of the tracheal system was monitored. Images were taken in the DIC (A, C-G) and GFP channels (A', C'-F'). (**A**) Dorsal TTCs without fluorescence. (**B**) Visceral TTCs without fluorescence. (**C**) Percentage of larvae showing GFP fluorescence in the DB and TTCs. (**D–G**) TTCs with expression of *Drs-GFP*. White arrows indicate shortened TTC branches (G). The black arrows mark a

melanization site and the arrowhead marks a translucent branch without air filling (G).

Dashed lines represent the proximal end of the TTCs. Scale bar, 50 µm.

Fig. 3: TTCs do not express the Imd receptor PGRP-LCx. (A–D) The secreted Imd

pathway receptor *PGRP-LE* is expressed in the main parts of the tracheal system

(*ppk4>PGRP-LE (GFP-Drs)*). The arrows indicate TTCs not expressing GFP. The

dashed lines represent the proximal end of the TTC. **(A–C)** An activated immune

response in the larvae is visualized by expression of GFP-tagged *Drosomycin* (*Drs*).

(D, E) Detailed TTCs were observed in fillet preparations (D) and in the dissected

intestine (E) in both the DIC (D, E) and GFP channels (D', E'). **(F)** Example TTC of

infected PGRP-LE animals with fluorescence. (**G**) Percentage of PGRP-LE^{OE} animals

with tracheal GFP expression. (**H, I**) Expression of GFP under the control of a *PGRP-*

LCx promoter (*PGRP-LCx-Gal4 > UAS-GFP*) revealed a lack of promoter activity,

and expression of GFP, in TTCs on the cuticle (H, H') and intestine (I, I'), which is in

contrast to the rest of the tracheal system. (**J, K**) The TTCs in the tracheal system are

visualized by GFP expression in the cuticle (J) and intestine (K) of wild-type larvae

(*btl-Gal4; UAS-GFP*). (**L**) Percentage of *PGRP-LCx* animals with tracheal GFP

expression. Scale bar, 50 µm. Dashed lines represent the proximal end of the TTCs. dt

= dorsal trunk.

Fig. 4: Expression of PGRP-LCx by TTCs leads to size reduction and loss of functionality. (A, B) Dorsal TTCs in the control (A) and *DSRF*-driven overexpression of *PGRP-LCx* in TTCs (B). **(C, D)** Measurement and quantification of the number (C) and length (D) of branches (n=22–45). Data are presented as the mean ± SD. **(E)** The

hypoxia sensitivity assay was conducted with control and *PGRP-LCx*-expressing 3rd

instar larvae under hypoxic conditions (2–3 % O_2, n = 11–14). Data are presented as

the means ± SEM. Statistical significance was tested using Mann-Whitney-U test. *

$p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. (**F**) Culture vials containing

control, hid;rpr, and Tak1 larvae (*DSRF > hid;rpr/Tak1*) at 4 days post-oviposition.

(**G, H**) Transmission light microscopy of dissected intestines from 3rd instar larvae (F,

G) or 2nd instar larvae (H) with the connected TTCs of control (G) showing expression

of *hid;rpr* (H) or *PGRP-LCx* (I). (**J, K**) Dissected intestines from control larvae (J) or

larvae expressing *PGRP-LCx* in TTCs (K) were stained with an antibody specific for

cleaved *Drosophila* Dcp-1 (purple). (**I', J'**) and then counterstained to detect GFP

(green). (**I'', J''**) Merged channels. Scale bars, 50 μm.

Fig. 5: JNK signaling is associated with impaired TTC branching. (A) Schematic showing

how the PGRP-LC-activated Imd signaling pathway is subdivided into the NF-κB

(Relish) and JNK (hep = JNKK, bsk = JNK) pathways. **(B–D)** Relish (B, *Rel68*), as

well as constitutively active hep (C, *hep*^{CA}) and bsk (D, *bsk*^{OE}), were expressed in

TTCs (*DSRF >*). **(E, F)** Measurement and quantification of the number (E) and length

(F) of branches in control (*w*^{1118}) versus *PGRP-LCx*-expressing TTCs (n=22–45). Data

are presented as the mean ± SD. Statistical significance was evaluated using Mann-Whitney-U test, ** p < 0.01, **** p < 0.0001, ns = not significant. **(G, H)** Dissected intestines from control (*DSRF > w¹¹¹⁸*) and *PGRP-LCx*-expressing flies (*DSRF > PGRP-LCx*), in which the TTCs were stained to detect Relish (Rel, G) and pJNK (purple, H). TTCs were counterstained with GFP (green). Arrows mark the TTC nucleus. Merged channels are shown. Scale bars, 50 μm.

Fig. 6: The TTC phenotype induced by PGRP-LCx is dependent on the transcription

factors kay and foxo. (**A**) Schematic illustration of the JNK signaling pathway

downstream of Tak1, which includes Ets21C, kay, and Jra (AP1). (**B–E**) *DSRF*-driven

PGRP-LCx^{OE} in TTCs was combined with the dominant-negative form of *Tak1, bsk*

(B), and *kay* (C), or with RNAi targeting *foxo* (D) or *Ets21C* (E). (**F**, **G**) Measurement

and quantification of the number (**F**) and length (**G**) of branches (n=16–45). (**H–K**)

Measurement and quantification of the number (H) and length (I) of GFP expressing

branches in control (w^{1118}) and PGRP-LCx-expressing cells (n=7–45). TTCs

overexpressing *foxo* (J, *foxo^{OE}*), and *kay* and *Jra* (K, *kay^{OE}* + *Jra^{OE}*). (**L**, **M**). TRE-RFP

expression in control (L) and PGRP-LCx-expressing TTCs (M). (**N**) *Foxo* promoter

activity in TTCs (*foxo-Gal4 > UAS-GFP*). Data are expressed as the mean ± SD.

Statistical significance was evaluated using Mann-Whitney-U test, * p < 0.05, *** p <

0.001, **** p < 0.0001, ns = not significant. The color of the asterisk indicates the

corresponding comparison. Dashed lines represent the mean control value. Scale bar,

50 µm.

Fig. 7: Targeted reduction of *foxo* expression in TTCs leads to hyperbranching. (A, B)

Representative tracheal branching in control (A) and *DSRF*-driven *foxo*^{RNAi} (B) cells

under normoxic conditions. **(C, D)** Representative images showing tracheal branching

in control (C) and *foxO*^{RNAi} (D) cells under hypoxic conditions. **(E)** Quantification of

branches in control and *DSRF > foxO*^{RNAi} TTCs under normoxic (21%) and hypoxic

(5%) conditions. Scale bar, 50 μm. n=21, Data are presented as the mean ± SD.

Statistical significance was evaluated using Mann-Whitney-U t-test, * $p < 0.05$, **** p

< 0.0001, ns = not significant.

Fig. 8: Schematic summarizing the JNK-mediated immune/stress response in the

trachea and TTCs. (A) Tracheal infection leads to an immune response involving

expression of antimicrobial peptides such as *Drosomycin* (*Drs*, green). In most cases,

the immune response is restricted to the tracheal trunks and the TTCs are unaffected

(bold arrow). In rare cases, TTCs express *Drs*, resulting in an impaired phenotype

(dashed arrow). (**B**) Imd signaling in the main tracheal trunks is induced by

peptidoglycan recognition receptors (PGRP)-LC and -LE. Downstream, the signaling

branches into a Relish (Rel) and a JNK signaling pathway. Activation of the pathways

mediates airway remodeling (*13*). However, activation in TTCs is avoided by the

absence of PGRP-LC, even though all other JNK signaling pathway components are

present. The pathway can be activated by external stressors, resulting in AP-1-

mediated cell death. The transcription factor foxo, which is component not only of the

JNK signaling pathway but also of the insulin signaling pathway, plays a role in TTC

homeostasis and their ability to branch.

Supplementary figures:

Fig. S1: Drs expression in infected larvae. GFP reporter larvae were infected with *P.*

carotovorum for 24 h, and GFP fluorescence was monitored. Drs expression was

visible in different larval structures like hemocytes and fat body (A-D). Scale bar, 50

µm. Drs = Drosomycin, Def = Defensin, Mtk = Metchnikowin, Dpt = Diptericin.

Fig. S2: PGRP-LCx mediated epithelial thickening of the dorsal trunks. PGRP-LCx

expression in the tracheal system was induced for 24 h in 3rd instar larvae. Epithelial

thickness of control larvae (A, *btl-Gal4, UAS-GFP; tub-Gal80ts > w^{1118}*) were

compared to PGRP-LCxOE larvae with PGRP-LCx on the third chromosome (III, B,

btl-Gal4, UAS-GFP; tub-Gal80ts > UAS-PGRP-LCx) and on the second chromosome

(II, C). Epithelial thickness was quantified (D). Scale bar, 50 μm. Data are presented

as the mean ± SD. Statistical significance was evaluated using unpaired t-test, **** p

< 0.0001.

Fig. S3: Increase in apoptotic signalling in PGRP-LC overexpression. The influence of PGRP-LC overexpression on apoptotic cells was investigated using -Dcp1 staining. For the analysis, the 8th tracheal metamere of the controls (**A**; *ppk4-Gal4 > w^1118* (Control_D), *w^1118 > UAS-PGRP-LC_x* (Control_R)) and PGRP-LE^OE (**B**; *ppk4-Gal4 > UAS-PGRP-LC_x*) were used. (**C**) Quantification was performed by measuring the fluorescence intensity and calculating the CTCF (corrected total cell fluorescence) of Control_D, Control_R and PGRP-LE^OE. The statistical significance was evaluated using the Mann-Whitney test. ****= p < 0.0001, ***= p < 0.001, ns = not significant. Scale bar, 50 µm.

Fig. S4: Overexpression of PGRP-LE and infection increased hypoxia sensitivity. The
influence of PGRP-LE[OE] and bacterial infection on hypoxia sensitivity was analyzed
based on the escape response of 3[rd] instar larvae exposed to 2.5 % O□. Control: w^{1118}
> *UAS-PGRP-LE* (*Drs-GFP*), PGRP-LE overexpression: *Btl-Gal4;Gal80 > UAS-*
PGRP-LE (Drs-GFP). Six independent replicates, each with 10 larvae of the control
(grey), infected control (black), PGRP-LE[OE] (pink) and infected PGRP-LE[OE]
(burgundy) were used for quantification. Values are presented as mean ± SEM.
Statistical analysis was performed using two-way ANOVA. **= $p < 0.01$, *= $p < 0.1$.

References

Abid MR, Guo S, Minami T, Spokes KC, Ueki K, Skurk C, Walsh K, Aird WC. 2004. Vascular endothelial
growth factor activates PI3K/Akt/forkhead signaling in endothelial cells. *Arterioscler Thromb
Vasc Biol* **24**: 294–300.
Abid MR, Shih SC, Otu HH, Spokes KC, Okada Y, Curiel DT, Minami T, Aird WC. 2006. A novel class of
vascular endothelial growth factor-responsive genes that require forkhead activity for
expression. *J Biol Chem* **281**: 35544–35553.
Adachi-Yamada T, Fujimura-Kamada K, Nishida Y, Matsumoto K. 1999. Distortion of proximodistal
information causes JNK-dependent apoptosis in Drosophila wing. *Nature* **400**: 166–169.
Akhouayri I, Turc C, Royet J, Charroux B. 2011. Toll-8/Tollo negatively regulates antimicrobial
response in the Drosophila respiratory epithelium. *PLoS Pathog* **7**: e1002319.
Amersfoort J, Eelen G, Carmeliet P. 2022. Immunomodulation by endothelial cells - partnering up
with the immune system? *Nat Rev Immunol* **22**: 576–588.

Bach FH, Hancock WW, Ferran C. 1997. Protective genes expressed in endothelial cells: a regulatory
response to injury. *Immunol Today* **18**: 483–486.
Bossen J, Knop M, Niu X, Thiedmann M, Prange R, Tahanzadeh N, Franzenburg S, Bruchhaus I, Heine
H, Roeder T. 2025. Chronic airway inflammation in Drosophila lacking the A20-like protein
Trabid. *Front Immunol* **16**: 1564386.
Bossen J, Kuhle JP, Roeder T. 2023. The tracheal immune system of insects - A blueprint for
understanding epithelial immunity. *Insect Biochem Mol Biol* **157**: 103960.
Burguete AS, Francis D, Rosa J, Ghabrial A. 2019. The regulation of cell size and branch complexity in
the terminal cells of the Drosophila tracheal system. *Dev Biol* **451**: 79–85.
Centanin L, Dekanty A, Romero N, Irisarri M, Gorr TA, Wappner P. 2008. Cell autonomy of HIF effects
in Drosophila: tracheal cells sense hypoxia and induce terminal branch sprouting. *Dev Cell* **14**:
547–558.
Centanin L, Gorr TA, Wappner P. 2010. Tracheal remodelling in response to hypoxia. *J Insect Physiol*
56: 447–454.
Chatterjee N, Bohmann D. 2012. A versatile PhiC31 based reporter system for measuring AP-1 and
Nrf2 signaling in Drosophila and in tissue culture. *PLoS One* **7**: e34063.
Chavakis E, Dimmeler S. 2002. Regulation of endothelial cell survival and apoptosis during
angiogenesis. *Arterioscler Thromb Vasc Biol* **22**: 887–893.
Cheung K, Ma L, Wang G, Coe D, Ferro R, Falasca M, Buckley CD, Mauro C, Marelli-Berg FM. 2015.
CD31 signals confer immune privilege to the vascular endothelium. *Proc Natl Acad Sci U S A*
112: E5815–5824.
Daniel S, Arvelo MB, Patel VI, Longo CR, Shrikhande G, Shukri T, Mahiou J, Sun DW, Mottley C, Grey
ST et al. 2004. A20 protects endothelial cells from TNF-, Fas-, and NK-mediated cell death by
inhibiting caspase 8 activation. *Blood* **104**: 2376–2384.
Davis MM, Engstrom Y. 2012. Immune response in the barrier epithelia: lessons from the fruit fly
Drosophila melanogaster. *J Innate Immun* **4**: 273–283.
DeVorkin L, Go NE, Hou Y-CC, Moradian A, Morin GB, Gorski SM. 2014. The Drosophila effector
caspase Dcp-1 regulates mitochondrial dynamics and autophagic flux via SesB. *Journal of Cell*
Biology **205**: 477–492.
Faisal MN, Hoffmann J, El-Kholy S, Kallsen K, Wagner C, Bruchhaus I, Fink C, Roeder T. 2014.
Transcriptional regionalization of the fruit fly's airway epithelium. *PLoS One* **9**: e102534.
Fernando MD, Kounatidis I, Ligoxygakis P. 2014. Loss of Trabid, a new negative regulator of the
drosophila immune-deficiency pathway at the level of TAK1, reduces life span. *PLoS Genet*
10: e1004117.
Ferrandon D, Jung AC, Criqui M, Lemaitre B, Uttenweiler-Joseph S, Michaut L, Reichhart J, Hoffmann
JA. 1998. A drosomycin-GFP reporter transgene reveals a local immune response in
Drosophila that is not dependent on the Toll pathway. *EMBO J* **17**: 1217–1227.
Fink C, Hoffmann J, Knop M, Li Y, Isermann K, Roeder T. 2016. Intestinal FoxO signaling is required to
survive oral infection in Drosophila. *Mucosal Immunol* **9**: 927–936.
Forrester JV, Xu H, Lambe T, Cornall R. 2008. Immune privilege or privileged immunity? *Mucosal*
Immunol **1**: 372–381.
Galea I, Bechmann I, Perry VH. 2007. What is immune privilege (not)? *Trends Immunol* **28**: 12–18.
Gervais L, Casanova J. 2011. The Drosophila homologue of SRF acts as a boosting mechanism to
sustain FGF-induced terminal branching in the tracheal system. *Development* **138**: 1269–
1274.
Grether ME, Abrams JM, Agapite J, White K, Steller H. 1995. The head involution defective gene of
Drosophila melanogaster functions in programmed cell death. *Genes Dev* **9**: 1694–1708.
Hafezi F, Steinbach JP, Marti A, Munz K, Wang ZQ, Wagner EF, Aguzzi A, Reme CE. 1997. The absence
of c-fos prevents light-induced apoptotic cell death of photoreceptors in retinal degeneration
in vivo. *Nat Med* **3**: 346–349.
Hill MA, Kwon JH, Gerry B, Hardy WA, Walkowiak OA, Kavarana MN, Nadig SN, Rajab TK. 2021.
Immune Privilege of Heart Valves. *Front Immunol* **12**: 731361.
Horowitz A, Simons M. 2008. Branching morphogenesis. *Circ Res* **103**: 784–795.

Hou YCC, Hannigan AM, Gorski SM. 2009. An executioner caspase regulates autophagy. *Autophagy* **5**:
530–533.
Jarecki J, Johnson E, Krasnow MA. 1999. Oxygen regulation of airway branching in Drosophila is
mediated by branchless FGF. *Cell* **99**: 211–220.
Jones TA, Metzstein MM. 2011. A novel function for the PAR complex in subcellular morphogenesis
of tracheal terminal cells in Drosophila melanogaster. *Genetics* **189**: 153–164.
League GP, Hillyer JF. 2016. Functional integration of the circulatory, immune, and respiratory
systems in mosquito larvae: pathogen killing in the hemocyte-rich tracheal tufts. *BMC Biol*
14: 78.
Lehmann M, Jiang C, Ip YT, Thummel CS. 2002. AP-1, but not NF-kappa B, is required for efficient
steroid-triggered cell death in Drosophila. *Cell Death Differ* **9**: 581–590.
Linneweber GA, Jacobson J, Busch KE, Hudry B, Christov CP, Dormann D, Yuan M, Otani T, Knust E, de
Bono M et al. 2014. Neuronal control of metabolism through nutrient-dependent modulation
of tracheal branching. *Cell* **156**: 69–83.
Liu L, Johnson WA, Welsh MJ. 2003. Drosophila DEG/ENaC pickpocket genes are expressed in the
tracheal system, where they may be involved in liquid clearance. *Proc Natl Acad Sci U S A*
100: 2128–2133.
Luo X, Puig O, Hyun J, Bohmann D, Jasper H. 2007. Foxo and Fos regulate the decision between cell
death and survival in response to UV irradiation. *EMBO J* **26**: 380–390.
Mai J, Virtue A, Shen J, Wang H, Yang XF. 2013. An evolving new paradigm: endothelial cells--
conditional innate immune cells. *J Hematol Oncol* **6**: 61.
Meijering E, Jacob M, Sarria JC, Steiner P, Hirling H, Unser M. 2004. Design and validation of a tool for
neurite tracing and analysis in fluorescence microscopy images. *Cytometry A* **58**: 167–176.
Mundorf J, Donohoe CD, McClure CD, Southall TD, Uhlirova M. 2019. Ets21c Governs Tissue Renewal,
Stress Tolerance, and Aging in the Drosophila Intestine. *Cell Rep* **27**: 3019–3033 e3015.
Naito H, Iba T, Wakabayashi T, Tai-Nagara I, Suehiro JI, Jia W, Eino D, Sakimoto S, Muramatsu F,
Kidoya H et al. 2019. TAK1 Prevents Endothelial Apoptosis and Maintains Vascular Integrity.
Dev Cell **48**: 151–166 e157.
O'Connor DS, Schechner JS, Adida C, Mesri M, Rothermel AL, Li F, Nath AK, Pober JS, Altieri DC. 2000.
Control of apoptosis during angiogenesis by survivin expression in endothelial cells. *Am J*
Pathol **156**: 393–398.
Potente M, Urbich C, Sasaki K, Hofmann WK, Heeschen C, Aicher A, Kollipara R, DePinho RA, Zeiher
AM, Dimmeler S. 2005. Involvement of Foxo transcription factors in angiogenesis and
postnatal neovascularization. *J Clin Invest* **115**: 2382–2392.
Ruff WE, Greiling TM, Kriegel MA. 2020. Host-microbiota interactions in immune-mediated diseases.
Nat Rev Microbiol **18**: 521–538.
Silverman N, Zhou R, Erlich RL, Hunter M, Bernstein E, Schneider D, Maniatis T. 2003. Immune
activation of NF-kappaB and JNK requires Drosophila TAK1. *J Biol Chem* **278**: 48928–48934.
Skurk C, Maatz H, Kim HS, Yang J, Abid MR, Aird WC, Walsh K. 2004. The Akt-regulated forkhead
transcription factor FOXO3a controls endothelial cell viability through modulation of the
caspase-8 inhibitor FLIP. *J Biol Chem* **279**: 1513–1525.
Spyridopoulos I, Brogi E, Kearney M, Sullivan AB, Cetrulo C, Isner JM, Losordo DW. 1997. Vascular
endothelial growth factor inhibits endothelial cell apoptosis induced by tumor necrosis
factor-alpha: balance between growth and death signals. *J Mol Cell Cardiol* **29**: 1321–1330.
Takatsu Y, Nakamura M, Stapleton M, Danos MC, Matsumoto K, O'Connor MB, Shibuya H, Ueno N.
2000. TAK1 participates in c-Jun N-terminal kinase signaling during Drosophila development.
Mol Cell Biol **20**: 3015–3026.
Takehana A, Katsuyama T, Yano T, Oshima Y, Takada H, Aigaki T, Kurata S. 2002. Overexpression of a
pattern-recognition receptor, peptidoglycan-recognition protein-LE, activates imd/relish-
mediated antibacterial defense and the prophenoloxidase cascade in Drosophila larvae. *Proc*
Natl Acad Sci U S A **99**: 13705–13710.
Takehana A, Yano T, Mita S, Kotani A, Oshima Y, Kurata S. 2004. Peptidoglycan recognition protein
(PGRP)-LE and PGRP-LC act synergistically in Drosophila immunity. *EMBO J* **23**: 4690–4700.

Tzou P, Ohresser S, Ferrandon D, Capovilla M, Reichhart JM, Lemaitre B, Hoffmann JA, Imler JL. 2000.
Tissue-specific inducible expression of antimicrobial peptide genes in Drosophila surface
epithelia. *Immunity* **13**: 737–748.

Uv A, Cantera R, Samakovlis C. 2003. Drosophila tracheal morphogenesis: intricate cellular solutions
to basic plumbing problems. *Trends Cell Biol* **13**: 301–309.

Wagner C, Isermann K, Fehrenbach H, Roeder T. 2008. Molecular architecture of the fruit fly's airway
epithelial immune system. *BMC Genomics* **9**: 446.

Wagner C, Isermann K, Roeder T. 2009. Infection induces a survival program and local remodeling in
the airway epithelium of the fly. *FASEB J* **23**: 2045–2054.

Wagner C, Uliczka K, Bossen J, Niu X, Fink C, Thiedmann M, Knop M, Vock C, Abdelsadik A, Zissler UM
et al. 2021. Constitutive immune activity promotes JNK- and FoxO-dependent remodeling of
Drosophila airways. *Cell Rep* **35**: 108956.

White K, Grether ME, Abrams JM, Young L, Farrell K, Steller H. 1994. Genetic control of programmed
cell death in Drosophila. *Science* **264**: 677–683.

Whitten JM. 1957. The Post-embryonic Development of the Tracheal System in Drosophila
melanogaster. *Journal of Cell Science* **s3-98**: 123–150.

Wilhelm K, Happel K, Eelen G, Schoors S, Oellerich MF, Lim R, Zimmermann B, Aspalter IM, Franco
CA, Boettger T et al. 2016. FOXO1 couples metabolic activity and growth state in the vascular
endothelium. *Nature* **529**: 216–220.

Wong DM, Shen Z, Owyang KE, Martinez-Agosto JA. 2014. Insulin- and warts-dependent regulation of
tracheal plasticity modulates systemic larval growth during hypoxia in Drosophila
melanogaster. *PLoS One* **9**: e115297.

Wu D, Wang Z, Zhang J, Robinson AG, Lyu B, Chen Z, Wang C, Wei B, Xia X, Zhang Q et al. 2022.
Apoptotic caspase inhibits innate immune signaling by cleaving NF-κBs in both Mammals and
Flies. *Cell Death & Disease* **13**: 731.

Yang J, Yan J, Liu B. 2018. Targeting VEGF/VEGFR to Modulate Antitumor Immunity. *Front Immunol* **9**:
978.

Yang L, Joseph S, Sun T, Hoffmann J, Thevissen S, Offermanns S, Strilic B. 2019. TAK1 regulates
endothelial cell necroptosis and tumor metastasis. *Cell Death Differ* **26**: 1987–1997.

825