Patient-derived colon epithelial organoids reveal lipid-related metabolic dysfunction in

pediatric ulcerative colitis

Babajide A. Ojo[1,2], Lyong Heo[3], Sejal R. Fox[4], Amanda Waddell[5,6], Maria E. Moreno-Fernandez[5,7], Marielle

Gibson[1,2], Tracy Tran[1,2], Ashley L. Dunn[1,2], Essam I.A Elknawy[1,2], Neetu Saini[8], Javier A. López-Rivera[1,2],

Senad Divanovic[5,6,9], Vinicio A. de Jesus Perez[10], Michael J. Rosen[1,2,#]

[1] Division of Pediatric Gastroenterology, Hepatology, and Nutrition, Stanford University School of Medicine,

Stanford, CA, USA

[2] Center for IBD and Celiac Disease, Stanford Medicine Children's Health, Palo Alto, CA, USA

[3] Stanford Center for Genomics and Personalized Medicine, Stanford University, Palo Alto, CA, USA

[4] Applied Gene and Cell Therapy Center, Cincinnati Children's Hospital Medical Center, Cincinnati, OH,

USA

[5] Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH, USA

[6] Division of Immunobiology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

[7] Division of Gastroenterology, Hepatology, and Nutrition, Cincinnati Children's Hospital Medical Center,

Cincinnati, OH, USA

[8] Division of Pediatric Hematology, Oncology, Stem Cell Transplantation and Regenerative Medicine,

Stanford University School of Medicine, Palo Alto, CA, USA

[9] Center for Inflammation and Tolerance, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

[10] Division of Pulmonary and Critical Care Medicine, Department of Medicine, Stanford University, Stanford,

CA, USA

[#] Address correspondence to Michael J. Rosen, MD, MSCI, Division of Gastroenterology, Hepatology, and

Nutrition, Department of Pediatrics, Stanford University, Stanford, CA (rosenm@stanford.edu)

Competing interests: The authors have no competing interests to disclose.

Abstract

Ulcerative colitis (UC) is associated with epithelial metabolic derangements which exacerbate gut inflammation. Patient-derived organoids recapitulate complexities of the parent tissue in health and disease; however, whether colon organoids (colonoids) model the metabolic impairments in the pediatric UC epithelium is unclear. Here, we developed colonoid lines from pediatric patients with endoscopically active UC, inactive UC, and those without endoscopic or histologic evidence of colon inflammation (non-IBD controls) to interrogate functional metabolic differences in the colon epithelia. We demonstrate that colonoids from active UC patients exhibit hypermetabolic features and cellular stress, specifically during differentiation. Hypermetabolism in differentiating active UC colonoids was driven, in part, by increased proton leak, and supported by enhanced glycolytic capacity and dysregulated neutral lipid accumulation. Transcriptomic and pathway analyses indicated a role for PPAR-α in lipid-induced hypermetabolism in aUC colonoids, which was validated by PPAR-α activation in non-IBD colonoids. Accordingly, limiting neutral lipid accumulation in active UC colonoids through pharmacological inhibition of PPAR-α induced a metabolic shift towards glucose utilization, suppressed hypermetabolism and chemokine secretion, and improved markers of cellular stress and epithelial differentiation. Taken together, we reveal a role for lipid-related metabolic dysfunction in the pediatric UC epithelium and support the advancement of colonoids as a preclinical human model for testing epithelial-directed therapies against such metabolic dysfunction.

Main

The restoration of normal intestinal epithelial function is critical for colon mucosa healing in inflammatory bowel diseases (IBD)[1,2]; however, current therapies predominantly target the immune system. As such, only 20-30% of patients achieve mucosal healing with today's therapies[3–6]. Ulcerative colitis (UC) is an IBD subtype characterized by the continuous chronic inflammation of the colon mucosa, extending proximally from the rectum[7]. Pediatric UC is marked by more severe and extensive disease than adult UC.[7–11]. Achieving mucosa healing is one of the best predictors of sustained disease remission.[11–13] Studies that enhance our understanding of the human intestinal epithelial function in health and disease may advance epithelial-directed therapies in pediatric UC.

Systemic and local dysregulation of metabolism is a hallmark of inflammation in human IBD[14,15]. Direct investigation of the inflamed human colon tissue in UC has proposed several metabolic impairments in UC, including impaired mitochondrial respiratory chain complex activity, enhanced epithelial mitochondria fission, deficient butyrate metabolism, and reduced epithelial mitochondrial membrane potential[16–19]. This may be critical to the course of disease since mitochondrial activity regulates several aspects of epithelial cellular homeostasis, including the maintenance of energetic balance, adequate reactive oxygen species (ROS) production, and regulating pro-inflammatory responses[20]. However, the cellular mechanisms underlying UC epithelial metabolic dysfunction are not known. Such a deeper understanding may reveal novel epithelial targets to promote mucosal healing.

The development of epithelial-directed therapies in UC is hindered by a lack of human preclinical models of the diseased epithelium. Over the last decade, advances in organoid biology have grown their promise as human pre-clinical models of disease[21]. In particular, patient-derived adult stem cell organoids are capable of self-organizing into structural and functional models that may reflect the cell and molecular complexities of parent tissues[22]. In UC, patient-derived colon organoids (colonoids) may reflect disease-specific disparities compared to healthy colonoids including alterations in transcription, epigenetics, and somatic

gene mutations[23–27]. Despite the promise of intestinal organoids in understanding human intestinal

diseases, they have not been used to interrogate the metabolic nature of disease in pediatric UC.

In this study, we used mucosal biopsy-derived colonoids to understand the metabolic features of colonoids

from non-IBD, active, and inactive pediatric UC patients before and during differentiation. We report an

inefficient bioenergetic response to differentiation in colonoids from active UC patients characterized by

enhanced oxygen consumption and proton leak. UC epithelial metabolic dysfunction was associated with

impaired differentiation, and increased cellular stress and chemokine secretion. Differentiating active UC

colonoids exhibited increased neutral lipid accumulation and transcriptomic evidence of heightened lipid

metabolism. We identified a PPAR-α causal network explaining differential gene expression in active UC

colonoids. Accordingly, inhibition of PPAR-α reduced aberrant neutral lipid accumulation, hypermetabolism,

cellular stress, and chemokine secretion and improved markers of differentiation in active UC colonoids.

Results

Derivation of pediatric colon organoids

We obtained rectal biopsies from pediatric patients undergoing clinically-indicated colonoscopies (see

Supplementary Table 1 for patient demographics). We dissociated fresh rectal biopsies and removed

epithelial cells under a dissecting microscope before culturing and expansion in organoid growth medium

(OGM, Extended Data Fig. 1a). We observed enhanced recovery of non-differentiated spheroids from

biopsies obtained from non-IBD controls compared to those from active UC (aUC). Control spheroids were

sufficiently mature for passaging and expansion by 7 days (Extended Data Fig. 1b), whereas aUC

spheroids required 10-14 days. After the extended growth period, fully established aUC spheroids were

morphologically comparable to controls and maintained similarity after multiple passages and a freeze-thaw

cycle (Extended Data Fig. 1c,d). The timeline and morphology of spheroids generated from inactive UC

(iUC) biopsies (patients with endoscopic healing from active disease) were similar to that of biopsies from

non-IBD patients (data not shown). To compare the regenerative capacity of spheroids post

cryopreservation from each patient group, we digested eight control and aUC spheroid lines into single cell

suspensions and measured spheroid diameter after 6 days in OGM. We found no differences in the

spheroid diameter between control and aUC spheroids (Extended Data Fig. 2a,b). Overall, we had a 93%,

82%, and 89% success rate in generating stable spheroids from colon biopsies obtained from control, aUC

and iUC patients, respectively.

Colonoids from active UC patients exhibit impaired differentiation and hypermetabolism

Due to the indifference in spheroid morphology, we wondered whether control and aUC spheroids had

analogous metabolic profiles. Hence, we subjected eight individual spheroid lines each from control and

aUC patients to extracellular flux analysis to measure the oxygen consumption rate (OCR) at baseline and

in response to mitochondrial inhibitors (Extended Data Fig. 2c,d). We found no differences in basal OCR

and OCR due to maximal respiration, proton leak, ATP-linked respiration, and non-mitochondrial respiration

(Extended Data Fig. 2d). Analysis of mitochondrial mass (mtMass) by fluorescence showed that the

undifferentiated control and aUC spheroids had similar mtMass (Extended Data Fig. 2e). Similarly, we

analyzed extracellular LDH release into the medium – a marker cellular cytotoxicity – and found no

differences in LDH release between control and aUC spheroids (Extended Data Fig. 2f). These results

suggest that control and aUC spheroids are morphologically and metabolically similar in their

undifferentiated states.

We then sought to examine the morphology and metabolic profiles of control and aUC spheroids during

differentiation. After 3 days in the differentiation condition, we observed morphologic differences between

aUC and control differentiating colon organoids (hereafter referred to as colonoids) as more aUC colonoids

remained in their cystic morphology compared to the largely budded structures of control colonoids (Fig.

1a,b). Further differentiation up to 7 days resulted in visibly reduced survival in aUC colonoids compared to

controls (Extended Data Fig. 2g). Apart from morphological features, complementary means to assess

differentiation of adult stem cell-derived organoids include their multipotency into mature cell types[28]. Thus,

we determined the gene expression of the secretory lineage progenitor, *ATOH1,* (ref: Yang et al, 2001), the

enteroendocrine cell marker, *CHGA* (ref), and the crypt-based goblet cell marker of the colon, *WFDC2*[29,30],

in C and aUC before and after 3-day differentiation (i.e spheroids vs colonoids). None of the examined

markers were significantly different in the control and aUC spheroids (Extended Data Fig 2h,i; Fig. 1c). As

expected, *ATOH1* and *CHGA* were upregulated in both control and aUC colonoids compared to their

respective undifferentiated spheroids (Extended Data Fig. 2h,i). In aUC colonoids, there was a failure to

upregulate WFDC2, consistent with the phenomenon of goblet cell depletion in UC (Fig. 1c)[29]. Furthermore,

there was also a reduction in upregulation of CHGA with differentiation as compared to control colonoids

(Extended Data Fig. 2i).

Following the evidence of impaired differentiation in aUC colonoids, we examined whether the metabolic

profiles of control and aUC colonoids diverged during the early stages of differentiation. We analyzed OCR

in eight patient-derived colonoids per group after 3-day differentiation (Fig. 1d,e). We found a 22% elevation

of basal OCR in aUC colonoids compared to the controls. In addition, we found a 75% significant elevation

in median OCR due to proton leak in aUC colonoids compared to control, with a non-significant increase in

non-mitochondrial and ATP-linked OCR compared to control colonoids (Fig. 1f). The observation that proton

leak explained much of the higher oxygen consumption in aUC colonoids suggested mitochondrial

inefficiency in aUC. Hence, we measured total ATP and found no significant difference in total cellular ATP

between aUC and control colonoids (Fig. 1g). Colon epithelial cells from treatment-naïve UC patients exhibit

lower mitochondrial membrane potential (MMP)[19], and we observed a similar reduction in MMP in aUC

colonoids compared to controls (Fig, 1h). In line with the hypermetabolic patterns in aUC colonoids, we

observed a significant elevation of mtMass during differentiation compared to controls (Fig. 1i).

Correspondingly, we found significant increases in mitochondrial ROS levels and LDH release in aUC

colonoids compared to control, suggesting that hypermetabolism in aUC colonoids is accompanied by

enhanced oxidative stress and cytotoxicity (Fig. 1j,k). Taken together, these data suggest a metabolic

inefficiency in aUC colonoids at the early stages of differentiation which may contribute to oxidative stress

and cytotoxicity.

Colonoids from inactive UC patients recall features of hypermetabolism upon inflammatory exposure

Following the dysregulated metabolic profile observed in aUC colonoids, we asked whether colonoids from

inactive UC (iUC) patients, those on treatment with no endoscopic disease activity, were metabolically

distinct from control colonoids. First, we observed no morphological differences between control and iUC

colonoids during differentiation (Extended Data Fig. 3 a,b). The bioenergetic profiles of control and iUC were

similar after a 3-day differentiation as we observed no differences in basal OCR, proton leak, maximal

respiration, and ATP-linked OCR; however, non-mitochondrial OCR was higher in iUC compared to control

colonoids (Extended Data Fig. 3c,d). Accordingly, and similar to our observation in aUC colonoids, we

observed enhanced ROS in iUC colonoids compared to control colonoids, suggesting that ROS generation

in our colonoids is not solely coupled to hypermetabolism.

Since biopsies for aUC colonoids were obtained from inflamed epithelial regions, we wondered whether

prior exposure to inflammation *in vivo* may explain the hypermetabolic nature of aUC colonoids *in vitro*. To

model this, we reasoned that the pre-exposure of spheroids to inflammation may recapitulate

hypermetabolic features in control and iUC colonoids during differentiation. Thus, before differentiation, we

exposed control and iUC spheroids to a modified inflammatory cocktail (40 ng/mL TNFa, 20 ng/mL IL1b,

and 500 ng/mL flagellin) previously suggested to re-induce the inflammatory nature of UC *ex vivo*[25]

(Extended Data Fig. 3f). We observed that unlike control colonoids, iUC spheroids showed reduced budded

morphology after a 3-day differentiation when pre-exposed to the inflammatory cocktail. (Extended Data Fig.

3g). We observed pre-exposure with the inflammatory cocktail led to enhanced mtMass in iUC but not

control colonoids (Extended Data Fig. 3h). Correspondingly, we found a significant increase in basal OCR

in iUC colonoids pre-exposed to the inflammatory cocktail compared to vehicle and exposed control.

However, there were no significant effects on maximal OCR and proton leak (Extended Data Fig. 3i,j).

Collectively, these data suggest that iUC spheroids are more susceptible to inflammatory challenge and this

leads to a partial recapitulation of aUC-related enhanced respiratory phenotypes during differentiation.

Metabolic uncoupling contributes to hypermetabolism in aUC colonoids

The proton leak and resulting metabolic inefficiency observed in aUC colonoids during differentiation may

be indicative of enhanced uncoupling of mitochondrial respiration[31]. Thus, we hypothesized that

endogenous mitochondrial uncoupling proteins would be upregulated in aUC compared to control. First, we

analyzed public bulk RNA-seq datasets from the RISK study (GSE117993)[32] and PROTECT Pediatric IBD

inception cohort studies (GSE109142)[19] and found that RNA for uncoupling protein 2 (*UCP2*) is the

dominant uncoupling protein RNA expressed in non-inflamed rectal tissue from non-IBD control patients

(Extended Data Fig. 4a). Rectal gene expression data from 56 and 206 UC patients in the RISK and Protect

Studies, respectively, showed that *UCP2* is significantly upregulated in the rectal mucosa of pediatric UC

patients compared to controls (Extended Data Fig 4b,c). In our patient-derived colonoid samples, we

confirmed enhanced UCP2 expression at the mRNA and protein levels (Extended Data Fig 4d).

Immunofluorescence imaging showed the coexpression of UCP2 protein with the mitochondrial marker

COX4, suggesting enhanced presence of UCP2 in the mitochondria during differentiation (Extended Data

Fig 4e).

To test whether metabolic uncoupling was sufficient to induce the altered colon epithelial differentiation and

metabolism observed aUC, we added a low-dose Fccp (protonophore that uncouples mitochondrial

OXPHOS), into the medium throughout a 3-day differentiation of control colonoids (Extended Data Fig. 4f).

Paired analysis showed a significant elevation of basal OCR and OCR due to maximal respiration, proton

leak, non-mitochondrial respiration, but also significantly increased ATP-linked respiration, suggesting that

the healthy colonoids may overcome chemical uncoupling to maintain ATP levels (Extended Data Fig. 4g,h).

Still, we observed elevated mtMass and mitochondrial ROS levels in Fccp-treated colonoids compared to

controls suggesting an increase in mitochondrial and oxidative stress during differentiation (Extended Data

Fig. 4i,j). In contrast, we did not see an increase in LDH release to suggest uncoupling alone induced

cytotoxicity (Extended Data Fig. 4k). Finally, we observed that Fccp uncoupling during differentiation did not

impact colonoid morphology or *WFDC2* gene expression (Extended Data Fig 4l-n). These data suggest that

upregulation of UCP2 and mitochondrial uncoupling likely contribute to hypermetabolism and oxidative

stress in aUC colonoids, but may be insufficient to impair differentiation or induce cytotoxicity.

Active UC colonoids exhibit an altered transcriptional profile and upregulated chemokine secretion

To understand transcriptional changes that may underly altered metabolism and function in aUC colonoids,

we conducted bulk RNA-seq on spheroids and colonoids differentiated for 3 days (n= 8 patient lines per

diagnosis, passage 5-7). Among 13,884 genes detected in spheroids after quality control, only seven were

significantly upregulated (*CYP3A4, KRT23, FLNC, APOL1, DAPP1, TNFSF15, GRHL3*) and four were

significantly downregulated (*EVC, NEFH, CDKN1C, SORBS1*) in aUC compared to control undifferentiated

spheroids (Log$_2$FC ≥ ±1, FDR P ≤ 0.1, Extended Data Fig 5a, Supplementary Table 2). These findings

suggest that in conditions promoting stemness, C and aUC spheroids are largely transcriptionally similar in

agreement with their metabolic similarity.

We found that upon differentiation, control and aUC colonoids had divergent transcriptomic profiles (Fig.

2a), in line with their distinct metabolic profiles. Specifically, 112 genes were significantly upregulated and 34

significantly downregulated (Log$_2$FC ≥ ±1, FDR P ≤ 0.1; Supplementary Table 2). The top upregulated

genes included well-known MHCII-related genes in active UC[19,33,34], *HLA-DRA, HLA-DMB, HLA-DRB, HLA-*

DMA (Fig. 2b, Supplementary Table 2); lipid-related genes (*FABP6, ABCA12, and APOL3*); and *CARD6* a

member of the caspase recruitment domain family (ref) (Fig. 2b, Supplementary Table 2). Top

downregulated genes included those involved in the development of cytoskeletal filaments (*NEFH, KRT12*),

goblet cells (*ZG16, WFDC2*), and those involved in cellular development and histone maintenance (*GATA2*

and *HIST1H4K*, respectively) (Fig. 2b, Supplementary Table 2). Consistent with the metabolic profile of

inactive UC colonoids, only two genes (*HLA-DRA* and *HLA-DMB*) were significantly upregulated in iUC

colonoids compared to control colonoids (Supplementary Table 2).

Chemokines recruit immune cells to the epithelium in UC, where they can incite damage. We observed

enhanced expression of chemokine genes involved in the recruitment of macrophages (CCL2), T-cells

(CXCL11), neutrophils, (CXCL1), and T- and B-cells (CCL28) in aUC colonoids compared to controls (Fig.

2c). Using medium collected during a 2-d differentiation, We confirmed by ELISA of colonoid media after 2

days of differentiation that aUC colonoids exhibit increased secretion of these chemokine proteins

compared to controls (Fig 2d-f). Due to the contribution of metabolic uncoupling to the hypermetabolism in

aUC colonoids (Extended Data Fig. 4), we asked whether metabolic uncoupling was sufficient to induce

chemokine secretion in control colonoids during differentiation Indeed, treatment of control colonoids with

Fccp during differentiation significantly upregulated the gene and protein secretion of CXCL1, CXCL11, and

CCL2 (Extended Data Fig. 4o,p). Thus, molecular events that stimulate metabolic uncoupling during

epithelial differentiation in UC may contribute to the chemoattraction of immune cells.

Active UC colonoids exhibit enhanced glycolytic capacity

To begin to understand the cellular underpinnings of hypermetabolism in aUC colonoids, we analyzed our

differentially expressed transcriptomic data using functional annotation analysis (Ingenuity Pathway Analysis

[IPA])[35] to predict the dominant molecular and cellular functional differences between aUC and control

colonoids. IPA implicated carbohydrate metabolism as the top mediator of cellular function during

differentiation (Fig. 3a). Other dominant functions include molecules involved in cell death and survival,

cellular assembly and organization, cellular development, cellular growth and proliferation, and lipid

metabolism (see details of all cellular and molecular functions in Supplementary Table 3).

Given that carbohydrate metabolism is the top cellular function in our data, we tested whether aUC

colonoids exhibited enhanced glycolysis during differentiation. In aerobic glycolysis, pyruvate is produced,

which enters the citric acid cycle. In anaerobic glycolysis, pyruvate is converted to lactate. We assessed

extracellular acidification rates (ECAR) that result predominantly from lactate secreted as an end product of

anaerobic glycolysis[36] by extracellular flux analysis in control and aUC colonoids beginning in a glucose-

depleted medium. After glucose injection, we did not observe any differences in ECAR due to anaerobic

glycolysis (Fig. 3b,c). However, we observed that aUC colonoids had an enhanced glycolytic capacity and

reserve compared to control colonoids after the suppression of OXPHOS by inhibiting mitochondrial

complex V (ATP Synthase) with oligomycin[37] followed by inhibition of glycolysis with 2-deoxy glucose (Fig.

3b,d-e). Still, non-glycolytic acidification in aUC was significantly higher compared to controls, which

suggests other pathways such as CO_2 from the TCA cycle contribute to extracellular acidification[36].

To complement the results of real-time extracellular flux analysis, we used medium collected over the first 2

days of differentiation to measure cumulative extracellular glucose consumption and lactate production

during the early process of differentiation. On average, aUC colonoids consumed 4.2 times more glucose

from the media and produced 2.8 times more lactate compared to controls (Fig. 3g,h). Collectively, the data

indicate that, during the early stages of differentiation, aUC colonoids consume more glucose and exhibit

increased capacity for glycolysis which could support the observed hypermetabolism.

Dysregulated lipid metabolic genes in aUC colonoids

Functional annotation analysis of aUC colonoid differential gene expression also identified lipid metabolism

as one of the top 10 dominant molecular and cellular functions (Fig. 3a). Upregulated lipid metabolism

genes in aUC included those related to fatty acid transport (*FABP6, ABCA12*), apolipoproteins (*APOL1,*

APOL3), sphingolipids and phospholipid biosynthesis (*SPTLC3* and *DGKG*, respectively), lipid-metabolizing

CYPs (*CYPC218*, *CYP3A4*), and lipolysis (*LPL, PLB1, AADAC, LIPA*). The only downregulated lipid-related

gene was *ALOX15B*, which plays a role in the generation of specialized pro-resolving lipid mediators in

inflammation (Supplementary Table 3, Fig. 4a) [38].

To confirm that increased expression of these lipid-related genes in aUC colonoids is relevant to the colon

epithelium in patients, we examined their expression in other bulk and single-cell RNA-seq UC data sets.

First, the PROTECT bulk RNA-seq data from 206 pediatric patients also showed enhanced expression of

ABCA12, APOL3, APOL1, LPL, DGKG, and *LIPA* in active UC patients compared to controls (Extended

Data Fig. 5a). Next, we analyzed a publically available UC single-cell RNA-seq data[39] to evaluate whether

inflamed epithelial cells exhibit similar upregulation of lipid-related genes in UC. Compared to healthy

samples, a higher proportion of the inflamed samples visibly expressed these identified lipid-related genes

including stem cells and other progenitor epithelial lineages (Extended Data Fig. 5b,c). Taken together,

these analyses of the complementary organoid, bulk tissue, and single-cell RNA-seq data sets indicate the

transcriptomic dysregulation of lipid-related genes in UC is modeled in our colonoids.

Aberrant lipid accumulation and oxidation in aUC colonoids during differentiation

Considering the altered expression of lipid metabolic genes in aUC, we sought to determine whether

enhanced lipid metabolism contributes to the observed hypermetabolic phenotype. First, we measured the

accumulation of BODIPY[+] neutral lipids in our colonoids before and during differentiation. Consistent with

their analogous metabolic states before differentiation, control and aUC spheroids had similar neutral lipid

content (Fig. 4b). However, we observed a significant increase in neutral lipid accumulation during

differentiation in aUC colonoids compared to their paired spheroids, while control spheroids and colonoids

still retained similar neutral lipid content. (Fig. 4b). Furthermore, differentiated aUC colonoids contained

elevated lipid content compared to the control colonoids (Fig. 4b,c). Extracellular and intracellular

triglycerides, cholesterol esters, and membrane phospholipids are hydrolyzed to free fatty acids, which

undergo fatty acid oxidation (FAO) and support OXPHOS. Our bulk RNA-seq data showed an upregulation

of 4 lipid hydrolysis genes (*LPL, PLB1, AADAC, LIPA*) in aUC colonoids (Fig. 4a), which is largely similar to

the analyzed single-cell RNA-seq data[39] (Extended Data Fig. 5c). Specifically, we observed that lipoprotein

lipase (*LPL)* was predominantly overexpressed in differentiating cycling transit amplifying (TA) cells in

inflamed samples (Extended Data Fig. 5c). Since TA cells are progenitors that are involved in cell cycle,

mitochondrial biogenesis and terminal differentiation into mature epithelial cell types[30,40], we reasoned that

for our colonoids to recapitulate *in vivo* lipid metabolic phenotypes, LPL activity would dissipate over time

during differentiation. Indeed, a 5-day differentiation protocol showed elevated LPL activity in aUC colonoids

compared to control on Day 2, however, by Day 5 of differentiation, LPL activity in both control and aUC

dropped to pointedly low levels with no significant difference between groups (Fig. 4d). These data further

suggest that active lipid cycling may be overamplified during differentiation in the UC colon epithelium and

our pediatric colonoid model of UC may recapitulate these lipid metabolic dynamics similar to the parent

tissues from whence they came.

Given the increased potential for neutral lipid accumulation and lipid hydrolysis in aUC colonoids, we

hypothesized that the oversupply of lipids contributes to elevated O_2 consumption in aUC colonoids during

differentiation. Thus, we exposed control and aUC colonoids to a nutrient-depleted medium with or without

etomoxir, a carnitine palmitoyl-transferase 1a (CPT1a) inhibitor that inhibitis fatty acid oxidation (Fig. 4e).

With negligible external sources of metabolic substrates, aUC colonoid mean basal OCR was 78% greater

than that of controls (Fig. 4f,g). Etomoxir reduced basal OCR in aUC colonoids to the level of control

colonoids (Fig. 4g). Further, etomoxir did not impact maximal respiration (Extended Data Fig. 6a). Etomoxir

significantly reduced proton leak in aUC but not control colonoids, suggesting that enhanced FAO drives

proton leak in aUC (Fig. 4h). Compared to control colonoids, we observed a significant increase in ATP-

linked OCR in aUC colonoids under this nutrient-depleted condition, with a significant decrease in ATP-

linked OCR in aUC with etomoxir inhibition (Fig. 4i). Non-mitochondrial respiration was elevated in aUC

colonoids compared to control with no effect observed with etomoxir treatment in either group, which

suggests that etomoxir specifically modulated mitochondrial respiration in this model (Extended Data Fig.

6b). These results indicate that there is enhanced lipid accumulation and oxidation in the differentiating

active UC epithelium which is necessary for the hypermetabolic phenotype during differentiation.

Lipotoxicity precedes *CXCL1* overexpression during colonoid differentiation

Lipid exposure induces stemness in intestinal progenitor cells[41]; hence, we wondered whether lipotoxicity

during colonoid differentiation may contribute to the impaired differentiation and chemokine expression

observed in aUC colonoids (Fig. 1c, Fig. 2c). Thus, we exposed control spheroids to BSA-bound palmitic

acid (BSA-PA) during the initial 24 hrs of differentiation. We found a dose-dependent increase in the gene

expression of the uncoupling protein *UCP2*, fatty acid transporter *FABP6*, and a decrease of the crypt-base

goblet cell marker, *WFDC2*[29] (Extended Data Fig. 6c) in response to BSA-PA exposure. Consequently, we

hypothesized that lipid-induced stress and toxicity to the colon epithelial progenitors in aUC may contribute

to the observed enhanced chemokine production in aUC colonoids under non-stimulated conditions *in vitro*.

Hence, we exposed non-IBD colonoids to BSA-PA in ODM for 3 days and continued differentiation without

PA until day 5 (Fig. 4k). Compared to control, PA exposure impaired colonoid morphology and induced a

significant upregulation of *CXCL1* in all colonoid lines (Fig 4j,m). Overall, these data suggest that fatty acid

exposure may impair differentiation and induce inflammation in differentiating colon epithelial cells.

PPAR-alpha contributes to lipid-induced hypermetabolic stress during colonoid differentiation

To understand the regulatory networks that may explain the hypermetabolic nature of differentiating aUC

colonoids, we applied Causal Network Analysis of our bulk transcriptomic colonoid data[35]. We identified the

PPARA-RXRA network as the most significant causal factor (Fig. 5a). Other notable networks include HIF3A

(endogenous lipid sensor[42]), tyrphostin (role in metabolic uncoupling[43]), MLXIPL (role in carbohydrate

sensing and glucose-induced lipogenesis[44]), and SH3GLB1 (contributes to lipolysis and lipid droplet

clearance[45]).

We focused on PPAR-α as it regulates a family of genes central to glucose and lipid metabolism[46]. Analyses

of epithelial lineage-specific PPARA expression in the scRNA-seq data from treatment-naïve UC colon

epithelium[39] showed the upregulation of PPARA majorly in progenitor cells (stem, TA2, cycling TA, secretory

TA) in the inflamed and non-inflamed colon epithelial regions compared to healthy samples (Extended Data

Fig. 7a). We did not see any differences in *PPARA* mRNA expression in our bulk RNA-seq data between

aUC and control colonoids differentiated for 3 days; however, we found an enhanced PPAR-α activity in

nuclear extracts from aUC colonoids compared to controls at the early stages (24 hr) of differentiation

(Extended Data Fig. 7b).

Given the upregulation of PPAR-α-induced genes and PPAR-α DNA binding in aUC colonoids, we

wondered whether PPAR-α activation is sufficient to induce the observed hypermetabolic phenotype.

Treatment of control colonoids with the PPAR-α agonist fenofibrate during differentiation significantly

reduced colonoid budded morphology in every patient colonoid line tested (Fig. 5b-d). Consistent with the

differentiation pattern in aUC colonoids, we found a significant downregulation of the crypt-base goblet cell

marker[29] *WFDC2,* in control colonoids exposed to fenofibrate (Fig. 5e). Fenofibrate enhanced metabolic

substrate utilization as we observed increased glucose consumption over two days in every colonoid line

compared to untreated control (Fig. 5f). Overall, glucose consumption was not accompanied by significant

lactate production suggesting a negligible induction of anaerobic glycolysis (Fig. 5g). Fenofibrate treatment

enhanced UCP2 expression during colonoid differentiation (Fig. 5h). Consistent with aUC lipid metabolic

phenotypes, we observed a significant accumulation of neutral lipids and enhancement of LPL activity in

control colonoids exposed to fenofibrate (Fig. 5i-k). We also observed enhanced mitochondrial mass, and

mitochondrial ROS in control colonoids upon fenofibrate treatment (Fig. 5l,m). Following these observations,

we hypothesized that fenofibrate treatment during differentiation would result in organoids with a

hypermetabolic bioenergetic profile similar to that seen previously in aUC. Indeed, differentiation of control

colonoids exposed to fenofibrate resulted in significantly higher basal, maximal, proton leak, and non-

mitochondrial OCR with no significant increase in ATP-linked OCR (Fig. 5n,o). Finally, fenofibrate treatment

induced LDH release (a marker of cytotoxicity) consistent with our observation in differentiating aUC

colonoids (Fig. 1k, Fig. 5p). These findings suggest that there is enhanced PPAR-α activity in active UC and

such activity is sufficient to induce to hypermetabolic stress and impair differentiation in colon epithelial

cells.

Pharmacological inhibition of PPAR-α reprograms epithelial metabolism in aUC colonoids

Following the observed contribution of PPAR-α activity to lipid accumulation and hypermetabolic stress

during differentiation, we reasoned that inhibition of PPAR-α during aUC colonoid differentiation may

prevent endogenous lipid accumulation and improve metabolic and cellular stress. We supplemented ODM

with the PPAR-α antagonist, GW6471 (GW) during differentiation of aUC colonoids. GW treatment in aUC

colonoids suppressed neutral lipid accumulation during differentiation in every colonoid line compared to

untreated aUC colonoids (Fig. 6a,b). GW treatment in aUC colonoids also downregulated *LPL* and *UCP2*

gene expression (Extended Data Fig. 8a, Fig. 6c). We then determined mitochondrial bioenergetics in aUC

colonoids differentiated with or without GW in substrate-enriched medium (10 mM glucose, 1 mM pyruvate,

2 mM glutamine). We observed a significant reduction in basal, maximal, proton leak, and ATP-linked OCR

in GW-treated colonoids (Fig. 6). Luminescence assay for total cellular ATP showed no significant

differences in untreated and GW-treated aUC colonoids. These data suggest that PPAR-α inhibition reduces

lipid accumulation and normalizes cellular bioenergetics in active UC colon epithelial cells.

To test whether PPAR-α inhibition improved UC colon epithelial bioenergetics through altering FAO, we

examined the effect of GW on colonoids in the presence or absence of etomoxir. Blockade of FAO with

etomoxir did not affect basal OCR in GW-treated aUC colonoids (Fig. 6g,h). Similarly, etomoxir did not have

any effect on maximal OCR, proton leak, ATP-linked OCR, or non-mitochondrial OCR in GW-treated aUC

colonoids (Extended Data Fig. 8b-e). These data suggest that GW treatment suppressed the

hypermetabolic bioenergetic profile of aUC colonoids by reducing FAO. Since GW treatment starved aUC

colonoids of lipid-based metabolic substrates during the energy-demanding differentiation, we reasoned that

GW may have induced a metabolic switch to the usage of other substrates to sustain differentiation. Hence,

we assayed for extracellular glucose and lactate in ODM over 2 days of differentiation in aUC colonoids with

or without GW. We observed a significant increase in glucose consumption in all GW-treated aUC colonoid

lines compared to their paired untreated aUC colonoids without changes in lactate production, suggesting

an increased reliance of glucose as a substrate for OXPHOS (Fig. 6i,j).

We next examined whether the metabolic reprogramming in aUC colonoids with PPAR-α inhibition impacted

cellular health markers. GW treatment in aUC colonoids did not impact mitochondrial mass or mitochondrial

ROS levels compared to untreated aUC colonoids (Extended Data Fig. 8f,g). In contrast, GW treatment

significantly reduced LDH release (cytotoxicity marker) in every colonoid line compared to untreated aUC

colonoids (Fig. 6k). Considering that lipotoxicity contributed to chemokine expression in differentiating

colonoids (Fig. 4n), we hypothesized that the prevention of neutral lipid accumulation by GW during

differentiation would impact epithelial chemokines secreted in aUC colonoids (Fig. 2d-f). Hence, we

collected ODM from colonoids differentiated for 2 or 5 days and found that GW treatment in aUC colonoids

significantly suppressed extracellular CXCL1 protein in day 2 and day 5 ODM collected from all colonoid

lines tested (Fig. 6l). There was no significant effect of GW treatment on extracellular CXCL11 and a trend

toward a reduction in extracellular CCL2 on day 2 (p=0.053) while there was no effect on day 5 (Extended

Data Fig. 8h,i).

Considering that aUC colonoids exhibited morphologic evidence of impaired differentiation and

corresponding downregulated expression of the crypt-base goblet cell marker, *WFDC2* (Fig. 1a-c), we

asked whether the remodeling of colonoid metabolism in aUC with GW treatment impacted differentiation markers. Blinded visual assessment of colonoid morphology showed no consistent impact of GW on colonoid budding (Fig. 6m,n). To complement these findings, we assessed the mRNA expression of stemness markers (*LGR5* and *ASCL2*) and goblet cell markers *WFDC2* and *MUC2*. The expression of stemness markers reduced over time in every aUC colonoid line with GW treatment, with significant downregulation in *ASCL2* (Fig. 8p). *WFDC2* significantly increased in GW-treated colonoid lines compared to untreated aUC colonoids while the increase in *MUC2* expression was not statistically significant (Fig. 8o). Overall, these data suggest that PPAR-α inhibition in aUC colonoids reduces chemokine secretion and augments molecular markers of epithelial differentiation.

Discussion

Leveraging clinical samples to develop human IBD pre-clinical models is important for determining mechanisms of human disease pathogenesis. We used patient-derived colon epithelial organoids as a human model to study epithelial metabolic dysfunction in pediatric UC. We report profound elevation in oxygen consumption in aUC colonoids during differentiation related primarily to increased proton leak. This hypermetabolic phenotype was associated with increased chemokine secretion, oxidative stress, cytotoxicity, and impaired differentiation. The aUC colonoids showed an elevated capacity for lipid accumulation during differentiation and their hypermetabolic phenotype required FAO. Our data indicate PPAR-α drives altered lipid metabolism and OXPHOS in aUC colonoids and that pharmacological inhibition of PPAR-α normalizes abnormal metabolism, chemokine expression, and differentiation in aUC epithelial cells.

In our study, colonoids from active UC patients showed increased mitochondrial respiration during differentiation without a concomitant change in ATP levels. The normal colon epithelium is less dependent on highly oxidative processes for energy generation[47,48]. Accordingly, our data in non-IBD control colonoids showed marginal increases in median oxygen consumption during the energy-demanding differentiation process. Hypermetabolism may channel energetic demands towards molecular programs that favor a

hypersecretory phenotype[49,50], which may explain the hypersecretion of chemokines in aUC colonoids

during differentiation. Furthermore, the oversupply of metabolic substrates may result in hypermetabolism

with associated mitochondrial and cellular stress[51]. Cellular dependency on lipids for energy generation is

inefficient since the oxidation of lipids consumes energy and induces metabolic uncoupling[52] as observed

with increased proton leak and UCP2 expression in our aUC colonoids. This notion is corroborated by our

finding of *UCP2* induction in healthy colonoids exposed to palmitic acid. Our transcriptomic and functional

data suggest a lipid-related hypermetabolism during differentiation in epithelial aUC colonoids, which may

increase the energetic cost of differentiation and contribute to enhanced epithelial cellular stress in aUC.

The large PROTECT pediatric UC inception cohort study[19] examined rectal mucosa bulk RNA expression in

children with active UC and non-IBD controls and found extensive downregulation of genes related to

mitochondria and OXPHOS. Several of our findings in colonoids were in line with the PROTECT study

tissue data including increased expression of the uncoupling gene UCP2, and multiple lipid-related genes

and decreased epithelial MMP in active UC. Notably, the PROTECT study did not examine colon epithelial

respiration directly. While we did not observe any significant changes in mitochondrial genes in our aUC

colonoids (data not shown), we observed abnormally high mitochondrial respiration in aUC colonoids. In a

dextran sodium sulfate (DSS) mouse model of colitis[53], enhanced epithelial OXPHOS activity was reported

during the active stage of the disease at day 7 despite the downregulation of some mitochondrial genes.

These metabolic and transcriptional disparities in the active epithelium largely resolved to the level of

control upon recovery at day 28[53]. It is possible that the downregulation of mitochondrial genes is driven by

the inflammatory milieu of the colon mucosa which is absent in the sterile culture of colonoids in our study.

Our study provided functional evidence of lipid metabolic dysregulation in the pediatric UC epithelium. Few

metabolomic and animal studies have suggested the potential contribution of lipid metabolic dysregulation

to IBD[54]. In adults, onset of UC is predicted 4 years earlier by upregulated lipid biosynthesis, elongation,

and degradation pathways as measured by plasma metabolomics[55]. Elevated fecal fat is apparent in UC

patients (as reviewed previously[56]), and lipid dysregulation occurs in the colon tissue of the DSS-induced

mouse model of colitis[57]. These studies support our findings as they suggest colon epithelia may be

exposed to excess lipid or abnormally process lipid during intestinal inflammation – even before disease onset. Neutral lipid accumulation in lipid droplets may be a cytoprotective mechanism against lipotoxic-free non-esterified fats[58]. Conversely, aberrant lipid accumulation or lipid droplet hydrolysis may trigger lipotoxicity-related pathways including apoptosis, inflammation, ER stress, and autophagy[59]. The impact of long-chain fatty acid lipotoxicity on *CXCL1* expression is documented in hepatoma cells[60] similar to our findings in colonoids. Hence, it is plausible that approaches that suppress aberrant epithelial lipid accumulation, hydrolysis, and oxidation may improve colon epithelial function in UC.

Our study also suggests a dysregulation of PPAR-α network in the UC epithelium and colonoids. PPARA is a transcriptional regulator of several genes involved in glucose and lipid homeostasis[46], and, indeed, we observed elevated glucose consumption and neutral lipid accumulation in control colonoids treated with a PPAR-α agonist. PPAR-α activation was also associated with hypermetabolic phenotype, morphologic and molecular evidence of impaired differentiation, similar to our observation in aUC colonoids. Consistent with our study, PPAR-α agonism through fenofibrate exaggerated colitis in DSS-induced mouse models with a concomitant increase in bioactive serum lipids such as sphingomyelins and ceramides[61,62]. Furthermore, activation of fatty acid oxidation partly via PPAR-α promotes intestinal stemness in high-fat diet models[63], suggesting a detrimental effect on intestinal stem cell differentiation and maintenance of epithelial homeostasis. Earlier studies in adult UC showed that natural agonists of PPAR-α such as PUFAs, are disproportionally accumulated in the inflamed colon mucosa of UC patients which correlated positively with endoscopic disease activity[64,65]. Thus, an understanding of PPAR-α-regulated molecules that are specifically dysregulated in pediatric aUC samples may provide specific mechanistic targets for future studies.

In contrast, pharmacological inhibition of PPAR-α in aUC colonoids blocked aberrant neutral lipid accumulation and thus, suppressed lipid-dependent energy generation during differentiation. PPAR-α inhibition also reduced aUC colonoid hypermetabolism and proton leak. In this scenario, aUC colonoids consumed more glucose likely due to a metabolic switch to compensate for the loss of lipid-dependent oxidation for energy. This lipid-starvation state and the resulting metabolic switch enhanced expression of genes related to differentiation including *WFDC2*. Further, the PPAR-α antagonist reduced cytotoxicity and

expression of the chemokine CXCL1, which is critical for neutrophil infiltration in IBD[66]. Hypermetabolism

may divert energy expenditure towards translational processes that upregulate cytokine and stress

responses[49]; hence, it is plausible that the blockade of lipid's contribution to the hypermetabolic state in aUC

colonoids by PPAR-α inhibition reduces such responses. Animal studies have suggested the potential of

PPAR-α inhibition in ameliorating colitis induced by epithelial injury[67,68]. Overall, our study in human patient-

derived colonoids suggests that PPAR-α antagonism in the epithelium may be a viable treatment strategy

for reducing lipotoxicity and hypermetabolic stress in the UC epithelium.

In conclusion, our data in patient-derived epithelial colon organoids indicate that hypermetabolism and lipid

metabolic dysregulation are significant contributors to the epithelial pathogenesis of pediatric UC.

Furthermore, epithelial PPAR-α may be a therapeutic target for restoring colon epithelial homeostasis by

reducing dysregulated lipid metabolism and normalizing cellular respiration. The use of patient-derived

colonoids in this study opens novel research directions in colon epithelial metabolism that may identify

metabolic targets for next-generation therapies specifically directed toward healing the colon epithelia in

pediatric UC.

Methods

Human Study Approval and Informed Consent

Pediatric patient recruitment occurred at Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, and Lucile Packard Children's Hospital Stanford, Palo Alto, California. Patient data and biopsy collection followed approved protocols by the Research Ethics Committees at each institution. Four to eight colon biopsies (2–5 mm^3) from patients with active UC, inactive UC, and non-IBD controls were collected into ice-cold DMEM/F12 (Corning 10-092-CV) + 50 µg/mL Normocin (InvivoGen #ant-nr) during clinically-indicated lower endoscopy and transported on ice to the lab for immediate processing for spheroid generation. Patient demographics and clinical data are presented in **Supplementary Table 1**.

Biopsy processing and spheroid culture

Fresh biopsies were transferred into 5 mL of 2 mM PBS-EDTA and incubated for 45 min on a rotator in a cold room. Biopsies were transferred into 2.5 mL ice-cold sorbitol buffer (2% sorbitol, 1% sucrose, 1% BSA, 50 µg/mL Normocin) in a petri dish and crypts were gently scraped off using a college cotton tweezer under a dissecting microscope (Leica S9E). Big tissue pieces were removed with a 20 µL pipette tip and the crypts and buffer were transferred into a 15 mL tube. Crypts were sheared briskly with a 1000 µL pipette (80X total with 1 min interval) and the suspension was made up to 14 mL with ice-cold PBS (Gibco #10010-023). The suspension was centrifuged (300 x g, 5 mins, 4°C) and the pellet was resuspended in organoid wash medium, OWM, (DMEM/F12, 10% FBS, 50 µg/mL Normocin, and washed twice (300 x g, 5 mins, 4°C). The visible pellet was resuspended in growth factor-reduced Matrigel (Corning #356231) and 50 µL drops of the Matrigel mixture were allowed to polymerize at 37°C for at least 15 mins on pre-warmed 12-well plates. Upon stabilization, 700 µL of Intesticult organoid growth medium, OGM, (STEMCELL Technologies, #06010) containing 10 µM TGF-beta receptor inhibitor, SB431542 (Cayman Chemical, #13031), 10 µM ROCK inhibitor, Y27632 (STEMCELL Technologies, #72304), and 50 µg/mL Normocin. Crypts were placed in a tissue culture incubator (37°C, 5% CO2) and OGM was replaced every 2-3 days for the first 7-14 days.

Spheroids were considered stable and mature enough for expansion when >70% of spheroids are ~100 µm in diameter.

For spheroid expansion, OGM was aspirated and 1mL of ice-cold PBS was added to each well. The Matrigel domes were dislodged from the bottom of the plate with a 1000 µL pipette and the PBS-Matrigel mixture was transferred into a 15 mL tube. The Matrigel was broken up inside the tube with a 1000 µL pipette, made up to 5 mL with ice-cold PBS, and incubated on ice for 5 min. The suspension was centrifuged (300 x g, 5 mins, 4°C), 350 uL of TrypLE Express (Gibco #12604-013) was added to the spheroid pellets and incubated in a 37 °C water bath for 4-5 mins. Spheroids were broken up by pipetting 50X with a 1000 µL pipette followed by the addition of 5 mL ice-cold OWM. Spheroids were washed twice in OWM (300 x g, 5 mins, 4°C), the resulting pellets were resuspended in 3 parts of Matrigel to 1 part of OWM, and 50 µL drops of the Matrigel suspension were plated in pre-warmed 12-well plates and allowed to polymerize as previously described. For spheroid expansion, 2-3 drops of this suspension were plated per well of a 12-well plate and 800 µL of L-WRN conditioned medium[69,70] was added and replaced every other day.

For cryopreservation, spheroids underwent at least 3 passages before storage in our pediatric organoid biobank. Matrigel domes with matured spheroids (~80 µm) were dislodged in ice-cold PBS and centrifuged as previously described. Pellets were resuspended in freezing medium (90% FBS, 10% DMSO), and 1 mL of the suspension was added to a cryotube. Samples were slow-frozen in a Mr. Frosty freezing container at -80°C and transferred to liquid N_2 for long-term storage. Overall, the pooling of matured spheroids from two Matrigel domes in one cryotube allows for sufficient post-thaw recovery of spheroids.
All experiments used cryopreserved spheroids. Withdrawal of cryotubes from the LN_2 tank followed a quick thaw in a 37°C water bath for ~2mins. Cryotube contents were quickly transferred into pre-warmed 6 mL of OWM and centrifuged (300 x g, 5 mins, 20°C). Spheroid pellets were immediately resuspended in 5 mL OWM, washed once, resuspended in a 3:1 mix of Matrigel and OWM, and plated in 50 µL domes as previously described. Cryopreserved spheroids were passaged at least once before usage for all analyses.

Spheroids processing and plating for analysis

After 5-7 days after passaging with TrypLE Express, Matrigel domes with matured spheroids were dislodged

with ice-cold PBS as previously described and filtered with a 100 μm strainer (pluriSelect). Filtrates were

pelleted and washed once in OWM, manually counted, and replated in domes containing a 1:1 mix of

Matrigel and OWM (8-15 spheroids per μL depending on the analysis) and maintained in OGM. For all

analyses requiring differentiation into colonoids, replated spheroids were allowed to acclimatize overnight in

OGM before switching to Intesticult human organoid differentiation medium, ODM, (STEMCELL

Technologies, #100-0214) for the indicated differentiation durations with ODM changed every 2 days. In

some experiments, matured spheroids were filtered through a 40 μm strainer (Pluriselect) immediately after

passaging and cultured in OGM for 6-7 days before counting, replating, and differentiation when necessary.

To measure drug-induced responses, 2 μM FCCP (Sigma, #C2920), 50 μM Fenofibrate (Abcam,

AB120832), and 1 μM GW6471 (Sigma, #G5045) were added to ODM throughout the differentiation as

indicated in figure legends. The solvents served as vehicle controls for respective experiments (DMSO for

FCCP, and 100% ethanol for Fenofibrate and GW6471). ODM with drugs were removed and colonoids were

processed for analyses.

Spheroid Diameter and Colonoid Budding

Spheroids were passaged as previously described and filtered through a 40 μm strainer (Pluriselect).

Spheroids were manually counted and 750 spheroids were replated per 50 μL dome in pre-warmed 12-well

plates and cultured in OGM for 6 days. Phase contrast images were obtained on the Keyence BZ-X800

microscope.

To assess colonoid budding, matured spheroids were counted, replated in 50 μL Matrigel (15 spheroids/μL),

and allowed to acclimatize overnight in OGM. Each well was washed with PBS at room temperature (RT) for

3 min before switching to ODM for the indicated number of days. Phase contrast images were obtained on

the Keyence BZ-X800 microscope. All measurements of spheroid diameter and enumeration of colonoid

budding were performed in a manner blinded to colonoid diagnosis group using the Keyence BZ-X800

analyzer. The number of buds on each colonoid was presented relative to the total number of visible

colonoids per field.

Extracellular Flux Analyses

The XF Seahorse MitoStress Test (Agilent) was used to measure oxygen consumption rates (OCR), at

baseline, and after the injection of mitochondrial inhibitors[71]. Each spheroid line was replated in

quadruplicates as described previously in 15 µL matrigel domes (8 spheroids/µL) containing a 1:1 mix of

Matrigel and OWM into Seahorse XF96 cell culture plates (Agilent #101085-004) and cultured overnight in

150 µL OGM. The four edges of the plate served as blank with only a 1:1 mix of Matrigel and OWM. For

some experiments, 50 µL of OGM was removed after 24 hrs into an empty 96-well plate and stored at -80°C

for later measurements of LDH activity. Where differentiation was needed, spheroids were washed once

with PBS on the following day and differentiated in 200 µL ODM into colonoids for the indicated number of

days in the figure legends. For some experiments, 50 µL of ODM was removed from all wells after 48 hr into

an empty 96-well plate and frozen at -80°C for later measurements of glucose, lactose, and LDH activity.

On the day of the assay, ODM was removed from colonoids, washed once with 200 µL PBS, and replaced

with 150 µL DMEM assay medium (Agilent, #103334-100), pH 7.4 containing 10 mM glucose, 2 mM

glutamine, and 1 mM pyruvate (all from Agilent). Plates were incubated in a non-CO_2 incubator for 1 hr and

OCR was measured at baseline and after a sequential injection of oligomycin (final concentration, 4µM),

FCCP (final concentration, 2 µM), and rotenone/antimycin (final concentration, 100 nM/1 µM) using the

XF96[e] Extracellular Flux assay kits on the XF96[e] analyzer.

The Glycostress test (Agilent) was used to measure extracellular acidification rates (ECAR) in colonoids

after a 3-day differentiation. On the day of the assay, ODM was replaced with base medium (Agilent

#103575-100) lacking glucose, pyruvate, and glutamine and incubated in a non-CO_2 incubator for 1 hr.

Baseline ECAR was assessed using the XF96^e analyzer (Agilent) followed by sequential injections of

glucose, oligomycin, and 2-deoxy glucose at final concentrations of 40 μM, 4 μM, and 500 mM, respectively.

Raw data were normalized with DNA content as described below.

For etomoxir experiments, colonoids were differentiated for 3 days. On the day of the assay, ODM was

replaced with assay medium (Agilent #103575) lacking glucose, pyruvate, and glutamine, and incubated in

a non-CO_2 incubator for 40 min. Etomoxir (Sigma, #E1905) was prepared in the assay medium, quickly

added to designated wells within 5 mins at a final concentration of 44 μM, and returned to the non-CO_2

incubator for 15 min. OCR assessment followed as described previously.

DNA extraction

For the normalization of colonoid extracellular flux data, DNA normalization protocol was used as previously

described with modifications[71]. DNA was extracted using the QIAamp DNA Micro kit (Qiagen #56304).

Briefly, the medium was removed from the cell culture plates and 45 μL of ATL lysis buffer was added per

well, followed by 5 μL of Protein kinase K. The plate was incubated at 56°C for 1 hr, quadruplicate wells

were pooled into 1.5 mL tubes and returned to 56°C incubator for 1 hr. DNA extraction continued according

to the manufacturer's instructions. The DNA content of each colonoid line was calculated relative to the

blank wells and used to normalize raw data.

Total ATP measurement

Spheroids were plated in clear flat bottom 96-well tissue culture plates (Falcon #353075) as described for

extracellular flux analyses and differentiated for the indicated number of days. Total ATP levels were

assayed with the luminescent ATP detection assay kit (Abcam, ab113849), according to the manufacturer's

instructions and luminescence was recorded using the Spectramax id5 plate reader (Molecular Devices).

ATP levels were normalized by measuring DNA content in a parallel plate as described for extracellular flux

analyses.

Mitochondrial Superoxide (MitoSOX) Analysis

Spheroids were plated in black 96-well tissue culture plates with clear bottoms (Corning, #3603) as

described for extracellular flux analyses and differentiated for the indicated number of days in the figure

legends. Mitochondrial ROS was estimated with the fluorescent probe MitoSOX red (Thermofisher,

#M36008). Briefly, a 5 mM MitoSOX stock solution was prepared in DMSO and diluted with HBSS to a

working concentration of 3 µM. Colonoids were washed twice with HBSS and the MitoSOX working solution

was added to the colonoids and incubated for 30 min (37^{o}C, 5% CO2). The colonoids were washed 3 times

with HBSS and MitoSOX fluorescence was measured at 396/610 nm using the Spectramax iD5 plate reader

(Molecular Devices). Data were normalized by measuring DNA content as described for extracellular flux

analyses.

Lactate Dehydrogenase (LDH) release

OGM and ODM stored from Seahorse XF96 cell culture plates as described above were thawed on ice and

extracellular LDH activity was measured using the LDH-Glo Cytotoxicity assay (Promega, #J2380).

Samples were diluted 40X in ice-cold storage buffer (200mM Tris-HCl (pH 7.3), 10% Glycerol, 1% BSA),

and LDH activity was measured according to the manufacturer's instructions. Luminescence was recorded

using the Spectramax id5 plate reader (Molecular Devices), and the data was normalized with DNA content

from the corresponding Seahorse plate as described previously. LDH release from spheroids and colonoids

was estimated relative to blank wells in the Seahorse XF96 cell culture plate.

Extracellular Glucose and Lactate

ODM from colonoids stored from Seahorse XF96 cell culture plates as previously described were thawed on

ice and extracellular glucose and lactate were measured using the Glucose-Glo (Promega, #J6021) and

Lactate-Glo (Promega, #J5021) assays, respectively. Samples were diluted 50X in ice-cold PBS and the

measurements continued according to the manufacturer's instructions. Luminescence was recorded using the Spectramax id5 plate reader (Molecular Devices), and the data was normalized with DNA content from the corresponding Seahorse plate as described previously. Glucose consumed or lactate produced were estimated relative to ODM in blank Seahorse XF96 cell culture wells.

Flow Cytometry

<u>Mitochondrial mass (mtMass) analyses</u>

For mtMass analyses, we labeled the mitochondria in spheroids and colonoids using the MitoTracker Green FM probe (Cell Signaling, #9074). Briefly, single cells from matured spheroids (15 spheroids/µL) and corresponding colonoids were obtained by the addition of 900 µL TrypLE Express followed by incubation at 37°C for 10 mins. Single cells were released by vortexing for 20 seconds with 10-second intervals, and washed in FACS buffer (5% FBS in PBS). Pellets were resuspended in 150 nM mtGreen solution prepared in phenol-red free DMEM/F12 + 5% FBS, followed by incubation for 30 mins at 37°C in a 5% CO_2 incubator. Cells were washed twice in 2 mL PBS and resuspended in 500 µL of ice-cold PBS. The cells were passed through Falcon FACS tubes with a cell strainer cap (Fisher, #0877123) and incubated with 5 µL of the live-dead stain, 7-Aminoactinomycin D (7-AAD), for 5 mins on ice. The mean fluorescence intensity (MFI) of mtGreen was measured in 5000 events using the FITC channel and following 7-AAD exclusion (PE-Cy5) on the Agilent NovoCyte Flow Cytometer. Analysis and images were obtained using the FlowJo software v.10.

<u>Mitochondrial Membrane Potential (MMP)</u>

Single cells from colonoids were processed as previously described and MMP was assessed with the MitoProbe JC-1 assay kit (Thermo Fisher Scientific, MP34152) following the manufacturer's protocol. Briefly, Cells were incubated with 1 µM JC-1 in PBS for 30 mins at 37°C, 5% CO_2. The cells were washed once with PBS and passed through Falcon FACS tubes with a cell strainer cap and incubated with 5 µL of 7-AAD, for 5 mins on ice. 10000 events from each line were recorded using the Agilent NovoCyte Flow Cytometer. After 7-AAD exclusion, MMP was calculated as the MFI ratio of the red (PE)/green (FITC) channels.

BODIPY Neutral lipids

BODIPY Neutral lipids

Matured spheroids from each line were replated (15 spheroids/µL) in two wells of a 24-well plate and

allowed to stabilize overnight in OGM. For each line, one well was switched to ODM to initiate differentiation

into colonoids for 2 days while the other well continued in OGM. Single cells were processed from spheroids

and colonoids from each line as previously described and cell pellets were fixed using the Cytofix/Cytoperm

kit (BD Biosciences, #555028) according to the manufacturer's instructions. Neutral lipid droplets in cells

were stained with 1µg/mL BODIPY 493/503 (Thermo Fisher Scientific, #D3922) in PBS for 1 hr at RT with

gentle mixing every 20 mins. Cells were washed once with 2 mL PBS and passed through Falcon FACS

tubes with a cell strainer cap. BODIPY MFI was recorded in 5000 events using the FITC channel on the

Agilent NovoCyte Flow Cytometer. Analysis and images were obtained using the FlowJo software v.10.

Immunofluorescence Imaging

Colonoid Embedding in Optical Cutting Temperature (OCT) compound

Colonoid Embedding in Optical Cutting Temperature (OCT) compound

Matured spheroids from each line were replated (15 spheroids/µL, 3:1 mixture of Matrigel and OWM) as

previously described in 6 domes per well of a 6-well plate. After a 3-day differentiation, colonoids were fixed

with 5 mL of 2% Paraformaldehyde and 0.1% Glutaraldehyde for 30 mins at RT. Colonoids were washed 3X

with 5 mL PBS for 10 min, and each Matrigel dome was carefully scooped into a 50 mL tube with 20%

sucrose in PBS. After 2-3 days following the sinking of all domes, 6 domes/line was embedded in Tissue

Tek OCT compound (Sakura Finetek, #4583) and frozen in liquid N_2 using the Seal'N Freeze cryotray and

box (Fisher Scientific, #NC1877501). From the Colonoid OCT block, 10 µm thick cryosections were cut on

slides and stored at -80ºC. For immunostaining, cryosections were washed once with PBS for 15 secs at RT

followed by autofluorescence quenching by incubation with 10 mM Sodium Borohydride in PBS twice for 5

mins at RT. Sections were washed thrice in PBS for 10 minutes each, followed by permeabilization with

0.15% Triton X-100 for 15 mins at RT. Slides were washed thrice with PBS and incubated with the blocking

buffer (3% BSA, 5% donkey serum, and 5% goat serum in PBS) for 1 hr at RT. This was followed by

overnight incubation with primary antibodies rabbit anti-UCP2 (1:100, Proteintech, #11081-1-AP) and mouse anti-COX4 (1:200, Invitrogen, GT6310) in the blocking buffer in a humidified chamber at 4°C. Slides were washed twice with PBS for 15 secs, followed by 2X wash in PBS for 5 mins. Slides were incubated in a humidified chamber for 1 hr at RT in secondary antibodies: donkey anti-rabbit Alexa Flour 647 (1:1000, Invitrogen, #A31573) and goat anti-mouse Alexa Flour 488 (1:1000, Invitrogen, #A11001) prepared in blocking buffer. Slides were washed 3X in PBS and counterstained with Phalloidin (1:250, ThermoFisher Scientific #T7471) in PBS for 30 min at RT. Slides were washed once in distilled water and mounted in Prolong Glass Antifade Mountant with NucBlue Stain containing the Hoechst 33342 DNA marker (ThermoFisher Scientific, #P36983). Images were acquired on the Keyence BZ-X800 microscope. Images were merged with the Keyence BZ-X800 software analyzer and cells expressing DAPI and those co-expressing UCP2 and COX4 were counted on at least four high-power fields (60X oil objective) per patient colonoid line.

Colonoid whole-mount staining

For whole-mount staining, matured spheroids were replated (15 spheroids/μL, 1:1 mixture of Matrigel and OWM) as previously described in 8-well chamber μ-slides (IBIDI, #80826). After differentiation, colonoids were washed with PBS at RT and fixed with 500 μL of 4% PFA for 30 mins at RT. For UCP2 staining, colonoids were washed 2X with 500 μL PBS for 5 mins and permeabilized with 0.2% Triton X-100 in PBS for 30 mins at RT followed by 2X wash in PBS. Samples were incubated with 50 mM ammonium chloride (NH_4Cl) in PBS for 30 mins at RT to quench potential autofluorescence. After a 2X wash in PBS, samples were incubated in blocking buffer (3% BSA, 5% donkey serum in PBS) for 1 hr at RT and incubated in primary antibody- rabbit anti-UCP2 (1:50, Proteintech, #11081-1-AP) overnight and secondary antibody- donkey anti-rabbit Alexa Flour 647 (1:500, Invitrogen, #A31573) for 1 hr at RT, and incubated with Phalloidin as previously described. Samples were mounted in 200 μL Fructose-glycerol clearing solution[72] and incubated for 30 mins at RT. Whole-mount images were acquired using the Stellaris 8 Inverted Confocal Microscope (Leica) using the 20x oil objective and individual channels were merged using the Leica LAS X software.

For visualizing neutral lipids, colonoids differentiated for 2 days in chamber slides were fixed in 4% PFA, washed, and incubated in 50 mM NH_4Cl as previously described. Then colonoids were stained with 1 µg/mL BODIPY 493/503 (Thermo Fisher Scientific, #D3922) in PBS for 1 hr at RT, washed twice in PBS, and incubated with Phalloidin as previously described. Chamber slides were washed once in distilled water and mounted in 2 drops of Prolong Glass Antifade Mountant with NucBlue Stain containing the Hoechst 33342 DNA marker (ThermoFisher Scientific, #P36983). Samples were allowed to cure overnight at RT in the dark. Whole-mount images were acquired using the Stellaris 8 Inverted Confocal Microscope (Leica) as previously described.

RNA processing and Bulk RNA sequencing

RNA was processed concurrently from eight patient organoid lines per diagnosis using the RNeasy Micro Kit (Qiagen, #74004). Briefly, matured spheroids from each line (n=8 per diagnosis) were filtered through a 100 µm strainer (Pluriselect), replated (15 spheroids/µL, 2:1 mixture of Matrigel and OWM) in two wells of a 12-well plate (3 domes/well) and allowed to stabilize overnight in 800 µL OGM. The following day, one well of spheroids/line was processed for RNA extraction, while the other well was switched to ODM for differentiation into colonoids for 3 days. For RNA extraction, the medium was removed, and Matrigel domes were dislodged with PBS from the bottom of the plate as previously described. To digest and remove any traces of Matrigel, the suspension was centrifuged (300 x g, 5 mins, 4°C), 350 µL of TrypLE Express (Gibco) was added to the spheroid pellets and incubated in a 37°C water bath for 3 mins. The tube was swirled briefly followed by the addition of 5 mL ice-cold OWM. Samples were washed (300 x g, 5 mins, 4°C), and the supernatant was aspirated. Any remnants of OWM were carefully removed with a 200 µL pipette and the resulting pellets were resuspended in the lysis buffer from the RNeasy Micro Kit and stored immediately at -80°C. Further processing of RNA followed the manufacturer's instructions. RNA sequencing of poly-A enriched RNA libraries was carried out on the Illumina NovaSeq PE150 platform (Novogene). RNA-seq reads were aligned to the human reference hg38 genome and the nf-core/rnaseq (v3.5) workflow was used for developing the expression matrix.

Ingenuity Pathway Analysis (IPA)

Fold changes from the colonoid bulk RNA-seq data (Log2FC and p-adj values) were uploaded into the IPA

software and used to predict canonical pathways that underlie transcriptional data. All upregulated and

downregulated genes were concurrently analyzed in IPA with an expression p-value cutoff of 0.1. The top

biological and cellular pathways and causal networks were selected based on the fisher's exact test p-value

which measures the overlap of observed and predicted regulatory gene sets[35].

Single Cell Data Analyses
818
819
We utilized data from the established single-cell transcriptomic atlas of 68 colon biopsies obtained from 18

UC patients and 12 healthy controls[39]. Processed single-cell expression matrices and metadata for human

colon epithelial cells were obtained from the Single Cell Portal

(https://singlecell.broadinstitute.org/single_cell, accession SCP259). Expression counts were normalized

using the R (v.4.3.3) package Seurat (v.5.1.0). The reported PPARA, RXR, and lipid-related genes were

visualized using the R package ComplexHeatMap (v.2.18.0).

Quantitative RT PCR

For most analyses, total RNA was processed from spheroids or colonoids by matrigel removal and digestion

as previously described. Due to the short half-life of *UCP2* (ref) samples for *UCP2* analysis were plated in a

1:1 mixture of Matrigel and OWM, differentiated, and lysed directly in the culture well without Matrigel

removal. For all analyses, RNA was processed (RNeasy Micro Kit, Qiagen, #74004) and cDNA was

synthesized using the High-Capacity cDNA Reverse Transcription kit (Applied Biosystems, #4368814).

Gene expression was quantified on the QuantStudio 3 thermocycler using Taqman probes (**Supplementary**

Table 5). Fold changes were quantified using the $2^{-\Delta\Delta CT}$ method using *ACTB* or *GAPDH* as the reference

gene where appropriate.

LPL Activity

Spheroids were plated, differentiated for 2 days or 5 days, and processed for Matrigel removal as previously

described for RNA processing. Total colonoid protein was isolated using the Cell Lytic buffer (Sigma, #2978)

with 1X Protease and Phosphatase inhibitor cocktail (ThermoFisher Scientific, #78440). Colonoids were

incubated on ice for 30 mins with brief vortexing every 10 mins. Samples were centrifuged (10000 x g, 10

mins, 4°C), and the supernatant was transferred into fresh 1.5 mL tubes. Protein concentration was

determined using the BCA protein Assay kit (ThermoFisher Scientific, #23227). LPL activity in each sample

was determined in duplicates using the LPL activity assay kit (Cell Biolabs, #STA-610) following the

manufacturer's instructions, and the data was normalized to total protein concentration.

Nuclear PPAR-α activity assay

Three spheroids per group were differentiated in ODM for 24 hrs, pooled, and processed for Matrigel

removal as previously described for RNA processing. Nuclei isolation followed the established Omni-ATAC

protocol used for nuclei preparations for ATAC-seq[73]. Nuclei protein concentration was determined using the

BCA protein Assay kit (ThermoFisher Scientific, #23227). PPAR-α activity was assessed in the nuclear

extracts with the PPAR-α activity kit (Abcam, #ab133107) following the manufacturer's instructions.

Extracellular Chemokine assay

Spheroids were plated and differentiated in ODM for the indicated days in the figure legend, as previously

described for RNA processing. ODM was collected at the indicated days and frozen at -80°C until analyses.

For analyses involving differentiation until day 5, ODM was changed on day 2 before collection and

analyses on day 5. The total colonoid protein was determined using the Cell Lytic buffer with 1X Protease

and Phosphatase inhibitor as previously described. Extracellular chemokines in ODM on the indicated days

were assessed using the following kits (CXCL1: Abcam #ab190805, CXCL11: ThermoFisher #EHCXCL11,

CCL2: ThermoFisher #88-7399-22, CCL28: ThermoFisher #EHENC1) following manufacturer's instructions.

BSA-PA assay

Palmitic acid (PA) complexed to bovine serum albumin (BSA) was prepared based on a previously

described protocol with few modifications[60]. Briefly, 50 mg/mL of PA (Sigma, #P0500) was dissolved in

100% ethanol (150 x g, 20 mins, 37°C) to make a 195 mM PA stock solution. A 3 mM PA solution was

prepared in 10% fat-free BSA (Sigma, #A8806) in PBS and incubated at 47°C for 20 mins on a shaker. For

controls, an equivalent amount of EtOH was added to 10% fat-free BSA and processed as previously

described for PA. The BSA-PA and BSA-EtOH solutions were distributed into aliquots and stored at -20°C.

For colonoid exposure experiments, the solutions were diluted in ODM to prepare the desired

concentrations as described in figures and legends.

Statistical analyses

For all data, statistical analysis was conducted in Prism (v.10.0). Due to the varied statistical approaches

used in this study, statistical methods for each figure are outlined in individual figure legends. For this study,

eight patient-derived colonoids were generated per diagnosis (control, active UC, inactive UC). Wherever

representative colonoids were used, the number of independent colonoids and technical replicates used

were described in the relevant figure legend or method section. For differentially expressed genes from the

RNA-seq data, transcripts with $Log_2FC \geq \pm1$, FDR $P \leq 0.1$ were considered statistically significant. For all

other data P values less than 0.05 were considered statistically significant.

Data Availability

Bulk RNAseq data sets from spheroids and colonoids in this study were deposited into Gene Expression

Omnibus (GEO) under the accession number GSE… . Other appropriate data will be provided by the

corresponding author upon request.

References

1. Reinink, A. R., Lee, T. C. & Higgins, P. D. R. Endoscopic Mucosal Healing Predicts Favorable Clinical Outcomes in Inflammatory Bowel Disease: A Meta-analysis. *Inflamm. Bowel Dis.* **22**, 1859–1869 (2016).

2. Colombel, J. F. *et al.* Early mucosal healing with infliximab is associated with improved long-term clinical outcomes in ulcerative colitis. *Gastroenterology* **141**, 1194–1201 (2011).

3. Hyams, J. S. *et al.* Clinical and biological predictors of response to standardised paediatric colitis therapy (PROTECT): a multicentre inception cohort study. *The Lancet* **393**, 1708–1720 (2019).

4. Danese, S. *et al.* Upadacitinib as induction and maintenance therapy for moderately to severely active ulcerative colitis: results from three phase 3, multicentre, double-blind, randomised trials. *The Lancet* **399**, 2113–2128 (2022).

5. Sandborn William J. *et al.* Tofacitinib as Induction and Maintenance Therapy for Ulcerative Colitis. *N. Engl. J. Med.* **376**, 1723–1736 (2017).

6. Sands, B. E. *et al.* Ustekinumab as Induction and Maintenance Therapy for Ulcerative Colitis. *N. Engl. J. Med.* **381**, 1201–1214 (2019).

7. Rosen, M. J., Dhawan, A. & Saeed, S. A. Inflammatory Bowel Disease in Children and Adolescents. *JAMA Pediatr.* **169**, 1053–1060 (2015).

8. Ye, Y., Manne, S., Treem, W. R. & Bennett, D. Prevalence of Inflammatory Bowel Disease in Pediatric and Adult Populations: Recent Estimates From Large National Databases in the United States, 2007–2016. *Inflamm. Bowel Dis.* **26**, 619–625 (2020).

9. Kappelman, M. D., Moore, K. R., Allen, J. K. & Cook, S. F. Recent trends in the prevalence of Crohn's disease and ulcerative colitis in a commercially insured US population. *Dig. Dis. Sci.* **58**, 519–525 (2013).

10. Kappelman, M. D. *et al.* Direct Health Care Costs of Crohn's Disease and Ulcerative Colitis in US Children and Adults. *Gastroenterology* **135**, 1907–1913 (2008).

11. Ardizzone, S. *et al.* Mucosal healing predicts late outcomes after the first course of corticosteroids for

newly diagnosed ulcerative colitis. *Clin. Gastroenterol. Hepatol. Off. Clin. Pract. J. Am. Gastroenterol.*

Assoc. **9**, 483-489.e3 (2011).

12. Rogler, G., Vavricka, S., Schoepfer, A. & Lakatos, P. L. Mucosal healing and deep remission: what does

it mean? *World J. Gastroenterol.* **19**, 7552–7560 (2013).

13. Neurath, M. F. & Travis, S. P. L. Mucosal healing in inflammatory bowel diseases: a systematic review.

Gut **61**, 1619–1635 (2012).

14. Adolph, T. E. *et al.* The metabolic nature of inflammatory bowel diseases. *Nat. Rev. Gastroenterol.*

Hepatol. **19**, 753–767 (2022).

15. Aldars-García, L., Gisbert, J. P. & Chaparro, M. Metabolomics Insights into Inflammatory Bowel

Disease: A Comprehensive Review. *Pharm. Basel Switz.* **14**, 1190 (2021).

16. Fu, S.-C. *et al.* Excessive Mitochondrial Fission Suppresses Mucosal Repair by Impairing Butyrate

Metabolism in Colonocytes. *Inflamm. Bowel Dis.* **30**, 114–124 (2024).

17. Sifroni, K. G. *et al.* Mitochondrial respiratory chain in the colonic mucosal of patients with ulcerative

colitis. *Mol. Cell. Biochem.* **342**, 111–115 (2010).

18. Santhanam, S. *et al.* Mitochondrial Electron Transport Chain Complex Dysfunction in the Colonic

Mucosa in Ulcerative Colitis. *Inflamm. Bowel Dis.* **18**, 2158–2168 (2012).

19. Haberman, Y. *et al.* Ulcerative colitis mucosal transcriptomes reveal mitochondriopathy and

personalized mechanisms underlying disease severity and treatment response. *Nat. Commun.* **10**, 38

(2019).

20. Novak, E. A. & Mollen, K. P. Mitochondrial dysfunction in inflammatory bowel disease. *Front. Cell Dev.*

Biol. **3**, (2015).

21. Clevers, H. Modeling Development and Disease with Organoids. *Cell* **165**, 1586–1597 (2016).

22. Ojo, B. A., VanDussen, K. L. & Rosen, M. J. The Promise of Patient-Derived Colon Organoids to Model

Ulcerative Colitis. *Inflamm. Bowel Dis.* **28**, 299–308 (2022).

23. Dotti, I. *et al.* Alterations in the epithelial stem cell compartment could contribute to permanent changes

in the mucosa of patients with ulcerative colitis. *Gut* **66**, 2069–2079 (2017).

24. Sarvestani, S. K. *et al.* Cancer-predicting transcriptomic and epigenetic signatures revealed for

ulcerative colitis in patient-derived epithelial organoids. *Oncotarget* **9**, 28717–28730 (2018).

25. Arnauts, K. *et al.* Ex Vivo Mimicking of Inflammation in Organoids Derived From Patients With

Ulcerative Colitis. *Gastroenterology* **159**, 1564–1567 (2020).

26. Nanki, K. *et al.* Somatic inflammatory gene mutations in human ulcerative colitis epithelium. *Nature* **577**,

254–259 (2020).

27. Howell, K. J. *et al.* DNA Methylation and Transcription Patterns in Intestinal Epithelial Cells From

Pediatric Patients With Inflammatory Bowel Diseases Differentiate Disease Subtypes and Associate

With Outcome. *Gastroenterology* **154**, 585–598 (2018).

28. Beumer, J. & Clevers, H. Hallmarks of stemness in mammalian tissues. *Cell Stem Cell* **31**, 7–24 (2024).

29. Parikh, K. *et al.* Colonic epithelial cell diversity in health and inflammatory bowel disease. *Nature* **567**,

49–55 (2019).

30. Burclaff, J. *et al.* A Proximal-to-Distal Survey of Healthy Adult Human Small Intestine and Colon

Epithelium by Single-Cell Transcriptomics. *Cell. Mol. Gastroenterol. Hepatol.* **13**, 1554–1589 (2022).

31. Nicholls, D. G. Mitochondrial proton leaks and uncoupling proteins. *Biochim. Biophys. Acta BBA -*

Bioenerg. **1862**, 148428 (2021).

32. Pelia, R. *et al.* Profiling non-coding RNA levels with clinical classifiers in pediatric Crohn's disease. *BMC*

Med. Genomics **14**, 194 (2021).

33. van Heel, D. A. *et al.* Inflammatory bowel disease susceptibility loci defined by genome scan meta-

analysis of 1952 affected relative pairs. *Hum. Mol. Genet.* **13**, 763–770 (2004).

34. Garrity-Park, M. M., Loftus, E. V., Sandborn, W. J., Bryant, S. C. & Smyrk, T. C. MHC Class II alleles in

ulcerative colitis-associated colorectal cancer. *Gut* **58**, 1226–1233 (2009).

35. Krämer, A., Green, J., Pollard, J. & Tugendreich, S. Causal analysis approaches in Ingenuity Pathway

Analysis. *Bioinforma. Oxf. Engl.* **30**, 523–530 (2014).

36. TeSlaa, T. & Teitell, M. A. Techniques to monitor glycolysis. *Methods Enzymol.* **542**, 91–114 (2014).

37. Althaher, A. R. & Alwahsh, M. An overview of ATP synthase, inhibitors, and their toxicity. *Heliyon* **9**,

e22459 (2023).

38. Benatzy, Y., Palmer, M. A. & Brüne, B. Arachidonate 15-lipoxygenase type B: Regulation, function, and

its role in pathophysiology. *Front. Pharmacol.* **13**, 1042420 (2022).

39. Smillie, C. S. *et al.* Intra- and Inter-cellular Rewiring of the Human Colon during Ulcerative Colitis. *Cell*

178, 714-730.e22 (2019).

40. Umar, S. Intestinal Stem Cells. *Curr. Gastroenterol. Rep.* **12**, 340–348 (2010).

41. Beyaz, S. *et al.* High fat diet enhances stemness and tumorigenicity of intestinal progenitors. *Nature*

531, 53–58 (2016).

42. Fala, A. M. *et al.* Unsaturated fatty acids as high-affinity ligands of the C-terminal Per-ARNT-Sim

domain from the Hypoxia-inducible factor 3α. *Sci. Rep.* **5**, 12698 (2015).

43. Soltoff, S. P. Evidence that tyrphostins AG10 and AG18 are mitochondrial uncouplers that alter

phosphorylation-dependent cell signaling. *J. Biol. Chem.* **279**, 10910–10918 (2004).

44. Abdul-Wahed, A., Guilmeau, S. & Postic, C. Sweet Sixteenth for ChREBP: Established Roles and

Future Goals. *Cell Metab.* **26**, 324–341 (2017).

45. Liu, Y. *et al.* Bif-1 deficiency impairs lipid homeostasis and causes obesity accompanied by insulin

resistance. *Sci. Rep.* **6**, 20453 (2016).

46. Bougarne, N. *et al.* Molecular Actions of PPARα in Lipid Metabolism and Inflammation. *Endocr. Rev.* **39**,

760–802 (2018).

47. Konjar, Š., Pavšič, M. & Veldhoen, M. Regulation of Oxygen Homeostasis at the Intestinal Epithelial

Barrier Site. *Int. J. Mol. Sci.* **22**, 9170 (2021).

48. Mas-Bargues, C. *et al.* Relevance of Oxygen Concentration in Stem Cell Culture for Regenerative

Medicine. *Int. J. Mol. Sci.* **20**, 1195 (2019).

49. Sturm, G. *et al.* OxPhos defects cause hypermetabolism and reduce lifespan in cells and in patients

with mitochondrial diseases. *Commun. Biol.* **6**, 1–22 (2023).

50. Sercel, A. J. *et al.* Hypermetabolism and energetic constraints in mitochondrial disorders. *Nat. Metab.* **6**, 192–195 (2024).

51. Picard, M. & Turnbull, D. M. Linking the metabolic state and mitochondrial DNA in chronic disease, health, and aging. *Diabetes* **62**, 672–678 (2013).

52. Maassen, J. A., Romijn, J. A. & Heine, R. J. Fatty acid-induced mitochondrial uncoupling in adipocytes as a key protective factor against insulin resistance and beta cell dysfunction: a new concept in the pathogenesis of obesity-associated type 2 diabetes mellitus. *Diabetologia* **50**, 2036–2041 (2007).

53. Lan, A. *et al.* Mitochondrial remodeling and energy metabolism adaptations in colonic crypts during spontaneous epithelial repair after colitis induction in mice. *Free Radic. Biol. Med.* **205**, 224–233 (2023).

54. Boldyreva, L. V., Morozova, M. V., Saydakova, S. S. & Kozhevnikova, E. N. Fat of the Gut: Epithelial Phospholipids in Inflammatory Bowel Diseases. *Int. J. Mol. Sci.* **22**, 11682 (2021).

55. Hua, X. *et al.* Inflammatory Bowel Disease Is Associated With Prediagnostic Perturbances in Metabolic Pathways. *Gastroenterology* **164**, 147-150.e2 (2023).

56. Mourad, F. H., Barada, K. A. & Saade, N. E. Impairment of Small Intestinal Function in Ulcerative Colitis: Role of Enteric Innervation. *J. Crohns Colitis* **11**, 369–377 (2017).

57. Hamabata, T., Nakamura, T., Masuko, S., Maeda, S. & Murata, T. Production of lipid mediators across different disease stages of dextran sodium sulfate-induced colitis in mice. *J. Lipid Res.* **59**, 586–595 (2018).

58. Jarc, E. & Petan, T. Lipid Droplets and the Management of Cellular Stress. *Yale J. Biol. Med.* **92**, 435–452 (2019).

59. Lipke, K., Kubis-Kubiak, A. & Piwowar, A. Molecular Mechanism of Lipotoxicity as an Interesting Aspect in the Development of Pathological States-Current View of Knowledge. *Cells* **11**, 844 (2022).

60. Dahlquist, K. J. V., Voth, L. C., Fee, A. J. & Stoeckman, A. K. An Autocrine Role for CXCL1 in Progression of Hepatocellular Carcinoma. *Anticancer Res.* **40**, 6075–6081 (2020).

61. Qi, Y. *et al.* PPARα-dependent exacerbation of experimental colitis by the hypolipidemic drug fenofibrate. *Am. J. Physiol. Gastrointest. Liver Physiol.* **307**, G564-573 (2014).

62. Gu, X. *et al.* GC-MS metabolomics on PPARα-dependent exacerbation of colitis. *Mol. Biosyst.* **11**,

1329–1337 (2015).

63. Mana, M. D. *et al.* High-fat diet-activated fatty acid oxidation mediates intestinal stemness and

tumorigenicity. *Cell Rep.* **35**, 109212 (2021).

64. Nishida, T. *et al.* Increased arachidonic acid composition of phospholipids in colonic mucosa from

patients with active ulcerative colitis. *Gut* **28**, 1002–1007 (1987).

65. Pearl, D. S. *et al.* Altered colonic mucosal availability of n-3 and n-6 polyunsaturated fatty acids in

ulcerative colitis and the relationship to disease activity. *J. Crohns Colitis* **8**, 70–79 (2014).

66. Huo, A. & Wang, F. Biomarkers of ulcerative colitis disease activity CXCL1, CYP2R1, LPCAT1, and

NEU4 and their relationship to immune infiltrates. *Sci. Rep.* **13**, 12126 (2023).

67. Wang, L. *et al.* rSj16 Protects against DSS-Induced Colitis by Inhibiting the PPAR-α Signaling Pathway.

Theranostics **7**, 3446–3460 (2017).

68. Yao, H. *et al.* Matrine protects against DSS-induced murine colitis by improving gut barrier integrity,

inhibiting the PPAR-α signaling pathway, and modulating gut microbiota. *Int. Immunopharmacol.* **100**,

108091 (2021).

69. Miyoshi, H. & Stappenbeck, T. S. In vitro expansion and genetic modification of gastrointestinal stem

cells in spheroid culture. *Nat. Protoc.* **8**, 2471–2482 (2013).

70. VanDussen, K. L., Sonnek, N. M. & Stappenbeck, T. S. L-WRN conditioned medium for gastrointestinal

epithelial stem cell culture shows replicable batch-to-batch activity levels across multiple research

teams. *Stem Cell Res.* **37**, 101430 (2019).

71. Ludikhuize, M. C., Meerlo, M., Burgering, B. M. T. & Rodríguez Colman, M. J. Protocol to profile the

bioenergetics of organoids using Seahorse. *STAR Protoc.* **2**, 100386 (2021).

72. Dekkers, J. F. *et al.* High-resolution 3D imaging of fixed and cleared organoids. *Nat. Protoc.* **14**, 1756–

1771 (2019).

73. Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and enables interrogation of

frozen tissues. *Nat. Methods* **14**, 959–962 (2017).

1046

Acknowledgments

1047

This work was funded by the National Institutes of Health (NIH) to M.J.R. under the award number

R21DK123691 and the Stanford Medicine Children's Health Center for IBD and Celiac Disease. B.A.O was

supported by a fellowship from the Stanford Medicine Children's Health Center for IBD and Celiac. This

work utilized computing resources provided by the Stanford Genetics Bioinformatics Service Center. V.DJ.P

was supported by R01HL139664, R01HL160018, R01 HL134776, R01HL59886, 1R01HL172449-01 from

the NIH. S.D was supported by the R01DKDK099222 award from the NIH. We thank L. Prince (Pediatric

Neonatology, Stanford Medicine) for resources for flow cytometry and confocal microscopy. We

acknowledge the support of the National Center for Advancing Translational Sciences of the NIH under

Award Number UL1TR003142. The content is solely the authors' response and does not necessarily

represent the views of the funding agencies.

1058

Competing interests

1060 The authors have no competing interests to disclose.

1061

Author contributions

1063

1064

Figures

Fig. 1

1066

1067

Fig. 1: Hypermetabolism and cellular stress in active UC colonoids during differentiation

a, Representative phase contrast images of pediatric control (C) and active (a)UC colonoids after 3-day

differentiation in unstimulated conditions in ODM. Scale bar, 200 μm. **b**, The number of buds/colonoid in

colonoids treated as in **a**. Circles represent the average measures of 20-30 colonoids from individual

colonoid line. n=8 patient colonoid line/group, two-sided Mann-Whitney test. **c**, Spheroids were cultured in

OGM, and paired samples were differentiated in ODM for 3 days (colonoids) and assayed for *WFDC2*

mRNA abundance using qRT-PCR. n=3 patient line/group, 2-way ANOVA, Fisher LSD test. **d**, Phase

contrast image of a Seahorse 96-well plate with a representative spheroid line before and after 3-day

differentiation. Scale bar, 200 μm **e**, Bioenergetic profile of C and aUC colonoids differentiated for 3 days as

in **a**, and subjected to the Seahorse MitoStress test. Lines represent average measures of individual

colonoid lines. n=8 patient colonoid line/group. **f**, MitoStress OCR response of C and aUC colonoids

differentiated for 3 days as in **a**, in 96-well Seahorse plate. Circles represent the average quadruplicate

wells of individual colonoid lines. n=8 patient colonoid lines/group, Mann-Whitney test. **g**, The colonoids

cultured and used in **f**, were subjected to total ATP luminescence assay. Circles represent the average

luminescence of duplicate wells of individual colonoid lines**.** n=8 patient colonoid line/group, two-sided

Mann-Whitney test. **h**, Colonoids were treated as in **a**, and mitochondrial membrane potential (MMP) was

estimated with the JC-1 dye by flow cytometry. Circles represent the average luminescence of duplicate

wells of individual colonoid lines**.** n=7-8 patient colonoid line/group, two-sided Mann-Whitney test. **i**,

Estimation of mitochondrial mass with Mitotracker green intensity in C and aUC colonoids treated as in **a**.

Circles represent the mean fluorescence intensities (MFI) of 5000 cells from individual colonoid lines. n=8

patient spheroid line/group, two-sided Mann-Whitney test. **j**, The same set of colonoids cultured and used in

f, were used to estimate mitochondrial ROS with the MitoSOX fluorescence assay. Circles represent the

average fluorescence of duplicate wells of individual colonoid lines**.** n=8 patient colonoid line/group, two-

sided Mann-Whitney test. **k**, LDH activity in C and aUC colonoids in ODM taken from the same plate as **f**

after a 2-day differentiation. Circles represent the average duplicate measures of individual spheroid lines.

n=8 patient spheroid line/group, unpaired, two-sided t-test. *=P < 0.05, **=P < 0.01, ***=P < 0.001. LDH,

lactate dehydrogenase; OCR, oxygen consumption rates; ODM, organoid differentiation medium; OGM,

organoid growth medium; ROS, reactive oxygen species

1096

1097

Fig. 2:

Fig. 2: Distinct transcriptome in epithelial colonoids from active UC pediatric patients

C and aUC pediatric epithelial colonoids differentiated for 3 days in ODM and were subjected to bulk RNA-

seq. **a**, Volcano plot of differentially expressed genes between aUC and C samples (Log2FC>1, FDR <0.1).

The top 10 dysregulated genes are labeled on the plot. **b,** Heat map of the top 10 dysregulated genes

showing the diversity and consistency of expression among each colonoid line n=8 patient colonoid

line/group. **c**, Chemokine genes overexpressed in aUC epithelial colonoids compared to C in the bulk RNA-

seq data (Log2FC>1, FDR <0.1), n=8 patient colonoid line/group. The full list of differentially expressed

genes is provided in Supplementary Table 2. **d**, Robust overexpression of chemokine protein, CCL2, **e**,

CXCL11, and **f**, CXCL1 in aUC colonoids using ODM collected after a 3-day differentiation. **d-f**, n=4 patient

colonoid lines/group. Circles represent the average of duplicate measures of individual colonoid lines, two-

sided unpaired t-test, *=P < 0.05, **=P < 0.01, ***=P < 0.001. ODM, organoid differentiation medium

Fig. 3

1112 **Fig. 3: Higher glycolytic capacity in active UC colonoids during differentiation**

1113 **a**, Ingenuity Pathway Analysis of differentially expressed RNAs between aUC and C colonoids. The top 10

1114 Molecular and Cellular Functions are shown. Full details of molecules involved under each function are

1115 presented in Supplementary Table 3. **b**, Bioenergetic profile of C and aUC colonoids differentiated in 96-

1116 well Seahorse plates for 3 days under non-stimulated conditions in ODM and subjected to the Seahorse

1117 Glycostress assay. Lines represent the average of quadruplicate ECAR measurements from individual

1118 colonoid lines. n=6 patient colonoid line/group. **c-f**, Glycostress extracellular acidification rates (ECAR) of C

1119 and aUC colonoids as in **b**, to determine real-time **c**, glycolysis, **d**, glycolytic capacity, **e**, glycolytic reserve,

1120 and **f**, non-glycolytic acidification rate. Circles in **c-f** represent the average quadruplicate wells of individual

1121 colonoid lines. n=6 patient colonoid lines/group, two-sided unpaired t-test. **g**, Extracellular glucose and **h**,

extracellular lactate concentration in 2-day ODM from C and aUC colonoids differentiated for two days.

Circles represent the average duplicate measures of individual colonoid lines relative to measures in blank

wells with ODM only. n=8 patient colonoid line/group, unpaired, two-sided t-test. *=P < 0.05, **=P < 0.01.

ECAR, extracellular acidification rates, ODM, organoid differentiation medium.

Fig. 4

Fig. 4: Dysregulated lipid metabolism supports hypermetabolism in active UC colonoids during differentiation

a, Significant differentially expressed lipid-related genes (FDR<0.1) in aUC vs C epithelial colonoids differentiated for 3 days. Dots represent the Log2 fold change in aUC compared to C colonoids. n=8 patient colonoid lines/group. **b**, Spheroids were cultured in OGM, and paired samples were differentiated in ODM for 2 days (colonoids) and assayed for BODIPY$^+$ neutral lipid estimation by flow cytometry. Open symbols represent the mean fluorescence intensities of BODIPY$^+$ cells from individual patient spheroids, while closed symbols represent the mean fluorescence intensities of BODIPY$^+$ cells from individual colonoid lines. Data presented relative to C-spheroids. n=5 patient lines/group, 2-way ANOVA, Tukey's *post hoc* test. **c**, Representative confocal microscopy images of C and aUC colonoids treated as in **b**, for visualizing BODIPY$^+$ neutral lipid accumulation. Scale bar, 50 µm . **d**, LPL protein activity in C and aUC colonoids after a 2-day and 5-day differentiation in ODM. Black and Red circles represent the average duplicate measures of individual colonoid lines. n=5 patient colonoid lines/group. Two-way ANOVA, Fisher's LSD test. **e**, Schematic of C and aUC colonoids differentiated for 3 days in ODM subjected to the Seahorse MitoStress test in a nutrient-deprived medium with or without etomoxir to inhibit fatty acid oxidation. **f**, Bioenergetic profile of C and aUC colonoids as in **e**. Lines represent the average of quadruplicate OCR measures from individual colonoid lines. n=3 patient colonoid lines/group. **g**, Basal OCR, **h**, proton leak, and **i**, ATP-linked OCR response of C and aUC colonoids as in **e**. Symbols represent the average triplicate wells of individual colonoid lines. n=3 patient colonoid lines/group. 2-way ANOVA, Tukey's *post hoc* test. **j**, Representative phase contrast images of colonoids exposed to 200 µM BSA-PA or BSA-EtOH control for the first 3 days before further differentiation up to 5 days. Scale bar, 200 µm. **m**, Gene expression of chemokines in colonoids treated as in **j**, assessed by qRT-PCR. Circles represent individual colonoid lines treated with BSA-PA relative to those treated with BSA-EtOH**.** n=4 patient colonoid lines/group. Two-sided paired t-test. *=P < 0.05, **=P < 0.01, ***=P < 0.001. ****=P < 0.0001 BSA, bovine serum albumin; OCR, oxygen consumption rates; ODM, organoid differentiation medium; PA, Palmitic acid

Fig. 5

Fig. 5: PPAR-α agonism in non-IBD colonoids induces hypermetabolic stress during differentiation

a, Ingenuity Pathway Analysis of bulk transcriptomic data showing the top 10 predicted Causal Networks of differential gene expression between aUC and C colonoids. The full details of molecules involved under each master network are presented in Supplementary Table 4. **b**, Schematic of PPARA agonist, fenofibrate, treatment in non-IBD colonoids during differentiation. **c**, Representative phase contrast images of pediatric C colonoids treated as in **c**. Scale bar, 200 μm. **d**, The number of buds/colonoid in colonoids treated as in **b**, and differentiated for 3 days. Circles represent the average measures of individual colonoid lines**.** n=8 patient colonoid lines/group, two-sided Wilcoxon signed-rank test. **e**, WFDC2 gene expression in colonoids treated as in **b**, assessed by qRT-PCR. Circles represent the average duplicate measures of individual colonoid lines**.** n=3 patient colonoid lines/group, two-sided paired t-test. **f**, Extracellular glucose and **g**, extracellular lactate concentration in 2-day ODM from C and aUC colonoids. Circles represent the average duplicate measures of individual colonoid lines relative to measures in blank wells with ODM only. n=8 patient colonoid line/group, two-sided Wilcoxon signed-rank test. **h**, Representative confocal immunofluorescence images of UCP2 expression in non-IBD colonoids treated as in b, for 3 days. Scale bar, 50 μm **i**, Colonoids treated as in **b**, differentiated for 2 days and assayed for BODIPY[+] neutral lipid estimation by flow cytometry. Open symbols represent the mean fluorescence intensities (MFI) of BODIPY[+] cells from individual patient colonoids, Data presented relative to the MFI of vehicle. n=5 patient lines/group, two-sided paired t-test. **j**, Representative confocal images of C colonoids treated as in **i**, for visualizing BODIPY[+] neutral lipid accumulation. Scale bar, 50 μm. **k**, LPL activity in C colonoids treated as in **b**, and differentiated for 2 days. Circles represent the average duplicate measures of individual colonoid lines. n=6 patient colonoid lines/group, two-sided, paired t-test. **l**, Estimation of mitochondrial mass with Mitotracker green intensity in C colonoids treated as in **b**, and differentiated in ODM for 3 days. Circles represent the MFI of 5000 cells from individual colonoid lines relative to vehicle. n=8 patient spheroid line/group, two-sided paired t-test. **m**, C colonoids treated as in **b**, and differentiated for 3 days were used to estimate ROS with the MitoSOX fluorescence assay. Circles represent the average fluorescence of duplicate wells of individual colonoid lines**.** n=8 patient colonoid line/group, two-sided Wilcoxon signed-rank test. **n**, Bioenergetic profile of C colonoids treated as in **b**, and subjected to the Seahorse MitoStress test after a 3-day differentiation in ODM. Lines represent average OCR measures of individual colonoid lines. n=8 patient

colonoid lines/group. **o**, MitoStress OCR responses of C colonoids in **n**, Circles represent the average

quadruplicate wells of individual colonoid lines. n=8 patient colonoid line/group, two-sided paired t-test. **p**,

LDH activity in ODM sampled from the same plate as **o** after a 2-day differentiation in ODM. Circles

represent the average duplicate measures of individual colonoid lines. n=8 patient colonoid lines/group, two-

sided Wilcoxon signed-rank test. n=8 colonoid lines/group. *=$P < 0.05$, **=$P < 0.01$, ***=$P < 0.001$. ****=$P <$

0.0001 OCR, oxygen consumption rates; ODM, organoid differentiation medium

1190

Fig. 6

Fig. 6: PPAR-α inhibition reprograms epithelial metabolism and ameliorates hypermetabolic stress in active UC colonoids during differentiation

aUC colonoids were treated with or without the PPAR-α antagonist, GW6471 (GW), during differentiation in ODM. **a**, GW effect on BODIPY$^+$ neutral lipid estimation by flow cytometry in aUC colonoids differentiated in ODM for 2 days. Open symbols represent the mean fluorescence intensities (MFI) of BODIPY$^+$ cells from individual patient colonoids. Data presented relative to the MFI of aUC controls. The dotted line indicates MFI of no stain control. n=5 patient lines/group, two-sided paired t-test. **b**, Representative confocal images of aUC colonoids treated as in **a**, for visualizing BODIPY$^+$ neutral lipid accumulation. Scale bar, 50 μm. **c**, Effect of GW on UCP2 gene expression by qRT-PCR. Open symbols represent the expression of individual colonoid lines. n=6 colonoid lines/group, two-sided paired t-test. **d**, Bioenergetic profile of aUC colonoids treated with or without GW and subjected to the Seahorse MitoStress test after a 3-day differentiation in ODM. Lines represent average OCR measures of individual colonoid lines. n=6 patient colonoid lines/group. **e**, MitoStress OCR responses of aUC colonoids treated as in **d**, Open circles represent the average quadruplicate wells of individual colonoid lines. n=6 patient colonoid line/group, two-sided paired t-test. The same colonoids cultured and used in **d**,**e**, were subjected to total ATP luminescence assay. Circles represent the average luminescence of duplicate wells of individual colonoid lines**.** n=6 patient colonoid line/group, two-sided Mann-Whitney test. **g**, Bioenergetic profile of aUC colonoids differentiated for 3 days in ODM with or without GW and subjected to the Seahorse MitoStress test in a nutrient-deprived medium with or without etomoxir. **h**, Basal OCR, aUC colonoids as in **g.** Symbols represent the average OCR from triplicate wells of individual colonoid lines. n=3 patient colonoid lines/group. 2-way ANOVA, Tukey's *post hoc* test. **i**, Extracellular glucose and **j**, lactate concentration in 2-day ODM from aUC colonoids with or without GW sampled from the same plate as **e**. Open circles represent the average duplicate measures of individual colonoid lines relative to measures in blank wells with ODM only. n=6 patient colonoid lines/group, two-sided Wilcoxon signed-rank test. **k**, LDH activity in ODM sampled from the same plate as **e** after a 2-day differentiation in ODM. Open circles represent the average duplicate measures of individual colonoid lines. n=6 patient colonoid lines/group, two-sided Wilcoxon signed-rank test. n=6 colonoid lines/group. **l,** aUC colonoids with or without GW were differentiated in ODM for 5 days. ODM collected on day 2 and on day 5 (representing the last 3 days of differentiation) was used to assess CXCL1 secretion. Open circles

represent the average duplicate measures of individual colonoid lines. n=6 patient colonoid lines/group, two-

sided paired t-test. **m**, Representative phase contrast images of aUC colonoids differentiated in ODM with

or without GW for 5 days. Scale bar, 200 μm.   **n**. The number of buds/colonoid in aUC colonoids treated as

in **m**. Open circles represent the average measures of individual colonoid lines. n=6 patient colonoid

lines/group, two-sided Wilcoxon signed-rank test. **o**, Effect of GW on gene expression of differentiation

markers assessed by qRT-PCR. Open symbols represent the expression of individual colonoid lines. n=6

colonoid lines/group, two-sided paired t-test. Dotted vertical lines represent the average of at least 2 non-

IBD control colonoids. $*=P < 0.05$, $**=P < 0.01$, $***=P < 0.001$. $****=P < 0.0001$ OCR, oxygen consumption

rates; ODM, organoid differentiation medium

1229

1230

1231 **Extended data**

1232

1233

Extended Data Fig. 1: Colon Epithelial Spheroid Generation

a, Schematic of epithelial spheroids derivation and maintenance from pediatric rectal biopsies.

Demographics of pediatric patients are provided in Supplementary Table 1. **b**, Phase-contrast images of C

and aUC spheroids after the first 7 days of biopsy derivation in OGM show that the development of aUC

spheroids from biopsies is slower than the C spheroids. Scale bar, 200 µm. **c**, Phase contrast images of C

and aUC spheroids in culture after 3 passages, showing that after initial slow growth rates, aUC spheroids

become more stable like C spheroids. Scale bar, 200 µm. **d**, High magnification images of C and aUC

spheroids, showing expected morphology with large lumens after freeze-thaw protocol. Scale bar, 50 µm.

OGM, Organoid Growth Medium; ODM, Organoid Differentiation Medium

1243

Extended Data Fig. 2

1244

Extended Data Fig. 2: Pediatric colon spheroids exhibit similar morphological and metabolic profiles

a, Phase contrast images of undifferentiated C and aUC spheroids filtered after trypsinization, passed

through a strainer, and monitored for 7 days. Scale bar, 200 µm. **b**, Diameter of C and aUC spheroids

described in **a.** Blinded measures of the diameter of all spheroids on a high-powered field were taken. Black

and Red circles represent the average measures of individual spheroids within a high-powered field of each

spheroid line**.** n=8 patient spheroid line/group, two-sided Mann-Whitney test. **c**, Bioenergetic profile of C and

aUC spheroids from Seahorse MitoStress test. n=8 patient spheroid line/group. Black and Red lines

represent average OCR measures of individual spheroid lines. **d**, OCR response of C and aUC spheroids

cultured in unstimulated conditions in OGM. Black and Red circles represent the average quadruplicate

wells of individual spheroid lines. n=8 patient spheroid line/group, unpaired, two-sided t-test. **e**, Estimation of

mitochondrial mass with Mitotracker green intensity in undifferentiated C and aUC spheroids. Black and

Red circles represent the mean fluorescence intensities (MFI) of 5000 cells from individual spheroid lines.

n=8 patient spheroid line/group, two-sided Mann-Whitney test. **f**, LDH activity in C and aUC spheroids in

OGM from the same plate as **c**, **d**. Black and Red circles represent the average duplicate measures of

individual spheroid lines. n=8 patient spheroid line/group, unpaired, two-sided t-test. **g**, Phase contrast

images of C and aUC differentiated spheroids (colonoids) on day 7 of differentiation. Scale bar, 200 µm. **h**,

Spheroids embedded in Matrigel were cultured in OGM and paired samples were differentiated in ODM for

3 days (colonoids) and assay for **h**, *ATOH1* and **i**, *CHGA* mRNA abundance using qRT-PCR. n=3 patient

line/group, 2-way ANOVA. *=$P < 0.05$, **=$P < 0.01$. OCR, oxygen consumption rates; ODM, organoid

differentiation medium; OGM, organoid growth medium

1265

1266

Extended Data Fig. 3

Extended Data Fig. 3: Colonoids from patients with inactive UC exhibit increased metabolic

susceptibility to inflammatory stimulus.

a, Phase contrast images of pediatric C and iUC colonoids after 3-day differentiation in unstimulated

conditions in ODM. Scale bar, 200 µm. **b**, The number of buds/colonoid in colonoids treated as in **a**. Black

and purple circles represent the average measures of individual colonoid lines**.** n=8 patient colonoid

line/group, two-sided Mann-Whitney test. **c**, Bioenergetic profile of C and iUC colonoids differentiated for 3

days as in **a**, and subjected to the Seahorse MitoStress test. Black and Purple lines represent average OCR

measures of individual colonoid lines. n=8 patient colonoid line/group. **d**, MitoStress OCR response of C

and iUC colonoids differentiated for 3 days as in **a**, in 96-well Seahorse plate. Black and Purple circles

represent the average quadruplicate wells of individual colonoid lines. n=8 patient colonoid line/group,

Mann-Whitney test. **e**, The same cohort of colonoids cultured and used in **d**, were used to estimate

mitochondrial ROS with the MitoSOX fluorescence assay. Black and Purple circles represent the average

fluorescence of duplicate wells of individual colonoid lines**.** n=8 patient colonoid line/group, two-sided Mann-

Whitney test. **f**, C, and iUC spheroids were exposed to an inflammatory cocktail (40 ng/mL TNFa, 20 ng/mL

IL-1b, and 500 ng/mL Flagellin) in OGM for 16 hrs before differentiation in ODM for 3 days. **g**, Phase

contrast images of C and iUC colonoids treated as in **f**. Scale bar, 200 μm. **h**, Estimation of mitochondrial

mass with Mitotracker green intensity in C and iUC colonoids treated as in **f**. Black and purple circles

represent the mean fluorescence intensities (MFI) of 5000 cells from individual spheroid lines. n=4 patient

colonoid line/group, 2-way ANOVA, with Tukey correction. **i**, Bioenergetic profile of C and iUC colonoids

treated as in **f**, and subjected to the Seahorse MitoStress test. Black and Purple lines represent average

measures of individual colonoid lines. n=4 patient colonoid line/group. **j**, MitoStress OCR response of C and

iUC colonoids treated as in **f**. Black and Purple circles represent the average quadruplicate wells of

individual colonoid lines. n=7 patient colonoid line/group, 2-way ANOVA with Tukey correction. *=$P < 0.05$,

**=$P < 0.01$, ***=$P < 0.001$. OCR, oxygen consumption rates; ODM, organoid differentiation medium; OGM,

organoid growth medium; ROS, reactive oxygen species

1293

Extended Data Fig. 4

1295 **Extended Data Fig. 4: Metabolic uncoupling contributes to hypermetabolic phenotypes during**

1296 **colonoid differentiation**

1297 **a**, Rectal gene expression of uncoupling proteins in bulk RNA-Seq data from non-IBD control participants in

1298 the NIH PROTECT study showing dominant UCP2 gene expression over other uncouplers (transcripts per

1299 million (TPM) values). n=20. **b**, Rectal UCP2 gene expression (TPM) from treatment-naïve participants with

UC in the RISK cohort. n=43-55/group. Two-tailed Mann-Whitney test. **c**, Rectal UCP2 gene expression

(TPM) from the NIH PROTECT study of treatment-naive pediatric UC patients. n=20(C) -206(aUC) /group.

Two-tailed Mann-Whitney test. **d**, (i) UCP2 gene expression assessed by qRT-PCR and (ii) UCP2 protein

expression in pediatric C and aUC colonoids differentiated for 3 days in ODM. n=4 patient colonoid

line/group i) two-tailed unpaired t-test. ii) two-sided Mann-Whitney test. **e,** C and aUC colonoids were

embedded in OCT after 3-day differentiation. Representative immunofluorescence images showing UCP2

expression and merged expression with the mitochondrial marker, COX4. Semi-quantitation of

COX4+UCP2+ cell counts is presented in d(ii). Scale bar, 50 μm. **f**, Morphological and metabolic effects of

mitochondrial uncoupling during colonoid differentiation were tested in C colonoids treated with or without

the mitochondrial protonophore, FCCP. **g**, Bioenergetic profile of C colonoids treated as in **f**, and subjected

to the Seahorse MitoStress test. Black and Red lines represent average OCR measures of individual

colonoid lines. n=8 patient colonoid lines/group. **h**, MitoStress OCR response of C colonoids treated as in **f**,

in 96-well Seahorse plate. Black and Red circles represent the average quadruplicate wells of individual

colonoid lines. n=8 patient colonoid line/group, two-sided paired t-test. **i**, Estimation of mitochondrial mass

with Mitotracker green intensity in C colonoids treated as in **f**. Black and Red circles represent the mean

fluorescence intensities (MFI) of 5000 cells from individual colonoid lines. n=8 patient spheroid line/group,

two-sided paired t-test. **j**, The same set of colonoids cultured and used in **f**, were used to estimate

mitochondrial ROS with the MitoSOX fluorescence assay. Black and Red circles represent the average

fluorescence of duplicate wells of individual colonoid lines**.** n=8 patient colonoid line/group, two-sided

Wilcoxon signed-rank test. **k**, LDH activity in ODM taken from the same plate as **h** after a 2-day

differentiation. Black and Red circles represent the average duplicate measures of individual colonoid lines.

n=8 patient colonoid lines/group, two-sided Wilcoxon signed-rank test. **l**, Representative phase contrast

images of pediatric C colonoids treated as in **f**. Scale bar, 200 μm. **m**, The number of buds/colonoid in

colonoids treated as in **f**. Black and Red circles represent the average measures of individual colonoid lines**.**

n=8 patient colonoid line/group, two-sided Wilcoxon signed-rank test. **n**, WFDC2 gene expression in

colonoids treated as in **f**, assessed by qRT-PCR. Black and Red circles represent the average measures of

individual colonoid lines**.** n=4 patient colonoid lines/group, paired t-test. **o**, Gene expression of chemokines

in colonoids treated as in **f**, assessed by qRT-PCR. Black and Red circles represent the average measures

1328 of individual colonoid lines**.** n=4 patient colonoid line/group, paired t-test. **p**, Colonoids were treated as in **f**,

1329 and Chemokine protein concentrations were assessed in ODM after a 2-day differentiation. Black and Red

1330 circles represent the average duplicate measures of individual colonoid lines. n=6 colonoid lines/group,

1331 paired t-test. *=$P < 0.05$, **=$P < 0.01$, ***=$P < 0.001$. LDH, lactate dehydrogenase; OCR, oxygen

1332 consumption rates; ODM, organoid differentiation medium; ROS, reactive oxygen species

1333

Extended Data Fig. 5

1334

Extended Data Fig. 5: Dysregulated lipid metabolism genes in the colon epithelia of active UC

a, Rectal gene expression (TPM) of lipid-related colonoid genes (as in Fig. 4**a**) from the NIH PROTECT

study[19] of treatment-naive pediatric UC patients. n=20(C) -206(aUC)/group. Kruskal-Wallis test with Dunn's

multiple comparisons. Black and Red symbols represent the TPM value of each patient's data. **b**,

Expression of colonoid lipid-related genes (as in Fig. 4**a**) by disease state in publicly available adult single-

cell (sc) RNA-seq data[39]. Data is colored by relative gene expression and the relative size of each dot

shows the percentage of cells expressing each marker per disease state. **c**, Expression of colonoid lipid-

related genes (as in Fig. 4**a**) in scRNA-seq data[39] by epithelial cluster. Data is colored by relative gene

expression and the relative size of each dot shows the percentage of cells expressing each marker per

epithelial cluster.*=$P < 0.05$, **=$P < 0.01$, ***=$P < 0.001$.

1345

1346

Extended Data Fig. 6

1347

Extended Data Fig. 6: Lipotoxicity modulates UCP2 and WFDC2 gene expression during

differentiation

a, Maximal respiration and, **b**, Non-mitochondrial respiration of C, and aUC colonoids differentiated for 3

days in ODM and subjected to the Seahorse MitoStress test in a nutrient-deprived medium with or without

Etomoxir. Black and Red symbols represent the average OCR from triplicate wells of individual colonoid

lines n=3 patient colonoid lines/group. 2-way ANOVA, Tukey's *post hoc* test. **c**, qRT-PCR gene expression

data of non-IBD colonoids differentiated in ODM for 24 hrs with the stated doses of BSA-PA. Symbols

represent 3 technical replicates in independent matrigel domes. One-way ANOVA and Dunnett tests. *=*P* <

0.05, **=*P* < 0.01. BSA, bovine serum albumin; OCR, oxygen consumption rates; ODM, organoid

differentiation medium; PA, Palmitic acid

Extended Data Fig. 7

Extended Data Fig. 7: Epithelial PPARA expression in active UC

a, Colon scRNA-seq data[39] showing the expression of PPARA and RXRA by disease state in colon epithelial

lineages. Data is colored by relative gene expression and the relative size of each dot shows the

percentage of cells expressing each marker per disease state. **b**, PPAR-α activity assay in nuclear extracts

of pooled C and aUC colonoids (n=3/group) differentiated for 24 hrs. Black and red symbols represent four

technical replicates from pooled nuclear extracts and are presented relative to C. Two-sided unpaired t-test.

*=$P < 0.05$.

1367

Extended Data Fig. 8

1368

Extended Data Fig. 8: Blockade of lipid synthesis suppresses hypermetabolic features in active UC colonoids

aUC colonoids were treated with or without the PPAR-α antagonist, GW6471 (GW), during differentiation in ODM. **a**, Effect of GW on *LPL* gene expression by qRT-PCR. Red and Blue open circles represent the expression of individual colonoid lines. n=6 colonoid lines/group, two-sided paired t-test. **b**, Maximal, **c**, Proton leak **d**, ATP-linked, and **e**, Non-mitochondrial OCR response of aUC colonoids differentiated for 3 days in ODM with or without GW and subjected to the Seahorse MitoStress test in a nutrient-deprived medium with or without Etomoxir. Red and Blue symbols represent the average OCR from triplicate wells of individual colonoid lines. n=3 patient colonoid lines/group. 2-way ANOVA, Tukey's *post hoc* test. **f**, Estimation of mitochondrial mass with Mitotracker green intensity in aUC colonoids treated with or without GW in ODM for 3 days. Red and Blue open circles represent the MFI of 5000 cells from individual colonoid lines relative to aUC. n=6 patient spheroid line/group, two-sided Wilcoxon signed-rank test. **g**, ROS estimate with the MitoSOX fluorescence assay in a UC colonoids differentiated for 3 days with or without GW. Red and Blue open circles represent the average fluorescence of duplicate wells of individual colonoid lines**.** n=6

patient colonoid line/group, two-sided Wilcoxon signed-rank test. **h,** aUC colonoids with or without GW were

differentiated in ODM for 5 days. ODM collected on day 2 and on day 5 (representing the last 3 days of

differentiation) was used to assess CXCL11 and **i**, CCL2 secretion. Red and Blue open circles represent the

average duplicate measures of individual colonoid lines. n=6 patient colonoid lines/group, two-sided paired

t-test. *=$P < 0.05$, **=$P < 0.01$, ***=$P < 0.001$. ****=$P < 0.0001$ OCR, oxygen consumption rates; ODM,

organoid differentiation medium; MFI, mean fluorescence intensity

1389

1390

1391   **Supplementary information**

1392

1393   Supplementary Table 1: Patient demographics *(Microsoft Word)*

1394

1395   Supplementary Table 2: Colonoid Bulk RNA-seq. *(Microsoft Excel file)*

1396

1397   Supplementary Table 3: Ingenuity Pathway Analysis: Molecular and Cellular Functions *(Microsoft Excel file)*

1398

1399   Supplementary Table 4: Ingenuity Pathway Analysis: Causal Networks *(Microsoft Excel file)*

1400

1401   Supplementary Table 5: Taqman Probes for qRT-PCR *(Microsoft Word file)*