The oncometabolite D-2-hydroxyglutarate promotes DNA hypermethylation at lineage-specific enhancers controlling microglial activation in IDH[mut] gliomas

Alice Laurenge[1,2*], Pietro Pugliese[1*], Quentin Richard[1], Bertrand Mathon[3], Stéphanie Jouannet[1], Yvette Hayat[1], Sarah Scuderi[1], Laurent Capelle[3Ŧ], Pauline Marijon[3], Karim Labreche[1], Agustí Alentorn[1,2], Maïté Verreault[1], Ahmed Idbaih[1,2], Cristina Birzu[1,2], Emmanuelle Huillard[1], Nina Pottier[1], Aurore Desmons[4], Inès Fayache[4], Garrett A. Kaas[5], Philip J. Kingsley[6], Lawrence J. Marnett[7], Eric Duplus[8], Elias El-Habr[9], Lucas A. Salas[10], Karima Mokhtari[1,11], Suzanne Tran[1,11], Mehdi Touat[1,2], Franck Bielle[1,11], Mario L. Suvà[12,13], Antonio Iavarone[14], Michele Ceccarelli[15], Michel Mallat[1], Marc Sanson[1,2,16], Luis Jaime Castro-Vega[1]

[1]Sorbonne Université UMRS1127, CNRS UMR7225, Inserm U1127, ICM - Paris Brain Institute, F-75013 Paris, France; Équipe Labellisée Par La Ligue Nationale Contre Le Cancer.
[2]Service de Neuro-Oncologie, AP-HP, Hôpital Pitié-Salpêtrière, F-75013 Paris, France.
[3]Department of Neurosurgery, AP-HP, Hôpital Pitié Salpêtrière, F-75013 Paris, France.
[4]Clinical Metabolomic Department, AP-HP, Hôpital Saint Antoine, Sorbonne Université, Paris, France.
[5]Division of Genetic Medicine, Vanderbilt University Medical Center, Nashville, TN, USA.
[6]Department of Biochemistry, Vanderbilt University, Nashville, TN, USA.
[7]Departments of Biochemistry, Chemistry, and Pharmacology, Vanderbilt Institute of Chemical Biology, Vanderbilt-Ingram Cancer Center, Vanderbilt University School of Medicine, Nashville, TN, USA.
[8]Sorbonne Université, IBPS, CNRS UMR8256, INSERM ERL1114, Biological Adaptation and Aging, Paris, France.
[9]Sorbonne Université, IBPS, CNRS UMR8246, Inserm U1130, Neuroscience Paris Seine Laboratory, Paris, France.
[10]Geisel School of Medicine at Dartmouth, Dartmouth Cancer Center, Lebanon, NH, 03756, USA.
[11]Department of Neuropathology, AP-HP, Hôpital Pitié Salpêtrière, F-75013 Paris, France.
[12]Department of Pathology and Krantz Family Center for Cancer Research, Massachusetts General Hospital and Harvard Medical School, Boston, MA, 02114, USA.
[13]Broad Institute of Harvard and MIT, Cambridge, MA 02142, USA.
[14]Department of Neurological Surgery and Department of Biochemistry, Sylvester Comprehensive Cancer Center, University of Miami Miller School of Medicine, Miami, FL, USA.
[15]Department of Public Health Sciences and Sylvester Comprehensive Cancer Center, University of Miami Miller School of Medicine, Miami, FL, USA.
[16]Onconeurotek Tumor Bank, AP-HP, Hôpital Pitié-Salpêtrière, F-75013 Paris, France.

*These authors contributed equally to this work
†Dr. Laurent Capelle passed away in June 2023. This work is dedicated to his memory.

Corresponding authors:

Marc Sanson, M.D., Ph.D.
Email address: marc.sanson@aphp.fr
Genetics & Development of Brain Tumors
ICM - Paris Brain Institute, 75013, Paris, France

Luis J. Castro-Vega, M.D., Ph.D.
Email address: luis.castrovega@icm-institute.org
Genetics & Development of Brain Tumors
ICM - Paris Brain Institute, 75013, Paris, France

Summary

Tumor-associated macrophages and microglia (TAMs) are highly abundant myeloid cells in gliomas, with their phenotype and immune response determined by ontogeny and microenvironment. TAMs display distinctive transcriptional programs according to the IDH mutation status but the underlying regulatory mechanisms remain largely unknown. Herein, we uncover that CD11B+ myeloid cells in human IDH^{mut} gliomas exhibit DNA hypermethylation predominantly at distal enhancers. This hypermethylation was linked to decreased expression of genes involved in inflammatory responses and glycolytic metabolism, and the inactivation of transcription factors that regulate microglial responses to environmental stimuli. Prolonged exposure of human primary microglia to D-2-hydroxyglutarate (D-2HG) inhibited TET-mediated 5mC oxidation, resulting in a reduced accumulation of global 5hmC levels. We confirmed high 5mC/5hmC ratios at lineage-specific enhancers, by analyzing CpGs at single-base resolution. D-2HG-treated microglia show reduced proinflammatory capacity and enhanced oxidative phosphorylation, consistent with the remodeled enhancer landscape. Conversely, depletion of D-2HG following treatment of a glioma patient with an IDH mutant inhibitor was associated with enhanced microglial responses, as assessed by snRNA-seq. Our findings provide a mechanistic rationale for the hyporesponsive state of microglia in IDH^{mut} gliomas and support the concept that oncometabolites may disrupt the function of immune cells residing in the tumor microenvironment.

Keywords: D-2HG, 5mC, 5hmC, Glioma, TAMs, IDH mutation, Tumor Microenvironment

Introduction

Diffuse gliomas are the most frequent malignant primary brain tumors and are refractory to standard therapies[1]. A high proportion of these tumors bear an IDH mutation, which confers a neomorphic ability to convert α-ketoglutarate (α-KG) to (D)-2-hydroxyglutarate (D-2HG)[2]. It is believed that D-2HG drives cellular transformation by inhibiting α-KG-dependent dioxygenases, including the TET family of DNA demethylases, KDMs histone demethylases, and FTO/ALKBH RNA demethylases, ultimately leading to widespread epigenomic dysregulations and blocking of cell differentiation[3]. Much less is known about the tumor cell-extrinsic effects of this oncometabolite, which accumulates in the tumor microenvironment (TME)[4,5]. Emerging evidence indicates that D-2HG released by IDH[mut] cells suppresses T cells and impairs the metabolism of infiltrating macrophages, dendritic cells, and CD8+ T cells by non-epigenetic mechanisms[6-10]. It remains to be determined whether D-2HG is able to reshape the epigenetic landscape of the TME, particularly in resident microglia, which are abundant in IDH[mut] gliomas and are exposed to this oncometabolite from tumor initiation.

Tumor-associated macrophages and microglia (TAMs) represent almost 40% of the tumor mass and play critical roles in tumor progression[11]. As such, these cells may represent prognostic biomarkers and therapeutic targets[12,13]. TAMs are plastic cells, with their phenotype and immune response determined by both ontogeny and microenvironment[14,15]. Previous studies in IDH[wt] glioma models uncovered that resident microglia and monocyte-derived macrophages (MDMs) display distinctive transcriptional and chromatin landscapes[16]. Transcriptional differences have also been observed in the myeloid compartment of gliomas as a function of the IDH mutation status[17,18]. In IDH[wt] tumors, MDMs and microglial cells exhibit expression changes associated with activation, including high expression of MHC class genes[17]. On the other hand, in IDH[mut] gliomas, these cells display a hyporeactive phenotype similar to that of homeostatic microglia in non-tumor tissues[17].

The relative proportion of MDMs and microglial cells differs according to the IDH[mut] status[17-21] and/or tumor grade[22,23]. Glioma progression entails a process of cellular competition, wherein infiltrating macrophages overtake the niche occupied by resident microglia[24,25]. Recent single-cell RNA-seq studies of the myeloid compartment in human gliomas have further unveiled subsets of MDMs with mesenchymal-like, lipid, and hypoxic/glycolytic signatures, as well as subsets of inflammatory and oxidative phosphorylation (OXPHOS)-related microglia[20-22,24,26], some of which have been associated with clinical outcomes. Additionally, it has been reported that myeloid cells express superimposed immunomodulatory activity programs, irrespective of the ontological origin[23]. The epigenetic changes, i.e., DNA methylation, that drive the transcriptional programs and cell states of TAMs in human glioma, and the effects of tumor-derived signals on the TAM methylome, remain largely uncharacterized.

Inhibition of TET-mediated demethylation by D-2HG in tumor cells accounts for DNA hypermethylation characteristic of IDH[mut] gliomas[27]. Considering the elevated concentrations of this oncometabolite in the TME and its uptake by myeloid cells[9,28], we hypothesized that such a mechanism could be also at play in CD11B+ myeloid cells of these tumors, thus contributing to both epigenetic intratumoral heterogeneity and altered immune functions. Herein, to investigate whether and how TAMs in IDH[mut] gliomas display functional DNA methylation changes, we first profiled the bulk transcriptome and methylome of CD11B+ myeloid cells isolated from a well-characterized series of human gliomas and non-tumor tissues. Our findings reveal that myeloid cells in IDH[mut] tumors exhibit DNA hypermethylation at enhancer elements that regulate gene networks involved in microglia activation. We demonstrate the direct effects of D-2HG on TET-mediated DNA demethylation at lineage-specific enhancers in primary cultures of human microglia, thereby providing a mechanistic foundation for the hyporesponsive behavior of these cells in IDH[mut] gliomas. These results have implications for therapeutics aimed at inhibiting the IDH mutant enzyme or repolarizing TAMs.

Results

Study design

To determine the mechanistic underpinnings of TAM phenotype linked to the IDH status, we analyzed bulk DNA methylome and/or transcriptome of CD11B+ myeloid cells isolated by magnetic sorting from adult-type human diffuse gliomas (n=36) comprising IDH[wt], IDH[mut] oligodendrogliomas and IDH[mut] astrocytomas. In addition, we analyzed available methylome data of CD11B+ cells from normal brains, isolated using a similar method and profiled with the same technology[29] (Fig. 1a). To adjust for potential confounding variables, samples were matched by age and sex while being fully characterized for their clinical, pathological and genetic features (Supplementary Table 1). The contamination of CD11B+ fractions by tumor cells was assessed by analysis of IDH1 R132H or TERT C228T/C250T promoter mutations through droplet digital PCR (ddPCR) and further verified by RNA-seq-based analysis of IDH variant allelic frequency (Extended Data Fig. 1a and 1b). The estimated purity of the CD11B+ samples included in downstream analyses reached ~95%. This analysis rules out the possibility that TAMs from our study cohort harbored the IDH1 R132H mutation to a significant extent, as previously reported[30].

CD11B+ myeloid cells in human IDH[mut] gliomas display global DNA hypermethylation

To confirm the DNA methylation status of the cohort in relation to the IDH mutation, we first profiled glioma cell-enriched CD11B- fractions isolated from both IDH[wt] and IDH[mut] tumors (n=17, Supplementary Table 1) using MethylationEPIC (EPIC) arrays. Analysis of methylation levels across all probes targeting CpG sites confirmed that IDH[mut] tumors display global hypermethylation, yet with a high degree of heterogeneity as previously reported[31-33] (Extended Data Fig. 2a and 2b). In total, there were 49,676 differentially methylated CpG sites (absolute $\Delta\beta > 0.2$, FDR < 0.05), of which 92.5% were hypermethylated in IDH[mut] compared with IDH[wt] glioma samples (Extended Data Fig. 2c). The distribution per functional genomic region revealed a high proportion of hypermethylated CpGs located in regulatory regions. These regions encompass promoters, defined as constant genomic regions within a distance of -1.5 kb to +0.5 kb respect to transcription start site (TSS), and enhancers,

defined as distal regulatory elements associated with FANTOM robust promoters within a range of up to 500 kb (FDR < 1e-5; Pearson's correlation), as outlined by the FANTOM5 consortium[34]. Our analysis confirms that enhancers are particularly sensitive to hypermethylation in IDH[mut] glioma cells[35] (Extended Data Fig. 2c and 2d).

We then looked for methylome changes in the fraction of CD11B+ myeloid cells isolated from gliomas (n=26) and from normal brain tissues (n=6) (Supplementary Table 1). Principal component analysis (PCA) revealed a distinct segregation of myeloid cells based on both tissue origin and IDH mutation status in component 1, which accounted for 48% of the variance (Fig. 1b). Despite the substantial heterogeneity of methylome data in CD11B+ samples from IDH[mut] tumors, we found 37,198 differentially methylated CpG sites (absolute $\Delta\beta$ > 0.2, FDR < 0.05) when compared with CD11B+ cells from IDH[wt] gliomas (Fig. 1c). Strikingly, similar to what was observed in glioma cells, CD11B+ samples from corresponding IDH[mut] tumors exhibited a bias towards hypermethylation, particularly in regulatory regions (Fig. 1c). It is known that IDH[wt] glioblastomas are enriched in MDMs, whereas the TME of IDH[mut] tumors is dominated by tissue-resident microglia[17,18]. However, the asymmetric distribution of the methylation changes we observed indicates that this was not simply an effect of differences in cell composition. Indeed, global hypermethylation was also observed when comparing the DNA methylome of CD11B+ cells isolated from IDH[mut] tumors with CD11B+ cells isolated from normal brain tissues, both predominantly composed of microglial cells (Fig. 1c). While in this comparison the number of differentially methylated CpG sites and genomic region was higher, there is a notable overlap between regulatory regions that exhibited hypermethylation in both comparisons (Fig. 1d). Of these, the number and methylation level of affected enhancers, along with their overlap between the two comparisons, were greater than those observed for promoters (Fig. 1d). Functional enrichment analyses of genes regulated by hypermethylated regulatory regions intersected in the two comparisons indicated roles in inflammatory responses such as TNF-α, IFN-γ and IL2/IL6 signaling, as well as in complement and hypoxia pathways (Fig. 1e and Supplementary Table 2).

Figure 1. DNA methylome profiling of CD11B+ cells underscores hypermethylation in the myeloid compartment of IDH^mut gliomas. (A) Overview of the *ex vivo* study workflow. Tumor samples (n=36) underwent enzymatic dissociation, followed by CD11B-based magnetic sorting. DNA and/or RNA extraction was performed on both CD11B+ and CD11B- fractions for downstream analyses. Available methylome data of CD11B+ cells from six normal brains were included as controls for comparisons. Clinical, pathological, and genetic annotations of the study cohort are detailed in Supplementary Table 1. **(B)** Principal component analysis performed using methylome data from around 600,000 CpG sites across the genome in CD11B+ samples from the study cohort. **(C)** Stacked bar charts (left) show absolute numbers and relative percentages of CpG sites and of four genomic regions with respect to the methylation status (absolute Δβ > 0.2, FDR < 0.05) for the indicated comparisons. Volcano plots (right), colored by methylation status, depict the magnitude and extent of differentially methylated enhancers and promoters for the indicated comparisons. **(D)** Venn diagrams illustrate significant overlaps between hypermethylated enhancers (upper panel) or promoters (bottom panel) in CD11B+ cells from IDH^mut gliomas, compared to IDH^wt gliomas or normal brain tissues. P-values were computed using over-representation analysis (ORA). **(E)** Bubble plots indicate the hallmark gene sets enriched in the lists of genes exhibiting hypermethylated enhancers (n=3027) and promoters (n=349) in CD11B+ samples from IDH^mut gliomas in comparison to CD11B+ samples from IDHwt gliomas and normal brain tissues (Fig. 1D). The size and color code of each dot indicate the number of genes with significant enrichment for the indicated categories (FDR < 0.05). **(F)** Significant sequence motifs (ranked by p-value) enriched in hypermethylated enhancers and promoters in CD11B+ samples from IDH^mut gliomas, in comparison to CD11B+ samples from IDH^wt gliomas and normal brain tissues (Fig. 1D), were determined using HOMER.

Transcription factors (TFs) bind to specific DNA motifs on promoters and enhancers to control cell-type specific transcriptional programs, and the affinity can be modulated by the methylation status[36-38]. Since a substantial proportion of hypermethylation observed in CD11B+ cells from IDH^mut tumors involved regulatory elements in both comparisons (Fig. 1c), we sought to examine enriched TF binding motifs located within hypermethylated regulatory regions. This analysis revealed an overrepresentation of 21 motifs at 349 hypermethylated promoters and 24 motifs at 3027 hypermethylated enhancers (p-value < 10^{-12}) (Fig. 1f). We found SPI1 (PU.1) as the most significantly enriched motif in both hypermethylated regulatory regions (36% of enhancers and 32% of promoters). Interestingly, the other top enriched motifs were bound by key TFs controlling an environment-dependent transcriptional network in microglial cells such as members of C/EBP, AP-1, RUNX, and IRF[39], alongside NF-κβ. Collectively, these results reveal extensive differences in the DNA methylome of TAMs as a function of the IDH mutational status.

Hypermethylation correlates with reduced expression of inflammatory regulons in CD11B+ cells from human IDH^{mut} gliomas

To investigate the transcriptome landscape of TAMs in gliomas, we performed RNA-seq on CD11B+ fractions (n=28, Supplementary Table 1). PCA indicated a good separation between samples derived from IDH^{wt} and IDH^{mut} cases (Extended Data Fig. 3a). Differential expression analysis identified 2714 genes upregulated and 1404 genes downregulated (log2FC > 1 or < -1, respectively; FDR < 0.05) in TAMs from IDH^{mut} compared with those from IDH^{wt} (Supplementary Table 3). We found significant overlaps between the list of downregulated genes in our CD11B+ sample series and those reported by Klemm et al., who further discriminated the expression changes displayed by microglia and MDMs (Extended Data Fig. 3b). Noteworthy, the largest overlap occurred in downregulated genes within microglial cells from IDH^{mut} tumors, with 156 out of the 357 genes falling into this category. Furthermore, significant proportions of genes defining normal microglial signatures also intersected with the list of downregulated genes from our series of CD11B+ fractions in IDH^{mut} gliomas (Extended Data Fig. 3c). Enrichment analysis revealed that these downregulated genes are involved in proinflammatory pathways such as INF-γ/α response and IL6/JAK/STAT3 signaling, cell cycle/proliferation, hypoxia, glycolysis and the acquisition of mesenchymal-like traits (Extended Data Fig. 3d).

DNA methylation is an important mechanism of gene expression regulation. To evaluate the impact of methylation on the observed transcriptional changes in CD11B+ cells isolated from IDH^{mut} gliomas, we intersected the lists of differentially expressed genes with the lists of differentially methylated promoters or enhancers. This analysis involved 36 CD11B+ samples, with 18 of them having both data available (Supplementary Table 1). Contrary to IDH^{mut} glioma cells, where methylation and expression changes have been shown to be uncoupled[35,40], myeloid cells from these tumors displayed confident overlaps in the expected direction: for instance, reduced gene expression was strongly associated with hypermethylation at promoters and enhancers (p-value=2.5e-14 and p-value=7.7e-18, respectively, over-representation analysis (ORA) (quadrant IV, Fig. 2a), while the occurrence

Figure 2. Integration of significant changes in the transcriptome and methylome displayed by CD11B+ myeloid cells in IDHmut gliomas. (A) Cartesian diagrams represent genes (dots) exhibiting significant changes in both expression (Y-axis) and methylation (X-axis) at promoters or enhancers in CD11B+ cells from IDHmut vs. IDHwt gliomas. The total number of genes displaying differential methylation and expression changes is indicated for each quadrant. Over-representation of canonical methylation/expression associations (quadrants II and IV) and under-representation of non-canonical associations (quadrants I and III) were evaluated using ORA and URA, respectively. Quadrants IV (highlighted in red) indicate genes of interest exhibiting downregulation and hypermethylation at either promoters or enhancers **(B)** Bubble plot indicates the "hallmark gene sets" enriched in the integrative gene list (n=370) obtained by combining quadrants IV from both promoters and enhancers analyses. The size and color code of each dot indicate the number of genes with significant enrichment in the corresponding categories (FDR < 0.05). **(C)** The plot illustrates a subset of genes (n=110) from the integrative list that are enriched (FDR < 0.05) in Gene Ontology biological processes related to immune response and metabolism. The hypermethylation status at promoter and enhancer regions is also indicated. Positive or negative hits are represented in green or gray, respectively.

of genes showing non-canonical associations such as overexpression coupled with hypermethylation at promoters and enhancers, was lower than expected (quadrants I, Fig. 2a) (p-value=8.4e-3 and 2.4e-15, under-representation analysis (URA). This integrative analysis of methylome and transcriptome changes revealed 370 unique genes with a hypermethylated enhancer and/or promoter linked to downregulated expression in CD11B+ cells from IDHmut tumors (Supplementary Table 4).

The integrative gene list contained key factors involved in TNF-α signaling via NF-κβ, IL6/JAK/STAT3 signaling, and response to IFN-γ/α (Fig. 2b and Supplementary Table 4). Biological processes such as chemotaxis, leukocyte activation, and positive regulation of metabolism also appeared enriched in this gene list (Fig. 2c). Among the genes whose downregulation was associated to hypermethylation at both regulatory regions, we found CIITA, the most important transactivator of MHC class II, the cytokine oncostatin M (OSM), and myeloid receptors such as formyl peptide receptor 1 (FPR1), and Toll-like receptor 2 (TLR2) (Fig. 2c). Hypermethylation at either promoter or enhancer together with reduced expression was observed for HLA genes, proinflammatory cytokines CXCL10, IL18, and CXCL8, chemokine receptors CCR2/5, FPR3 and TLR1/8, master regulators of glycolysis/hypoxia HIF1A, PKM, ENO1, CD44, and signaling components or enzymes

involved in cytokine production and NAD biosynthesis, such as TYROBP and NAMPT, respectively (Fig. 2c). Therefore, hypermethylation appears to affect various levels of signaling regulation within inflammatory pathways and, possibly, the activation of myeloid cells.

To assess the influence of DNA hypermethylation on TF binding at the enhancers highlighted in quadrants I and IV (Fig. 2a), we estimated TF activity by examining high-confidence TF-gene interactions (n=1,209 human TFs)[41,42] in our RNA-seq data of CD11B+ samples. Analysis of regulons identified 45 TFs having differential activity (FDR <0.05; Wilcoxon's test) according to the IDH mutation status of corresponding tumors, with the majority of TFs showing a decreased activity in IDH^mut gliomas (Fig. 3a). Consistent with the binding motifs enriched in the entire repertoire of hypermethylated enhancers (Fig. 1f), this analysis emphasizes that lineage-determining and stimulus-related microglial TFs are repressed from DNA binding by methylated CpGs in CD11B+ cells from IDH^mut gliomas. Among these, SPI1 (PU.1), AP-1 complex (ATF2, FOS and JUN), and CEBPB, have previously been shown to be sensitive to methylation[43-45], while NFATC2, RELA and STAT3 have been associated with microglial super-enhancers[39]. Similar results were observed when using the whole set of hypermethylated promoters (Extended Data Fig. 4). In addition, these analyses revealed low activity of MYC and HIF1A (Fig. 3a and Extended Data Fig. 4).

On the other hand, a few TFs appeared activated in these cells, including TP63, TP73, SNAI1, CTNNB1, EPAS1/HIF2A and KLF4. In these cases, methylated CpGs could promote TF binding[46]. Upon closer examination of specific TFs inactivated in CD11B+ cells from IDH^mut gliomas, a network inference analysis also suggests that TF-target genes implicated in inflammation and antigen presentation are effectively downregulated as a result of disrupted cooperative effects of core microglial TFs (Fig. 3b). Similarly, targets that are commonly regulated in a positive manner by both MYC and HIF1A, which play roles in the response to hypoxia and glycolysis, were more enriched than the targets regulated by each individual TF in the list of downregulated genes in CD11B+ cells from IDH^mut vs. IDH^wt tumors (Fig. 3c).

Figure 3. Gene regulatory network analysis reveals lower activity of core microglial TFs linked to hypermethylated enhancers in CD11B+ cells from IDH^mut gliomas. (A) The heatmap depicts TFs (n=45) exhibiting differential activity (FDR <0.05; Wilcoxon's test) in CD11B+ cells according to the IDH mutation status of corresponding tumors. The TFs targets used for this analysis, retrieved from the CollecTRI database were searched among the genes displaying upregulated/downregulated expression and hypermethylation at enhancers in myeloid cells from IDH^mut gliomas (quadrants I and IV, Fig. 2A). The number of targets regulated positively and negatively by each TF is indicated in brackets. Unsupervised clustering was performed using the z-scores of TF activity, where red and blue colors indicate high and low activity, respectively, and non-significant activity is colored in gray. **(B)** Gene regulatory network of four representative TFs (blue background) having lower activity in CD11B+ cells from IDH^mut compared to IDH^wt gliomas. TFs targets have a red, orange, pink or green background if they are controlled by four, three, two or one TF, respectively. Targets with or without black box come from quadrant I (hypermethylated enhancers and upregulated expression) or quadrant IV (hypermethylated enhancers and downregulated expression) of Fig. 2A, respectively. **(C)** Gene set enrichment analysis of positive gene targets of HIF1A, MYC (retrieved from CollecTRI database) and those in common between these TFs. Genes are ranked using log2 fold change from differential expression analysis of CD11B+ samples in human gliomas (IDH^mut, n=14 vs. IDH^wt, n=8).

Of note, shared targets of these TFs involved in glycolysis (CD44, ENO1, GAPDH, HIF1A, ID2), and targets of HIF1A involved in NADH regeneration and mitochondrial function (ENO2, NAMPT, PKM, PFKFB4, SDHB) also appear in the integrative gene list (Fig. 2c). These findings were validated in published RNA-seq data of microglia and MDMs from IDH^mut vs. IDH^wt gliomas, with the most marked differences being displayed by microglia respect to MDMs (Extended Data Fig. 5). Overall, these results highlight enhancers as main targets of microenvironmental perturbations affecting microglial activation in IDH^mut gliomas and raise questions about the underlying mechanisms driving hypermethylation.

D-2HG inhibits TET-mediated 5mC oxidation and promotes DNA hypermethylation at lineage-specific enhancers in human microglia

We hypothesized that D-2HG, the oncometabolite produced by the IDH^mut glioma cells and released into the TME, could impact the DNA methylome of CD11B+ cells. Ten-eleven translocation (TET) enzymes are α-KG-dependent dioxygenases that mediate the first step of active DNA demethylation by catalyzing the conversion of 5-methylcytosine (5mC) into 5-hydroxymethylcytosine (5hmC)[47] (Extended Data Fig. 6a). Inhibition of TET-mediated DNA demethylation by D-2HG results in decreased levels of 5hmC along with increased levels of 5mC after several rounds of cell replications, ultimately establishing the G-CIMP-

high phenotype of IDH[mut] gliomas[27,48]. To explore whether this mechanism is at play in the myeloid cells of these tumors, we analyzed global levels of 5hmC and 5mC, as well as the cumulative cell replication history, in an independent series of CD11B+ and CD11B- fractions isolated from human gliomas, and from normal tissues used as controls (Extended Data Fig. 6b and 6c). We confirmed a higher 5mC/5hmC ratio in CD11B- cells of IDH[mut] gliomas compared to non-tumor tissues (Extended Data Fig. 6b). Remarkably, a significant increase in 5mC/5hmC ratio was also observed in corresponding myeloid cells along with a higher cumulative number of cell divisions as indicated by epigenetic clock analysis (Extended Data Fig. 6c). These results suggest that TET inhibition contributes to the global DNA hypermethylation observed in myeloid cells (most likely microglia) from IDH[mut] gliomas.

To ascertain the effect of D-2HG on TET-mediated DNA demethylation in human microglia, we set up primary cultures obtained from tissue aspirates of patients who underwent surgery for glioma or drug-resistant epilepsy (Fig. 4a). Microglial cells, maintained in a serum-free defined culture medium[49], are viable for up to 15 days. Over 95% of these cells express the markers ionized calcium-binding adapter molecule 1 (IBA1) and transmembrane protein 119 (TMEM119) (Fig. 4b), whereas less than 5% of cells were proliferating (KI67+) (not shown). Furthermore, these cells maintain a stable RNA-seq microglial signature over time (Extended Data Fig. 7a) and are functional, as evidenced by the proinflammatory response upon lipopolysaccharide (LPS) stimulation at 48 hours or 14 days (Extended Data Fig. 7b and 7c). To evaluate the uptake of D-2HG by human microglial cells, we measured intracellular levels after 48 hours of treatment at a concentration estimated to occur in the TME[4,5], which does not affect cell viability. We found that the levels of D-2HG reached by microglial cells were comparable to those observed in *bona fide* IDH[mut] cells (Extended Data Fig. 8a and 8b, and Fig. 4c) and were sufficient to induce a significant decrease in TET enzymatic activity (Fig. 4d and Extended Data Fig. 8c). We then exposed primary microglia cultures to D-2HG for a prolonged period and observed that global levels of 5hmC in treated and control microglia have opposite trends over time, leading to a significant decrease of 5hmC levels in treated cells at 14 days (Fig. 4e and Extended Data Fig. 8d).

Figure 4. D-2HG uptake by human primary microglial cells inhibits TET function. (A) Overview of the experimental workflow for establishing primary microglia cultures treated with D-2HG, and analyses assessing the inhibition of TET-mediated DNA demethylation. The tissue collection procedures and culture conditions are thoroughly detailed in the Methods section. **(B)** Representative immunofluorescence images show the preserved expression of microglial markers TMEM119 (confocal microscopy) and IBA1 (Apotome) in primary cells at 12 days in culture (scale bars: 100 µm). Estimated percentages of TMEM119+ and IBA1+ cells in primary cultures at different time points are indicated on the right. **(C)** The quantification of intracellular levels of 2HG was determined using LC/MS in primary human microglia obtained from five surgical aspirates with IDH^{wt} (n=2) or IDH^{mut} (n=3) status. Data are represented as the mean with a 95% CI; p-value was calculated using a one-sided paired t-test. **(D)** TET enzymatic activity measured as detection of hydroxymethylated product (ng) after 1-hour incubation of the substrate with protein extracts from microglia untreated and treated with 5 mM D-2HG over 48 hours. Primary cultures were generated from surgical aspirates with IDH^{wt} status (n=3). The p-value was calculated using one-sided paired t-test. **(E)** Assessment of global levels of 5hmC, determined by LC-MS/MS, in primary human microglia obtained from six surgical aspirates with IDH^{wt} (n=5) or IDH^{mut} (n=1) status after 48 hours and 14 days of treatment with 5 mM D-2HG (added every 2-3 days). Data are represented as the mean with a 95% CI; p-value was calculated using a one-sided paired t-test. **(F)** DNA methylation profiling of primary microglia was conducted with EPIC array. Stacked bar chart shows absolute numbers and relative percentages of functional genomic regions respect to the methylation status (absolute Δβ > 0.03, FDR < 0.25) in D-2HG treated vs. untreated primary microglia cultured for 14 days. Samples were obtained from eight surgical aspirates with IDH^{wt} (n=5) or IDH^{mut} (n=3) status. **(G)** Volcano plot illustrates the magnitude and extent of differentially methylated enhancers in D-2HG treated vs. untreated primary microglia. **(H)** Motif enrichment analysis of hypermethylated enhancers in D-2HG-treated primary microglia. Significant sequence motifs (ranked by p-value) were determined using HOMER.

No concomitant changes in 5mC levels were observed in D-2HG-treated microglia (data not shown). These results suggest that TET-mediated demethylation in non-dividing human microglia is inhibited by D-2HG. However, contrary to the *ex vivo* situation (Extended Data Fig. 6b and 6c), the low proliferation rate of cultured microglia and the reduced time window (only 2 weeks) precluded our ability to demonstrate D-2HG-induced global hypermethylation. Nonetheless, further analysis of DNA methylome in microglial cells treated with D-2HG for 14 days, using the same EPIC array for the analysis of *ex vivo* CD11B+ cells in gliomas, revealed subtle yet reproducible hypermethylation at 283 enhancers (absolute Δβ > 0.03, FDR < 0.25) (Fig. 4f and 4g). Importantly, these enhancers showed significant enrichment for motifs of lineage-determining and stimulus-responsive TFs as in the *ex vivo* study, including AP-1, C/EBP and NF-κβ, but not SPI1 (PU.1) (Fig. 4h).

While EPIC arrays offer sufficient coverage of enhancers, only a single probe captures methylation changes across an enhancer region[50]. Moreover, this profiling method does not discriminate 5hmC from 5mC, which could provide valuable insights into impaired TET function. Considering the high turnover of DNA methylation at enhancers, which relies on TET activity rather than replicative passive dilution of 5mC[51], we wondered whether D-2HG affects TET-mediated demethylation at enhancers genome-wide. To investigate this possibility, we performed a DNA methylome and hydroxymethylome profiling of microglial cells treated with D-2HG for 48 hours or 14 days. For this experiment, we used microglial preparations from normal brain tissues of three individuals with drug-resistant epilepsy to exclude tumor influences other than released D-2HG, and we applied a reduced representation and non-bisulfite sequencing approach[52]. The extensive coverage enables a more confident assessment of 5mCG and 5hmCG levels in enhancers, yet only across 20% of the enhancers analyzed with EPIC arrays, which cover ~58% of FANTOM annotated enhancers[50]. Changes of 5mCG and 5hmCG levels in the entire enhancer regions were most evident at 14 days of treatment respect to 48 hours, with a marked difference for 5hmCG, reflecting its accumulation over time in D-2HG-treated cells (Fig. 5a). This result indicates that D-2HG-related changes in genomic 5hmC content, initially detected by LC-MS/MS, mostly occur in non-CpG contexts, and shows that TET function is not affected in the entire enhancers regions, at least *in vitro*. Moreover, while EPIC arrays revealed some lineage-specific enhancers that underwent hypermethylation following D-2HG treatment (Fig. 4f-4h), only a few of them were covered by the reduced representation sequencing approach.

To ascertain whether inhibition of TET-mediated demethylation by D-2HG affects a subset of lineage-specific enhancers, we pursued the following analytical strategy with our sequencing data (Fig. 5b). We aimed to use as ground truth the enhancers in CD11B+ cells from normal brain tissues, which eventually become hypermethylated in myeloid cells from IDH[mut] gliomas, likely corresponding to microglia (Fig. 1c). Considering that the highest level of TET activity occurs around TFs binding motifs located in enhancers[51,53], we then focused our analysis on TFs binding motifs within hypermethylated enhancers found in both CD11B+

cells from IDH^{mut} gliomas (Fig. 1f) and D-2HG-treated microglia (Fig. 4h), potentially impacting TF occupancy (Fig. 3a and 3b).

Figure 5. Reduced representation non-bisulfite sequencing reveals high 5mCG/5hmCG ratios at lineage-specific enhancers in D-2HG-treated primary microglia. (A) Bar plots show the percentages of 5mCG and 5hmCG at the whole set of covered enhancers (full-length regions) in primary human microglia treated with D-2HG for 48 hours or 14 days and in paired untreated controls. The number of paired enhancers for each comparison is indicated. Only common enhancers across patients were analyzed. **(B)** Workflow analysis of 5mCG and 5hmCG levels quantification in defined windows around TFs binding motifs, located in lineage-specific enhancers, in D-2HG-treated microglia for 14 days and in paired untreated controls. **C)** Plots showing a separate analysis of 5mCG and 5hmC percentages (upper) around TFs binding motifs for the indicated TFs in D-2HG-treated microglia and paired untreated controls. Only common enhancers across patients were analyzed. The size of each dot indicates the number of enhancers analyzed in a paired manner between conditions. The 5mCG/5hmCG ratios (bottom), which are higher in D-2HG-treated microglia than in paired controls, are also shown. The TFs binding motifs for this analysis are located in enhancers exhibiting hypermethylation in CD11B+ cells from IDH[mut] gliomas vs. normal brain tissues in the *ex vivo* study (Fig 1C).

Notably, higher 5mCG/5hmCG ratios were detected around binding motifs of members of the AP-1, C/EBP and NF-κβ TFs families in D-2HG-treated primary microglia compared to paired untreated controls with these motifs being located in hypermethylated enhancers of CD11B+ cells from IDH[mut] gliomas vs. normal brain tissue (Fig. 5c). Similar results were obtained when this analysis was extended to motifs of SPI1 (PU.1), IRF4/8, RUNX3, HIF1A and MYC, which also appeared to be relevant in the *ex vivo* situation (Fig. 5c). In sharp contrast, 5mCG and 5hmCG levels in the proximity of binding motifs located in the whole set of covered enhancers, exhibit opposite trends (Extended Data Fig. 9a). Notably, the genes regulated by the selected enhancers bound by key TFs (Fig 5c), that exhibited hypermethylation/hypohydroxymethylation, were enriched among the downregulated genes, indicating a direct impact of D-2HG-induced hypermethylation on gene expression (Extended Data Fig. 9b and Supplementary Table 5). Taken together, these results strongly support that D-2HG has the potential to impede TET-mediated DNA demethylation selectively at lineage-specific enhancers, leading to a blunted microglial cell response.

D-2HG dampens the LPS-induced inflammatory response and increases basal oxidative metabolism in human microglial cells

Lineage-specific enhancers that were affected by D-2HG treatments in our *in vitro* experiments have been shown to be transcribed upon macrophage stimulation[54]. Thus, we

hypothesized that the methylome primed by D-2HG could hinder the activation of enhancers governing the proinflammatory capacity of human microglia. To address this issue, we evaluated transcriptional changes in cells pre-treated with D-2HG for 14 days and subsequently challenged with LPS for 6 hours (Fig. 6a).

Gene set enrichment analysis (GSEA) indicated that treatment with D-2HG attenuates LPS-induced microglial activation (Fig. 6b). Specifically, these cells exhibit an M2-like phenotype characterized by enhanced OXPHOS and fatty acid metabolism alongside marked suppression of the inflammatory response pathways including TNF-α signaling via NF-κβ (Fig. 6b). Additionally, in RNA-seq data from the *ex vivo* study, we find reduced expression of proinflammatory cytokines in CD11B+ cells from IDH^mut compared to IDH^wt gliomas (Extended Data Fig. 10a). We then prompted to evaluate the effect of D-2HG on the expression of LPS-induced cytokines using RT-qPCR in an independent series of primary microglia preparations. We confirmed that the expression of TNFA and IL6 was abrogated by D-2HG in LPS-challenged cells compared to paired controls (p-value=0.02 and 0.07, respectively) (Extended Data Fig. 10b). Similarly, assessment of mitochondrial respiration parameters demonstrated that D-2HG-treated microglia exhibit an increase in basal and maximal respiration even after LPS stimulation (Extended Data Fig. 10c). This observation suggests that this oncometabolite impedes the glycolytic shift that accompanies cellular activation and instead favors OXPHOS metabolism. All in all, these results indicate that D-2HG dampens microglia proinflammatory response to LPS, likely through direct effects on TET-mediated demethylation at enhancers governing microglial activation.

Lastly, to determine whether suppressed microglial cell signaling linked to D-2HG-driven enhancer hypermethylation can be restored upon targeted therapy, we analyzed microglial cells from single-nucleus RNA-seq (snRNA-seq) data obtained from one oligodendroglioma patient treated with ivosidenib, an inhibitor of the IDH1 mutant enzyme that reduces D-2HG levels in tumor bulk (Fig 6c and Supplementary Table 1)[55].

Figure 6. D-2HG impairs microglial activation. (A) Experimental workflow for the LPS challenge experiment in primary microglia pre-treated with D-2HG for 14 days. RNA-seq data was generated from human primary microglia obtained from four surgical aspirates with IDH^{wt} (n=3) or IDH^{mut} (n=1) status. Patient-paired approach was adopted for RNA-seq data analysis. **(B)** GSEA ridge plot illustrates "hallmark gene sets" in D-2HG-treated microglia for 14 days, 6 hours after LPS stimulation. Gene sets with positive or negative normalized enrichment score are associated with up or down regulated genes, respectively (FDR < 0.05). **(C)** Uniform Manifold Approximation and Projection (UMAP) plots depict microglial cells analyzed with snRNA-seq from one oligodendroglioma patient pre- and on-treatment with the mutant IDH inhibitor ivosidenib. **(D)** The heatmap illustrates the significantly upregulated genes (log2FC > 1, Bonferroni-adjusted p-value < 0.05) in microglia upon ivosidenib treatment. **(E)** Normalized counts of selected genes in microglia pre and on-treatment. Data are represented as the mean with a 95% CI.

After removing low quality cells, 178 microglial cells remained, as determined by the high expression of markers TAL1 and SLC2A5, and the low/absence of expression of MDMs markers CD163 and TGFBI. These cells encompassed two well-defined clusters corresponding almost entirely to on- (n=87) and pre-treatment (n=91) microglia, on which the analysis was focused (Fig 6c). We noticed a transcriptional shift towards gene upregulation in cluster 1 (on-treatment) compared to cluster 2 (pre-treatment) (1683 genes with log2FC >1, 501 genes with log2FC < -1, respectively). Furthermore, we found 24 upregulated (including the master regulators TFs FOS and CEBPB) (Fig 6d) and 3 downregulated genes (Bonferroni-adjusted p-value < 0.05, log2FC > 1 or < -1, respectively) with a strong enrichment of the upregulated genes in the TNF-α signaling via NF-κβ gene set (FDR=1.5e-4; ORA). Additionally, microglial cells in cluster 1 displayed overexpression of other TFs governing activation, including AP-1 family members (FOSB, ATF3, JUNB, and JUND), REL, and HIF1A, as well as related target genes highlighted in the *ex vivo* and *in vitro* analyses, such as OSM, IL18, IL1B, FPR1, TLR2, NAMPT, TNF, TYROBP, and SLC2A3 (GLUT-3) (Fig 6d and 6e). This result suggested that decreased levels of D-2HG in the TME may reverse the hypermethylation phenotype observed in microglia from IDH^{mut} gliomas, thereby enabling their activation program. Supporting this hypothesis, a significant proportion of genes upregulated following ivosidenib treatment are regulated by enhancers that were found to be hypermethylated in CD11B+ cells from IDH^{mut} gliomas (p-value = 0.0012, ORA).

Discussion

The development of new therapeutic options to cure glioma requires a deep understanding of the epigenetic mechanisms governing transcriptional programs, cell states, and proinflammatory capacities in TAMs. Differences in the transcriptome and chromatin landscapes have been associated with the IDH[mut] status[17,18,21], but the extent and the role of DNA methylation in TAMs have remained uncharacterized. Herein, through genome-wide profiling of the DNA methylome, we uncovered global and differential distribution of 5mC and 5hmC changes in TAMs according to the IDH[mut] status of gliomas. Our findings revealed how this differential methylation affects microglial enhancers involved in the regulation of proinflammatory and metabolic pathways.

We found that CD11B+ cells residing in an IDH[mut] environment display a striking bias towards DNA hypermethylation. TAMs comprise MDMs and microglia in proportions and states of activation that differ according to glioma IDH[mut] status, subtype, and grade[17-23]. Previous studies reported that most of the TAMs in IDH[mut] tumors are microglia, whereas IDH[wt] gliomas are marked by an influx of MDMs[17-20]. In this regard, we do not rule out the possibility that differentially methylated changes observed in CD11B+ cells partly reflect a decreased recruitment of MDMs in IDH[mut] compared to IDH[wt] tumors. However, the hypermethylation we observed holds true even when comparing CD11B+ cells from IDH[mut] glioma to non-tumor brain tissues, both of which are highly enriched in microglia. In addition, a recent study indicated that immunomodulatory activity programs are expressed across myeloid cell types, and demonstrated convergent phenotypes of resident microglia and infiltrating MDMs, with the latter acquiring a microglia-like phenotype in tumors[23]. Moreover, mouse studies indicate that MDMs can adopt a DNA methylation profile very similar to that of microglia after colonizing the central nervous system niche[56]. Therefore, our bulk profiling of CD11B+ cells strongly suggests that microglia undergo a hypermethylated state in IDH[mut] tumors. This assumption is further substantiated by our analyses of methylation across gene regulatory regions, revealing functional implications related to proinflammatory activation and metabolism in microglia.

We highlighted that the occurrence and extent of differentially methylated changes were more prominent in distal enhancers. DNA methylation at CpG sites within or flanking motifs may prevent TF binding at regulatory regions by inducing steric hindrance, altering DNA shape and rigidity, or influencing nucleosome positioning, thereby affecting the formation of active chromatin[36,38,43,57-59]. A recent study showed that only a minority of enhancers in a given cell type are sensitive to DNA methylation[60]. In line with this, our integrative analysis of methylome and transcriptome changes demonstrates that, among the hypermethylated enhancers displayed by CD11B+ cells from IDH[mut] vs. IDH[wt] comparison, 23% may account for the downregulated expression of target genes. We also observed that hypermethylated enhancers were enriched for motifs of lineage-determining TFs, including SPI1 (PU.1) and members of the C/EBP and RUNX families, as well as motifs for stimulus-induced TFs, such as AP-1, IRF, and NF-κB. These TFs are known to regulate the functional properties of myeloid cells. Our interrogation of gene regulatory networks further supports the hypothesis that hypermethylated enhancers are responsible for the reduced activity of core TFs that govern the identity and activation program in human microglial cells[39]. Specifically, hypermethylation seems to disrupt cooperative regulatory functions involving lineage-determining TFs SPI1 (PU.1) and C/EBP, and stimulus-responsive TFs AP-1 and NF-κβ, which in turn affect the expression of key genes involved in inflammatory responses. Hypermethylated enhancers also appear to influence the expression of activating receptors, as well as of genes involved in antigen presentation, an observation previously reported in bulk methylome analysis of IDH[mut] gliomas[61]. These findings indicate that microglia in these tumors undergo a profound alteration of promoter-enhancer interactions that characterize these cells[62].

In addition to microglial TFs, our data revealed the inactivation of HIF1A and MYC, which possibly reinforces the silencing of shared targets essential for the glycolytic shift of stimulated macrophages[63-66]. Along with a marked downregulation of several target genes, there was low expression of HIF1A, accompanied by hypermethylation at regulatory regions

of HIF1A and some of its targets. In particular, reduced expression of HIF1A targets such as LDHA, ENO1, HK2, PKM, VIM, ITGB2, and PGK1 has also been reported in bulk IDH[mut] tumors and/or cells[67], which may be attributable to hypermethylation driving the silencing of these genes in addition to the effect of D-2HG on HIF-1α protein destabilization[68]. Collectively, our findings provide evidence that DNA hypermethylation may contribute to a reduced proinflammatory microglial response by targeting enhancers that orchestrate synergistic responses to environmental stimuli driving myeloid cell activation[69-71]. This assumption is supported by our experiments, which demonstrate how oncometabolite signaling modulates the methylome and subsequently affects the activation of human microglia.

Our analysis of single-nucleus RNA sequencing data of two paired oligodendroglioma samples revealed that treatment with an inhibitor of the IDH1 mutant enzyme could unleash the signaling that suppresses microglia activation. This observation supports the notion that D-2HG production plays a major role in the suppressive signaling that affects microglia function and highlights the prognostic and therapeutic value of microglial activation in IDH[mut] glioma. By establishing primary cultures of human microglia, we have shown that D-2HG is taken up by these cells, inhibits TET-mediated DNA demethylation, and hampers the accumulation of 5hmC genomic content after a 14 days treatment. Although D-2HG-driven global hypermethylation is challenging to reproduce in our microglial cultures, our analysis of methylome and hydroxymethylome at nucleotide resolution clearly demonstrates higher 5mCG/5hmCG ratios at lineage-specific enhancers that were also hypermethylated in *ex vivo* CD11B+ cells from IDH[mut] gliomas. This result is consistent with the previous demonstration that the highest TET activity is localized to enhancer elements[51,53]. Our observation that the 5hmC turnover is limited in cultures of non-proliferating microglial cells points to a passive demethylation mechanism that could unfold under successive rounds of replication. Nevertheless, a comprehensive characterization of other TET-mediated oxidized derivatives, such as 5fC and 5caC, and an assessment of the activity and the role of thymidine DNA glycosylase[47], will be required to determine the relative contribution of active vs. passive demethylation mechanisms in human microglia.

Regardless of the demethylation mechanism, we speculate that inhibition of TET2 in slowly proliferating microglial cells of IDH[mut] gliomas, which are exposed to D-2HG over several years, may lead to full-blown hypermethylation, as opposed to the limited changes observed in our non-proliferating microglial cultures exposed to this oncometabolite for no more than two weeks. It is also possible that additional mechanisms contribute to the hypermethylation highlighted in the *ex vivo* context. Nevertheless, our *in vitro* system was instrumental in demonstrating that D-2HG readily affects TET-mediated demethylation at specific enhancers, leading to suppressed expression of targeted genes. A high turnover of DNA methylation at enhancers is crucial for cellular plasticity, differentiation, and responsiveness to environmental changes, for instance, during macrophage activation[72,73]. Thus, the significant overlap of enhancer elements impacted by D-2HG *in vitro* with those exhibiting hypermethylation in myeloid cells from IDH[mut] gliomas strongly suggests that the methylome of microglial cells primed by D-2HG negatively affects the transcriptional response to inflammatory stimuli.

Consistent with this hypothesis, we observed that D-2HG-treated microglia, when challenged with LPS, display attenuation of the inflammatory response while favoring an OXPHOS metabolism that marks anti-inflammatory M2-like macrophage phenotypes[74,75]. While the hypermethylation we observed may affect the glycolytic shift—a metabolic response documented in iPSC-derived human microglia treated with LPS[76]— our data indicate that hypermethylation could also impact genes involved in cytokine production. Considering the time-dependent effects of D-2HG on the microglia methylome, we propose that an epigenetic drift—involving the selection of hypermethylated enhancers over time— underlies the silencing of master regulators, which impedes not only the proinflammatory response but also the glycolytic reprogramming, both of which are essential for microglial cell activation. This does not exclude the possibility of contributions from chromatin remodeling and non-epigenetic mechanisms to the LPS hyporesponsiveness of D-2HG-treated microglia, including signaling pathways targeted by D-2HG in other cell types[6,9,10,77].

In particular, it has been reported that acute octyl-D-2HG treatment can dampen LPS-triggered cytokine expression in the immortalized mouse BV2 microglial cell line through a mechanism involving AMP-activated protein kinase (AMPK)-mTOR-mediated reduction of NF-κB activity[77]. Whether this mechanism also applies to human primary microglia remains undetermined.

In conclusion, we provide evidence that D-2HG reshapes the enhancer landscape of resident microglia in IDH^mut gliomas, thus impairing the regulatory function of core TFs. This priming mechanism precludes the capacity of these cells to mount the inflammatory response and possibly the shift towards a glycolytic metabolism. We speculate that D-2HG effects on myeloid cells promote a cold immune TME that favors tumor progression. Given that the reduced concentration of D-2HG in bulk tumors, induced by inhibitors of the IDH mutant enzyme, is associated with inflammatory and metabolic changes[78], it is important to evaluate the extent of these effects in myeloid cells and to correlate them with the clinical response to this therapeutic option.

Limitations of the study

We acknowledge certain limitations in our study that should be taken into consideration. Our comparative analyses of CD11B+ myeloid fractions from gliomas and non-tumor tissues remained at the bulk level. Moreover, the methylome profiling is limited to CpG residues in the EPIC array and to global levels assessed by LC-MS/MS, whereas no exhaustive analysis of 5hmC at nucleotide resolution was conducted *ex vivo*. While our results indicate that hypermethylation affects the dominant population of resident microglia cells in IDH^mut gliomas, employing single-cell multimodal approaches will be necessary to better associate enhancer landscapes with their regulatory roles in specific myeloid cell subsets/states. Lastly, we did not analyze the potential effects of D-2HG on the local chromatin environment, which could also influence enhancer function.

Acknowledgements

The authors express their deepest gratitude to the memory of Prof. Dr. Laurent Capelle for his insightful feedback and invaluable contribution to the collection of neurosurgical specimens for research purposes, including the present study. The authors also thank the personnel working at the ICM platforms iGenSeq (sequencing), Celis (cell culture), and Histomics (histology) for their technical support, and Julie Jardon, Coralie Gimonnet and Giovanni Scala for their assistance in experiments and bioinformatics. This work is supported by the Ligue Nationale contre le Cancer (Équipe Labellisée), as well as by the following grants: INCa-DGOS-Inserm 12560, Site de Recherche Intégré sur le Cancer (SiRIC CURAMUS), French National Cancer Institute (INCa-PLBIO22-243), Fondation Bristol Myers Squibb pour la Recherche en Immuno-Oncologie (BMS 2104009NA), French National Cancer Institute (INCa), Fondation ARC and Ligue National contre le cancer (LNCC) grant (PAIR TUMC21-001, INCa-16280), and Entreprises contre le Cancer Paris-Île-de-France (GEFLUC R20202DD). A.L. is supported by a Fondation Recherche Médicale (FRM) scholarship, Q.R. is supported by a French Ministry of Education and Research scholarship, and Y.H. is supported by a La Ligue Nationale contre le Cancer scholarship.

Author contributions

A.L: Conceptualization, optimization of primary microglial cultures, designed and performed *in vitro* experiments, data collection, analysis and interpretation of results, draft editing. P.P: Conceptualization, data curation, computational and statistical analyses, interpretation of results, data visualization, draft editing. B.M: Provided surgical resection specimens, optimization of primary microglial cultures, draft editing. Q.R., S.J: Performed *ex vivo* experiments and data visualization. Y.H., S.S: Conducted validation experiments, data analysis and interpretation, data visualization, draft editing. P.M: Provided surgical resection specimens. K.L., A.A: Bioinformatics assistance, draft editing. M.V., A.I., C.B., E.H., N.P., K.M., S.T., M.L.S., M.T., F.B: Provided material, data curation. A.D., I.F: Performed LC/MS analysis of 2HG. G.A.K., P.J.K., L.J.M: Performed LC-MS/MS and data analysis of global 5mC and 5hmC levels. E.D., E.E-H: Performed seahorse experiments and data analysis. L.A.S: analysis of

epigenetic clocks and draft editing. A.I: Data analysis and interpretation, draft editing. M.C: Computational analyses, draft editing. M.M: Conceptualization, experimental design, data analysis and interpretation, draft editing. M.S: Conceptualization, experimental design, data analysis and interpretation, draft editing, funding acquisition, supervision. L.J.C-V: Conceptualization, experimental design, data analysis and interpretation, data visualization, writing–original draft, funding acquisition, project administration and supervision. All authors listed revised the manuscript critically for important intellectual content and approved it for publication.

Conflict statement

L.A.S. is a scientific advisor and co-founder of Cellintec L.L.C., which had no role in this research. M.L.S. is equity holder, scientific co-founder, and advisory board member of Immunitas Therapeutics. The other authors declare no competing or financial interests.

STAR Methods

Tumor samples and CD11B sorting

Human brain tissues were obtained with informed consent approved by the institutional ethical committee board in accordance with the Declaration of Helsinki. Samples for analyses were selected from the Pitié-Salpêtrière tumor bank Onconeurotek based on clinical information and validation by expert neuropathologists (K.M., S.T., F.B) of both histological features, and molecular diagnosis as previously described[79]. Cryopreservation was as follows: tumors taken from the neurosurgery room were immediately transported on ice in HBSS (Gibco), cut into 2-5 mm diameter pieces, and submerged in cryotubes containing 1 mL of DMEM/F-12 (Gibco), 20% FBS, and 10% DMSO (Sigma-Aldrich). The cryotubes were placed at -80°C in alcohol-free freezing containers (Corning® CoolCell®). The cryopreserved samples were then stored in liquid nitrogen until they were used. On the day of analysis, tumor pieces were thawed and rinsed in DMEM/F-12. Next, the tissue was enzymatically and mechanically digested for 5-10 minutes at 37°C in an HBSS-papain-based lysis buffer

(Worthington) containing DNAse (0.01%, Worthington) and L-Cysteine (124 μg/mL, Sigma). Enzymatic digestion was stopped with ovomucoid (700 μg/mL, Worthington). The homogenates were sequentially filtered using 100 μm, 70 μm, and 30 μm SmartStrainers (Miltenyi) to remove residual clumps. HBSS was added, and samples were then centrifuged at 300 g for 10 minutes at 4°C. Cell pellets were resuspended in an appropriate volume of cold MACS debris removal solution following the manufacturer's instructions (Miltenyi). The suspension was gently mixed by pipetting 10-12 times, and ice-cold DPBS without calcium and magnesium was added very slowly above the cell suspension to create a transparent gradient. The suspension was centrifuged at 4 °C and 3000 g for 10 min. After centrifugation, the top liquid and solid interphases were discarded, and cell suspensions were resuspended with cold MACS Buffer (0,5% BSA and 2 mM EDTA in PBS). Tubes were gently inverted 3-5 times and then centrifuged at 1000 g for 10 min at 4°C. After centrifugation, the supernatant was completely removed, and red blood cells lysis was performed with ACK buffer (Gibco) for 5 min at room temperature. Next, cells were centrifuged at 4°C and pellets were washed with cold DPBS. After centrifugation, cells were resuspended in MACS buffer and counted. Next, cell pellets were labeled with an appropriate volume of CD11B MicroBeads (Miltenyi) in MACS buffer and incubated for 15 min at 4°C. Subsequently, cells were resuspended in cold MACS buffer and centrifuged at 300 g for 10 min at 4°C. The supernatant was aspirated, and cells were resuspended in cold MACS buffer. LS MACS Separation Columns were placed in the magnetic field of a suitable MACS Separator, and the cell suspension was applied onto the columns. CD11B+ and CD11B- fractions were separately collected on ice. The dry pellets were stored at -80°C until nucleic acid extraction.

Nucleic acids extractions

RNA and DNA of CD11B+ and CD11B- samples were co-eluted using the AllPrep DNA/RNA Micro Kit (Qiagen, 80284) or AllPrep DNA/RNA Mini Kit (Qiagen, 80204) following the manufacturer's instructions. In some cases, nucleic acids were purified separately using the Maxwell RSC simplyRNA Cells Kit (Promega, AS 1390) and the Maxwell RSC Blood DNA Kit

(Promega, AS 1400), following the manufacturer's instructions. The yield and quality of total RNA was assessed using the Tapestation 2200 (Agilent) instrument.

Digital droplet PCR

IDH1 R132H, TERT C228T, and TERT C250T mutations were analyzed in CD11B+ and CD11B- fractions from human gliomas using ddPCR. Droplet generation and partition was performed using the Bio-Rad system QX200. Primer sets, probes, and ddPCR™ Supermix for Probes (No dUTP) were used for quantification of absolute copy number in 3 ng of input DNA following the manufacturer's instructions. All primers and probes were obtained from Bio-Rad. Sequence information is available at www.bio-rad.com using the ID numbers: IDH1 p.R132H Hsa, Human (Ref: 10031246 UniqueAssayID: dHsaCP2000055), IDH1 WT for p.R132H Hsa, Human (Ref: 10031249 UniqueAssayID: dHsaCP2000056), TERT C228T_113 Hsa, Human (Ref: 12009308 UniqueAssayID: dHsaEXD72405942), TERT C250T_113 Hsa, Human (Ref: 12003908 UniqueAssayID: dHsaEXD46675715). The data were then analyzed using Poisson statistics to determine the target DNA template concentration in the original sample. Positive controls consisted of tumor DNA from the CD11B- fractions and negative controls contained water instead of DNA.

DNA methylation profiling

Genomic DNA was quantified by Quant-iT dsDNA Broad range Assay (Thermo Fisher Scientific, Waltham, MA, USA) in a Tecan SPARK microplate Reader (TECAN, Switzerland). Total DNA (500 ng) from each sample was sodium bisulfite converted using Zymo EZ-96 DNA Methylation kit and following the manufacturer's recommendations for Infinium assay (Zymo Research, catalog number: D5004). After conversion, DNA concentration of each sample was adjusted to 50 ng/µL with M-elution Buffer or concentrated using speed vacuum. In total 300 ng of converted DNA for each sample was used as a template on the Infinium MethylationEPIC BeadChip following the manufacturer's recommendations. Briefly, bisulfite-converted DNA is whole genome amplified, fragmented and hybridized to the array. After hybridization, unhybridized and no specifically hybridized DNA is washed away, and the

captured product is extended with fluorescent labels coupled to nucleotides. Finally, the arrays are scanned with a high-resolution Illumina scanner (iScan), which acquires light images emitted from fluorophores. The intensities are measured and the methylation signals are extracted and recorded as raw data (IDAT). These assays were performed at the P3S platform (Sorbonne University). DNA methylation data were analyzed using the RnBeads package. In short, after the removal of bad quality samples, data were aligned to the hg19 assembly using the beta Mixture Quantile dilation (BMIQ) method for normalization and adjusted for sex as a covariate. Sex chromosomes were included and sites having a standard deviation below 0.01 across all samples were filtered out. Together with default annotations, we included the enhancers coordinates (entire enhancers regions) as defined in the FANTOM5 enhancer-robust promoter associations. Briefly, using Pearson's correlation, enhancers having and FDR < 1e-5 in an enhancer-promoter pair within 500 kb over cell types and organs were retained. The enhancer-promoter associations are computed between enhancers and all FANTOM robust promoters. Epigenetic clock analysis of CD11B+ cells from the *ex vivo study* was conducted using "epiTOC2" (Epigenetic Timer Of Cancer), which calculates the mitotic age of the cells using the updated total number of divisions per stem cell[80]. This new method approximates age as a weighted estimate based on 163 CpG sites interrogated with EPIC arrays.

RNA sequencing

Library preparations were performed with Kapa mRNA Hyper prep (Roche) following the manufacturer's instructions and sequenced (paired end) with the Illumina Novaseq 6000 Sequencing system with 200 cycles cartridge, to obtain 2*60 million reads 100bases/RNA. Sequencing was performed at the iGenseq platform of the ICM – Paris Brain Institute. For the *in vitro* experiments, we combined Illumina Stranded Total RNA Prep and Ribo-Zero Plus. The quality of raw data was evaluated with FastQC. Sequences were trimmed or removed with Fastp software to retain only good quality paired reads. Star v2.5.3a was used to align reads on the GRCh38 reference genome using default parameters except for the maximum number of multiple alignments allowed for a read which was set to 1. Gene level transcript

quantification was done with rsem 1.2.28. Differential expression analysis was conducted using the quasi-likelihood F-test from the edgeR package. Normalized data were computed using the cpm function. Multiple hypothesis testing was corrected using the False Discovery Rate Benjamini-Hochberg method. Samples from all the *in vitro* experiments were analyzed in a paired manner per individual.

Sequence motif enrichment analysis

Motif enrichment was performed using Hypergeometric Optimization of Motif EnRichment (HOMER) v4.11[81], enabling masking repeats, setting the parameter size to "given" and searching for motifs with a length within the range of 7-14 bp. We used HOCOMOCO v12 CORE as the reference dataset (matrix with threshold level corresponding to a p-value of 0.001)[82].

Transcription factor activity

We used CollecTRI database[41] to infer TF activity, which contains regulons comprising signed TF-target gene interactions. To estimate TF activity at hypermethylated enhancers, we took into account genes with associated overexpression or downregulation (falling in quadrants I and IV, respectively, Fig. 2a). These genes were then interrogated among the targets of each TF of the database. We retained the TFs having at least 5 total targets regulated in a positive or negative manner. Then, we computed the TF activity using the Normalized Enrichment Score (NES) of the Mann-Whitney-Wilcoxon genes set test (MWW-GST)[42]. We applied MWW-GST to each sample normalized data and retained TFs having an FDR < 0.05 for the NES in at least 18 samples and an FDR < 0.05 in the comparison of the activity between groups (Wilcoxon's test). For promoters, we took into account all genes associated with hypermethylation in the IDH^{mut} vs. IDH^{wt} comparison without testing for differential activity between TFs. Heatmaps depicting TF activity at hypermethylated enhancers and promoters were generated using a z-score of the activity.

Primary cultures of human microglia

Primary microglial cultures were prepared using surgical aspirates obtained from patients with either brain tumors or drug-resistant epilepsy, following established surgical protocols. For gliomas, tissue collection involved the removal of both the adjacent "entry cortex" (the tissue surrounding the surgical access point) and the tumor core, whereas in cases of epilepsy surgery, only the normal tissue at the entry point was collected. Upon receipt (averaging 2 hours), tissues were promptly processed following the protocol specified for human tumor samples. After enzymatic digestion and MACS purification, CD11B+ cells were counted and directly seeded in non-coated 12-well plates at a density of 750-1000 K cells/well in DMEM/F12, GlutaMAX medium (Gibco Ref# 10565018) supplemented with N-acetyl-cysteine and insulin [5 μg/mL], apo-Transferrin [100 μg/mL], sodium selenite [100 ng/mL], human TGFβ2 [2 ng/mL], CSF-1 [20 ng/mL], ovine wool cholesterol [1.5 μg/mL], ascorbic acid [75 μM] and penicillin/streptomycin[49]. All reagents were purchased from Sigma and growth factors from Preprotech. Non-permeable D-2HG acid disodium salt (Sigma) was used at a final concentration of 5 mM for all the experiments, with treatments starting at 48 hours culture. Half of the culture medium was renewed every 2-3 days. For the 6 hours LPS challenge experiments, 1% FBS was added to all conditions.

IDH^mut and IDH^wt cell lines

Human glioma cell lines IDH^wt were established by the GlioTEx team (Glioblastoma and Experimental Therapeutics) in the Paris Brain Institute (ICM) laboratory. Cultured cells were maintained at 37 °C, 5% CO2 under neurosphere growth conditions using DMEM/F12 medium (Gibco) supplemented with 1% penicillin/streptomycin, B27 diluted 1:50 (Gibco), EGF (20 ng/mL), and FGF (20 ng/mL) (Preprotech). Human glioma cell lines IDH^mut BT88, BT54 were kindly provided by Artee Luchman and Sam Weiss (University of Calgary); BT237, BT138 were kindly provided by Keith Ligon (Dana-Farber Cancer Institute). Cells were grown using DMEM/F12 medium (Gibco) supplemented with B27 (1/100; Gibco), 20ng/ml EGF and FGF (Peprotech). IDH^wt and IDH^mut cell lines were used as negative and positive controls, respectively, in assays applied to quantify intracellular D-2HG levels. The Idh1 R132H

conditional knock-in (cKI) mouse line (Bardella et al., 2016) crossed with *Rosa26^{LSL-YFP}*
(Srinivas et al., 2001) was used to generate postnatal (P0-P1) neural progenitor cells. Cells
were grown in DMEM/F-12 with GlutaMAX™ Supplement (61965-026, ThermoFisher),
Penicillin-Streptomycin 1X (15140122, Life Technologies), B27 supplement 1X (17502048,
ThermoFisher), N2 supplement 1X (17502048, ThermoFisher), 0.6% w/v D-glucose (G8769,
Sigma), 5mM HEPES (15630056, Life Technologies) and 20µg/ mL insulin (I5500, Sigma).
Growth factors (20 ng/ mL EGF GMP100-15, Peprotech, and 10 ng/ mL bFGF GMP100-18B,
Peprotech) were added to the cultures and replaced every 2-3 days. Cells were maintained
in a humidified incubator at 37 °C with 5% CO2. Cultures were infected with a Cre expressing
adenovirus (Ad:cre, VB180308-1398jmt, VectorBuilder) after 1 passage, enabling
recombination at the Idh1 and Rosa26LSL-YFP locus leading to expression of Idh1 R132H
and YFP. Efficient DNA recombination was assessed through YFP visualization in NSC. Cells
were cultured for 3-6 passages and assayed for D-2HG levels using a fluorimetric test, TET
enzymatic activity, and 5mC/5hmC genomic content using LC-MS/MS.

Immunofluorescence staining

Sterile coverslips were placed in 12 multiple wells and the surface was covered with poly-D-
lysine and laminin (Sigma). The plate was incubated at room temperature (RT) for 2 hours
and the solution was aspirated; then, it was rinsed with sterile water and allowed to dry. Cells
were seeded at a density of 750K cells/well and maintained as described for 3, 10 and 17
days. Cells were washed with PBS (Gibco) and fixed with 4% paraformaldehyde (Electron
Microscopy Sciences) in PBS for 10 min at RT. Next, cells were washed with PBS and
permeabilized with 0,1% Triton X-100 in 1x PBS for 5 min at RT. Cells were quickly washed
with PBS and blocked for non-specific binding sites with Human Fc Block (#564219, BD
Pharmingen) for 10 min at RT, and subsequently with 5% donkey in PBS for 30-60 min at RT.
Incubation with primary antibodies was overnight at 4°C using anti-Iba1 (#19-19741, Wako;
diluted 1/800), Tmem119 (#A16075D, BioLegend; diluted 1/100), or Ki-67 (#AB16667-1001,
Abcam; diluted 1/250) in a humidified and light protected chamber. Coverslips were washed
with PBS and incubated with secondary antibodies (anti-rabbit #A21206 or anti-mouse

#A21203; diluted 1/1000) in blocking solution for 30 min at RT and protected from light. Subsequently, the coverslips were washed with PBS and mounted with Fluoroshield-DAPI solution (Sigma) and read under an Apotome inverted microscope (ZEISS Apotome 3) and a confocal microscope (A1R HD25 Nikon Inverted). Negative controls (background fluorescence) were obtained omitting primary antibodies. The percentages of TMEM119+ and IBA1+ cells were determined in primary cultures grown in uncoated wells by counting 3 to 5 fields, covering 0.22% of the well surface, at different time points over a two-week period. The fields were randomly selected in the area where the cell density, as assessed by DAPI, was adequate.

Intracellular quantifications of D-2HG

Cultured cells were pelleted by centrifugation to remove the media, followed by two washes with ice-cold DPBS to ensure complete removal of all D-2HG-containing media. Microglial cells were collected by trypsinization (0.25% for 5 min), whereas glioma cells and neurospheres were collected using accutase. Corresponding culture medium supplemented with 10% FBS was added for inactivation. After centrifugation (300 g at 4°C for 5 min) the resulting pellet was washed with cold DPBS. Cell pellets were then preserved at -80°C until used. The concentration of D-2HG in cell lysates was determined using the D-2-Hydroxyglutarate (D-2HG) Assay Kit (Ref MAK320, Sigma) and with LC/MS for validation. For the fluorometric assay, cellular extracts were prepared by adding 75 µl of CelLytic MT Cell Lysis Reagent (Sigma) with Protease Inhibitor Cocktail (Sigma). An aliquot was kept for protein quantification with the Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific). The collected supernatants were deproteinized by perchloric acid precipitation and neutralized with KOH. A standard curve was prepared with serial dilutions of D-2HG. The equivalent of 5000 pmol was added to a cell extract and used as spike-in internal control. The plate is incubated at 37°C for 30–60 minutes. This assay is based on the oxidation of D-2HG to α-ketoglutarate (αKG) by the enzyme (D)-2-hydroxyglutarate dehydrogenase (HGDH), coupled with the reduction of NAD+ to NADH[83]. The amount of NADH formed is then quantitated by the diaphorase-mediated reduction of resazurin to the fluorescent dye resorufin (λex =

540 nm/λem = 590 nm) using Spectramax i3x microplate reader. The concentration of D-2HG for each sample is estimated as pmol/µg of protein. For the quantification of 2HG using LC-MS, dry cell pellets were lysed in water and the solution was divided in half for 2-HG measurement and protein quantification. NaCl was added and samples were acidified using HCl. A liquid-liquid extraction of organic acids with ethylacetate was performed. Three extractions were performed, organic phases were pooled, dried under nitrogen stream at 30 °C. Then samples were derivated by a standard silylaltion protocol N,O-Bis [trimethylsilyl] trifluoroacetamide and 1% TMCS (Trimethylchlorosilane) under anhydrous conditions using pyridine. Chromatographic separation was performed with a TR-5MS (30m x 0.25 mm x 0.25 mm) column from Thermo Scientific (Waltham, Massachusets, USA). Spectral data acquisitions were performed using XCalibur software (Thermo Electron Corporation, Austin, TX, USA). Sample were placed 30 min at 80°C, and then injected into the GC system. Quantifications were performed using internal standard 2-hydroxyglutaric acid-D3 from Cambridge isotope laboratories (Tewksbury, Massachusetts, USA).

LC-MS/MS assessment of genomic 5mC and 5hmC content
Genomic DNA (gDNA) was isolated as indicated and concentrated by ethanol precipitation. Samples were then digested at 37°C for 12 h in 25 µL reactions containing 1-4 µg of gDNA, 7.5 U of DNA Degradase Plus (Zymo Research), reaction buffer and isotope labeled internal standards [700 pmol 2'-deoxycytidine-15N3 (Cambridge Isotope Laboratories, Inc.); 1.75 pmol 5-methyl-2′-deoxy cytidine-d3 (Toronto Research Chemicals); 1 pmol 5-hydroxymethyl-2′-deoxycytidine-d3 (Toronto Research Chemicals)] for quantification. Prior to LC/MS analysis, samples were diluted 1:1 in mobile phase component A (see below). The LC-MS system consisted of a Shimadzu Nexera UPLC system in-line with a SCIEX 6500 QTrap mass spectrometer equipped with a TurboV Ion spray source and operated in positive ion mode. The analytes were chromatographed on a gradient elution system where mobile phase A was water with 10 mM ammonium formate and 0.05% trifluoroacetic acid (TFA) (Sigma Aldrich); and mobile phase B was 7:1 methanol:acetonitrile (ThermoFisher Scientific), with 10 mM ammonium formate and 0.05% TFA. The column was a Phenomenex Curosil-

PFP column (10 x 0.2 cm 1.7 um) (Phenomenex Inc) held at 42°C. The flow rate was 300 µL/min and the gradient started at % B = 1.0% (held for 30 s) and was then increased to 19% over 5 min. All analytes were detected using selected reaction monitoring (SRM). Analyst software (version 1.6.2) was used to acquire and process the data. Supplementary Table 6 gives the Q1 and Q3 m/z values, collision energy (CE) and declustering potential (DP) for all analytes and their internal standards. Analytes were quantified by stable isotope dilution against their stably labeled internal standard. Data represents a percentage for each analyte relative to the total cytosine pool (dC+5mC+5hmC) in the same sample.

Reduced Representation Enzymatic Methyl sequencing (RREM-seq)

Library preparation was performed using 100 ng of genomic DNA is spiked with 0.95 ng of unmethylated Lambda DNA and 0.05 ng of CpG methylated pUC19. The mixture is digested with 400 units of MspI (New England Bioloabs) in NEB ultra II buffer (New England Biolabs) in a final volume of 57 µL for 2 hours at 37°C. After enzymatic DNA digestion, fragments are repaired and A-tailed using 15 units of Klenow 3'-5' exo- (New England Biolabs). After End-repair and A-tailing, Illumina universal methylated adapters are ligated to DNA fragments using the NEB ultra II ligation module, following the provider's recommendations. After ligation step, DNA fragments are purified and size-selected with a two steps SPRI beads purification in order to select fragments from 40bp to 400bp. Clean-up ligated DNA is eluted in 10 mM Tris pH 8. Purification product is then converted using the Enzymatic methyl-seq module (New England Biolabs) following the provider's recommendations. Briefly, 5mC are converted to 5hmC with TET2, then 5hmC are specifically protected from APOBEC deamination with a T4-BGT glycosylation. After 5hmC/5mC glycosylation, libraries are denatured during 10min at 85°C in 20% Formamide. Then unmodified cytosines are de-aminated during APOBEC reaction. The converted product is amplified by PCR using indexing primers and polymerase KAPA HiFi Hotstart U+ Ready Mix 2X in a final volume of 50 µL for 13 cycles. PCR products are finally purified with a 1.2X SPRI ratio, then controlled by capillary electrophoresis on Fragment Analyzer. After quantification by qPCR libraries are sequenced on an Illumina Novaseq 6000 as paired-end 100b reads.

Reduced Representation hydroxyMethyl sequencing (RREhM-seq)

We used the RREhM-seq protocol published by Sun, Zhiyi et al[52]. with slight modifications. Library preparation was performed using an input of 200 ng with a spike of 1.9 ng of unmethylated Lambda DNA and 0.1 ng of CpG methylated pUC19. Following MspI digestion and reparation with Klenow 3'-5' exo-, pyrrolo-dC Illumina adapters are used instead of 5mC adapters during the ligation step. For the TET2 step during the conversion, TET2 enzyme and Fe2+ are replaced by H_2O to avoid 5mC protection. The product is deaminated, amplified, qualified and sequenced in the same conditions as RREM-seq libraries. RREM-seq and RREhM-seq were performed by Integragen SA (Evry, France).

RREM-seq & RREhM-seq data analysis

FastQ files were aligned using BSseeker2 (https://github.com/BSSeeker/BSseeker2) on hg19 genome with bowtie2. The hg19 index was built with the rrbs option on and selected fragments size between 40bp and 400 bp. The adapters sequence and CCGG motif were removed during the alignment steps. After alignment, methylation calling is performed in order to obtain methylation level and covering for each cytosine positions from the library. Options to remove overlap and bad quality sequences were enabled. For RREM-seq libraries, methylation signal corresponds to 5mC and 5hmC, whereas the methylation signal only concerns 5hmC for RREhM-seq libraries[84]. Levels of total 5mC (methylation and hydroxymethylation) or 5hmC were defined as percentage of the number of reads having the corresponding modification on a specific cytosine divided by the number of reads covering that cytosine (modified or not modified). Methylation-only level was obtained subtracting 5hmC level from the total 5mC level. CpG sites were analyzed with methylKit package[85] using a minimum coverage of 10 reads per base. For the defined window analysis, TFs motifs were retrieved from HOCOMOCO v12 CORE and searched inside entire enhancers regions using Biostrings package[86] with a maximum mismatch of 1 base. The region of interest was defined as a window of 600 bp centered around the first base of the motif. For each patient, only common enhancers windows between treated and control samples were analyzed. Microglia samples used for analyses of 5mC and 5hmC at 2 days (2D) and 14 days

(14D) were obtained from the aspirates of normal tissue from three patients with epilepsy who underwent surgery. One technical outlier from the 5hmC analysis at 14D was excluded from the study. 5mC-only and 5hmC levels are reported as the mean across patients for each time point/condition, using only the common enhancer regions or defined windows shared among patients.

TET enzymatic assay

Nuclei isolation was performed using the nuclear extraction kit (cat. Ab113474, Abcam, Cambridge, MA, USA). Cells were collected by trypsinization (0.25% for 5 min) and counted. Then, cells were resuspended in an appropriate volume of pre-extraction buffer containing dithiothreitol and protease inhibitor cocktail, and incubated on ice for 2 minutes. After centrifugation at 12,000 rpm for 3 minutes, the cytoplasmic extract (supernatant) was carefully removed, leaving behind the nuclear pellet. Two volumes of extraction buffer containing dithiothreitol and protease inhibitor cocktail were added to the nuclear pellet. The extract was incubated on ice for 15 minutes with vortexing for 5 seconds every 3 minutes. After incubation, the suspension was centrifuged for 10 minutes at 14,000 rpm at 4°C, and the resulting supernatant was transferred to a new microcentrifuge vial. Protein quantification from the nuclear extract was performed using a Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific) following the manufacturer's instructions. Technical triplicates for this assay were performed using 12 µg of protein from nuclear extracts of microglial cells (untreated and treated with D-2HG), as well as from external controls (neurospheres obtained from the cKI Idh R132H mouse model). To determine TET enzymatic activity, we employed the TET Hydroxylase Activity kit from Abcam (ab156913). In this fluorimetric assay, a methylated substrate is stably coated onto microplate wells. Active TET enzymes bind to the substrate and convert the methylated substrate into hydroxymethylated products. The hydroxymethylated products generated by TET activity can be recognized with a specific antibody. The ratio or amount of hydroxymethylated products, which is proportional to the enzyme activity, was measured fluorometrically using a fluorescent microplate reader with

excitation at 530 nm and emission at 590 nm. The TET enzyme activity is proportional to the relative fluorescent units measured.

RT-qPCR

Total RNA from cultured microglia was purified using the Maxwell RSC simplyRNA Cells Kit (Promega). Next, 300 ng of RNA was reverse transcribed to complementary DNA using the Maxima First Strand cDNA Synthesis Kit (Thermo Fisher Scientific). Assays were run in triplicate on a Light Cycler 480 instrument (Roche) using the LightCycler 480 SYBR Green Master 2X (Roche). Primer sequences were as follow: TNFA, forward: 5'-GAGCCAGCTCCCTCTATTTA-3', reverse: 5'-GGGAACAGCCTATTGTTCAG-3'; IL-6, forward: 5'-CCTTCCAAAGATGGCTGAAA-3', reverse: 5'-TGGCTTGTTCCTCACTACT-3'; PPIA, forward: 5'-ATGCTGGACCCAACACAAAT-3', reverse: 5'-TCTTTCACTTTGCCAAACACC-3'; ACTB, forward: 5'-CCAACCGCGAGAAGATGA-3', reverse: 5'- CCAGAGGCGTACAGGGATAG-3'. Linearity and efficiency, as well as specific amplifications were verified for all primer sets. Relative quantifications of cytokine expression were normalized to the mean of the two control genes (PPIA and ACTB) and expressed as fold change[87].

Mitochondrial respiration measurement

Real-time measurement of oxygen consumption rate (OCR) was performed using XFe extracellular flux analyzer (Seahorse Bioscience, Billerica, MA) following the manufacturer's instructions. Primary isolated microglia (as detailed in materials and methods) were plated at a density of 1,000,000 cells/well in XFe24 cell culture microplates pre-coated with poly-L-lysine (Sigma). Cells were maintained in 200 µl DMEM/F-12 medium supplemented as previously indicated and treated with 5 mM D-2HG for a period of 12 days. Half of the culture medium was renewed every 2-3 days. The day of the experiment, cells were stimulated with LPS (1µg/mL) for 6 hours. FBS 1% was also added to all experimental conditions. Next, the medium was replaced by Seahorse XF base medium (Agilent, 102353-100) supplemented with 17.5 mM D-Glucose (Dextrose), 0.5 mM sodium pyruvate and 2.5 mM L-Glutamine and incubated for 1 h in CO2 free incubator at 37 °C. The utility plate was hydrated with XF calibrant solution (pH 7.4) (Agilent, 100840-000; 1 mL/well) and incubated overnight (37°C,

CO2-free). The day after, the utility plate with the cartridge containing injector ports and sensors was run on the Seahorse for calibration. The assay medium (Seahorse XF DMEM assay medium, pH 7.4) was prepared immediately before assay. OCR was measured under basal conditions and after sequential injection of the following inhibitors: Oligomycin (1 μM) to stop ATP-synthase, FCCP (1 μM) to dissipate proton gradient, and Rotenone/Antimycin A (0.5 μM each to block electron transport chain (ETC). All the following measurements were carried out with 2 min mix, 2 min delay, and 3 min measure. Three baseline OCR measurements were recorded, followed by assessment of mitochondrial metabolism by injection of oligomycin, FCCP, and a combination of rotenone and antimycin A. The following parameters were deduced: basal respiration (OCR values, used to provide ATP under baseline conditions), ATP-linked respiration (following oligomycin injection, a reduction in OCR values represents the part of basal respiration used to produce ATP), and maximal respiration (the maximal OCR values following FCCP injection). Basal respiration was calculated from the subtraction between the last rate measurement before oligomycin injection and the non-mitochondrial respiration rate (NMRT) determined as the minimum rate measurement after rotenone/antimycin A injection. Maximal respiration was calculated by subtracting the maximum measurement after FCCP injection and the NMRT. ATP-linked respiration was calculated by subtracting the last rate measurement before oligomycin injection and the minimum rate measurement after oligomycin injection. OCR data were normalized by the number of viable cells estimated at the end of the experiment. To do this, the cells were fixed with paraformaldehyde 4%, followed by washes with PBS and DAPI [10 μg/mL] staining. Nuclei were automatically counted in a CellInsight NXT platform.

Single-nucleus RNA sequencing analysis

We downloaded raw data of oligodendroglioma (patient BWH445) from Spitzer et al. study on IDHi[55] GSE260928 and used Seurat (version 5.0.3)[88] to perform data analysis. First, we performed data quality control excluding cells having less than 1500 or more than 3400 genes expressed and less than 2200 or more than 12000 unique molecular identifiers. These stringent thresholds allowed us to have high-quality nuclei also avoiding doublets.

Subsequently, we normalized the entire raw gene count matrix by regularized negative binomial regression using the sctransform function version 2 and clustering cells using 30 principal components. We selected the microglia cluster based on the expression of TAL1, P2RY12, SLC2A5 and the absence of the infiltrating myeloid marker CD163. Selected microglial cells were then analyzed using the raw expression matrix and normalized with the sctransform function version 2 and clustered using 20 principal components. Differential expression analysis was performed using the function FindMarkers.

Statistical analysis and plots generation

Sample sizes and statistical tests are provided in the figure legends. Analyses were performed with R. Normal distribution of data was verified with Shapiro-Wilk's normality test (p-value > 0.05). Plots and analyses were generated using ggplot2[89], the extension ggsignif[90], ComplexHeatmap[91], Cytoscape[92], clusterProfiler[93] and eulerr[94]. Hallmark gene set collection (v2023.1.Hs) for GSEA analyses[95] was downloaded from the Molecular Signatures Database (MSigDB)[96].

Data and code availability

Data and code are available upon request.

Supplementary Tables

Supplementary Table 1: Clinical, pathological and genetic features of the study cohort.

Supplementary Table 2: Genes regulated by hypermethylated enhancers and promoters in CD11B+ cells from IDH^{mut} gliomas.

Supplementary Table 3: Differentially expressed genes in CD11B+ cells from IDH^{mut} vs. IDH^{wt} gliomas.

Supplementary Table 4: Genes regulated by hypermethylated enhancers and/or promoters linked to downregulated expression in CD11B+ cells from IDH^{mut} vs. IDH^{wt} gliomas.

Supplementary Table 5: Genes regulated by the selected enhancers bound by TFs of Fig 5c exhibiting hypermethylation/hypohydroxymethylation in D-2HG-treated microglia.

Supplementary Table 6: Analyte and internal standard Q1 and Q3 *m/z* values, collision energy (CE), and declustering potential (DP) for LC-MS/MS quantifications.

References

1. van den Bent, M. J., Geurts, M., French, P. J., Smits, M., Capper, D., Bromberg, J. E. C., & Chang, S. M. (2023). Primary brain tumours in adults. The Lancet, 402(10412), 1564–1579. https://doi.org/10.1016/S0140-6736(23)01054-1.

2. Dang, L., White, D. W., Gross, S., Bennett, B. D., Bittinger, M. A., Driggers, E. M., Fantin, V. R., Jang, H. G., Jin, S., Keenan, M. C., Marks, K. M., Prins, R. M., Ward, P. S., Yen, K. E., Liau, L. M., Rabinowitz, J. D., Cantley, L. C., Thompson, C. B., vander Heiden, M. G., & Su, S. M. (2009). Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. Nature, 462(7274), 739–744. https://doi.org/10.1038/nature08617.

3. Richard, Q., Laurenge, A., Mallat, M., Sanson, M., & Castro-Vega, L. J. (2022). New insights into the Immune TME of adult-type diffuse gliomas. Current Opinion in Neurology, 35(6), 794–802. https://doi.org/10.1097/WCO.0000000000001112.

4. Linninger, A., Hartung, G. A., Liu, B. P., Mirkov, S., Tangen, K., Lukas, R. v, Unruh, D., James, C. D., Sarkaria, J. N., & Horbinski, C. (2018). Modeling the diffusion of D-2-hydroxyglutarate from IDH1 mutant gliomas in the central nervous system. Neuro-Oncology, 20(9), 1197–1206. https://doi.org/10.1093/neuonc/noy051.

5. Pickard, A. J., Sohn, A. S. W., Bartenstein, T. F., He, S., Zhang, Y., & Gallo, J. M. (2016). Intracerebral Distribution of the Oncometabolite d-2-Hydroxyglutarate in Mice Bearing Mutant Isocitrate Dehydrogenase Brain Tumors: Implications for Tumorigenesis. Frontiers in Oncology, 6. https://doi.org/10.3389/fonc.2016.00211.

6. Bunse, L., Pusch, S., Bunse, T., Sahm, F., Sanghvi, K., Friedrich, M., Alansary, D., Sonner, J. K., Green, E., Deumelandt, K., Kilian, M., Neftel, C., Uhlig, S., Kessler, T., von Landenberg, A., Berghoff, A. S., Marsh, K., Steadman, M., Zhu, D., … Platten, M. (2018). Suppression of antitumor T cell immunity by the oncometabolite (R)-2-hydroxyglutarate. Nature Medicine, 24(8), 1192–1203. https://doi.org/10.1038/s41591-018-0095-6.

7. Böttcher, M., Renner, K., Berger, R., Mentz, K., Thomas, S., Cardenas-Conejo, Z. E., Dettmer, K., Oefner, P. J., Mackensen, A., Kreutz, M., & Mougiakakos, D. (2018). D-2-hydroxyglutarate interferes with HIF-1α stability skewing T-cell metabolism towards oxidative phosphorylation and impairing Th17 polarization. OncoImmunology, 7(7), e1445454. https://doi.org/10.1080/2162402X.2018.1445454.

8. Ugele, I., Cárdenas-Conejo, Z., Hammon, K., Wehrstein, M., Bruss, C., Peter, K., Singer, K., Gottfried, E., Boesch, J., Oefner, P., Dettmer, K., Renner, K., & Kreutz, M. (2019). D-2-Hydroxyglutarate and L-2-Hydroxyglutarate Inhibit IL-12 Secretion by Human Monocyte-Derived Dendritic Cells. International Journal of Molecular Sciences, 20(3), 742. https://doi.org/10.3390/ijms20030742.

9. Friedrich, M., Sankowski, R., Bunse, L., Kilian, M., Green, E., Ramallo Guevara, C., Pusch, S., Poschet, G., Sanghvi, K., Hahn, M., Bunse, T., Münch, P., Gegner, H. M., Sonner, J. K., von Landenberg, A., Cichon, F., Aslan, K., Trobisch, T., Schirmer, L., … Platten, M. (2021). Tryptophan metabolism drives dynamic immunosuppressive myeloid states in IDH-mutant gliomas. Nature Cancer, 2(7), 723–740. https://doi.org/10.1038/s43018-021-00201-z.

10. Notarangelo, G., Spinelli, J. B., Perez, E. M., Baker, G. J., Kurmi, K., Elia, I., Stopka, S. A., Baquer, G., Lin, J.-R., Golby, A. J., Joshi, S., Baron, H. F., Drijvers, J. M., Georgiev, P., Ringel, A. E., Zaganjor, E., McBrayer, S. K., Sorger, P. K., Sharpe, A. H., … Haigis, M. C. (2022). Oncometabolite R-2HG alters T cell metabolism to impair CD8 + T cell function. Science, 377(6614), 1519–1529. https://doi.org/10.1126/science.abj5104.

11. Ruiz-Moreno, C., Marco Salas, S., Samuelsson, E., Brandner, S., Kranendonk, M.E., Nilsson, M., Stunnenberg, H.G. Harmonized single-cell landscape, intercellular crosstalk and tumor architecture of glioblastoma. doi: https://doi.org/10.1101/2022.08.27.505439.

12. Müller, S., Kohanbash, G., Liu, S. J., Alvarado, B., Carrera, D., Bhaduri, A., Watchmaker, P. B., Yagnik, G., di Lullo, E., Malatesta, M., Amankulor, N. M., Kriegstein, A. R., Lim, D. A., Aghi, M., Okada, H., & Diaz, A. (2017). Single-cell profiling of human gliomas reveals macrophage ontogeny as a basis for regional differences in macrophage

activation in the tumor microenvironment. Genome Biology, 18(1), 234. https://doi.org/10.1186/s13059-017-1362-4.

13. Mantovani, A., Allavena, P., Marchesi, F., & Garlanda, C. (2022). Macrophages as tools and targets in cancer therapy. Nature Reviews Drug Discovery, 21(11), 799–820. https://doi.org/10.1038/s41573-022-00520-5.

14. Park, M. D., Silvin, A., Ginhoux, F., & Merad, M. (2022). Macrophages in health and disease. Cell, 185(23), 4259–4279. https://doi.org/10.1016/j.cell.2022.10.007.

15. Amit, I., Winter, D. R., & Jung, S. (2016). The role of the local environment and epigenetics in shaping macrophage identity and their effect on tissue homeostasis. Nature Immunology, 17(1), 18–25. https://doi.org/10.1038/ni.3325.

16. Bowman, R. L., Klemm, F., Akkari, L., Pyonteck, S. M., Sevenich, L., Quail, D. F., Dhara, S., Simpson, K., Gardner, E. E., Iacobuzio-Donahue, C. A., Brennan, C. W., Tabar, V., Gutin, P. H., & Joyce, J. A. (2016). Macrophage Ontogeny Underlies Differences in Tumor-Specific Education in Brain Malignancies. Cell Reports, 17(9), 2445–2459. https://doi.org/10.1016/j.celrep.2016.10.052.

17. Klemm, F., Maas, R. R., Bowman, R. L., Kornete, M., Soukup, K., Nassiri, S., Brouland, J.-P., Iacobuzio-Donahue, C. A., Brennan, C., Tabar, V., Gutin, P. H., Daniel, R. T., Hegi, M. E., & Joyce, J. A. (2020). Interrogation of the Microenvironmental Landscape in Brain Tumors Reveals Disease-Specific Alterations of Immune Cells. Cell, 181(7), 1643-1660.e17. https://doi.org/10.1016/j.cell.2020.05.007.

18. Friebel, E., Kapolou, K., Unger, S., Núñez, N. G., Utz, S., Rushing, E. J., Regli, L., Weller, M., Greter, M., Tugues, S., Neidert, M. C., & Becher, B. (2020). Single-Cell Mapping of Human Brain Cancer Reveals Tumor-Specific Instruction of Tissue-Invading Leukocytes. Cell, 181(7), 1626-1642.e20. https://doi.org/10.1016/j.cell.2020.04.055.

19. Venteicher, A. S., Tirosh, I., Hebert, C., Yizhak, K., Neftel, C., Filbin, M. G., Hovestadt, V., Escalante, L. E., Shaw, M. L., Rodman, C., Gillespie, S. M., Dionne, D., Luo, C. C., Ravichandran, H., Mylvaganam, R., Mount, C., Onozato, M. L., Nahed, B. v., Wakimoto, H., … Suvà, M. L. (2017). Decoupling genetics, lineages, and

microenvironment in IDH-mutant gliomas by single-cell RNA-seq. Science, 355(6332). https://doi.org/10.1126/science.aai8478.

20. Hara, T., Chanoch-Myers, R., Mathewson, N. D., Myskiw, C., Atta, L., Bussema, L., Eichhorn, S. W., Greenwald, A. C., Kinker, G. S., Rodman, C., Gonzalez Castro, L. N., Wakimoto, H., Rozenblatt-Rosen, O., Zhuang, X., Fan, J., Hunter, T., Verma, I. M., Wucherpfennig, K. W., Regev, A., … Tirosh, I. (2021). Interactions between cancer cells and immune cells drive transitions to mesenchymal-like states in glioblastoma. Cancer Cell, 39(6), 779-792.e11. https://doi.org/10.1016/j.ccell.2021.05.002.

21. Blanco-Carmona, E., Narayanan, A., Hernandez, I., Nieto, J. C., Elosua-Bayes, M., Sun, X., Schmidt, C., Pamir, N., Özduman, K., Herold-Mende, C., Pagani, F., Cominelli, M., Taranda, J., Wick, W., von Deimling, A., Poliani, P. L., Rehli, M., Schlesner, M., Heyn, H., & Turcan, Ş. (2023). Tumor heterogeneity and tumor-microglia interactions in primary and recurrent IDH1-mutant gliomas. Cell Reports Medicine, 4(11), 101249. https://doi.org/10.1016/j.xcrm.2023.101249.

22. Gupta, P., Dang, M., Oberai, S., Migliozzi, S., Trivedi, R., Kumar, G., … & Bhat, K. P. (2024). Immune landscape of isocitrate dehydrogenase-stratified primary and recurrent human gliomas. Neuro-Oncology, noae139. doi: https://doi.org/10.1093/neuonc/noae139.

23. Miller, E., El Farran, C.A., Couturier, C.P., Chen, Z., D'Antonio, J.P., Verga, J., Villanueva, M.A., Gonzalez Castro, L.N., Tong, Y.E., Saadi, T.A., Chiocca, A.N., Fischer, D.S., Heiland, D.H., Guerriero, J.L., Petrecca, K., Suva, M.L., Shalek, A.K., Bernstein, B.E. (2023). Programs, Origins, and Niches of Immunomodulatory Myeloid Cells in Gliomas. doi: https://doi.org/10.1101/2023.10.24.563466.

24. Pombo Antunes, A. R., Scheyltjens, I., Lodi, F., Messiaen, J., Antoranz, A., Duerinck, J., Kancheva, D., Martens, L., de Vlaminck, K., van Hove, H., Kjølner Hansen, S. S., Bosisio, F. M., van der Borght, K., de Vleeschouwer, S., Sciot, R., Bouwens, L., Verfaillie, M., Vandamme, N., Vandenbroucke, R. E., … Movahedi, K. (2021). Single-cell profiling of myeloid cells in glioblastoma across species and disease stage

reveals macrophage competition and specialization. Nature Neuroscience, 24(4), 595–610. https://doi.org/10.1038/s41593-020-00789-y.

25. Yeo, A. T., Rawal, S., Delcuze, B., Christofides, A., Atayde, A., Strauss, L., Balaj, L., Rogers, V. A., Uhlmann, E. J., Varma, H., Carter, B. S., Boussiotis, V. A., & Charest, A. (2022). Single-cell RNA sequencing reveals evolution of immune landscape during glioblastoma progression. Nature Immunology, 23(6), 971–984. https://doi.org/10.1038/s41590-022-01215-0.

26. Abdelfattah, N., Kumar, P., Wang, C., Leu, J.-S., Flynn, W. F., Gao, R., Baskin, D. S., Pichumani, K., Ijare, O. B., Wood, S. L., Powell, S. Z., Haviland, D. L., Parker Kerrigan, B. C., Lang, F. F., Prabhu, S. S., Huntoon, K. M., Jiang, W., Kim, B. Y. S., George, J., & Yun, K. (2022). Single-cell analysis of human glioma and immune cells identifies S100A4 as an immunotherapy target. Nature Communications, 13(1), 767. https://doi.org/10.1038/s41467-022-28372-y.

27. Noushmehr, H., Weisenberger, D. J., Diefes, K., Phillips, H. S., Pujara, K., Berman, B. P., Pan, F., Pelloski, C. E., Sulman, E. P., Bhat, K. P., Verhaak, R. G. W., Hoadley, K. A., Hayes, D. N., Perou, C. M., Schmidt, H. K., Ding, L., Wilson, R. K., van den Berg, D., Shen, H., … Aldape, K. (2010). Identification of a CpG Island Methylator Phenotype that Defines a Distinct Subgroup of Glioma. Cancer Cell, 17(5), 510–522. https://doi.org/10.1016/j.ccr.2010.03.017.

28. Chuntova, P., Yamamichi, A., Chen, T., Narayanaswamy, R., Ronseaux, S., Hudson, C., Tron, A. E., Hyer, M. L., Montoya, M., Mende, A. L., Nejo, T., Downey, K. M., Diebold, D., Lu, M., Nicolay, B., & Okada, H. (2022). Inhibition of D-2HG leads to upregulation of a proinflammatory gene signature in a novel HLA-A2/HLA-DR1 transgenic mouse model of IDH1R132H-expressing glioma. Journal for ImmunoTherapy of Cancer, 10(5), e004644. https://doi.org/10.1136/jitc-2022-004644.

29. de Witte, L. D., Wang, Z., Snijders, G. L. J. L., Mendelev, N., Liu, Q., Sneeboer, M. A. M., Boks, M. P. M., Ge, Y., & Haghighi, F. (2022). Contribution of Age, Brain Region, Mood Disorder Pathology, and Interindividual Factors on the Methylome of Human

Microglia. Biological Psychiatry, 91(6), 572–581.

https://doi.org/10.1016/j.biopsych.2021.10.020.

30. Zheng, P.-P., van der Weiden, M., van der Spek, P. J., Vincent, A. J. P. E., & Kros, J. M. (2012). Isocitrate dehydrogenase 1R132H mutation in microglia/macrophages in gliomas. Cancer Biology & Therapy, 13(10), 836–839. https://doi.org/10.4161/cbt.20836.

31. Ceccarelli, M., Barthel, F. P., Malta, T. M., Sabedot, T. S., Salama, S. R., Murray, B. A., Morozova, O., Newton, Y., Radenbaugh, A., Pagnotta, S. M., Anjum, S., Wang, J., Manyam, G., Zoppoli, P., Ling, S., Rao, A. A., Grifford, M., Cherniack, A. D., Zhang, H., … Zmuda, E. (2016). Molecular Profiling Reveals Biologically Discrete Subsets and Pathways of Progression in Diffuse Glioma. Cell, 164(3), 550–563. https://doi.org/10.1016/j.cell.2015.12.028.

32. Klughammer, J., Kiesel, B., Roetzer, T., Fortelny, N., Nemc, A., Nenning, K.-H., Furtner, J., Sheffield, N. C., Datlinger, P., Peter, N., Nowosielski, M., Augustin, M., Mischkulnig, M., Ströbel, T., Alpar, D., Ergüner, B., Senekowitsch, M., Moser, P., Freyschlag, C. F., … Bock, C. (2018). The DNA methylation landscape of glioblastoma disease progression shows extensive heterogeneity in time and space. Nature Medicine, 24(10), 1611–1624. https://doi.org/10.1038/s41591-018-0156-x.

33. Binder, H., Willscher, E., Loeffler-Wirth, H., Hopp, L., Jones, D. T. W., Pfister, S. M., Kreuz, M., Gramatzki, D., Fortenbacher, E., Hentschel, B., Tatagiba, M., Herrlinger, U., Vatter, H., Matschke, J., Westphal, M., Krex, D., Schackert, G., Tonn, J. C., Schlegel, U., … Loeffler, M. (2019). DNA methylation, transcriptome and genetic copy number signatures of diffuse cerebral WHO grade II/III gliomas resolve cancer heterogeneity and development. Acta Neuropathologica Communications, 7(1), 59. https://doi.org/10.1186/s40478-019-0704-8.

34. Andersson, R., Gebhard, C., Miguel-Escalada, I., Hoof, I., Bornholdt, J., Boyd, M., Chen, Y., Zhao, X., Schmidl, C., Suzuki, T., Ntini, E., Arner, E., Valen, E., Li, K., Schwarzfischer, L., Glatz, D., Raithel, J., Lilje, B., Rapin, N., … Sandelin, A. (2014). An

atlas of active enhancers across human cell types and tissues. Nature, 507(7493), 455–461. https://doi.org/10.1038/nature12787.

35. Chaligne, R., Gaiti, F., Silverbush, D., Schiffman, J. S., Weisman, H. R., Kluegel, L., Gritsch, S., Deochand, S. D., Gonzalez Castro, L. N., Richman, A. R., Klughammer, J., Biancalani, T., Muus, C., Sheridan, C., Alonso, A., Izzo, F., Park, J., Rozenblatt-Rosen, O., Regev, A., … Landau, D. A. (2021). Epigenetic encoding, heritability and plasticity of glioma transcriptional cell states. Nature Genetics, 53(10), 1469–1479. https://doi.org/10.1038/s41588-021-00927-7.

36. Stadler, M. B., Murr, R., Burger, L., Ivanek, R., Lienert, F., Schöler, A., Nimwegen, E. van, Wirbelauer, C., Oakeley, E. J., Gaidatzis, D., Tiwari, V. K., & Schübeler, D. (2011). DNA-binding factors shape the mouse methylome at distal regulatory regions. Nature, 480(7378), 490–495. https://doi.org/10.1038/nature10716.

37. Aran, D., Sabato, S., & Hellman, A. (2013). DNA methylation of distal regulatory sites characterizes dysregulation of cancer genes. Genome Biology, 14(3), R21. https://doi.org/10.1186/gb-2013-14-3-r21.

38. Blattler, A., & Farnham, P. J. (2013). Cross-talk between Site-specific Transcription Factors and DNA Methylation States. Journal of Biological Chemistry, 288(48), 34287–34294. https://doi.org/10.1074/jbc.R113.512517.

39. Gosselin, D., Skola, D., Coufal, N. G., Holtman, I. R., Schlachetzki, J. C. M., Sajti, E., Jaeger, B. N., O'Connor, C., Fitzpatrick, C., Pasillas, M. P., Pena, M., Adair, A., Gonda, D. D., Levy, M. L., Ransohoff, R. M., Gage, F. H., & Glass, C. K. (2017). An environment-dependent transcriptional network specifies human microglia identity. Science, 356(6344). https://doi.org/10.1126/science.aal3222.

40. Court, F., le Boiteux, E., Fogli, A., Müller-Barthélémy, M., Vaurs-Barrière, C., Chautard, E., Pereira, B., Biau, J., Kemeny, J.-L., Khalil, T., Karayan-Tapon, L., Verrelle, P., & Arnaud, P. (2019). Transcriptional alterations in glioma result primarily from DNA methylation–independent mechanisms. Genome Research, 29(10), 1605–1621. https://doi.org/10.1101/gr.249219.119.

41. Müller-Dott, S., Tsirvouli, E., Vazquez, M., Ramirez Flores, R. O., Badia-i-Mompel, P., Fallegger, R., Türei, D., Lægreid, A., & Saez-Rodriguez, J. (2023). Expanding the coverage of regulons from high-confidence prior knowledge for accurate estimation of transcription factor activities. Nucleic Acids Research, 51(20), 10934–10949. https://doi.org/10.1093/nar/gkad841.

42. Frattini, V., Pagnotta, S. M., Tala, Fan, J. J., Russo, M. v., Lee, S. B., Garofano, L., Zhang, J., Shi, P., Lewis, G., Sanson, H., Frederick, V., Castano, A. M., Cerulo, L., Rolland, D. C. M., Mall, R., Mokhtari, K., Elenitoba-Johnson, K. S. J., Sanson, M., … Iavarone, A. (2018). A metabolic function of FGFR3-TACC3 gene fusions in cancer. Nature, 553(7687), 222–227. https://doi.org/10.1038/nature25171.

43. Yin, Y., Morgunova, E., Jolma, A., Kaasinen, E., Sahu, B., Khund-Sayeed, S., Das, P. K., Kivioja, T., Dave, K., Zhong, F., Nitta, K. R., Taipale, M., Popov, A., Ginno, P. A., Domcke, S., Yan, J., Schübeler, D., Vinson, C., & Taipale, J. (2017). Impact of cytosine methylation on DNA binding specificities of human transcription factors. Science, 356(6337). https://doi.org/10.1126/science.aaj2239.

44. Suzuki, T., Maeda, S., Furuhata, E., Shimizu, Y., Nishimura, H., Kishima, M., & Suzuki, H. (2017). A screening system to identify transcription factors that induce binding site-directed DNA demethylation. Epigenetics & Chromatin, 10(1), 60. https://doi.org/10.1186/s13072-017-0169-6.

45. Yang, J., Zhang, X., Blumenthal, R. M., & Cheng, X. (2020). Detection of DNA Modifications by Sequence-Specific Transcription Factors. Journal of Molecular Biology, 432(6), 1661–1673. https://doi.org/10.1016/j.jmb.2019.09.013.

46. Spruijt, C. G., Gnerlich, F., Smits, A. H., Pfaffeneder, T., Jansen, P. W. T. C., Bauer, C., Münzel, M., Wagner, M., Müller, M., Khan, F., Eberl, H. C., Mensinga, A., Brinkman, A. B., Lephikov, K., Müller, U., Walter, J., Boelens, R., van Ingen, H., Leonhardt, H., … Vermeulen, M. (2013). Dynamic Readers for 5-(Hydroxy)Methylcytosine and Its Oxidized Derivatives. Cell, 152(5), 1146–1159. https://doi.org/10.1016/j.cell.2013.02.004

47. Wu, X., & Zhang, Y. (2017). TET-mediated active DNA demethylation: mechanism, function and beyond. Nature Reviews Genetics, 18(9), 517–534. https://doi.org/10.1038/nrg.2017.33.

48. Turcan, S., Rohle, D., Goenka, A., Walsh, L. A., Fang, F., Yilmaz, E., Campos, C., Fabius, A. W. M., Lu, C., Ward, P. S., Thompson, C. B., Kaufman, A., Guryanova, O., Levine, R., Heguy, A., Viale, A., Morris, L. G. T., Huse, J. T., Mellinghoff, I. K., & Chan, T. A. (2012). IDH1 mutation is sufficient to establish the glioma hypermethylator phenotype. Nature, 483(7390), 479–483. https://doi.org/10.1038/nature10866.

49. Tewari, M., Khan, M., Verma, M., Coppens, J., Kemp, J. M., Bucholz, R., Mercier, P., & Egan, T. M. (2021). Physiology of Cultured Human Microglia Maintained in a Defined Culture Medium. ImmunoHorizons, 5(4), 257–272. https://doi.org/10.4049/immunohorizons.2000101.

50. Pidsley, R., Zotenko, E., Peters, T. J., Lawrence, M. G., Risbridger, G. P., Molloy, P., van Djik, S., Muhlhausler, B., Stirzaker, C., & Clark, S. J. (2016). Critical evaluation of the Illumina MethylationEPIC BeadChip microarray for whole-genome DNA methylation profiling. Genome Biology, 17(1), 208. https://doi.org/10.1186/s13059-016-1066-1.

51. Ginno, P. A., Gaidatzis, D., Feldmann, A., Hoerner, L., Imanci, D., Burger, L., Zilbermann, F., Peters, A. H. F. M., Edenhofer, F., Smallwood, S. A., Krebs, A. R., & Schübeler, D. (2020). A genome-scale map of DNA methylation turnover identifies site-specific dependencies of DNMT and TET activity. Nature Communications, 11(1), 2680. https://doi.org/10.1038/s41467-020-16354-x.

52. Sun, Z., Vaisvila, R., Hussong, L.-M., Yan, B., Baum, C., Saleh, L., Samaranayake, M., Guan, S., Dai, N., Corrêa, I. R., Pradhan, S., Davis, T. B., Evans, T. C., & Ettwiller, L. M. (2021). Nondestructive enzymatic deamination enables single-molecule long-read amplicon sequencing for the determination of 5-methylcytosine and 5-hydroxymethylcytosine at single-base resolution. Genome Research, *31*(2), 291–300. https://doi.org/10.1101/gr.265306.120.

53. Yu, M., Hon, G. C., Szulwach, K. E., Song, C.-X., Zhang, L., Kim, A., Li, X., Dai, Q., Shen, Y., Park, B., Min, J.-H., Jin, P., Ren, B., & He, C. (2012). Base-Resolution Analysis of 5-

Hydroxymethylcytosine in the Mammalian Genome. Cell, 149(6), 1368–1380. https://doi.org/10.1016/j.cell.2012.04.027.

54. Kaikkonen, M. U., Spann, N. J., Heinz, S., Romanoski, C. E., Allison, K. A., Stender, J. D., Chun, H. B., Tough, D. F., Prinjha, R. K., Benner, C., & Glass, C. K. (2013). Remodeling of the Enhancer Landscape during Macrophage Activation Is Coupled to Enhancer Transcription. Molecular Cell, 51(3), 310–325. https://doi.org/10.1016/j.molcel.2013.07.010.

55. Spitzer, A., Gritsch, S., Nomura, M., Jucht, A., Fortin, J., Raviram, R., Weisman, H. R., Gonzalez Castro, L. N., Druck, N., Chanoch-Myers, R., Lee, J. J. Y., Mylvaganam, R., Lee Servis, R., Fung, J. M., Lee, C. K., Nagashima, H., Miller, J. J., Arrillaga-Romany, I., Louis, D. N., ... Tirosh, I. (2024). Mutant IDH inhibitors induce lineage differentiation in IDH-mutant oligodendroglioma. Cancer Cell, 42(5), 904-914.e9. https://doi.org/10.1016/j.ccell.2024.03.008.

56. Lund, H., Pieber, M., Parsa, R., Han, J., Grommisch, D., Ewing, E., Kular, L., Needhamsen, M., Espinosa, A., Nilsson, E., Överby, A. K., Butovsky, O., Jagodic, M., Zhang, X.-M., & Harris, R. A. (2018). Competitive repopulation of an empty microglial niche yields functionally distinct subsets of microglia-like cells. Nature Communications, 9(1), 4845. https://doi.org/10.1038/s41467-018-07295-7.

57. Keshet, I., Lieman-Hurwitz, J., & Cedar, H. (1986). DNA methylation affects the formation of active chromatin. Cell, 44(4), 535–543. https://doi.org/10.1016/0092-8674(86)90263-1.

58. Lee, J. Y., & Lee, T.-H. (2012). Effects of DNA Methylation on the Structure of Nucleosomes. Journal of the American Chemical Society, 134(1), 173–175. https://doi.org/10.1021/ja210273w.

59. Rao, S., Chiu, T.-P., Kribelbauer, J. F., Mann, R. S., Bussemaker, H. J., & Rohs, R. (2018). Systematic prediction of DNA shape changes due to CpG methylation explains epigenetic effects on protein–DNA binding. Epigenetics & Chromatin, 11(1), 6. https://doi.org/10.1186/s13072-018-0174-4.

60. Kreibich, E., Kleinendorst, R., Barzaghi, G., Kaspar, S., & Krebs, A. R. (2023). Single-molecule footprinting identifies context-dependent regulation of enhancers by DNA methylation. Molecular Cell, 83(5), 787-802.e9. https://doi.org/10.1016/j.molcel.2023.01.017.

61. Luoto, S., Hermelo, I., Vuorinen, E. M., Hannus, P., Kesseli, J., Nykter, M., & Granberg, K. J. (2018). Computational Characterization of Suppressive Immune Microenvironments in Glioblastoma. Cancer Research, 78(19), 5574–5585. https://doi.org/10.1158/0008-5472.CAN-17-3714.

62. Nott, A., Holtman, I. R., Coufal, N. G., Schlachetzki, J. C. M., Yu, M., Hu, R., Han, C. Z., Pena, M., Xiao, J., Wu, Y., Keulen, Z., Pasillas, M. P., O'Connor, C., Nickl, C. K., Schafer, S. T., Shen, Z., Rissman, R. A., Brewer, J. B., Gosselin, D., … Glass, C. K. (2019). Brain cell type–specific enhancer–promoter interactome maps and disease - risk association. Science, 366(6469), 1134–1139. https://doi.org/10.1126/science.aay0793.

63. Cheng, S.-C., Quintin, J., Cramer, R. A., Shepardson, K. M., Saeed, S., Kumar, V., Giamarellos-Bourboulis, E. J., Martens, J. H. A., Rao, N. A., Aghajanirefah, A., Manjeri, G. R., Li, Y., Ifrim, D. C., Arts, R. J. W., van der Veer, B. M. J. W., Deen, P. M. T., Logie, C., O'Neill, L. A., Willems, P., … Netea, M. G. (2014). mTOR- and HIF-1α–mediated aerobic glycolysis as metabolic basis for trained immunity. Science, 345(6204). https://doi.org/10.1126/science.1250684.

64. Palsson-McDermott, E. M., Curtis, A. M., Goel, G., Lauterbach, M. A. R., Sheedy, F. J., Gleeson, L. E., van den Bosch, M. W. M., Quinn, S. R., Domingo-Fernandez, R., Johnston, D. G. W., Jiang, J., Israelsen, W. J., Keane, J., Thomas, C., Clish, C., Vander Heiden, M., Xavier, R. J., & O'Neill, L. A. J. (2015). Pyruvate Kinase M2 Regulates Hif-1α Activity and IL-1β Induction and Is a Critical Determinant of the Warburg Effect in LPS-Activated Macrophages. Cell Metabolism, 21(1), 65–80. https://doi.org/10.1016/j.cmet.2014.12.005.

65. Corcoran, S. E., & O'Neill, L. A. J. (2016). HIF1α and metabolic reprogramming in inflammation. Journal of Clinical Investigation, 126(10), 3699–3707. https://doi.org/10.1172/JCI84431.

66. Sangineto, M., Ciarnelli, M., Cassano, T., Radesco, A., Moola, A., Bukke, V. N., Romano, A., Villani, R., Kanwal, H., Capitanio, N., Duda, L., Avolio, C., & Serviddio, G. (2023). Metabolic reprogramming in inflammatory microglia indicates a potential way of targeting inflammation in Alzheimer's disease. Redox Biology, 66, 102846. https://doi.org/10.1016/j.redox.2023.102846.

67. Chesnelong, C., Chaumeil, M. M., Blough, M. D., Al-Najjar, M., Stechishin, O. D., Chan, J. A., Pieper, R. O., Ronen, S. M., Weiss, S., Luchman, H. A., & Cairncross, J. G. (2014). Lactate dehydrogenase A silencing in IDH mutant gliomas. Neuro-Oncology, 16(5), 686–695. https://doi.org/10.1093/neuonc/not243.

68. Koivunen, P., Lee, S., Duncan, C. G., Lopez, G., Lu, G., Ramkissoon, S., Losman, J. A., Joensuu, P., Bergmann, U., Gross, S., Travins, J., Weiss, S., Looper, R., Ligon, K. L., Verhaak, R. G. W., Yan, H., & Kaelin Jr, W. G. (2012). Transformation by the (R)-enantiomer of 2-hydroxyglutarate linked to EGLN activation. Nature, 483(7390), 484–488. https://doi.org/10.1038/nature10898

69. Lavin, Y., Winter, D., Blecher-Gonen, R., David, E., Keren-Shaul, H., Merad, M., Jung, S., & Amit, I. (2014). Tissue-Resident Macrophage Enhancer Landscapes Are Shaped by the Local Microenvironment. Cell, 159(6), 1312–1326. https://doi.org/10.1016/j.cell.2014.11.018.

70. Gosselin, D., Link, V. M., Romanoski, C. E., Fonseca, G. J., Eichenfield, D. Z., Spann, N. J., Stender, J. D., Chun, H. B., Garner, H., Geissmann, F., & Glass, C. K. (2014). Environment Drives Selection and Function of Enhancers Controlling Tissue-Specific Macrophage Identities. Cell, 159(6), 1327–1340. https://doi.org/10.1016/j.cell.2014.11.023.

71. Troutman, T. D., Kofman, E., & Glass, C. K. (2021). Exploiting dynamic enhancer landscapes to decode macrophage and microglia phenotypes in health and disease. Molecular Cell, 81(19), 3888–3903. https://doi.org/10.1016/j.molcel.2021.08.004.

72. Parry, A., Rulands, S., & Reik, W. (2021). Active turnover of DNA methylation during cell fate decisions. Nature Reviews Genetics, 22(1), 59–66. https://doi.org/10.1038/s41576-020-00287-8.

73. Onodera, A., González-Avalos, E., Lio, C.-W. J., Georges, R. O., Bellacosa, A., Nakayama, T., & Rao, A. (2021). Roles of TET and TDG in DNA demethylation in proliferating and non-proliferating immune cells. Genome Biology, 22(1), 186. https://doi.org/10.1186/s13059-021-02384-1.

74. Vats, D., Mukundan, L., Odegaard, J. I., Zhang, L., Smith, K. L., Morel, C. R., Greaves, D. R., Murray, P. J., & Chawla, A. (2006). Oxidative metabolism and PGC-1β attenuate macrophage-mediated inflammation. Cell Metabolism, 4(1), 13–24. https://doi.org/10.1016/j.cmet.2006.05.011.

75. Huang, S. C.-C., Everts, B., Ivanova, Y., O'Sullivan, D., Nascimento, M., Smith, A. M., Beatty, W., Love-Gregory, L., Lam, W. Y., O'Neill, C. M., Yan, C., Du, H., Abumrad, N. A., Urban, J. F., Artyomov, M. N., Pearce, E. L., & Pearce, E. J. (2014). Cell-intrinsic lysosomal lipolysis is essential for alternative activation of macrophages. Nature Immunology, 15(9), 846–855. https://doi.org/10.1038/ni.2956.

76. Sabogal-Guáqueta, A. M., Marmolejo-Garza, A., Trombetta-Lima, M., Oun, A., Hunneman, J., Chen, T., Koistinaho, J., Lehtonen, S., Kortholt, A., Wolters, J. C., Bakker, B. M., Eggen, B. J. L., Boddeke, E., & Dolga, A. (2023). Species-specific metabolic reprogramming in human and mouse microglia during inflammatory pathway induction. Nature Communications, 14(1), 6454. https://doi.org/10.1038/s41467-023-42096-7.

77. Han, C., Zheng, J., Sun, L., Yang, H., Cao, Z., Zhang, X., Zheng, L., & Zhen, X. (2019). The oncometabolite 2-hydroxyglutarate inhibits microglial activation via the AMPK/mTOR/NF-κB pathway. Acta Pharmacologica Sinica, 40(10), 1292–1302. https://doi.org/10.1038/s41401-019-0225-9.

78. Mellinghoff, I. K., Lu, M., Wen, P. Y., Taylor, J. W., Maher, E. A., Arrillaga-Romany, I., Peters, K. B., Ellingson, B. M., Rosenblum, M. K., Chun, S., Le, K., Tassinari, A., Choe, S., Toubouti, Y., Schoenfeld, S., Pandya, S. S., Hassan, I., Steelman, L., Clarke, J. L., &

Cloughesy, T. F. (2023). Vorasidenib and ivosidenib in IDH1-mutant low-grade glioma: a randomized, perioperative phase 1 trial. Nature Medicine, 29(3), 615–622. https://doi.org/10.1038/s41591-022-02141-2.

79. di Stefano, A. L., Picca, A., Saragoussi, E., Bielle, F., Ducray, F., Villa, C., Eoli, M., Paterra, R., Bellu, L., Mathon, B., Capelle, L., Bourg, V., Gloaguen, A., Philippe, C., Frouin, V., Schmitt, Y., Lerond, J., Leclerc, J., Lasorella, A., … Sanson, M. (2020). Clinical, molecular, and radiomic profile of gliomas with FGFR3-TACC3 fusions. Neuro-Oncology, 22(11), 1614–1624. https://doi.org/10.1093/neuonc/noaa121.

80. Teschendorff, A. E. (2020). A comparison of epigenetic mitotic-like clocks for cancer risk prediction. Genome Medicine, 12(1), 56. https://doi.org/10.1186/s13073-020-00752-3.

81. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh, H., & Glass, C. K. (2010). Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. Molecular Cell, 38(4), 576–589. https://doi.org/10.1016/j.molcel.2010.05.004.

82. Vorontsov, I. E., Eliseeva, I. A., Zinkevich, A., Nikonov, M., Abramov, S., Boytsov, A., Kamenets, V., Kasianova, A., Kolmykov, S., Yevshin, I. S., Favorov, A., Medvedeva, Y. A., Jolma, A., Kolpakov, F., Makeev, V. J., & Kulakovskiy, I. V. (2024). HOCOMOCO in 2024: a rebuild of the curated collection of binding models for human and mouse transcription factors. Nucleic Acids Research, 52(D1), D154–D163. https://doi.org/10.1093/nar/gkad1077.

83. Balss, J., Pusch, S., Beck, A.-C., Herold-Mende, C., Krämer, A., Thiede, C., Buckel, W., Langhans, C.-D., Okun, J. G., & von Deimling, A. (2012). Enzymatic assay for quantitative analysis of (d)-2-hydroxyglutarate. Acta Neuropathologica, 124(6), 883–891. https://doi.org/10.1007/s00401-012-1060-y.

84. Ewing, A. D., Smits, N., Sanchez-Luque, F. J., Faivre, J., Brennan, P. M., Richardson, S. R., Cheetham, S. W., & Faulkner, G. J. (2020). Nanopore Sequencing Enables

Comprehensive Transposable Element Epigenomic Profiling. Molecular Cell, 80(5), 915-928.e5. https://doi.org/10.1016/j.molcel.2020.10.024.

85. Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., & Mason, C. E. (2012). methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. Genome Biology, 13(10), R87. https://doi.org/10.1186/gb-2012-13-10-r87

86. Pagès H, Aboyoun P, Gentleman R, DebRoy S (2024). Biostrings: Efficient manipulation of biological strings. R package version 2.72.0, https://bioconductor.org/packages/Biostrings

87. Schmittgen, T. D., & Livak, K. J. (2008). Analyzing real-time PCR data by the comparative CT method. Nature Protocols, 3(6), 1101–1108. https://doi.org/10.1038/nprot.2008.73.

88. Hao, Y., Stuart, T., Kowalski, M. H., Choudhary, S., Hoffman, P., Hartman, A., Srivastava, A., Molla, G., Madad, S., Fernandez-Granda, C., & Satija, R. (2024). Dictionary learning for integrative, multimodal and scalable single-cell analysis. Nature Biotechnology, 42(2), 293–304. https://doi.org/10.1038/s41587-023-01767-y.

89. Wickham, H., & Wickham, H. (2016). Data analysis (pp. 189-201). Springer International Publishing. doi.org/10.1007/978-3-319-24277-4_9.

90. Ahlmann-Eltze, C., & Patil, I. (2021). ggsignif: R Package for Displaying Significance Brackets for 'ggplot2'. PsyArxiv. doi.org/10.31234/osf.io/7awm6.

91. Gu, Z. (2022). Complex Heatmap Visualization, iMeta. doi.org/10.1002/imt2.43.

92. Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., ... & Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome research, 13(11), 2498-2504. doi.org/10.1101/gr.1239303.

93. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., ... & Yu, G. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The innovation, 2(3). doi.org/10.1016/j.xinn.2021.100141.

94. Larsson J. (2024). eulerr: Area-Proportional Euler and Venn Diagrams with Ellipses. R package version 7.0.0, https://CRAN.R-project.org/package=eulerr.

95. Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., ... & Mesirov, J. P. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proceedings of the National Academy of Sciences, 102(43), 15545-15550. doi.org/10.1073/pnas.0506580102.

96. Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J. P., & Tamayo, P. (2015). The molecular signatures database hallmark gene set collection. Cell systems, 1(6), 417-425. doi.org/10.1016/j.cels.2015.12.004.

97. Khanna, A., Larson, D., Srivatsan, S., Mosior, M., Abbott, T., Kiwala, S., ... & Miller, C. (2022). Bam-readcount - rapid generation of basepair-resolution sequence metrics. The Journal of Open Source Software, 7(69), 3722. doi.org/10.21105/joss.03722.

98. Aaronson, J., Beaumont, V., Blevins, R. A., Andreeva, V., Murasheva, I., Shneyderman, A., ... & Vogt, T. F. (2021). HDinHD: A rich data portal for Huntington's disease research. Journal of Huntington's Disease, 10(3), 405-412. doi.org/10.3233/JHD-210491.

Extended data

Extended Data Fig. 1. Purity of CD11B+ myeloid fractions isolated from human gliomas. (A) Digital droplet PCR (ddPCR) performed with specific assays to evaluate IDH1 R132H, TERT promoter mutations C228T/C250T, and corresponding wildtype alleles in available DNAs from CD11B+ samples (n=12) isolated from human gliomas with known

mutation status. Less than 5% copies/µL of the indicated mutations relative to their corresponding wildtype alleles were detected in the same reaction for each analyzed sample. Positive and negative controls from bulk tumors for each assay are also shown. The detection of IDHmut and TERT mutations in positive controls differs, possibly reflecting regional clonal heterogeneity. **(B)** Expression frequency of IDH mutated alleles was assessed in RNA-seq data from CD11B+ samples (n=20) isolated from human gliomas with known mutation status using bam-readcount[97]. Compared to ddPCR, this analysis slightly overestimated the contamination in samples P9 and P10.

Extended Data Fig. 2. CD11B- fractions enriched of glioma cells display global DNA hypermethylation in IDH^{mut} gliomas. (A) Density plot of DNA methylation levels (β-values) in glioma cells-enriched CD11B- fractions isolated from IDH^{mut} (n=13) and IDH^{wt} (n=4) tumors. **(B)** Principal component analysis performed using methylome data from around 600,000 CpG sites across the genome in CD11B- samples from the study cohort. **(C)** Stacked bar charts show absolute numbers and relative percentages of CpG sites and of four functional genomic regions respect to the methylation status in the CD11B- samples from IDH^{mut} vs. IDH^{wt} tumors. **(D)** Volcano plots depict the magnitude and extent of differentially methylated changes (colored red and blue) located at enhancers and promoters in CD11B- samples from IDH^{mut} vs. IDH^{wt} tumors.

A

B

Downregulated genes in TAMs (*IDH*-mut vs. *IDH*-wt)

C

Downregulated genes (N = 1404) in CD11B+ cells (*IDH*-mut vs. *IDH*-wt)

D

Downregulated genes (N = 1404) in CD11B+ cells (*IDH*-mut vs. *IDH*-wt)

Extended Data Fig. 3. Genes with reduced expression in CD11B+ cells from IDH^{mut} **human gliomas belong to microglial signatures. (A)** Principal component analysis performed using RNA-seq data from *ex vivo* CD11B+ myeloid cells isolated from IDH^{mut} (n=20) and IDH^{wt} (n=8) tumors. **(B)** Venn diagrams illustrate significant overlaps between downregulated genes displayed by CD11B+ cells fraction from IDH^{mut} gliomas (this study), and downregulated genes displayed by MDMs (left) or microglial cells (right) in Klemm et al. study. P-values were computed using overrepresentation analysis (ORA). **(C)** Enrichment analysis of the downregulated genes in CD11B+ cells fraction from IDH^{mut} gliomas in the present study using HDinHD microglia signatures dataset[98]. **(D)** Enrichment analysis of the downregulated genes in CD11B+ cells fraction from IDH^{mut} gliomas (this study) using hallmark signatures dataset.

Extended Data Fig. 4. Transcription factors activity controlling genes with hypermethylated promoter in CD11B+ cells from human gliomas. The heatmap depicts TFs (n=26) exhibiting differential activity (FDR <0.05; Wilcoxon's test) in CD11B+ cells according with the IDH mutation status of corresponding tumors. The TF targets used for this analysis, retrieved from the CollecTRI database, were searched among the genes displaying hypermethylation at promoters in CD11B+ cells fraction from IDH^{mut} vs. IDH^{wt} comparison (Fig. 1D). The number of targets regulated positively and negatively by each TF is shown in brackets. Unsupervised clustering was performed using the z-scores of TF activity, where red and blue colors indicate high and low activity, respectively. Non-significant activity is shown in gray.

Extended Data Fig. 5. Positive targets of HIF1A/MYC display low expression in microglia and MDMs from IDH^{mut} gliomas. Box plots showing normalized RNA-seq expression data for representative targets of HIF1A/MYC in CD11B+ myeloid cells from the present study cohort (IDH^{mut}, n=20 vs. IDH^{wt}, n=8), as well as in myeloid cells from the Klemm et al. study comprising MDMs (IDH^{mut}, n=4 vs. IDH^{wt}, n=11) and microglia (IDH^{mut}, n=6 vs. IDH^{wt}, n=13) used as validation set. P-values were computed using Wilcoxon's test.

Extended Data Fig. 6. LC-MS/MS analysis of 5mC and 5hmC genomic content and cell divisions estimate in CD11B+ myeloid cells from IDH^{mut} gliomas and normal brain tissues. **(A)** Schematic representation of α-KG-dependent enzymatic oxidation of 5mC by TET2 and effect of competitive inhibition of TET2 by D-2HG over time in bulk IDH^{mut} gliomas. **(B)** Global levels of 5mC and 5hmC (left) and 5mC/5hmC ratios (right) were measured in CD11B- (upper) and CD11B+ (bottom) fractions from IDH^{mut} gliomas (n=4) and from normal tissues collected during epilepsy surgery (n=5), which served as controls for comparisons. Data are represented as the mean with a 95% CI; p-values were calculated using one-sided Welch's test except for comparison of ratios in CD11B- (one-sided Wilcoxon's test). **(C)** Epigenetic mitotic clock analysis was conducted by applying the epiTOC2[80] model to methylome data (EPIC arrays) of CD11B+ fractions from IDH^{mut} gliomas (n=19) and from normal brain tissues (n=6). The model (adjusted by age) estimates the cumulative number of stem cell divisions in the myeloid compartment. The p-value was calculated using Welch's test.

A

Gosselin et al.–6H	P45–14D	P47–14D	P51–14D	P44–14D	P48–14D	P46–14D	P51–2D	P44–2D	P49–14D	P50–14D	P45–2D	P47–2D	P48–2D	P50–2D	P46–2D	P49–2D
10.9	14.1	17.4	19.5	19.6	20.3	21.4	21.5	21.8	23.2	23.4	26.5	27.8	27.8	29.1	32.2	32.7

Microglia score
30
25
20
15

Microglia sample
Epilepsy
IDH–mut
IDH–wt
Normal

Sex
F
M

B

6H LPS challenge of microglia at day 2

log–fold–change vs Average log CPM
- NotSig
- Up
- Down

6H LPS challenge of microglia at day 14

log–fold–change vs Average log CPM
- NotSig
- Up
- Down

C

Left (day 2):
INTERFERON GAMMA RESPONSE
INFLAMMATORY RESPONSE
TNFA SIGNALING VIA NFKB
INTERFERON ALPHA RESPONSE
IL6 JAK STAT3 SIGNALING
ALLOGRAFT REJECTION
EPITHELIAL MESENCHYMAL TRANSITION
KRAS SIGNALING UP
COMPLEMENT
IL2 STAT5 SIGNALING
APOPTOSIS
HYPOXIA
KRAS SIGNALING DN
COAGULATION
UV RESPONSE UP
P53 PATHWAY
MITOTIC SPINDLE
MYC TARGETS V2
E2F TARGETS

FDR
0.01
0.02
0.03
0.04

Normalized enrichment score

Right (day 14):
TNFA SIGNALING VIA NFKB
INTERFERON GAMMA RESPONSE
INFLAMMATORY RESPONSE
INTERFERON ALPHA RESPONSE
IL6 JAK STAT3 SIGNALING
ALLOGRAFT REJECTION
IL2 STAT5 SIGNALING
KRAS SIGNALING UP
EPITHELIAL MESENCHYMAL TRANSITION
COMPLEMENT
APOPTOSIS
KRAS SIGNALING DN
UV RESPONSE UP
COAGULATION
HYPOXIA
ANDROGEN RESPONSE
XENOBIOTIC METABOLISM
P53 PATHWAY
E2F TARGETS
MITOTIC SPINDLE

FDR
0.01
0.02
0.03
0.04

Normalized enrichment score

Extended Data Fig. 7. Assessment of the functionality of primary microglial cultures. **(A)** Analysis of a human microglia signature at 2 days (2D) and 14 days (14D) using RNA-seq data from eight independent primary cultures. A microglial score was determined as follows: we took the microglia signature genes and the raw gene expression matrix from Gosselin et al[39]. Subsequently, after normalizing the expression data, we calculated the median of the differences of each gene from the microglia signature in our samples with respect to the five negative samples used as controls in Gosselin et al. study. Microglia score was defined as the average of these five medians. The plot (upper), showing such microglia score per sample, demonstrates that the signature is nearly twice as high, even at 14 days of culture, when using defined culture conditions (as in our study), in comparison to the study by Gosselin et al. study (culture shock at 6 hours), irrespective of tissue origin and sex (as indicated at the bottom). **(B)** The plots illustrate expression changes using four primary microglial cultures analyzed with RNA-seq in a paired manner, 6 hours after LPS stimulation at 2 days (left) and 14 days (right). Red and blue colors indicate upregulation and downregulation, respectively, with FDR <0.05, in LPS-stimulated microglia vs. control. **(C)** Ridge plots illustrate hallmark gene sets GSEA analyses in LPS-stimulated microglia at 2 days (left) and 14 days (right) of culture. Activated (>0) and inactivated (<0) gene sets with FDR < 0.05 are shown.

A NSC *Idh*-wt: *Idh1*$^{wt/wt}$; Rosa26$^{LSL-YFP}$
NSC *Idh*-mut: *Idh1*$^{fl(R132H)/wt}$; Rosa26$^{LSL-YFP}$

P0   P1   P3   P6

P1 Ad:Cre

D2HG intracellular levels
TET enzymatic activity
5hmdC and 5mdC levels

B

C

D

Extended Data Fig. 8. Assessment of D-2HG levels and TET function in external controls. (A) Workflow procedure for inducing the *in vitro* expression of Idh1 R132H in neural progenitor cells (NSCs) obtained from a conditional knock-in IDH^{mut} mouse model (cKI Idh1), used as external isogenic controls in three assays. Cells collected at day 14 for analyses underwent 3-6 passages. **(B)** Intracellular levels of D-2HG were determined using a fluorimetric assay in primary microglia treated with D-2HG [5 mM] for 48 hours. Microglia were obtained from samples with IDH^{wt} status (n=3). External controls for comparison (cKI model Idhwt n=5; Idhmut n=4, and patient-derived glioma cells IDH^{wt} n=4; IDH^{mut} n=3) are also shown. The levels of D-2HG in each sample were normalized to the corresponding micrograms of protein to enable comparisons among cell types. Data are represented as the mean with a 95% CI. **(C)** TET enzymatic activity assessed in nuclei extracts from NSCs after 14 days in culture. The plot shows a dramatic reduction of TET activity in IDH^{mut} cells with respect to the IDH^{wt} control. Data are represented as the mean with a 95% CI from technical replicates (n=2). A negative internal control, consisting of D-2HG with a final concentration of 400 μM, added to the IDH^{wt} extract during the assay, is also indicated (green color). **(D)** Quantification of 5hmC and 5mC global levels determined by LC-MS/MS in IDH^{mut} and IDH^{wt} NSCs controls after 14 days in culture. Data are represented as the mean with a 95% CI from technical replicates (n=4). P-values were calculated using one-sided Welch's test.

A

AP-1 family – all covered enhancers

CEBP family – all covered enhancers

Other TFs – all covered enhancers

B

Extended Data Fig. 9. Single-base resolution analysis of 5mCG and 5hmCG in the proximity of TFs motifs in all covered enhancers of D-2HG-treated primary microglia. **(A)** Plots showing separate analysis of 5mCG and 5hmCG percentages (upper) in the proximity of binding motifs, residing in all covered enhancers, for the indicated TFs in D-2HG-treated microglia and paired untreated controls. The size of each dot indicates the number of motifs analyzed in a paired manner between conditions. The 5mCG/5hmCG ratios at analyzed regions are also shown. The whole set of covered enhancers (used as controls) comprises the selected enhancers of Fig. 5c. **(B)** GSEA analysis using ranked genes by decreasing log2FC of D-2HG-treated vs. control samples of RNA-seq. The plot shows a significant bias towards decreased expression of genes regulated by D-2HG-driven hypermethylated/hypohydroxymethylated enhancers bound by TFs depicted in Fig 5c. Both methylome/hydroxymethylome and RNA-seq analyses were conducted using the same samples and in a paired manner.

A

B

C

Extended Data Fig. 10. Validation experiments on primary microglia treated with D-2HG for 14 days and challenged with LPS. (A) Heatmap depicting the expression of proinflammatory cytokines in CD11B+ cells from IDH^mut^ (n=20) and IDH^wt^ (n=8) human gliomas using RNA-seq data. **(B)** The plots indicate the expression of TNFA and IL6 (relative to housekeeping controls) in paired D-2HG-treated and control samples as determined by RT-qPCR. Primary microglia were obtained from surgical aspirates with IDH^mut^ status (n=4). P-values were calculated using one-sided paired t-tests. **(C)** Assessment of oxygen consumption rate (OCR) in paired D-2HG-treated and control samples as determined by XFe extracellular flux analyzer. Basal respiration (BR), maximal respiration (MR) and ATP-linked respiration (ALR) parameters are indicated for the comparisons control vs. D-2HG, and LPS vs. LPS+D-2HG. Primary microglia were obtained from surgical aspirate of one IDH^wt^ meningioma. Cells were maintained, treated, and collected under the same conditions as previously shown (Fig. 6a).