Determining the culturability of the bovine rumen bacterial and archaeal microbiota

Alice M. Buckner[1]* (ORCID ID: 0000-0002-7393-9882), Laura Glendinning[1] (ORCID ID: 0000-0003-4789-6644), Juan M. Palma Hidalgo[2] (ORCID ID: 0000-0002-6237-1822), Jolanda M. van Munster[2] (ORCID ID: 0000-0003-0412-7001), Mark Stevens[1] (ORCID ID: 0000-0003-2174-5039), Mick Watson[2] (ORCID ID: 0000-0003-4211-0358), and C. Jamie Newbold[2] (ORCID ID: 0000-0001-9561-300X)

[1] The Roslin Institute & Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush, Midlothian, EH25 9RG, United Kingdom

[2] Scotland's Rural College (SRUC), King's Buildings, West Mains Road, Edinburgh, EH9 3JG, United Kingdom.

* a.m.buckner@sms.ed.ac.uk

Abstract

Ruminants play an important part in global food security, but also emit methane which contributes to global warming. Microbes in the rumen play a key role in energy retention from the host's plant-based diet and the produce methane as a by-product of this digestion. While many thousands of novel microbial genomes have been assembled from metagenome sequence data, the extent to which these can be cultured is ill-defined. Here different media were used to study their effect on the bacterial and archaeal populations isolated; two non-selective media, one containing rumen fluid and the other not, and one selective medium which contained rumen fluid as well as a range of other substrates not present in the non-selective. *Bacteroidota* (52.53% ± 5.10) was the most common phylum found in the rumen fluid used as an inoculum, followed by *Bacillota* (41.00% ± 3.96), the archaeal phylum *Euryarchaeota* (5.12% ± 1.94), and *Pseudomonadota* (1.22% ± 0.78). The most common genera were bacterial, and included *Prevotella* (29.13% ± 4.16), *Butyrivibrio* (18.21% ± 2.08), *Succiniclasticum* (15.57% ± 5.03), unclassified *Bacteroidota* (13.91% ± 1.67), and unclassified *Prevotellaceae* (9.50% ± 2.01). All the operational taxonomic units in the rumen fluid were also found in the cultures. However, there was a significant difference between the communities of microbes grown on the different media, irrespective of the source cow. Of the 34 taxa isolated, 31 were found in a significantly different mean relative abundance in at least one medium compared to the others. These data further emphasise the importance of using defined media to selectively enrich for different microbial taxa, and only when used in combination is the true diversity of the rumen microbiota likely to be explored. This is essential in order to understand the complex workings of the rumen microbes, and how they interact, to enhance digestion efficiency and reduce the loss of energy as methane.

Introduction

The majority of ruminant digestion occurs in the forestomach termed the reticulorumen [1], referred to as the rumen in this paper. This non-secretory stomach compartment contains a community of microorganisms that aid the host in its digestion of plant matter by breaking down and fermenting ingesta before it reaches the 'true' secretory stomach, the abomasum [1]. The microbes in the rumen are key to digestion [1], and are composed of bacteria, archaea, protozoa, fungi, and bacteriophages [2]. This microbial community consists of a number of groups that are common across a wide variety of ruminants, often referred to as the 'core rumen microbiome' [3]; however, other members vary by factors such as species, breed, diet, or location [4], with this composition having a great influence on the efficiency of ruminant digestion [5,6].

Around 18.4% of global greenhouse gas emissions are produced by the agricultural industry, with 5.8% of global emissions coming from livestock and manure alone [7]. There is currently a high, and increasing, demand for animal products [2,8–10]. However the negative environmental impacts of the production of greenhouse gases during digestion need to be mitigated, and to do this the efficiency of the ruminant digestive system needs to be optimised [2].

Studying the rumen microbiome to find methods to ensure it is as efficient as possible is key to this goal, with traditional research methods involving culturing the organisms *in vitro* [11,12]. However, rumen microbes are fastidious and require strict anaerobic techniques to isolate them [13]. Following the development of Next Generation Sequencing (NGS) techniques, the use of metagenomics and metabarcoding strategies to analyse the community composition has become increasingly popular due

to the development of new techniques and technology, and the wealth of information it can provide [14]. There are advantages and disadvantages to each of these methods [15,16], however it has been recognised that the combination of sequencing and culturing can provide a more complete view of the rumen microbiome [13,17,18].

Members of the rumen microbiome currently isolated in culture do not represent the full diversity and complexity of the community, and taxonomic classification of those currently in culture is as yet incomplete [19–21], with many rumen microbes likely to have been misclassified [19]. The fact that some microbes seem to resist cultivation indicates a gap in knowledge of their basic metabolism and interactions with their environment [13]. The phyla *Bacillota* (heterotypic synonym; *Firmicutes)*, *Pseudomonadota* (homotypic synonym; *Proteobacteria)*, and *Actinomycetota* (homotypic synonym; *Actinobacteria)* are found most often in culture, with *Bacteroidota* (homotypic synonym; *Bacteroidetes)* being under represented [19]. The estimated proportion of culturable microbes in the rumen varies, spanning from < 1% [22] to 23% [18]. Methods to increase this proportion include a higher number of technical replicates, multiple media, and multiple dilutions of rumen fluid in the variety of media [18]. Here this is tested to help elucidate factors to be considered when trying to isolate novel bacteria from the rumen. The aim of this study was to determine; i) which rumen bacteria and archaea could be grown on the three media types chosen, ii) if any bacteria and archaea were enriched on any medium/media, and iii) the extent to which the rumen microbiome differed between individual cattle.

Results

Identification of operational taxonomic units (OTUs) present in rumen fluid and cultured on different media

Rumen fluid, diluted 1/100 in anaerobic diluent, was added to three distinct types of anaerobic liquid media (Med2, Med10, and MedTC) and incubated at 39°C for 48 hours. After the rumen fluid had been cultured, the microbes incubated, and their DNA extracted; their 16S V4 regions were amplified using unique barcoded primers, pooled, and sequenced. The raw sequencing data reads were then processed to generate operational taxonomic units (OTUs) which were classified to genus level. Of the 8152 OTUs generated, the majority were filtered out of the dataset, either due to there being a low number of reads for that OTU (< 0.1% abundance across all samples) or a low number of reads for a particular sample (< 3075 reads or a Good's Coverage of < 92.3%) (Supplementary Figure 1). At the end of the filtering steps, and after removing any OTUs not present in the original rumen fluid, 34 OTUs were present in the cultures. The 8118 OTUs removed only accounted for 11.7% of the entire number of reads in the dataset.

Microbial DNA in the basal medium

The three media chosen to cultivate the rumen microbes were based on those described by McSweeney et al. (2005) [22]. The media Med2 and MedTC contained sterile clarified rumen fluid as a medium constituent, whereas the medium Med10 did not. After sequencing the DNA extracted from all the samples, including medium-only controls, it was determined that there was rumen microbe DNA present in the medium from this sterile clarified rumen fluid. However there was a minimal effect on the presence of this DNA on the results (Supplementary Figure 2). On further comparison between each medium and its respective cultures, there were sufficient significant differences between the OTUs in the basal medium and those in the inoculated cultures to indicate the selective growth of different rumen microbes (Supplementary Figure 3, Supplementary Figure 4, Supplementary Figure 5).

Microbes grown in different media

For all the relative abundances below, the variation metric given is the standard deviation. Of the 34 taxa found across all the cultured samples, *Bacillota* (75.28% ± 6.34) was the most common phylum cultured, based on their relative abundance, followed by *Bacteroidota* (19.99% ± 4.85), *Pseudomonadota* (2.46% ± 2.01), and *Actinomycetota* (2.09% ± 1.07) (Figure 1a). The most common genera cultured across all media, based on their relative abundance, were *Selenomonas* (28.08% ± 11.71), *Streptococcus* (22.67% ± 6.06), *Prevotella* (18.71% ± 4.02), and unclassified *Lachnospiraceae* (11.50% ± 2.54) (Figure 1b).

Figure 1 The relative abundances of phyla (a) and genera (b) found in each replicate culture by cow and medium. Any phyla or genera with a relative abundance < 5% are included in the 'Other' category.

The α-diversity, or species richness, of a community is defined as the variation of taxa within a community, and can be measured in a multitude of ways [23]. Common indices of α-diversity include the Observed diversity which takes into account only the number of taxa and the number of individuals, Chao1 diversity which also includes the number of singletons and doubletons in that community to calculate species richness, and Shannon diversity which takes into account the relative and absolute abundance of different taxa combining both the diversity and evenness of the community [23,24]. When comparing the media Med2 and MedTC, there was a significant difference between the α-diversity for all three indices using a one-way ANOVA and post-hoc Tukey HSD test (Observed: p = 3.48E-03, 95% C.I. = 1.58,9.18; Shannon: p = < 1.00E-06, 95% C.I. = 0.52,0.79; Chao1: p = 0.03, 95% C.I. = 0.27,7.78) (Figure 2). For the other media comparisons, there was no significant difference between the Observed and Chao1 diversities, however there was a significant difference between the Shannon diversity between the media Med2 and Med10 (p < 1.00E-06, 95% C.I. = 0.53, -0.26) and Med10 and MedTC (p = 1.29E-04, 95% C.I. = 0.12,0.39). When looking at the difference in α-diversity metrics between the microbial communities grown from different individual cows, irrespective of media, there were no significant differences between the Observed, Shannon, or Chao1 diversities.

Figure 2 Comparison of α-diversity metrics between the microbial communities grown on the three media. a) Observed diversity, b) Shannon diversity, c) Chao1 diversity. Statistical tests used are one-way ANOVAs followed by post-hoc Tukey HSD tests if p < 0.05, with the results indicated with P < 0.05 = *, P < 0.01 = **, P < 0.001 = ***.

β-diversity is another method used to measure the diversity variation between communities [25], with the UniFrac metric along with Principal Coordinates Analysis (PCoA) often used to investigate differences between microbial populations, taking into account the phylogeny of the microbial taxa [26]. There was a significant difference in the β-diversity in the microbial communities found in the different media (PERMANOVA, F(2, 54, 56) = 2.67, p = 0.03), though post-hoc pairwise comparisons using a pairwise-PERMANOVA with Bonferroni correction for multiple testing only indicated a significant difference between the media Med2 and MedTC (F(1, 37, 38) = 4.11, p = 0.001) (Figure 3a). Using the same tests, there was no significant difference in the β-diversity between samples from different cattle (Figure 3b).

Figure 3 PCoA plot of weighted UniFrac distances of the cultures highlighted by medium (a) and source cow (b). Statistical tests used are one-way PERMANOVAs followed by post-hoc pairwise-PERMANOVAs with Bonferroni correction tests if p < 0.05, with the results indicated with P < 0.05 = *, P < 0.01 = **, P < 0.001 = ***.

Of the 34 OTUs classified to genus level found in the cultures, 31 were found in a significantly different mean relative abundance in at least one medium compared to the others (Kruskal-Wallis rank sum test with Bonferroni correction for multiple comparisons, Supplementary Table 1). The remaining three were not found to be enriched in any of the media. For those OTUs found in a significantly different relative abundance by medium, a post-hoc Dunn test was carried out (Supplementary Table 2). The 8 most abundant OTUs found across all the cultured samples are shown in Figure 4, which, when combined, make up 80% of the OTUs in terms of relative abundance.

Figure 4 The eight most abundant OTUs (classified to genus level) and their differential relative abundances in the cultured samples by medium. The taxa included are; a) Selenomonas, b) Streptococcus, c) Prevotella, d) Clostridium XIVa, e) Prevotella, f) unclassified Lachnospiraceae, g) unclassified Lachnospiraceae, h) Prevotella. Genera from the Bacillota phylum are indicated with 'Bcl', and those from the Bacteroidota phylum are indicated with 'Bct'. Statistical tests used are Kruskal-Wallis tests followed by post-hoc pairwise Dunn tests with Bonferroni correction if p < 0.05, with the results indicated with P < 0.05 = *, P < 0.01 = **, P < 0.001 = ***.

Microbes identified in the source rumen fluid inoculum

In addition to sequencing the 16S rRNA found in the cultures, the rumen fluid from each cow used as the culture inoculum were also sequenced. After going through the same filtering steps as the cultures, the four samples of rumen fluid contained 34 OTUs (classified to genus level). Of these, *Bacteroidota* (52.53% ± 5.10) was the most common phylum found in the four cows rumen fluid, based on their relative abundance, followed by *Bacillota* (41.00% ± 3.96), the archaeal phylum *Euryarchaeota* (5.12% ± 1.94), *Pseudomonadota* (1.22% ± 0.78), and finally *Actinomycetota* (0.12% ± 0.08) (Figure 5a). The most common genera found in the four cows rumen fluid, based on their relative abundance, were *Prevotella* (29.13% ± 4.16), *Butyrivibrio* (18.21% ± 2.08), *Succiniclasticum* (15.57% ± 5.03), unclassified *Bacteroidetes* (13.91% ± 1.67), and unclassified *Prevotellaceae* (9.50% ± 2.01) (Figure 5b).

Figure 5 The relative abundances of the phyla (a) and genera (b) that make up the microbial communities found in each of the rumen fluid samples from the source cows. All phyla are shown, however any genera with a relative abundance < 5% are included in the 'Other category.

The difference between the cultured microbes and those in the source rumen fluid inoculum

When comparing the microbial communities found in the rumen fluid from the source cows to those grown in culture, vast differences were observed (Figure 6, Supplementary Table 3). The same OTUs were found in both datasets, however in different relative abundances. This was quantified using the Log2-fold change of the mean relative abundance in the different media compared to the rumen fluid. Based on this, the phyla (Figure 6a) *Actinomycetota* (Log2-fold change compared to rumen fluid; Med10 = 3.66, Med2 = 4.60, MedTC = 3.84) and *Bacillota* (Med10 = 0.92, Med2 = 0.96, MedTC = 0.74) were overrepresented in the cultures on all media, and *Pseudomonadota* was overrepresented in Med2 (0.29) and MedTC (2.02), but not Med10 (-0.40). *Euryarchaeota* archaea (Med10 = -6.93, Med2 = -9.92, MedTC = -3.32) and *Bacteroidota* (Med10 = -1.39, Med2 = -1.73, MedTC = -1.11) were found in lower relative abundances in the media compared to the rumen fluid. Regarding genera (Figure 6b), the most underrepresented in the media compared to the rumen fluid were the methanogenic *Methanomassiliicoccus* in Med2 (-9.92), *Succiniclasticum* in Med2 (-8.47), and unclassified *Prevotellaceae* in Med2 (-8.26). The genera most overrepresented in culture were *Clostridium XIVa* (Med10 = 11.23, Med2 = 9.91, MedTC = 9.17), *Megasphaera* (Med10 = 8.34, MedTC = 8.99), *Selenomonadales* in Med10 (7.16), and *Streptococcus* (Med10 = 7.87, Med2 = 7.31, MedTC = 7.64).

Figure 6 The mean relative abundances of the phyla (a) and genera (b) that make up the microbial communities found in the cultures grown on each of the types of media compared to the rumen fluid samples, along with the Log2 fold change of the mean relative abundance of each phylum and genus compared to the respective mean relative abundance found in the rumen fluid. Those taxa found in higher mean relative abundances in the rumen fluid are highlighted in red, those in lower mean relative abundances are shown in blue, and those with little change are shown in white.

Discussion

Culturing rumen microbes is essential to explore the full diversity of the rumen microbiome, and to investigate how different microbes interact and react to different conditions. However the majority of rumen microbes are fastidious obligate anaerobes, making isolating them in culture very difficult. In addition, many of the microbes in the rumen, identified through metagenomic methods, cannot be classified and have not been isolated in pure culture, meaning it is difficult to develop culture media that will support their growth. In this study, using a range of commonly used media to isolate rumen microorganisms, 34 rumen microbes were cultivated in co-culture. Different rumen microbial genera, such as *Selenomonas*, *Streptococcus*, and *Prevotella*, grew in different relative abundances on the different media, and a number of microbes were grown that could not be classified to genus level. This shows the value of different media for selecting for the growth of different rumen microbes, and the potential for further optimisation to enrich for the growth of target, as-yet-unclassified, microbes.

The majority of the OTUs identified in the cultures belonged to the genus *Selenomonas*, a Gram-negative, obligately anaerobic, flagellated genus of bacteria, with those found in, and isolated from, the rumen usually belonging to the species *S. ruminantium* [27,28]. They primarily break down less complex carbohydrates to produce acetate, propionate and carbon dioxide, though some can also break down lactate and glycerol [27,28], and are one of the key lactate-degraders in the rumen [27]. Their ability to utilise lactate and succinate means they are involved in the succinate pathway of propionate production [29,30], increased production of which is associated with reduced methane formation [29]. *Selenomonas* can also degrade proteins, amino acids, urea, and ammonia [27,28], and are involved in nitrite reduction [27,28,31,32]. In this study, the single OTU identified as belonging to the genus *Selenomonas* grew in the highest relative abundance in the medium with no VFAs (Med2). However, the difference in absolute amount of this microbe between the three media could only be ascertained with additional qPCR. This medium did, however, contain the highest amount of yeast extract and also contained many less complex carbohydrates such as cellobiose, maltose, and glucose. This, along with the presence of sodium lactate could explain why this microbe grew in the highest relative abundance on this medium.

The next most abundant microbe cultured in this study were *Streptococcus spp.*, with members of this genus being Gram-positive facultative aerobes, able to function in both aerobic and anaerobic conditions [33]. The bovis species group includes, amongst others, *S. bovis* (now *S. equinus*) and various subspecies of *S. gallolyticus*, and can be classified as Lancefield serogroup Group D [33]. They are non-motile, oval cocci and are found in pairs or chains [33], and produce L-(+)lactate from glucose [4,33]. They appear in the highest relative abundance on Med2 and MedTC, which may be due to the presence of sterile clarified rumen fluid, mineral solution 5, sodium lactate, $NaHCO_3$, and lack of VFAs compared to Med10. *Prevotella*, another microbe cultured in high abundance, is a common rumen microbe. *Prevotella spp.* are rod-shaped and vary in length, being non-motile, non-sporing, and strictly anaerobic [34,35]. This genus is highly diverse with saccharolytic, xylanolytic, pectinolytic, proteolytic, hemicellulolytic, and amylolytic activity [34–36], and are able to break down complex carbohydrates [37]. To grow they require haemin, menadione, and peptides such as those found in trypticase, peptone, and yeast extract [35]. The three most relatively abundant OTUs belonging to *Prevotella* were enriched on one of each of the media. This could be due to them belonging to different species or strains and therefore having different substrate requirements from the medium.

Following this, the next most abundant microbe cultured was a form of unclassified *Lachnospiraceae*, potentially belonging to one of the existing genera, or a new lineage entirely. The family *Lachnospiraceae* includes the genera *Butyrivibrio*, *Lachnospira*, and *Roseburia* [28]. These genera are all obligately anaerobic, motile, curved rods, with various Gram staining characteristics [27,28]. The main *Butyrivibrio* species found in the rumen is *B. fibrisolvens*, though the bacteria classified to this species are likely as-yet-unclassified sub-groups, species, and strains due to their varying colony morphology and fermentation patterns [28]. They are all xylanolytic and mainly produce butyrate, though they can ferment a wide range of carbohydrates and use a variety of nitrogen sources and produce a range of VFAs and vitamins [28,38]. Some also have proteolytic activity and produce bacteriocins [28,38]. When cultured in broth they produce a sediment, and grow well on media containing xylan and fine cellulose, along with trypticase and peptone [28]. *Butyrivibrio spp.* are often found in higher amounts in ruminants on high concentrate diets [28]. *Lachnospira* form filamentous and 'woolly' cultures in broth and on plates, and don't require cellulose, xylans, trypticase, or peptones [28]. They are pectinolytic and grow better if ammonia is present, producing lactate, formate and acetate, with little butyrate and no propionate produced [28]. They are found in higher amounts in ruminants on high pectin diets, for example clover [29].

Roseburia ferment glucose and acetate to butyrate and produce H_2 and CO_2 [27,39], and are able to use a variety of carbohydrates as energy sources [27]. They grow best in media containing VFAs, along with yeast extract, trypticase, peptone, and haemin [27]. The two most relatively abundant OTUs belonging to this unclassified taxon were also enriched in different media, suggesting they are different species with different metabolic requirements.

In the rumen, higher abundances of *Selenomonas spp*. are often associated with high grain [27,28] or high concentrate diets [40]. Xue et al. (2022) found that more efficient animals showed interactions in the rumen between these *Selenomonas spp*. and bacteria of the *Succinivibrionaceae* family [41], with Smith et al. (2022) observing higher levels of *Selenomonas* in animals with low residual methane emissions [42]. High levels of the lactate degrading species, often named *Selenomonas ruminantium subsp. Lactilytica*, are often found in higher numbers in the rumen when the rumen pH is reduced to the extent it leads to acidosis and bloat. As a genus involved in propionate production [29], further study of this genus in culture could lead to a better understanding of how this pathway could be used to reduce methane production. In addition, a probiotic dose of a strain of *Selenomonas ruminantium subsp. Lactilytica* was able to reduce the effects of lactic acidosis [31], and could act as a potential treatment or prophylactic for this issue. Ensuring a number of strains are characterised and reliably cultured in vitro is essential, with this study showing three media able to grow this microbe in high relative abundances, with Med2 being particularly effective at this. High levels of *Streptococcus spp*. in the rumen can lead to acidosis caused by the accumulation of lactate [43], though ruminal *Streptococcus spp*. have been shown to produce bacteriocins that are inhibitory to other *Streptococci* [44], which could provide scope for the control of ruminal acidosis using bacteriocins or bacteriocin-producing bacteria. *Streptococcal* species are also one of the causative agents of mastitis [45], another disease these potential bacteriocins could help treat [46]. *Prevotella spp*. also has the potential to promote the stimulation of alternative hydrogen sinks and reduce the amount of methane produced from ruminal fermentation [37]. In the literature, a high abundance of *Prevotella spp*. seems to be linked to low methane production due to the shift in fermentation profile towards propionate production [37,47]. However, *Prevotella spp*. are greatly under-represented in pure culture collections [48]. Similarly to *Selenomonas spp*., this may have the potential to improve ruminant efficiency if the proportion of *Prevotella spp*. are increased in the rumen, however in-depth study of different species and strains in vitro is required first. Many unclassified *Lachnospiraceae* have been identified in the rumen, and are likely to be important [49]. *Lachnospiraceae* have been shown to be associated with a high residual feed intake[50] and low feed conversion ratio [51], so the *in vitro* study and further classification of these isolates would help elucidate the as-yet-undetermined characteristics of these microbes, with the media used in this research able to facilitate this.

The complete bacterial and archaeal microbiomes of the cattle used in this study were similar, aligning with evidence suggesting that similarly treated animals of the same breed raised in the same location will have a similar microbiome [4]. The majority of the OTUs identified in the full rumen fluid belonged to the phyla *Bacteroidota*, *Bacillota*, unclassified *Bacteria*, and the archaeal *Euryarchaeota*. This coincides with the generally agreed structure of the core microbiome outlined in reviews such as Terry et al. (2019) [52], Castillo-González et al. (2014) [53], and Anderson et al. (2021) [54]. However the rumen microbiome structure between ruminant populations can differ due to the location of the animals [55] and their breed [56]. The diet also affects the rumen microbiome, with cattle used in this study grazing on perennial ryegrass with a supplementary garlic lick. A grass diet is often associated with higher levels of *Bacteroidota* [57], and garlic licks are associated with reduced levels of methanogens [58] and increased levels of *Prevotella* [59].

The microbes cultured here coincide with others who have used Med10 [60,61], Med2 [62–66], and MedTC media [67], however the majority of the rumen microbiome is uncultured and members of it were not able to be identified or classified using current databases [18–21,49,68]. At the moment, isolates of *Bacillota*, *Pseudomonadota*, and *Actinomycetota* are found most often in culture collections with *Bacteroidota* being under represented [19]. This is also observed in this study with a higher proportion of the bacteria cultured belonging to *Bacillota*, than those in the rumen fluid where a large proportion belong to *Bacteroidota*. In addition, the proportion of archaea cultured, compared to that of the original rumen fluid, was minimal. Despite them having a lower relative abundance and level of diversity compared to the rumen bacteria, it is essential to further investigate methods to isolate them in culture [69–71]. Because of this, there is vast potential for the development of new innovative techniques and media formulations to culture and classify rumen microbes. New and improved protocols, equipment, and techniques could aid in overcoming many of the barriers that currently exists when attempts to bring new microbes into culture are made. These can include changes in sample collection, processing, and inoculation methods with innovations around media composition, incubation conditions, length of time incubated, and the

type of anaerobic atmosphere needed. There is also some current bias and accuracy of database choice on OTU classification [72], and the choice of the V4 16S region will also have meant that microbes such as fungi will have been missed in this study, and may have skewed some of the taxonomic classification [73]. To work towards increasing the number of rumen microbes in culture collections and improving the understanding of the rumen microbiome, future work with these cultures will include using dilution-to-extinction and streak plating in order to isolate the bacteria and archaea in pure culture and to characterise these.

There is great scope for expanding the number of cultured rumen microbes using different media and anaerobic techniques, as demonstrated in this study. Increasing numbers of these isolates in pure culture, and their availability in culture collections, is essential to improve knowledge surrounding the intricacies of the rumen microbiome, microbial degradation of feed products, and reduction of methane production

Methods

Ethics statement

Animal studies were carried out under the Animal (Scientific Procedures) Act 1986. The maintenance, sampling, and testing regime of these animals were covered by the Home Office project licence PP7153972.

Sample collection and microbial culture

Samples of rumen fluid, for the purpose of inoculation and culturing, were collected from four fistulated, female, Jersey cows, aged 5 to 12 years, grazing a sward of perennial ryegrass. Rumen fluid was collected via the fistula, squeezed into a pre-warmed thermos, and transported back to the laboratory for inoculation within two hours of collection. The samples were squeezed through two layers of muslin into a CO_2 gassed container and kept in a water bath set to 39°C. This was then serially diluted in anaerobic diluent to achieve a 10^{-2} dilution of the original rumen fluid. The anaerobic diluent was based on that from Bryant and Burkey (1952) [74], however the concentration of cysteine was reduced to match that present in the media (0.51 g/L). In addition to the dilution series, 8 mL of the strained rumen fluid was spun at 3018 g for 20 minutes before the supernatant was removed and the pellet stored at -80°C.

McSweeney et al. (2005) described five types of anaerobic culture media commonly used for isolation of rumen bacteria and archaea [22]. Based on a pilot experiment (data not shown) three media were chosen based on their ability to support microbial growth (based on the level of change in optical density over time after inoculation with rumen fluid). These three media were composed of two non-selective media, one containing rumen fluid (Med2 [75]) and the other not (Med10 [76]), and one selective medium (MedTC [22]) which contained rumen fluid as well as a range of other substrates not present in the non-selective media. The non-selective media aimed to mimic the entire rumen environment to culture a wide range of rumen microbes, while the selective media contained additional substrates to select for additional microbes, such as ones that are xylanolytic, pectinolytic, or that rely on the presence of trace elements.

All stock solutions that were used to make up the media were made in aerobic conditions. The three media and anaerobic diluent were made using these component solutions under anaerobic conditions using 100% CO_2 gas. The water was added to a heat proof flask and put on a hot plate with magnetic stirrer and temperature probe. The hot plate was set to 95°C and stirrer to 300 rpm. The liquid and solid media ingredients were added. When the mixture reached 95°C the CO_2 line was added and the hot plate set to 55°C. When the mixture reached 55°C the reducing agents were added. When the mixture had no red tint, the media was dispensed into 10 mL Hungate tubes for the media or 100 mL Wheaton tubes for the anaerobic diluent. The tube was gassed with CO_2, the anaerobic solution added, and a cap and bung quickly used to seal the tube. The tubes were then autoclaved at standard conditions, allowed to cool to room temperature, and the caps sterilised with ethanol before the heat unstable ingredients were added using a needle and syringe. An overview of the components of the media and anaerobic diluent is given in Supplementary Table 4.

For each cow and each medium, four controls were processed and incubated alongside their respective samples. Two of the controls consisted of sterile media only, while two were mock inoculated with 1 mL of sterile anaerobic diluent. Ethanol was used to sterilise the caps of the tubes of media, and 1 mL of the 10^{-2} dilution was added to five replicate tubes of media. The optical density (590 nm) of the cultures

was measured just after inoculation (time 0), after 24 hours, and after 48 hours of incubation at 39°C. After the 48-hour measurement 8 mL of each sample was removed and spun at 4,500 rpm for 20 minutes. The supernatant was discarded and the pellet stored at -80°C for DNA extraction. 2 mL of sterile 25% glycerol was added to the remaining 2 mL of culture, and was stored at -80°C.

Determining microbial composition

The pellets of the original rumen fluid sample and the cultures (along with negative and reagent-only controls) were subject to DNA extraction, 16S rDNA amplification, and massively-parallel sequencing before bioinformatics was used to determine the OTUs present. A microbial standard community (ZymoBIOMICS Microbial Community Standard) was also used to determine how effective and accurate these steps were. The relevant results are described in Supplementary Figure 1.

DNA from the pellet was extracted using a modified protocol based on the Qiagen DNeasy® PowerLyzer® PowerSoil® Kit and Qiagen QIAamp® Fast DNA Stool Mini Kits. The majority of the protocol followed the QIAamp® Fast DNA Stool Mini Kit protocol, with an additional glass bead beating step on centrifuged culture samples with the supernatant removed (as described in DNeasy® PowerLyzer® PowerSoil® with modifications described in Supplementary Note 1), and eluting the DNA in a lower volume (50 µL). DNA was extracted from the Zymo microbial standard with InhibitX added to 75ul of the standard before the protocol was carried out as per the other samples.

16S rDNA amplification was then carried out on the eluted DNA. Barcoded primers (5′–TATGGTAATTGTGTGCCAGCMGCCGCGGTAA–3′ and 5′–AGTCAGTCAGCCGGACTACHVGGGTWTCTAAT–3′) were used to target the V4 region before being cycled through a PCR machine with a heated lid, running at 95°C for 2 minutes followed by 30 cycles of 95°C for 20 seconds, 55°C for 15 seconds, 72°C for 5 minutes followed by 72°C for 10 minutes. 12 µL of Q5 Hot Start High-Fidelity 2X Master Mix, 9 µL of nuclease free water, 1.25 µL of the forward primer (10nM), and 1.25 µL of the reverse primer (10nM) was added to either 1 µl or 0.2 µl of the template DNA, depending on which provided the highest quantity of amplicon product. The product was then purified using Ampure magnetic bead purification in a 1:1 ratio, quantified using Nanodrop and Qubit, and visualised using gel electrophoresis. Using the HS Qubit values for the concentration of amplified DNA in each sample, the DNA was pooled to form a pool with equimolar concentrations of DNA from each sample.

Sequencing of the single pool was carried out by Edinburgh Genomics using Illumina MiSeq V2 chemistry, with 500 cycles producing an output of around 11 million 300 base pair paired-end reads. As very few reads were produced for three of the four rumen fluid samples, due to the importance of having these samples in the dataset, they were sequenced again in another identical Miseq run and then substituted into the initial dataset.

Bioinformatic analysis was carried out using Mothur v. 1.48.0., and follows the Miseq SOP (https://mothur.org/wiki/miseq_sop/) [77]. Steps in the sections 'Reducing sequencing and PCR errors' and 'Processing improved sequences' were followed exactly, with the exception of Archaea not being removed in the remove.lineage command. OTUs were generated using the commands given in the 'OTU' section of the 'Preparing for analysis' section (without the use of cluster.split), before consensus sequences for each OTU were generated using the classify.otu command given in the 'Phylotypes' section. Databases used were Version 138.1 of the SILVA database (https://mothur.org/wiki/silva_reference_files/) with the 'full-length sequences and taxonomy references' modified to contain just the V4 region, and Version 18 of the '16S rRNA reference (RDP)' database (https://mothur.org/wiki/rdp_reference_files/).

R v. 4.2.2. was used with the package Phyloseq v. 1.42.0. [78] to collate the metadata, OTU table, and taxonomy table, and the protocol followed was based on that as recommended by the package developer [79] and others including [80–82]. The OTU table, taxonomy table, metadata, and tree were imported from MOTHUR into R. They were combined using the Phyloseq package to give 8152 taxa by 6 taxonomic ranks from 96 samples by 9 sample variables.

Sampling coverage was calculated to determine which samples to remove by calculating the number of reads per sample (Supplementary Figure 1). From the histogram, many samples had very few reads, with the scatterplot showing a large cluster of samples at the bottom at around 1e+02 with the rest scattered around 1e+05. This is also echoed in the line graph showing many low abundance samples

followed by significant step up to around 3,000 reads. The Good's coverage was then calculated, which quantifies the relationship between the number of sequences and the number of OTUs only present once in each sample, with 7 of the 60 samples having a Good's coverage of < 99%. Due to this, samples with fewer than (or equal to) 3075 reads or 92.3% Good's coverage were removed to give 8129 taxa across 82 samples.

Sampling depth was calculated to determine which low abundance OTUs to remove by looking at the total number of reads per OTU across all samples. The total number of reads was 5,256,392. If any OTUs had a relative abundance of less than 0.1% across all cultures, they were removed. These OTUs are unlikely to be able to be reliably cultured again as, for a bacteria and archaea with a relative abundance of < 0.1%, over 1000 tubes of media would have to be inoculated for these bacteria and archaea to be cultured once. This resulted in a dataset containing 43 taxa from 82 samples. Finally, any OTUs present in the cultures or controls, but not present in any of the rumen fluid samples, were removed, giving 34 taxa. Despite this step removing 8118 OTUs, only 11.7% of the total reads were discarded.

To determine if there was a difference in the OTUs found in the medium-only controls between the three media, the α-diversity (Observed, Shannon, and Chao1 calculated using the Microbiome R package) and the β-diversity (weighted UniFrac) were calculated. An ANOVA was carried out to determine if there was a significant difference between the α-diversity metrics by medium, with a post-hoc Tukey test used if $p < 0.05$. For the β-diversity a PERMANOVA (using adonis2 from the Vegan package) was used to determine if there was a significant difference between the β-diversity by medium, with a post-hoc pairwise PERMAVOVA run if $p < 0.05$. Following this, each medium-only control was compared to the cultures grown on this medium to determine if the OTU diversity had changed. A two-sample T Test was carried out to determine the difference between the α-diversities of the medium-only controls and the cultures grown on that medium, and a PERMANOVA was used for the β-diversity. Next, the α-diversity metrics and β-diversity was compared between the cultures by medium, with an ANOVA (and post-hoc Tukey) and PERMANOVA (and post-hoc pairwise PERMANOVA with manual Bonferroni adjustment for multiple testing) used as described above. Finally, the difference in OTU relative abundance was determined using the same method above using a Kruskal–Wallis test (along with a post-hoc Dunn test if $p < 0.05$). The 8 most abundance OTUs across all cultures were shown in a boxplot as they make up over 80% of the dataset.

Data availability statement

The raw reads produced through sequencing can be found through the European Nucleotide Archive (ENA), with the primary accession code PRJEB78542 (https://www.ebi.ac.uk/ena/browser/view/PRJEB78542).

Acknowledgements

The authors would like to thank the farm staff for facilitating sample collection.

Author contribution statement

AB led the work on the project and LG, JPH, JvM, MS, MW, and JN co-supervised. All were involved in experimental design. AB, LG, JPH, and JvM carried out experimental procedures, and AB, LG, and MW led the bioinformatics and data analysis. AB wrote the main body of the manuscript, and all authors were involved in editing the manuscript and preparing it for publication.

Additional information

Funding declaration

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC) grant number BB/T00875X/1 and Institute Strategic Programme Grant BBS/E/RL/230001A. Sequencing was carried out by Edinburgh Genomics, the University of Edinburgh, which is partly supported with core funding from NERC (UKSBS PR18037). For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Competing interests

The authors declare no competing interests.

References

1. Czerkawski, J. W. *An Introduction to Rumen Studies*. (Pergamon Press, Great Britain, 1985).
2. Huws, S. A. *et al.* Addressing global ruminant agricultural challenges through understanding the rumen microbiome: Past, present, and future. *Front. Microbiol.* **9**, (2018).
3. Wallace, R. J. *et al.* A heritable subset of the core rumen microbiome dictates dairy cow productivity and emissions. *Sci Adv* **5**, 8391–8394 (2019).
4. Hungate, R. *The Rumen and Its Microbes*. (Academic Press, New York and London, 1966).
5. Hernandez-Sanabria, E. *et al.* Impact of feed efficiency and diet on adaptive variations in the bacterial community in the rumen fluid of cattle. *Appl. Environ. Microbiol.* **78**, 1203–1214 (2012).
6. Zhou, M., Hernandez-Sanabria, E. & Le, L. G. Assessment of the microbial ecology of ruminal methanogens in cattle with different feed efficiencies. *Appl. Environ. Microbiol.* **75**, 6524–6533 (2009).
7. Ritchie, H., Roser, M. & Rosado, P. CO_2 and Greenhouse Gas Emissions. *Our World in Data* https://ourworldindata.org/co2-and-greenhouse-gas-emissions (2020).
8. Delgado, C. L. Rising Consumption of Meat and Milk in Developing Countries Has Created a New Food Revolution. *Am. Sciety Nutr. Sci.* 3907S-3910S (2003).
9. Ritchie, H. & Roser, M. Meat and Dairy Production. *Our World in Data* https://ourworldindata.org/meat-production (2017).
10. FAO. *Shaping the Future of Livestock: Sustainably, Responsibly, Efficiently. The 10th Global Forum for Food and Agriculture (GFFA)*. https://www.fao.org/3/i8384en/I8384EN.pdf (2018).
11. Holdeman, L. V., Cato, E. P. & Moore, W. E. C. *Anaerobe Laboratory Manual*. 152 (V.P.I. Anaerobe Laboratory, Virginia Polytechnic Institute and State University, 1977).
12. Sebald, M. & Hauser, D. Pasteur, oxygen and the anaerobes revisited. *Anaerobe* **1**, 11–16 (1995).
13. Ó Cuív, P., McSweeney, C. & Mackie, R. A question of culture: bringing the gut microbiome to life in the -omics era. in *Improving Rumen Function* 29–54 (Burleigh Dodds Science Publishing, 2020). doi:10.19103/AS.2020.0067.03.
14. Kothari, R. K. *et al.* Comprehensive Exploration of the Rumen Microbial Ecosystem With Advancements in Metagenomics. in *Metagenomics: Perspectives, Methods, and Applications* 215–229 (Elsevier Inc., 2018). doi:10.1016/B978-0-08-102268-9.00011-2.
15. Lema, N. K., Gemeda, M. T. & Woldesemayat, A. A. Recent Advances in Metagenomic Approaches, Applications, and Challenges. *Curr. Microbiol.* **80**, 347 (2023).
16. Liu, Y.-X. *et al.* A practical guide to amplicon and metagenomic analysis of microbiome data. *Protein Cell* **12**, 315–330 (2021).
17. Greub, G. Culturomics: A new approach to study the human microbiome. *Clin. Microbiol. Infect.* **18**, 1157–1159 (2012).
18. Zehavi, T., Probst, M. & Mizrahi, I. Insights into culturomics of the rumen microbiome. *Front. Microbiol.* **9**, (2018).
19. Creevey, C. J., Kelly, W. J., Henderson, G. & Leahy, S. C. Determining the culturability of the rumen bacterial microbiome. *Microb. Biotechnol.* **7**, 467–479 (2014).
20. Seshadri, R. *et al.* Cultivation and sequencing of rumen microbiome members from the Hungate1000 Collection. *Nat. Biotechnol.* **36**, 359–367 (2018).
21. Matthews, C. *et al.* The rumen microbiome: a crucial consideration when optimising milk and meat production and nitrogen utilisation efficiency. *Gut Microbes* **10**, 115–132 (2019).
22. Mcsweeney, C. S., Denman, S. E. & Mackie, R. I. Rumen bacteria. in *Methods in Gut Microbial Ecology for Ruminants* (eds. Makkar, H. P. S. & McSweeney, C. S.) 23–27 (IAEA, Netherlands, 2005).
23. Thukral, A. K. A review on measurement of Alpha diversity in biology. *Agric. Res. J.* **54**, 1 (2017).
24. Xia, Y. & Sun, J. *Bioinformatic and Statistical Analysis of Microbiome Data: From Raw Sequences to Advanced Modeling with QIIME 2 and R*. (Springer International Publishing, Cham, 2023). doi:10.1007/978-3-031-21391-5.
25. Anderson, M. J. *et al.* Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist: Roadmap for beta diversity. *Ecol. Lett.* **14**, 19–28 (2011).
26. Lozupone, C., Lladser, M. E., Knights, D., Stombaugh, J. & Knight, R. UniFrac: an effective distance metric for microbial community comparison. *ISME J.* **5**, 169–172 (2011).
27. *Bergey's Manual of Systematic Bacteriology, Volume 3 (The Firmicutes)*. vol. 3 The Firmicutes (Springer, New York, 2009).

28. *The Prokaryotes: Volume 4: Bacteria: Firmicutes, Cyanobacteria.* (Springer US, New York, NY, 2006). doi:10.1007/0-387-30744-3.

29. Lan, W. & Yang, C. Ruminal methane production: Associated microorganisms and the potential of applying hydrogen-utilizing bacteria for mitigation. *Sci. Total Environ.* **654**, 1270–1283 (2019).

30. Ungerfeld, E. M. & Kohn, R. A. The role of thermodynamics in the control of ruminal fermentation. in *Ruminant physiology: Digestion, metabolism and impact of nutrition on gene expression, immunology and stress* (eds. Sejrsen, K., Hvelplund, T. & Nielsen, M. O.) (Wageningen Academic Publishers, The Netherlands, 2006). doi:10.3920/978-90-8686-566-6.

31. Wiryawan, K. & Brooker, J. Probiotic control of lactate accumulation in acutely grain-fed sheep. *Aust. J. Agric. Res.* **46**, 1555 (1995).

32. Yoshii, T., Asanuma, N. & Hino, T. Number of nitrate- and nitrite-reducing *Selenomonas ruminantium* in the rumen, and possible factors affecting its growth. *Anim. Sci. J.* **74**, 483–491 (2003).

33. Toit, M. D., Huch, M., Cho, G. & Franz, C. M. A. P. The genus Streptococcus. in *Lactic Acid Bacteria* (eds. Holzapfel, W. H. & Wood, B. J. B.) 457–505 (Wiley, 2014). doi:10.1002/9781118655252.ch28.

34. Flint, H. J. & Duncan, S. H. Bacteroides and Prevotella. in *Encyclopedia of Food Microbiology* 203–208 (Elsevier, 2014). doi:10.1016/B978-0-12-384730-0.00031-8.

35. *Bergey's Manual of Systematic Bacteriology, Volume 4 (The Bacteroidetes, Spirochaetes, Tenericutes (Mollicutes), Acidobacteria, Fibrobacteres, Fusobacteria, Dictyoglomi, Gemmatimonadetes, Lentisphaerae, Verrucomicrobia, Chlamydiae, and Planctomycetes).* vol. 4 The Bacteroidetes, Spirochaetes, Tenericutes (Mollicutes), Acidobacteria, Fibrobacteres, Fusobacteria, Dictyoglomi, Gemmatimonadetes, Lentisphaerae, Verrucomicrobia, Chlamydiae, and Planctomycetes (Springer New York, New York, NY, 2010).

36. Zhou, M., Chen, Y. & Guan, L. Rumen Bacteria. in *Rumen Microbiology: From Evolution to Revolution* (eds. Puniya, A. K., Singh, R. & Kamra, D. N.) (Springer, 2015).

37. Betancur-Murillo, C. L., Aguilar-Marín, S. B. & Jovel, J. Prevotella: A Key Player in Ruminal Metabolism. *Microorganisms* **11**, 1 (2022).

38. Sengupta, K. *et al.* Genomic architecture of three newly isolated unclassified Butyrivibrio species elucidate their potential role in the rumen ecosystem. *Genomics* **114**, 110281 (2022).

39. McLoughlin, S. *et al.* Rumen Microbiome Composition Is Altered in Sheep Divergent in Feed Efficiency. *Front. Microbiol.* **11**, (2020).

40. Wang, L., Zhang, G., Li, Y. & Zhang, Y. Effects of High Forage/Concentrate Diet on Volatile Fatty Acid Production and the Microorganisms Involved in VFA Production in Cow Rumen. *Animals* **10**, 223 (2020).

41. Xue, M.-Y. *et al.* Integrated meta-omics reveals new ruminal microbial features associated with feed efficiency in dairy cattle. *Microbiome* **10**, 32–32 (2022).

42. Smith, P. E., Kelly, A. K., Kenny, D. A. & Waters, S. M. Differences in the Composition of the Rumen Microbiota of Finishing Beef Cattle Divergently Ranked for Residual Methane Emissions. *Front. Microbiol.* **13**, 855565 (2022).

43. Herrera, P., Kwon, Y. M. & Ricke, S. C. Ecology and pathogenicity of gastrointestinal Streptococcus bovis. *Anaerobe* **15**, 44–54 (2009).

44. Mantovani, H. C., Kam, D. K., Ha, J. K. & Russell, J. B. The antibacterial activity and sensitivity of Streptococcus bovis strains isolated from the rumen of cattle1. *FEMS Microbiol. Ecol.* **37**, 223–229 (2001).

45. Kabelitz, T., Aubry, E., Van Vorst, K., Amon, T. & Fulde, M. The Role of Streptococcus spp. in Bovine Mastitis. *Microorganisms* **9**, 1497 (2021).

46. Pieterse, R. & Todorov, S. D. Bacteriocins: exploring alternatives to antibiotics in mastitis treatment. *Braz. J. Microbiol.* **41**, 542–562 (2010).

47. Aguilar-Marin, S. B., Betancur-Murillo, C. L., Isaza, G. A., Mesa, H. & Jovel, J. Lower methane emissions were associated with higher abundance of ruminal Prevotella in a cohort of Colombian buffalos. *BMC Microbiol.* **20**, 364 (2020).

48. Bekele, A. Z., Koike, S. & Kobayashi, Y. Genetic diversity and diet specificity of ruminal Prevotella revealed by 16S rRNA gene-based analysis: Genetic diversity and diet specificity of rumen Prevotella. *FEMS Microbiol. Lett.* **305**, 49–57 (2010).

49. Kim, M., Morrison, M. & Yu, Z. Status of the phylogenetic diversity census of ruminal microbiomes: Status of microbial census in rumen. *FEMS Microbiol. Ecol.* **76**, 49–63 (2011).

50. Li, F., Zhou, M., Ominski, K. & Guan, L. L. Does the rumen microbiome play a role in feed efficiency of beef cattle?1. *J. Anim. Sci.* **94**, 44–48 (2016).

51. Hernandez-Sanabria, E. *et al.* Correlation of particular bacterial pcr-denaturing gradient gel electrophoresis patterns with bovine ruminal fermentation parameters and feed efficiency traits. *Appl. Environ. Microbiol.* **76**, 6338–6350 (2010).

52. Terry, S. A., Badhan, A., Wang, Y., Chaves, A. V. & McAllister, T. A. Fibre digestion by rumen microbiota — a review of recent metagenomic and metatranscriptomic studies. *Can. J. Anim. Sci.* **99**, 678–692 (2019).

53. Castillo-González, A. R., Burrola-Barraza, M. E., Domínguez-Viveros, J. & Chávez-Martínez, A. *Rumen Microorganisms and Fermentation*. vol. 46 349–361 (2014).

54. Anderson, C. J., Koester, L. R. & Schmitz-Esser, S. Rumen Epithelial Communities Share a Core Bacterial Microbiota: A Meta-Analysis of 16S rRNA Gene Illumina MiSeq Sequencing Datasets. *Front. Microbiol.* **12**, 625400 (2021).

55. Wu, D. *et al.* Host and altitude factors affect rumen bacteria in cattle. *Braz. J. Microbiol.* **51**, 1573–1583 (2020).

56. Martínez-Álvaro, M. *et al.* Bovine host genome acts on rumen microbiome function linked to methane emissions. *Commun. Biol.* **5**, 350 (2022).

57. Cholewińska, P., Czyż, K., Nowakowski, P. & Wyrostek, A. The microbiome of the digestive system of ruminants – a review. *Anim. Health Res. Rev.* **21**, 3–14 (2020).

58. Patra, A. K. & Yu, Z. Effects of garlic oil, nitrate, saponin and their combinations supplemented to different substrates on *in vitro* fermentation, ruminal methanogenesis, and abundance and diversity of microbial populations. *J. Appl. Microbiol.* **119**, 127–138 (2015).

59. Zhu, W. *et al.* Garlic skin induces shifts in the rumen microbiome and metabolome of fattening lambs. *Animal* **15**, 100216 (2021).

60. Rabee, A. E., Sayed Alahl, A. A., Lamara, M. & Ishaq, S. L. Fibrolytic rumen bacteria of camel and sheep and their applications in the bioconversion of barley straw to soluble sugars for biofuel production. *PLOS ONE* **17**, e0262304 (2022).

61. Rabee, A. E., Forster, R. & Sabra, E. A. Lignocelluloytic activities and composition of bacterial community in the camel rumen. *AIMS Microbiol.* **7**, 354–367 (2021).

62. Accetto, T. & Avguštin, G. Inability of Prevotella bryantii to Form a Functional Shine-Dalgarno Interaction Reflects Unique Evolution of Ribosome Binding Sites in Bacteroidetes. *PLoS ONE* **6**, e22914 (2011).

63. Atasoglu, C., Valdés, C., Walker, N. D., Newbold, C. J. & Wallace, R. J. De Novo Synthesis of Amino Acids by the Ruminal Bacteria *Prevotella bryantii* B$_1$4, *Selenomonas ruminantium* HD4, and *Streptococcus bovis* ES1. *Appl. Environ. Microbiol.* **64**, 2836–2843 (1998).

64. Avgustin, G., Wright, F. & Flint, H. J. Genetic Diversity and Phylogenetic Relationships among Strains of Prevotella (Bacteroides) ruminicola from the Rumen. *Int. J. Syst. Bacteriol.* **44**, 246–255 (1994).

65. Maia, M. R. G., Chaudhary, L. C., Figueres, L. & Wallace, R. J. Metabolism of polyunsaturated fatty acids and their toxicity to the microflora of the rumen. *Antonie Van Leeuwenhoek* **91**, 303–314 (2007).

66. Paillard, D. *et al.* Quantification of ruminal *Clostridium proteoclasticum* by real-time PCR using a molecular beacon approach. *J. Appl. Microbiol.* **103**, 1251–1261 (2007).

67. Hivarkar, S. S., Vasudevan, G., Dhakephalkar, P. K. & Dagar, S. S. *Description of Sporanaerobium Hydrogeniformans Gen. Nov., Sp. Nov., an Obligately Anaerobic, Hydrogen Producing Bacterium of Lachnospiraceae Family Isolated from Aravali Hot Spring in India*. https://www.researchsquare.com/article/rs-2832220/v1 (2023) doi:10.21203/rs.3.rs-2832220/v1.

68. McCabe, M. S. *et al.* Illumina MiSeq Phylogenetic Amplicon Sequencing Shows a Large Reduction of an Uncharacterised Succinivibrionaceae and an Increase of the Methanobrevibacter gottschalkii Clade in Feed Restricted Cattle. *PLOS ONE* **10**, e0133234 (2015).

69. Cersoimo, L. M. & Wright, A.-D. G. Rumen Methanogens. in *Rumen Microbiology: From Evolution to Revolution* (Springer India, 2015).

70. Patra, A., Park, T., Kim, M. & Yu, Z. Rumen methanogens and mitigation of methane emission by anti-methanogenic compounds and substances. *J. Anim. Sci. Biotechnol.* **8**, 13 (2017).

71. Janssen, P. H. & Kirs, M. Structure of the archaeal community of the rumen. *Appl. Environ. Microbiol.* **74**, 3619–3625 (2008).

72. Smith, R. H., Glendinning, L., Walker, A. W. & Watson, M. Investigating the impact of database choice on the accuracy of metagenomic read classification for the rumen microbiome. *Anim. Microbiome* **4**, 57 (2022).

73. Johnson, J. S. *et al.* Evaluation of 16S rRNA gene sequencing for species and strain-level microbiome analysis. *Nat. Commun.* **10**, 5029 (2019).

74. Bryant, M. P. & Burkey, L. A. Cultural methods and some characteristics of some of the more numerous groups of bacteria in the bovine rumen. *J. Dairy Sci.* **26**, 205–217 (1952).
75. Hobson, P. N. Rumen Bacteria. *Methods Microbiol.* **3b**, 133–149 (1969).
76. Caldwell, D. R. & Bryant, M. P. Medium Without Rumen Fluid for Nonselective Enumeration and Isolation of Rumen Bacteria. *Appl. Microbiol. Sept* **14**, (1966).
77. Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl. Environ. Microbiol.* **79**, 5112–20 (2013).
78. McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLoS ONE* **8**, e61217 (2013).
79. phyloseq_ Explore microbiome profiles using R.pdf.
80. Haverkamp, T. H. A. Analyzing the Mothur MiSeq SOP dataset with Phyloseq. *Repository for bioinformatics training at the Norwegian Veterinary Institute: Analyzing the Mothur MiSeq SOP dataset with Phyloseq* https://norwegianveterinaryinstitute.github.io/BioinfTraining/phyloseq_tutorial.html (2018).
81. Hui, Y. Tutorial for microbiome analysis in R. https://www.yanh.org/2021/01/01/microbiome-r/ (2021).
82. Ollberding, N. Introduction to the Statistical Analysis of Microbiome Data in R. https://www.nicholas-ollberding.com/post/introduction-to-the-statistical-analysis-of-microbiome-data-in-r/ (2019).