Dynamic convergence of neurodevelopmental disorder risk genes across
neurodevelopment

AUTHORS

Meilin Fernandez Garcia[1*], Kayla Retallick-Townsley[1,2*], April Pruitt[3*], Elizabeth
Davidson[4*], Novin Balafkan[1,5*], Jonathan Warrell[6], Tzu-Chieh Huang[1,16], Alfred
Kibowen[1], Zhiyuan Chu[6], Yi Dai[7], Sarah E. Fitzpatrick[3,8], Ran Meng[6], Annabel Sen[1],
Sophie Cohen[1], Olivia Livoti[1], Suha Khan[8], Charlotte Becker[1], Andre Luiz Teles e
Silva[1,9], Jenny Liu[1], Grace Dossou[8], Jen Cheung[1], Susanna Liu[6], Sadaf Ghorbani[1], P.J.
Michael Deans[1], Marisa DeCiucis[10,11,16], Prashant Emani[6], Huanyao Gao[1], Hongying
Shen[10,11,16], Mark Gerstein[6,12], Zuoheng Wang[7,12], Laura M. Huckins[1,2,3], Ellen J.
Hoffman[4,13#], Kristen Brennand[1,2,3,14,15,16#]

AFFILIATIONS

[1] Department of Psychiatry, Division of Molecular Psychiatry, Yale University School of
Medicine, New Haven, CT 06511

[2] Pamela Sklar Division of Psychiatric Genomics, Department of Genetics and
Genomics, Icahn Institute of Genomics and Multiscale Biology, Nash Family
Department of Neuroscience, Friedman Brain Institute, Icahn School of Medicine at
Mount Sinai, New York, NY 10029

[3] Interdepartmental Neuroscience Program, Yale School of Medicine, New Haven, CT
06511

[4] Child Study Center, Yale University School of Medicine, New Haven, CT 06511

[5] Mohn Research Centre for Regenerative Medicine, Haukeland University Hospital,
Bergen, Norway.

[6] Department of Molecular Biophysics & Biochemistry and Program in Computational
Biology and Bioinformatics, Yale University, New Haven, CT 06511

[7] Department of Biostatistics, Yale University School of Public Health, New Haven, CT
06510

[8] Medical Scientist Training Program, Yale School of Medicine, New Haven, CT, US

[9] Hospital Israelita Albert Einstein, São Paulo, SP, Brazil

[10] Department of Cellular & Molecular Physiology, Yale School of Medicine, New Haven,
CT, 06511

[11] Systems Biology Institute, Yale West Campus, West Haven, CT 06516

[12] Department of Biomedical Informatics and Data Science, Yale School of Medicine;
New Haven, CT 06511

[13] Department of Neuroscience, Yale University School of Medicine, New Haven, CT
06511

[14] Department of Genetics, Yale University School of Medicine, New Haven, CT 06511

[15] Wu Tsai Institute, Yale University School of Medicine, New Haven, CT 06511

[16] BD2: Breakthrough Discoveries for thriving with Bipolar Disorder, Santa Monica, CA

* These authors contributed equally

Co-corresponding author

CORRESPONDENCE

kristen.brennand@yale.edu and ellen.hoffman@yale.edu

KEYWORDS

ABSTRACT

Over three hundred and seventy-three risk genes, broadly enriched for roles in neuronal
communication and gene expression regulation, underlie risk for autism spectrum
disorder (ASD) and developmental delay (DD). Functional genomic studies of subsets
of these genes consistently indicate a convergent role in neurogenesis, but how these
diverse risk genes converge on a smaller number of biological pathways in mature
neurons is unclear. To uncover shared downstream impacts between
neurodevelopmental disorder (NDD) risk genes, here we apply a pooled CRISPR
approach to contrast the transcriptomic impacts of targeting 29 NDD loss-of-function
genes across human induced pluripotent stem cell (hiPSC)-derived neural progenitor
cells, glutamatergic neurons, and GABAergic neurons. Points of convergence vary
between the cell types of the brain and are greatest in mature glutamatergic neurons,
where they broadly target not just synaptic and epigenetic, but unexpectedly,
mitochondrial biology. The strongest convergent networks occur between NDD genes
with common co-expression patterns in the post-mortem brain, biological annotations,
and clinical associations, suggesting that convergence may one-day inform patient
stratification and treatment. Towards this, ten out of eleven drugs tested that were
predicted to reverse convergent signatures in human cells and/or arousal and sensory
processing behaviors in zebrafish ameliorated at least one behavioral phenotype *in*
vivo. Altogether, robust convergence in post-mitotic neurons represents a clinically
actionable therapeutic window.

INTRODUCTION

Autism spectrum disorder (ASD) and related developmental delay (DD) are highly heritable[1]. The aggregate impact of common variants of small effect reflects most genetic risk[2], but in as many as a quarter of cases, potentially damaging rare inherited and *de novo* mutations in risk genes are detected[3]. There is significant overlap between those genes affecting ASD[4] and those more broadly affecting developmental[5,6] and psychiatric[7,8] disorders. Altogether, neurodevelopmental disorder (NDD) risk genes are typically expressed during cortical development[9], particularly the excitatory and inhibitory lineages[4], and broadly split between two functional classes: neuronal communication (e.g., synaptic function) and gene expression regulation (e.g., chromatin regulators and transcription factors)[4,10-15]. Over half of NDD genes have roles in gene expression regulation[4], sharing substantial overlap in genomic binding sites in the brain[16], and with targets enriched for NDD risk genes[17-20]. Yet, evidence to support the parsimonious explanation that regulatory NDD genes preferentially target synaptic NDD genes, is lacking[4]. It remains unclear how disrupting NDD genes with distinct functions yields similar outcomes.

Many NDD genes seem to have broad roles outside their annotated function; for example, some chromatin regulators (e.g., *CHD8, CHD2,* and *POGZ*) localize to microtubules in the centrosome[21], mitotic spindle[22], and cilia[23,24], suggesting the possibility that they function directly in neurogenesis and/or synaptic biology. Indeed, both regulatory and synaptic genes impact proliferation and patterning of progenitors (e.g., *ARID1B*[25,26], *CHD8*[27,28], *NRXN1*[29,30], *SYNGAP1*[31]), excitatory transmission by glutamatergic neurons (e.g., *CHD8*[32,33], *NRXN1*[34], *SHANK3*[35], *SYNGAP1*[36]), and inhibitory transmission by GABAergic neurons (e.g., *ARID1B*[37], *CHD8*[32], *NRXN1*[38], *SHANK3*[39]). Do overlapping downstream impacts explain how heterogeneous gene mutations result in similar neuronal phenotypes and clinical outcomes[40]?

Many have proposed that diverse ASD genes are convergent[41-43]. Indeed, NDD genes are co-expressed in the brain[44-46], suggesting that they are regulated together and involved in related biological processes, and result in highly interconnected protein-protein interactomes[47-50], indicating functional relationships between NDD proteins. Even as the number of NDD genes grows, risk genes continue to converge on a finite number of biological pathways, developmental stages, brain regions and cell types[41]. Disentangling these complex etiologies remains an outstanding challenge.

Excitatory-inhibitory (E:I) imbalance is widely believed to underlie NDD[51-53], whether arising from altered proportions of neuronal lineage cell types in the developing brain or synaptic deficits in glutamatergic or GABAergic neurons. Indeed, knockdown of subsets of NDD genes in human neural progenitor cells (NPCs)[22,54,55], cerebral organoids[27,56,57], and developing mouse[58], tadpole[59] and zebrafish[60] brains reveal overlapping impacts on neurogenesis. Despite synaptic dysfunction being a hallmark of NDD, the extent to which downstream impacts of NDD genes also converge in mature neurons is largely unknown.

Given emerging evidence that epigenetic NDD genes have diverse and interconnected roles[21-24], we tested the hypothesis that the nature of convergence is dynamic, influenced by developmental and cell-type contexts. We report a pooled CRISPR-

knockout (KO) strategy targeting loss-of-function (LoF) mutations to 29 NDD genes,
most with roles in chromatin biology (*ANK3, ARID1B, ASH1L, ASXL3, BCL11A, CHD2,*
CHD8, CREBBP, DPYSL2, FOXP2, KMT5B (SUV420H1), KDM5B, KDM6B, KMT2C,
MBD5, MED13L, NRXN1, PHF12, PHF21A, POGZ, PPP2R5D, SCN2A, SETD5,
SHANK3, SIN3A, SKI, SLC6A1, SMARCC2, WAC) in induced NPCs, glutamatergic
neurons, and GABAergic neurons *in vitro*. We describe convergent networks that are
unique between cell types, and in neurons, enriched not just for synaptic biology, but
also epigenetic regulation and, unexpectedly, mitochondrial function. Novel applications
of machine learning allowed us to extend our analyses *in silico* across all known NDD
genes, resolving how the degree of convergence between risk genes was influenced by
clinical associations, biological function, and co-expression patterns in the post-mortem
brain. Convergent analyses resolved the genes and cell types that underlie *in vivo*
behavioral stratification and successfully predicted drugs capable of suppressing
phenotypes in mutant zebrafish, suggesting that precision medicine-based approaches
can successfully target shared downstream gene targets between multiple NDD genes.
Novel points of convergence in post-mitotic neurons represent exciting new therapeutic
targets occurring within a clinically actionable therapeutic window.

RESULTS

A systematic comparison of NDD gene effects across neuronal cell types

From 102 highly penetrant loss-of-function (LoF) gene mutations associated with NDD
(previously described as 58 gene expression regulation, 24 neuronal communication,
and 20 other)[4], we used gene ontology and primary literature to identify 21 epigenetic
modifiers specifically involved in chromatin organization, rearrangement, and
modification (*ASH1L, ARID1B, ASXL3, BCL11A, CHD2, CHD8, CREBBP, PPP2R5D,*
KDM5B, KDM6B, KMT2C, KMT5B (SUV420H1), MBD5, MED13L, PHF12, PHF21A,
SETD5, SIN3A, SKI, SMARCC2, WAC), as well as two transcription factors with
putative roles as chromatin regulators (*FOXP2, POGZ*). Three extensively studied
synaptic genes (*NRXN1, SCN2A, SHANK3*) and three under-explored neuronal
communication genes (*ANK3, DPYSL2, SLC6A1*) strongly associated with NDD were
added (**SI Fig. 1A**). Many of these 29 genes differed in relative frequency of LoF gene
mutations between ASD (n=16) and DD (n=4)[61], schizophrenia[62], and epilepsy[63,64] (**Fig.**
1A-B, SI Fig. 1B), as well as general associations with GWAS for many
neuropsychiatric disorders (MAGMA[65]) (**Fig. 1C; SI Fig. 1C**), indicating a pleotropic
effect consistent with the shared genetic liability across neuropsychiatric disorders[66].
iNPCs, iGLUTs, and iGABAs (**SI Fig. 2A**), as well as their *in vivo* fetal counterparts (**SI**
Fig. 2B), expressed all genes prioritized herein[67].

Towards resolving whether regulatory genes confer continuous or distinct periods of
susceptibility across neurodevelopment, we knocked out (KO) regulatory NDD genes in
neural progenitor cells (SNaPs[68], here termed iNPCs), immature and mature
glutamatergic neurons (iGLUTs)[69], and mature GABAergic neurons (iGABAs)[70] (**Fig.**
1D). A pooled CRISPR approach (ECCITE-seq[71]) combined direct detection of sgRNAs
and single-cell RNA sequencing to compare loss-of-function effects across 29 NDD
genes. The CRISPR-KO library was generated from pre-validated gRNAs (three to four

gRNAs per gene; **SI Table 1**). Sequencing of the gRNA library confirmed the presence of gRNAs targeting 24 genes (*ANK3, ARID1B, ASH1L, ASXL3, BCL11A, CHD2, CHD8, DPYSL2, FOXP2, KMT5B (SUV420H1), KDM5B, KDM6B, KMT2C, MBD5, MED13L, NRXN1, PHF12, PHF21A, SCN2A, SETD5, SIN3A, SKI, SMARCC2, WAC*), but three (*DPYSL2, FOXP2, SCN2A*) were present at lower frequency (**SI Fig. 3B-C**).

Control hiPSCs were induced to iNPCs, iGLUTs, and iGABAs (**SI Fig. 3A**), transduced first with lentiviral-Cas9v2 (Addgene #98291) and subsequently with the pooled lentiviral gRNA library three days before harvest, at day 7 (iNPC and immature iGLUT), day 21 (iGLUT), and day 36 (iGABA) (experimental workflow **SI Fig. 4A;** computational workflow **SI Fig. 4B;** experimental validation of CRISPR editing efficiency in **SI Fig. 5**). After filtering and QC (**SI Fig. 4C-E**), we resolved NDD transcriptomes for 118,436 single cells: 25,402 iNPC, 38,097 immature (d7) iGLUT, 28,388 mature (d21) iGLUT, and 26,549 mature (d36) iGABA. Because original gene-expression based clustering was driven by cellular heterogeneity, cell quality, and sequencing lane effects (**SI Fig. 6A**), independent of gRNA identity, we removed cells with high expression of subtype markers and adjusted for cellular heterogeneity (**SI Fig. 6B,C**; **SI Tables 2-3**). 'Weighted-nearest neighbor' (WNN) analysis assigned clusters based on both gRNA identity class and gene expression to ensure that cells assigned to a gRNA identity class demonstrated successful perturbation of the targeted NDD gene[72]. For those WNN clusters where most cells were assigned to a single KO target, the transcriptomic signatures were compared to non-targeting scramble control clusters. Altogether, 35,777 cells were used for downstream analyses: 12,107 iNPC, 3,171 immature iGLUT, 11,802 mature iGLUT, and 8,697 mature iGABA). An average of 474 cells were assigned to each individual sgRNA (757 iNPC, 227 immature iGLUT, 562 mature iGLUT, 414 mature iGABA), totaling 33,150 perturbed cells and 2,627 controls (882 iNPC, 90 immature iGLUT, 1,258 mature iGLUT, and 397 mature iGABA). The gene expression patterns of non-perturbed iNPCs and iNeurons (>30% of all pooled cells) were significantly correlated with fetal brain cells and cortical adult neurons.

Successful perturbations (scCRISPR-KO) were identified for 23 NDD genes (**SI Fig. 6,7**): 16 in iNPCs, 14 in immature iGLUT neurons, and 21 in mature iGLUT and iGABA neurons (**SI Fig. 6**). Nine NDD genes were perturbed in all four cell types (*ARID1B, ASH1L, CHD2, MED13L, NRXN1, PHF21A, SETD5, SIN3A, SMARCC2*; **SI Fig. 7A,B**). For most NDD genes, KO in mature iGLUTs yielded the largest number of differentially expressed genes (DEGs, pFDR<0.05) (**SI Fig. 7B**), an effect that was not driven by differences in the extent of perturbation of the NDD gene itself between cell types (**SI Fig. 7Ci**). The transcriptomic effects of individual NDD genes cluster by cell type: the strongest NDD gene correlations are in mature iGLUTs (i.e., all nominally significant (p<0.01) log2FC DEGs are most highly correlated with each other and least correlated with the other cell types, whether relative to all scramble control cells (**Fig. 1Ei,ii; SI Fig. 7Cii**) or random subsets of scramble control cells (**SI Fig. 8A,B**). DEGs across individual NDDs shared significant gene ontology enrichments (**SI Fig. 8C**), with mature iGLUTs frequently enriched for SCZ GWAS genes (12 of 21 NDD genes), whereas mature iGABAs for migraine GWAS genes (8 of 21) (**SI Fig. 9**).

Unsurprisingly, given the greater within cell-type correlations between NDD genes and the unique pathway enrichments across cell-types, very few DEGs shared significance

and direction of effect for the same NDD gene perturbation across all four cell-types
(FDR adjusted p_{meta}<0.05, Cochran's heterogeneity Q-test p_{Het} > 0.05; computational
workflow, **SI Fig. 10A**); in fact, the only common DEG between cell types was frequently
the targeted NDD gene itself. With a more relaxed statistical threshold (nominal p-value
<0.05), modest shared effects of individual NDD genes could be resolved across cell
types. These effects rarely resulted in perturbation of the other NDD genes themselves
(**SI Fig. 10B**), showed very little overlap between NDD genes (**SI Fig. 10C**), and no
significant enrichments with psychiatric GWAS after multiple testing correction (**SI Fig.**
10D).

NDD gene knockouts resulted in cell-type-specific convergent genes and networks that
were strongest in glutamatergic neurons.

"Convergent genes" (**Fig. 2**) are those DEGs with significant and shared direction of
effect across all NDD gene perturbations (FDR adjusted p_{meta}<0.05, Cochran's
heterogeneity Q-test p_{Het} > 0.05)[73,74] (computational workflow, **Fig. 2A**). Across the nine
NDD genes perturbed in all four cell types (*ARID1B, ASH1L, CHD2, MED13L, NRXN1,*
PHF21A, SETD5, SIN3A, SMARCC2), convergence was highly cell-type specific (**Fig.**
2; SI. Fig. 11A-C; SI Data 2). Although the strength of convergence correlated across
cell types (**Fig. 2C,ii**), it was greatest in mature iGLUTs (quantified as the ratio of
convergent genes to the average number of DEGs across all 152 unique two-to-five
gene combinations of these nine NDD genes) (**Fig. 2C,i**).The unique "top" convergent
genes (**Table 1**) showed little overlap across all cell-types, with mature iGLUTs (11,473)
having the largest absolute number of convergent genes (**Fig. 2D**). Convergent genes
were enriched for schizophrenia GWAS loci (MAGMA[65], FDR <0.05) (**Fig. 2Ei**), rare
ASD and FMRP target genes (FDR <0.05) (**Fig. 2E,ii**), and pathways involved in
neurodevelopment, mitochondrial function, and translational regulation **(SI Fig. 12).**
When tested again across the 21 NDD genes perturbed in both iGLUTs and iGABAs,
mature iGLUTs again showed the largest absolute number of convergent genes
(iGLUTs, 10,557, **SI Fig. 13A;** iGABAs, 892, **SI Fig. 13B**). Intriguingly, although
convergent genes were highly cell-type-specific, those NDD gene combinations that
were highly convergent in one cell type were likely to be convergent in others; in
neurons, top convergent sets most frequently included *ARID1B, SETD5* and *NRXN1* **(SI**
Fig. 11D).

Given that the biological impact of convergence is likely to be impacted by the strength
of shared gene regulatory relationships and functions, we re-examined convergence
within the framework of co-expression networks (Bayesian bi-clustering). "Convergent
networks" (**Fig. 3**) are co-expressed genes that share similar expression patterns
across NDD gene perturbations[73,74] (computational workflow, **Fig. 3A**). The network
connectivity score ("network convergence") informs the strength and composition across
cell types (i.e., networks with more interconnectedness and containing genes with
greater functional similarity have increased convergence). Convergent networks
generated from the 9 NDD genes perturbed in all four cell types (**Fig. 3B**) or across the
21 NDDs genes in both iGLUTs and iGABAs (**SI Fig. 13C**) revealed the greatest
convergent network strength in iGLUTs. Network-level convergence was weakly
correlated between cell types **(Fig. 3C)**; the number of convergent unique network
nodes was greatest in iGLUTs, distinct across cell types (**Fig. 3D; Tables 2-4; SI Data**

2), and significantly enriched for rare variants linked to schizophrenia and ASD (**Fig. 3E;**
Tables 2-4). Convergent networks in iNPCs highlighted pathways associated with
neurogenesis (e.g., cell cycle, cell division, EPO signaling) (**Fig. 3F)**, while in mature
iGLUTs they were enriched for synaptic function (transmembrane transport and
receptor signaling, secretory vesicles, SNARE complex) (**Fig. 3G)**.

Convergent networks are strongest between NDD genes with shared co-expression
patterns in the post-mortem brain, biological annotations (synaptic or epigenetic), and
clinical outcomes (ASD or DD).

To resolve the extent to which functional similarity and co-expression patterns between
NDD genes predicted convergence, we trained a prediction model (random forest linear
regression)[75] using 70% of our data, evaluated it using 30% of our data, and validated in
an external dataset[73] (computational workflow, **Fig. 4A**; model predictor variables, **Fig.**
6B; more information **SI Fig. 14,15**). Cell type, brain co-expression (dorsolateral
prefrontal cortex, DLPFC), and functional similarity (i.e., gene ontology) correlate with
convergence (**Fig. 4C**) and well-predicted gene level convergence (97% variance
explained; mean of squared residuals (RMSE)=0.021) (**Fig. 4Di**) and moderately
predicted network-level convergence (53% variance explained; RMSE=0.73) (**Fig. 4Dii**).
Our trained model accurately predicted gene-level (Pearson's R=0.998, P<0.001,
RMSE=0.15) (**Fig. 4Ei; SI Fig. 15C**) and network-level convergence in our testing set
(R=0.72, P<2.2e-16, RMSE=0.85) (**Fig. 4Eii; SI Fig. 15D),** and performed moderately
well in predicting network-level convergence (R=0.26, P<0.001, RMSE=0.68) (**Fig. 4Fii;**
SI Fig. 15Eii) and to a lesser extent gene-level convergence (R=0.14, P<0.001
RMSE=1.75) (**Fig. 4Fi; SI Fig. 15Ei)** in the external dataset.

To query whether convergence reflected clinical associations to ASD or DD, we again
quantified convergence as the ratio of convergent genes to the average number of
DEGs (see **Fig. 2E**), here across all (2-5 gene) combinations of all NDD genes
perturbed in each cell type (e.g., 27,824 unique combinations of 21 NDD genes in
iGLUTs and iGABAs; **SI Fig. 16A**). Convergence, both gene-level (**SI Fig. 16A,C**) and
network-level (**SI Fig. 16B,D**)**,** was greater between genes with stronger associations to
ASD compared to DD[61]**,** particularly in mature neurons (**SI Fig. 16E-F**). Yet this analysis
was limited by the relatively small number of predominantly ASD (n=16) and DD (n=4)
included in our dataset (**Fig. 1B**).

To extend our comparisons of convergence across larger sets of NDD genes,
particularly those clinically defined as predominantly ASD or DD genes[61], or those with
biologically annotated synaptic or epigenetic roles[4], we asked if it was possible to train a
machine learning model to predict cell-type-specific impacts of CRISPR knockout of all
102 NDD genes[4]. An integrative Linear Network of Cell Type Phenotypes (**LNCTP**)
model, previously trained on >2.8 million nuclei from the prefrontal cortex across 388
individuals, accurately imputes single-cell expression following simulated
perturbations[76]. By retraining the LNCTP model using our scCRISPR-KO data (**Fig. 5A**),
we resolved convergent genes within three *in silico* post-mortem brain network models
(bulk prefrontal cortex (PFC) tissue, excitatory neurons only, and inhibitory neurons
only), noting that the LNCTP model better replicates experimental iGLUT data (**Fig. 5B**).

Expanded LNCTP *in silico* comparisons across all 102 NDD genes (**Fig. 2; SI. Fig. 17**)
predicted greater convergence in excitatory neurons compared to inhibitory neurons,
consistent with our *in vitro* findings (**Fig. 2C, 3D**), even more so for synaptic NDD genes
(n=24) relative to regulatory genes (n=58) (**Fig. 5C**). Predominantly ASD genes (n=50)
had greater predicted convergence in excitatory neurons (**Fig. 5C**), whereas
predominantly DD genes (n=40) in inhibitory neurons (**Fig. 5C**). Overall, across
functional or clinical categories, despite limited overlap in specific convergent genes
(**Fig. 5D**) and terms (**Fig. 5E, F**), there was overall enrichment for synaptic, epigenetic,
and mitochondrial biology (**Fig. 5G**), consistent with *in vitro* scCRISPR-KO (**Fig. 2F**).

Convergent genes and networks in glutamatergic neurons targeted synaptic, epigenetic,
and mitochondrial biology.

Convergent genes and networks revealed cell-type-specific disease (**Fig. 2E**) and
functional enrichments (**Fig. 2F, 5G,6A-B**), many consistent with established NDD
etiology in neurogenesis[22,27,54-60] and synaptic biology[47-50]. For example, iNPCs were
significantly enriched for pathways involved in proliferation and differentiation, whereas
mature iGLUTs showed unique enrichments in neuronal communication (e.g., pre-
synaptic function) and regulation of gene expression (e.g., mRNA processing and
protein translation). Unexpectedly, both mature iGLUT and iGABA neurons were
enriched for mitochondrial biology (e.g., oxidative phosphorylation: mature iGLUTs:
NEs=2.8, $p<2.2e-16$, FDR<0.001; mature iGABAs: NES=1.67, p=0.023, FDR<0.05)**.**

Functional validation of five NDD genes (*KMT5B, NRXN1, CHD8, ASH1L, ARID1B*) in
inducible Cas9 (iCas9)[77] NPCs (CD184$^+$/CD133$^-$ NPCs) in arrayed format revealed
effects on proliferation (Ki67; **Fig. 6C; SI Fig. 18A**), neurogenesis (NPCs:
CD184+/CD44-/CD24+, neurons: CD184-/CD44-/CD24+*; **SI Fig. 18B**), and gliogenesis
(astrocytes: CD184+/CD44+; **SI Fig. 18C)** that varied between genes. Likewise, a
pooled CRISPR analysis in iCas9 cortical organoids confirmed effects on neurogenesis,
again with variable effects between NDD genes (**SI Fig. 19**).

To assess how loss of NDD-associated genes affects mitochondrial function, we
performed a pooled CRISPR knockout screen using a nearly identical library (same
backbone, guide density, and control set) in the H1-iCas9 line. Transduced cells were
differentiated into NPCs and iGLUTs by day 21, stained with the Δψm-sensitive dye JC-
1, and sorted by fluorescence-activated cell sorting (FACS) into high- (PE-high) and
low- (FITC-high) membrane-potential fractions, following amplicon sequencing to
quantify gRNA representation in each fraction (**Fig. 6D**). Of the fifteen KOs, ten resulted
in elevated mitochondrial membrane potential (MPP) in both NPCs and iGLUTs, the
remaining five caused cell-type-specific impacts on mitochondrial membrane potential.
Pathway enrichment of the ten NDD genes that increased mitochondrial membrane
revealed a convergence on chromatin remodeling complexes, microRNAs, and
transcription factors.

For three NDD KOs (*ASH1L, ARID1B, NRXN1)*, we validated mitochondrial effects in
arrayed format, using a platform with the ability to resolve dose-dependent changes in
mitochondrial fragmentation following pharmacological insults **(SI Fig. 20**). By high
content imaging, we analyzed and quantified 1×10^4 mitochondria per genotype, with
morphological measurements taken for mitochondrial (TOMM20-positive) volume,

surface area, and sphericity (roundness) as well as total OXPHOS complex, within
neuronal dendrites (MAP2-positive) of mature (d21) iGLUTs. Among the three NDD
KOs, *ARID1B* resulted in increased mitochondrial networking (indicated by decreased
mitochondrial sphericity and increased branch length; one-way ANOVA, Šidák's
adjusted p=0.0213 and p=0.0081 respectively) concomitant with increased levels of
OXPHOS proteins (one-way ANOVA, Šidák's, adjusted p=0.0024) (**Fig. 6E; SI Fig.**
21A), overall consistent with increased mitochondrial efficiency. Second, we tested
oxygen consumption using Seahorse Cell Mito Stress test. NRXN1 KO resulted in
increased coupled and maximal respiration in iGLUTs (one-way ANOVA, p<0.05; **Fig.**
6F); increased mitochondrial reliance, in the absence of fused mitochondria with
elevated OXPHOS protein levels point to a possible metabolic overload due to reduced
mitochondrial efficiency (**Fig. 6E**). In contrast, ARID1B and ASH1L KOs did not show
significant changes in these Seahorse parameters (**SI Fig. 21B–C**). Taken together,
both ARID1B and NRXN1 KO neurons show evidence of increased mitochondrial
activity, ARID1B KO through enhanced fusion and elevated expression of OXPHOS
complexes, whereas NRXN1 KO by increasing OXPHOS activity to meet ATP
demands. As observed for neurogenesis in iNPCs, single gene knockouts iGLUTs
confirmed convergent effects on mitochondrial biology, finding distinct but related
phenotypes between NDD genes.

Pharmacological targeting of convergent genes reversed behavioral phenotypes in
mutant zebrafish

By design, *in vitro* models substantially limit the complexity of the observed impact of
NDD genes, lacking higher circuit-level effects. Towards applying molecular
convergence *in vitro* to explore the mechanisms of phenotypic convergence *in vivo,* the
convergence of sets of NDD genes were next explored on the basis of shared
behavioral effects in zebrafish mutants (**Fig. 7; SI Tables 4-5**). A comprehensive *in vivo*
high-throughput, automated behavioral analysis in larval zebrafish[60] revealed clear
stratification of NDD genes based on basic arousal and sensory processing behaviors in
the developing vertebrate brain (**Fig. 7A; SI Fig. 22**). Given that zebrafish brain gene
expression was significantly correlated with *in vitro* human-derived mature neurons (**Fig.**
7B; SI Fig. 23), we asked whether behavioral stratification of NDD mutants in larval
zebrafish can be attributed to molecular convergence. For fifteen NDD genes for which
we have matched behavioral and molecular analyses, zebrafish stable mutant lines and
CRISPR F0 mutants were clustered based on 24 sleep-wake and visual-startle
parameters, yielding four distinct clusters of genes: set 1 (*nrxn1a, mbd5, kdm5bab*),
set2 (*phf12ab, skiab, chd2, smarcc2*), set 3 (*kdm6bab, kmt5b, kmt2cab*), and set 4
(*wacab, arid1b, phf21aab, chd8, ash1l*) (**Fig. 7A; SI Data 3)**. Gene-level convergence
between NDD genes in these sets was distinct, largely non-overlapping between cell-
types, and stronger in mature iGLUTs than mature iGABAs (**Fig. 7C**). Across behavioral
sets, rare ASD, SCZ, and ID LoF genes were enriched primarily in iGLUTs, with all sets
converging on FMRP targets, highly intolerant CNVs, and ASD variants (**Fig. 7D**).
Phenotypes related to developmental delay, behavior, and motor function showed
unique enrichments by set, predominately in the iGLUTs, whereas all sets were
enriched for seizure, hypertonia, and abnormal skeletal muscle morphology (**Fig. 7E**).
Candidate drugs predicted to reverse convergent genes (i.e., drugs with anticorrelating
transcriptomic signatures) in iGLUTs and iGABAs were prioritized from the 776 cMAP[78]

drugs with matched clinical and experimental zebrafish data. Top enriched drugs
included antidepressants, antipsychotics, and statins (**SI Data 2; SI Fig. 24A**). Whereas
some drugs were broadly predicted to reverse convergent signatures in all four NDD
gene sets (e.g., the antipsychotic perphenazine), others uniquely targeted specific sets
(e.g., naltrexone in set 2 iGLUTs, sirolimus in set 3 iGLUTs, and valsartan in set 3
iGABAs). Sets 3 and 4 showed the greatest number of cMAP enrichments. By
considering existing pharmacological effects of the top drugs on zebrafish behavior,[60]
some of the predicted drug reversers were shown to oppose effects on NDD related
phenotypes in zebrafish (**S**l **Fig. 24B**). Yet, the direction of effect predicted based on
transcriptomic convergence in human neurons did not always align with anti-correlating
behavioral effects in zebrafish (e.g., moxifloxacin, perphenazine).

The top negatively enriched drugs for iGLUT convergence from cMAP and anti-
correlating drugs predicted from a pharmaco-behavioral screen of 376 drugs in larval
zebrafish were empirically tested in representative mutants from sets 2-4, which showed
the strongest cMAP enrichments (**Fig. 7F**). We determined whether the phenotypic
impact of mutant-x-drug combinations led to partial rescue, rescue, over-correction, or
exacerbation of the mutant phenotype across significant arousal and startle behavioral
parameters (**Fig. 7G**). Ten out of eleven drugs rescued at least one dysregulated
behavioral parameter (**Fig. 7G, SI Fig. 24C-E**). Paclitaxel robustly rescued behavioral
parameters in *kdm6bab* F0 mutants and pravastatin partially and completely rescued
select parameters in *chd2$^{\Delta7/\Delta7}$* mutants (**Fig. 7Gi**), including nighttime sleep bouts in
kdm6bab F0 mutants and responses to lights-ON stimuli in *chd2$^{\Delta7/\Delta7}$* mutants (**SI Fig.
24Fi-ii**). Interestingly, we also observed over-correction of the *phf21aab* F0 mutant
phenotype by fluvoxamine (**Fig. 7Gii**), such as increased sleep bouts that were
significantly decreased following fluvoxamine treatment (**SI Fig. 24Fiii**). Taken together,
in vivo behavioral profiling of NDD genes in zebrafish overlaps with *in vitro*-defined
convergent networks and identifies pharmacological suppressors of specific behavioral
phenotypes.

DISCUSSION

Towards empirically resolving the common pathways converged upon by NDD risk gene
effects, 29 NDD genes were targeted through a pooled CRISPR-KO strategy. The
molecular points of convergence across NDD risk genes varied between the cell types
of the brain, being greatest in mature glutamatergic neurons, where they were enriched
not just for pathways with well-established links to ASD etiology (e.g., gene regulation,
synaptic biology), but also mitochondrial function[79]. While downstream effects of
epigenetic NDD genes unexpectedly targeted mitochondrial genes, in fact, five percent
of NDD cases meet diagnostic criteria for classic mitochondrial disorders[80].
Mitochondrial DNA mutations[81,82], haplotypes[83] and heteroplasmy[81,84] have all been
associated with NDD. Not only do mitochondrial mutations cause synaptic and
behavioral phenotypes[85], but multiple lines of human and animal evidence link NDDs to
mitochondrial deficits and oxidative stress[10,86-91], with neuronal and/or behavioral
phenotypes reversed by antioxidant treatment[87,89-91]. Conversely, knockout of NDD
genes in NPCs primarily alter neurogenesis[54,57,59] and developmental dynamics[27,92]. Put
simply, perturbations of the same NDD genes resulted in different convergent networks

across cell types. This observation connects the pleiotropic nature of many NDD genes
and pathophysiological evidence linking multiple cell types and distinct cellular functions
to NDD.

What explains phenotypic convergence between NDD genes with distinct annotated
functions? The strength of convergence was most highly correlated to common clinical
associations, biological annotations, and co-expression patterns in the post-mortem
brain. Critically, these factors are inter-dependent. NDD risk genes most strongly
implicated in DD are enriched for expression in progenitor cells and immature neurons,
and those in ASD in mature neurons[61]. Indeed, cellular identities and biological
pathways are captured by patterns of gene co-expression[93,94]. Transcriptomic and
epigenomic analyses of post-mortem brain from NDD cases likewise indicate
convergent molecular signatures[95] and subtypes of NDD[96]. Thus, we posit that shared
clinical and phenotypic effects of distinct NDD genes in fact reflect the patterns of co-
expression in the developing brain.

Personalized medicine seeks to tailor treatments to individual patients[97]; for example,
cancer[98] and monogenic disease[99] patients with specific genetic mutations receive
targeted treatments. Previous efforts to classify genes that predict NDD clinical features
or treatment response applied gene ontology[4,61] or differential neurodevelopmental KO
effects *in vitro*[54] or *in vivo*[59]. Here, we proposed to stratify risk genes based on
convergent molecular impacts in human neurons. Our overarching hypothesis, in doing
so, was that by resolving shared downstream gene targets between multiple NDD
genes, we might inform a precision medicine-based approach that did not necessarily
need to target risk genes one-at-a-time. Although convergent networks did not predict
behavioral stratification of zebrafish mutants, they did inform drug prediction, with ten
out of eleven drugs tested found to ameliorate at least one mutant behavioral phenotype
in vivo. This ability to reverse, rather than prevent, a behavioral phenotype, indicates
that targeting convergent networks in post-mitotic neurons may represent a clinically-
actionable neurodevelopmental window that persists through symptom onset.   The
extent to which convergent downstream targets, whether associated with risk or
resilience, can be manipulated to prevent or ameliorate NDD signatures and
phenotypes warrants future investigation.

Although rare LoF NDD gene mutations tend to confer large effects in the individuals
who carry them, the small effects of common variants account for much of the genetic
risk for NDD at the population level[2,100]. The differences in expressivity and incomplete
penetrance of high effect-size rare variants is frequently attributed to diversity across
polygenic backgrounds[101]; *in vitro*, NDD gene effects are indeed influenced by the
individual genomic context[27]. In psychiatry, common genetic variants are more
associated with cross-disorder behavioral dimensions[102] and rare variants with co-
occurring intellectual disability[103]. Common risk variants interact with rare mutations to
determine individual-level liability in ASD[104-106], schizophrenia[107,108], epilepsy[109],
Huntington's disease[110] and more[111]. Our results, highlighting that convergence
downstream of NDD gene effects are enriched for cross-disorder GWAS variants and
rare LoF genes, inform pleiotropy of genetic risk for psychiatric disorders. Moving
forward, we argue that it is critical that empirical functional genomic studies
systematically consider the impact of common and rare variants together, including

screening the impact of LoF genes in hiPSC lines derived from donors with high and low
polygenic risk scores[112]. Intriguingly, even susceptibility to environmental risk factors for
NDD (e.g., valproic acid[113]) seems to be mediated by genetic background[114]. Deeper
phenotypic characterization of NDD effects across donors will be critical in determining
how complex genetic (or environmental) interactions shape cellular phenotypes, circuit
function, and human behavior in the clinic.

In the post-mortem brain, NDD gene signatures are not just associated with
downregulation of co-expression modules involving synaptic signalling[115], but also
upregulation of microglial and astrocyte gene modules[88,96,115-120]. The extent to which
increased neuroimmune activity in NDD is a response to cellular or environmental
sources of inflammation, or indicative of a role for glia cells in risk is unclear; evidence
supports both possibilities. Consistent with a model of maternal immune activation
during neurodevelopment[121], glucocorticoids and inflammatory cytokines perturb the
expression of psychiatric risk genes[122,123], altering the regulatory activity of psychiatric
risk loci[124], and interfering with neuronal maturation in brain organoids[125]. Yet, *in vivo*
analysis of NDD genes in zebrafish revealed global increases in microglia[60] and *in vitro*
screening in human microglia uncovered roles in endocytosis and uptake of synaptic
material[126]. Indeed, given the reciprocal relationships between neuronal activity and glial
function, epigenetic state, and gene expression[127-130], it seems probable that both cell-
autonomous and non-cell-autonomous effects underlie and/or exacerbate NDD gene
effects.

In summary, we demonstrate that convergent effects of NDD risk genes vary between
cell types. Our analyses suggest that clinical convergence between regulatory and
synaptic genes in the etiology of NDD is driven more so by co-expression patterns of
risk genes then direct regulation of epigenetic genes on synaptic targets. If the
convergence of multifold risk genes on a smaller number of shared molecular pathways
indeed explains how genetically heterogeneous mutations result in similar clinical
features, then genetic stratification of cases will inform novel therapeutic targets. We
predict that such individualized points of therapeutic intervention may be most effective
when targeting mature glutamatergic neurons, which not only harbor the strongest
convergent effects but also represent a therapeutic window that is actionable after
diagnosis.

MATERIALS AND METHODS

Generation of neural cells: Informed consent was obtained at the National Institute of
Mental Health, under the review of the Internal Review Board of the NIMH. hiPSC work
was reviewed by the Internal Review Board of the Icahn School of Medicine at Mount
Sinai as well as by the Embryonic Stem Cell Research Oversight Committee at the
Icahn School of Medicine at Mount Sinai and Yale University. Fibroblasts were
genotyped by IlluminaOmni 2.5 bead chip genotyping[131,132], PsychChip[133], and exome
sequencing[133]; hiPSCs[133] were validated by G-banded karyotyping (Wicell
Cytogenetics) and genome stability monitored by Infinium Global Screening Array v3.0
(Illumina). SNP genotype was inferred from all RNAseq data using the Sequenom
SURESelect Clinical Research Exome (CRE) and Sure Select V5 SNP lists to confirm

that neuron identity matched donor. Control hiPSCs were cultured in StemFlex media
(Gibco, #A3349401) supplemented with Antibiotic-Antimycotic (Gibco, #15240062) on
Geltrex-coated plates (Gibco, #A1413302). Cells were passaged at 80-90% confluence
with 5mM EDTA (Life Technologies #15575-020) for 3 min at room temperature (RT).
EDTA was aspirated and cells dissociated in fresh StemFlex media. Media was
replaced every 48-72 hours for 4-7 days until the next passage.

Transient transcription factor overexpression from stable clonal hiPSCs was used to
induce control hiPSCs to iNPCs (here SNaPs)[68], iGLUTs[69], and iGABAs[70]. iNPCs are
rapidly generated by 48-hour induction with *NGN2*[68,134]. iGLUTs are induced via
transient overexpression of *NGN2*, and are >95% glutamatergic neurons, robustly
express excitatory genes, and show spontaneous excitatory synaptic activity by three-
to-four weeks *in vitro*[29,34,35,67,69,135-141]. iGABA neurons are induced via transient
overexpression of *ASCL1* and *DLX2,* and are >95% GABAergic neurons, robustly
express inhibitory genes, and show spontaneous inhibitory synaptic activity by five-to-
six weeks[38,70,137,142,143]. iNPCs, iGLUTs, and iGABAs express most NDD genes,
including all genes prioritized herein[67].

We transduced hiPSCs from two control donors (553-3, karyotypic XY; 3182-3,
karyotypic XX) with lentiviral *pUBIQ-rtTA* (Addgene #20342) and *tetO-NGN2-eGFP-*
NeoR (Addgene #99378) for iNPCs and iGLUTs, or *pUBIQ-rtTA* (Addgene #20342),
tetO-ASCL1-PuroR (Addgene #97329)*,* and *tetO-DLX2-HygroR* (Addgene #97330) for
iGABAs. Following transduction by spinfection at 1000g for 1 hour at 37^{o}C, hiPSCs
were subjected to 48-hour antibiotic selection (1mg/mL neomycin G418 (Thermo
#10131027), 0.5µg/mL puromycin (Thermo #A1113803), and/or 250µg/mL hygromycin
(Thermo, #10687010) and then clonalized by expansion from single colonies.
Ultimately, clonal and inducible iNPC/iGLUT 3182-3-clone5 (XX) and iGABA 553-3-
clone34 (XY) hiPSCs were validated lentiviral genome integration by PCR, doxycycline
induced transcription factor expression by qPCR, and robust and consistent neuronal
induction confirmed by RNA-seq and immunocytochemistry for relevant cell type
markers. Analyses throughout reflect data from iGLUT 3182-3-clone5 (iNPC, d7 iGLUT
and d21 iGLUT) and iGABA 553-3-clone34 (d36 iGABA).

iNPCs: At DIV0, 3182-3-clone5 hiPSCs were dissociated and plated at 1.5 x 10^{6} cells
per well onto Geltrex-coated 6-well plates (1:250 dilution coating) in SNaP Induction
Media (DIV0): DMEM/F12 with Glutamax (ThermoFisher, 11320082), Glucose (0.3%
v/v), N2 Supplement (1:100, ThermoFisher, 17502048), Doxycycline (2 µg/mL; Sigma-
Aldrich, D9891), LDN-193189 (200 nM; Stemgent, 04-0074), SB431542 (10 µM; Tocris,
1614), and XAV939 (2 µM; Stemgent, 04-00046) supplemented with 25 ng/mL Chroma I
ROCK2 Inhibitor. After 24 hours, DIV2, cells were fed with Selection Media: DMEM/F12
with Glutamax, Glucose (0.3% v/v), N2 Supplement (1:100), Doxycycline (2 µg/mL),
Geneticin (0.5 mg/mL; Thermofisher, 10131035), LDN-193189 (100 nM), SB431542 (5
µM), and XAV939 (1 µM). After 48 hours post induction (DIV2), SNaPs were dissociated
with Accutase for 10 minutes at 37°C, quenched in DMEM, pelleted at 800g for 5
minutes, and replated at 1.5x10^{6} cells per well onto Geltrex-coated 6-well plates in
SNaP Selection Media supplemented with Geneticin (0.5 mg/mL). After 16-18 hr (DIV3),
medium was switched to SNaP maintenance Medium: DMEM/F12 with Glutamax,
Penn/Strep (1:100), MEM-NEAA (1:100; Life Technologies, 10370088), B27 minus

Vitamin A (1:50; Life Technologies, 12587010), N2 Supplement (1:100; Life
Technologies, 17502048), recombinant human EGF (10 ng/mL; R&D Systems, 236-EG-
200), recombinant human basic FGF (10 ng/mL; Life Technologies, 13256029),
Geneticin (0.5 mg/mL), and Chroman I (25 ng/mL). Cells were fed every 48 hours with
SNaP maintenance medium lacking Chroman I and Geneticin. Cells were dissociated
and seeded weekly at a density of 1.25-1.5x10^6 cells per well onto Geltrex-coated 6-well
plates until NPC morphology was observed and persistent. Cells were expanded and
cryofrozen.

DIV7 iGLUTs: 3182-3-clone5 iNPCs were thawed and seeded at 1x 10^6 cells per well
onto Geltrex-coated 12-well plates. *NGN2* expression was induced with Doxycycline (2
µg/mL) for 24 hrs (DIV0) with antibiotic selection for 48 hrs (DIV1-3) in SNaP
maintenance medium. At DIV 4 SNaPs were dissociated with Accutase, switched into
Neuronal Medium: Brainphys (Stemcell, 05790), Glutamax (1:100), Sodium Pyruvate (1
mM), Anti-Anti (1:100), N2 (1:100), B27 without vitamin A (1:50), BDNF (20 ng/mL;
R&D, 248-BD-025), GDNF (20 ng/mL; R&D, 212), dibutyryl cAMP (500 µg/mL; Sigma,
D0627), L-ascorbic acid (200 µM; Sigma, A4403), Natural Mouse Laminin (1.2 µg/m;
Thermofisher, 23017015) and seeded in Geltrex-coated (1:120 dilution coating) 12-well
plates. Medium was changed every 24 hrs until DIV7 harvest.

D21 iGLUTs: hiPSCs were harvested in Accutase (Innovative Cell Technologies, AT-
104) for 5 minutes 37°C, dissociated into a single-cell suspension, quenched in DMEM,
pelleted via centrifugation for five minutes at 1000 rcf and resuspended in StemFlex
containing 25 ng/mL Chroma I ROCK2 Inhibitor and 2.0 µg/mL doxycycline (DIV0),
seeded 1 x 10^6 cells per well onto Geltrex-coated 6-well plates (1:250 dilution coating),
and incubated overnight at 37°C. The next day, DIV1, hiPSCs were subjected to 48-
hour antibiotic selection by medium replacement with Induction Media: DMEM/F12
(Thermofisher, 10565018), Glutamax (1:100; Thermofisher, 10565018), N-2 (1:100;
Thermofisher, 17502048), B27 without vitamin A (1:50; Thermofisher, 12587010),
Antibiotic-Antimycotic (1:100) with 1.0µg/mL doxycycline and 0.5mg/ml Geneticin. At
DIV3, cells were treated with 4.0µM cytosineβ-D-arabinofuranoside hydrochloride (Ara-
C) and 1.0µg/mL doxycycline to arrest proliferation and eliminate non-neuronal cells in
the culture. At DIV4 immature neurons were dissociated with Accutase and 5 units/mL
DNAse I at 37°C for 7-10 min, quenched in DMEM, centrifuged for five minutes at 1,500
rpm and resuspended in 25 ng/mL Chroma I ROCK2 Inhibitor, 1.0 µg/mL doxycycline
and 4.0µM Ara-C and switched to Neuron Medium: Brainphys (Stemcell, 05790),
Glutamax (1:100), Sodium Pyruvate (1 mM), Anti-Anti (1:100), N2 (1:100), B27 without
vitamin A (1:50), BDNF (20 ng/mL; R&D, 248-BD-025), GDNF (20 ng/mL; R&D, 212),
dibutyryl cAMP (500 µg/mL; Sigma, D0627), L-ascorbic acid (200 µM; Sigma, A4403),
Natural Mouse Laminin (1.2 µg/mL; Thermofisher, 23017015) and seeded 7 x 10^5 cells
per well onto Geltrex-coated (1:60 dilution coating) 12-well plates and incubated
overnight at 37°C. The next day, DIV 6, Chroman I was removed from culture and Ara-C
lowered to 2.0 µM with a full Neuronal medium change. At DIV 7 a full Neuronal Medium
change was performed to remove doxycycline and Ara-C from culture, to allow for
antibiotic resistant genes silencing. From DIV7 onwards, half neuronal medium changes
were performed every 72 – 96 hrs until mature DIV 21 for harvest.

DIV36 iGABAs: hiPSCs were harvested in Accutase (Innovative Cell Technologies, AT-
104) for 5 minutes 37°C, dissociated into a single-cell suspension, quenched in DMEM,
pelleted via centrifugation for five minutes at 1000 rcf and resuspended in StemFlex
containing 25 ng/mL Chroma I ROCK2 Inhibitor and 2.0 µg/mL doxycycline (DIV0),
seeded 1.5-2x 10^6 cells per well onto Geltrex-coated 6-well plates (1:250 dilution
coating), and incubated overnight at 37°C. The next day, DIV1, hiPSCs were subjected
to 48-hour antibiotic selection by medium replacement with Induction Media: DMEM/F12
(Thermofisher, 10565018), Glutamax (1:100; Thermofisher, 10565018), N-2 (1:100;
Thermofisher, 17502048), B27 without vitamin A (1:50; Thermofisher, 12587010),
Antibiotic-Antimycotic (1:100) with 1.0µg/mL doxycycline, 1.0 µg/mL puromycin (Sigma,
P7255) and 250 µg/mL hygromycin (Sigma, 10687010). At DIV3, cells were treated with
4.0µM cytosineβ-D-arabinofuranoside hydrochloride (Ara-C) and 1.0µg/mL doxycycline
to arrest proliferation and eliminate non-neuronal cells in the culture. At DIV5 immature
neurons were dissociated with Accutase and 5 units/mL DNAse I at 37°C for 7-10 min,
quenched in DMEM, centrifuged for five minutes at 1,500 rpm and resuspended in 25
ng/mL Chroma I ROCK2 Inhibitor, 1.0 µg/mL doxycycline and 4.0µM Ara-C and
switched to Neuron Medium: Brainphys (Stemcell, 05790), Glutamax (1:100), Sodium
Pyruvate (1 mM), Anti-Anti (1:100), N2 (1:100), B27 without vitamin A (1:50), BDNF (20
ng/mL; R&D, 248-BD-025), GDNF (20 ng/mL; R&D, 212), dibutyryl cAMP (500 µg/mL;
Sigma, D0627), L-ascorbic acid (200 µM; Sigma, A4403), Natural Mouse Laminin (1.2
µg/mL; Thermofisher, 23017015) and seeded 7 x 10^5 cells per well onto Geltrex-coated
(1:60 dilution coating) 12-well plates and incubated overnight at 37°C. The next day,
DIV 6, Chroman I was removed from culture and Ara-C lowered to 2.0 µM with a full
Neuronal medium change. At DIV 7 a full Neuronal Medium change was performed to
remove doxycycline and Ara-C from culture, to allow for antibiotic resistant genes
silencing. From DIV7 onwards, half neuronal medium changes were performed every
72-96 hrs until mature DIV 36 for harvest.

CRISPR knockout gRNA library design (Thermofisher) and validation
From the 102 highly penetrant loss-of-function (LoF) gene mutations associated with
ASD (58 gene expression regulation, 24 neuronal communication genes, 9 cytoskeletal
genes, and 11 multifunction genes)[4], gene ontology and primary literature research
identified 26 epigenetic modifiers specifically involved in chromatin organization,
rearrangement, and modification. ASD gene expression (RNA-seq RPKM in iGLUTs)
was plotted against significance of ASD association (TADA FDR Values), to ensure
selection of genes with the highest expression and highest clinical association. Gene
expression was confirmed across development in the brain (BrainSpan[144]), and in bulk
and scRNA-seq. 21 epigenetic modifiers (*ASH1L, ASXL3, ARID1B, CHD2, CHD8,*
CREBBP, KDM5B, KDM6B, KMT2C, KMT5B, MBD5, MED13L, PHF12, PHF21A,
POGZ, PPP2R5D, SETD5, SIN3A, SKI, SMARCC2, WAC,) as well as two transcription
factors with putative roles as chromatin regulators (*FOXP2, BCL11A*) were selected.
Gene regulatory transcription factors, general transcription factors, and DNA replication
genes were excluded. Three extensively studied synaptic genes (*NRXN1, SCN2A,*
SHANK3) with roles in ASD were included as positive controls and three under-explored
genes for ASD role in neuronal communication genes (*ANK3, DPYSL2, SLC6A1*) were
also included in the library.

Individual DNA from glycerol stocks of Invitrogen™ LentiArray™ Human CRISPR
Library gRNAs-PuroR (ThermoFisher, A31949) (3-4 individual gRNAs per gene, see **SI**
Table 1) were prepared using GeneJET Plasmid Miniprep Kit (K0503) and pooled at an
equimolar ratio and a 5-fold ratio of scramble control gRNA plasmid. Library quality was
confirmed by restriction enzyme digest (10x Cutsart NEB), agarose gel purification
using QIAquick Gel Extraction Kit (#28706) to check library purity, followed by Mi-seq
for gRNA count distribution. Based on the abundance of gRNAs from Mis-seq, 4 NDD
gene targets were highly unlikely to be resolved in the final experiments – *POGZ,*
PP2R5D, SHANK3, SLC6A1 – and 3 with low abundance and less likely to be resolved
(*SCNA2, FOXP2, DYPSL2*).

Lentiviral Cas9v2-HygroR (Addgene, 98291) and pooled LentiArray-gRNA-PuroR
CRISPR-KO library were packaged as high-titer lentiviruses (Boston Children's Hospital
Viral Core) and experimentally titrated in each cell type. Highest viable MOI was used
for Cas9v2 and MOI < 0.5 for lentivirus gRNAs pool library.

CRISPR and gRNA delivery: Lentiviral Cas9v2-HygroR (Addgene #98291) transduction
of iNPCs, day 4 (iGLUTs), or day 5 (iGABAs) occurred via spinfection (one hour at
1,000 g) and followed by 72 hr hygromycin (250 µg /mL) (except for iGABAs, which
express inducible hygromycin resistance at this stage). Pooled Invitrogen™
LentiArray™ Human gRNA-PuroR CRISPR-KO Library gRNAs (ThermoFisher
#A31949) (MOI 0.3-0.5) were transduced via spinfection three days prior to harvest
(e.g., d4 for D7 iGLUTs, d18 for D21 iGLUTs, d33 for d36 iGABA), with fresh medium
containing puromycin (1 µg/mL) added 16-24 hours post transduction of gRNAs. For
mature iGLUTs and iGABAs, as doxycycline was removed from medium at DIV7, and
by DIV18 neurons had lost transcription factor linked antibiotic resistance, at 24 hours
post-transduction (DIV19 or DIV34) puromycin (1 µg/mL) and hygromycin (250 µg /mL)
were added to media for 48-hr antibiotic selection prior to harvest.

Dissociation of different neural cell types to single cells for scRNAseq assays: Cells
were dissociated 72 hrs post gRNA library delivery for single cell sequencing, as iNPCs,
DIV7 and DIV21 iGLUTs, or DIV36 iGABAs as follows:

iNPCs and DIV7 iGLUTs were dissociated in accutase for 5min @37°C, washed with
DMEM/10%FBS, centrifuged at 1,000xg for 5 min, gently resuspended, and counted.

DIV21 iGLUTs and DIV36 iGABAs were dissociated with papain. Papain was pre-
warmed (39°C) for 30 minutes in HBSS (ThermoFisher, 14025076), HEPES (10 mM,
pH 7.5) EDTA (0.5 mM), Papain (0.84 mg/mL; Worthington-Biochem, LS003127). The
cells were washed with PBS-EDTA (0.5 mM) and 300 uL of papain solution and 5 units
of DNAse I was added per well of 12-well plate and incubated at 37°C for 10-15
minutes, 125 rpm. Dissociation was quenched with DMEM-10%FBS. Detached neurons
were broken by gentle manual pipetting, pelleted at 600 g for 5 minutes, resuspended in
DMEM-10%FBS, filtered through a cell strainer and counted and submitted for 10X
sequencing.

Cells were loaded into 10X in four lanes per cell type, targeting 20,000 cells per lane for
a total of ~80,000 targeted cells per cell type. scRNA-seq was performed at Yale
Genomics Core with the 10X single cell 5' v2 HT with CRISPR barcode kit.

Bulk RNAseq and CRISPR-editing efficiency evaluation: The H1 hESC line with iCas9
(NIHhESC-10-0043), generously provided by the Huangfu Lab, was used to assess the
editing efficiency of the gRNAs[77,145] and conduct the mitochondrial pooled and arrayed
experiments. NPCs were generated using the dual SMAD inhibition approach per the
STEMdiff SMADi Neural Induction Kit protocol (STEMCell Technologies, #08581). To
validate gene KO, NPCs were transduced with LV particles carrying four gRNAs per
target gene. After 48 h of selection with 1 µg/mL puromycin, Cas9 expression was
induced by adding dox at 2 µg/mL for 72 h. Following induction, cells were collected for
bulk RNA-seq. Total RNA was extracted using TRIzol™ reagent (Invitrogen). PolyA
RNA-seq library preparation and sequencing were conducted at the Yale Center for
Genomic Analysis (YCGA). Raw fastq files were quality-checked by FastQC, then
mapped to human genome reference hg38 (STAR[146]). GRNA targeted-loci for each
sample were extracted (SAMtools[147]). Variation/small insertion/deletion at site of interest
and mutation efficiency at corresponding loci was called (CrispRVariants R package[148]),
after excluding possible germline variants from Cas9-non-induced samples.

Proliferation and neurogenesis analysis: For proliferation analysis using Ki-67, NPCs
were seeded into 24-well plates and either treated with doxycycline (induced) to activate
Cas9 or left untreated (uninduced). The cells were cultured for 7 days, representing
approximately three NPC generations. On day 7, cells were collected, and ~1 × 10^6
cells were stained with Ki-67-FITC (#130-117-803, Miltenyi Biotec) using the
Foxp3/Transcription Factor Staining Buffer Set (#00-5523, Invitrogen), following the
manufacturer's protocol.

To evaluate the effects of gene KOs on neurogenesis and gliogenesis, transduced
NPC-iCas9 lines were spontaneously differentiated into human cortical neurons and
glial cells. Briefly, 1 × 10^6 cells were seeded in GelTrex-coated (1:5) 6-well plates and
cultured in complete neuronal media containing BrainPhys™ Neuronal Medium,
Glutamax (100X), Sodium Pyruvate (100 mM), B-27 (-RA) supplement (50X), N2
(100X), Anti-Anti (100X), Natural Mouse Laminin (1 mg/ml), dbcAMP (500 mg/ml), L-
Ascorbic Acid (200 µM), BDNF (20 µg/ml), and GDNF (20 µg/ml). Media was refreshed
every three days. On day 25, cells were collected and stained for FACS analysis using
surface markers previously described[149] to differentiate NPCs (CD184+/CD44-/CD24+),
neurons (CD184-/CD44-/CD24+), and glia (CD184+/CD44+). CD271, a marker for
mesenchymal stem cells, was excluded from the original panel as NPCs were pre-
purified via FACS using CD133+/CD184+/CD271- markers before differentiation. A
minimum of 50,000 cells per gate were acquired using a BD LSRFortessa™ Cell
Analyzer at the Yale Flow Cytometry Core. Flow cytometry data were analyzed using
FlowJo™ v10.10 Software (BD Life Sciences).

All statistical analyses for flow cytometry assessment were conducted using GraphPad
Prism version 9.5.1 (528) for macOS (GraphPad Software, San Diego, CA). Each well
was treated as an independent replicate. Differences between knockout (induced) and
control (uninduced) groups were assessed by comparing the mean fluorescence
intensity (MFI) of the target fluorophore using an unpaired t-test with Welch correction to
account for individual group variance. Multiple comparisons were corrected using the
False Discovery Rate (FDR) method with a two-stage step-up procedure (Benjamini,
Krieger, and Yekutieli) at an FDR threshold of 5%.

 FACS Analysis of Mitochondrial Membrane Potential and CRISPR Screen Read-out via
Amplicon Sequencing

For our mitochondrial assays we used a nearly identical library (same backbone, guide
density, and control set) screened exclusively in the H1 inducible Cas9 (H1-iCas9)
hPSC line. Mitochondrial inner membrane potential (Δψm) was measured in H1-iCas9,
following differentiation to NPCs or iGlut on day 21. Cells were harvested, counted, and
aliquoted at 1 × 10^6 cells per sample. JC-1 dye (MitoProbe™ JC-1 Assay Kit;
Invitrogen #M34152) was dissolved in DMSO at a stock concentration of 200 µM and
added to each sample to achieve a final concentration of 2 µM, then incubated for 30
min at 37 °C in 5% CO_2. A 50 µM CCCP control was included to induce complete
mitochondrial depolarization. After staining, cells were washed once in their respective
culture medium, resuspended in FACS buffer (Invitrogen eBioscience Staining Buffer
#00422226), and analyzed immediately on a Thermo Fisher "Bigfoot" spectral cell sorter
using 488 nm excitation with 525/50 nm (FITC) and 585/40 nm (PE) emission filters.
Debris and doublets were excluded by forward/side scatter gating, and CCCP-treated
samples were used to define FITC and PE gates. Approximately 1 × 10^6 events per
sample were recorded. Cells were then pelleted (300 × g, 5 min) and genomic DNA
extracted using the Qiagen DNeasy Blood & Tissue Kit (#69504).

UMI-tagged amplicon libraries were generated in three PCR steps. In PCR-1, genomic
DNA was amplified with Platinum™ II Hot-Start PCR Master Mix (Invitrogen,
#14000012) and UMI-containing primers (Forward: 5′-
ACACTCTTTCCCTACACGACGCTCTTCCGATCTACGTGACGTAGAAAGTAATAATTT
CTTGGGT-3′; Reverse: 5′-
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTN(25)NNNNNNNNNNACTCGGTGC
CACTTTTTCAA-3′) under the following conditions: 94 °C for 2 min; 4–6 cycles of 98 °C
for 5 s, 60 °C for 15 s, 60 °C for 30 s. The resulting ~180 bp products were purified and
concentrated using the Zymo DNA Clean & Concentrator-5 kit (#D4013) and eluted in
10 µL nuclease-free water (Thermo Fisher #AM9938). In PCR-2, purified product was
amplified with adaptor primers (Forward: 5′-
ACACTCTTTCCCTACACGACGCTCTTCCGATCT-3′; Reverse: 5′-
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT-3′) for 22 cycles under identical
cycling conditions in a ~20 µL reaction. A seven-cycle indexing PCR (PCR-3) was
performed by the sequencing facility Yale Center for Genome Analysis (YCGA) prior to
sequencing. Final libraries were sequenced on an Illumina NovaSeq platform (paired-
end 150 bp, 5 million reads per sample).

Flanking sequences on both sides of each gRNA were trimmed using BBDuk, and
reads were then mapped to gRNA reference sequences and counted using
MAGeCK[150]. Raw counts for each gRNA were normalized to counts of scrambled
gRNA. Abundance of each target gene was then calculated by summing of all gRNAs
targeting that gene. Log2-transformed fold changes of gRNA-targets abundance were
compared between PE-high samples and FITC-high samples.

Immunostaining. Cells were fixed with fixative solution (4 % sucrose and 4 %
paraformaldehyde prepared in Dulbecco's Phosphate Buffered Saline (DPBS)) for 10
min at room temperature (RT). Following this, cells were washed twice with DPBS and
incubated in blocking solution (2% normal donkey serum prepared in DPBS)

supplemented with 0.1% Triton for two hours at RT. After this, cells were incubated overnight at 4 °C in the primary antibody solution prepared in blocking solution. Cells were washed three times with DPBS, incubated at RT in secondary antibody prepared in blocking solution, then washed three times with DPBS. In the second wash, cells were incubated in DBPS supplemented DAPI (Sigma D9542,1 µg/mL) for 2 min at RT.

Antibody	Species	Vendor	Catalog #	Dilution
anti-MAP2	chicken	Invitrogen, Abcam	PA1-10005, ab5392	1:1000
anti-Nestin	rabbit	Millipore	ABD69	1:200
anti-vGLUT1	rabbit	Synaptic systems	135-303	1:200
anti-GABA	rabbit	Sigma-Aldrich	A2052	1:200
TOMM20	mouse	Santa Cruz Biotechnology	sc-17764	1:200
Total OXPHOS	n/a	Abcam	AB-317270	1:500
anti-mouse	donkey	Jackson ImmunoResearch	715-605-151	1:500
anti-rabbit	donkey	Jackson ImmunoResearch	711-545-152	1:500
anti-chicken	donkey	Jackson ImmunoResearch	715-605-150, 703-545-155	1:500

Fixed cultures were acquired using a DragonFly Confocal Dual Spinning Disk confocal, at 60x magnification and 1.4 numerical aperture. All images were acquired with a fixed laser intensity and exposure time across experimental conditions. Four images were acquired per well, and 4-10 wells were acquired per experimental condition. Each well represents a biological replicate and statistical datapoint. Therefore, each replicate represents hundreds of μm^2 of neuronal area and tens of thousands of individual mitochondria.

Mitochondria morphology features were determined using the Surface module of Imaris 10.2. Likewise, OXPHOS complex features were determined using the surface module of Imaris 10.2. The Volume, Area and Sphericity features of the Surface modules were selected for analysis. Mitochondria networking features were determined using published, open-source methods[179]. A one-way ANOVA with a Šidák's multiple comparisons test was performed on data on GraphPad Prism 10.

To validate robustness and sensitivity of the microscopy assay, we treated D14 iGluts overnight with carbonyl cyanide 4-(trifluoromethoxy) phenylhydrazone/FCCP (Sigma-Aldrich, SML2959) at 5 µM, 10 µM and 50 µM doses. Following this, we conducted the immunostaining, mitochondrial structural analysis and statistical analyses outlined above.

Seahorse XF Mito Stress Test: Day 5 iGLUTs were plated at 1.65 × 10 cells/well in XF24 microplates (Agilent, 100777-004) and cultured to day 21. One hour prior to measurement, growth medium was removed, leaving 50 µL per well, and replaced with 1 mL of pre warmed Seahorse XF DMEM (Agilent, 103575-100) supplemented with 25 mM glucose (Agilent, 103577-100) and 0.23 mM pyruvate (Agilent, 103578-100). Plates were equilibrated for 1 h at 37 °C in a non CO$_2$ incubator. Immediately before the assay, the medium was replaced with 500 µL of fresh assay buffer. Oxygen consumption rate (OCR) was recorded on a Seahorse XFe24 Analyzer (Agilent) using the standard Mito Stress Test. The program consisted of three sequential injections—1.5 µM oligomycin (Sigma-Aldrich, 75351), 1.5 µM carbonyl cyanide 4-(trifluoromethoxy) phenylhydrazone/FCCP (Sigma-Aldrich, C2920), and a mix of 0.5 µM rotenone (Sigma-Aldrich, R8875) + 0.5 µM antimycin A (Sigma-Aldrich, A8674)—separated by four measurement phases (baseline plus post injection 1–3). Each phase comprised three cycles of 3 min mixing, 2 min waiting, and 3 min measurement. After the assay, cells were lysed using M-PER™ Mammalian Protein Extraction Reagent (ThermoFisher, 78501) supplemented with cOmplete™ Mini Protease Inhibitor Cocktail (Sigma-Aldrich, 11836153001) and PhosSTOP™ (Sigma-Aldrich, 4906845001), according to the manufacturer's instructions. Total protein concentrations were determined using the Pierce™ Dilution-Free™ Rapid Gold BCA Protein Assay (ThermoFisher, A55860), and OCR values were normalized to total protein content.

CRISPR organoid assays: H1-hESC-iCas9 cells were transduced with a pooled gRNA library containing four gRNAs per target gene, with 20% of the library comprising non-targeting gRNAs. Following selection with 1 µg/mL puromycin, the established cell line was used to generate cortical organoids following a well-established protocol[151] with slight modifications. In brief, embryoid bodies (EBs) were generated using AggreWell plates (Stemcell Technologies) according to the manufacturer's instructions. Once formed, EBs were transferred to ultralow-attachment 10 cm plates (Corning) for further culture. Patterning was initiated using StemFlex base media (A3349401, Gibco) supplemented with 100 nM LDN193189 (x) and 10 µM SB431542 (x). The media was refreshed daily. Organoids were cultured on an orbital shaker at 53 rpm for the duration of the protocol. On Day 6, the patterning media was replaced with growth media; Neurobasal A medium (10888022, Gibco), 1× GlutaMAX (35050061, Gibco), and 1× B27 (12587010, Gibco), supplemented with 20 ng/mL FGF (PeproTech) and 20 ng/mL EGF (PeproTech). On Day 14, Cas9 expression was induced by treating the organoids with 2 µg/mL doxycycline (Sigma-Aldrich) for 72 hours. From Day 25, FGF and EGF were replaced with 20 ng/mL BDNF (PeproTech) and 20 ng/mL NT-3 (PeproTech). Media changes were performed every other day. Starting from Day 42, organoids were maintained in growth media without additional supplements. Media was refreshed 2–3 times per week.

The organoids were maintained in culture for ~80 days, at which point five organoids from three biological replicates were collected for DNA extraction using the DNeasy Blood & Tissue Kit (#69504, Qiagen). Extracted DNA was subjected to PCR amplicon sequencing with unique molecular identifiers (UMIs) using a three-step PCR protocol. In the first step (PCR-1), UMI-containing primers (5'-

ACACTCTTTCCCTACACGACGCTCTTCCGATCTACGTGACGTAGAAAGTAATAATTT
CTTGGGT-3') and (5'-
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTN(25252525)NNNNNNNNNACTC
GGTGCCACTTTTTCAA-3') were used for 4 cycles. PCR-2 utilized adaptor primers (5'-
ACACTCTTTCCCTACACGACGCTCTTCCGATCT-3') and (5'-
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT-3') for 22 cycles. PCR-3,
performed by the sequencing facility, added sample-specific indexing in 7 additional
cycles. The prepared libraries were sequenced on a NovaSeq platform with paired-end
150 bp reads, generating 10 million reads per sample at the Yale Center for Genomic
Analysis (YCGA).

Fragments amplified by PCR were sequenced on NovaSeq 6000 sequencer pair end at
150bp with ~10 million reads per sample. Flanking sequence on both side of gRNAs
were trimmed using BBDuk, and reads were then mapped to gRNA reference
sequences and counted using MAGeCK package[150]. Raw counts for each gRNA were
normalized to counts of scrambled gRNA. Abundance of each gRNA-target genes were
then calculated by sum of all gRNAs targeting that gene after excluding gRNAs with low
KO-efficiency (<5%). Average Log2-transformed fold change of gRNA-targets
abundance were compared between doxycycline-induced versus uninduced samples on
day 77 samples.

Analysis of single-cell CRISPRko screens in NPCs, DIV 7, DIV 21 iGLUTs and DIV 36
iGABAs. mRNA sequencing reads were mapped to the GRCh38 reference genome
using the *Cellranger* Software. To generate count matrices for GDO (gRNA) libraries,
the kallisto indexing and tag extraction (kite) workflow were used. Count matrices were
used as input into the R/*Seurat* package[152] to perform downstream analyses, including
QC, normalization, cell clustering, GDO demultiplexing, and covariate regression[71,153].

Normalization and downstream analysis of RNA data were performed using the Seurat
R package (v.5.1.0), which enables the integrated processing of multimodal single-cell
datasets. CRISPR-screen experiments in each cell-type were processed independently.
Within each cell-type, ~100-80,000 cells were sequenced across 4 lanes. gRNA and
RNA UMI feature counts were filtered removing the top and bottom decile of cells based
on distribution of counts in each cell-type. The percentage of all the counts belonging to
the mitochondrial, ribosomal, and hemoglobin genes calculated using
Seurat::PercentageFeatureSet were filtered with cell-type specific thresholds, given the
relatively high proportion of mitochondrial genes expressed in neurons. Mitochondrial,
ribosomal, and hemoglobin genes as well as MALAT1 were removed
(^RP[SL][[:digit:]]|^RPLP[[:digit:]]|^RPSA|^HB[AEGQ][[:digit:]]|^HB[ABDMQ]|^MT-
|^MALAT1$). Lowly expressed genes, those that had at fewer than 2 read counts in
90% of samples were also removed. Hashtag and guide-tag raw counts were
normalized using centered log ratio transformation, where counts were divided by the
geometric mean of the corresponding tag across cells and log-transformed. gRNA
demultiplexing was performed using the Seurat::MULTIseqDemux function for each
lane individually and then counts were merged across lanes (**SI Fig. 3B**). In NPCs,
94,363 cells were retained after filtering and removal of negatively assigned cells with
62,7% classified as doublets and 37.3% classified as singlets. In DIV7 and DIV21
iGLUTs, 57,685 and 31,473 cell were retained with 34% and 9.8% doublets and 66%

and 90.2% singlets respectively. In DIV35 iGABAs, 64,462 cells were retained with
48.3% doublets and 51.7% singlets. For all downstream analysis only cells with "singlet"
gRNA classification were used (26,549-38,097 cells per experiment) (**SI Fig. 4C-E**).
Number of singlet cells by gRNA per cell-type shown in **SI Fig. 6AB**.

Cell-type specific population heterogeneity correction. Gene-expression based
clustering was largely driven by cellular heterogeneity, cell quality, and sequencing lane
effects. gRNA identity was not correlated with these covariates (**SI. Fig. 7**), so we
adjusted for transcriptomic variability arising from cellular heterogeneity by applying
maturity and cellular subtype scores across both perturbed and non-perturbed cells.
First, variation related to cell-cycle phase of individual cells was accounted for by
assigning cell cycle scores using Seurat::CellCycleScoring which uses a list of cell cycle
markers[154] to segregate by markers of G2/M phase and markers of S phase. Second, to
address variance due to cellular heterogeneity within a single experiment, we adapted
the method applied by Seurat::CellCycleScoring to calculate a "Maturity. Score" and
"Subtype.Score" for each cell based on cellular subtype (more variable in mature
GABAergic neurons) and developmental time-point specific markers (mora variable in
NPCs and immature iGLUTs) (**SI Table 2-3**). Cells with outlier maturity scores and
subtype scores were removed from downstream analyses. RNA UMI count data were
then normalized, log-transformed and the percent mitochondrial, hemoglobulin, and
ribosomal genes (markers of cell quality), lane, cell cycle scores (Phase), and maturity
scores regressed out using Seurat::SCTransform. The scaled residuals of this model
represent a 'corrected' expression matrix, that was used for all downstream analyses.

Although demultiplexing assigned the correct guide identity to each cell, to remove
"false positives" whereby gRNAs were assigned but gene expression was unperturbed,
the transcriptomes of gRNA clusters were evaluated relative to scramble gRNAs,
ensuring that cells assigned to a guide-tag identity class demonstrated successful
perturbation of the targeted NDD gene. To remove subsequent "false negatives",
whereby a successful CRISPR-KO may not result in significant down-regulation of the
targeted gene[71] yet still achieve an overall transcriptomic profile distinct from scramble
populations, we performed 'weighted-nearest neighbor' (WNN) analysis to assign
clusters based on both guide-tag identity class and gene expression[72]. To identify
successfully perturbed cells, the transcriptomes of gRNA clusters were compared to
Scramble-gRNA control clusters by differential gene expression analysis (Wilcoxon
Rank Sum) comparing each cluster to all other clusters. Non-targeting WNN clusters
and KO gRNA WNN clusters were filtered by setting a quantile base average
expression threshold of target genes based on the distribution of target gene average
expression across all other clusters. Clusters were the collapsed by gRNA identity;
gRNAs with less than 75 cells were removed from analysis. These cells were then used
for downstream differential gene-expression analyses[155]. For each cell-type individually,
single-cell gene expression matrices were PseudoBulked using
scuttle::aggregateAcrossCells function across lanes (4 pseudo-bulk samples per
perturbation), lowly-expressed genes were removed (leaving 18-22,000 genes) followed
by edgeR/limma differential gene expression analysis. Concordance between Wilcox-
rank sum differential gene expression analysis using single-cell data and limma:voom
using PseudoBulked data was assessed for each gene.

Altogether, Wilcoxon Rank Sum was applied to measure NDD gene knockdown from
single-cell DEG analysis. Given the concordance between the DEG results using single-
cell Wilcox and pseudo-bulk limma:voom (**SI Fig. 6C**), all main figure and all SI figures
thereafter applied pseudobulked data analyzed with limma.

To validate whether the high correlation within cell type was due to exactly the same
scramble control cells, we re-performed DEGs using random selection of subset of
scramble cells for each cell type (**SI Fig. 8**). Briefly, for each gene, 50% (if number of
pseudobulked sample cells > 50) or 80% (if number of pseudobulked sample cells < 50)
of scramble cells were randomly selected using sample function from R. DEGs were
then performed as described above using limma/dreamlet package between KOs and
subset of scrambles different among genes. The process was repeated three times to
avoid random selection bias and median of each gene logFC was used as the final
logFC. Average overlap of random scramble cells across different genes is
approximately 50%.

Meta-analysis of gene expression across perturbations[73]. Across NDD KOs, DEGs were
meta-analyzed (METAL[156]), and "convergent" genes were defined as those with
significant and shared direction of effect across all NDD gene perturbations and with
non-significant heterogeneity (FDR adjusted $p_{meta} < 0.05$, Cochran's heterogeneity Q-test
$p_{Het} > 0.05$). To test convergence between NDD-KOs, meta-analyses were performed
across all possible combinations of 2-5 KO perturbations with and without sub-setting
for those shared across cell types (>40,000 combinations across cell-types) (**SI Data 1**).

Bayesian Bi-clustering to identify Convergent Networks[73]. Across NDD KOs, convergent
networks were generated by Bayesian bi-clustering[157] and undirected gene co-
expression network reconstruction from the NDD KOs. Not constrained by statistical
cut-offs, and able to capture the effect of more lowly expressed genes, convergent
networks may be a more sensitive measure of convergence. Networks were built based
on bi-clustering (BicMix)[158] using log2CPM expression data from all the replicates
across each of the NDD gene sets and Scramble gRNA jointly. We performed 40 runs
of BicMix on these data and the output from iteration 400 of the variational Expectation-
Maximization algorithm was used. Target Specific Network reconstruction[159] was
performed to identify convergent networks across all possible combinations of the 9
NDD gene KO perturbations shared across cell-types (n=502 combinations/cell-type)
and randomly sampled combinations of 2-21 KO perturbations without sub-setting for
those shared across cell types (n=1400-2300 combinations).

Influence of Functional Similarity on Convergence Degree. To test the influence of
functional similarity and brain co-expression between KOs on convergence and
compare the degree of convergence between the same KOs in different cell-types we
established two methods for defining and measuring convergence. First, gene-level
convergence using meta-analysis as described above, with the strength of convergence
for each set defined as ratio of convergent genes to the average number of DEGs.

$$gene\ level\ convergence = \frac{nConvergent\ Genes}{mean(\sum_1^N\ nDEGs)}$$

Second, network-level convergence based on undirected network reconstruction from
Bayesian bi-clustering as described above. Bi-clustering identifies co-expressed genes

shared across the downstream transcriptomic impacts of any given set of KO
perturbations, thus, the resolved networks are the transcriptomic similarities between
distinct perturbations (convergence)**.** We calculated the "degree of convergence" for
each network based on previously described metric[73]. Briefly, convergence scores are
based on (1) network connectivity as defined by the sum of the clustering coefficient
(Cp) and the difference in average length path (Lp) from the maximum average length
path resolved across all possible sets [(max)Lp-Lp] and (2) similarity of network genes
based on biological pathway membership scored by taking the sum of the mean
semantic similarity scores [160] between all genes in the network and (3) minimum
percent duplication rate across 40 runs. Duplication thresholds are network-dependent
and a metric of confidence in the connections.

$$network\ level\ convergence$$
$$= Cp + [(Lp) - Lp] + mean\left(\sum_{1}^{N} MFsemsim + BPsemsim + CCsemsim\right)$$
$$+ nDuplication\ runs/nTotal\ Runs$$

Functionally similarity scores across the NDD KO genes represented in each set was
calculated using (1) Gene Ontology Semantic Similarity Scores: the average semantic
similarity score based on Gene Ontology pathway membership within Biological
Pathway (BP), Cellular Component (CC), and Molecular Function (MF) between NDD
genes in a set[160] and (2) brain expression correlation (BEC) score: based on the
strength of the correlation in NDD gene expression in the CMC (n=991 after QC) post-
mortem dorsa-lateral pre-frontal cortex (DLPFC) gene expression data,.

We performed Pearson's correlation analysis (Holm's adjusted P) on similarity scores
and the degree of network convergence to determine the influence of the similarity of
the initial KO genes on downstream convergence. We compared the average strength
of convergence across cell-types using a parametric Welch's F-test and pairwise
Games-Howell test.

Enrichment analysis of convergence for risk loci using MAGMA. We intersected cross
cell-type perturbation specific and cross perturbation cell-type-specific gene-level
convergence with genetic risk of psychiatric and neurological disorders/traits [attention-
deficit/hyperactivity disorder (ADHD)[161], anorexia nervosa (AN)[162], autism spectrum
disorder (ASD)[2], alcohol dependence (AUD)[163], bipolar disorder (BIP)[164], cannabis use
disorder (CUD)[165], major depressive disorder (MDD)[166], obsessive-compulsive disorder
(OCD)[167], post-traumatic stress disorder (PTSD)[168], and schizophrenia (SCZ)[169], Cross
Disorder (CxD)[170], Alzheimer disease (AD)[171], Parkinson disease (PD)[172], amyotrophic
lateral sclerosis (ALS)[173], Tourette's[174], migraine[175], chronic pain[176], and neurotic
personality traits[177] GWAS summary statistics] using multi-marker analysis of genomic
annotation (MAGMA)[65]. SNPs were mapped to genes based on the corresponding build
files for each GWAS summary dataset using the default method, snp-wise□=□mean (a
test of the mean SNP association). A competitive gene set analysis was then used to
test enrichment in genetic risk for a disorder across gene sets with an FDR<0.05.

To test if observed effects were due to the differential size of the gene sets for each
GWAS or owing to the fact that DEGs are more likely to include neural genes, which are

more likely to be associated with brain disorder, GWAS sets were filtered for genes
expressed in each cell-type prior to enrichment testing and enrichment tests were
performed after randomly down-sampling GWAS Gene Sets to 100, 250, 500, 750, and
1000 genes (**SI Fig. 9)**, performed ten times within each set size (i.e., 50 tests for each
GWAS)."

Over-representation analysis, functional enrichment annotation, and biological theme
comparison of convergence. To identify pathway enrichments unique to individual KOs,
convergent genes, and convergent networks based on zebrafish behavioral subgroups
(see zebrafish methods below), we performed biological theme comparison and GSEA
using ClusterProfiler[178]. Using FUMAGWAS: GENE2FUNC, the 102 ASD genes were
functionally annotated and overrepresentation gene-set analysis for each convergent
gene set was performed[179]. Using WebGestalt (WEB-based Gene SeT AnaLysis
Toolkit)[180], over-representation analysis (ORA) was performed on all convergent gene
sets against publicly available genset lists GeneOntology, KEGG, DisGenNet, Human
Phenotype Ontology, and a curated gene list of rare-variant targets associated with
ASD,SCZ, and ID[67].

Random forest prediction model of convergence strength. To determine how well
functional similarity between KOs can predict gene-level and network-level convergence
we trained a random forest model[75] (randomForest package in R) for each type of
convergence, evaluated the model in an independent internal dataset, and validated the
model in an external CRISPRa activation screen[73]. Data from randomly tested gene
combinations (2-5 KO sets at the gene level and 2-10 KO sets at the network level)
tested across cell-types were randomly down-sampled into a training set (70%) and
testing set (30%) – all with comparable proportions of data by cell-type. The random
forest model was trained with bootstrap aggregation using C.C, M.F, B.P semantic
similarity scores, brain expression correlation, number of genes, and cell-type as
predictors. The Random Forest linear regression model was evaluated in the testing
data by comparing actual values to predicted values, estimating the root mean squared
error and performing Pearson's correlations. Predictor models were validated using an
external dataset of 10 CRISPR-activation perturbations of SCZ common variant target
genes with multifunctional annotations broadly grouped as signaling/cell communication
(*CALN1, NAGA, FES, CLCN3, PLCL1*) and epigenetic/regulatory (*SF3B1, TMEM219,*
UBE2Q2L, ZNF804A, ZNF823)[73], and assessed based the root mean squared error and
Pearson's correlation between actual and predicted convergence strength.

LNCTP in silico model

To investigate the perturbation of ASD genes in silico, we adapt the Linear Network of
Cell-Type Phenotypes (LNCTP) model[76] to predict the effects of changes in gene
expression in the prefrontal cortex, across neuronal and non-neuronal cell-types. The
LNCTP is defined as an energy model representing the joint distribution of a collection
of phenotypes of interest conditioned on the genotype. Since we are interested
primarily in the effects of gene expression perturbations on the expression of other
genes, we use only the imputation segment of the LNCTP model (excluding the
prediction of higher-order phenotypes and cell-cell interactions).

The probabilistic model for the imputation-based LNCTP may be expressed as:

$$p_{LNCTP}(x_i|z_i) = \exp(-E(x_i|z_i))$$

1070

$$E(x_i|z_i) = x_{i0.}^T J x_{i0.} + \sum_g x_{i0g}^T b(z_i, \beta_g) + \sum_c (x_{ic.}^T J_c x_{ic.} + x_{ic.}^T b_c)$$
$$+ \lambda \sum_g (x_{i0g} - f(z_i)^T x_{i,1...C,g})^2 .$$

(1)

Here, z_i represents the genotype of individual i, and x_i represents bulk and cell-type
specific gene expression from individual i. We further index the gene expression by C
cell-types (which are here: Excitatory Neurons, Inhibitory Neurons, Oligodendrocytes,
Astrocytes, Oligodendrocyte Precursor Cells, Endothelial Cells and Microglia), which will
be denoted $x_1, x_2, ... x_C$, and we will use x_0 to denote the bulk expression. The variables
$f_{1...C}$ represent the estimated cell-fractions in the bulk observations (predicted from the
genotype, z). The parameters of the model are $\theta = \{\beta_{1...G}, J_{0...C}\}$ and λ acts as a
hyperparameter. The parameters $\beta_{1...G}$ and $J_{0...C}$ reflect the gene specific expression
biases and pairwise interactions respectively, whose non-zero elements are determined
by the sparsity structure arising from eQTLs and Gene Regulatory Network (GRN)
linkages respectively; the non-zero elements of J_c occur only between genes connected
in the GRN of cell-type c.

Further details on the training of the model in Eq. (1) can be found in[76]; here, we outline
the specific differences in the training for the purposes of our analysis. As in[76], we use
genetics and expression data from post-mortem PFC samples from the PsychENCODE
consortium. However, we group together samples from all higher-order phenotypes
during training (control (CTR), schizophrenia (SCZ), bipolar disorder (BPD) and autism
spectrum disorder (ASD)), and split the data into three partitions of size 760, 100 and
100 for training, validation and testing respectively (each including samples from all
higher-order phenotypes). Further, we include all 29 CRISPR targeted genes, 102 NDD
genes[61], Transcription Factors[76] and neuropsychiatric TWAS-selected genes[76], and the
top 100 up and down regulated CRISPR convergent genes in iGLUT and iGABA cells
(400 genes in total), in the model, generating 1325 genes in total. The eQTL and GRN
linkages from PsychENCODE are then restricted to this subset of genes.

LNCTP Simulating Perturbations

To perform perturbations in this model corresponding to the 29 CRISPR targeted genes,
we use the following perturbation-conditioned version of the LNCTP model:

$$p_{LNCTP}(x_{i,\neg(c^*,g^*)}|z_i, x_{i,c^*,g^*} = \{k, -k\}) = \exp(x_{i,\neg(c^*,g^*)}|z_i, x_{i,c^*,g^*} = \{k, -k\})$$

1099

$$E\left(x_{i,\neg(c^*,g^*)}\middle|z_i, x_{i,c^*,g^*} = \{k, -k\}\right)$$
$$= x_{i0\cdot}^T J x_{i0\cdot} + \sum_g x_{i0g}^T b(z_i, \beta_g) + \sum_c (x_{ic\cdot}^T J_c x_{ic\cdot} + x_{ic\cdot}^T b_c)$$
$$+ \lambda \sum_g \left(x_{i0g} - f(z_i)^T x_{i,1\ldots C,g}\right)^2 + K\delta\left(x_{i,c^*,g^*} = \{k, -k\}\right).$$

(2)

where (c^*, g^*) denotes the perturbed gene and cell type, whose expression is set to k or
$-k$, $\delta(a)$ is a delta function whose value is 0 if expression a is true, and 1 otherwise,
and K is an arbitrarily large value. We perturb each of the CRISPR targeted genes in
turn in the bulk network, using $k = 2$, and applying a negative perturbation to mimic the
effect of the CRISPR perturbation. We note that, since the model is trained on Z-scored
log-normalized expression counts, this corresponds to introducing a large negative fold-
change to the selected gene. The *in silico* predicted log fold-changes per individual
across all genes (per cell-type) are then calculated by comparing the expected values
before and after perturbation:

$$\Delta_{i,c,g} = \mathbb{E}_{p_{LNCTP}(\cdot|z_i)}\left[x_{i,c,g}\right] - \mathbb{E}_{p_{LNCTP}\left(\cdot\middle|z_i, x_{i,c^*,g^*}=\{k,-k\}\right)}\left[x_{i,c,g}\right]$$

(3)

and the final predicted log fold-changes are calculated by taking the expectation across
individuals. We use the sampling approach in[76] to evaluate the expectations in Eq. (3).

To perform perturbations across all 102 NDD genes, for efficiency we learn a reduced
model by remove the dependency on z_i in Eq. (1). We sample cell-type specific
expression values for each individual from the full model, and then fit the reduced model
by refitting the model parameters to maximize the likelihood of the full data vectors
(consisting of the original bulk and sampled cell-specific expression vectors for each
individual). Perturbations are performed in the reduced model as in Eq. (2) and fold-
changes are calculated as in Eq. (3), while removing the dependency on z_i and the i
subscripts respectively.

LNCTP in silico convergent genes

To identify *in silico* convergent genes for a set of perturbations, $S = $
$\{(c_1^*, g_1^*), \ldots, (c_N^*, g_N^*)\}$, we calculate $\Delta_{c,g}$ using Eq. (3) for each perturbation, writing $\Delta_{c,g}^{c^*,g^*}$
for the log fold-change to (c, g) generated by applying perturbation (c^*, g^*), and $\Delta_{c,g}^S$ for
the set of log fold-changes by applying all perturbations in S. Then, the set of *in silico*
convergent genes for S is found by selecting those for which $p_{\text{sign}}\left(\Delta_{c,g}^S \cdot \left[\Delta_{c,g}^S \geq \tau\right]\right) <$
0.1, where $p_{\text{sign}}(.)$ is the p-value from a 2-tailed one-sample sign-test. The threshold τ
is introduced to reduce noise from perturbations which are estimated to generate small
log fold-changes, and throughout we set $\tau = 0.3$.

For the comparison of *in silico* convergent genes derived from different perturbation sets
S, we apply two-sided hypergeometric tests to the gene sets defined as above (using all
1325 genes in our model as the background set). For Gene Set Enrichment Analysis of

convergent genes derived from S, we apply clusterprofiler[178] to the full set of genes in
our model, ranked by $p_{\text{sign}}\left(\Delta_{c,g}^{S} \cdot \left[\Delta_{c,g}^{S} \geq \tau\right]\right)$ as defined above.

LNCTP semantic distance test

To test the semantic distance between enriched terms for two sets of perturbations S_1
and S_2, we generate the set of enriched terms T_1 and T_2 by applying GSEA to each set
as described above (using Benjamini Höchberg correction and an FDR threshold of 0.2
to select enriched terms T_1 and T_2). We then calculate the similarity between terms t_1
and t_2 by evaluating $s(t_1, t_2) = |G(t_1) \cap G(t_2)|/|G(t_1) \cup G(t_2)|$, where $G(t)$ denotes the
set of genes occurring in the leading edge of term t. We test for a significant semantic
distance between S_1 and S_2 by evaluating $s(t_1, t_2)$ between all pairs $t_1 \in S_1$, $t_2 \in S_2$,
versus all pairs $t_1 \in S_1$, $t_2 \in S_1$ and $t_1 \in S_2$, $t_2 \in S_2$, and applying a one-sided rank-sum
test for the for a smaller similarity in the former pairs versus the latter.

Transcriptional correlations between hiPSC-derived neural cells, fetal and adult brain
cell types, and the zebrafish brain.

We compared wild-type (WT) zebrafish brain expression to gene expression in our
hiPSC-derived models and to sign-cell expression data for the fetal and adult PFC
(PsychENCODE[181,182]:
http://resource.psychencode.org/Datasets/Derived/SC_Decomp/DER-
20_Single_cell_expression_processed_TPM.tsv). We first filtered zebrafish gene names
and converted them to the appropriate *Homo sapiens* orthologs using the R package
orthogene (v3.2.1[183]); genes without matched orthologs were dropped from both
species. Pseudo-bulk expression data from scramble control cells were used as the
baseline expression across NPCs, D7 iGLUTs, D21 iGLUTs, and D36 iGABAs.
Pearson's correlation coefficients between *in vitro* cells, fetal and adult postmortem
brain cells, and zebrafish brain were calculated and a Bonferroni correction applied.

Zebrafish

All procedures involving zebrafish were conducted in accordance with Institutional
Animal Care and Use Committee (IACUC; Protocol #2024-20054) regulatory standards
at Yale University. Zebrafish larvae were raised at 28°C on a 14:10 hour light:dark
cycle. Larvae were grown in 150 mm Petri dishes in blue water (0.3g/L Instant Ocean, 1
mg/L methylene blue, pH 7.0) at a density of 60-80 larvae per dish. Behavioral assays
were conducted in zebrafish larvae at 5-7 dpf. At these developmental stages, sex is not
yet determined.

Zebrafish mutant generation

We performed automated, high-throughput, quantitative behavioral profiling of larval
zebrafish to measure arousal and sensorimotor processing as a readout of circuit-level
deficits resulting from gene perturbation.[60] We quantified 24 parameters across sleep-
wake activity and visual-startle responses in 18 stable homozygous mutant or F0
mosaic crispant lines for 15 NDD genes (**SI Tables 4-5**). Stable zebrafish lines were
generated by our lab (*arid1b$^{\Delta7/\Delta7}$, chd2$^{\Delta7/\Delta7}$, chd8$^{\Delta7/\Delta7}$, chd8$^{\Delta5/\Delta5}$, kdm5ba$^{\Delta17/\Delta17}$b$^{\Delta14/\Delta14}$*,
kdm5ba$^{4i/4i}$b$^{\Delta4/\Delta4}$)[60] or provided as a generous gift from the Thyme lab
(*ash1l$^{1i,\Delta60,19i/1i,\Delta60,19i}$*, *kmt5b$^{\Delta208,1i,\Delta5/\Delta208,1i,\Delta5}$*, *kmt2ca$^{\Delta82,17i/\Delta82,17i}b^{\Delta6,\Delta29/\Delta6,\Delta29}$*,

nrxn1a$^{\Delta218/\Delta218}$)[184,185]. F0 crispants for the following genes were generated according to
ref. [186]: *chd2, kdm6bab, mbd5, phf12ab, phf21aab, skiab, smarcc2, wacab*. Briefly, we
designed two CRISPR crRNAs per allele, prioritizing early exons for targeting. CRISPR
RNPs were assembled individually and then combined prior to injection at the one-cell
stage. The number of scrambled guides injected into the control group was matched to
the number of CRISPR guides used for the experimental group. Injected embryos were
raised to 5 dpf at which point the behavioral assays (described below) were conducted.
We identified unique behavioral fingerprints for each NDD gene mutant, revealing
convergent and divergent phenotypes across mutants (**SI Fig. 22B**). To classify
convergent behavioral subgroups that may share circuit-level functions, we performed
correlation analyses with hierarchical clustering across mutants. We identified four
distinct subgroups of NDD genes with highly correlated behavioral features (**Fig. 7A**).

Behavioral assays

Larvae were placed into individual wells of a 96 well plate (7701-1651; Whatman,
Clifton, NJ) containing 650 µL of standard embryo water (0.3 g/L Instant Ocean, 1 mg/L
methylene blue, pH 7.0) per well within a Zebrabox (Viewpoint LifeSciences; Viewpoint
Life Sciences, Montreal, Quebec, Canada). Locomotion was quantified with automated
video-tracking system (Zebrabox and ZebraLab software). The visual-startle assay was
conducted at 5 days post fertilization (dpf) as described[60]. To assess larval responses to
lights-off stimuli (VSR-OFF), larvae were acclimated to white light for 1 hour, and
baseline activity was tracked for 30 minutes followed by five 1-second dark flashes with
intermittent white light for 29 seconds. To evaluate larval responsivity to lights-on stimuli
(VSR-ON), the assay was reversed, where larvae were acclimated to darkness for 1
hour, and baseline activity was tracked for 30 minutes followed by five 1-second white
light flashes with intermittent darkness for 29 seconds. For VSR-OFF and VSR-ON, six
behavioral parameters were quantified using custom MATLAB code[60] (available on
github at https://github.com/ehoffmanlab/Weinschutz-Mendes-et-al-2023-behavior;
DOI:10.5281/zenodo.7644898): (i) average intensity of all startle responses; (ii) average
post-stimulus activity; (iii) average activity after first stimulus; (iv) stimulus versus post-
stimulus activity; (v) intensity of responses to the first stimulus; (vi) intensity of
responses to the final stimulus. The sleep-wake paradigm was conducted between 5-7
dpf, following the VSR-OFF and VSR-ON assays. During a 14h:10h white light:darkness
cycle, larvae activity and sleep patterns were tracked within the Zebrabox and analyzed
with custom MATLAB code[60] (available on github at (https://github.com/JRihel/Sleep-
Analysis/tree/Sleep-Analysis-Code; DOI: 10.5281/zenodo.7644073). Six behavioral
parameters were quantified for daytime and nighttime: (i) total activity; (ii) total sleep; (iii)
waking activity; (iv) rest bouts; (v) sleep length; (vi) sleep latency. Across VSR-OFF,
VSR-ON, and sleep-wake assays, we analyzed 24 parameters.

Behavioral analysis

Linear mixed models (LMM) were used to compare phenotypes of each behavioral
parameter between homozygous mutant versus wild-type or crispant versus scramble-
injected fish for each gene of interest. Variations of behavioral phenotypes across
experiments were accounted for by including the date of the experiment as a random
effect in LMM. Hierarchical clustering analysis was performed to cluster mutants and
behavioral parameters based on signed -log10-transformed p-values from LMM, where

sign indicates direction of the difference in behavioral phenotype when comparing
stable mutant to wild-type or crispant to scrambled-injected. Pearson correlation
analysis was used to assess correlations between mutants based on the difference in
the 24 parameters. Difference was evaluated using signed -log10-transformed p-values.

Drug prioritization based on zebrafish pharmaco-behavioral profiles

NDD gene-associated mutant and crispant behavioral phenotypes were compared to a
dataset of 376 U.S. FDA-approved drugs that were screened for their behavioral effects
in larval zebrafish using the visual-startle and sleep-wake assays described above.
These drugs have a significant effect on at least two behavioral parameters (LMM,
$p<0.05/3$, corrected for three behavioral assays). Pearson's correlation analysis was
used to identify drugs that significantly correlate (correlation >0.5, $p<0.05$, t-statistic) or
anti-correlate (correlation <-0.5, $p<0.05$, t-statistic) with mutant behavioral signatures (**SI**
Data 2-3).

Drug prioritization based on perturbation signature reversal in LiNCs Neuronal Cell
Lines. To identify drugs that could reverse cell-type specific convergence across
different KOs, we used the Query tool from The Broad Institute's Connectivity Map
(Cmap) Server[78]. Briefly, the tool computes weighted enrichment scores (WTCS)
between the query set and each signature in the Cmap LiNCs gene expression data
(dose, time, drug, cell-line), normalizes the WTCS by dividing by the signed mean within
each perturbation (NCS), and computes FDR as fraction of "null signatures" (DMSO)
where the absolute NCS exceeds reference signature. We prioritized drugs that were
negatively enriched for convergent signatures specifically in neuronal cells (either
neurons (NEU) or neural progenitor cells (NPCs) with NCS $<= -1.00$, FDR$<=0.05$) and
filtered for drugs that had clinical data in humans and paired behavioral phenotyping in
zebrafish (**SI Data 2**).

Targeted drug rescue of behavioral phenotypes in zebrafish

For mutant-x-drug experiments, larval activity was monitored from 5-7 dpf using the
behavioral assays described above. Individual wild-type zebrafish larvae were added to
each well of a 96-well plate containing 650 µl of standard embryo water. A 5 mM stock
solution of each compound dissolved in DMSO or DMSO alone (control) was pipetted
directly into each well after which the visual-startle and sleep-wake assays were
performed. Drugs were tested at a final concentration of 10 µM (0.1% DMSO final
concentration) in 12-24 background-matched homozygous or wild-type larvae or 24
crispant or scrambled control-injected larvae with genotyping conducted after each
experiment to confirm genotypes for stable mutant lines and confirm on-target mutations
in crispants.

For behaviors that were nominally significantly different between mutant+DMSO and
WT+DMSO ($p<0.06$), we characterized the effect of the mutant-x-Drug on behavior as:
i) "exacerbated" [significant effect mutant+Drug-v-WT > significant effect mutant-v-WT] if
mutant behavior $p<=0.06$ and mutant-x-drug behavior p.value $<=$ mutant behavior
p.value with increased absolute beta values (i.e., stronger p-value with appreciable
difference in the magnitude of effect but not direction); ii) "unchanged" [significant effect
mutant+drug-v-WT = significant effect mutant-v-WT]; iii) "partial rescue" [significant
effect mutant+Drug-v-WT < significant effect mutant-v-WT], if mutant behavior $p<=0.06$

and mutant-x-drug behavior p>0.06 or if mutant behavior p.value <= mutant-x-drug
behavior p.value with reduced effects on the absolute beta value; iv) "rescued" [sig.
effect mutant-v-WT, no sig. effect mutant+Drug-v-WT], mutant behavior p<=0.06 and
mutant-x-drug behavior p>0.06; v) "over-corrected" [mutant+Drug-v-WT opposite
direction of sig. effect mutant-v-WT]. mutant behavior p<=0.06 and mutant-x-drug
behavior p<=0.06, with opposing directions of effect. Note "drug specific/side-
effects" indicate significant mutant-by-drug effects.

1271

STATEMENT OF ETHICS

Yale University Institutional Review Board waived ethical approval for this work. Ethical
approval was not required because the hiPSC lines, lacking association with any
identifying information and widely accessible from a public repository, are thus not
considered to be human subject research. Post-mortem brain data are similarly lacking
identifiable information and are not considered human subject research.

All procedures involving zebrafish were conducted in accordance with Institutional
Animal Care and Use Committee (IACUC; Protocol #2024-20054) regulatory standards
at Yale University.

1281

1282 **CONFLICT OF INTEREST STATEMENT**

1283 The authors declare no conflict of interest.

1284

FUNDING SOURCES

This work was supported by F31MH130122 (K.R.T), HHMI Gilliams Fellowship (A.P.),
Autism Science Foundation (A.P.), T32MH014276 (M.F.G.), T32GM136651 (E.D., S.F.)
R01MH123155 (K.J.B.), RM1MH132648 (K.J.B. and E.J.H.), R01MH121074 (K.J.B.),
R01MH116002 (E.J.H.) R21MH133245 (E.J.H.), and R01ES033630 (L.H., K.J.B.),
R01MH124839 (LMH), R01MH118278 (L.MH.), Simons Foundation (#1012863KB,
#573508EH and #345993EH), Spector Fund, (E.J.H. and Swebilius Foundation (E.J.H.);
Kavli Foundation (E.J.H.); BD2: Breakthrough Discoveries for thriving with Bipolar
Disorder (#DG230102 H.S., M.D., T.C.H., K.J.B.), the European Union's Horizon 2020
research and innovation programme under the Marie Skłodowska-Curie grant
(#101065629 N.B.); and Interdepartmental Neuroscience Program at Yale (A.P).

1296

AUTHOR CONTRIBUTIONS

MFG designed and executed the NDD gene ECCITEseq in iNPCs, iGLUTs, and
iGABAs, with technical assistance from SC, OL, and JC and support from PJMD. KRT
conducted all bioinformatic analyses and generated all figures, with technical assistance
from AS. Arrayed KO lines and neuronal cultures were generated, validated,
phenotyped, and analyzed by NB. NB, T-CH, CB, ALTS, and JL conducted
mitochondrial phenotypic analyses. SBT, AP, ED, YD, SEF, and SK, generated,
phenotyped, and analyzed zebrafish mutants with technical assistance from GD and

bioinformatic supervision from ZW. JW, RM, ZC, and MG conducted LNCTP analyses. Funding and mentorship provided by LH, EJH, and KJB. Manuscript was written by KJB with extensive feedback from LH, EJH, and KRT as well as contributions for all authors.

Special thanks to Michael Talkowski and Douglas Ruderfer for countless discussions on convergence and to Summer Thyme and the Thyme lab for sharing zebrafish mutant lines.

INCLUSION AND DIVERSITY

One or more of the authors of this paper self-identifies as an under-represented ethnic minority in their field of research or within their geographical location. One or more of the authors of this paper self-identifies as living with a disability. One or more of the authors of this paper self-identifies as a gender minority in their field of research. One or more of the authors of this paper self-identifies as a member of the LGBTQIA+ community. One or more of the authors of this paper received support from a program designed to increase minority representation in their field of research.

DATA AVAILABILITY

All source donor hiPSCs have been deposited at the Rutgers University Cell and DNA Repository (study 160; http://www.nimhstemcells.org/).

sc-RNA sequencing data reported in this paper will be uploaded to Gene expression omnibus (GEO) prior to publication. Previously published SCZ-CRISPRa screen datasets that were used for external validation of random forest models are available on the GEO (GSE200774) and on Synapse (syn27819129).

CODE AVAILABILITY

The full analysis pipeline (including code and processed data objects) used for analysis of single-cell CRISPR-KO data, evaluation and characterization of gene-level and network level convergence, and predictive modeling using random forest will be publicly available through Synpase prior to publication.

Custom MATLAB software developed by the Hoffman Lab to analyze visual-startle response parameters is available on github at https://github.com/ehoffmanlab/Weinschutz-Mendes-et-al-2023-behavior; https://doi.org/10.5281/zenodo.7644898. Custom MATLAB sofyware developed by Jason Rihel to analyze sleep-wake assays is available on github at https://github.com/JRihel/Sleep-Analysis/tree/ Sleep-Analysis-Code; https://doi.org/10.5281/zenodo.7644073.

REFERENCES

1   Sandin, S. *et al.* The Heritability of Autism Spectrum Disorder. *JAMA* **318**, 1182-1184 (2017). https://doi.org/10.1001/jama.2017.12141

2   Grove, J. *et al.* Identification of common genetic risk variants for autism spectrum
disorder. *Nat Genet* **51**, 431-444 (2019). https://doi.org/10.1038/s41588-019-
0344-8
3   Mahjani, B. *et al.* Prevalence and phenotypic impact of rare potentially damaging
variants in autism spectrum disorder. *Mol Autism* **12**, 65 (2021).
https://doi.org/10.1186/s13229-021-00465-3
4   Satterstrom, F. K. *et al.* Large-Scale Exome Sequencing Study Implicates Both
Developmental and Functional Changes in the Neurobiology of Autism. *Cell* **180**,
568-584 e523 (2020). https://doi.org/10.1016/j.cell.2019.12.036
5   Kaplanis, J. *et al.* Evidence for 28 genetic disorders discovered by combining
healthcare and research data. *Nature* **586**, 757-762 (2020).
https://doi.org/10.1038/s41586-020-2832-5
6   Coe, B. P. *et al.* Neurodevelopmental disease genes implicated by de novo
mutation and copy number variation morbidity. *Nat Genet* **51**, 106-116 (2019).
https://doi.org/10.1038/s41588-018-0288-4
7   Singh, T. *et al.* Rare coding variants in ten genes confer substantial risk for
schizophrenia. *Nature* **604**, 509-516 (2022). https://doi.org/10.1038/s41586-022-
04556-w
8   Palmer, D. S. *et al.* Exome sequencing in bipolar disorder identifies AKAP11 as a
risk gene shared with schizophrenia. *Nat Genet* **54**, 541-547 (2022).
https://doi.org/10.1038/s41588-022-01034-x
9   Willsey, A. J. *et al.* Coexpression networks implicate human midfetal deep
cortical projection neurons in the pathogenesis of autism. *Cell* **155**, 997-1007
(2013). https://doi.org/10.1016/j.cell.2013.10.020
10   De Rubeis, S. *et al.* Synaptic, transcriptional and chromatin genes disrupted in
autism. *Nature* **515**, 209-215 (2014). https://doi.org/10.1038/nature13772
11   O'Roak, B. J. *et al.* Recurrent de novo mutations implicate novel genes
underlying simplex autism risk. *Nat Commun* **5**, 5595 (2014).
https://doi.org/10.1038/ncomms6595
12   Talkowski, M. E. *et al.* Sequencing chromosomal abnormalities reveals
neurodevelopmental loci that confer risk across diagnostic boundaries. *Cell* **149**,
525-537 (2012). https://doi.org/10.1016/j.cell.2012.03.028
13   Neale, B. M. *et al.* Patterns and rates of exonic de novo mutations in autism
spectrum disorders. *Nature* **485**, 242-245 (2012). https://doi.org/nature11011 [pii]
10.1038/nature11011
14   O'Roak, B. J. *et al.* Multiplex targeted sequencing identifies recurrently mutated
genes in autism spectrum disorders. *Science* **338**, 1619-1622 (2012).
https://doi.org/10.1126/science.1227764
15   Sanders, S. J. *et al.* De novo mutations revealed by whole-exome sequencing
are strongly associated with autism. *Nature* **485**, 237-241 (2012).
https://doi.org/nature10945 [pii] 10.1038/nature10945
16   Fazel Darbandi, S. *et al.* Five autism-associated transcriptional regulators target
shared loci proximal to brain-expressed genes. *Cell reports* **43**, 114329 (2024).
https://doi.org/10.1016/j.celrep.2024.114329

Markenscoff-Papadimitriou, E. *et al.* Autism risk gene POGZ promotes chromatin accessibility and expression of clustered synaptic genes. *Cell reports* **37**, 110089 (2021). https://doi.org/10.1016/j.celrep.2021.110089

Notwell, J. H. *et al.* TBR1 regulates autism risk genes in the developing neocortex. *Genome Res* **26**, 1013-1022 (2016). https://doi.org/10.1101/gr.203612.115

Sugathan, A. *et al.* CHD8 regulates neurodevelopmental pathways associated with autism spectrum disorder in neural progenitors. *Proc Natl Acad Sci U S A* **111**, E4468-4477 (2014). https://doi.org/10.1073/pnas.1405266111

Cotney, J. *et al.* The autism-associated chromatin modifier CHD8 regulates other autism risk genes during human neurodevelopment. *Nat Commun* **6**, 6404 (2015). https://doi.org/10.1038/ncomms7404

O'Neill, A. C. *et al.* Spatial centrosome proteome of human neural cells uncovers disease-relevant heterogeneity. *Science* **376**, eabf9088 (2022). https://doi.org/10.1126/science.abf9088

Sun, N. *et al.* Autism genes converge on microtubule biology and RNA-binding proteins during excitatory neurogenesis. *bioRxiv*, 2023.2012.2022.573108 (2024). https://doi.org/10.1101/2023.12.22.573108

Kostyanovskaya, E. *et al.* Convergence of autism proteins at the cilium. *bioRxiv* (2025). https://doi.org/10.1101/2024.12.05.626924

Teerikorpi, N. *et al.* Ciliary biology intersects autism and congenital heart disease. *bioRxiv* (2024). https://doi.org/10.1101/2024.07.30.602578

Li, C. *et al.* Single-cell brain organoid screening identifies developmental defects in autism. *bioRxiv*, 2022.2009.2015.508118 (2022). https://doi.org/10.1101/2022.09.15.508118

Martins-Costa, C. *et al.* ARID1B controls transcriptional programs of axon projection in an organoid model of the human corpus callosum. *Cell stem cell* **31**, 866-885 e814 (2024). https://doi.org/10.1016/j.stem.2024.04.014

Paulsen, B. *et al.* Autism genes converge on asynchronous development of shared neuron classes. *Nature* **602**, 268-273 (2022). https://doi.org/10.1038/s41586-021-04358-6

Villa, C. E. *et al.* CHD8 haploinsufficiency links autism to transient alterations in excitatory and inhibitory trajectories. *Cell reports* **39**, 110615 (2022). https://doi.org/10.1016/j.celrep.2022.110615

Flaherty, E. *et al.* Neuronal impact of patient-specific aberrant NRXN1alpha splicing. *Nat Genet* **51**, 1679-1690 (2019). https://doi.org/10.1038/s41588-019-0539-z

Sebastian, R. *et al.* Schizophrenia-associated NRXN1 deletions induce developmental-timing- and cell-type-specific vulnerabilities in human brain organoids. *Nature Communications* **14**, 3770 (2023). https://doi.org/10.1038/s41467-023-39420-6

Birtele, M. *et al.* Non-synaptic function of the autism spectrum disorder-associated gene SYNGAP1 in cortical neurogenesis. *Nat Neurosci* **26**, 2090-2103 (2023). https://doi.org/10.1038/s41593-023-01477-3

Ellingford, R. A. *et al.* Cell-type-specific synaptic imbalance and disrupted homeostatic plasticity in cortical circuits of ASD-associated Chd8

haploinsufficient   mice.   *Mol   Psychiatry*   **26**,   3614-3624   (2021).
https://doi.org/10.1038/s41380-021-01070-9
33 Shi, X. *et al.* Heterozygous deletion of the autism-associated gene CHD8 impairs
synaptic function through widespread changes in gene expression and chromatin
compaction.   *Am   J   Hum   Genet*   **110**,   1750-1768   (2023).
https://doi.org/10.1016/j.ajhg.2023.09.004
34 Pak, C. *et al.* Cross-platform validation of neurotransmitter release impairments
in schizophrenia patient-derived NRXN1-mutant neurons. *Proc Natl Acad Sci U S*
A **118** (2021). https://doi.org/10.1073/pnas.2025598118
35 Yi, F. *et al.* Autism-associated SHANK3 haploinsufficiency causes Ih
channelopathy   in   human   neurons.   *Science*   **352**,   aaf2669   (2016).
https://doi.org/10.1126/science.aaf2669
36 Vermaercke, B. *et al.* SYNGAP1 deficiency disrupts synaptic neoteny in
xenotransplanted   human   cortical   neurons   in   vivo.   *Neuron*   (2024).
https://doi.org/10.1016/j.neuron.2024.07.007
37 Jung, E. M. *et al.* Arid1b haploinsufficiency disrupts cortical interneuron
development and mouse behavior. *Nat Neurosci* **20**, 1694-1707 (2017).
https://doi.org/10.1038/s41593-017-0013-0
38 Fernando, M. B. *et al.* Phenotypic complexities of rare heterozygous neurexin-1
deletions. *bioRxiv* (2024). https://doi.org/10.1101/2023.10.28.564543
39 Chen, Q. *et al.* Dysfunction of cortical GABAergic neurons leads to sensory
hyper-reactivity in a Shank3 mouse model of ASD. *Nat Neurosci* **23**, 520-532
(2020). https://doi.org/10.1038/s41593-020-0598-6
40 Geschwind, D. H. Autism: many genes, common pathways? *Cell* **135**, 391-395
(2008). https://doi.org/10.1016/j.cell.2008.10.016
41 Willsey, H. R., Willsey, A. J., Wang, B. & State, M. W. Genomics, convergent
neuroscience and progress in understanding autism spectrum disorder. *Nat Rev*
Neurosci **23**, 323-341 (2022). https://doi.org/10.1038/s41583-022-00576-7
42 Bicks, L. K. & Geschwind, D. H. Functional neurogenomics in autism spectrum
disorders: A decade of progress. *Curr Opin Neurobiol* **86**, 102858 (2024).
https://doi.org/10.1016/j.conb.2024.102858
43 Quesnel-Vallieres, M., Weatheritt, R. J., Cordes, S. P. & Blencowe, B. J. Autism
spectrum disorder: insights into convergent mechanisms from transcriptomics.
Nat Rev Genet **20**, 51-63 (2019). https://doi.org/10.1038/s41576-018-0066-2
44 Parikshak, N. N. *et al.* Integrative functional genomic analyses implicate specific
molecular pathways and circuits in autism. *Cell* **155**, 1008-1021 (2013).
https://doi.org/10.1016/j.cell.2013.10.031
45 Voineagu, I. *et al.* Transcriptomic analysis of autistic brain reveals convergent
molecular pathology. *Nature* **474**, 380-384 (2011). https://doi.org/nature10110
[pii]
10.1038/nature10110
46 Liao, C. *et al.* Convergent coexpression of autism-associated genes suggests
some novel risk genes may not be detectable in large-scale genetic studies. *Cell*
Genom **3**, 100277 (2023). https://doi.org/10.1016/j.xgen.2023.100277

47   Pintacuda, G. *et al.* Protein interaction studies in human induced neurons indicate convergent biology underlying autism spectrum disorders. *Cell Genom* **3**, 100250 (2023). https://doi.org/10.1016/j.xgen.2022.100250

48   Wang, B. *et al.* A foundational atlas of autism protein interactions reveals molecular convergence. *bioRxiv* (2023). https://doi.org/10.1101/2023.12.03.569805

49   Murtaza, N. *et al.* Neuron-specific protein network mapping of autism risk genes identifies shared biological mechanisms and disease-relevant pathologies. *Cell reports* **41**, 111678 (2022). https://doi.org/10.1016/j.celrep.2022.111678

50   Gao, Y. *et al.* Proximity analysis of native proteomes reveals phenotypic modifiers in a mouse model of autism and related neurodevelopmental conditions. *Nat Commun* **15**, 6801 (2024). https://doi.org/10.1038/s41467-024-51037-x

51   Rubenstein, J. L. & Merzenich, M. M. Model of autism: increased ratio of excitation/inhibition in key neural systems. *Genes, brain, and behavior* **2**, 255-267 (2003). https://doi.org/10.1034/j.1601-183x.2003.00037.x

52   Antoine, M. W., Langberg, T., Schnepel, P. & Feldman, D. E. Increased Excitation-Inhibition Ratio Stabilizes Synapse and Circuit Excitability in Four Autism Mouse Models. *Neuron* **101**, 648-661 e644 (2019). https://doi.org/10.1016/j.neuron.2018.12.026

53   Nelson, S. B. & Valakh, V. Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders. *Neuron* **87**, 684-698 (2015). https://doi.org/10.1016/j.neuron.2015.07.033

54   Cederquist, G. Y. *et al.* A Multiplex Human Pluripotent Stem Cell Platform Defines Molecular and Functional Subclasses of Autism-Related Genes. *Cell stem cell* **27**, 35-49 e36 (2020). https://doi.org/10.1016/j.stem.2020.06.004

55   Lalli, M. A., Avey, D., Dougherty, J. D., Milbrandt, J. & Mitra, R. D. High-throughput single-cell functional elucidation of neurodevelopmental disease-associated genes reveals convergent mechanisms altering neuronal differentiation. *Genome Res* **30**, 1317-1331 (2020). https://doi.org/10.1101/gr.262295.120

56   Meng, X. *et al.* Assembloid CRISPR screens reveal impact of disease genes in human neurodevelopment. *Nature* **622**, 359-366 (2023). https://doi.org/10.1038/s41586-023-06564-w

57   Li, C. *et al.* Single-cell brain organoid screening identifies developmental defects in autism. *Nature* **621**, 373-380 (2023). https://doi.org/10.1038/s41586-023-06473-y

58   Jin, X. *et al.* In vivo Perturb-Seq reveals neuronal and glial abnormalities associated with autism risk genes. *Science* **370** (2020). https://doi.org/10.1126/science.aaz6063

59   Willsey, H. R. *et al.* Parallel in vivo analysis of large-effect autism genes implicates cortical neurogenesis and estrogen in risk and resilience. *Neuron* **109**, 1409 (2021). https://doi.org/10.1016/j.neuron.2021.03.030

60   Weinschutz Mendes, H. *et al.* High-throughput functional analysis of autism genes in zebrafish identifies convergence in dopaminergic and neuroimmune

pathways. *Cell reports* **42**, 112243 (2023).
https://doi.org/10.1016/j.celrep.2023.112243

61 Fu, J. M. *et al.* Rare coding variation provides insight into the genetic architecture
and phenotypic context of autism. *Nat Genet* **54**, 1320-1331 (2022).
https://doi.org/10.1038/s41588-022-01104-0

62 Marshall, C. R. *et al.* Contribution of copy number variants to schizophrenia from
a genome-wide study of 41,321 subjects. *Nat Genet* **49**, 27-35 (2017).
https://doi.org/10.1038/ng.3725

63 Johannesen, K. M. *et al.* Defining the phenotypic spectrum of SLC6A1 mutations.
Epilepsia **59**, 389-402 (2018). https://doi.org/10.1111/epi.13986

64 Heyne, H. O. *et al.* De novo variants in neurodevelopmental disorders with
epilepsy. *Nat Genet* **50**, 1048-1053 (2018). https://doi.org/10.1038/s41588-018-
0143-7

65 de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: generalized
gene-set analysis of GWAS data. *PLoS Comput Biol* **11**, e1004219 (2015).
https://doi.org/10.1371/journal.pcbi.1004219

66 Sullivan, P. F. & Geschwind, D. H. Defining the Genetic, Genomic, Cellular, and
Diagnostic Architectures of Psychiatric Disorders. *Cell* **177**, 162-183 (2019).
https://doi.org/10.1016/j.cell.2019.01.015

67 Schrode, N. *et al.* Synergistic effects of common schizophrenia risk variants. *Nat*
Genet **51**, 1475-1485 (2019). https://doi.org/10.1038/s41588-019-0497-5

68 Wells, M. F. *et al.* Natural variation in gene expression and viral susceptibility
revealed by neural progenitor cell villages. *Cell stem cell* **30**, 312-332 e313
(2023). https://doi.org/10.1016/j.stem.2023.01.010

69 Zhang, Y. *et al.* Rapid single-step induction of functional neurons from human
pluripotent stem cells. *Neuron* **78**, 785-798 (2013).
https://doi.org/10.1016/j.neuron.2013.05.029

70 Yang, N. *et al.* Generation of pure GABAergic neurons by transcription factor
programming. *Nat Methods* (2017). https://doi.org/10.1038/nmeth.4291

71 Mimitou, E. P. *et al.* Multiplexed detection of proteins, transcriptomes, clonotypes
and CRISPR perturbations in single cells. *Nat Methods* **16**, 409-412 (2019).
https://doi.org/10.1038/s41592-019-0392-0

72 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587 e3529 (2021). https://doi.org/10.1016/j.cell.2021.04.048

73 Deans, P. J. M. *et al.* Convergent impact of schizophrenia risk genes. *bioRxiv*,
2022.2003.2029.486286 (2025). https://doi.org/10.1101/2022.03.29.486286

74 Deans, P. M. *et al.* Non-additive effects of schizophrenia risk genes reflect
convergent downstream function. *medRxiv*, 2023.2003.2020.23287497 (2023).
https://doi.org/10.1101/2023.03.20.23287497

75 Breiman, L. Random Forests. *Machine Learning* **45**, 5-32 (2001).
https://doi.org/10.1023/A:1010933404324

76 Emani, P. S. *et al.* Single-cell genomics and regulatory networks for 388 human
brains. *Science* **384**, eadi5199 (2024). https://doi.org/10.1126/science.adi5199

77 Gonzalez, F. *et al.* An iCRISPR platform for rapid, multiplexable, and inducible
genome editing in human pluripotent stem cells. *Cell stem cell* **15**, 215-226
(2014). https://doi.org/10.1016/j.stem.2014.05.018

78   Subramanian, A. *et al.* A Next Generation Connectivity Map: L1000 Platform and
the First 1,000,000 Profiles. *Cell* **171**, 1437-1452 e1417 (2017).
https://doi.org/10.1016/j.cell.2017.10.049
79   Manji, H. *et al.* Impaired mitochondrial function in psychiatric disorders. *Nat Rev*
Neurosci **13**, 293-307 (2012). https://doi.org/10.1038/nrn3229
80   Rossignol, D. A. & Frye, R. E. Mitochondrial dysfunction in autism spectrum
disorders: a systematic review and meta-analysis. *Mol Psychiatry* **17**, 290-314
(2012). https://doi.org/10.1038/mp.2010.136
81   Wang, Y., Picard, M. & Gu, Z. Genetic Evidence for Elevated Pathogenicity of
Mitochondrial DNA Heteroplasmy in Autism Spectrum Disorder. *PLoS genetics*
12, e1006391 (2016). https://doi.org/10.1371/journal.pgen.1006391
82   Varga, N. A. *et al.* Mitochondrial dysfunction and autism: comprehensive genetic
analyses of children with autism and mtDNA deletion. *Behav Brain Funct* **14**, 4
(2018). https://doi.org/10.1186/s12993-018-0135-x
83   Chalkia, D. *et al.* Association Between Mitochondrial DNA Haplogroup Variation
and Autism Spectrum Disorders. *JAMA Psychiatry* **74**, 1161-1168 (2017).
https://doi.org/10.1001/jamapsychiatry.2017.2604
84   Wang, Y. *et al.* Association of mitochondrial DNA content, heteroplasmies and
inter-generational transmission with autism. *Nat Commun* **13**, 3790 (2022).
https://doi.org/10.1038/s41467-022-30805-7
85   Yardeni, T. *et al.* An mtDNA mutant mouse demonstrates that mitochondrial
deficiency can result in autism endophenotypes. *Proc Natl Acad Sci U S A* **118**
(2021). https://doi.org/10.1073/pnas.2021429118
86   Shen, M. *et al.* Reduced mitochondrial fusion and Huntingtin levels contribute to
impaired dendritic maturation and behavioral deficits in Fmr1-mutant mice. *Nat*
Neurosci **22**, 386-400 (2019). https://doi.org/10.1038/s41593-019-0338-y
87   Fernandez, A. *et al.* Mitochondrial Dysfunction Leads to Cortical Under-
Connectivity and Cognitive Impairment. *Neuron* **102**, 1127-1142 e1123 (2019).
https://doi.org/10.1016/j.neuron.2019.04.013
88   Gandal, M. J. *et al.* Shared molecular neuropathology across major psychiatric
disorders parallels polygenic overlap. *Science* **359**, 693-697 (2018).
https://doi.org/10.1126/science.aad6469
89   Kanellopoulos, A. K. *et al.* Aralar Sequesters GABA into Hyperactive
Mitochondria, Causing Social Behavior Deficits. *Cell* **180**, 1178-1197 e1120
(2020). https://doi.org/10.1016/j.cell.2020.02.044
90   Li, J. *et al.* Mitochondrial deficits in human iPSC-derived neurons from patients
with 22q11.2 deletion syndrome and schizophrenia. *Translational psychiatry* **9**,
302 (2019). https://doi.org/10.1038/s41398-019-0643-y
91   Li, J. *et al.* Association of Mitochondrial Biogenesis With Variable Penetrance of
Schizophrenia. *JAMA Psychiatry* **78**, 911-921 (2021).
https://doi.org/10.1001/jamapsychiatry.2021.0762
92   Schafer, S. T. *et al.* Pathological priming causes developmental gene network
heterochronicity in autistic subject-derived neurons. *Nat Neurosci* **22**, 243-255
(2019). https://doi.org/10.1038/s41593-018-0295-x

93 Rajarajan, P. *et al.* Neuron-specific signatures in the chromosomal connectome
associated with schizophrenia risk. *Science* **362** (2018).
https://doi.org/10.1126/science.aat4311
94 Wen, C. *et al.* Cross-ancestry atlas of gene, isoform, and splicing regulation in
the developing human brain. *Science* **384**, eadh0829 (2024).
https://doi.org/10.1126/science.adh0829
95 Wong, C. C. Y. *et al.* Genome-wide DNA methylation profiling identifies
convergent molecular signatures associated with idiopathic and syndromic
autism in post-mortem human brain tissue. *Hum Mol Genet* **28**, 2201-2211
(2019). https://doi.org/10.1093/hmg/ddz052
96 Ramaswami, G. *et al.* Integrative genomics identifies a convergent molecular
subtype that links epigenomic with transcriptomic differences in autism. *Nat*
Commun **11**, 4873 (2020). https://doi.org/10.1038/s41467-020-18526-1
97 Bell, J. Stratified medicines: towards better treatment for disease. *Lancet* **383**
Suppl 1, S3-5 (2014). https://doi.org/10.1016/S0140-6736(14)60115-X
98 Tsimberidou, A. M. *et al.* Molecular tumour boards - current and future
considerations for precision oncology. *Nature reviews. Clinical oncology* **20**, 843-
863 (2023). https://doi.org/10.1038/s41571-023-00824-4
99 Zhang, H., Colclough, K., Gloyn, A. L. & Pollin, T. I. Monogenic diabetes: a
gateway to precision medicine in diabetes. *J Clin Invest* **131** (2021).
https://doi.org/10.1172/JCI142244
100 Gaugler, T. *et al.* Most genetic risk for autism resides with common variation. *Nat*
Genet **46**, 881-885 (2014). https://doi.org/10.1038/ng.3039
101 Schaaf, C. P. *et al.* A framework for an evidence-based gene list relevant to
autism spectrum disorder. *Nat Rev Genet* **21**, 367-376 (2020).
https://doi.org/10.1038/s41576-020-0231-2
102 Shi, Y. *et al.* Multi-polygenic scores in psychiatry: From disorder specific to
transdiagnostic perspectives. *Am J Med Genet B Neuropsychiatr Genet* **195**,
e32951 (2024). https://doi.org/10.1002/ajmg.b.32951
103 Andreassen, O. A., Hindley, G. F. L., Frei, O. & Smeland, O. B. New insights
from the last decade of research in psychiatric genetics: discoveries, challenges
and clinical implications. *World Psychiatry* **22**, 4-24 (2023).
https://doi.org/10.1002/wps.21034
104 Weiner, D. J. *et al.* Statistical and functional convergence of common and rare
genetic influences on autism at chromosome 16p. *Nat Genet* **54**, 1630-1639
(2022). https://doi.org/10.1038/s41588-022-01203-y
105 Weiner, D. J. *et al.* Polygenic transmission disequilibrium confirms that common
and rare variation act additively to create risk for autism spectrum disorders. *Nat*
Genet **49**, 978-985 (2017). https://doi.org/10.1038/ng.3863
106 Klei, L. *et al.* How rare and common risk variation jointly affect liability for autism
spectrum disorder. *Mol Autism* **12**, 66 (2021). https://doi.org/10.1186/s13229-
021-00466-2
107 Bergen, S. E. *et al.* Joint Contributions of Rare Copy Number Variants and
Common SNPs to Risk for Schizophrenia. *Am J Psychiatry* **176**, 29-35 (2019).
https://doi.org/10.1176/appi.ajp.2018.17040467

Akingbuwa, W. A., Hammerschlag, A. R., Bartels, M., Nivard, M. G. & Middeldorp, C. M. Ultra-rare and common genetic variant analysis converge to implicate negative selection and neuronal processes in the aetiology of schizophrenia. *Mol Psychiatry* **27**, 3699-3707 (2022). https://doi.org/10.1038/s41380-022-01621-8

Oliver, K. L. *et al.* Common risk variants for epilepsy are enriched in families previously targeted for rare monogenic variant discovery. *EBioMedicine* **81**, 104079 (2022). https://doi.org/10.1016/j.ebiom.2022.104079

Genetic Modifiers of Huntington's Disease, C. Identification of Genetic Factors that Modify Clinical Onset of Huntington's Disease. *Cell* **162**, 516-526 (2015). https://doi.org/10.1016/j.cell.2015.07.003

Kingdom, R., Beaumont, R. N., Wood, A. R., Weedon, M. N. & Wright, C. F. Genetic modifiers of rare variants in monogenic developmental disorder loci. *Nat Genet* **56**, 861-868 (2024). https://doi.org/10.1038/s41588-024-01710-0

Dobrindt, K. *et al.* Publicly Available hiPSC Lines with Extreme Polygenic Risk Scores for Modeling Schizophrenia. *Complex Psychiatry* **6**, 68-82 (2021). https://doi.org/10.1159/000512716

Bjork, M. H. *et al.* Association of Prenatal Exposure to Antiseizure Medication With Risk of Autism and Intellectual Disability. *JAMA Neurol* **79**, 672-681 (2022). https://doi.org/10.1001/jamaneurol.2022.1269

Anton-Bolanos, N. *et al.* Brain Chimeroids reveal individual susceptibility to neurotoxic triggers. *Nature* (2024). https://doi.org/10.1038/s41586-024-07578-8

Gandal, M. J. *et al.* Broad transcriptomic dysregulation occurs across the cerebral cortex in ASD. *Nature* **611**, 532-539 (2022). https://doi.org/10.1038/s41586-022-05377-7

Wamsley, B. *et al.* Molecular cascades and cell type-specific signatures in ASD revealed by single-cell genomics. *Science* **384**, eadh2602 (2024). https://doi.org/10.1126/science.adh2602

Yap, C. X. *et al.* Brain cell-type shifts in Alzheimer's disease, autism, and schizophrenia interrogated using methylomics and genetics. *Sci Adv* **10**, eadn7655 (2024). https://doi.org/10.1126/sciadv.adn7655

Zhang, P. *et al.* Neuron-specific transcriptomic signatures indicate neuroinflammation and altered neuronal activity in ASD temporal cortex. *Proc Natl Acad Sci U S A* **120**, e2206758120 (2023). https://doi.org/10.1073/pnas.2206758120

Bhattacharya, A. *et al.* Isoform-level transcriptome-wide association uncovers genetic risk mechanisms for neuropsychiatric disorders in the human brain. *Nature genetics* **55**, 2117-2128 (2023). https://doi.org/10.1038/s41588-023-01560-2

Gandal, M. J. *et al.* Transcriptome-wide isoform-level dysregulation in ASD, schizophrenia, and bipolar disorder. *Science* **362** (2018). https://doi.org/10.1126/science.aat8127

Han, V. X., Patel, S., Jones, H. F. & Dale, R. C. Maternal immune activation and neuroinflammation in human neurodevelopmental disorders. *Nat Rev Neurol* **17**, 564-579 (2021). https://doi.org/10.1038/s41582-021-00530-8

122 Seah, C. *et al.* Modeling gene x environment interactions in PTSD using human
neurons reveals diagnosis-specific glucocorticoid-induced gene expression. *Nat*
Neurosci **25**, 1434-1445 (2022). https://doi.org/10.1038/s41593-022-01161-y
123 Seah, C. *et al.* Common genetic variation impacts stress response in the brain.
bioRxiv, 2023.2012.2027.573459 (2023).
https://doi.org/10.1101/2023.12.27.573459
124 Retallick-Townsley, K. G. *et al.* Dynamic stress- and inflammatory-based
regulation of psychiatric risk loci in human neurons. *bioRxiv*,
2024.2007.2009.602755 (2024). https://doi.org/10.1101/2024.07.09.602755
125 Cruceanu, C. *et al.* Cell-Type-Specific Impact of Glucocorticoid Receptor
Activation on the Developing Brain: A Cerebral Organoid Study. *Am J Psychiatry*,
appiajp202121010095 (2021). https://doi.org/10.1176/appi.ajp.2021.21010095
126 Teter, O. M. *et al.* CRISPRi-based screen of Autism Spectrum Disorder risk
genes in microglia uncovers roles of ADNP in microglia endocytosis
and uptake of synaptic material. *bioRxiv*, 2024.2006.2001.596962 (2024).
https://doi.org/10.1101/2024.06.01.596962
127 Ma, Y. *et al.* Activity-Dependent Transcriptional Program in NGN2+ Neurons
Enriched for Genetic Risk for Brain-Related Disorders. *Biol Psychiatry* **95**, 187-
198 (2024). https://doi.org/10.1016/j.biopsych.2023.07.003
128 Roussos, P., Guennewig, B., Kaczorowski, D. C., Barry, G. & Brennand, K. J.
Activity-Dependent Changes in Gene Expression in Schizophrenia Human-
Induced Pluripotent Stem Cell Neurons. *JAMA Psychiatry* **73**, 1180-1188 (2016).
https://doi.org/10.1001/jamapsychiatry.2016.2575
129 Sanchez-Priego, C. *et al.* Mapping cis-regulatory elements in human neurons
links psychiatric disease heritability and activity-regulated transcriptional
programs. *Cell reports* **39**, 110877 (2022).
https://doi.org/10.1016/j.celrep.2022.110877
130 Boulting, G. L. *et al.* Activity-dependent regulome of human GABAergic neurons
reveals new patterns of gene regulation and neurological disease heritability. *Nat*
Neurosci **24**, 437-448 (2021). https://doi.org/10.1038/s41593-020-00786-1
131 Ahn, K. *et al.* High rate of disease-related copy number variations in childhood
onset schizophrenia. *Mol Psychiatry* **19**, 568-572 (2014).
https://doi.org/10.1038/mp.2013.59
132 Ahn, K., An, S. S., Shugart, Y. Y. & Rapoport, J. L. Common polygenic variation
and risk for childhood-onset schizophrenia. *Mol Psychiatry* (2014).
https://doi.org/10.1038/mp.2014.158
133 Hoffman, G. E. *et al.* Transcriptional signatures of schizophrenia in hiPSC-
derived NPCs and neurons are concordant with post-mortem adult brains. *Nat*
Commun **8**, 2225 (2017). https://doi.org/10.1038/s41467-017-02330-5
134 Guss, E. J. *et al.* Protocol for neurogenin-2-mediated induction of human stem
cell-derived neural progenitor cells. *Star Protoc* **5**, 102878 (2024).
https://doi.org/10.1016/j.xpro.2024.102878
135 Wang, M. *et al.* Transformative Network Modeling of Multi-omics Data Reveals
Detailed Circuits, Key Regulators, and Potential Therapeutics for Alzheimer's
Disease. *Neuron* **109**, 257-272 e214 (2021).
https://doi.org/10.1016/j.neuron.2020.11.002

136　Ho, S. M. *et al.* Rapid Ngn2-induction of excitatory neurons from hiPSC-derived
neural progenitor cells. *Methods* **101**, 113-124 (2016).
https://doi.org/10.1016/j.ymeth.2015.11.019

137　Marro, S. G. *et al.* Neuroligin-4 Regulates Excitatory Synaptic Transmission in
Human Neurons. *Neuron* **103**, 617-626 e616 (2019).
https://doi.org/10.1016/j.neuron.2019.05.043

138　Zhang, Z. *et al.* The fragile X mutation impairs homeostatic plasticity in human
neurons by blocking synaptic retinoic acid signaling. *Science translational*
medicine **10** (2018). https://doi.org/10.1126/scitranslmed.aar4338

139　Meijer, M. *et al.* A Single-Cell Model for Synaptic Transmission and Plasticity in
Human iPSC-Derived Neurons. *Cell reports* **27**, 2199-2211 e2196 (2019).
https://doi.org/10.1016/j.celrep.2019.04.058

140　Zhang, S. *et al.* Allele-specific open chromatin in human iPSC neurons elucidates
functional disease variants. *Science* **369**, 561-565 (2020).
https://doi.org/10.1126/science.aay3983

141　Sun, Y. *et al.* A deleterious Nav1.1 mutation selectively impairs telencephalic
inhibitory neurons derived from Dravet Syndrome patients. *Elife* **5** (2016).
https://doi.org/10.7554/eLife.13073

142　Barretto, N. *et al.* ASCL1- and DLX2-induced GABAergic neurons from hiPSC-
derived NPCs. *J Neurosci Methods* **334**, 108548 (2020).
https://doi.org/10.1016/j.jneumeth.2019.108548

143　Powell, S. K. *et al.* Induction of dopaminergic neurons for neuronal subtype-
specific modeling of psychiatric disease risk. *Mol Psychiatry* **28**, 1970-1982
(2023). https://doi.org/10.1038/s41380-021-01273-0

144　Miller, J. A. *et al.* Transcriptional landscape of the prenatal human brain. *Nature*
508, 199-206 (2014). https://doi.org/10.1038/nature13185

145　Shi, Z. D. *et al.* Genome Editing in hPSCs Reveals GATA6 Haploinsufficiency
and a Genetic Interaction with GATA4 in Human Pancreatic Development. *Cell*
stem cell **20**, 675-688 e676 (2017). https://doi.org/10.1016/j.stem.2017.01.001

146　Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-
21 (2013). https://doi.org/10.1093/bioinformatics/bts635

147　Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**
(2021). https://doi.org/10.1093/gigascience/giab008

148　Lindsay, H. *et al.* CrispRVariants charts the mutation spectrum of genome
engineering experiments. *Nat Biotechnol* **34**, 701-702 (2016).
https://doi.org/10.1038/nbt.3628

149　Yuan, S. H. *et al.* Cell-surface marker signatures for the isolation of neural stem
cells, glia and neurons derived from human pluripotent stem cells. *PLoS One* **6**,
e17540 (2011). https://doi.org/10.1371/journal.pone.0017540

150　Li, W. *et al.* MAGeCK enables robust identification of essential genes from
genome-scale CRISPR/Cas9 knockout screens. *Genome biology* **15**, 554 (2014).
https://doi.org/10.1186/s13059-014-0554-4

151　Birey, F. *et al.* Assembly of functionally integrated human forebrain spheroids.
Nature **545**, 54-59 (2017). https://doi.org/10.1038/nature22330

152 Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-
cell transcriptomic data across different conditions, technologies, and species.
Nat Biotechnol **36**, 411-420 (2018). https://doi.org/10.1038/nbt.4096
153 Papalexi, E. *et al.* Characterizing the molecular regulation of inhibitory immune
checkpoints with multimodal single-cell screens. *Nat Genet* **53**, 322-331 (2021).
https://doi.org/10.1038/s41588-021-00778-2
154 Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by
single-cell   RNA-seq.   *Science*   **352**,   189-196   (2016).
https://doi.org/10.1126/science.aad0501
155 Tian, R. *et al.* CRISPR Interference-Based Platform for Multimodal Genetic
Screens   in   Human   iPSC-Derived   Neurons.   *Neuron*   (2019).
https://doi.org/10.1016/j.neuron.2019.07.014
156 Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of
genomewide   association   scans.   *Bioinformatics*   **26**,   2190-2191   (2010).
https://doi.org/10.1093/bioinformatics/btq340
157 Saha, A. *et al.* Co-expression networks reveal the tissue-specific regulation of
transcription   and   splicing.   *Genome   Res*   **27**,   1843-1858   (2017).
https://doi.org/10.1101/gr.216721.116
158 Gao, C., McDowell, I. C., Zhao, S., Brown, C. D. & Engelhardt, B. E. Context
Specific   and   Differential   Gene   Co-expression   Networks   via   Bayesian
Biclustering.   *PLoS   Comput   Biol*   **12**,   e1004791   (2016).
https://doi.org/10.1371/journal.pcbi.1004791
159 Gao C, B. C., Engelhardt BE. A latent factor model with a mixture of sparse and
dense factors to model gene expression data with confounding effects. *arXiv*
(2013).
160 Yu, G. Gene Ontology Semantic Similarity Analysis Using GOSemSim. *Methods
in molecular biology* **2117**, 207-215 (2020). https://doi.org/10.1007/978-1-0716-
0301-7_11
161 Demontis, D. *et al.* Discovery of the first genome-wide significant risk loci for
attention   deficit/hyperactivity   disorder.   *Nat   Genet*   **51**,   63-75   (2019).
https://doi.org/10.1038/s41588-018-0269-7
162 Duncan, L. *et al.* Significant Locus and Metabolic Genetic Correlations Revealed
in Genome-Wide Association Study of Anorexia Nervosa. *Am J Psychiatry* **174**,
850-858 (2017). https://doi.org/10.1176/appi.ajp.2017.16121402
163 Walters, R. K. *et al.* Transancestral GWAS of alcohol dependence reveals
common genetic underpinnings with psychiatric disorders. *Nat Neurosci* **21**,
1656-1669 (2018). https://doi.org/10.1038/s41593-018-0275-1
164 Mullins, N. *et al.* Genome-wide association study of more than 40,000 bipolar
disorder cases provides new insights into the underlying biology. *Nat Genet* **53**,
817-829 (2021). https://doi.org/10.1038/s41588-021-00857-4
165 Johnson, E. C. *et al.* A large-scale genome-wide association study meta-analysis
of   cannabis   use   disorder.   *Lancet   Psychiatry*   **7**,   1032-1045   (2020).
https://doi.org/10.1016/S2215-0366(20)30339-4
166 Howard, D. M. *et al.* Genome-wide meta-analysis of depression identifies 102
independent variants and highlights the importance of the prefrontal brain

regions. *Nat Neurosci* **22**, 343-352 (2019). https://doi.org/10.1038/s41593-018-
0326-7

167   International Obsessive Compulsive Disorder Foundation Genetics, C. & Studies,
O. C. D. C. G. A. Revealing the complex genetic architecture of obsessive-
compulsive disorder using meta-analysis. *Mol Psychiatry* **23**, 1181-1188 (2018).
https://doi.org/10.1038/mp.2017.154

168   Nievergelt, C. M. *et al.* International meta-analysis of PTSD genome-wide
association studies identifies sex- and ancestry-specific genetic risk loci. *Nat*
Commun **10**, 4558 (2019). https://doi.org/10.1038/s41467-019-12576-w

169   Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology
in schizophrenia. *Nature* **604**, 502-508 (2022). https://doi.org/10.1038/s41586-
022-04434-5

170   Cross-Disorder Group of the Psychiatric Genomics Consortium. Genomic
Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric
Disorders. *Cell* **179**, 1469-1482 e1411 (2019).
https://doi.org/10.1016/j.cell.2019.11.020

171   Marioni, R. E. *et al.* GWAS on family history of Alzheimer's disease. *Translational*
psychiatry **8**, 99 (2018). https://doi.org/10.1038/s41398-018-0150-6

172   Nalls, M. A. *et al.* Identification of novel risk loci, causal insights, and heritable
risk for Parkinson's disease: a meta-analysis of genome-wide association
studies. *The Lancet. Neurology* **18**, 1091-1102 (2019).
https://doi.org/10.1016/S1474-4422(19)30320-5

173   Nicolas, A. *et al.* Genome-wide Analyses Identify KIF5A as a Novel ALS Gene.
Neuron **97**, 1268-1283 e1266 (2018).
https://doi.org/10.1016/j.neuron.2018.02.027

174   Yu, D. *et al.* Interrogating the Genetic Determinants of Tourette's Syndrome and
Other Tic Disorders Through Genome-Wide Association Studies. *Am J*
Psychiatry **176**, 217-227 (2019). https://doi.org/10.1176/appi.ajp.2018.18070857

175   Hautakangas, H. *et al.* Genome-wide analysis of 102,084 migraine cases
identifies 123 risk loci and subtype-specific risk alleles. *Nat Genet* **54**, 152-160
(2022). https://doi.org/10.1038/s41588-021-00990-0

176   Johnston, K. J. A. *et al.* Genome-wide association study of multisite chronic pain
in UK Biobank. *PLoS genetics* **15**, e1008164 (2019).
https://doi.org/10.1371/journal.pgen.1008164

177   Lo, M. T. *et al.* Genome-wide analyses for personality traits identify six genomic
loci and show correlations with psychiatric disorders. *Nat Genet* **49**, 152-156
(2017). https://doi.org/10.1038/ng.3736

178   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for
comparing biological themes among gene clusters. *OMICS* **16**, 284-287 (2012).
https://doi.org/10.1089/omi.2011.0118

179   Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional
mapping and annotation of genetic associations with FUMA. *Nat Commun* **8**,
1826 (2017). https://doi.org/10.1038/s41467-017-01261-5

180   Wang, J. & Liao, Y. *WebGestaltR: Gene Set Analysis Toolkit WebGestaltR. R*
package version 0.4.3., <https://CRAN.R-project.org/package=WebGestaltR>
(2020).

181 Darmanis, S. *et al.* A survey of human brain transcriptome diversity at the single
cell level. *Proc Natl Acad Sci U S A* **112**, 7285-7290 (2015).
https://doi.org/10.1073/pnas.1507125112
182 Lake, B. B. *et al.* Neuronal subtypes and diversity revealed by single-nucleus
RNA sequencing of the human brain. *Science* **352**, 1586-1590 (2016).
https://doi.org/10.1126/science.aaf1204
183 Schilder, B. M. & Skene, N. G. orthogene: an R package for easy mapping of
orthologous genes across hundreds of species. *Bioconductor* (2022).
https://doi.org/https://doi.org/doi:10.18129/B9.bioc.orthogene.
184 Capps, M. E. S. *et al.* Disrupted diencephalon development and neuropeptidergic
pathways in zebrafish with autism-risk mutations. *Proc Natl Acad Sci U S A* **122**,
e2402557122 (2025). https://doi.org/10.1073/pnas.2402557122
185 Calhoun, C. C. S. *et al.* Removal of developmentally regulated microexons has a
minimal impact on larval zebrafish brain morphology and function. *bioRxiv*,
2024.2008.2019.608697 (2024). https://doi.org/10.1101/2024.08.19.608697
186 Kroll, F. *et al.* A simple and effective F0 knockout method for rapid screening of
behaviour and other complex phenotypes. *Elife* **10** (2021).
https://doi.org/10.7554/eLife.59683
179 Chaudhry, A., Shi, R. & Luciani, D. S. A pipeline for multidimensional confocal
analysis of mitochondrial morphology, function, and dynamics in pancreatic β-
cells. *Am J Physiol Endocrinol Metab* **318**, E87–E101 (2020).

TABLES

Table 1. Disorder and behavioral associations of top convergent up and down-regulated genes by cell-type from MalaCards, OMIM, and GWAS catalogue.

Cell-type	Top Meta Gene	Meta P	Z-score	Rare Disorders *MalaCards*	GWAS and behavioral associations *GWAS Catalog*
iNPCs	AGAP4	1.2e-14	7.71		
	CYTL1	7.8e-9	-5.77	Muscular Trophy (AD)	T1D, UC, CD, T2D, psoriasis, celiac, autoimmune disease, thyroid disease, ankylosing spondylitis
immature iGLUTs	MBD2	1.2e-15	8.01	Cerebellar Ataxia, Deafness, Narcolepsy (AD), Breast Cancer	Allergic disease, psoriasis, neuroticism, hoarding disorder, executive function measurement, memory function
	SHOX	2.9e-13	-7.30	Turner Syndrome, Dysplasia	
mature iGLUTs	MAP3K14	6.2e-36	12.52	Immunodeficiency (AR; X-linked), Ectodermal Dysplasia, Noonan Syndrome	PD, neuroticism, neuroimaging, unipolar depression, mood disorder, anxiety, cognitive function, MS, asthma, allergic disease,
	JMY	2.4e-44	-13.97	Galloway Mowat Syndrome (AR; X-linked)	T2D
mature iGABAs	GPR83	1.3e-17	-8.54		testosterone measurement, free androgen index, age at menarche, anxiety like-behaviors
	UBE2D4	3.9e-21	9.44	Brachydactyly (AD)	
Autosomal dominant (AD), Autosomal recessive (AR), Parkinson's disease (PD), multiple sclerosis (MS), ulcerative colitis (UC), type-I diabetes (T1D), type-2 diabetes (T2D), Crohn's disease (CD)					

Table 2. Disorder and behavioral associations of top nodes by cell-type from MalaCards, OMIM, and GWAS catalogue.

Cell-type	Top protein-coding nodes across all networks	Rare disorders *MalaCards*	GWAS and behavioral associations *GWAS Catalog*
iNPCs	KIAA2012		ADHD/conduct disorder (rs1521882), educational attainment (rs12623702, rs4675248, rs2160317, rs2177083, rs34189321,rs58100125), migraine/T2D (rs6748072), amygdala volume (rs72936662)
immature iGLUTs	MYH15	Deafness (AR); de novo SCZ CNV	Social interaction measurement (rs13082569), cognitive function (rs3860537), unipolar depression (rs1531188), MDD (rs113689582), insomnia (rs62266174, rs6768511, rs6786515, rs6795280), BIP (rs1531188), ANX (rs4855559), educational attainment (rs115910830, rs2290601, rs3860537, rs60785803)
mature iGLUTs	GALNTL5	ASD, Spastic paraplegia 48 (AR)	
mature iGABAs	EREG	Epilepsy, Immune Deficiency Disease	ADHD (rs1350666), wellbeing measurement (rs112444088), learning & memory (pathway)
Autosomal recessive (AR), Anxiety disorder (ANX), Bipolar Disorder (BIP), Type-2 diabetes (T2D)			

Table 3. Convergent nodes that overlap with CNV and rare variant target genes for
each cell-type.

Celltype	Rare variant gene targets for ASD, SCZ, BIP, ID, and Epilepsy
iNPCs	AATK, DLC1, PAK6
immature iGLUTs	ACMSD, HCST, MYH15, PAH, RSPO1, SFTPC, SH3RF2, SLC28A2, SNAI2, ACOT6, CSPG4, PYCARD, SLC5A7, SULT1B1, TBXA2R, TEKT5 ,TEX15, ARG1, ASB14, CACNA1D, OPLAH, GRM4, KCNT1, FKBP6, NPAP1, OCA2, ATP10A
mature iGLUTs	S100G, TRIM50, FOLR1, COX7B2, KIF23
mature iGABAs	CHRND, RETN, PAX6, RIMBP3, SPDYE5, TSKS

Table 4. Disorder and behavioral associations of top nodes by cell-type and behavioral
set from MalaCards, OMIM, and GWAS catalogue.

Cell type		Top Node	Gene name	Meta P	Z-score	Rare Disorders	GWAS
Mature iGLUT		CNBP	Sterol-mediated transcriptional repression	2.9e-16	8.18	Myotonic Dystrophy, Neuromyotonia & axonal neuropathy (AR), Creutzfeldt-Jakob Disease, Deafness (AR), Fragile-X tremor/ataxia, ALS	
	1	ANKRD36		1.95e-18	8.8		
		FOXJ3		1.7e-17	8.5		
	2	ATP6V0C		5.3e-44	-13.9	ASD, Epilepsy, NDD	
		FOXJ3	Gene expression regulation/chromatin	1.9e-19	9.02		EA (rs35011283)
	3	SOX12	Implicated in cell fate decisions during development	1.6e-17	-8.52	Microphtalmia Syndromic 12 (neurological)	Memory performance (rs78532272)
	4	FOXJ3		7.9e-22	9.6		
		ANKRD36	Ankyrin Repeat Domain 36	6.6-e26	-10.5	Neuropathy, Giant Axonal Nonsyndromic Deafness (AR)	SCZ (rs9631085), neuroimaging (rs11692435, rs167684, +6)
Mature iGABA	1	UBE2D4	Ubiquitin-Conjugating Enzyme E2 D4	1.2e-09	6.1		
		PRKAG2	Protein Kinase AMP-Activated Non-Catalytic Subunit	6.7e-7	-4.97	Wolff-Parkinson-White Syndrome, Neuromuscular Disease, Specific Language Impairment, Microencephaly, Epilepsy	EA (rs1860735, rs2538046, rs4726070), ASD/SCZ (rs115136442), impulse control (rs2302532), BIP (rs7795096), memory (rs2536058)
	2	MYL3	BBB & immune cell trasnmigration	4e-12	6.9	Wolff-Parkinson-White Syndrome, Microencephaly, Epiliepsy	
		TNFSF11		3.6e-10	-6.3	ID, NDD, developmental & epileptic encephalopathy, IBS,	
	3	KIF1A	Axonal Transporter Of Synaptic Vesicles	8.8e-12	6.8	ID, ASD, neuropathy, Epilepsy, Spastic Ataxia, Cortical Dysplasia, Alacrima, Achalasia, and Impaired ID Syndrome	Insomnia (rs10196604,rs4455151)
		UNC5C	Netrin Receptor	3.98e-8	-5.5	Alzheimer Disease, Familial 1	Insomnia (rs371745379), wellbeing (rs116984513), OCD (rs17384439), cognition (rs3846455), unipolar depression (rs6822806)
	4	UBE2D4		8.5e-13	7.2		
		UBC		1.7e-10	-6.4	Parkinson's disease, epilepsy, muscular dystrophy	Sleep duration (rs10773112), AD (rs78184510),

1935

Figure 1. Knock-out (KO) effects of 21 NDD risk genes are most strongly correlated in mature neurons. (A) List of rare-variant target risk genes associated neurodevelopmental disorders (NDD) separated by chromatin modifiers and neuronal communication genes. Bold gene names indicate strong assocciations with ASD based on Fu et al. 2022. Gene targets of rare variants associated with schizophrenia (SCZ), epilepsy (EPI) and bipolar disorder (BIP) are annotated. **(B)** Strength of association with ASD, as estimated by distribution of posterior probability (p.p.) scores from Fu et al. 2022; 4 out of 29 NDD genes were more strongly associated with developmental delay (DD) (blue; p.p.<=0.1) while 16 out of 29 were more strongly associated with ASD (red; p.p.>=0.9). Further annotation of individual risk genes are shown in **SI Figures 1-2**. **(C)** MAGMA enrichments of targeted genes across GWAS for anorexia nervosa (AN), chronic pain, amyotrophic lateral sclerosis (ALS), SCZ, and BIP, BIP-I (bipolar subtype 1), and BIP-II (bipolar subtype 2). #nominal p-value<0.05, *FDR<0.05, **FDR<0.01, *FDR<0.001 **(D)** Schematic of hiPSC-derived cell-type specific scCRISPR-KO screen. Representative immunofluorescence for markers of NPCs (DAPI/Nestin), mature iGLUTs (DAPI/MAP2/vGLUT), and mature iGABAs (DAPI/MAP2/GABA). **(E)** Transcriptomic impact of NDD gene KO represented as the number of nominally significant (p<0.01) differentially expressed genes (DEGs). **(i)** Pearson's correlation

1954   matrix of log2FC DEGs across all NDDs and cell-types. **(ii)** Cross cell-type correlation
1955   network diagram across NDD perturbations (number of NDD gene knockout (KO)
1956   perturbations resolved indicated in parentheses); the mature iGLUT cluster was most
1957   dense, and the iNPC most sparse.

1958

1959

Figure 2. Gene-level convergence is greatest in mature glutamatergic neurons. In total, nine NDD genes showed evidence of knockout across all four cell types: *ARID1B, ASH1L, CHD2, MED13L, NRXN1, PHF21A, SETD5, SIN3A, SMARCC2*. For these nine, "convergent genes" are defined as those differentially expressed genes (DEGs) with significant and shared direction of effect across all NDD gene perturbations. **(A)** Schematic explaining cell-type specific convergence at the individual gene level via differential gene expression meta-analysis (FDR adjusted pmeta<0.05, Cochran's heterogeneity Q-test pHet > 0.05). **(B)** Convergence across 9 NDD genes is unique to each cell type, using rank-rank hypergeometric (RRHO) test to explore correlation of convergent genes shared across 9 NDD perturbations (RRHO score = -log10*direction of effect) between cell-types. The top right quadrant represents down-regulated genes (meta-analysis z-score >0) for the y-axis and x-axis cell-type. The bottom left quadrant represents up-regulated convergent genes (meta-analysis z-score <0) for the y-axis and x-axis cell-type. Significance is represented by color, with red regions representing significantly convergent gene expression. **(C) (i)** The average strength of convergence, measured as the ratio of convergent genes to the average number of DEGs across all 152 unique combinations of 2-5 genes from the nine NDD genes, was highest in iGLUTs. **(ii)** The magnitude of convergence between the same NDDs tested in different cell types was highly correlated (Pearson's correlation, P_{holm}<2.2e-16); with the strongest relationship between immature and mature iGLUTs. **(D)** Venn diagram representing the absolute overlap (regardless of direction of dysregulation) of cell-type

specific convergent genes shared across 9 NDDs. **(E) (i)** MAGMA enrichment –log10(p-value) of cell-type-specific (color of points) convergence and GWAS-risk associated genes with significance after multiple testing correction indicated as follows: [#]unadjusted p-value=<0.05, *FDR<=0.05, **FDR<0.01, ***FDR<0.001. The direction of the triangles indicates a positive (upwards triangle) or negative (downwards triangle) enrichment beta. **(ii)** Over-representation analysis (ORA) enrichment ratios of cell-type-specific (color of bars) convergence and rare variant target genes. Significance after multiple testing correction indicated as follows: [#]unadjusted p-value=<0.05, *FDR<=0.05, **FDR<0.01, ***FDR<0.001. **(F)** Gene set enrichment analysis (GSEA) identified downstream pathways involved in neural proliferation, neurite outgrowth, synaptic vesicle transport, and mitochondrial function as cell-type specific targets of convergent genes across 9 NDDs. Results were filtered for pathways with nominal p-values <0.05. Normalized GSEA enrichment scores represent the direction of enrichment based on the meta-analyzed Z-score for each convergent gene. Cell-type is represented by shape and the size of each point represents the –log10(FDR).

1997

Figure 3. Network-level convergence resolves cell-type-specific and developmental-specific node genes. "Convergent networks" are co-expressed genes that share similar expression patterns across NDD gene perturbations, here resolved for the nine NDD knockouts resolved across all four cell types: *ARID1B, ASH1L, CHD2, MED13L, NRXN1, PHF21A, SETD5, SIN3A, SMARCC2*. **(A)** Schematic explaining cell-type specific convergence at the network level using Bayesian bi-clustering and unsupervised network reconstruction. **(B)** Strength of network convergence across all random combinations of 9 NDD KO perturbations by cell-type. **(i)** The mean strength of network convergence is significantly different by cell-type, with the highest convergence present in immature iGLUTs. The same KO combinations tested in one cell type may not resolve convergence in another cell type. Each point represents a resolved network, and its calculated convergence strength. Dots that represent the same combinations of KO perturbations, but tested in each cell type, are connected by a line. **(C)** Convergent network strength was most correlated between mature iGLUTs and iGABAs (Pearson's Correlation Coefficient (PCC) = 0.6, P_{Holm} <2.2e-16). Convergent network strength in iNPCs was not correlated with network strength in neurons. **(D)** Venn diagrams of the

total number of unique node genes within convergent networks for each cell-type. The
lack of overlapping node genes between cell types **(D),** as well as the weak correlations
of convergence strength between immature and mature cell-types **(C),** suggest greater
cell-type specificity in the magnitude of network-level convergence compared to gene-
level convergence. **(E)** Enrichment ratios from over-representation analysis (ORA) of
cell-type specific (color of bars) convergent node genes for rare variant targets.
([#]unadjusted p-value=<0.05, *FDR<=0.05, **FDR<0.01, ***FDR<0.001). **(F, G)**
Representative cell-type specific network plots for convergence across 15 genes
(*ARID1B, ASH1L, ASXL3, BCL11A, KDM5B, CHD2, MBD5, MED13L, NRXN1, PHF12,*
PHF21A, SETD5, SIN3A, SKI, SMARRC2) from **(F)** iNPCs and **(G)** mature iGLUTs.
Network genes were filtered for protein-coding genes, clustered, and annotated based
on the primary node gene for each cluster. Gene set enrichment analysis of the
networks identified unique functions by cell type. Convergent networks in iNPCs were
enriched for pathways associated with neurogenesis (e.g., cell cycle, cell division, EPO
signaling), while in mature iGLUTs for pathways associated with synaptic function
(transmembrane transport and receptor signaling, secretory vesicles, SNARE complex).

2030
2031 **Figure 4. Functional similarity and brain co-expression between NDD genes**
2032 **predict gene-level and network-level convergence, with unique influences by cell-**
2033 **type. (A)** Schematic for training random forest models for gene and network-level
2034 convergence with external validation in a SCZ CRISPRa screen. **(B)** Predictor variables
2035 included in the model include scores of functional similarity, dorsolateral prefrontal
2036 cortex (DLPFC) brain co-expression, cell-type, and the number of KOs. (**B.P score** =
2037 semantic similarity of GO: Biological Process membership between KO genes; **C.C.**
2038 **score** = semantic similarity of GO: Cellular Component membership between KO
2039 genes; **M.F score** = semantic similarity of GO: Molecular functions membership
2040 between KO genes; **B.E.C** = dorsolateral prefrontal cortex expression correlations
2041 between KO genes; **nKOs** = number of KO genes tested for convergence). **(C)**
2042 Pearson's correlations of predictor variables and gene-level and network-level

convergence (PBonferroni<=0.01**, PBonferroni<=0.01***). **(D)** Functional similarity, brain co-expression, cell-type, and the number of KOs assayed strongly predicted gene-level convergence (97% variance explained by the model; mean of squared residuals=0.02) and moderately predicted network-level convergence (53% variance explained; mean of squared residuals=0.73). **(i-ii)** Importance of each of the predictor variables was assessed by two metrics: the percent mean increase in squared residuals (%IncMSE) and the increase in node purity. In the model – number of KO genes in a set is the most important predictor of convergence based on %Inc MSE, but not node purity. However, the impact of nKOs on gene-level convergence is much stronger – likely an artifact of the method used for measuring convergence. For network level convergence, each variable has a IncMSE between 20-30%. **(E)** Internal evaluation of the model using 30% of the original data resulted in high concordance between convergence predicted by the model and the measured convergence. Predicted gene-level **(i)** [gene-level convergence: n=19,823; Pearson's R=0.984; p<2.2e-16; root mean squared error (RMSE) =0.15] and network-level **(ii)** convergence [network-level convergence: n=962; rho=0.722; p<2.2e-16; RMSE=0.85)] by the model strongly correlated with the measured convergence in the testing sets. Correlation of predicted vs. accrual convergence values are color-coded by cell-type with corresponding color-coded correlations and p-values listed in the upper right corners of the scatterplots. **(F)** External validation in an independent scCRISPRa screen of SCZ target genes predicted showed moderate, but significant, correlation between convergence predicted by the model and the measured convergence. **(i)** gene-level (n=1013, R=0.14, p=1.1e-05, RMSE=1.748) and **(ii)** network-level convergence (n=826, R=0.26, p=2.9e-14, RMSE=0.68).

2067
2068 **Figure 5. LNCTP predicts effects of convergent genes *in silico*.** (**A**) LNCTP

imputation and perturbation model: an energy-based network model is trained to impute bulk and cell-type specific expression data in the prefrontal cortex over a population of post-mortem individuals from PsychENCODE using a panel of 1325 genes and embedded cell-type specific Gene Regulatory Networks (GRNs) (*LNCTP in silico model*); a chosen gene is then perturbed by fixing its expression, and the effects on other genes are predicted by the model; *in silico* category-specific convergent genes are then identified by analyzing the fold-changes across subjects (*LNCTP Simulating Perturbations*). **(B)** Predicted *in silico* log fold-changes for the *in vitro* positive and negative convergent genes across the 29 CRISPR perturbations, in Bulk, Excitatory and Inhibitory neuron networks (*LNCTP Simulating Perturbations*, 2-tailed t-test p-values shown). **(C)** Proportion of genes showing same direction fold-changes in *in silico* and *in vitro* perturbations across classes of perturbation and cell-type (left), and the intersection of convergent in silico genes across classes of perturbation (*LNCTP in silico convergent genes*, synaptic-epigenetic genes reduced and ASD-DD genes enriched, p<1e-3, 2-tailed hypergeometric test). **(D)** Venn diagram of *in silico* convergent genes across all categories by clinical (ASD vs DD) or functional (synaptic vs epigenetic) annotation. **(E)** Number of terms enriched for convergent genes across all categories for 102 in silico perturbations. **(F)** Semantic distance of pairs of enriched terms within or between sets determined by synaptic and epigenetic convergent gene rankings (*LNCTP semantic distance test*, 2-tailed Mann Whitney test) (G) Percent of concordant genes in each perturbation and ontology category within the leading-edge enriched genes (*LNCTP in silico convergent genes*).

Figure 6. NDD knock-outs converge on mitochondrial function. (A) Gene set
enrichment analysis (GSEA) identified downstream pathways involved in neurogenesis,
neurite outgrowth, synaptic biology, and mitochondrial function as cell-type specific
targets of convergent genes across 15 NDD KOs **(*ARID1B, ASH1L, ASXL3, BCL11A,
KDM5B, CHD2, MBD5, MED13L, NRXN1, PHF12, PHF21A, SETD5, SIN3A, SKI,
SMARRC2)*** in iNPCs and mature iGLUTs. Results were filtered for pathways with

nominal p-values <0.05. Normalized GSEA enrichment scores represent the direction of
enrichment based on the meta-analyzed Z-score for each convergent gene. Cell-type is
represented by shape and the size of each point represents the –log10(FDR). **(B)**
Summary of network and gene-level pathway enrichments (from **Fig. 2-3)** for shared
effects of nine and fifteen NDD KOs in iNPCs and mature iGLUTs. **(C)** Proliferation
assessment of NPCs using Ki-67 median fluorescence intensity (MFI) measured with
flow cytometry, wildtype (purple: no iCas9 induction) versus knockout (green:
doxycycline to induce iCas9). 4-6 replicates per condition, unpaired t-test with Welch
correction; p-values corrected for multiple comparisons using FDR. **(D)** Scatter plot of
gRNA $\log_2$ fold-change (high- (PE-high) and low- (FITC-high) Δψm-sensitive dye JC-1
membrane-potential fractions) in NPCs (x-axis) and mature iGLUT neurons (y-axis),
with points colored by enrichment category (shared NPC and iGLUT in red; distinct
between NPC and iGLUT in blue). Right: Bar chart of $-\log_{10}$(FDR) for over-represented
gene sets in the tene gene KOs enriched in both lineages. **(E)** (i) High resolution, high-
throughput microscopy of mitochondrial morphology (scale bar 10 μm): an isolated
dendrite labelled with a dendritic marker (MAP2), mitochondrial marker (TOMM20) and
marker of the OXPHOS complex (Total OXPHOS) (scale bar 5 μm). **(ii)** Effect of
ARID1B-KO on mitochondrial sphericity and branch length independent of changes in
mitochondrial volume and surface area **(SI Fig. 20-21). (iii)** Effect of ARID1B-KO on
average fluorescence intensity of OXPHOS proteins. Each datapoint indicates one well
of a 96-well, representing hundreds of μm^2 of neuronal area and tens of thousands of
individual mitochondria (*adjusted p<0.05, ** adjusted p<0.01). **(F)** Effect of NRXN1-KO
on maximal respiration and coupled respiration in iGLUTs. Oligo: oligomycin; FCCP:
carbonyl cyanide 4-(trifluoromethoxy) phenylhydrazone; R+A: rotenone and antimycin
A. Data are presented as mean ± SEM. Statistical analysis was performed using one-
way ANOVA. *p<0.05. Each datapoint represents one well of a 24-well Seahorse assay
plate. The experiment was independently replicated twice.

Figure 7. NDD gene mutants with shared behavioral phenotypes in zebrafish resolve unique and cell-type-specific gene-level convergent signatures and are rescued by predicted medications. (A) NDD risk genes uniquely cluster based on sleep-wake/visual-startle behavioral responses in zebrafish mutants. set 1: *nrxn1a, mbd5, kdm5bab;* set 2: *phf12ab, skiab, chd2, smarcc2;* set 3: *kdm6bab, kmt5b, kmt2cab;* set 4: *wacab, arid1b, phf21aab, chd8, ash1l.* **(B)** Gene expression in human mature iGLUTs and iGABAs correlate with expression in the zebrafish brain. Cellular deconvolution of wild-type larval zebrafish brain expression based on adult human single-cell brain reference identifying neurons as the largest proportion of cells in the

fish brain. Gene expression in wild-type zebrafish brain significantly positively correlates with gene expression of mature iGLUTs (rho=0.39, Holm's adj.P<0.001) and iGABAs (rho=0.39, Holm's adj.P <0.001). **(C)** For each of the four behaviorally defined sets, gene-level convergence (DEGs with significant and shared direction of effect across all NDD genes within each of the four sets (FDR adjusted p_{meta}<0.05, Cochran's heterogeneity Q-test p_{Het} > 0.05)) is largely non-overlapping between mature iGLUTs and iGABAs, with unique enrichments for common psychiatric risk gene targets. Number of convergent genes that are up (+) or down (-) regulated for each NDD set are indicated. **(D)** In both iGABAs and iGLUTs, all four behavioral sets were enriched for FMRP targets. Gene targets of neurodevelopmental rare variants were only significantly enriched for convergent signatures in mature iGLUTs; behavioral set 4 uniquely significantly enriched for secondary targets of ASD loss-of-function variant and set 3 uniquely enriched for primary targets of SCZ non-synonymous variants. **(E)** In iGLUTs, NDD related behaviors were only enriched in sets 1 and 3, with enrichments for language, speech, and intellectual delays in sets 1,3 and 4. All sets were enriched for seizure and hypertonia. **(F)** Potential "rescue" drugs for these 4 phenotypic groups were selected from enrichment scores using cMAP and filtered for drugs included in a screen of 376 compounds for behavioral effects in zebrafish. Top candidates that were significantly negatively enriched for iGLUT convergence from cMAP and negatively correlated with mutant behavioral features were tested in mutant lines representative of sets 2-4. n.p. indicates that the drug repaglinide was not present in the cMAP dataset. Mutant-x-Drug combinations were as follows: $chd2^{\Delta7/\Delta7}$-x-pravastatin; $kdm6bab$ F0-x-paclitaxel; $kdm6bab$ F0-x-sirolimus; $kmt5b^{\Delta208,1i,\ \Delta5/\Delta208,1i,\ \Delta5}$-x-paclitaxel; $kmt5b^{\Delta208,1i,\ \Delta5/\Delta208,1i,\ \Delta5}$-x-sirolimus; ; $ash1l^{1i,\ \Delta60,19i/\ 1i,\ \Delta60,19i}$-x-ezetimibe; $ash1l^{1i,\ \Delta60,19i/\ 1i,\ \Delta60,19i}$-x-repaglinide; $ash1l^{1i,\ \Delta60,19i/\ 1i,\ \Delta60,19i}$-x-rosuvastatin; $ash1l^{1i,\ \Delta60,19i/\ 1i,\ \Delta60,19i}$-x-sunitinib; $phf21aab$ F0-x-amiodarone; $phf21aab$ F0-x-fluvoxamine. **(G)** For behaviors that were significantly different between mutant+DMSO and WT+DMSO (p<0.05), we characterized the effect of the mutant-x-Drug on behavior as either (a) *exacerbated* [sig. effect mutant+Drug-v-WT > sig. effect mutant-v-WT], (b) *unchanged* [sig. effect mutant+drug-v-WT = sig. effect mutant-v-WT], (c) *partial rescue* [effect mutant+Drug-v-WT < effect mutant-v-WT], (d) *rescued* [sig. effect mutant-v-WT, no sig. effect mutant+Drug-v-WT], (e) *over-corrected* [mutant+Drug-v-WT opposite direction of sig. effect mutant-v-WT]. All drugs reversed at least one dysregulated behavior except for sirolimus in *kmt5b*. **(i)** Comparison of the magnitude of effect (beta) on behavior between the mutant+DMSO compared to mutant+Drug groups shows rescue of select behavioral features in *kdm6b* and *chd2* mutants by paclitaxel (Shapiro Wilk's Normality p= , Student T statistic=-3.533, p=0.0017788, df=23) and pravastatin (Student T statistic=-3.533, p=0.0017788, df=23), respectively. **(ii)** the *phf21a* mutant phenotype was strongly opposed by fluvoxamine (Pearson's correlation=-0.58, p=0.0028).