Title:

Declining juvenile survival of Adélie penguins in Antarctica

Authors:

Téo Barracho[*], Gaël Bardon, Rémi Choquet, Aymeric Houstin, Alexander Winterl, Michaël Beaulieu, Pierrick Blanchard, Cindy C. Cornet, Robin Cristofari, Thierry Raclot, Sebastian Richter, Nicolas Chatelain, Julien Courtecuisse, Matthieu Brucker, Daniel P. Zitterbart, Nicolas Lecomte[*†] and Céline Le Bohec[*†]

[†] Co-last authors

* tbarracho@centrescientifique.mc

* nicolas.lecomte@umoncton.ca

* celine.le-bohec@cnrs.fr

Affiliations :

TB : Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France ; Canada Research Chair in Polar and Boreal Ecology and Centre d'Études Nordiques, Department of Biology, University of Moncton, Moncton, NB, Canada ; CEFE, Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France ; Centre Scientifique de Monaco, Département de Biologie Polaire, Monaco, Principality of Monaco

GB : Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France ; Centre Scientifique de Monaco, Département de Biologie Polaire, Monaco, Principality of Monaco

RCH : CEFE, Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France

AH : Woods Hole Oceanographic Institution, Applied Ocean Physics and Engineering Department, Woods Hole, Massachusetts, USA ; Centre Scientifique de Monaco, Département de Biologie Polaire, Monaco, Principality of Monaco

AW : Friedrich-Alexander-University Erlangen-Nürnberg, Department of Physics, Erlangen, Germany

MBE : Wild Animal Initiative 5123 W 98th St, 1204 Minneapolis, MN 55437, USA

PB : Centre de Recherche sur la Biodiversité et l'Environnement (CRBE), Université de Toulouse, CNRS, IRD, Toulouse INP, Université Toulouse 3 – Paul Sabatier (UT3), Toulouse, France

CCC : Centre for Blue Governance, Faculty of Business and Law, University of Portsmouth, UK ; Centre Scientifique de Monaco, Département de Biologie Polaire, Monaco, Principality of Monaco

RCR : Institute of Biotechnology, University of Helsinki, Helsinki, Finland

TR : Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

SR : Friedrich-Alexander-University Erlangen-Nürnberg, Department of Physics, Erlangen, Germany

NC : Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

JC : Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

MBR : Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

DPZ : Woods Hole Oceanographic Institution, Applied Ocean Physics and Engineering Department, Woods Hole, Massachusetts, USA ; Friedrich-Alexander-University Erlangen-Nürnberg, Department of Physics, Erlangen, Germany

NL : Canada Research Chair in Polar and Boreal Ecology and Centre d'Études Nordiques, Department of Biology, University of Moncton, Moncton, NB, Canada

CLB : CEFE, Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France ; Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France ; Centre Scientifique de Monaco, Département de Biologie Polaire, Monaco, Principality of Monaco

Abstract:

As summer sea ice around Antarctica reaches modern lows, quantifying the demographic response of polar species to such environmental changes becomes critical. To achieve this, synthesizing results across species' ranges and elucidating the environmental factors driving population dynamics are key. Adélie penguins are considered reliable indicators of changes in Antarctica but the processes through which sea ice and other environmental factors shape their population dynamics are still unclear, especially for critical age groups such as juveniles. Using a 17-year dataset of Adélie penguins electronically tagged in Adélie Land (Antarctica), we found that juvenile survival probability was most impacted by sea ice concentration near their natal colony right after fledging, with lower ice concentrations detrimental to survival. Importantly, we found that juvenile survival declined by 32% from 2007 to 2020, mirroring trends at other distant colonies. The emergence of similar patterns at opposite ends of the continent may be an early signal for shifts in population trends expected from climate change.

Introduction:

Antarctica is at the center of large and rapid environmental shifts, with both accelerating climate changes[1–3] and increasing exploitation pressure by fisheries[4]. Quantifying the response of animal populations to such environmental forcing is essential to determine their potential dynamics and persistence[5,6]. Yet, our understanding of how Antarctic species respond to such changes is still limited. Typically, the spatially and temporally heterogeneous dynamics of climate[7,8] and of environmental parameters such as sea ice[9] across species distributions makes identifying drivers of any demographic rate challenging[10–12]. This hampers our ability to anticipate species response to environmental changes, especially over scales relevant to ecosystem management[13,14].

The Adélie penguin (*Pygoscelis adeliae*), an iconic Antarctic species, exemplifies this challenge particularly well. Early observations of contrasting population trends between the Ross Sea and the Antarctic Peninsula in response to sea ice declines have fostered a population dynamics model, where abundance is maximal at intermediate sea ice concentrations[15,16]. This model has been instrumental in projecting the response of Adélie penguins to climate variability across large spatial and temporal scales[17–19]. However, more localized and shorter-term predictions are still hindered by the inherent variability and noise in abundance time series, which most population studies are constrained to rely on[19–21]. Precise quantification of

key demographic rates (e.g. juvenile survival) is critical for elucidating population dynamics driven by complex environmental interactions[22,23]. The need for such an approach is twofold; it can provide short-term feedback for ecological management at the local scale and be integrated at larger spatial scales as more studies become available across the species range[24–27].

In long-lived species, adult survival typically holds the greatest potential to influence population dynamics, but actual population growth rates can often be driven by other demographic rates such as fecundity or juvenile survival[28]. In long-lived Adélie penguins, juvenile survival indeed appears to play a prominent role in driving population fluctuations. For example, Adélie penguin's abundance (as measured by the number of breeding pairs) correlates with sea ice conditions with a 5-years lag[29]; which is consistent with effects of sea ice on juvenile survival during their first months at sea[30]. In the Antarctic Peninsula[31,32] and East Antarctica[33], population size decreased when juvenile survival also did, further emphasizing its central role in shaping overall population trends. Despite such findings, we still lack clear evidence linking juvenile survival to environmental parameters known to affect population dynamics, such as sea ice concentration[13]. Although adult survival has been convincingly linked to environmental variables (e.g., Southern Oscillation Index, winter sea ice concentration) in several populations around Antarctica[31,34–37], the drivers of juvenile survival remain unclear[31,33,36]. Given the importance of this demographic parameter, evaluating its trends across the species' range and elucidating its environmental drivers are required to adequately predict the species' response to future environmental changes[38].

Here, we quantify juvenile survival probabilities for the first time in the Western Pacific Ocean sector (90°E to 160°E, Fig. S1) using a 17-year-long dataset of known-age, electronically tagged Adélie penguins from Pointe Géologie archipelago, Adélie Land. By investigating trends in survival for this population located thousands of kilometers away from other studied colonies and synthesizing all previously published studies on this vital rate, we provide a comparative picture of trends in Adélie penguin juvenile survival across Antarctica. To uncover the mechanisms behind variations in juvenile survival, we also investigate the drivers of juvenile survival at Pointe Géologie by linking survival probabilities to intrinsic and environmental parameters such as body mass at fledging and sea ice concentrations during their first winter at sea.

Results:

We used a multi-state capture-recapture framework accounting for short-range dispersion within a 10-km radius to model the survival probability of juvenile Adélie penguins (0-2 years old). Based on 17 years of tagging and detection data of 2807 individuals (2007-2023, see methods and Table S4), this model enabled us to quantify juvenile survival probabilities over 14 years (2007-2020). Over this period, juvenile survival estimates averaged 0.42 ± 0.18 and showed high inter-annual variability (CV = 44%, Fig. 2A). Notably, juvenile survival probabilities exhibited a negative temporal trend (Fig. 2A; Table 1, model 36, analysis of deviance: p-ANODEV = 0.016). The annual rate of change was of -2.5% ($\pm$ 95%CI: -3.1, -2.0), translating into a 32% loss between 2007 and 2020. Juvenile survival has previously been quantified at 5 other locations around Antarctica[31,33,35,39,40] (summarized in Fig. 2). Temporal variation was investigated at only two of them, and juvenile survival probabilities declined at both (Admiralty Bay, King George Island, Western Antarctic Peninsula, -1.3% per year for 1982-2000 [31]; Béchervaise Island, Mac. Robertson Land, East Antarctica, -1.8% per year for 1992-2015 [33]; Fig. 2).

To understand the drivers behind the decline observed at our study site, we used our previously developed model to test the effect of various intrinsic (mass at fledging, cohort size) and environmental covariates such as sea ice concentrations (SIC), windchill temperatures, and global climatic indices (Southern Oscillation Index, Southern Annular Mode, Table S2) on juvenile survival probabilities. Local and regional variables (Table S2) explained more of the variation in juvenile survival rates than global indices of climate variability (Table 1). Autumn sea ice concentrations (SIC.autumn) within 200 km of the natal colony (where juveniles are expected to be following fledging) correlated positively with juvenile survival rates (Fig. 3A, Table 1, model 29, %DEV = 28.6, slope $\pm$ 95%CI = 0.95 (0.94, 0.95)). We also found a quadratic effect of winter windchill temperatures (Table 1, model 26, %DEV = 24.7), with warmer temperatures associated with higher juvenile survival.

To better understand how SIC could affect juvenile survival at the onset of winter, we decomposed SIC.autumn into monthly data (SIC.march and SIC.april), and only SIC.april explained more than 20% DEV (Table 1, model 31, %DEV = 34.8). By decomposing SIC.april into the fraction of total ice area (15-100% SIC) covered by the marginal ice zone (15-80% SIC, MIZFRAC.april) and the fraction of total ice area covered by dense sea ice (80-100% SIC,

DENSEFRAC.april), we then found that the amount of loose sea ice in the vicinity of the colony (MIZFRAC.april) was negatively related to juvenile survival rates (slope ± 95%CI = - 0.96 (- 0.98, - 0.93), Fig. 3B) and accounted for 71.4% of the temporal variation (Table 1, model 33). Conversely, the amount of dense sea ice (DENSEFRAC.april) was positively related to juvenile survival rates (slope ± 95%CI = 0.83 (0.82, 0.84), Fig. 3C) and accounted for 33.8% of the temporal variation (Table 1, model 32). This amount of April dense sea ice near the colony declined by 13% between 2007 and 2020 (linear model: F = 7.65, p = 0.016, R^2 = 0.37, Fig. 3D).

Chick body mass at tagging (fledging mass), an important driver of post-fledging survival in birds, averaged 3.57 ± 0.62 kg and varied more within (average of annual mass SD = 0.54 kg) than across cohorts (SD of annual average mass = 0.33 kg, Fig. S8). At the individual level, our multi-state capture-recapture modeling showed that fledging mass was positively related to the probability of surviving the juvenile phase, with equal support for linear and quadratic relationships (Table 1, models 16 and 17, LRT: both p < 0.001). The quadratic model predicts that juvenile survival plateaus at ca. 4 kg body mass (Fig. 4), and this relationship did not differ among years (Table 1). At the population level however, inter annual variations in juvenile survival were poorly explained by the average mass of the cohort (mean annual fledging mass, MFM, Table 1, model 35, %DEV = 7.2, slope ± 95%CI = 0.06 (0.04, 0.07)).

Discussion:

Using 17 years of individual electronic monitoring (2007-2023), we observed a loss of 2.5% per year in Adélie penguin juvenile survival over 14 years (2007-2020) in an Adélie Land population. This decline echoes those already observed in the only two other populations for which juvenile survival time series are also available (in Mac. Robertson Land, East Antarctica, -1.8% per year[31], and in the Western Antarctic Peninsula, -1.3% per year[33]). By quantifying the survival of juvenile Adélie penguins for the first time in a region located thousands of kilometers away from other monitored colonies (3,400 km and 6,300 km, respectively), our results point to emerging negative trends across the continent. We also highlight potential mechanisms at play, since the decline in juvenile survival probability recorded in Adélie Land was associated with decreasing sea ice concentrations in the vicinity of the natal colony within two months after fledging. As sea ice declines are predicted to shrink Adélie penguin

populations in the latter part of the century[17,18], steady declines in juvenile survival may already foreshadow such shifts.

- *Range-wide declines in juvenile survival*

Hinke et al.[31] were the first to report an annual decline in juvenile survival probability, with a decline of -1.3% per year between 1982 and 2000 for Admiralty Bay (King George Island, Western Antarctic Peninsula). Survival probability at this site then seemed to stabilize up to 2011, although a change in marking methodology (from aluminum to stainless steel flipper bands) and high inter-annual variability in survival probabilities made it difficult to compare the two periods[31,32]. More recently, Emmerson and Southwell[33] extended the analysis of the capture-recapture dataset from Béchervaise Island (Mac. Robertson Land, East Antarctica[36]) to reveal that juvenile survival had also declined (-1.8% per year, 1992-2015) in this population located halfway between the Antarctic Peninsula and Adélie Land (Fig. 2).

With an annual decrease of -2.5% between 2007 and 2020 in Adélie Land, our study adds to the picture of declining juvenile survival across the species range (Fig. 2). Juvenile survival has now declined at all three locations where time trends have been investigated (Fig 2). The time series used here (14 years) is shorter than those for Admiralty Bay (18 years) and Béchervaise Island (24 years), but it is worth noting that the -2.5% annual decline at Pointe Géologie exceeds the average inter-annual variability in survival probabilities (1.79%) as in Béchervaise Island (1.80% vs. 1.57%) but unlike Admiralty Bay (1.30% vs. 2.90%).

Permanent dispersion of marked individuals away from study sites can bias the results of capture-recapture studies when incorrectly accounted for[41]. Here, our deployment of a mobile RFID antenna network, radiating 10 kilometers around the main study colony (Fig. 1), allowed a robust estimation of survival probability despite local dispersion. Long-distance permanent dispersion of juveniles, which could bias our survival estimates low, remains nonetheless largely unknown and should be addressed in the future.

- *Drivers of juvenile survival*

The existence of a relationship between Adélie penguin population dynamics and sea ice is well established[13,18,40,42]. However, despite the general importance of juvenile survival in

driving population trends[33,43,44], previous studies have found relatively limited evidence for a relationship between SIC and juvenile survival[31,33,36]. Here, unlike for the two populations previously studied[31,33], we found that local sea ice concentration within 200 km of the natal colony in the timeframe of one to two months after fledging (April) accounted for up to 71% of the temporal variation in survival probability (Fig. 3). Specifically, higher SIC in April had a positive effect on juvenile survival. The timing of this relationship is consistent with a survival bottleneck occurring in the weeks following fledging, as identified for Adélie and other *Pygoscelis* penguins in the Antarctic Peninsula[45]. A mortality peak immediately following fledging is also consistent with previous studies in mammals[46,47] and birds[48,49].

Our finding that SIC in April, but not in March, was correlated to juvenile survival may be particularly informative on the mechanisms linking sea ice to the survival of young Adélie penguins. Although the winter freeze-up of the Antarctic Ocean begins in March, sea ice usually consolidates markedly and expands equatorward in April[50,51] (see also Fig. S10). At the onset of winter, newly fledged Adélie penguins could then rely on such sea ice formation, especially dense landfast ice (Fig. 3, Fig. S11). By decomposing April SIC into its loose (the marginal ice zone, between 15 and 80% SIC) and dense (80-100%) fractions, we were able to show that juveniles survived better in years when sea ice was made of large, fully frozen areas, rather than patchy ice (Fig. 3B, 3C, Fig. S11).

Although only biologging studies may be able to determine how juvenile penguins exploit sea ice features, we hypothesize that dense sea ice could provide more suitable resting grounds for fledglings, in a context where they are at high risk of energy reserves depletion[45]. Furthermore, extensive fast ice areas contribute to the formation of polynyas[52], where primary productivity is highest at the end of the summer[53] and could thus concentrate penguin's prey. Large amounts of loose sea ice may therefore impair the detection and exploitation of food-abundant areas by inexperienced penguins, likely to display lower foraging abilities compared to adults as shown in king penguins (*Aptenodytes patagonicus*)[54]. Finally, patchier sea ice may favor predation, especially by leopard seals (*Hydrurga leptonyx*)[55], although predation pressure during fall and winter has never been investigated. During the winter, adult Adélie penguins tend to be associated with dense sea ice areas[56], with higher SIC promoting higher survival to the next breeding season[37]. For juveniles, however, only two studies so far have investigated their post-fledging movements[45,57], while their diving behavior and association with specific sea ice types remain completely unexplored.

While the major challenge for juvenile Adélie penguins may occur soon after fledging[45], we also found that colder windchill temperatures during their first winter at sea (April-September) were associated with lower survival rates. A likely explanation involves the progressive build-up of thermoregulatory abilities that can continue up to several months into the juvenile (post-fledging) phase[58]. Although we cannot exclude that indirect rather than direct effects are linking winter windchill temperatures to juvenile survival, our results corroborate previous findings from another colony located several thousands of kilometers away from Adélie Land[33]. More spatial coverage may be needed before confirming the generality of this driver, but weather conditions during winter (as proxied by windchill temperatures), could emerge as an important predictor of juvenile survival in this species.

Body mass at fledging is a common predictor for juvenile survival in birds[61], including penguins[62–64]. Here, we found that survival probabilities increased steadily with chick mass, up to a plateau at ca. 4 kg body mass. While this echoes previous findings for the species[62,65], this is demonstrated for the first time in a setup unbiased by flipper banding[66] and accounting for imperfect detection. Importantly, the benefits of fledging at a higher mass were similar across years, suggesting that this advantage may be maintained across a range of environmental conditions (see also[62]). While in the case of Pointe Géologie post-fledging sea ice conditions effects clearly superseded those of fledging mass in explaining interannual variations in juvenile survival, fledging mass could connect changes at other temporal scales to survival variations. For instance, both the diet[67,68] and the foraging efficiency of breeders[69,70] have been linked to sea ice conditions during the summer season or the previous winter, and these parameters may directly affect chick fledging mass[60,62]. Similarly, land-based, food-independent factors such as heavier precipitation and stronger winds in summer negatively affect chick fledging mass[59], likely because of higher thermoregulatory costs[60]. In the Western Antarctic Peninsula region, where sea ice loss is associated with a poorer chick-diet[60] and is stronger than in East Antarctica or in the Ross Sea[71], these factors could act independently or combine with post-fledging sea ice conditions to reduce juvenile survival, thereby negatively affecting population trends[72].

● *Demographic consequences and perspectives*

In long-lived species, the survival of young age classes can have major impacts on population dynamics[73]. For Adélie penguins, this is exemplified by declines in abundance of breeders that followed declines in juvenile survival probabilities in two distinct populations from the Western Antarctic Peninsula[31,32] and East Antarctica[33]. As in the present study, it was not possible to identify the onset of the decline in juvenile survival in these two populations, since trends were observable right from the beginning of the monitoring (Fig. 2). Nonetheless, evidence from East Antarctica suggests that declining juvenile survival may not cascade into an abundance decline immediately, with a ~15 years lag between the start of the juvenile survival time series and the breakup in the abundance time series[33] (Fig. 2). Furthermore, events of low adult survival and repeatedly low breeding productivity may have acted in conjunction to generate the decline in abundance at Admiralty Bay[32] and Béchervaise Island[33], respectively. This may help explain why abundance at Pointe Géologie has not yet shown signs of decline[74] (Fig. 2). However, given the occurrence of two massive breeding failures in the past decade alone[75] and the stronger rate of decline in juvenile survival observed at Pointe Géologie compared to other studies and to local inter-annual variability in this vital rate, it is key to be closely monitoring the trends of the Pointe Géologie population in the coming years.

While Adélie penguin population declines for the Western Antarctic Peninsula are well documented[76–79] and consistent with the stronger rate of warming in the region[80], declines in other regions and notably in East Antarctica were not expected before the later part of the 21st century[18], with the Ross sea likely being the main refugia for the species[17]. However, our findings in the Western Pacific Ocean sector of Antarctica, combined with recent evidence that an entire metapopulation elsewhere in East Antarctica has already declined by almost half in the past decade alone[33], suggest that changes could happen earlier than expected. Such rapid changes may challenge the adaptive abilities of the Adélie penguin. One of the main response mechanisms to these changes could be dispersal towards areas where environmental conditions remain or become favorable[18,81,82]. Although mid-range dispersion (< 120 km) is possible in Adélie penguins[83,84], we still lack estimations of long-distance dispersion over time scales relevant with the current rate of environmental change.

In conclusion, our study has highlighted the emergence of similar negative trends in juvenile survival across the Adélie penguin range. While cross-population drivers behind these trends

remain to be explored, recent range-wide studies[13,18,19] provide a relevant template to achieve this goal. The next steps would first be to harmonize demographic databases across study sites and analyze them collectively. Specifically, quantifying the temporal variation in juvenile survival for the two Ross sea populations [35,83,85] is key. Completing the transition from flipper bands to RFID-tags[66] could also facilitate comparisons of demographic rates among populations. Second, conducting biologging studies of juvenile movement and diving behavior, in conjunction with sea ice analyses (akin to[37]), is essential to uncover how inexperienced penguins exploit their environment. Third, rigorous efforts should be made to assess the dispersal capabilities of juvenile Adélie penguins under climate change, such as deploying autonomous RFID systems (Fig. S2) in populations adjacent to the main colonies where this method is used. With the forthcoming fifth International Polar Year (2032-2033) and the likely large-scale decline in the survival of juvenile Adélie penguins, a significant window of opportunity emerges for closely monitoring the demographic trends of vulnerable species and unraveling the underlying factors driving these trends.

Methods:

- *Study area and design*

Fieldwork was carried out at Pointe Géologie archipelago (Adélie Land, Antarctica; Fig. 1), close to the Dumont d'Urville research station (66°40′S, 140°01′E). This archipelago has hosted 30,000-50,000 breeding pairs of Adélie penguins annually for the last two decades[74].

From the 2006-2007 breeding season onwards (breeding season will be referred to by the fledging year, e.g., 2007 in this case), Radio Frequency IDentification tags (RFID-tags) were implanted in chicks from a colony of about 270 breeding pairs located in a natural canyon (hereafter study colony). Unlike flipper bands[66,86,87], RFID-tags are not known to negatively affect penguin vital rates. All chicks still alive right before fledging were tagged (n = 2,883 individuals). A few RFID-tagged chicks (n = 76) were found dead within the colony prior to fledging and were consequently excluded from subsequent analyses, bringing the final dataset to 2,807 individuals (range = 0-350/year, annual average = 187 from 2007 to 2021).

The study colony was fenced off in 2009 and instrumented with RFID gateways similar to those used in other penguin populations (Adélie penguins[88–90]; King penguins[91,92]; Southern rockhopper penguins[93]; Macaroni penguins[94]; Little penguins[95]). This setting made it impossible to miss tagged individuals visiting the colony. To account for dispersal from the study colony to neighboring colonies within a 10-km radius (Fig. 1), a grid of mobile RFID detection units (2-8 units, Fig. S2) was deployed every year from November to March, starting in 2013. Mobile RFID units were positioned at natural bottlenecks frequently used by penguins traveling between colonies and the sea. Units were moved every one to two weeks to maximize the number of detections. Among the 1061 individuals re-observed at least once after their tagging year, only 19 (1.8%) were detected exclusively by mobile RFID antennas, i.e. away from their natal colony. The mobile RFID-units resighting effort was more intensive closer to the study colony (< 200 m) but was performed up to 10 km from it (Fig. 1).

● *Colony and individual-scale parameters*

Colony-scale breeding productivity was defined as the number of fledglings, divided by the annual maximum number of adults inside the colony divided by two (a proxy for the number of breeding pairs). Individuals present in the colony were photo-counted weekly during the whole breeding season, starting in 2011. Cohort size was taken as the total number of RFID-tagged chicks that fledged. We ensured that observations from the study colony were representative of the larger-scale Pointe Géologie archipelago by comparing our local estimate of breeding productivity with data available from the literature for Pointe Géologie (2011-2017)[96]. Similarly, we ensured that our monitoring setup did not impact breeding abundance at the colony, by comparing the trend in number of breeding pairs in the study colony with that of two other undisturbed colonies located on the same island for the 2011-2024 period (Fig. S7).

All RFID-tagged chicks were weighed during tagging to the nearest 0.05 kg. The timing of tagging was adjusted to annual chick departure phenology (i.e. starting when the first molting chick was observed in the colony), ensuring mass was comparable across years. Each year, tagging occurred 10-15 days prior to fledging, making chick mass at tagging a reliable and consistent proxy for fledging mass[62]. In Pointe Géologie archipelago, fledging occurs from late February to early March.

 ● *Capture Mark Recapture (CMR) modeling and statistical analyses*

To model apparent survival and recapture probabilities of juvenile Adélie penguins, we

constructed capture histories for all individuals (n = 2,807) fledged between 2007 and 2021

and subsequently observed between the 2009 and 2023 breeding seasons. Out of these 2,807

fledglings, 1,061 were observed again in subsequent years.

Given the dispersal of some birds from the study colony to neighboring colonies, and the higher

resighting effort on and near the study colony, we implemented a multi-state framework to

model survival, recapture, and transition probabilities[97] using program MARK[98] v.9.0.

Different resighting probabilities among colonies were accounted for using a two-state

classification: a local state (1) for the study colony and adjacent colonies within 200 meters,

and a distant state (2) for all colonies located more than 200 meters from the study colony. This

allowed closing the system and enhancing the robustness of survival estimates despite short-

range dispersal dynamics. As the detection probability of 1-year-old individuals was near 0

(only a single bird was detected at age 1, Fig. S3), it was not possible to compute survival

probability between ages 0 and 1. The first recapture occasion following tagging was thus

deleted for all individuals (supplementary text). Consequently, juvenile survival was estimated

between 0 and 2 years old.

We used U-CARE software[99] to evaluate the goodness of fit (GOF) of the two-states time-

dependent (JollyMoVe, JMV) model to our dataset[100]. The JMV model did not fit the original

dataset because of transients (i.e. individuals never reencountered after tagging, Table S1a).

We thus tested the GOF of the JMV model to a dataset where the first capture was set to 0 for

each individual (i.e. effectively making capture history of each individual start at first

reencounter following tagging, see supplementary text), thereby allowing to test the fit of a

model with extended age-dependence on survival and recapture probabilities. The fit of the

JMV model to this dataset was satisfactory (Table S1), albeit with indications of further age

structure (i.e. > two age classes) in survival probabilities (significant test 3G.SM, Table S1b)

and capture probabilities (almost significant Test M.ITEC, Table S1b). This validated the use

of an initial umbrella model incorporating ≥ two age classes on recapture, survival, and

transition probabilities (model 1, Table S3).

In Adélie penguins, presence at the breeding colony (and thus availability for detection) is

conditional on age, with reproductively mature individuals more likely to visit colonies than

pre-breeders. Survival is also age-dependent, with lower survival probability in juveniles compared to adults[35]. Earlier studies of known-age Adélie penguins have therefore accounted for age-structured recapture and survival probabilities, with either two[31], three[36], or five age classes[35]. Here, we started from a model with five age classes on survival, recapture, and transition probabilities, and explored all the possibilities in each submodel down to only two age classes. Finally, our starting (umbrella) model was adjusted to account for the low number of individuals detected in distant colonies. This model was framed with age variation only (no time effects) for all transition probabilities, and recapture probabilities in the distant colonies (state 2). The survival probabilities in both states (i.e. local and distant) were also set to be equal in that model, which appears as a reasonable assumption considering that all colonies belong to the same population of Pointe Géologie archipelago (Fig. 1). We successively looked for the most appropriate age and time structure for recapture, transition, and survival probabilities, using the previously retained best structure for each submodel. This selection process yielded an optimal model (model 13, Table S3) including two age classes (0-2 and 3+) and time effects within each age class for survival probabilities. Recapture probabilities were best modeled using three age classes in both states. The best structure included additional time effects in each of these three age classes for the local state only (Table S3). Transition probabilities were best modeled with state and age effects (five age classes in each state, Table S3). Because survival and recapture probabilities are not separable for the last year (2021) in time-dependent CMR models, this model allowed to estimate juvenile (0-2) survival from 2007 to 2020 only. Estimates of recapture probability in the local state were low and displayed high inter-annual variability for the first age class (mean ± SD: 0.33 ± 0.24, Fig. S9), but were high and stable in the last age class (mean ± SD: 0.96 ± 0.05, Fig. S9). Recapture probabilities in the distant state were estimated for the last age class only and were particularly low (0.02, SE $= 0.009$, Fig. S9). This model was then used as a basis for testing the effect of selected covariates on juvenile survival (see below).

- *Covariate selection and definition*

Candidate environmental variables were considered based on relevance to the Adélie penguin's ecology and previous studies of Antarctic seabirds (Table S2). We first considered two large-scale indices, indicators of broad climatic variations in Antarctica and the surrounding Southern Ocean: the Southern Oscillation Index (SOI) and the Southern Annular Mode (SAM). Together, they affect wind patterns, temperatures, and sea ice dynamics, with complex

regional-specific variability[7,101]. Such variability has been linked to opposite responses both among and within species. For instance, positive SOI values were previously linked to lower survival in adult Adélie penguins in Adélie Land[34] but no support for such an effect was found in Mac. Robertson Land[36], the Western Ross Sea[35], or the Antarctic Peninsula[31]. High SAM values were also found to affect the survival of other Antarctic seabirds, either positively (adult Cape petrel, *Daption capense*[102]; juvenile Emperor penguins, *Aptenodytes forsteri*[103] or negatively (juvenile Snow petrels, *Pagodroma nivea*[104]). Monthly SOI data were downloaded from the NOAA National Centers for Environmental Information (https://www.cpc.ncep.noaa.gov/data/indices/soi) and daily SAM data from the NOAA Climate Prediction Center (https://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily_ao_index/aao/aao.shtml).

Regional sea ice extent anomalies (SIEA) were also considered because of the quadratic relationship previously observed between SIEA and the survival of adult Adélie penguins in the Western Ross Sea[35]. Monthly data for the Western Pacific sector (90°E to 150°E), which encompasses our study site, were downloaded from the National Snow and Ice Data Center (NSDIC, https://nsidc.org/data/g02135/versions/3). To capture the conditions experienced by juveniles after fledging and during their first winter at sea, each index (SOI, SAM, SIEA) was averaged from March to September[36], resulting in one value per year per index.

At the local scale, we considered both linear and quadratic effects of the average windchill temperatures (WCT) between April and September, based on the quadratic relationship previously observed between WCT and the survival of juvenile Adélie penguins[33]. Ten-day average data of minimum temperature, average wind-speed, and average relative humidity from the Dumont D'Urville weather station were downloaded from Meteo France (https://meteo.data.gouv.fr/) and used to calculate WCT following the Australian Bureau of Meteorology calculation (as in[33]).

Finally, we considered sea ice concentrations (SIC, average sea ice concentration in a specific area) at three different spatio-temporal scales (Fig. S4). As tracking data for juveniles is currently not available in our focal region, we considered broadly similar migration routes as those of adults previously studied at Pointe Géologie and found to gradually move westward after reproduction following the Antarctic coastal current[105]. This pattern is corroborated by the tracking of both juveniles and adults in another east Antarctica population[36,57] and therefore

likely in our study area. To reflect post-fledging (autumn) conditions experienced by birds near their natal colony, we considered March-April SIC between 135°E and 145°E (SIC.autumn). We also considered winter (May-July, SIC.winter) and late winter (August-September, SIC.latewinter) conditions, both between 110°E and 135°E. Sea ice concentrations were computed by averaging daily concentration values from 25×25 km grids downloaded from the NSDIC[106] (https://nsidc.org/data/g02202/versions/4) over the spatio-temporal windows defined above. Following initial analyses, we further adjusted our sea ice metrics by first decomposing SIC.autumn into March and April SIC (SIC.march, SIC.april). Because different types of ice may affect seabirds differently[107], we then calculated the April fraction of the total ice area (15-100% SIC) covered by the marginal ice zone (loose sea ice between 15-80% SIC, MIZFRAC.april) and the April fraction of the total ice area covered by dense pack ice (80-100% SIC, DENSEFRAC.april).

We also considered cohort size and average mass at tagging (as a proxy for fledging mass[62]) as likely intrinsic drivers of juvenile survival in Adélie penguins. Larger cohorts could reflect favourable growth conditions or be associated with lower predation pressure, as suggested in a previous study[33], while fledging mass is a regular predictor for juvenile survival in birds[61], including penguins[62–64]. The effect of fledging mass on inter-annual differences in survival probability was investigated by using mean fledging mass (MFM) for each cohort. Inter-individual, within-season effects of fledging mass were investigated, using chick mass as an individual covariate.

To avoid including covariates correlated with each other (multicollinearity)[25], we checked for pair-wise correlations among our initial set of covariates (SOI, SAM, SIEA, SIC.autumn, SIC.winter, SIC.latewinter, WCT, MFM, cohort size). Only SIC.winter was correlated strongly ($r > 0.7$)[108] with two other closely related sea ice metrics (SIEA and SIC.latewinter) so that we decided to drop this covariate from analyses. Remaining covariates were scaled to improve model convergence and facilitate parameters comparisons. We first investigated the effect of individual fledging mass on juvenile survival using Likelihood Ratio Tests (LRT), and included this parameter in further models if it improved model fit. We then tested annual covariates in isolation using ANODEV and considered them influential when the 95% confidence interval (CI) of the effect size did not overlap 0 and when they accounted for $\geq 20\%$ of the temporal variation in survival rates[25]. The support for a linear trend in juvenile survival was assessed using ANODEV[25] on a model set (i.e. constant, time, and trend models) that did not include

fledging mass as an individual covariate in order to account for any contribution of fledging mass in driving potential temporal trends.

Homoscedasticity and normality assumptions were assessed for linear models. Values are reported as means $\pm$ SD unless mentioned otherwise. Statistical analyses other than the CMR modeling were conducted in R[109] version 4.3.2. Timelines for the different datasets used in the study (RFID-tagging of chicks, resighting of RFID-tagged individuals, number of breeding pairs and breeding productivity data) are summarized in Table S4.

References:

1. Constable, A. *et al.* Climate change and Southern Ocean ecosystems I: How changes in physical habitats directly affect marine biota. *Global change biology* **20**, (2014).

2. Chown, S. L. *et al.* Antarctic Climate Change and the Environment.

3. Purich, A. & Doddridge, E. W. Record low Antarctic sea ice coverage indicates a new sea ice state. *Commun Earth Environ* **4**, 1–9 (2023).

4. Brooks, C. M. *et al.* Antarctic fisheries: factor climate change into their management. *Nature* **558**, 177–180 (2018).

5. Urban, M. C. *et al.* Improving the forecast for biodiversity under climate change. *Science* **353**, aad8466 (2016).

6. Lee, H. *et al.* IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (Eds.)]. IPCC, Geneva, Switzerland. https://www.ipcc.ch/report/ar6/syr/ (2023) doi:10.59327/IPCC/AR6-9789291691647.

7. Fogt, R. L. & Marshall, G. J. The Southern Annular Mode: Variability, trends, and climate impacts across the Southern Hemisphere. *WIREs Climate Change* **11**, e652 (2020).

8. Convey, P. & Peck, L. S. Antarctic environmental change and biological responses. *Sci. Adv.* **5**, eaaz0888 (2019).

9. Eayrs, C., Li, X., Raphael, M. N. & Holland, D. M. Rapid decline in Antarctic sea ice in

recent years hints at future change. *Nat. Geosci.* **14**, 460–464 (2021).

10.   Olivier, F., Franeker, J. A. van, Creuwels, J. C. S. & Woehler, E. J. Variations of

snow petrel breeding success in relation to sea-ice extent: detecting local response to

large-scale processes? *Polar Biol* **28**, 687–699 (2005).

11.   Descamps, S. *et al.* Large-scale oceanographic fluctuations drive Antarctic petrel

survival and reproduction. *Ecography* **39**, 496–505 (2016).

12.   Labrousse, S. *et al.* Landfast ice: a major driver of reproductive success in a polar

seabird. *Biology Letters* **17**, 20210097 (2021).

13.   Iles, D. T. *et al.* Sea ice predicts long-term trends in Adélie penguin population

growth, but not annual fluctuations: Results from a range-wide multiscale analysis. *Global

Change Biology* **26**, 3788–3798 (2020).

14.   Şen, B. *et al.* Spatio-temporal transferability of environmentally-dependent population

models: Insights from the intrinsic predictabilities of Adélie penguin abundance time

series. *Ecological Indicators* **150**, 110239 (2023).

15.   Fraser, W. R. & Trivelpiece, W. Z. *Factors Controlling the Distribution of Seabirds:*

Winter-Summer Heterogeneity in the Distribution of Adélie Penguin Populations. vol. 70

(American Geophysical Union, Washington, D. C., 1996).

16.   Smith, R. C. *et al.* Marine Ecosystem Sensitivity to Climate Change: Historical

observations and paleoecological records reveal ecological transitions in the Antarctic

Peninsula region. *BioScience* **49**, 393–404 (1999).

17.   Ainley, D. *et al.* Antarctic penguin response to habitat change as Earth's troposphere

reaches 2°C above preindustrial levels. *Ecological Monographs* **80**, 49–66 (2010).

18.   Cimino, M. A., Lynch, H. J., Saba, V. S. & Oliver, M. J. Projected asymmetric

response of Adélie penguins to Antarctic climate change. *Scientific Reports* **6**, 28785

(2016).

19. Che-Castaldo, C. *et al.* Pan-Antarctic analysis aggregating spatial estimates of Adélie

penguin abundance reveals robust dynamics despite stochastic noise. *Nature*

Communications **8**, 832 (2017).

20. Lynch, H. J. & LaRue, M. A. First global census of the Adélie Penguin. *tauk* **131**,

457–466 (2014).

21. Humphries, G. R. W. *et al.* Predicting the future is hard and other lessons from a

population time series data science competition. *Ecological Informatics* **48**, 1–11 (2018).

22. Talis, E. J., Che-Castaldo, C., Şen, B., Krumhardt, K. & Lynch, H. J. Variability,

skipped breeding and heavy-tailed dynamics in an Antarctic seabird. *Journal of Animal*

Ecology **91**, 2437–2450 (2022).

23. Iles, D. & Jenouvrier, S. Projected population consequences of climate change. in

Effects of Climate Change on Birds 147–164 (Oxford University Press, 2019).

doi:10.1093/oso/9780198824268.003.0012.

24. Frederiksen, M., Harris, M. P. & Wanless, S. Inter-population variation in

demographic parameters: a neglected subject? *Oikos* **111**, 209–214 (2005).

25. Grosbois, V. *et al.* Assessing the impact of climate variation on survival in vertebrate

populations. *Biological Reviews* **83**, 357–399 (2008).

26. Irons, D. *et al. Circumpolar Seabird Monitoring Plan.* (2015).

27. Nicol-Harper, A., Doncaster, C. P., Hilton, G. M., Wood, K. A. & Ezard, T. H. G.

Conservation implications of a mismatch between data availability and demographic

impact. *Ecology and Evolution* **13**, e10269 (2023).

28. Sæther, B.-E. & Bakke, Ø. Avian Life History Variation and Contribution of

Demographic Traits to the Population Growth Rate. *Ecology* **81**, 642–653 (2000).

29. Southwell, C. *et al.* Spatially Extensive Standardized Surveys Reveal Widespread,

Multi-Decadal Increase in East Antarctic Adélie Penguin Populations. *PLOS ONE* **10**,

e0139877 (2015).

30. Kappes, P. J. *et al.* Age-related reproductive performance of the Adélie penguin, a long-lived seabird exhibiting similar outcomes regardless of individual life-history strategy. *Journal of Animal Ecology* **90**, 931–942 (2021).

31. Hinke, J. T., Trivelpiece, S. G. & Trivelpiece, W. Z. Adélie penguin (Pygoscelis adeliae) survival rates and their relationship to environmental indices in the South Shetland Islands, Antarctica. *Polar Biol* **37**, 1797–1809 (2014).

32. Hinke, J. T., Trivelpiece, S. G. & Trivelpiece, W. Z. Variable vital rates and the risk of population declines in Adélie penguins from the Antarctic Peninsula region. *Ecosphere* **8**, e01666 (2017).

33. Emmerson, L. & Southwell, C. Environment-triggered demographic changes cascade and compound to propel a dramatic decline of an Antarctic seabird metapopulation. *Global Change Biology* **28**, (2022).

34. Jenouvrier, S., Barbraud, C. & Weimerskirch, H. Sea ice affects the population dynamics of Adélie penguins in Terre Adélie. *Polar Biol* **29**, 413–423 (2006).

35. Ballerini, T., Tavecchia, G., Olmastroni, S., Pezzo, F. & Focardi, S. Nonlinear effects of winter sea ice on the survival probabilities of Adélie penguins. *Oecologia* **161**, 253–265 (2009).

36. Emmerson, L. & Southwell, C. Adélie penguin survival: age structure, temporal variability and environmental influences. *Oecologia* **167**, 951–965 (2011).

37. Schmidt, A. E. *et al.* Sea ice concentration decline in an important Adélie penguin molt area. *Proc. Natl. Acad. Sci. U.S.A.* **120**, e2306840120 (2023).

38. Rintoul, S. R. *et al.* Choosing the future of Antarctica. *Nature* **558**, 233–241 (2018).

39. Reid, B. An interpretation of the age structure and breeding status of an Adélie penguin population. *Notornis Quarterly Journal of the Ornithological Society of New Zealand* 193 (1968).

40. Ainley, D. G. *The Adélie Penguin: Bellwether of Climate Change*. (Columbia University Press, New York, 2002).

41. Badia-Boher, J. A. *et al.* Joint estimation of survival and dispersal effectively corrects the permanent emigration bias in mark-recapture analyses. *Sci Rep* **13**, 6970 (2023).

42. Fraser, W. R., Trivelpiece, WayneZ., Ainley, D. G. & Trivelpiece, SusanG. Increases in Antarctic penguin populations: reduced competition with whales or a loss of sea ice due to environmental warming? *Polar Biol* **11**, 525–531 (1992).

43. Wilson, P. *et al.* Adélie penguin population change in the pacific sector of Antarctica: relation to sea-ice extent and the Antarctic Circumpolar Current. *Mar. Ecol. Prog. Ser.* **213**, 301–309 (2001).

44. Hinke, J. T., Salwicka, K., Trivelpiece, S. G., Watters, G. M. & Trivelpiece, W. Z. Divergent responses of Pygoscelis penguins reveal a common environmental driver. *Oecologia* **153**, 845–855 (2007).

45. Hinke, J. T., Watters, G. M., Reiss, C. S., Santora, J. A. & Santos, M. M. Acute bottlenecks to the survival of juvenile Pygoscelis penguins occur immediately after fledging. *Biology Letters* **16**, 20200645 (2020).

46. Cox, S. L., Authier, M., Orgeret, F., Weimerskirch, H. & Guinet, C. High mortality rates in a juvenile free-ranging marine predator and links to dive and forage ability. *Ecology and Evolution* **10**, 410–430 (2020).

47. Gaillard, J.-M. & Toïgo, C. Temporal variation in fitness component and population dynamics of large herbivores. **31**, 367–393 (2000).

48. Naef-Daenzer, B. & Grüebler, M. U. Post-fledging survival of altricial birds: ecological determinants and adaptation. *Journal of Field Ornithology* **87**, 227–250 (2016).

49. McIntyre, C. L., Collopy, M. W. & Lindberg, M. S. Survival Probability and Mortality of Migratory Juvenile Golden Eagles from Interior Alaska. *Journal of Wildlife Management* **70**, 717–722 (2006).

50. Massom, R. *et al.* Change and Variability in East Antarctic Sea Ice Seasonality, 1979/80–2009/10. *PLoS ONE* **8**, e64756 (2013).

51. Eayrs, C. *et al.* Understanding the Seasonal Cycle of Antarctic Sea Ice Extent in the Context of Longer-Term Variability. *Reviews of Geophysics* **57**, 1037–1064 (2019).

52. Nihashi, S. & Ohshima, K. I. Circumpolar Mapping of Antarctic Coastal Polynyas and Landfast Sea Ice: Relationship and Variability. (2015) doi:10.1175/JCLI-D-14-00369.1.

53. Arrigo, K. R., van Dijken, G. L. & Strong, A. L. Environmental controls of marine productivity hot spots around Antarctica. *Journal of Geophysical Research: Oceans* **120**, 5545–5565 (2015).

54. Enstipp, M. R. *et al.* The early life of king penguins: ontogeny of dive capacity and foraging behaviour in an expert diver. *Journal of Experimental Biology* **224**, jeb242512 (2021).

55. Ainley, D. G., Ballard, G., Karl, B. J. & Dugger, K. M. Leopard seal predation rates at penguin colonies of different size. *Antartic science* **17**, 335–340 (2005).

56. Takahashi, A. *et al.* Migratory movements and winter diving activity of Adélie penguins in East Antarctica. *Mar. Ecol. Prog. Ser.* **589**, 227–239 (2018).

57. Clarke, J. *et al.* Post-fledging and winter migration of Adélie penguins Pygoscelis adeliae in the Mawson region of East Antarctica. *Mar. Ecol. Prog. Ser.* **248**, 267–278 (2003).

58. Enstipp, M. R. *et al.* Apparent changes in body insulation of juvenile king penguins suggest an energetic challenge during their early life at sea. *Journal of Experimental Biology* **220**, 2666–2678 (2017).

59. Cimino, M. A. *et al.* Long-term patterns in ecosystem phenology near Palmer Station, Antarctica, from the perspective of the Adélie penguin. *Ecosphere* **14**, e4417 (2023).

60. Chapman, E., Hofmann, E., Patterson, D., Ribic, C. & Fraser, W. Marine and terrestrial factors affecting Adélie-penguin Pygoscelis adeliae chick growth and recruitment off the western Antarctic Peninsula. *Mar. Ecol. Prog. Ser.* **436**, 273–289 (2011).

61. Maness, T. J. & Anderson, D. J. Predictors of Juvenile Survival in Birds - Predictores de la Supervivencia de Aves Juveniles. *Ornithological Monographs* **78**, 1–55 (2013).

62. Ainley, D. G. *et al.* Post-fledging survival of Adélie penguins at multiple colonies: chicks raised on fish do well. *Marine Ecology Progress Series* **601**, 239–251 (2018).

63. Horswill, C. *et al.* Survival in macaroni penguins and the relative importance of different drivers: individual traits, predation pressure and environmental variability. *Journal of Animal Ecology* **83**, 1057–1067 (2014).

64. Le Scornec, E. *et al.* Intrinsic and extrinsic drivers of juvenile survival in emperor penguins. *Polar Biology* **48**, (2025).

65. Chapman, E. W., Hofmann, E. E., Patterson, D. L. & Fraser, W. R. The effects of variability in Antarctic krill (Euphausia superba) spawning behavior and sex/maturity stage distribution on Adélie penguin (Pygoscelis adeliae) chick growth: A modeling study. *Deep Sea Research Part II: Topical Studies in Oceanography* **57**, 543–558 (2010).

66. Dugger, K. M., Ballard, G., Ainley, D. G. & Barton, K. J. Effects of Flipper Bands on Foraging Behavior and Survival of Adélie Penguins (Pygoscelis Adeliae). *The Auk* **123**, 858–869 (2006).

67. Ainley, D. G. *et al.* Spatial and Temporal Variation of Diet Within a Presumed Metapopulation Of Adélie Penguins. *The Condor* **105**, 95–106 (2003).

68. Beaulieu, M. *et al.* When sea-ice clock is ahead of Adélie penguins' clock. *Functional Ecology* **24**, 93–102 (2010).

69. Watanabe, Y. Y., Ito, K., Kokubun, N. & Takahashi, A. Foraging behavior links sea ice to breeding success in Antarctic penguins. *Science Advances* **6**, eaba4828 (2020).

70. Le Guen, C. *et al.* Reproductive performance and diving behaviour share a common sea-ice concentration optimum in Adélie penguins (Pygoscelis adeliae). *Global Change Biology* **24**, 5304–5317 (2018).

71. Parkinson, C. L. & Cavalieri, D. J. Antarctic sea ice variability and trends, 1979–2010. *The Cryosphere* **6**, 871–880 (2012).

72. Gorman, K. B., Ruck, K. E., Williams, T. D. & Fraser, W. R. Advancing the Sea Ice Hypothesis: Trophic Interactions Among Breeding Pygoscelis Penguins With Divergent Population Trends Throughout the Western Antarctic Peninsula. *Front. Mar. Sci.* **8**, 526092 (2021).

73. Sæther, B.-E. *et al.* How Life History Influences Population Dynamics in Fluctuating Environments. *The American Naturalist* **182**, 743–759 (2013).

74. Barbraud, C. *et al.* Population trends of penguins in the French Southern Territories. *Polar Biol* **43**, 835–850 (2020).

75. Ropert-Coudert, Y. *et al.* Two Recent Massive Breeding Failures in an Adélie Penguin Colony Call for the Creation of a Marine Protected Area in D'Urville Sea/Mertz. *Front. Mar. Sci.* **5**, (2018).

76. Forcada, J., Trathan, P. N., Reid, K., Murphy, E. J. & Croxall, J. P. Contrasting population changes in sympatric penguin species in association with climate warming. *Global Change Biology* **12**, 411–423 (2006).

77. Trivelpiece, W. Z. *et al.* Variability in krill biomass links harvesting and climate warming to penguin population changes in Antarctica. *Proceedings of the National Academy of Sciences* **108**, 7625–7628 (2011).

78. Lynch, H. J., Naveen, R., Trathan, P. N. & Fagan, W. F. Spatially integrated assessment reveals widespread changes in penguin populations on the Antarctic Peninsula. *Ecology* **93**, 1367–1377 (2012).

79. Wethington, M., Flynn, C., Borowicz, A. & Lynch, H. J. Adélie penguins north and east of the 'Adélie gap' continue to thrive in the face of dramatic declines elsewhere in the Antarctic Peninsula region. *Sci Rep* **13**, 2525 (2023).

80. Turner, J. *et al.* Antarctic temperature variability and change from station data. *International Journal of Climatology* **40**, 2986–3007 (2020).

81. Lecomte, N. The great melt will shape unprotected ecosystems. *Nature* **620**, 499–500 (2023).

82. Gorman, K. B. *et al.* Population genetic structure and gene flow of Adélie penguins (Pygoscelis adeliae) breeding throughout the western Antarctic Peninsula. *Antarctic Science* **29**, 499–510 (2017).

83. Dugger, K. M., Ainley, D. G., Lyver, P. O., Barton, K. & Ballard, G. Survival differences and the effect of environmental instability on breeding dispersal in an Adélie

penguin meta-population. *Proceedings of the National Academy of Sciences* **107**, 12375–12380 (2010).

84. LaRue, M. A. *et al.* Climate Change Winners: Receding Ice Fields Facilitate Colony Expansion and Altered Dynamics in an Adélie Penguin Metapopulation. *PLoS ONE* **8**, e60568 (2013).

85. Morandini, V. *et al.* Sex-specific recruitment rates contribute to male-biased sex ratio in Adélie penguins. *Ecology and Evolution* **14**, e10859 (2024).

86. Saraux, C. *et al.* Reliability of flipper-banded penguins as indicators of climate change. *Nature* **469**, 203–206 (2011).

87. Gauthier-Clerc, M. *et al.* Long-term effects of flipper bands on penguins. *Proc Biol Sci* **271**, S423–S426 (2004).

88. Kerry, K., Clarke, J. & Else, G. The use of an automated weighing and recording system for the study of the biology of Adélie penguins. *Proc. NIPR Symp. Polar Biol.* **6**, 62–75 (1993).

89. Olmastroni, S., Corsolini, S., Pezzo, F., Focardi, S. & Kerry, K. The first five years of the Italian-Australian joint programme on the Adélie Penguin: An overview. *Italian Journal of Zoology* **67**, 141–145 (2000).

90. Lescroël, A., Ballard, G., Grémillet, D., Authier, M. & Ainley, D. G. Antarctic Climate Change: Extreme Events Disrupt Plastic Phenotypic Response in Adélie Penguins. *PLOS ONE* **9**, e85291 (2014).

91. Gendner, J.-P., Gauthier-Clerc, M., Le Bohec, C., Descamps, S. & Le Maho, Y. A new application for transponders in studying penguins. *Journal of Field Ornithology* **76**, 138–142 (2005).

92. Bardon, G. *et al.* RFIDeep: Unfolding the potential of deep learning for radio-frequency identification. *Methods in Ecology and Evolution* **14**, 2814–2826 (2023).

93. Dehnhard, N., Ludynia, K., Poisbleau, M., Demongin, L. & Quillfeldt, P. Good Days, Bad Days: Wind as a Driver of Foraging Success in a Flightless Seabird, the Southern Rockhopper Penguin. *PLOS ONE* **8**, e79487 (2013).

94. Afanasyev, V. *et al.* Increasing Accuracy: A New Design and Algorithm for Automatically Measuring Weights, Travel Direction and Radio Frequency Identification (RFID) of Penguins. *PLOS ONE* **10**, e0126292 (2015).

95. Robinson, S., Chiaradia, A. & Hindell, M. A. The effect of body condition on the timing and success of breeding in Little Penguins Eudyptula minor. *Ibis* **147**, 483–489 (2005).

96. Barreau, E., Ropert-Coudert, Y., Delord, K., Barbraud, C. & Kato-Ropert, A. Scale matters: sea ice and breeding success of Adélie penguins. *Polar Biol* **42**, 1405–1410 (2019).

97. Lebreton, J., Nichols, J. D., Barker, R. J., Pradel, R. & Spendelow, J. A. Chapter 3 Modeling Individual Animal Histories with Multistate Capture–Recapture Models. in *Advances in Ecological Research* vol. 41 87–173 (Elsevier, 2009).

98. White, G. C. & Burnham, K. P. Program MARK: survival estimation from populations of marked animals. *Bird Study* **46**, S120–S139 (1999).

99. Choquet, R., Lebreton, J.-D., Gimenez, O., Reboulet, A.-M. & Pradel, R. U-CARE: Utilities for performing goodness of fit tests and manipulating CApture–REcapture data. *Ecography* **32**, 1071–1074 (2009).

100. Pradel, R., Wintrebert, C. M. A. & Gimenez, O. A Proposal for a Goodness-of-Fit Test to the Arnason-Schwarz Multisite Capture-Recapture Model. *Biometrics* **59**, 43–53 (2003).

101. Kwok, R. & Comiso, J. C. Southern Ocean Climate and Sea Ice Anomalies Associated with the Southern Oscillation. *Journal of Climate* **15**, 487–501 (2002).

102. Sauser, C., Delord, K. & Barbraud, C. Demography of cape petrels in response to environmental changes. *Population Ecology* **65**, 25–37 (2021).

103. Abadi, F., Barbraud, C. & Gimenez, O. Integrated population modeling reveals the impact of climate on the survival of juvenile emperor penguins. *Global Change Biology* **23**, 1353–1359 (2017).

104. Sauser, C., Delord, K. & Barbraud, C. Increased sea ice concentration worsens fledging condition and juvenile survival in a pagophilic seabird, the snow petrel. *Biology Letters* **14**, 20180140 (2018).

105. Thiebot, J.-B. *et al.* Adélie penguins' extensive seasonal migration supports dynamic Marine Protected Area planning in Antarctica. *Marine Policy* **109**, 103692 (2019).

106. Meier, W., Fetterer, F., Windnagel, A. & Stewart, S. NOAA/NSIDC Climate Data Record of Passive Microwave Sea Ice Concentration, Version 4. NSIDC https://doi.org/10.7265/EFMZ-2T65 (2021).

107. Sauser, C. *et al.* Ice type matters: impacts of landfast and drift ice on body condition in a high Arctic seabird community. *Frontiers in Marine Science* **11**, 1326658 (2024).

108. Dormann, C. F. *et al.* Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* **36**, 27–46 (2013).

109. R Core Team (2023). _R: A Language and Environment for Statistical Computing_. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

110. Matsuoka, K. *et al.* Quantarctica, an integrated mapping environment for Antarctica, the Southern Ocean, and sub-Antarctic islands. *Environmental Modelling & Software* **140**, 105015 (2021).

Acknowledgements:

We are deeply grateful to all the members of Project 137, including Benjamin Friess, Yvon Le Maho (former PI of 137-ECOPHY), Victor Planas-Bielsa, Claire Saraux, and all the wintering and summering field teams since the inception of this project in the field in 2005. We would also like to thank all the members of the missions in Dumont D'Urville since then, and the French Polar Institute-IPEV logistics team in Dumont d'Urville for their important and continuous support in the field.

Funding:

This study was supported by the Institut Polaire Français Paul-Emile Victor (IPEV) within the framework of the Project 137-ANTAVIA, by the Centre Scientifique de Monaco with additional support from the LIA-647 and RTPI-NUTRESS (CSM/CNRS-UNISTRA), by the Centre National de la Recherche Scientifique (CNRS) through the Programme Zone Atelier Antarctique et Terres Australes (ZATA), and by the Deutsche Forschungsgemeinschaft (DFG) grants FA336/5-1 and ZI1525/3-1 in the framework of the priority program ''Antarctic research with comparative investigations in Arctic ice areas''.

Author contribution:

Author contributions follow the CRediT guidelines. In each section, authors are listed by order of appearance in the author list.

Conceptualization: TB, NL, CLB

Methodology: TB, RCH, AH, MBE, RCR, TR, NC, JC, MBU, NL, CLB

Software: TB, GB, RCH

Validation: TB, NL, CLB

Formal analysis: TB, GB, RCH, NL

Investigation: All authors

Resources: CLB

Data Curation: TB, GB, CCC, CLB

Writing – original draft preparation: TB, NL, CLB

Writing – review and editing: All authors

Visualization: TB

Supervision: NL, CLB

Project administration: CLB

Funding acquisition: DPZ, NL, CLB

Competing interests:

We declare no competing interests.

Data and material availability:

Data and codes will be made available upon acceptance of the manuscript.

 Figures and Tables:

Fig. 1. Overview of the Adélie penguin individual monitoring system operating in Pointe Géologie archipelago (Adélie Land) and location of similar monitoring sites across Antarctica still active as of 2024. (**A**) Main panel: Satellite imagery of Pointe Géologie archipelago (Pléiades Neo, Airbus DS 2021, 12/10/2021) with locations of the mobile RFID-antennas deployment sites (red), undisturbed control colonies (blue) and the main study colony (yellow). Top-left inset: location of study sites across Antarctica where longitudinal monitoring of marked Adélie penguins is still carried out (ADMI: Admiralty Bay, BECH: Béchervaise Island, EDMO: Edmonson Point, PGEO: Pointe Géologie, ROSS: Ross Island (4 monitored colonies including one on neighboring Beaufort Island)). **B**) Study colony in Pointe Géologie where all fledged chicks have been implanted with RFID-tags since 2007. (**C**) Penguins can only access or exit through two RFID-equipped passageways (only one shown here).

Fig. 2. Summary of the six locations around Antarctica where Adélie penguin juvenile survival (ϕ, 0-2 years old) has been quantified. Among them, only three have investigated temporal variations. **A)** Survival of juvenile Adélie penguins from Pointe Géologie archipelago, Adélie Land (PGEO, electronic tagging, 2007-2020). The dots and intervals represent annual survival probabilities ± SE. The regression line and associated SE are from a linear regression (p = 0.036, R^2 = 0.37); **B)** Adélie penguin abundance (number of breeding pairs) at Pointe Géologie Archipelago (2007-2019, data reproduced from[74]); **C-D)** Juvenile survival (1982-2000 aluminum bands, 1998-2011 stainless steel bands) and abundance (1982-2011) at Admiralty Bay, King George Island, Western Antarctic Peninsula (ADMI, data reproduced from[31,32]); **E-F)** Juvenile survival (electronic tagging, 1992-2015) and abundance (1992-2020) at Béchervaise Island, Mac. Robertson Land (BECH, data reproduced from[33]); **G)** Only mean estimates were available for three populations in the Ross Sea: Cape Hallett, Victoria Land (HALL, metal bands, 1959-1962, data extracted from[39]), Cape Crozier, Ross Island (CROZ, aluminum bands, 1961-1976, data extracted from[40]), Edmondson Point, Victoria Land (EDMO, electronic tagging, 1994-2002, data extracted from[35]). Survival estimates from HALL and to a lesser extent CROZ should be compared to other estimates with caution because they were not computed using modern capture-recapture techniques accounting for imperfect detection as in the other studies. Juvenile survival time series were fitted with linear regressions when authors reported a linear trend (**A,C,E**). Abundance time series were fitted with Generalized Additive Models (GAM).

Fig. 3. Relationships between the survival (φ) of juvenile Adélie penguins from Pointe Géologie archipelago (Adélie Land, East Antarctica)

and several sea ice metrics within 200 kilometers of the natal colony for the month of April. **A)** Average sea ice concentration (SIC), **B)**

Fraction of the total ice area (15-100% SIC) covered by the marginal ice zone (15-80% SIC), **C)** Fraction of total ice area (15-100% SIC) covered

by dense ice (80-100% SIC). **D)** Time series (2007-2021) of the fraction of total ice area (15-100% SIC) covered by dense ice (80-100% SIC).

Black thick lines and associated SE (gray shadow) on each panel are from linear regressions (respectively: p = 0.009, R² = 0.51 ; p < 0.001, R² =

0.79 ; p = 0.049, R² = 0.33 ; p = 0.016, R² = 0.37).

Fig. 4. Model-predicted relationship between Adélie penguin juvenile survival (φ) and individual body mass at tagging. Modeling was carried out using the program MARK[98]. The predicted relationship is drawn from model 17 (Table 1). The density probability for each fate (reobserved vs. never seen again) is superimposed (shaded areas) as well as individuals (green and violet dots).

Table 1. Comparison of the effect of covariates on Adélie penguin juvenile survival probability. Abbreviations: AICc, Akaike Information Criterion corrected for small sample sizes ; ΔAICc, AICc difference between current model and time-dependant model ; K, number of parameters ; Dev, Model deviance ; p ANODEV, p-value of the analysis of deviance ; %DEV, percentage of the time-dependent model's deviance explained by the inclusion of covariate[25] ; LRT, Likelihood Ratio Test. Covariates abbreviations: see footnote and Table S2.

Footnote Table 1: MIZFRAC.april (April fraction of total ice area between 135°E-145°E covered by the marginal ice zone (15-80% SIC)) ; DENSEFRAC.april (April fraction of total ice area between 135°E-145°E covered by dense ice (80-100% SIC)) ; SIC.april (April average SIC in area between 135°E-145°E) ; SIC.autumn (Mar-Apr average SIC in area between 135°E-145°E) ; WCT (Windchill temperatures for Dumont D'Urville in Apr-Sept) ; SAM (Southern Annular Mode, Mar-Sept) ; SIC.march (March average SIC in area between 135°E-145°E) ; SIC.latewinter (Aug-Sept average SIC in area between 110°E-135°E) ; SOI (Southern Oscillation Index, Mar-Sept) ; SIEA (Sea Ice Extent Anomalies, Mar-Sept), SIC.winter (May-Jul average SIC in area between 110°E-135°E).

Index	Individual covariate	Annual covariate	AICc	ΔAICc	AICc Weights	K	Dev	p ANODEV	Slope [95%CI]	%DEV	χ2 LRT	p LRT
Individual covariates												
19	mass^2 + time	-	12030.8	0.0	0.661	77	11875.2	-	-	-	19.976	<0.001
18	mass + time	-	12032.2	1.3	0.339	76	11878.6	-	-	-	14.598	<0.001
13	time	-	12044.7	13.9	0.001	75	11893.2	-	-	-	-	-
20	mass × time	-	12048.9	18.0	0.000	89	11868.7	-	-	-	24.499	0.0398
21	mass^2 × time	-	12160.8	129.9	0.000	102	11953.9	-	-	-	-60.734	NA
17	mass^2	-	12219.2	188.3	0.000	70	12077.8	-	-	-	50.743	<0.001
16	mass	-	12219.8	189.0	0.000	69	12080.6	-	-	-	48.014	<0.001
15	constant	-	12265.8	235.0	0.000	68	12128.6	-	-	-	-	-
Linear trend												
13	time	-	12044.7	0.0	0.001	75	11893.2	-	-	-	-	-
36	-	linear trend	12209.3	164.5	0.000	69	12070.0	0.016	-0.03 [-0.04 , -0.01]	0.31	-	-
15	constant	-	12265.8	221.1	0.000	68	12128.6	-	-	-	-	-
Annual covariates												
19	mass^2	time	12030.8	0	1	77	11875.2	-	-	-	-	-
33	mass^2	MIZFRAC.april	12076.5	45.6	0	71	11933.1	<0.001	-0.96 [-0.98 , -0.93]	0.71	-	-
31	mass^2	SIC.april	12150.6	119.8	0	71	12007.3	0.010	0.98 [0.97 , 1.00]	0.35	-	-
32	mass^2	DENSEFRAC.april	12152.8	121.9	0	71	12009.4	0.011	0.83 [0.82 , 0.84]	0.34	-	-
29	mass^2	SIC.autumn	12163.2	132.3	0	71	12019.8	0.022	0.95 [0.95 , 0.95]	0.29	-	-
26	mass^2	WCT^2	12173.2	142.4	0	72	12027.8	0.036	NA	0.25	-	-
25	mass^2	WCT	12179.4	148.5	0	71	12036.0	0.058	0.08 [0.06 , 0.10]	0.21	-	-
30	mass^2	SIC.march	12195.3	164.4	0	71	12051.9	0.145	0.77 [0.76 , 0.78]	0.13	-	-
23	mass^2	SAM	12205.6	174.8	0	71	12062.3	0.267	-0.09 [-0.16 , -0.02]	0.08	-	-
35	mass^2	MFM	12206.7	175.8	0	71	12063.3	0.283	0.06 [0.04 , 0.07]	0.07	-	-
34	mass^2	cohort.size	12217.8	187.0	0	71	12074.5	0.614	0.00 [-0.03 , 0.03]	0.02	-	-
17	mass^2	constant	12219.2	188.3	0	70	12077.8	-	-	-	-	-
22	mass^2	SOI	12219.2	188.3	0	71	12075.8	0.695	0.02 [0.00 , 0.05]	0.01	-	-
24	mass^2	SIEA	12221.2	190.3	0	71	12077.8	1.000	0.01 [-0.01 , 0.04]	0.00	-	-
28	mass^2	SIC.latewinter	12222.7	191.9	0	71	12079.4	1.000	0.1 [0.08 , 0.11]	0.00	-	-

  SUPPLEMENTARY MATERIAL:

Supplementary Text

Manipulating capture histories to account for near-zero recapture probability at
age 1 and for Goodness-Of-Fit (GOF) testing.

1. Accounting for near-zero recapture probability at age 1

In capture-recapture modeling, the capture history (CH) of an individual is coded as a
suite of numbers for each capture occasion. For example, if 0 = not seen ; 1 = seen in state 1 ;
and 2 = seen in state 2, the individual A presented in the table below was tagged in state 1 in
year 2, not seen in year 3, seen in state 1 in year 4 and seen in state 2 in year 5.

Year/Capture occasion	1	2	3	4	5
Capture history of individual A	0	1	0	1	2

The CH of individual A can be written as: "01012". Because recapture probabilities in
the year following tagging (year 3 in the case of individual A) were near 0 in our study (only
one individual was detected at one year old over our entire study period, Fig. S3), we rewrote
CH of all individuals by removing the year after tagging. For example, the CH of individual
A was transformed from "01012" into "0112".

This allows accounting for the fact that with so few individuals seen again one year after
fledging, it is not possible to estimate survival probability (ϕ) from age 0 to age 1 ($\phi\ 0\rightarrow1$).
Instead, we estimated survival from age 0 to age 2 ($\phi\ 0\rightarrow2$). This is the typical "juvenile
survival" in capture-recapture studies where individuals are not available for detection before
two years after fledging. Under the specific (but untestable in this case) hypothesis that ϕ
$0\rightarrow1 = \phi\ 1\rightarrow2$, survival from age 0 to age 1 ($\phi\ 0\rightarrow1$) can nonetheless be estimated as sqrt(ϕ
$0\rightarrow2$).

2. GOF testing

There is no direct GOF test for age-dependent, multistate capture-recapture models[99].
However, the presence of transients (individuals never seen again after tagging) is a common
source of age-dependence in recapture probabilities. To quantify how much of the lack of fit
of a time-dependent (JollyMoVe, JMV) model can be attributed to transients, a common
strategy is to test the GOF of the JMV model to a dataset where transients are removed by
replacing the first capture of each individual by 0 in their capture histories. For example, in
the dataset where transients are removed, the CH of individual A would be "00012".

When the GOF of the dataset without transients is satisfactory, the original dataset (i.e.
including transients) can be used but recapture probabilities must be specified as age-
dependent in the starting model, therefore accounting for transience.

908
909
910 **Fig. S1.**
911 **Sectors of Antarctica used for large-scale sea ice analyses.** Figure extracted from[71] .

912
Fig. S2.
A mobile RFID detection unit being deployed on an Adélie penguin (*Pygoscelis adeliae*) passageway near a breeding colony in Adélie
Land. This system allows the passive detection of RFID-tagged individuals away from the main study site where tagging is conducted.
Detection data and batteries are stored in the acquisition box (1), to which an antenna is affixed and buried in the snow on penguin passageways
(2, burying in process in this picture). To increase running time up to several weeks during the summer season, the system can be fitted with
solar panels (3). © Gregory Tran (main panel), Téo Barracho (inset), Institut Polaire Français - French Polar Institute

Fig. S3.
Age of first return at the colony after fledging for RFID-tagged Adélie penguins in
Pointe Géologie archipelago, Adélie land, Antarctica (2007-2020)

926
927
928 **Fig. S4.**
929 **Spatial windows considered for estimating the sea ice concentrations covariates for autumn (orange), and winter/late winter (green).** The
930 yellow line depicts the average April sea ice extent and the blue line the average maximal sea ice extent (both over the 1981-2010 period). Map
931 produced using Quantarctica[110] (https://www.npolar.no/quantarctica/)

Fig. S5.
Time series of breeding productivity for the study colony (2011-2024, average ± SD =
0.63 ± 0.44 chicks per breeding pair). Breeding productivity was calculated as the number
of chicks fledged per breeding pair. The number of breeding pairs was proxied as the annual
maximum number of adults inside the colony (from weekly photo counts), divided by two.
Breeding productivity may be superior to 1 in some years because Adélie penguins can raise
two chicks[40].

943
944
Fig. S6.
The annual breeding productivity (chicks per pair) of the study colony correlates with
that of the whole population of Pointe Géologie archipelago, Adélie Land, Antarctica
(2011-2017, Pearson correlation: r = 0.99, p < 0.001). Data for Pointe Géologie was
extracted from[96]. Breeding productivity may be superior to 1 in some years because Adélie
penguins can raise two chicks[40].

Fig. S7.
Number of breeding pairs (as proxied by the annual maximal number of adults counted during weekly surveys) in three neighboring Adélie penguin colonies in Pointe Géologie archipelago, Adélie Land, Antarctica. These three colonies include the study colony (Antavia) and two control colonies located on the same island (Hall Fusée and Isabelle) where no monitoring activities are conducted except for weekly photo counts. The number of breeding pairs in the study colony (Antavia) averaged 273 ± 62 between 2011 and 2024, and interannual variation remained comparable with that of the two control colonies (Spearman rank correlations, both p < 0.022).

Fig. S8.
Chick body mass at the time of RFID-tagging across cohorts (2007-2021) for Adélie penguins born in the study colony (Pointe Géologie archipelago, Adélie Land). Red dots represent annual means. Tagging consistently occurred 10-15 days prior to fledging, thus making mass at tagging a reliable proxy for fledging mass[62].

Fig. S9.
Estimated recapture probabilities of known-age RFID-tagged Adélie penguins in the
local (A) and distant (B) states. States refer to spatial locations of colonies where
individuals are detected. The local state includes the study colony and adjacent colonies (<
200 m), while the distant state includes colonies elsewhere in Pointe Géologie archipelago
(see Fig. 1 in the main text). Error bars indicate ± SE.

Fig. S10.

Annual increase in the proportion of dense sea ice (80-100% sea ice concentration) in the vicinity of Pointe Géologie archipelago (~200 km) between March and May of each year (2007-2021). The gray shaded area represents the month of April, where the increase in dense ice fraction is strongest. Daily sea ice concentration data over 25×25 km grids was downloaded from the NSDIC[106] (https://nsidc.org/data/g02202/versions/4).

Sea-ice scape near Dumont D'Urville - 29 April 2008

Sea-ice scape near Dumont D'Urville - 23 April 2018

996
997
Fig. S11.
Sea ice-scape at the end of April near Pointe Géologie archipelago in two years of
contrasting survival probabilities for juvenile Adélie penguins (2008, high survival
probability; 2018, low survival probability. MODIS satellite images were downloaded
from the NASA Worldview application (https://worldview.earthdata.nasa.gov), part of the
NASA Earth Observing System Data and Information System (EOSDIS).

 Table S1:

 Results of Goodness-of-fit tests for the multistate capture-recapture dataset used to
 estimate Adélie penguin juvenile survival probability at Pointe-Géologie archipelago,
 Adélie Land, Antarctica). df, Degrees of freedom; ĉ, Deviance inflation factor.

Test	$\chi 2$	*pval*	df	ĉ =$\chi 2$/Df
A) Original dataset				
WBWA	2.469	0.7	4	0.617
3G.SR	1827.802	<0.001	12	152.317
3G.Sm	1393.189	<0.001	35	39.805
M.ITEC	18.37	0.005	6	3.062
M.LTEC	6.831	0.2	5	1.366
JMV model	3248.661	<0.001	62	52.398
B) Dataset where 1st capt. set to 0 (no transients)				
WBWA	3.428	0.5	4	0.857
3G.SR	14.744	0.324	13	1.134
3G.Sm	47.702	0.0	31	1.539
M.ITEC	12.541	0.051	6	0.805
M.LTEC	6.015	0.3	5	0.919
JMV model	84.429	0.017	59	1.431

 Table S2.

 Annual covariates considered in the present study for explaining temporal variation in Adélie penguin juvenile survival.

Variable type	Variable	Spatial scale	Temporal scale	Hypothesized mechanisms and previous results	Expected effects on Adélie penguin juvenile survival
Global climatic index	**Southern Oscillation Index (SOI)**	global	March-September	Low SOI values linked to higher winter temperatures were negatively related to adult survival in Adélie penguins in Adélie Land[34]	Negative
	Southern Annular Mode (SAM)	global	March-September	Positive SAM values were related to higher juvenile survival in Emperor penguins in Adélie Land, possibly through increased food availability[103]. Also in Adélie Land, this relationship was negative for snow petrels, possibly	Positive

Sea ice metrics				because positive SAM increases fast ice extent and reduces food availability for this surface-feeding species[104].	
	Sea ice extent anomalies (SIEA)	regional (90°E - 150°E)	March-September	SIEA were quadratically related to Adélie penguin adult survival in the Ross sea[35] and positively related to juvenile survival in the WAP, possibly because increased ice extent provides more resting habitat[31].	Positive
	SIC.autumn (average SIC in area)	local (135°E - 145°E)	March-April	Increased autumn/winter sea ice concentrations were positively related to population growth rate across Antarctica, consistent with effects on juvenile survival[13]. Furthermore, tracking data from the Western Antarctic Peninsula (WAP) suggest a	Positive

				survival bottleneck occurs right after fledging[45].	
	SIC.march (average SIC in area)	local (135°E - 145°E)	March	Increased autumn/winter sea ice concentrations were positively related to population growth rate across Antarctica, consistent with effects on juvenile survival[13]. Furthermore, tracking data from the Western Antarctic Peninsula (WAP) suggest a survival bottleneck occurs right after fledging[45].	Positive
	SIC.april (average SIC in area)	local (135°E - 145°E)	April	Increased autumn/winter sea ice concentrations were positively related to population growth rate across Antarctica, consistent with effects on juvenile survival[13]. Furthermore, tracking data from the Western	Positive

				Antarctic Peninsula (WAP) suggest a survival bottleneck occurs right after fledging[45].	
	SIC.winter (average SIC in area)	local/regional (110°E - 135°E)	May-July	Increased autumn/winter sea ice concentrations were positively related to population growth rate across Antarctica, consistent with effects on juvenile survival[13]. Higher winter sea ice concentrations in the Weddell sea were positively related to juvenile survival in the WAP[31].	Positive
	SIC.latewinter (average SIC in area)	local/regional (110°E - 135°E)	August-September	Sea ice concentrations in Aug-Sept were weakly but negatively related to Adélie penguin juvenile survival in another east Antarctica population, possibly because too much ice pushes	Negative

				individuals away from foraging areas and/or affects predation pressure[36].	
	MIZFRAC.april (% of ice area (15-100% SIC) covered by the marginal ice zone (15-80% SIC))	local (135°E - 145°E)	April	As different types of sea ice may affect seabird vital rates differently[107], we decomposed the April sea ice concentration into its loose fraction (MIZFRAC.april) and its solid fraction (DENSEFRAC.april).	Unknown
	DENSEFRAC.april (% of ice area (15-100% SIC) covered by dense ice (80-100% SIC))	local (135°E - 145°E)	April	As different types of sea ice may affect seabird vital rates differently[107], we decomposed the April sea ice concentration into its loose fraction (MIZFRAC.april) and its solid fraction (DENSEFRAC.april).-	Unknown
Local climatic and intrinsic variables	**Windchill temperature (WCT)**	local/regional	April-September	WCT were quadratically or positively related to Adélie penguin juvenile	Positive

				survival in another east Antarctica population possibly because higher temperatures decreased thermoregulatory costs[33].	
	Cohort size	local	Annual	Density-dependent effects (positive relationship between cohort size and juvenile survival) were reported in another east Antarctica Adélie penguin population[33], possibly acting through diluted predation[55] pressure or increased social foraging.	Positive
	Average chick mass at tagging	local	Annual	Higher mass at fledging buffers inexperienced fledglings against depletion of energy reserve and is a general predictor of juvenile survival in birds[61].	Positive

1014

Table S3:

Model selection for recapture, transition, and survival probabilities. Abbreviations: AICc, Akaike Information Criterion corrected for small sample sizes ; ΔAICc, AICc difference between current model and time-dependent model ; K, number of parameters ; Dev, Model deviance.

Submodel set	Index	Model	AICc	ΔAICc (for each submodel set)	AICc Weights	K	Dev
Recapture (p)							
	3	Φ (5 age classes × time) **p(state1: 3 age classes × time; state2: 3 age classes)** Ψ(5 age classes ×	12076.5	0.0	0.902	104	2276.1
	2	Φ (5 age classes × time) **p(state1: 4 age classes × time; state2: 4 age classes)** Ψ(5 age classes ×	12081.8	5.3	0.063	114	2260.8
	1	Φ (5 age classes × time) **p(state1: 5 age classes × time; state2: 5 age classes)** Ψ(5 age classes ×	12083.0	6.5	0.035	123	2243.4
	5	Φ (5 age classes × time) **p(state1: 5 age classes + time; state2: 5 age classes)** Ψ(5 age classes ×	12095.7	19.2	0.000	74	2356.7
	6	Φ (5 age classes × time) **p(state1: 4 age classes + time; state2: 4 age classes)** Ψ(5 age classes ×	12098.2	21.7	0.000	73	2361.3
	4	Φ (5 age classes × time) **p(state1: 2 age classes + time; state2: 2 age classes)** Ψ(5 age classes ×	12135.6	59.1	0.000	93	2357.8
State transition (Ψ)							
	3	Φ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) **Ψ(5 age classes ×**	12076.5	0.0	0.991	104	2276.1
	8	Φ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) **Ψ(3 age classes ×**	12086.9	10.3	0.006	100	2294.7
	7	Φ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) **Ψ(4 age classes ×**	12087.7	11.2	0.004	102	2291.4
	9	Φ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) **Ψ(2 age classes ×**	12096.4	19.9	0.000	98	2308.3
	10	Φ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) **Ψ(state)**	12098.0	21.4	0.000	97	2311.9
Survival (Φ)							
	13	**Φ (2 age classes × time)** p(state1: 3 age classes × time; state2: 3 age classes) Ψ(5 age classes ×	12044.7	0.0	0.997	75	2303.7
	12	**Φ (3 age classes × time)** p(state1: 3 age classes × time; state2: 3 age classes) Ψ(5 age classes ×	12056.1	11.4	0.003	85	2294.7
	11	**Φ (4 age classes × time)** p(state1: 3 age classes × time; state2: 3 age classes) Ψ(5 age classes ×	12064.6	19.9	0.000	95	2282.7
	3	**Φ (5 age classes × time)** p(state1: 3 age classes × time; state2: 3 age classes) Ψ(5 age classes ×	12076.5	31.8	0.000	104	2276.1
	16	**Φ (2 age classes + time)** p(state1: 3 age classes × time; state2: 3 age classes) Ψ(5 age classes ×	12127.1	82.3	0.000	65	2406.4
	15	**Φ (2 age classes: cst/time)** p(state1: 3 age classes × time; state2: 3 age classes) Ψ(5 age classes ×	12264.6	219.8	0.000	68	2537.9
	14	**Φ (5 age classes)** p(state1: 3 age classes × time; state2: 3 age classes) Ψ(5 age classes × state)	12280.2	235.5	0.000	55	2580.0

1030 **Table S4:**
1031 **Timeline summary for the datasets used in the study** (datasets concerning the study colony only).
1032

Dataset	Time series duration	Explanation
Annual RFID-tagging of chicks ; chick mass data	2007-2021	The 2022 and 2023 cohorts were not included because there were not enough resighting opportunities for these cohorts.
Resighting of RFID-tagged birds at the study colony	2009-2023	RFID antennas at the study colony were set up in 2009, coincident with the return of the first cohort (2007).
Resighting of RFID-tagged birds outside the study colony	2013-2023	The grid of mobile RFID antennas covering neighboring colonies was set up annually from 2013 to 2023.
Juvenile survival probability of cohorts	2007-2020	The juvenile survival rate for the cohort of 2021 cannot be estimated because the last estimates of survival and recapture probabilities are confounded in fully time dependent CJS models (inherent constraint of these models).
Number of breeding pairs in the study colony and two control colonies ; Breeding productivity (BP) in the study colony	2011-2024	The weekly photographs of the study colony and two control colonies used to determine the number of breeding pairs and BP started in 2011.

Adélie penguin BP for Pointe Géologie archipelago	2011-2017	This data used to confirm that BP from the study colony was representative of the wider population was extracted from Barreau et al. (2019). Data from this study covers 1994-2017 but only 2011-2017 was used here to match our dataset for the study colony.