Odoribacter splanchnicus mitigates pathogen-induced gut inflammation and its
associated pathogenesis via its secreted bacteriocin

Dipasree Hajra[1^]**, Debapriya Mukherjee** [1#]**, Rhea Vij** [1#]**, Raju S Rajmani** [2],
Venkatareddy Dadireddy[3]**, Deepakash Das**[4]**, Tanweer Hussain**[4]**, Mahipal Ganji**[3,] **Utpal**
Tatu [3]**, Dipshikha Chakravortty** [1,5] *****

[1]Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India

^ Current Address: Department of Molecular and Cell Biology, University of California,
Berkeley, USA

[2]Centre of Infectious Disease Research, Indian Institute of Science, Bangalore, India

[3]Department of Biochemistry, Indian Institute of Science, Bangalore, India

[4]Department of Developmental Biology and Genetics, Indian Institute of Science,

Bangalore, India

[5]Adjunct Faculty, School of Biology, Indian Institute of Science Education and Research,
Thiruvananthapuram

*Corresponding author: Prof. Dipshikha Chakravortty

#Equal Contribution

Email: dipa@iisc.ac.in

Tel. No. 0091 80 22932842

Abstract

Foodborne pathogens pose a major global health hazard, worsened by drug-resistant strains.

Therefore, designing novel therapeutic strategies is crucial. Here, we identified the protective

role of gut-commensal _Odoribacter splanchnicus_ (OS) against _Salmonella_ in mice by

promoting colonization resistance, preserving gut barrier integrity, and preventing acute

inflammation and biofilm formation. _In vitro,_ both OS and their culture supernatant inhibited

Salmonella biofilm formation, intracellular proliferation in human intestinal cells, and

virulence gene expression. Further, our results depicted that OS's protective role acts over a broad spectrum as it confers protection against Gram-positive, *Listeria monocytogenes* and Gram-negative, *Salmonella* Typhimurium. Notably, OS conferred protection even when administered post-infection in mice, highlighting its therapeutic potential. Using several biochemical and proteomics approaches, we characterized key OS-secreted molecules that limit intracellular *Salmonella* and *Listeria* replication in human intestinal epithelial cells by regulating key virulence effectors and flagella. Collectively, our study establishes OS as the broad-spectrum protective agent against *Salmonella* and *Listeria* infections, with promising therapeutic potential.

Keywords: *Salmonella* pathogenesis, **Gut vascular barrier permeability**, **inflammation**, **inter-species bacterial regulation**, *Listeria*, *Odoribacter splanchnicus*, **gut commensal**, **biofilm, flagella**

Author Summary

Foodborne pathogens such as *Salmonella* and *Listeria* continue to threaten global health, especially with the rise of antibiotic resistance. In this study, we identify the gut commensal bacterium *Odoribacter splanchnicus* (OS) as a potent protectant against these pathogens. Using a mouse model, we show that OS enhances colonization resistance, maintains gut barrier integrity, and prevents both inflammation and biofilm formation during infection. Remarkably, OS also limits pathogen proliferation and virulence gene expression in human intestinal cells. These protective effects are mediated by both live OS and its secreted molecules within culture supernatant. Importantly, OS administration post-infection also conferred protection, indicating strong therapeutic potential. Biochemical and proteomic analyses revealed OS-secreted factors that suppress key virulence and motility pathways in pathogens. Our findings establish *Odoribacter splanchnicus* as a promising broad-spectrum biotherapeutic agent against major foodborne infections.

INTRODUCTION

The existence of the gut vascular barrier (GVB) acts as a protective barrier against the entry of microorganisms into systemic blood circulation (1, 2). Enteric pathogens such as *Salmonella enterica* can cross the mucus and intestinal epithelial barrier, ultimately leading to GVB damage and systemic dissemination (1). The gut microbiota aids in the maintenance of the intestinal epithelial barrier by the production of several metabolites such as short-chain fatty acids (SCFAs), indoles, and polyphenol metabolites that contribute to epithelial barrier integrity and turnover by regulating the expression of tight-junction (TJ) genes (3). Additionally, these SCFAs have the potential to exert anti-inflammatory properties by regulating cytokine production, mucosal immune cell migration, and secretion of antimicrobial peptides (AMPs) by intestinal epithelial cells and macrophages (4-6).

Salmonella enters the body through the ingestion of contaminated food and water, resisting the acidic stomach pH with acid tolerance response (ATR) genes, and finally reaches the lumen of the small intestine(7). The pathogen gains entry into the non-phagocytic epithelial cells via *Salmonella* Pathogenicity Island-1 (SPI-1) effectors to breach the intestinal barrier (8). Once traversing through the mucosal lining of the gut epithelium, it reaches the lamina propria (8). In the lamina propria, *Salmonella* encounters various host immune cells like neutrophils, macrophages, and dendritic cells. Macrophages, dendritic cells, and neutrophils are responsible for successful dissemination throughout the body through the reticulo-endothelial system (RES) to the secondary organs such as Mesenteric Lymph Node (MLN), liver, and spleen(8). *Salmonella* relies on the SPI-2 secretion for its intracellular and intravacuolar life by injecting various virulence factors into the host cell cytosol (9, 10).

Odoribacter splanchnicus (OS, belonging to the order Bacteroidales, is a common member of

the human intestinal microbiota. *Bacteroidetes* is one of the major commensal phyla

dominating the healthy mouse as well as the human gut (11). It is a Gram-negative, anaerobic,

non-motile and non-spore-forming bacterium. A decrease in *Odoribacter* abundance has been

linked to different microbiota-associated diseases, such as non-alcoholic fatty liver disease,

cystic fibrosis, and inflammatory bowel disease (IBD)(12). A recent report indicated the ability

of OS and its effectors transported in Outer membrane vesicles (OMVs) to exhibit anti-

inflammatory effects on the gut epithelium(12). Further, OS has been implicated in conferring

resistance against colitis and colorectal cancer(13). This led us to hypothesize that *O.*

splanchnicus might ameliorate the inflammatory disease conditions associated with *Salmonella*

infection and dissemination. Here, in the present study, we aim to understand whether

colonization with the gut commensal OS, could ameliorate *Salmonella*-induced inflammation

and pathogenic dissemination.

Our study shows that OS colonization promotes colonization resistance of *Salmonella*, and

ameliorates systemic dissemination. OS ameliorates acute *Salmonella*-induced inflammation

and its associated pathogenesis by promoting colonization resistance, resulting in decreased

Salmonella invasion of gut tissue, the preservation of tight junctions, and GVB maintenance.

Further, we demonstrate that not only live OS but its culture supernatant inhibits *Salmonella in*

vitro biofilm formation, intracellular proliferation in human intestinal epithelial cells, and

virulence gene expression. Subsequently, our results depict that the specific protective role of

OS acts over a broad spectrum as it confers protection against both flagellated Gram-positive,

Listeria monocytogenes and Gram-negative, *Salmonella* Typhimurium. Furthermore, OS

conferred protection even after its administration to mice post-establishment of infection

highlighting its therapeutic potential. Subsequently, we identified the key OS-secreted

bacteriocin protein responsible for limiting intracellular *Salmonella* and *Listeria* replication in

human intestinal cells by regulating virulence genes such as flagella. Overall, our study implicates the broad-spectrum protective role of OS in mitigating foodborne infections caused by *Salmonella* and *Listeria* pathogenesis.

MATERIALS AND METHODS

Bacterial Strains and Culture Conditions

Salmonella enterica serovar Typhimurium (STM) strain ATCC 14028S, STM ATCC 14028S constitutively expressing mCherry (pFPV25.1-mCherry-Amp^R, RFP), *Salmonella enterica* serovar Typhi CT18 (STY), laboratory stock of *Enterococcus faecalis* strain API 142 MTCC 439 (ATCC 35550) (Ef), *Listeria monocytogenes* MTCC 657 (LM) expressing mCherry (pFPV25.1-mCherry-Amp^R), and *Odoribacter splanchnicus* (OS) strain NCTC 10825 [1651/6] from ATCC were used in this study. 4% paraformaldehyde-fixed-OS was used as killed bacteria control. The STM, LM, and Ef bacterial strain was grown overnight in Miller LB broth (Himedia, M1245) at 37 ^{0}C under 160 rpm shaking conditions. OS was cultured under anaerobic conditions (anaerobic chamber) in tryptic soy broth (Himedia, M-011) at 37ºC for 48-72h after its revival from glycerol stock.

Preparation of Cell-free Bacterial Culture Supernatant

The overnight grown OS culture in tryptic soy broth was centrifuged at 10,000rpm for 10 min and the supernatant was harvested. The collected supernatant was further subjected to another two rounds to centrifugation to remove any residual cells. The supernatant was subsequently filtered using a 0.2 µm filter (Cole-Parmer PTFE Syringe Filters) and stored at -20˚C.

Animal Experiments

The animal experiments were conducted in accordance with the approved guidelines of the institutional animal ethics committee at the Indian Institute of Science, Bangalore, India

(Registration No: 48/GO/ReRcBiBt-S/Re-L/99/CPCSEA). All procedures involving the use of animals were performed according to the Institutional Animal Ethics Committee (IAEC)-approved protocol (CAF/Ethics/853/2021). 6–8 weeks old adult C57BL/6 male mice were used in this study (bred, housed, and provided (purchased) by Central Animal Facility, Indian Institute of Science, Bangalore). Male mice were used to minimize variability associated with estrous cycle–dependent hormonal fluctuations that can modulate innate and adaptive immune responses and influence infection outcomes. Using a single sex reduced experimental variability and improved statistical power to detect infection-induced phenotypes within the scope of this study. After one day of OS colonization (10^5 CFU/mice via oral gavage), the mice were infected with 10^6 (organ burden) CFU or 10^8 CFU (for assessing survival) of *S.* Typhimurium, and 10^8 (organ burden) or 10^9 (survival) CFU of *L. monocytogenes* via oral gavage. The colonization of *O. splanchnicus* was confirmed via RT-qPCR mediated quantitation of OS 16S rRNA from the mice fecal (3 fecal pellets harvested from the intestine and resuspended in Trizol) and ileal samples at 6^{th} day post-OS colonization. For post-infection OS administration studies, OS was orally administered (10^5 CFU/ mice) post 1 or 3dpi. 5-day post-infection, mice were euthanized, dissected out, and blood and organs were harvested. The organs were homogenized using a bead-beater. The bacterial organ load was enumerated by plating the tissue homogenates on *Salmonella-Shigella* (SS) agar plates for *Salmonella* or LB plates containing 50µg/ml of Ampicillin for LM. The harvested tissues and the fecal samples were collected in protein lysis buffer (RIPA) and TRIzol (Takara) for protein and RNA extraction respectively and were stored at -80ºC. For calculating percentage survival, the infected mice cohorts were monitored till fatality and fatalities were reported.

Blood collection and serum preparation

Blood collection was performed on the 5^{th} day post-infection via cardiac puncture post-euthanization of the mouse. The collected blood was allowed to clot by incubating the vials at

room temperature for 45 minutes. After allowing the standing time of 45 mins, centrifugation was performed at 2000g for 15 minutes in a refrigerated centrifuge. Following the centrifugation, the clear, yellowish supernatant was transferred to an autoclaved microfuge tube and stored at -80ºC till further downstream processing.

ELISA

Cytokines in mice serum were quantified according to the manufacturer's instructions(10, 14). Briefly, 96-well ELISA plates of IL-6 ELISA kit (BD Bioscience- Cat No-555240) were coated overnight with capture antibody at 4°C. Following day, plates were washed with 0.1% Tween-20 containing PBS and blocked with 10% FBS for 1 h. Following blocking, wells were washed and incubated with 50µl of serum samples in duplicates for 2 h at room temperature. Subsequently, plates were washed and incubated with detection antibody and enzyme reagent for 1 h at room temperature (BD Bioscience). TMB (Sigma) was used as a substrate and reactions were stopped with 2 N H_2SO_4.

Absorbance was measured at 450 nm wavelength in Tecan Plate reader and the concentration of cytokines were interpolated from a standard curve.

Quantitative Real Time PCR

Total RNA was isolated from the ileal section of the intestinal tissues or feces harvested in TRIzol (Takara) as per manufacturer's protocol. RNA was quantified using Nano Drop (Thermo-Fischer scientific).3 µg of total isolated RNA was subjected to DNaseI (Thermo Fischer Scientific) treatment at 37°C for 1 hr followed by heat inactivation at 75°C in the presence of 5mM concentration of EDTA for 10 mins. The mRNA was reverse transcribed to cDNA using oligo $(dT)_{18}$ primer, random hexamer primer, buffer, dNTPs and reverse transcriptase (Takara) as per manufacturer's protocol. The expression profile of target gene was evaluated using specific primers (Table-S2) by using SYBR green RT-PCR master mix

175 (Takara) in Applied Biosystems (Quantstudio 5) or BioRad Real time PCR instrument. β-actin

176 was used as an internal control for mammalian genes and for bacterial genes 16S rRNA was

177 used. All the reaction was setup in 384 well plate with three replicates for each sample.

Western blotting

The harvested tissue samples were lysed in ice-cold RIPA buffer (50 mM Tris-Cl, pH 8.0, 150

mM NaCl, 5mM EDTA, 1% NP-40, 1 % Sodium deoyxycholate, 1% SDS) containing protease

inhibitor cocktail (Roche). Total protein was estimated using Bradford reagent (BioRad).

Protein samples were denatured in **Laemmli SDS loading buffer** at 95ºC for 10 min and then

subjected to separation on 12% SDS-PAGE gels followed by transfer onto activated 0.45μm

PVDF membrane. Finally, the membrane was blocked with 5% skimmed milk in TBST (Tris

Buffered Saline containing 0.1% Tween-20) and probed with appropriate primary (Table-S3)

and secondary HRP-conjugated antibodies. The blots were developed using ECL substrate

(Bio-Rad) using a Chemidoc (BioRad). All densitometric analysis was performed using

ImageJ software.

Immunohistochemistry

The harvested tissues were fixed in 4% paraformaldehyde, embedded in paraffin and 5 μm

thick paraffin sections were collected on poly-lysine coated glass slides. The sectioned tissue

samples were deparaffinized using standard protocol. Briefly, the slides containing the

paraffinized tissues were immersed in xylene for 10 minutes and this step was repeated twice.

After the xylene treatment, the tissues were subjected to absolute alcohol treatment for 5

minutes twice, followed by 95% and 70% alcohol treatment for 5 minutes. The slides were

rinsed in water followed by boiling step in water for 2 minutes and then dried. For in vivo

estimation of biofilm, deparaffinized tissue samples were stained with Calcofluor White

(Sigma) stain for 20 min at RT. All immunofluorescence images were obtained using Zeiss

multi-photon 880 microscope and were analyzed using ZEN black 2012 software.

Haematoxylin and Eosin Staining

The harvested ileal tissues were collected and fixed using 3.5% paraformaldehyde. The fixed

liver was then dehydrated using a gradually increasing concentration of ethanol and embedded

in paraffin. 5μm sections were collected on coated plates. Sections were further rehydrated and

then stained with hematoxylin and eosin. Images were collected in a Leica microscope.

The scoring of the pathological changes is graded as 0-4; for each of the pathological lesions

(Severe/ marked -4; Moderate -3; Mild -2; Minor/minimum -1; No pathology -0) considering

the intestinal inflammation and associated villi injury as described previously (15, 16).

Cell Culture

210 Human colorectal adenocarcinoma cell line Caco-2 were maintained in Dulbecco's Modified

211 Eagle's Media (DMEM) (Lonza) and supplemented with 10% FCS (Fetal calf serum, Gibco)

212 and 1% non-essential amino acid solution (Sigma Aldrich) at 37ºC temperature in the presence

213 of 5% CO_2 in humidified incubator. Prior to each experiment, cells were seeded into 24 well

214 or 6 well plate as per requirement at a confluency of 60%.

Intracellular proliferation or gentamicin protection assay

Caco-2 epithelial were infected with 3 to 4 h old log phase STM or LM or STY bacterial culture,

STM (or LM or STY) with OS, STM (or LM or STY) in presence of *O. splanchnicus*

supernatant and STM along with PFA-fixed OS at an MOI of 10. For attachment of bacteria to

the cells, tissue culture plates were subjected to centrifugation at 600xg for 5 min and incubated

at 37 ^{0}C humified incubator with 5% CO_2 for 25 min. Following infection of the cells with

STM at an MOI of 10, cells were treated with DMEM (Sigma) + 10% FBS (Gibco) containing

100 µg/ml gentamicin for 1 hr. Subsequently, the gentamicin concentration was reduced to 25

µg/ml and maintained until the specified time point. Post 2hr and 16hr post-infection, cells

were lysed in 0.1% triton-X-100. Lysed cells were serially diluted and plated on *Salmonella-*

Shigella (SS) or LB agar to obtain colony-forming units (CFU). Percent invasion was

determined using the following formula (17, 18):

Percent invasion= [CFU at 2 h]/ [CFU of pre-inoculum] X 100

Fold proliferation was calculated as CFU at 16hr divided by CFU at 2hr.

Fold Proliferation= [CFU at 16h] / [CFU at 2h]

Immunofluorescence confocal microscopic studies

At the specified time points post infection with mCherry-tagged STMAS, cells were fixed with

3.5% paraformaldehyde for 15 min. Primary antibody staining was performed with specific

primary antibody in the presence of a permeabilizing agent, 0.01% saponin (Sigma) dissolved

in 2.5% BSA containing PBS at 4°C for overnight or for 6hr at room temperature (RT).

Following this, cells were washed with PBS stained with appropriate secondary antibody

tagged with fluorochrome for 1 hr at RT. This was followed by DAPI staining and mounting of

the coverslip onto a clean glass slide using the mounting media containing the anti-fade agent.

The coverslip sides were sealed with a transparent nail paint. All immunofluorescence images

were obtained using Zeiss Multiphoton 880 and were analyzed using ZEN Black 2012 software.

Growth Curve analysis

For *in vitro* growth curve experiments a single colony of STM and OS was inoculated in 5mL

of LB broth singly or in co-culture and grown overnight at 37°C in anaerobic jar. In another

set, STM was inoculated in 5mL of LB in presence of OS supernatant. Overnight-grown

stationary phase bacteria were sub-cultured at a ratio of 1: 100 in freshly prepared LB in

presence or absence of OS supernatant and kept at 37ºC. At different time intervals, aliquots

were taken for measurement of [OD]600nm by TECAN 96 well microplate reader and were

plated for CFU enumeration.

Biofilm-inducing growth media and conditions

249 LB medium lacking NaCl and containing 1% casein enzyme hydrolysate (Hi-Media, India) and

250 0.5% yeast extract (Hi-Media, India)) was used as biofilm media as per previous literature (19).

251 Overnight grown culture was subcultured in 2 ml biofilm medium at the dilution of 1:100 in a

252 flat-bottom 24-well plate (Tarsons, India) and incubated at 28˚C for 72 h without shaking.

Biofilm inoculation with OS culture supernatant

254 20 µl of STM overnight culture and indicated volume (1% v/v or as indicated) of the OS culture

255 supernatants were added to 2 ml of biofilm media and incubated under biofilm-promoting

256 conditions(19). For co-culture experiments, the overnight cultures of the STM and OS or STM

257 and PFA-fixed OS were added in a 1:1 ratio to 2 ml of biofilm media and incubated under

258 biofilm-inducing conditions.

Crystal Violet Staining

Post 72 h of biofilm induction, the media was discarded, and the plates were thoroughly washed

in water. The plates were dried, and 2 ml of 1% Crystal Violet (CV) was added to each well

and incubated for 15 min. Post 15 min incubation, the CV stain was removed, and the plate was

thoroughly rinsed in water to remove excess stain. The stained biofilm was destained with 70%

ethanol, and the intensity of the colour of the destained solution was quantified at OD_{595} in the

Tecan plate reader (Infinite Pro 200) (19).

Glass bead assay

1 mm glass beads were added onto the pellicle to assess the strength of the pellicle at the air-

liquid interface. The initial weight of the glass beads was noted. The glass beads were added

until the pellicle collapsed, and the final weight was noted down and plotted.

Gut Permeability Assay

The infected mice cohorts were orally administered with FITC Dextran (25mg/kg of mice body

weight, FD-4-100MG, Sigma) 4h before euthanization. The blood plasma was collected post-

euthanization by cardiac puncture followed by centrifugation. The blood plasma was evaluated

for mean fluorescence via plate reader (TECAN) and flow cytometry(CytoFlex, Beckman).

MTT Assay

Post-16h treatment of pre-seeded Caco-2 cells with 10µg/ml of bacteriocin, MTT (3-(4,5-

dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) staining solution (0.5mg/ml) was

added to each well followed by incubation at 37°C CO_2incubator for 4h. 50µl of DMSO was

added per well for solubilizing the formazan crystals. Absorbance was measured at 570nm

using a plate reader (TECAN).

Transmission electron microscopy

Overnight STM or LM-grown cultures (in the presence or absence of OS supernatant or

bacteriocin) were subcultured into fresh media (1:10) ratio and incubated at 37°C for 6 hours

under shaking conditions. These samples were centrifugated at 1000rpm for 10 min to obtain

the bacterial pellet. The bacterial pellet was subjected to PBS wash twice via centrifugation at

1000 rpm for 10 min.5 µl of the finally resuspended cell suspension was added on a glow-

discharged copper-grid, air dried, stained with 1% uranyl acetate for 30 s, and visualized under

a transmission electron microscope (JEM-100 CX II; JEOL).

Supernatant Treatment

The supernatant was treated with 7.5 mM concentrations of EDTA and incubated at 65 °C for 1 h. 10 µl of proteinase-K (NEB, stock 20 mg/ml) was added to the 200 µl supernatant and incubated at 37 °C for 1 h. The proteinase was inactivated with 0.5 mM PMSF and incubated at 28 °C for 1 h. The supernatant was subjected to heating at 37°C (60 min), 65°C (15 min) or 95°C (15 min) and immediately frozen at −20°C (19). Post-treatment, the supernatants were used to inoculate the biofilm to test the nature of the biofilm inhibitory molecules.

Supernatant concentration and molecular weight-based separation

4 ml of the supernatants were transferred to the 3k MWCO or 10k MWCO Amicon ultra filter centrifugal device (Amicon® Ultra-4 centrifugal filter device, 3k MWCO, UFC800324; Amicon® Ultra-4 Centrifugal filter Unit, 10k MWCO, UFC801024) and centrifuged at 4000 g for 30 min in a swing bucket rotor(19). The concentrated solute from the bottom of the filter device and the flowthrough were collected and subjected to biofilm inoculation.

Size Exclusion Chromatography

The bacterial supernatant was concentrated to 0.5ml using 3k MWCO-Amicon-Ultra-4-centrifugal unit and loaded onto the Superdex 200pg size exclusion column (XK16/60, 124mL; Cytiva, USA), pre-equilibrated Tris-Cl buffer (pH-8) containing 100mM NaCl and 20mM Tris-Cl. The sample was eluted with the same buffer at a flow rate of 0.8mL/min, and 1mL fractions were collected. The fractions corresponding to the elution peaks were analyzed for biofilm inhibition activity and subjected to mass spectrometry.

LC/MS analysis of the supernatant fractions

The concentrated gel filtration fractions of supernatants were subjected to in-solution trypsin

digestion for LC Q-TOF MS analysis. Briefly, 100 µg of total protein was resuspended in

100 µl of 7-8M of urea buffer. To this, 5 µl 200mM DTT was added and incubated at room

temperature for 1 hr, followed by alkylation with 20µl of 200 mM iodoacetamide and

incubation at room temperature for 1h in the dark. This was followed by the addition of 1mM

of Calcium chloride and incubation at RT for 10 min. For protein digestion, trypsin solution

was added to the sample to a final trypsin to protein ratio of 1:30 (w/w) and incubated at 37˚C

for 16 h, with frequent shaking. The reaction was stopped by adding 0.1% formic acid to adjust

the pH to 3-4. The digested protein samples were desalted using a reverse phase C-18 column

(Acclaim prep EASY-Spray (C18-RP)) using mobile phase solvent A-2% acetonitrile and

solvent B- 80% acetonitrile. The sample injection volume was 10µl. All the samples were

analyzed using Thermo Scientific SII for Xcalibur method. The MS scan was acquired using

Orbitrap detector with 60000 resolution in the positive polarity and scan range (*m/z*) 350-2000.

The ion source type was NSI with static voltage; positive ion (V)- 2200V and negative ion (V)-

600 along with temperature settings at 275ºC. This was followed by ddMS^2 IT HCD with 30%

fragmentation HCD collision energy using ion trap detector. The raw data was analyzed using

Proteome Discoverer 2.5 against the *O. splanchnicus* proteome available in UniProt (Proteome

ID: UP000006657).

Purification of bacteriocin

O. splanchnicus culture supernatant was concentrated within the range of >3kDa and <10kDa

using the Amicon Ultra-4 Centrifugal filter units of 10MWCO and 3MWCO. The concentrated

cell-free supernatant containing the bacteriocin (~7kDa) was precipitated by ammonium

sulphate (80% to 98%). The precipitate was collected by centrifugation at 12000rpm for 30

min and dissolved in 1X PBS. The dissolved precipitate was again filtrated using the 3KDa

cutoff Amicon centrifugal filter units. This concentrated fraction was further subjected to Fast

Performance Liquid Chromatography using Superdex 75 Increase 10/300 GL- 24ml volume

(GE Healthcare) column. The column was equilibrated with 1X Phosphate Buffered Saline

buffer and 0.5ml of concentrated sample (23.6 mg/ml) was injected into the column. Based on

the increase in Absorbance units at 280nm from baseline, 0.3ml fractions were collected at a

retention volume of 13.3ml to 21.8ml (Fig. S8). The fractions were analyzed by 20% SDS-

PAGE electrophoresis and stained with Coomassie staining solution.

In vitro **swim motility assay**

2 µl ($2*10^6$ CFU/ml) of treated or untreated stationary phase bacterial samples were spotted on

the 0.3% agar (LB-STM, BHI-LM) plates supplemented with 0.5% yeast extract, 1% casein

enzyme hydrolysate, 0.5% NaCl, and 0.5% glucose. The plates were incubated at 37˚C and

images were taken every 6 h (STM) or 24h (LM) using the camera of Biorad illuminator.

Statistical analysis

Data were analyzed and graphed using the GraphPad Prism 8 software (San Diego, CA).

Statistical significance was determined by Student's t-test or One-way ANOVA followed by

Dunnett's Multiple Comparisons test to obtain p values. ANOVA was used during comparison

among three or more groups whereas Student's t-test was used to compare the difference

between two groups. Mann Whitney t-test was used as a non-parametric counterpart of Student

t-test when variances differed. The log-rank test also known as Mantel-Cox test is used to

compare the survival percentages of the mice groups. Adjusted p-values below 0.05 are

considered statistically significant (****$p<0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$). The

results are expressed as mean ± SD or SEM of three independent experiments.

RESULTS

O. splanchnicus hinders *Salmonella* burden and its systemic dissemination

To denote the role of OS in *Salmonella* pathogenesis, we colonized C57BL/6 mice with 10^5 CFU of OS 1 day prior to *Salmonella* infection via the oral route and validated the colonization by undertaking OS 16S rRNA qPCR analysis (Fig 1, A-B). OS pre-colonized mice showed decreased susceptibility to *Salmonella* infection in comparison to the non-colonized mice cohort which succumbed to death by day 9 post-infection (p.i.). Contrastingly, 100% of the OS colonized mice were alive even at day 11 post *Salmonella* infection (Fig 1, C). *Salmonella* colonization of the feces and intestinal tissue and dissemination to the MLN, blood, and liver were reduced upon OS pre-colonization (Fig 1, D-K). OS non-colonized mice exhibited increased weight loss percentage upon *Salmonella* infection and OS pre-colonization prevented weight loss (Fig 1, H). Our result suggests OS could ameliorate *Salmonella* colonization, organ burden and dissemination (Fig 1).

O. splanchnicus colonization ameliorate the acute *Salmonella-induced* inflammation and its associated pathogenesis

Heightened inflammatory responses facilitate *Salmonella* dissemination in mice[20]. Thus, we hypothesized that OS gut colonization might ameliorate the *Salmonella*-induced inflammation. ELISA-mediated IL-6 serum quantification revealed reduced IL-6 production upon *S.* Typhimurium (STM) infection in the OS colonized mice group to the non-colonized cohort (Fig 2, A). To combat the enteric pathogen challenge, the host employs one of its major defence arsenals, that is, the production of antimicrobial peptides (AMPs)[21, 22]. STM infection induces enhanced mRNA expression of AMPs in mice with respect to the uninfected control as per previous literature findings [17]. We found that OS colonization in mice prevented STM infection-induced expression of several AMP genes such as Lysozyme-C1 precursor,

Angiogenin-4 precursor, CCL20, and Regenerating-islet-derived 3 beta-precursor (Fig 2, B).

The haematoxylin eosin-stained intestinal sections suggested increased intestinal tissue

damage with irregularly organized villi, necrotic submucosa layer, neutrophil infiltration, and

detached intestinal cells in all the STM-infected non-OS-colonized cohorts. However, OS pre-

colonization restored the intestinal tissue architecture in all the mice cohorts (Fig 2, C-D).

O. splanchnicus restricts Salmonella-mediated GVB damage and promotes tight junction

gene expression

Salmonella breaches the gut vascular barrier (GVB) by inducing tight junction disruption.

Salmonella like other food-borne antigens enters into the Mesenteric Lymph Node (MLN) by

accessing the lymphatics and subsequently gains entry into the liver via the bloodstream(23).

The translocation of the pathogen across the blood endothelial cells is governed by the integrity

of the GVB(1). The GVB refers to the endothelial barrier and the microvasculature

characterized by the elaborate framework of the junctional complexes (1). The GVB

architecture is maintained by several tight junctions and adherens junction proteins that control

paracellular trafficking(24, 25). Other cell types associated with the microvasculature of the

GVB forming a vascular unit includes pericytes and fibroblasts(26). PV-1 (plasmalemma-

vesicle associated protein-1) expression is a marker of GVB damage and heightened PV-1

expression denotes EC permeability(27, 28). Our data show reduced *PV-1* marker expression,

alongside enhanced tight junction protein mRNA expression, such as *Tjp1* and *Claudin5*, upon

OS colonization during *Salmonella* infection in comparison to the non-OS-colonized cohorts.

(Fig. 3, A). Previous studies in *Mycobacterium marinum* have shown that transcriptional

induction of pro-angiogenic markers like *Vegfa* contributes to increased bacterial burden and

dissemination(29). In our study, OS pre-colonization in the gut resulted in reduced expression

of the pro-angiogenic *Vegfa* in the infected liver tissues, which coincided with the reduced *S.*

Typhimurium burden in the bloodstream and liver (Fig S1, A). Previous literature suggested

reduced expression of *Axin2,* a β catenin target gene, upon wildtype *Salmonella* infection, even

after Wnt3a stimulation suggesting that *Salmonella* affects the GVB by inhibiting canonical

Wnt signalling (1). Here, our results suggest that OS pre-colonization restores the *Salmonella-*

triggered attenuated Wnt signalling response with increased Axin2 expression in the

Salmonella-infected intestinal tissues (Fig. S1, B). The integrity of the infected mice gut was

estimated by the gut permeability assay using FITC-Dextran (Fig. 3, B). Our results indicate

increased gut vascular permeability upon *Salmonella* challenge which declined upon OS pre-

colonisation, implicating the ability of OS to prevent gut barrier disruption.

***O. splanchnicus* culture supernatant restricts *Salmonella* biofilm formation and**

intracellular proliferation in human intestinal cells

To delineate the mechanism behind the protective role of OS against *Salmonella* pathogenesis,

we performed *Salmonella* infection studies in human intestinal epithelial cell line Caco-2 in

the presence of OS. The intracellular survival assay suggested decreased STM invasion and

subsequently attenuated intracellular replication within the infected Caco-2 cells upon co-

infection with live OS and in the presence of OS culture supernatant (Fig 4, A-B). However,

STM co-infection with fixed OS showed percent invasion and intracellular replication like that

of STM infection (Fig 4, A-B). Together, these results hinted at the presence of some secreted

inhibitory components in the OS culture supernatant that has the propensity to limit STM

infection. The reduced invasion and proliferation of STM within Caco-2 cells in presence of

OS was subsequently validated in immunofluorescence studies, wherein STM showed a

significant reduction in invasion at 2hr p.i. and intracellular proliferation at 16hr p.i. in case of

OS co-infection (Fig 4, C-D). Similar results were obtained upon infection with the human-

restricted serovar STY to the Caco-2 cells wherein OS and its supernatant attenuated STY

invasion and intracellular replication (Fig. 4, E-F). Further, we assessed the *in vitro* growth of

the STM in co-culture conditions with OS or in the presence of OS culture supernatant in

anaerobic conditions to mimic the microaerophilic or hypoxic environment of the gut or in

aerobic culture conditions. We observed no significant delay *in vitro* growth of STM at its

initial lag to log phase of its growth in the presence of OS supernatant or with live OS under

anaerobic culture conditions (Fig S2, A-B). In aerobic growth conditions as well, *Salmonella*

growth remained unperturbed even in the presence of OS or its culture supernatant (Fig S2, C-

D). Several previous studies have corroborated the ability of commensal bacterial culture

supernatants to inhibit colonization and the biofilm formation of several pathogens. For

instance, a commensal bacterium, *Staphylococcus epidermis*, could inhibit biofilm formation

and nasal colonization of its pathogenic counterpart, *Staphylococcus aureus* (30). Moreover,

biofilm formation is also regulated at the interspecies level. Indole is an inter-species biofilm

regulator that decreases *E. coli* biofilm formation and facilitates biofilm formation of

pseudomonads (31). STM forms biofilm to colonize the host, evade host defence arsenals and

antibiotics, persist during chronic infections, and transmit to new hosts (32, 33). We, therefore,

examined the biofilm pellicle strength and biofilm formation of STM via crystal violet staining

on the solid-liquid interface in the presence or absence of live, PFA-fixed-OS or OS culture

supernatant by inoculating STM in biofilm-inducing media with low salinity. STM exhibited

biofilm formation defect in the presence of live OS or OS supernatant (Fig 4, G). However, the

STM biofilm inhibitory property of OS was diminished upon treatment with paraformaldehyde

(PFA)-fixed OS (Fig 4, G). Further, qPCR-mediated gene expression studies of *Salmonella*

biofilm genes revealed decreased expression of genes regulating biofilm (*csgD*, *csgA*, *bapA*)

under treatment with live OS or its culture supernatant (Fig.4, H-J). Flagella-mediated

attachment and several virulence factors such as SPI-1 and SPI-2 genes are responsible for

Salmonella pathogenesis and its biofilm formation (19, 34, 35). Therefore, we evaluated a few

of the *Salmonella* flagellar and SPI-1 and SPI-2 gene expression in the presence or absence of

OS or its culture supernatant via quantitative real-time PCR. Our results depicted attenuated

expression of flagellar gene (*fliC*), SPI-1 (*sipC*) and SPI-2 (*pipB2*) genes both in the presence

of live OS and its spent media but not in presence of PFA fixed-OS (Fig. 4, K-M). The loss in

Salmonella flagellar motility was further validated via TEM imaging wherein OS or its

supernatant reduced the flagella formation of *Salmonella* considerably (Fig.4, N). Together,

our results implicate the ability of OS to inhibit STM biofilm formation via active secretion of

certain inhibitory molecules in the supernatant.

***O. splanchnicus* inhibitory effect on *Salmonella* is specific as *Enterococcus faecalis*,**

another gut resident microbe, fail to inhibit *Salmonella* intracellular proliferation or

biofilm formation

The resident gut microbes provide colonization resistance against invading pathogens by

deploying several mechanisms, including the production of inhibitory molecules, nutrient

competition, maintenance of the intestinal epithelial integrity, immune activation, and via the

deployment of bacteriophages (36, 37). To understand whether our observed inhibitory trait of

OS against *Salmonella* is due to a common colonization restriction phenomenon of gut

microbes via competitive inhibition, or it is a specific phenomenon, we evaluated the role of

another gut resident microbe, *Enterococcus faecalis* on *Salmonella* intracellular replication in

human intestinal epithelial cells and *Salmonella* biofilm formation. *Enterococcus sp.* belong to

the phylum of Firmicutes in the family Enterococcaceae. Approximately, 10^6-10^7 *Enterococcus*

sp. reside in the human intestine with major abundance of *E. faecalis* (10^5-10^7 CFU/gm feces)

and *E. faecium* (10^4-10^5/gm feces) (38). *E. faecalis*, *E. faecium*, *Bifidobacterium sp.*

Lactobacillus sp., and *E. coli* colonize the gut of healthy breastfed infants during the initial 7-

10 days post birth (39). Our intracellular survival assay data showed no defect in *Salmonella*

invasion or intracellular replication capability in Caco-2 cells in presence or absence of *E.*

faecalis, its culture supernatant or PFA-fixed *E. faecalis* (Fig. S3, A-B). Moreover, *E. faecalis*

or its *E. faecalis* supernatant did not inhibit *Salmonella* biofilm formation (Fig. S3, C).

Together, our results suggest the inability of *E. faecalis* to inhibit *Salmonella* invasion,

intracellular proliferation or its *in vitro* biofilm formation, thereby highlighting the specificity

of OS in restricting *Salmonella* pathogenesis.

O. splanchnicus* confers protection against another foodborne pathogen *Listeria

monocytogenes

To understand whether OS exerts a common broad-spectrum protection or colonization

resistance against other foodborne pathogens such as *Listeria monocytogenes*, we orally

infected 6-8-week-old C57Bl/6 mice with 10^8 CFU of *L. monocytogenes* (LM) 1 day post

colonization with 10^5 CFU of *O. splanchnicus* for organ burden evaluation via oral gavage. For

survival study, mice were orally gavaged with 10^9 CFU of *L. monocytogenes*. OS-colonized

mice exhibited increased survival rates upon LM challenge compared to the non-colonized

mice cohort (Fig. 5, A). In line with the survival data, the increased bacterial burden was

observed in the blood, feces, intestinal tissue, MLN, spleen and liver alongside enhanced

percent loss in mice body weight in the case of *Listeria* infection in non-colonized mice group

with respect to OS-precolonized- mice group (Fig. 5, B-J). In a similar line to *Salmonella*

infection, OS mitigated host antimicrobial response with reduced expression of antimicrobial

peptide transcripts during LM infection, amounting to reduced intestinal tissue damage.

Further, OS promoted tight junction functions during *Listeria* infection with reduced

expression of GVB damage marker, *PV-1* and angiogenesis marker, *Vegf* expression (Fig. S4,

A-C). Moreover, OS colonization could restrict the *in vivo Salmonella* and *Listeria* biofilm-

like microcolonies within the spleens of the infected mice cohorts as opposed to the non-

colonized mice cohorts (Fig.S4, D-E). We estimated the communities of bacterial cells within

the spleen tissue via Calcofluor white staining (40). Calcofluor white stains the biofilm matrix

or extracellular polymeric substances (EPS) comprised of cellulose (40, 41) and

exopolysaccharides that help in the adherence of the bacterial cells. This is the first study

demonstrating biofilm-like bacterial colonies of *Salmonella* or *Listeria* secreting EPS in

spleens. Like *Salmonella*, *in vitro* culture of *Listeria* with OS or its culture supernatant led to a

loss in flagella as evidenced via the TEM imaging (Fig.S4, F). Since flagella is one of the

factors facilitating biofilm formation and invasion, OS could inhibit both *Salmonella* and

Listeria colonisation and biofilm formation. Further, LM exhibited attenuated intracellular

proliferation within human intestinal Caco-2 cells upon live OS or OS supernatant treatment

(Fig. 6). Altogether, this result hints at the potency of OS to act broadly against Gram-positive

and Gram-negative bacteria.

The protective role of *O. splanchnicus* against *Salmonella* Typhimurium and *Listeria* monocytogenes persists even on administration post-establishment of infection

Next, we sought to examine whether the protective role of OS against *Salmonella* and *Listeria*

is sustained when the infection has been fully developed and whether it can serve as a treatment

regimen against foodborne diseases. Our results indicate the protective ability of OS against

both foodborne pathogens in their intestinal phase of infection when administered at 1-day

post-infection (dpi) as depicted by lower pathogen loads in all the organs and blood along with

reduced percent weight loss at 5th-day post-*Salmonella* or *Listeria* challenge in the OS-

administered (1-dpi) cohort when compared to the untreated cohorts (Fig. 7,A-K, Fig. S5, A-

C). Although the protective ability of OS persists even upon establishment of infection when

administered at 3rd-day post-*Salmonella* or *Listeria* challenge as indicated by reduced organ

loads and percent weight loss in the OS-colonized cohorts, the extent of protection is lower

compared to its administration at 1-dpi (Fig. 7, L-W, Fig. S5, D-S), Further, the OS challenge

at 1-dpi reduced the mice's susceptibility to the *Salmonella* and *Listeria* challenge(Fig. 7, L,R).

However, administration of OS at 3-dpi delayed the death of the *Salmonella* and *Listeria*-

infected mice cohorts by 1 day (Fig. 7, L,R). In line with the survival and organ burden data,

OS treatment at both 1st and 3rd dpi reduced both *Salmonella* and *Listeria*-induced GVB

damage with reduced PV-1 expression, AMPs*, and *Vegf* expression, alongside promoting tight

junction protein (*Tjp1* and *Claudin5*) and *Axin2* expression (Fig. S5, D-M). However, OS

administration at 3-dpi conferred less protection than at 1-dpi as indicated by the protein

expression profiles of PV-1, and tight junction protein ZO-1 (Fig. S5, N-O). Additionally, OS

colonization mitigated gut vascular permeability during *Salmonella* and *Listeria* challenges as

indicated by reduced FITC-dextran fluorescence units in the plasma both upon pre or post-OS

colonization (Fig. S5, P). Moreover, the hematoxylin- eosin-stained intestinal sections revealed

reduced intestinal tissue damage with restoration in the villi architecture in the 1- and 3-dpi

OS-administered mice cohorts compared to the non-OS-treated groups (Fig.S5, Q).

Moreover, OS colonization could restrict the *in vivo Salmonella* and *Listeria* biofilm within the

spleens of the infected mice cohorts as opposed to the non-colonized mice cohorts both when

administered at 1-day and 3-day post-pathogen challenge. (Fig.S5, R-S).

Characterization of the inhibitory active molecule within *O. splanchnicus* culture

supernatant

Our results suggested that not only live OS but also its spent culture supernatant could inhibit

Salmonella invasion and intracellular proliferation in intestinal human epithelial cells.

Moreover, the OS and its supernatant could inhibit STM biofilm formation *in vitro.* This

inhibitory effect disappeared upon co-culture with paraformaldehyde-fixed-OS indicating the

presumable role of certain inhibitory microbial products secreted by metabolically live OS. To

delineate the nature of the secreted molecule/s, we subjected the supernatant to various

conditions such as high temperature, divalent cationic chelator, EDTA, and protease (proteinase

K), and subsequently assessed the biofilm formation (Fig. 8, A-C). Our biofilm assay suggested

that the active molecule is resistant to EDTA treatment and is stable at higher temperatures

(Fig. 8, A-B). Nevertheless, the active molecules within the fraction were sensitive to the

proteinase K treatment, suggesting the possibility of being protein in nature (Fig. 8, C). SDS-PAGE of the OS supernatant revealed the size of the differential proteins with respect to the media control ranged below 65kDa. (Fig. 8, D). Concentration-dependent estimation of the biofilm activity of the supernatant revealed that the active molecules retain their activity at a minimal concentration of 0.4% v/v (Fig. 8, E). We further fractionated the supernatant using an Amicon 3k MWCO and 10k MWCO ultra filter device (Fig. 8, F-G). We found that the active molecule/s is/ are higher than 3kDa cut-off range eliminating the role of small molecules and ions (Fig. 8, F-G). A summary of the conditions tested for active molecule characterisation and their effective outcome has been summarized in Fig. 8, H. Subsequently, we subjected the culture supernatant to gel filtration chromatography (Fig. S6, A). Out of all the fractions tested, around 24 fractions showed a biofilm inhibitory role. These 24 fractions were further subjected to a titration experiment to zero down upon the three fractions with the highest inhibitory activity (A82,A83,A84) (S6, B-D). The gel filtration standard curve run suggested that the active molecules fall in the size range of <60kDa and >3kDa (Fig. S6, A). These three fractions along with the total concentrated supernatant (serving as positive control) and fractions with no or least inhibitory activity (serving as negative control-A31,B12) were ultimately subjected to Reverse Phase LC-MS mass spectrometry. Upon analysis of the proteomics dataset (Fig. S6, E), we obtained 364 proteins showing enrichment in the positive fractions. Among the 364 proteins, 74 proteins exhibited known or predicted function as per UniProt and STRING database and 11 proteins were linked to antibacterial, bacteriostatic, flagella, and biofilm regulation based on data curation and literature survey (Table-S1). We selected bacteriocin proteins exhibiting direct association with antibacterial activity for further analysis. These proteins include bacteriocins (A0A412WSU3 and A0A412WSS0) which are antimicrobial peptides that can kill or inhibit closely related or unrelated bacterial strains(42, 43). We purified the bacteriocins via ammonium sulphate precipitation followed by gel filtration

chromatography to obtain a single fraction comprising of bacteriocins with close-range

molecular weight (6.6 kDa and 7.156kDa, respectively) (Fig. S7).

O. splanchnicus*-derived purified peptide, bacteriocin, limits both *Salmonella* and *Listeria

pathogenesis *in vitro*

Treatment of the Caco-2 cells with the purified bacteriocin inhibited STM, STY, and LM

invasion and fold proliferation without compromising host-cell viability (Fig. 9, A-D).

Additionally, the purified bacteriocin treatment inhibited STM biofilm formation in a dose-

dependent manner alongside reduced transcript level expression of *Salmonella* biofilm

regulatory genes (Fig. S8, A-D). Moreover, bacteriocin treatment (10µg/ml) attenuated

Salmonella virulence gene expression such as SPI-1 gene- *sipC*, SPI-2 gene-*pipB2,* and

flagellar gene *fliC* (Fig. S8, E-G). Subsequently, TEM imaging revealed a complete loss of

Listeria flagella upon treatment with the OS-derived bacteriocin in contrast to *Salmonella*

which retained few flagella even on bacteriocin treatment (Fig. 9, E-H). Furthermore, the

motility assay demonstrated a significant swimming defect of *Listeria* and *Salmonella* upon

treatment with either OS supernatant or bacteriocin to the untreated control (Fig. 9, I-J).

Overall, our study depicts the novel role of OS-derived bacteriocin in limiting *Salmonella* and

Listeria invasion and pathogenesis by regulating virulence effectors such as SPI-1, SPI-2

genes, biofilm regulatory genes, and flagella.

DISCUSSION

Salmonella gains access to the systemic sites by traversing the gut intestinal barrier via utilizing

several routes. Its primary site of entry is the distal ileum wherein it induces the SPI-1 Type

Three Secretion System (T3SS) effector proteins to facilitate its uptake. After its initial entry

into the enterocytes and M-cells overlying the Peyer's patches, *Salmonella* colonizes the

draining MLN and eventually disseminates to the liver and spleen via the circulating

lymphatics and the bloodstream (44). *Salmonella* uptake by M cells facilitates its stealthy entry via avoidance of the mucus barrier. Additionally, the presence of flagella allows them to propel through the viscous mucus and activate the immune responses(45). The onset of systemic dissemination coincides with GVB disruption and loss of tight-junction protein expression (1). Several reports have suggested the protective role of gut commensals in maintaining gut epithelial health against food-borne pathogen insults. *E. coli* provide colonization resistance against STM in the presence of a specific microbial consortium by limiting galactitol wherein *Lachnospiraceae* consume the free sugar, conferring colonization resistance (46). Moreover, *E. coli* contribute to colonization resistance by competing with *S.* Enteritidis for aerobic respiration (47). Similarly, the commensal *Klebsiella michiganensis* retention or transmission provided colonization resistance to Enterobacteriaceae by exhibiting nutrient competition (48).

Previous literature hinted toward the beneficial effects of OS in healthy human gut wherein reduced OS abundance have been linked to obese gut signature (49). Furthermore, OS colonization in mice have been shown to enhance Foxp3$^+$/RORγt$^+$ regulatory T cells, interleukin IL-10 and short chain fatty acids production, limiting colitis in mouse models (50). OS have been depicted to confer protection against colitis and colorectal cancer in mice by inducing intestinal Th17 development(13).

In this study, we report the broad-spectrum protective role of the gut commensal of bacteroidales order *O. splanchnicus* that limits *Salmonella* and *Listeria* invasion, pathogenesis, and systemic dissemination by promoting colonization resistance and regulating virulence effectors such as SPI-1, SPI-2, flagella, and biofilm regulatory genes. Additionally*,* not only OS but its culture supernatant could exert its inhibitory effect on *Salmonella* invasion and intracellular proliferation in intestinal human epithelial cells. Moreover, the OS and its supernatant could inhibit STM biofilm formation by regulating flagella formation. This STM biofilm inhibitory effect was however lost upon co-culture with paraformaldehyde-fixed-OS

indicating the presumable role of certain inhibitory microbial products secreted by

metabolically live OS. Further, our studies identified OS-produced bacteriocins responsible for

restricting intracellular replication of *Salmonella* and *Listeria* in intestinal epithelial cells by

regulating bacterial virulence arsenals such as biofilm regulatory genes, SPI-1, SPI-2 genes,

and flagellar machinery. However, the presence of flagella on *Salmonella* post-bacteriocin

treatment suggests the potential involvement of additional secreted accessory molecules within

the OS supernatant and indicates that bacteriocins might act in concert with additional factors

through diverse mechanisms, warranting future research. Bacteriocin-mediated suppression of

several virulence-associated phenotypes with minimal impact on overall bacterial growth

suggests that the bacteriocin may interfere with regulatory pathways controlling virulence. One

possible mechanism is the modulation of global regulatory networks that coordinate virulence

gene expression, potentially through subtle physiological perturbations or cellular signalling

pathways. Such physiological changes could attenuate virulence programs including motility

required for host cell invasion and intracellular survival. Alternatively, bacteriocin exposure

may induce stress responses that indirectly downregulate energy-intensive virulence processes.

However, determining the precise molecular mechanisms underlying these effects will require

dedicated mechanistic studies, such as transcriptomic, proteomic, or regulatory pathway

analyses, which are beyond the scope of the present study but warrant further investigation in

future work.

Although the bacteriocin demonstrated significant anti-virulence activity in vitro, its

proteinaceous nature presents potential challenges for oral therapeutic applications due to

susceptibility to degradation in the gastrointestinal environment. Consequently, oral

administration experiments in mice were not pursued in this study. Future work should

therefore explore strategies to enhance stability and delivery, such as encapsulation and

protective formulations. Addressing these challenges will be important for translating

bacteriocin-based approaches into practical therapeutic applications.

Previous studies have suggested that bacteriocin expression by commensal bacteria can

influence gut niche competition and bacteriocin delivery can serve as an effective therapeutic

strategy to eliminate colonization of multi-drug resistant bacteria in the gastrointestinal tract

without disrupting the indigenous microbiota (51, 52). Several studies have highlighted the role

of commensal-produced bacteriocins. Bacteriocin-producing *Streptococcus salivarius* has been

shown to effectively target the CRC-causing pathogen *Fusobacterium nucleatum* in a model of

the human distal colon(53). *Lactobacillus plantarum* bacteriocin has been linked to intestinal

and systemic improvements in diet-induced obese mice and confers epithelial barrier integrity

(54). Further studies show that a class IIa bacteriocin, Pediocin PA-1, has an anti-bacterial

effect against *L. monocytogenes* (55).

Bacteriocins produced by commensal bacteria are typically characterized by strong bactericidal

activity that directly inhibits the growth of competing microorganisms. In contrast, the OS-

derived bacteriocin described in this study primarily suppressed virulence-associated

phenotypes including motility, SPI-1 and SPI-2 gene expression, and intracellular proliferation

without markedly impairing bacterial growth. This observation suggests that, rather than acting

solely as a classical antimicrobial compound, the OS-bacteriocin may interfere with regulatory

pathways controlling virulence. One possible mechanism is the modulation of global regulatory

networks that coordinate virulence gene expression, potentially through subtle physiological

perturbations or cellular signalling pathways. Alternatively, bacteriocin exposure may induce

stress responses that indirectly downregulate energy-intensive virulence processes. However,

determining the precise molecular mechanisms underlying these effects will require dedicated

mechanistic studies, such as transcriptomic, proteomic, or regulatory pathway analyses, which

are beyond the scope of the present study but warrant further investigation in the future. OS-

derived bacteriocin's virulence-modulating activity is conceptually distinct from many

previously described commensal-derived bacteriocins that mainly exert growth-inhibitory

effects. The ability to attenuate virulence while exerting limited pressure on bacterial viability

may represent a potentially advantageous strategy, as anti-virulence approaches are thought to

impose reduced selective pressure for resistance compared with conventional bactericidal

agents. Nevertheless, further mechanistic studies will be required to determine whether this

effect arises from direct interference with virulence regulatory networks or from broader

physiological perturbations induced by bacteriocin exposure.

This study provides evidence about inter-species regulation of gut colonization in maintaining

gut homeostasis and protecting against several invading foodborne pathogens. Moreover, the

long-term protective ability conferred by OS during pre-or post-infection colonization to mice

against broad-spectrum foodborne pathogens such as *Salmonella* and *Listeria* highlights its

therapeutic potential as a prebiotic for prophylaxis or post-infection treatment option, paving

the way for future therapeutic interventions to thwart food-borne infections.

ACKNOWLEDGEMENTS
We thank Nikhita Kirthivasan for her support during the commencement of the project.
We also thank our Divisional Mass Spectrometry facility, TEM facility, Departmental
Confocal Facility, Divisional Flowcytometry Facility and Central Animal Facility at IISc.
FUNDING
This work was supported by the DAE SRC fellowship (DAE00195) and DBT-IISc partnership
umbrella program for advanced research in biological sciences and Bioengineering to DC.
Infrastructure support from ICMR (Centre for Advanced Study in Molecular Medicine), DST
(FIST), UGC (special assistance) and TATA fellowship is highly acknowledged.
DH sincerely acknowledges the CSIR- SPM fellowship for her financial support.
DM and RV are supported by MHRD-GATE and CSIR fellowships respectively.
The funders had no role in study design, data collection and analysis, decision to publish, or
preparation of the manuscript.

AUTHOR CONTRIBUTION

Conceptualization: DH, DC

Methodology: DH, DM, RV, VD (Gel Filtration Chromatography), DD (Fast Performance Liquid Chromatography)

Investigation: DH, DM, RV, VD (Gel Filtration Chromatography), DD (Size exclusion chromatography for bacteriocin purification), RSR (Scoring of haemoxylin and eosin-stained samples)

Visualization: DH, DM, RV

Formal analysis: DH

Funding acquisition: DC

Project administration: DH, DC

Supervision: DC, UT (Proteomics), TH (Size Exclusion Chromatography),

MG (Size Exclusion Chromatography)

Writing – original draft: DH

Writing – review & editing: DH, DM, RV, RSR, VD, DD, TH, MG, UT, DC

COMPETING INTERESTS Authors declare that they have no competing interests.

DATA AND MATERIALS AVAILABILITY

All data are available in the main text or the supplementary materials.

The article has been submitted to the preprint server (56).

REFERENCES

1.   Spadoni I, Zagato E, Bertocchi A, Paolinelli R, Hot E, Di Sabatino A, et al. A gut-vascular barrier controls the systemic dissemination of bacteria. Science (New York, NY). 2015;350(6262):830-4.

2.   Brescia P, Rescigno M. The gut vascular barrier: a new player in the gut–liver–brain axis. Trends in Molecular Medicine. 2021;27(9):844-55.

3.   Ghosh S, Whitley CS, Haribabu B, Jala VR. Regulation of Intestinal Barrier Function by Microbial Metabolites. Cellular and Molecular Gastroenterology and Hepatology. 2021;11(5):1463-82.

4.   Zhao Y, Chen F, Wu W, Sun M, Bilotta AJ, Yao S, et al. GPR43 mediates microbiota metabolite SCFA regulation of antimicrobial peptide expression in intestinal epithelial cells via activation of mTOR and STAT3. Mucosal immunology. 2018;11(3):752-62.

5.   Schulthess J, Pandey S, Capitani M, Rue-Albrecht KC, Arnold I, Franchini F, et al. The short chain fatty acid butyrate imprints an antimicrobial program in macrophages. Immunity. 2019;50(2):432-45. e7.

6.   Hajra D, Kirthivasan N, Chakravortty D. Symbiotic Synergy from Sponges to Humans: Microflora-Host Harmony Is Crucial for Ensuring Survival and Shielding against Invading Pathogens. ACS infectious diseases. 2024;10(2):317-36.

7.   Garcia-del Portillo F, Foster JW, Finlay BB. Role of acid tolerance response genes in Salmonella typhimurium virulence. Infection and immunity. 1993;61(10):4489-92.

8.   Haraga A, Ohlson MB, Miller SI. Salmonellae interplay with host cells. Nat Rev Microbiol. 2008;6(1):53-66.

9.   Hensel M. Evolution of pathogenicity islands of Salmonella enterica. International journal of medical microbiology : IJMM. 2004;294(2-3):95-102.

10.   Hajra D. Understanding the mechanism of host deacetylase SIRT1 and SIRT3 in the modulation of Salmonella pathogenesis. 2024.

11.   Wang J, Lang T, Shen J, Dai J, Tian L, Wang X. Core Gut Bacteria Analysis of Healthy Mice. Frontiers in microbiology. 2019;10:887.

12.   Hiippala K, Barreto G, Burrello C, Diaz-Basabe A, Suutarinen M, Kainulainen V, et al. Novel
Odoribacter splanchnicus Strain and Its Outer Membrane Vesicles Exert Immunoregulatory Effects in
vitro. Frontiers in microbiology. 2020;11:575455.
13.   Xing C, Wang M, Ajibade AA, Tan P, Fu C, Chen L, et al. Microbiota regulate innate immune
signaling and protective immunity against cancer. Cell host & microbe. 2021;29(6):959-74.e7.
14.   Gogoi M, Chandra K, Sarikhani M, Ramani R, Sundaresan NR, Chakravortty D. Salmonella
escapes adaptive immune response via SIRT2 mediated modulation of innate immune response in
dendritic cells. PLoS pathogens. 2018;14(11):e1007437.
15.   Schaefer JS, Montufar-Solis D, Nakra N, Vigneswaran N, Klein JR. Small intestine inflammation
in Roquin-mutant and Roquin-deficient mice. PloS one. 2013;8(2):e56436.
16.   Erben U, Loddenkemper C, Doerfel K, Spieckermann S, Haller D, Heimesaat MM, et al. A
guide to histomorphological evaluation of intestinal inflammation in mouse models. International
journal of clinical and experimental pathology. 2014;7(8):4557-76.
17.   Hajra D, Nair AV, Roy Chowdhury A, Mukherjee S, Chatterjee R, Chakravortty D. Salmonella
Typhimurium U32 peptidase, YdcP, promotes bacterial survival by conferring protection against in
vitro and in vivo oxidative stress. Microbial pathogenesis. 2022;173(Pt B):105862.
18.   Roy Chowdhury A, Hajra D, Mukherjee D, Nair AV, Chakravortty D. Functional OmpA of
Salmonella Typhimurium provides protection from lysosomal degradation and inhibits autophagic
processes in macrophages. The Journal of infectious diseases. 2024.
19.   Chandra K, Nair AV, Chatterjee R, Muralidhara P, Singh A, Kamanna S, et al. Absence of
proline-peptide transporter YjiY in Salmonella Typhimurium leads to secretion of factors which
inhibits intra-species biofilm formation. Microbiological research. 2023;273:127411.
20.   Hajra D, Rajmani RS, Chaudhary AD, Gupta SK, Chakravortty D. Salmonella-induced SIRT1 and
SIRT3 are crucial for maintaining the metabolic switch in bacteria and host for successful
pathogenesis. eLife Sciences Publications, Ltd; 2024.
21.   Balakrishnan A, Schnare M, Chakravortty D. Of men not mice: bactericidal/permeability-
increasing protein expressed in human macrophages acts as a phagocytic receptor and modulates
entry and replication of Gram-negative bacteria. Frontiers in immunology. 2016;7:455.
22.   Chatterjee R, Chowdhury AR, Nair AV, Hajra D, Kar A, Datey A, et al. Salmonella Typhimurium
PgtE is an essential arsenal to defend against the host resident antimicrobial peptides.
Microbiological research. 2023;271:127351.
23.   Goubier A, Dubois B, Gheit H, Joubert G, Villard-Truc F, Asselin-Paturel C, et al. Plasmacytoid
Dendritic Cells Mediate Oral Tolerance. Immunity. 2008;29(3):464-75.
24.   Müller-Redetzky HC, Suttorp N, Witzenrath M. Dynamics of pulmonary endothelial barrier
function in acute inflammation: mechanisms and therapeutic perspectives. Cell and tissue research.
2014;355(3):657-73.
25.   Daneman R, Rescigno M. The gut immune barrier and the blood-brain barrier: are they so
different? Immunity. 2009;31(5):722-35.
26.   Yuan SY, Rigor RR. Integrated Systems Physiology: From Molecule to Function to Disease.
Regulation of Endothelial Barrier Function. San Rafael (CA): Morgan & Claypool Life Sciences

Copyright © 2011 by Morgan & Claypool Life Sciences.; 2010.
27.   Wang Y, Rattner A, Zhou Y, Williams J, Smallwood PM, Nathans J. Norrin/Frizzled4 signaling in
retinal vascular development and blood brain barrier plasticity. Cell. 2012;151(6):1332-44.
28.   Liebner S, Corada M, Bangsow T, Babbage J, Taddei A, Czupalla CJ, et al. Wnt/beta-catenin
signaling controls development of the blood-brain barrier. The Journal of cell biology.
2008;183(3):409-17.
29.   Oehlers SH, Cronan MR, Scott NR, Thomas MI, Okuda KS, Walton EM, et al. Interception of
host angiogenic signalling limits mycobacterial growth. Nature. 2015;517(7536):612-5.
30.   Iwase T, Uehara Y, Shinji H, Tajima A, Seo H, Takada K, et al. Staphylococcus epidermidis Esp
inhibits Staphylococcus aureus biofilm formation and nasal colonization. Nature.
2010;465(7296):346-9.

31.   Lee J, Jayaraman A, Wood TK. Indole is an inter-species biofilm signal mediated by SdiA. BMC
microbiology. 2007;7(1):1-15.
32.   Fux CA, Costerton JW, Stewart PS, Stoodley P. Survival strategies of infectious biofilms. Trends
in microbiology. 2005;13(1):34-40.
33.   Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent
infections. Science (New York, NY). 1999;284(5418):1318-22.
34.   Hajra D, Nair AV, Chakravortty D. An elegant nano-injection machinery for sabotaging the
host: Role of Type III secretion system in virulence of different human and animal pathogenic
bacteria. Physics of life reviews. 2021;38:25-54.
35.   Jennings ME, Quick LN, Ubol N, Shrom S, Dollahon N, Wilson JW. Characterization of
Salmonella type III secretion hyper-activity which results in biofilm-like cell aggregation. PloS one.
2012;7(3):e33080.
36.   Ducarmon QR, Zwittink RD, Hornung BVH, van Schaik W, Young VB, Kuijper EJ. Gut Microbiota
and Colonization Resistance against Bacterial Enteric Infection. Microbiology and molecular biology
reviews : MMBR. 2019;83(3).
37.   Caballero-Flores G, Pickard JM, Núñez G. Microbiota-mediated colonization resistance:
mechanisms and regulation. Nature Reviews Microbiology. 2023;21(6):347-60.
38.   Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, et al. A human gut microbial
gene catalogue established by metagenomic sequencing. Nature. 2010;464(7285):59-65.
39.   Gewolb IH, Schwalbe RS, Taciak VL, Harrison TS, Panigrahi P. Stool microflora in extremely
low birthweight infants. Archives of disease in childhood Fetal and neonatal edition.
1999;80(3):F167-73.
40.   Chakraborty P, Bajeli S, Kaushal D, Radotra BD, Kumar A. Biofilm formation in the lung
contributes to virulence and drug tolerance of Mycobacterium tuberculosis. Nature communications.
2021;12(1):1606.
41.   Yamamoto K, Tsujimura Y, Ato M. Catheter-associated Mycobacterium intracellulare biofilm
infection in C3HeB/FeJ mice. Sci Rep. 2023;13(1):17148.
42.   Heilbronner S, Krismer B, Brötz-Oesterhelt H, Peschel A. The microbiome-shaping roles of
bacteriocins. Nature reviews Microbiology. 2021;19(11):726-39.
43.   Cotter PD, Ross RP, Hill C. Bacteriocins - a viable alternative to antibiotics? Nature reviews
Microbiology. 2013;11(2):95-105.
44.   Watson KG, Holden DW. Dynamics of growth and dissemination of Salmonella in vivo.
Cellular microbiology. 2010;12(10):1389-97.
45.   McGuckin MA, Lindén SK, Sutton P, Florin TH. Mucin dynamics and enteric pathogens. Nature
reviews Microbiology. 2011;9(4):265-78.
46.   Eberl C, Weiss AS, Jochum LM, Durai Raj AC, Ring D, Hussain S, et al. E. coli enhance
colonization resistance against Salmonella Typhimurium by competing for galactitol, a context-
dependent limiting carbon source. Cell host & microbe. 2021;29(11):1680-92.e7.
47.   Litvak Y, Mon KKZ, Nguyen H, Chanthavixay G, Liou M, Velazquez EM, et al. Commensal
Enterobacteriaceae Protect against Salmonella Colonization through Oxygen
Competition. Cell host & microbe. 2019;25(1):128-39.e5.
48.   Oliveira RA, Ng KM, Correia MB, Cabral V, Shi H, Sonnenburg JL, et al. Klebsiella
michiganensis transmission enhances resistance to Enterobacteriaceae gut invasion by nutrition
competition. Nat Microbiol. 2020;5(4):630-41.
49.   Chanda D, De D. Meta-analysis reveals obesity associated gut microbial alteration patterns
and reproducible contributors of functional shift. Gut Microbes. 2024;16(1):2304900.
50.   Lima SF, Gogokhia L, Viladomiu M, Chou L, Putzel G, Jin WB, et al. Transferable
Immunoglobulin A-Coated Odoribacter splanchnicus in Responders to Fecal Microbiota
Transplantation for Ulcerative Colitis Limits Colonic Inflammation. Gastroenterology.
2022;162(1):166-78.

51.   Kommineni S, Bretl DJ, Lam V, Chakraborty R, Hayward M, Simpson P, et al. Bacteriocin
production augments niche competition by enterococci in the mammalian gastrointestinal tract.
Nature. 2015;526(7575):719-22.
52.   Kommineni S, Kristich CJ, Salzman NH. Harnessing bacteriocin biology as targeted therapy in
the GI tract. Gut Microbes. 2016;7(6):512-7.
53.   Lawrence GW, McCarthy N, Walsh CJ, Kunyoshi TM, Lawton EM, O'Connor PM, et al. Effect of
a bacteriocin-producing Streptococcus salivarius on the pathogen Fusobacterium nucleatum in a
model of the human distal colon. Gut Microbes. 2022;14(1):2100203.
54.   Heeney DD, Zhai Z, Bendiks Z, Barouei J, Martinic A, Slupsky C, et al. Lactobacillus plantarum
bacteriocin is associated with intestinal and systemic improvements in diet-induced obese mice and
maintains epithelial barrier integrity in vitro. Gut Microbes. 2019;10(3):382-97.
55.   Kuniyoshi TM, O'Connor PM, Lawton E, Thapa D, Mesa-Pereira B, Abulu S, et al. An oxidation
resistant pediocin PA-1 derivative and penocin A display effective anti-Listeria activity in a model
human gut environment. Gut Microbes. 2022;14(1):2004071.
56.   Hajra D, Mukherjee D, Vij R, Rajmani RS, Dadireddy V, Das D, et al. Odoribacter splanchnicus
mitigates Salmonella-induced gut inflammation and its associated pathogenesis via its secreted
bacteriocin. bioRxiv. 2024:2024.08.23.609322.

870
871

MAIN FIGURES AND LEGENDS

Figure 1. *Odoribacter splanchnicus* **colonization protects mice from** *Salmonella*

colonization and dissemination

A-Schematic representation showing animal experimental strategy of infection.

B-OS colonization validation via qPCR quantification of *O. splanchnicus* 16S rRNA from fecal

samples of mice at 6^{th} day post-OS colonization .

C-Survival percentage of non-colonized-STM-infected mice versus OS-colonized-STM-

infected mice cohorts. Data is representative of N=3, n=5 where N denotes number of

biological replicates and n denotes the number of mice per group. Mantel Cox or the log-rank

test was used to determine the p values.

D-G- STM organ loads in feces at 1^{st} day post-infection (D), 3^{rd} day post-infection (E), 5^{th}

post-infection (F) and in ileal tissues (contents and wall) at 5^{th} day post-infection (G) in OS

colonized and non-colonized mice cohorts. Data is representative of N=2, n=6 where N

denotes number of biological replicates and n denotes the number of mice per group. Mann

Whitney Test was performed to obtain p-values.

H-Percent of initial weight (at Day 0) exhibited by OS colonized and non-colonized mice upon

STM infection at 5^{th} day post-infection. Data is representative of N=3,n=6 where N denotes

number of biological replicates and n denotes the number of mice per group.

I-K- STM organ loads in MLN (I), liver (J) and blood (K) at 5^{th} post-infection in OS colonized

and non-colonized mice cohorts. Data is representative of N=3, n=6 where N denotes number

of biological replicates and n denotes the number of mice per group. Mann Whitney Test was

performed to obtain p-values.

Figure 2. *O. splanchnicus* **reduces** *Salmonella***-induced inflammation and intestinal**
damage

A-Estimation of serum IL-6 production via ELISA upon STM infection in OS non-colonized
and colonized mice cohorts at 5th day post-*Salmonella* infection. Unpaired t-test is performed
to obtain the p-values. Data is representative of N=3, n=3.

B-Quantitative gene expression data of several antimicrobial peptides (AMPs) precursor genes

in uninfected (UI) and STM-infected mice intestinal tissues harvested at 5th dpi from OS

colonized or non-colonized cohorts. Unpaired t-test is performed to obtain the p-values. Data

is representative of N=3, n=4.

C- **D**-Haematoxylin-eosin-stained intestinal sections harvested from STM-infected OS non-

colonized or colonized mice cohorts at 5th dpi. 0-represents normal pathology, 1- represents-

mild pathology, 2-3-represents severe pathology.

Fig 3. *Odoribacter splanchnicus* gut colonization prevents *Salmonella*-induced GVB

damage and promotes tight junction gene expression.

A-Quantitative gene expression studies of tight junction protein complexes (*Tjp1*, *Claudin5*)

and PV-1 gene (marker of GVB damage) in STM-infected mice intestinal (ileal) tissue in

presence or absence of OS gut colonization at 5^{th} day post-infection. Un-paired t-test was used

to obtain the p-values. Data is representative of N=3, n=4.

B- Estimation of FITC fluorescence in mice plasma via gut permeability assay in infected mice

cohorts upon oral administration of FITC Dextran (25mg/kg) 4h before euthanization at 5^{th} day

post *Salmonella* infection. Un-paired t-test was used to obtain the p-values. Data is

representative of N=4.

Fig 4. *O. splanchnicus* exhibits its inhibitory effect on *Salmonella* biofilm formation and inhibits invasion and intracellular replication in human intestinal cells

A-Percent invasion of STM in human Caco-2 human intestinal cells in presence or absence of live OS, OS culture supernatant, PFA-fixed-OS. Data is representative of N=3, n=4. One-way ANOVA was used to determine the p-values.

B-Intracellular proliferation of STM Caco-2 cells in presence or absence of live OS, OS culture supernatant, PFA-fixed-OS. Data is representative of N=3, n=4. One-way ANOVA was used to determine the p-values.

C- Representative immunofluorescence images depicting STM invasion into Caco-2 cells in the presence or absence of OS at 2hr post-infection. Data is representative of N=3, n=100.

D- Representative immunofluorescence images depicting STM invasion into Caco-2 cells in presence or absence of OS at 2hr post-infection. Data is representative of N=3, n=100.

E-Percent invasion of STY in Caco-2 human cells in presence or absence of OS, OS culture supernatant, PFA-fixed OS. Data is representative of N=3, n$\geq$9. One-way ANOVA was used to determine the p-values.

F-Intracellular proliferation of STY in Caco-2 cells in presence or absence of live OS, OS culture supernatant, PFA-fixed-OS. Data is representative of N=3, n$\geq$9. One-way ANOVA was used to determine the p-values.

G- Biofilm assay showing the biofilm formation ability of STM-WT or STM-WT grown in OS supernatant after 72h of inoculation in presence or absence of live OS, OS culture supernatant, PFA-fixed-OS. Data is representative of N=4, n=4.One-way ANOVA was used to determine the p-values. Image of the STM biofilm-formed plate in the above stated conditions has been shown.

H-M- RT-qPCR mediated gene expression of STM biofilm regulatory genes (*csgD*) (H), *csgA* (I), *bapA* (J), along with flagellar gene (*fliC*) (K), SPI-1 (*sipC*) (L) and SPI-2 (*pipB2*) (M) genes in presence or absence of live OS or its spent media or PFA-fixed-OS. Data is representative of N=4, n=4. One-way ANOVA was used to determine the p-values.

N- Transmission Electron Microscopic (TEM) images of STM in presence or absence of *Salmonella* WT supernatant, OS and its supernatant to assess the possession of flagella. The quantification plot exhibits the percentage of bacterial flagellated or non-flagellated cells (number of cells quantified n$\geq$ 100)

Fig 5. *Odoribacter splanchnicus* **exerts its protective role against** *Listeria monocytogenes*

in mice

A- Survival percentage of non-colonized LM-infected mice versus OS-colonized *Listeria*-infected mice cohorts. Data is represantative of N=2, n=8, where N denotes number of biological replicates and n denotes the number of mice per group. Mantel Cox or the log-rank test was used to determine the p values.

B-Percent of initial weight (at Day 0) exhibited by OS colonized and non-colonized mice upon LM infection at 5th dpi. Data is representative of N=2,n=8, where N denotes number of biological replicates and n denotes the number of mice per group. Mann Whitney Test was performed to obtain p-values.

C-F- LM organ loads in feces at 1st day post-infection (C), 3rd day post-infection (D), 5th post-

infection (E) and in ileal tissues (contents and wall) at 5th day post-infection (F) in OS pre-

colonized and non-colonized mice cohorts. Data is representative of N=2, n=7 where N

denotes number of biological replicates and n denotes the number of mice per group. Mann

Whitney Test was performed to obtain p-values.

G-J- LM organ loads in Mesenteric lymph Node (MLN) (G), liver (H), spleen (I) and blood

(J) at 5th dp. in OS colonized and non-colonized mice cohorts Data is representative of N=2,

n=8, where N denotes number of biological replicates and n denotes the number of mice per

group. Mann Whitney Test was performed to obtain p-values.

Fig. 6 *Odoribacter splanchnicus* and its culture supernatant restricts *Listeria*

***monocytogenes* pathogenesis in human intestinal cells**

A- Percent invasion of LM in Caco-2 cells in presence or absence of live OS, OS sup, PFA-

fixed-OS. Data is representative of N=3, n=4. One-way ANOVA and Multiple Comparison test

are used to obtain the p-values.

B-Intracellular fold proliferation of LM in Caco-2 cells in the presence or absence of live OS,

OS sup, PFA-fixed-OS. Data is representative of N=3, n=4. One-way ANOVA and Multiple

Comparison test are used to obtain the p-values.

Fig. 7 *O. splanchnicus* **even confers protection against foodborne pathogens when administered post-pathogenic challenge**

A- Schematic representation showing the animal experimental strategy of infection.

B- Percent of initial weight (at Day 0) exhibited by OS colonized (1-dpi) and non-colonized mice upon LM infection at 5th dpi. Data is representative of N=2,n=8, where N denotes number of biological replicates and n denotes the number of mice per group. Mann Whitney Test was performed to obtain p-values.

C-F- LM organ loads in MLN (C), liver (D), spleen (E) and blood (F) at 5th dpi upon OS administration at 1-dpi. Data is representative of N=2, n=8, where N denotes number of biological replicates and n denotes the number of mice per group. Mann Whitney Test was performed to obtain p-values.

G- Percent of initial weight (at Day 0) exhibited by OS colonized (1-dpi) and non-colonized mice upon STM infection at 5th dpi. Data is representative of N=2,n=8, where N denotes number of biological replicates and n denotes the number of mice per group. Mann Whitney Test was performed to obtain p-values.

H-K- STM organ loads in MLN (H), liver (I), spleen (J) and blood (K) at 5th-dpi upon OS administration at 1-dpi. Data is representative of N=2, n=8, where N denotes number of biological replicates and n denotes the number of mice per group. Mann-Whitney Test was performed to obtain p-values.

L- Survival percentage of non-colonized LM-infected mice versus OS-colonized (1 dpi or 3 dpi) *Listeria*-infected mice cohorts. Data is representative of N=2, n=8, where N denotes number of biological replicates and n denotes the number of mice per group. Mantel Cox or the log-rank test was used to determine the p values.

M- Percent weight loss exhibited by OS colonized (3-dpi) and non-colonized mice upon LM infection at 5th-day post-infection. Data is representative of N=2,n=8, where N denotes number of biological replicates and n denotes the number of mice per group.Mann-Whitney Test was performed to obtain p-values.

N-Q- LM organ loads in MLN (N), liver (O), spleen (P) and blood (Q) at 5th-dpi upon OS administration at 3-dpi. Data is representative of N=2, n=8, , where N denotes number of biological replicates and n denotes the number of mice per group. Mann-Whitney Test was performed to obtain p-values.

R- Survival percentage of non-colonized *Salmonella*-infected mice versus OS-colonized (1 dpi or 3 dpi) *Listeria*-infected mice cohorts. Data is represantative of N=2, n=8, where N denotes

number of biological replicates and n denotes the number of mice per group. Mantel Cox or

the log-rank test was used to determine the p values.

S- Percent weight loss exhibited by OS-colonized (3-dpi) and non-colonized mice upon STM

infection at 5th-day post-infection. Data is representative of N=2,n=8, where N denotes number

of biological replicates and n denotes the number of mice per group. Mann-Whitney Test was

performed to obtain p-values.

T-W-STM organ loads in MLN (T), liver (U), spleen (V), and blood (W) at 5^{th}-dpi upon OS

administration at 3-dpi. Data is representative of N=2, n=8, where N denotes number of

biological replicates and n denotes the number of mice per group. Mann-Whitney Test was

performed to obtain p-values.

1041

1042

1043 **Fig. 8- Characterization of the nature of inhibitory active molecule within *O. splanchnicus***

1044 **culture supernatant**

A- Biofilm assay showing the biofilm formation ability of STM after 72h of inoculation in presence of OS culture supernatant upon EDTA treatment. Data is representative of N=4, n=4. One-way ANOVA is used to obtain the p-values. Statistical test is performed with respect to STM.

B- Biofilm assay showing the biofilm formation ability of STM after 72h of inoculation in presence of OS culture supernatant at high temperature conditions. Data is representative of N=4, n=4. One-way ANOVA is used to obtain the p-values. Statistical test is performed with respect to STM.

C- Biofilm assay showing the biofilm formation ability of STM after 72h of inoculation in presence of OS culture supernatant under Proteinase K treatment. Data is representative of N=4, n=4. One-way ANOVA is used to obtain the p-values. Statistical test is performed with respect to STM.

D- Coomassie-stained SDS-PAGE demonstrating the size range of the differential protein bands present in OS supernatant as opposed to the tryptic soy broth media control.

E- Biofilm assay showing the biofilm formation ability of STM after 72h of inoculation in presence of varying concentration OS culture supernatant. Data is representative of N=4, n=4. One-way ANOVA is used to obtain the p-values. Statistical test is performed with respect to STM.

F- Biofilm assay showing the biofilm formation ability of STM after 72h of inoculation in presence of OS culture supernatant fractions fractionated using 3kDa Amicon filter. Data is representative of N=4, n=4. One-way ANOVA is used to obtain the p-values. Statistical test is performed with respect to STM.

G- Biofilm assay showing the biofilm formation ability of STM after 72h of inoculation in

presence of OS culture supernatant fractions fractionated using 10kDa Amicon filter. Data is

representative of N=4, n=4. One-way ANOVA is used to obtain the p-values. Statistical test is

performed with respect to STM.

H- Summary table demonstrating the conditions tested for active molecule characterization and

their effective outcome.

1074

1075

1076

Fig. 9- *O. splanchnicus* **bacteriocin restricts both** *Salmonella* **and** *Listeria* **pathogenesis** *in*

vitro

A- Percent invasion and intracellular fold proliferation of STM in Caco-2 cells under OS

bacteriocin (10µg/ml) treatment. Data is representative of N=2, n=3. Unpaired t-test is used to

obtain the p-values.

B- Percent invasion and intracellular fold proliferation of LM in Caco-2 cells under OS

bacteriocin (10µg/ml) treatment. Data is representative of N=2, n=3. Unpaired t-test is used to

obtain the p-values.

C- Percent invasion and intracellular fold proliferation of STY in Caco-2 cells under OS

bacteriocin (10µg/ml) treatment. Data is representative of N=2, n=3. Unpaired t-test is used to

obtain the p-values.

D-MTT assay evaluating the viability of Caco-2 cells upon 16h of mock or bacteriocin

(10µg/ml) treatment. Data is representative of N=2, n=4.

E-F-TEM images of STM under untreated (D) or bacteriocin (10µg/ml) (E) treatment to assess

the possession of flagella.

G-H- TEM images of LM under untreated (D) or bacteriocin (10µg/ml) (E) treatment to assess

the possession of flagella.

I- Swimming motility assay of STM under OS supernatant or bacteriocin treatment.

J- Swimming motility assay of LM under OS supernatant or bacteriocin treatment.

1096

GRAPHICAL ABSTRACT

1097

1098

1099

1100

SUPPLEMENTARY FIGURES AND LEGENDS

Fig S1. *O. splanchnicus* **gut colonization prevents** *Salmonella* **dissemination by restricting**

intestinal GVB damage and angiogenesis

A- qPCR mediated gene expression of pro-angiogenic marker *Vegf* in STM-infected mice liver

tissues in presence or absence of OS pre-colonization. Data is representative of N=2, n=4. Un-

paired t-test was used to obtain the p-values.

B- qPCR mediated gene expression of *Axin2* in STM-infected mice intestinal tissues in the

presence or absence of OS pre-colonization. Data is representative of N=2, n=4. Un-paired t-

test was used to obtain the p-values.

1113

Fig. S2 In vitro growth kinetics of *Salmonella* Typhimurium in presence of live or dead

***O. splanchnicus* or its culture supernatant**

A-B Growth kinetics of STM-WT, OS, STM-WT in presence of OS, STM-WT in presence of

STM or OS culture supernatant and STM-WT in presence of PFA-fixed OS in LB under

anaerobic growth conditions.

C-D- Growth kinetics of STM-WT, OS, STM-WT in presence of OS, STM-WT in presence of

STM or OS culture supernatant, STM-WT supernatant, and STM-WT in presence of PFA-fixed

OS in LB under aerobic growth conditions.

1122

Fig S3. Unlike *O. splanchnicus, E. faecalis* fail to exert any inhibitory effect on *Salmonella in vitro* biofilm formation or *Salmonella* invasion and intracellular replication in human intestinal cells

A-Percent invasion of STM in Caco-2 cells in presence or absence of live *E. faecalis* (Ef), Ef culture supernatant, PFA-fixed-Ef. Data is representative of N=3, n=3. One-way ANOVA is used to obtain the p-values.

B-Intracellular proliferation of STM in Caco-2 cells in presence or absence of live Ef, Ef culture supernatant, PFA-fixed-Ef. Data is representative of N=3, n=3. One-way ANOVA is used to obtain the p-values.

C- Biofilm assay depicting the biofilm formation ability of STM-WT or STM-WT grown in Ef supernatant after 72h of inoculation in presence or absence of live Ef, Ef or STM culture supernatant, PFA-fixed. Crystal-violet staining was performed to assess the biofilm formation on the solid-liquid interphase and the $[OD]_{595}$ absorbance reading was recorded. Data is representative of N=3, n=3. One-way Anova is used to obtain the p-values.

1137

1138

1141

1142

Fig. S4. *Odoribacter splanchnicus* **gut colonization protects mice against** *Listeria*
monocytogenes **pathogenesis similar to** *Salmonella* **infection**

A-Haematoxylin-eosin-stained intestinal sections harvested from OS non-colonized or colonized mice cohorts at 5^{th}-day-post-LM-infection. 0-represents normal pathology, 1-represents- mild pathology, 2-3-represents severe pathology.

B-Quantitative gene expression data of several antimicrobial peptides (AMPs) precursor genes in LM-infected mice intestinal tissues harvested at 5^{th} day post-infection from OS colonized or non-colonized cohorts. Unpaired Student's t-test is performed to obtain the p-values. Data is representative of N=2, n=9.

C- Quantitative gene expression study of tight junction protein genes *Tjp1*, *Claudin5*, GVB damage marker *PV-1*, angiogenesis marker-*Vegf,* and *Axin2* in LM- infected mice intestinal tissues harvested at 5^{th}-dpi from OS colonized or non-colonized cohorts. Unpaired Student's t-test is performed to obtain the p-values. Data is representative of N=2, n=5.

D-Representative Calcofluor white (CW) staining of STM-infected mice spleen tissues at 5^{th}-dpi. Data is representative of N=2, n=8 (tissue samples). Un-paired t-test was used to obtain the p-values.

E- Representative CW staining of LM-infected mice spleen tissues at 5^{th}-dpi. Data is representative of N=2, n=8 (tissue samples). Un-paired t-test was used to obtain the p-values.

F- Transmission Electron Microscopic (TEM) images of LM in presence or absence of LM supernatant, OS and its supernatant to assess the possession of flagella. The quantification plot exhibits the percentage of bacterial flagellated or non-flagellated cells (number of cells quantified n$\geq$ 100)

1165

1166

1167

1168

1169

1170

1172

1173

Fig. S5 *Odoribacter splanchnicus* administration post-establishment of infection protected mice against *Listeria monocytogenes* and *Salmonella* Typhimurium

A-C- STM and LM organ loads in feces at 3rd day post-infection (A), 5th post-infection (B) and in ileal tissues (contents and wall) at 5th day post-infection (C) upon OS administration at 1-dpi or 3-dpi. Data is representative of N=2, n=6 where N denotes number of biological replicates and n denotes the number of mice per group. Mann-Whitney Test was performed to obtain p-values.

D-H-Quantitative gene expression data of several antimicrobial peptides (AMPs) precursor genes in LM and STM-infected mice intestinal tissues harvested at 5th dpi upon OS colonization

at 1-dpi or 3-dpi. Unpaired Student's t-test is performed to obtain the p-values. Data is

representative of N=2, n=4.

I-M- Quantitative gene expression study of tight junction protein genes *Tjp1*, *Claudin5*, GVB

damage marker *PV-1*, angiogenesis marker-*Vegf,* and *Axin2* in LM and STM-infected mice

intestinal tissues harvested at 5^{th}-dpi upon OS colonization at 1-dpi or 3-dpi. Unpaired

Student's t-test is performed to obtain the p-values. Data is representative of N=2, n=6.

N-O- Immunoblotting representing the protein expression profile of ZO-1, and PV-1 in

Salmonella (N) or *Listeria* (O)-infected mice intestinal tissue harvested on 5^{th}-dpi with or

without OS pre-or post-colonization (1-dpi or 3-dpi). Data is representative of N=3.

P- Estimation of FITC fluorescence in mice plasma via gut permeability assay in *Salmonella*

or *Listeria*-infected mice cohorts with or without OS pre-or post-colonization (1dpi or 3dpi)

upon oral administration of FITC Dextran (25mg/kg) 4h before euthanization at 5^{th}-dpi.

Unpaired Student's t-test is performed to obtain the p-values. Data is representative of N=2,

n=4 (UI and OS, n=2).

Q- Haematoxylin-eosin-stained intestinal sections harvested at 5^{th}-dpi from Uninfected (UI) or

Salmonella or *Listeria*-infected mice cohorts having non-administered or administered OS (at

1^{st} or 3^{rd} dpi). 0-represents normal pathology, 1- represents- mild pathology, 2-3-represents

severe pathology. Data is representative of N=2, n=6.

R- Representative Calcofluor white (CW) staining of STM-infected mice spleen tissues at 5^{th}-

day post-infection with or without OS administration at 1^{st} or 3^{rd} dpi depicting the

quantification of the length of the fibril architecture of CW-stained sections. One-way ANOVA

and multiple comparison test were used to obtain the p-values.

S- Representative CW staining of LM-infected mice spleen tissues at 5^{th}-dpi with or without

OS administration at 1^{st} or 3^{rd} dpi depicting the quantification of the length of the fibril

architecture of CW-stained sections. One-way ANOVA and multiple comparison test were used

to obtain the p-values.

1209

1210

1211

1212

1213

Fig. S6 Characterization of the inhibitory active molecule within *O. splanchnicus* culture

supernatant

A-Gel filtration chromatogram of protein standards and the OS culture supernatant. Red shaded

region shows the fractions with biofilm inhibitory activity. The dark shaded region indicates

the fractions with the highest inhibitory activity.

B-C- Biofilm assay showing the biofilm formation ability of STM after 72h of inoculation in

presence of OS supernatant gel filtration fractions. Data is representative of N=2, n=2. One-

way ANOVA is used to obtain the p-values.

D- Biofilm assay showing the biofilm formation ability of STM after 72h of inoculation in

presence of varying concentration of OS supernatant biofilm inhibitory gel filtration fractions.

Data is representative of N=2, n=1.

E-Schematic representation of the workflow of mass spectrometric analysis of the gel filtration

fractions (A82, A83, A84 and Concentrated supernatant).

1227

1228   **Fig. S7 Purification procedure of *O. splanchnicus* produced bacteriocin via ammonium**

1229   **sulphate precipitation method followed by Fast Performance Liquid Chromatography**

1230   **A**-Size exclusion chromatogram of OS-produced bacteriocin fractions post 95-98%

1231   ammonium sulphate precipitation.

 B- Coomassie-stained SDS PAGE showing the purified and concentrated OS bacteriocin.

1233

1234

Fig. S8 *O. splanchnicus* bacteriocin attenuates *Salmonella* pathogenesis *in vitro*

A- Biofilm assay showing the biofilm formation ability of STM after 72h of inoculation under varying concentration of OS bacteriocin. Data is representative of N=2, n=2. Unpaired t-test is used to obtain the p-values.

B-G- RT-qPCR mediated gene expression of STM biofilm regulatory genes *csgD* (B), *csgA*(C), *bap* (D), along with flagellar gene *fliC*(E), SPI-1-*sipC* (F) and SPI-2-*pipB2*(G) in presence or absence of OS bacteriocin treatment. Data is representative of N=2, n≥5. Unpaired Student's t-test was used to obtain the p-values.

1243

1244

1245

Table S1- Selected List of 11 proteins from Mass Spectrometric Analysis based on bibliographic search

Accession ID	Protein Name and	Referenced Function
A0A412WBT8	Immunity protein Imm5 domain-containing protein	Uncharacterized
F9Z8G5	Peptidase	Modulate biofilm formation and host defense.
R6F650	Carboxyl-terminal protease	Critical for post-translational processing and associated with regulation of gene expression, stress response, maintenance of cell envelope integrity and bacterial virulence.
A0A412TJE3	ATP-dependent Clp protease proteolytic subunit	Influence virulence and infectivity of pathogenic bacteria. ClpP enhance biofilm formation in *Enterococcus faecalis*.
F9Z604	Peptidase C39 bacteriocin processing	The cysteine peptidase cleave the double glycine leader peptides from the precursors of bacteriocin
A0A412TJV3	UDP-2,4-diacetamido-2,4, 6-trideoxy-beta-L-altropyranose hydrolase	It catalyzes the third step in the biosynthesis of pseudaminic acid. Pseudaminic acid is a sialic-like sugar that modifies flagellin.
A0A412WSU3	Bacteriocin	Antimicrobial peptides that can kill or inhibit closely related or unrelated bacterial strains.
A0A413I832	Serine Protease	Extracellular proteases regulate biofilm formation and growth of competitors.
A0A412W3J2	Glutamate dehydrogenase	Involved in bacterial response to stress adaptation and virulence.
A0A412WSS0	Bacteriocin	Antimicrobial peptides that can kill or inhibit closely related or unrelated bacterial strains.
F9ZBU9	Peptidylprolyl isomerase	Promote biofilm formation in *Mycobacterium tuberculosis*.

1248

Table S2-List of primers

Primer Name	Sequence (5'---3')
O. splanchnicus 16S rRNA FP	ATGTAATGATGAGCACTCTAACGG
O. splanchnicus 16S rRNA RP	GGCTTTTGAGATTGGCATCC
PV-1 FP	CGAACTGGAGGCGCGTAA
PV-1 RP	GCAGCGACTTGGCCTTGAC
Claudin5 FP	CCCAGTTAAGGCACGGGTAG
Claudin5 RP	GGCACCGTCGGATCATAGAA

ßactin FP	CAGCAAGCAGGAGTACGATG
ßactin RP	GCAGCTCAGTAACAGTCCG
Vegf FP	GCAGCGACAAGGCAGACTAT
Vegf RP	AACCTCCTCAAACCGTTGGC
Tjp1 FP	TCTTGCAAAGTATCCCTTCTGT
Tjp1 RP	CAGAAATCGTGCTGATGTGCC
Reg3b FP	GCTTGTCAAGAGCTTCTGGATTTC
Reg3b RP	TGTCCCTTGTCCATGATGCTC
Reg3g FP	CGCTGAAGCTTCCTTCCTGT
Reg3g RP	GGGTTCATAGCCCAGTGTCG
Lysozyme C FP	GGTCTACAATCGTTGTGAGTTG
Lysozyme C RP	TGAGCTAAACACACCCAGTCA
CCL 20 FP	GCAACTACGACTGTTGCCTC
CCL20 RP	TTCTTGACTCTTAGGCTGAGGA
Ang4 FP	TGGTTGTGATTCCTCCAACTCT
Ang4 RP	TGTTGACATCTTTGCAAGGCG
fliC FP	CTAAACAAACTGGGTGGCGC
fliC RP	GCACCCAGGTCAGAACGTAA
bapA FP	ACGGTGACTACGCCTAAAAC
bapA RP	CTTCCGCCTCAGTCACTTTT
csgD FP	ACGATTATCCCTACCGTGAA
csgD RP	GCGGACTCGGTGCTGTTGTA
Salmonella 16S rRNA FP	GTGAGGTAACGGCTCACCAA
Salmonella 16S rRNA RP	TAACCGCAACACCTTCCTCC

sipC FP	GCAGTACGCCGCTACTCAGG
sipC RP	AGCGCGAATATTGCCTGCGA
pipB2 FP	TGGCGCCAATTTACAACACGC
pipB2 RP	CTCGCCTTTCGCAGGTTTGC
csgA FP	TCGCAGCAATCGTAGTT
csgA RP	CGGAACCGTACTGATAA

1250

Table S3-List of Antibodies

Antibody Name	Catalogue Number	Source
PV-1	**A15906-20ul (Abclonal)**	**Rabbit**
ZO-1	**R26.4C DHSB, CRL 1597**	**Mouse**
β actin	**A3854 (Sigma), Clone AC-15**	**Mouse**

1252

1253