1 **Paracrine rescue of MYR1-deficient *Toxoplasma gondii* mutants reveals**
2 **limitations of pooled *in vivo* CRISPR screens**
3

4 Francesca Torelli [1,2,a], Diogo M. da Fonseca [1,2,a], Simon Butterworth [1,b], Joanna C.
5 Young [1,c], Moritz Treeck [1,b,*]
6
[1] Signalling in Apicomplexan Parasites Laboratory, The Francis Crick Institute, London, UK NW1 1AT
[2] These authors contributed equally: Francesca Torelli, Diogo M. da Fonseca
[a] Current address: Host-Pathogen Interactions Laboratory, Gulbenkian Institute for Molecular Medicine,
Oeiras, Portugal 2780-156
[b] Current address: Whitehead Institute, Massachusetts Institute of Technology, Cambridge, MA 02142-
1479
[c] Current address: Institute of Immunology and Infection Research, University of Edinburgh, Edinburgh,
UK EH9 3FL
[*] Correspondence and requests for materials should be addressed to M.T. (mtreeck@igc.gulbenkian.pt)

Classifications: Biological Sciences/Microbiology
Keywords: CRISPR-Cas9 screen, *Toxoplasma gondii*, MYR1, host-pathogen interaction, paracrine effect

19
ABSTRACT

Toxoplasma gondii is an intracellular parasite that subverts host cell functions via
secreted virulence factors. Up to 70% of parasite-controlled changes in the host
transcriptome rely on the MYR1 protein, which is required for the translocation of
secreted proteins into the host cell. Mice infected with MYR1 knock-out (KO) strains
survive infection, supporting a paramount function of MYR1-dependent secreted
proteins in *Toxoplasma* virulence and proliferation. However, we have previously shown
that MYR1 mutants have no growth defect in pooled *in vivo* CRISPR-Cas9 screens in
mice, suggesting that the presence of parasites that are wild-type at the *myr1* locus in
pooled screens can rescue the phenotype. Here, we demonstrate that MYR1 is not
required for the survival in IFN-γ-activated murine macrophages, and that parasites
lacking MYR1 are able to expand during the onset of infection. While ΔMYR1 parasites
have restricted growth in single-strain murine infections, we show that the phenotype is
rescued by co-infection with wild-type (WT) parasites *in vivo,* independent of host
functional adaptive immunity or key pro-inflammatory cytokines. These data show that
the major function of MYR1-dependent secreted proteins is not to protect the parasite
from clearance within infected cells. Instead, MYR-dependent proteins generate a
permissive niche in a paracrine manner, which rescues ΔMYR1 parasites within a pool
of CRISPR mutants in mice. Our results highlight an important limitation of otherwise
powerful *in vivo* CRISPR screens and point towards key functions for MYR1-dependent
Toxoplasma-host interactions beyond the infected cell.

41
42

SIGNIFICANCE STATEMENT

Pooled CRISPR screens are powerful tools to interrogate gene function in a high-throughput manner. Genes conferring fitness advantages or disadvantages upon disruption can be identified by sequencing. However, in *Toxoplasma gondii* pooled CRISPR screens in mice, fitness defects for some selected mutants drastically diverge from those observed in single-strain infections. Here, we show that a growth defect of a single *Toxoplasma* gene deletion mutant is rescued if co-infected with wildtype parasites. These results shine light on *Toxoplasma*'s ability to subvert the host response beyond the infected cell, and highlight an important limitation of pooled CRISPR screens in mice. This limitation is probably encountered in CRISPR screens in general where paracrine effects occur.

INTRODUCTION

Pooled CRISPR screens have been an extraordinarily powerful genetic tool to identify gene function in an unbiased manner using negative or positive selection. They have been applied in various cell culture conditions and *in vivo*, in combination with different genetic or chemical bottlenecks to identify genes in a specific setting (1). Fitness-conferring genes are identified by assessing the relative abundance of cells with different genetic perturbations within a pool of mutants upon selective pressure.

We have previously performed pooled CRISPR screens in mice with the intracellular parasite *Toxoplasma gondii*, to identify exported parasite proteins that are important for survival in his natural host (2, 3). For the majority of tested genes, the phenotypes observed in pooled CRISPR screens are concordant to those observed in single-strain infections using clonal mutants for those genes. To our surprise, however, we found that while some *Toxoplasma* genes are required for survival in single-strain mutant KO murine infections, this fitness defect phenotype is lost when the same mutants are part of a heterogenous mutant pool used for CRISPR screens in mice. We hypothesised that individual mutants can be rescued by paracrine effects triggered by other parasites in the pool, pointing towards a potential limitation of pooled CRISPR screens to detect a subset of important virulence factors in *Toxoplasma.* This limitation likely is not restricted to *Toxoplasma*, or other infectious contexts, but CRISPR screens in general where paracrine effects are operating.

Toxoplasma is a ubiquitous parasitic pathogen in all warm-blooded animals. *Toxoplasma* infection is widespread in livestock and wild animals, as well as in humans, where one third of the population is estimated to be seropositive (4). Following oral infection by oocysts or latent stage tissue cysts, the parasite grows as tachyzoites in intermediate hosts, before disseminating to distal organs. The host immune response is

pivotal to reduce parasite burden caused by rapidly proliferating tachyzoites. However, it
is not sufficient to completely clear infection, as surviving parasites can differentiate into
the chronic bradyzoite forms and establish tissue cysts in the central nervous system
and skeletal muscle cells (5). During the onset of infection, recognition of *Toxoplasma*-
derived pathogen-associated molecular patterns (PAMPs) by host innate immune cells
triggers the production of IL-12, amongst other pro-inflammatory mediators (6). IL-12
has a protective role during toxoplasmosis, primarily by triggering IFN-γ production in
lymphocytes and thereby linking innate and adaptive immunity during infection (7). IFN-
γ is a pivotal cytokine in conferring resistance against *Toxoplasma* infection since it
induces expression of interferon-stimulated genes (ISGs) that limit intracellular parasite
proliferation and curtail infection (8–10). These antimicrobial effects are enhanced by
other pro-inflammatory cytokines. An example is TNF, which acts as a co-stimulatory
signal to trigger IFN-γ production by NK cells exposed to the parasite, and boosts the
antimicrobial activity of IFN-γ-activated macrophages (11–13). Following *Toxoplasma*
infection, host cells can also secrete chemokines such as CCL2, which drives
recruitment of $CCR2^+Ly6C^{high}$ inflammatory monocytes from the bone marrow to the
infected sites (14, 15). There, they can differentiate into monocyte-derived dendritic
cells or macrophages and act as an extra line of defence against the parasite (16, 17).
101
In order to survive clearance by host immune cells and to disseminate within the
infected host, *Toxoplasma* relies on an array of over 250 secreted proteins (18).
Following invasion, *Toxoplasma* replicates inside a parasitophorous vacuole (PV)
separated from the host cytoplasm by the PV membrane (PVM). During and after host
cell invasion, *Toxoplasma* secretes proteins from the rhoptries and the dense granules.
Secreted proteins from these organelles are not only pivotal for vacuole establishment,
but also nutrient uptake, reprogramming of the infected cell and protection against the
host immune response (19) . In order to exert their effect, some *Toxoplasma* proteins
secreted from dense granules must cross the PVM, likely via a multi-protein translocon
that depends on MYR1 (20). GRA16, a dense granule effector that relies on MYR-
dependent export to exit the PV and reach the host cytosol, was shown to drive
upregulation of the transcription factor c-Myc in host cells (21). After the identification of
MYR1 other putative components of the MYR translocon – MYR2, MYR3, MYR4,
ROP17 – were identified, using their ability to trigger GRA16-dependent c-Myc induction
in infected cells as a surrogate (22–24). It was subsequently suggested that most, if not
all exported dense granule proteins that reach the host cell cytosol might depend on
functional MYR1 for translocation (23, 25). To date, seven virulence factors have been
shown to be MYR1-dependent: IST, NSM, HCE1/TEEGR, GRA16, GRA18, GRA24 and
GRA28. Once exported, these parasite proteins can interfere with host cell transcription
and contribute to the establishment of permissive niches for *Toxoplasma* proliferation
and survival via multiple downstream mechanisms. Examples include boosting

resistance against IFN-γ-dependent antimicrobial mechanisms (IST (26, 27)), arresting
the host cell cycle (HCE1/TEEGR (28, 29)), inhibiting programmed host cell death
pathways (NSM (30)) and modulating cytokine/chemokine secretion (GRA16, GRA18,
GRA24, GRA28 (31–34).
127
Given the combined importance of exported dense granule effectors, it is not surprising
that the vast majority of transcriptional changes in *Toxoplasma*-infected human
fibroblasts are dependent on MYR1 (35), and that isogenic MYR1-deficient strains are
avirulent in mice (20). In pooled *in vivo* CRISPR screens, however, we have shown that
parasites lacking MYR1 had no fitness defect in a five-day infection experiment within
the mouse peritoneum (2, 3). This is in agreement with results from *in vivo* CRISPR
screens performed in other labs using different parasite strains and mouse backgrounds
where no (25, 36), or only very mild growth defects (37) have been observed for MYR1.
This divergence of results between isogenic infection, which clearly shows an important
role for MYR1 in murine infections, and the pooled CRISPR screens, which show no
defect of MYR1 mutants, led us to hypothesise that: 1) MYR1, and therefore proteins
that rely on the MYR1 complex for translocation, are not essential for the cell-
autonomous survival of parasites in macrophages, the main cell type infected in the first
days of a peritoneal infection (38); and 2) ΔMYR1 parasites within a pool of mutants
may be rescued by MYR1-competent parasites via a paracrine effect, setting up a
parasite-permissive immune environment in which MYR1-deficient mutants can thrive.
144
Here, we verify both hypotheses, showing that MYR1 is not important for the
Toxoplasma cell-autonomous survival within macrophages, and deploy co-infection
strategies proving that MYR1-competent parasites can trans-rescue the growth defect
of ΔMYR1 parasites *in vivo*. This rescue does not depend on the host adaptive immune
response and surprisingly still occurs despite high levels of key pro-inflammatory
cytokines. This knowledge is paramount to understand the biological function of MYR1-
driven rewiring of the host cell, and consequently the function of MYR1-dependent
effector proteins. It also highlights an important limitation of otherwise powerful *in vivo*
pooled CRISPR screens in *Toxoplasma,* where loss-of-function of a protein in one
parasite can be rescued by other parasites in the pool. This limitation likely extends to
CRISPR screens in other biological contexts in which paracrine effects operate.
156
RESULTS
MYR1 is not essential for survival within IFN-γ-stimulated macrophages
To test whether MYR1 was required for the parasite cell-autonomous survival in
immune cells, we infected IFN-γ-primed bone marrow-derived macrophages (BMDMs)
with WT or MYR1 KO parasites established in the type II Prugniaud (Pru) genetic
background, and quantified infected cells after 24 h. As GRA12 was shown to be

required for parasite survival in IFN-γ-stimulated macrophages (39) and had a strong
growth defect in our pooled *in vivo* CRISPR screen (2), we included a ΔGRA12 strain in
these experiments as positive control. As negative control, we included parasites
lacking IST (SI Appendix, Fig. S1b), a known MYR1-dependent factor that inhibits the
induction of interferon-stimulated genes. IST protects the parasites against intracellular
restriction when the infection precedes IFN-γ activation, but not if *Toxoplasma* infects
primed cells (26, 27). As expected, ΔGRA12 parasites showed a significant reduction in
the proportion of infected cells in the presence of IFN-γ (Fig. 1a). Infection with ΔMYR1
was comparable to ΔIST and not significantly different to that of WT parasites, although
a slight increase in restriction was observed for ΔIST and ΔMYR1. Our results are in
line with findings from Wang and colleagues where deletion of MYR1 in a more virulent
type I strain similarly causes no fitness defect in IFN-γ-activated macrophages (40).
175
To further validate the role of virulence factors in conferring parasite resistance against
IFN-γ-mediated host defence, we assessed replication of ΔMYR1 or ΔGRA12 in
competition with WT parasites over two lytic cycles in IFN-γ-treated or unprimed
BMDMs, and in HFFs as control (Fig. 1b). As expected, ΔGRA12 was outcompeted by
WT parasites exclusively in BMDMs pre-stimulated with IFN-γ, confirming its role to
survive the cytokine-mediated clearance in macrophages (Fig. 1c). On the contrary,
ΔMYR1 parasites displayed a fitness defect compared to WT parasites, regardless of
the infected cell type and independently of IFN-γ treatment (Fig. 1c). This result is in line
with the smaller size of the ΔMYR1 plaques established in HFFs compared to the
parental strain (Fig. 1d), which recapitulates published data (22).
186

187
Fig 1. MYR1 and MYR1-dependent factors are not contributing to *in vitro Toxoplasma* survival in
IFN-γ-activated macrophages. (a) BMDMs were stimulated with 100 U/ml IFN-γ for 24 h or left
untreated, before infection with mCherry-expressing *Toxoplasma* strains for 24 h. Cells were fixed and
imaged by high-content imaging, and the percentage of infected cells in IFN-γ-treated BMDMs compared
to untreated controls is shown. WT refers to PruΔUPRT. The box-plot shows the median value and the
whiskers show minimum and maximum values. Significance was tested with the One-way Anova test with
the Benjamini, Krieger and Yekutieli FDR correction, n=5. (b) Schematic of the flow cytometer-based
growth competition assay of an IFN-γ-dependent effector (e.g. GRA12). HFFs and BMDMs were co-
infected with equal amounts of colourless WT and mCherry-expressing ΔGRA12 or ΔMYR1 strains.
BMDMs were stimulated with 100 U/ml IFN-γ for 24 h or left untreated before infection. The mCherry
signal was quantified by flow cytometry analysis and the ratio between the strains after two passages
(output) was compared to the input. (c) Growth of competing mutant and WT parasites was assessed by
flow cytometry. The normalised ratios of output versus input are shown. The average of two independent
experiments with technical triplicates is shown. (d) Violin plot of the plaque size of parental PruΔKU80 and
derived PruΔMYR1 strains. The bar represents the median value and representative images are reported
on the right. Significance was tested using a two-tailed unpaired Welch's t-test, ** p<0.01, **** p<0.0001,
n=2.

MYR1 mutants expand during the course of infection and can form tissue cysts *in*
vivo

To investigate the growth of ΔMYR1 parasites *in vivo* and follow infection over time, we
generated parasite strains expressing firefly luciferase (Luc) in the wild-type (WT-Luc)
or ΔMYR1 backgrounds (ΔMYR1-Luc, SI Appendix Fig. S1c). We injected mice with
25,000 tachyzoites and monitored parasite growth over 7 days by intravital imaging (Fig.

2a). As expected from the reduced *in vitro* growth phenotype in HFFs and BMDMs,
ΔMYR1-Luc showed reduced bioluminescent signal compared to WT-Luc parasites
(Fig. 2b-c), confirming that the growth defect *in vitro* results in analogous phenotypes *in*
vivo. Nevertheless, MYR1-deficient parasites were still able to expand in mice, as
bioluminescence signal increased from day 3 to day 5 post infection (Fig. 2b-c). This
initial increase of growth is similar to what was observed for ΔIST parasites (26, 27).
ΔMYR1-Luc-infected mice that survived the acute phase of infection for four weeks p.i.
carried a low number of cysts in the brain (3.4 cysts/brain on average; 0-13 cysts/brain
detected, Fig. 1d). This is similar to previous observations (2) indicating that expression
of luciferase does not decrease the ability to produce cysts and further supports the
notion that ΔMYR1 parasites are able to form cysts, albeit at low numbers. As mice
infected with WT parasites succumbed during the acute stage of infection, we cannot
compare cyst numbers between the two strains.
225

226
Fig. 2. ΔMYR1 parasites expand in the murine host and form cysts *in vivo*. (a) Mice were infected
i.p. with luciferase-expressing *Toxoplasma* strains, WT-Luc or ΔMYR1-Luc, and whole-body intravital
imaging was performed at days 3, 5 and 7 p.i. (b) Representative whole-body intravital imaging. (c) Graph
shows the total bioluminescence signal converted to a logarithmic scale from two independent
experiments. Statistical differences between the two strains at each timepoint were tested with the Mixed-
effects model (REML) with Sidak post-hoc tests (continued lines). Significance for the ΔMYR1-Luc growth
over time was tested with a One-Way Anova test (dotted line). Number of mice per group: day 3 p.i.:
n=14; day 5 p.i.: n=11; day 7 p.i.: n=8. * $p < 0.05$, *** $p < 0.001$. (d) Representative fluorescent image of a
brain cyst from a mouse infected with ΔMYR1-Luc. mCherry-expressing ΔMYR1-Luc parasites were
detected by microscopy. Cyst wall was stained with the FITC-conjugated lectin DBA. The scale bar
represents 25 μm.
238
MYR1-dependent secreted factor(s) rescue ΔMYR1 *in vivo* growth defect via a
paracrine mechanism, independent of host adaptive immunity

We have shown that ΔMYR1 parasites have reduced *in vitro* and *in vivo* growth, despite
not having any significant growth defect when in a pool with other mutants. Therefore,
we hypothesised that MYR1-competent parasites within the KO pool generate a
permissive environment that ultimately promotes growth of the ΔMYR1 mutants in a
paracrine fashion. To test this hypothesis, we performed co-infection experiments where
mice were injected with an inoculum containing a 20:80 ratio of ΔMYR1-Luc parasites
with either WT parasites or ΔMYR1 mutants not expressing luciferase (Fig. 3a). This
setting allows us to assess if the presence of WT parasites affects growth of ΔMYR1-
Luc parasites within the peritoneum. Our results show that ΔMYR1-Luc parasites
proliferate better in mice co-infected with WT parasites than with ΔMYR1 parasites (Fig.
3b-c). These results support a paracrine role of MYR1-mediated effectors *in vivo*.
252
To understand what mediates this trans-rescue phenotype, we assessed the production
of selected pro-inflammatory cytokines important during *Toxoplasma* infection. Mice
infected with a mix of ΔMYR1:WT or ΔMYR1:ΔMYR1 parasites display comparable
levels of IL-12p40 in both peritoneum and serum at day 7 p.i. However, ΔMYR1:WT
infections elicited higher levels of CCL2/MCP-1, TNF and especially IFN-γ compared to
the ΔMYR1:ΔMYR1 mix (Fig. 3d, SI Appendix Fig. S1d-f).
259
Considering that ΔMYR1-Luc growth is rescued even when high levels of IFN-γ are
produced in the peritoneal cavity, we wanted to assess whether disrupting IFN-γ-
producing cell populations would impact the trans-rescue phenotype. Lymphocytes, in
particular CD8[+], Th1-committed CD4[+] and γδ T cells, are the main sources of IFN-γ
during *Toxoplasma* infection (41). Thus, we applied the same mixed infection strategy in
Rag2-deficient mice that do not produce mature T and B cells, and therefore fail to
deploy adaptive immune responses (42) (Fig. 3e). As expected, an overall higher
parasitaemia was detected in *Rag2*[-/-] mice when compared to WT mice, due to the
known role of the T and B cells to control infection. Nevertheless, in both *Rag2*[-/-] and
WT control mice ΔMYR1-Luc parasites proliferate more when mixed with WT parasites
than with ΔMYR1 parasites (Fig. 3f). These results confirm that the host adaptive
response is not essential for the trans-rescue of MYR1-deficient *Toxoplasma* during
acute infection.
273

Fig 3. The *in vivo* growth defect of ΔMYR1 is rescued by the presence of MYR1-competent parasites, independently of functional host adaptive immunity. (a) Mice were infected i.p. with a mixed inoculum of *Toxoplasma* tachyzoites, containing a 20:80 ratio of luciferase-expressing ΔMYR1-Luc to WT or ΔMYR1 strains that do not express luciferase. Growth of ΔMYR1-Luc was monitored at 3, 5 and 7 days p.i. by whole-body intravital imaging. (b) Representative whole-body intravital imaging. (c) Total bioluminescence signal converted to a logarithmic scale from mice co-infected with ΔMYR1-Luc:WT (n=18) or ΔMYR1-Luc:ΔMYR1 (n=17). Graph shows cumulative data from four independent experiments. Significance was tested with a Two-way Repeated Measures ANOVA with Sidak post-hoc tests. (d) IL-12p40, CCL2/MCP-1, TNF and IFN-γ levels were detected in peritoneal lavage and serum of animals infected with mixed *Toxoplasma* inoculum (20:80 ratio) at day 7 p.i. by ELISA. The violin-plots show the median (continued black line) and quartiles (dotted grey lines). Significance was tested with the Welch's t-test (TNF, CCL2/MCP-1 and IL-12p40 ELISAs) and Mann-Whitney test (IFN-γ ELISAs). Samples from 2 to 3 independent experiments were assayed. Number of samples for CCL2: WT n=13 and ΔMYR1 n=12. Number of samples for other cytokines: peritoneal lavage samples: WT n=8 and ΔMYR1 n=7; serum samples: WT n=4 and ΔMYR1 n=3. Data with higher differences between inocula (CCL2 and IFN-γ) are

displayed in a logarithmic scale in SI Appendix Fig. S1d-f. (e) Representative whole-body intravital
imaging at day 7 p.i. of C57BL/6Ntac (WT) or RAG2-deficient (*Rag2⁻/⁻*) mice infected i.p. with a mixed
inoculum containing a 20:80 ratio of luciferase-expressing ΔMYR1-Luc tachyzoites to WT or ΔMYR1
strains that do not express luciferase. (f) Total bioluminescence signal converted to a logarithmic scale
from mice co-infected with ΔMYR1-Luc:WT (n=10) or ΔMYR1-Luc:ΔMYR1 (n=8). Graph shows
cumulative data from two independent experiments. Significance was tested with the Multiple Mann-
Whitney test with a False Discovery Rate approach by a Two-stage step-method. * p<0.05, ** p<0.01, ***
p<0.001, **** p<0.0001.
298

299
300 Fig 4. Model illustrating paracrine rescue of ΔMYR1 parasites in pooled CRISPR screens
Left panel: In single-strain infections ΔMYR1 parasites grow slowly within the peritoneum and are cleared.
Right panel: Growth of ΔMYR1 parasites within a CRISPR pool is rescued by the presence of MYR-
competent parasites. In cells infected with MYR1 competent parasites, MYR-dependent effectors enter
the host cell, leading to secretion of cytokines that promote a favourable immune environment. MYR1
mutants are able to proliferate under these conditions (i.e are rescued *in trans*). However, *Toxoplasma*
virulence factor mutants required for survival within a cell (cell-autonomous), such as GRA12, can not be
rescued in trans.
308

DISCUSSION

In this work we show that deletion of MYR1, and by extension MYR1-dependent
effectors, does not impact the ability of *Toxoplasma* to initiate an infection in mice and
survive in IFN-γ-stimulated murine macrophages. While clonal ΔMYR1 mutants have a
significantly reduced growth compared to WT parasites *in vivo*, co-infection with WT
parasites increased their ability to proliferate. This finding supports results from pooled
CRISPR-Cas9 screens, where loss-of-function mutants of *Myr1* and other genes
previously shown to be involved in PV translocation of effector proteins, show no (2, 3,
25, 36) or only relatively minor fitness defects in mice (37). The rescue phenotype
observed in mixed infections is very unlikely to occur through co-infection of host cells
by WT and ΔMYR1 parasites, as in the peritoneal exudate from infected mice less than
2% of all infected cells were co-infected (data not shown). As such, our data indicates
that some MYR1-dependent effectors cause changes in infected murine cells, that in
turn provide a favourable environment for parasite growth in a paracrine manner. This is
important as: 1) It shows that the major transcriptional changes caused by MYR-

dependent effectors are not required for *Toxoplasma* to survive cell-autonomous
immune responses in IFN-γ-primed cells; 2) Paracrine rescue in pooled CRISPR-Cas9
screens may mask a significant amount of proteins required for *Toxoplasma* survival; 3)
It is likely that this "paracrine masking effect" can be found in CRISPR screens in other
biological contexts, e.g.: other host-microbe interaction screens, and possibly even in
non-infectious biological contexts, such as pooled CRISPR screen approaches to study
cancer immunity and to discover regulators of innate and adaptive immunity crosstalk
(reviewed in (43, 44)).
332
We found that the presence of WT parasites in a mixed inoculum with ΔMYR1 mutants
elicited significantly higher levels of IFN-γ and TNF in the peritoneum than ΔMYR1
mutants alone. As IFN-γ was shown to be a major cytokine to limit *Toxoplasma* growth
in a cell-autonomous manner, and that TNF acts mainly by enhancing the antimicrobial
effects in IFN-γ-activated cells, one would expect that ΔMYR1-Luc parasites would be
more restricted when injected in combination with WT parasites. However, we observed
the opposite, as higher ΔMYR1-Luc parasitaemia was observed in ΔMYR1:WT than in
ΔMYR1:ΔMYR1 mixes. These data provide further support that MYR1, and by
extension MYR1-dependent effectors, do not protect *Toxoplasma* from the IFN-γ-
mediated intracellular clearance in mice.
343
What could be the driving force of the rescue? We show that the adaptive immune
system plays no role, pointing towards cells of the innate immune system. Higher CCL2
levels in mice infected with ΔMYR1:WT suggests higher inflammatory Ly6C^{high}
monocytes recruitment to infected tissues (14, 15). These cells could be responsible for
the high levels of TNF and IFN-γ detected. While monocytes have been shown in
multiple reports to be important for limiting *Toxoplasma* upon exposure to IFN-γ in
infected niches (14, 17, 45), it is possible that recruited monocytes also provide an
important reservoir for parasites to grow in the peritoneum. However, how ΔMYR1
parasites are eventually cleared in homogenous infections in the absence of high levels
of CCL2 is not yet known.
354
Individual MYR-related effectors that may be responsible for the paracrine rescue have
not been investigated here and we hypothesise that the phenotype is likely the
concerted result of multiple effectors that affect cytokine secretion. For example,
previous studies showed that both GRA18 and GRA28 can induce release of CCL22
from infected cells (32, 46), while GRA16 and HCE1/TEEGR impair NF-kB signalling
and the potential release of pro-inflammatory cytokines such as IL-6, IL-1β and TNF
(29, 47). Regardless of the effector(s), our results highlight an important novel function
of MYR1-dependent effectors by establishing a supportive environment in *trans* for
Toxoplasma growth within the peritoneum.

364
We further confirm previous results using luciferase-expressing ΔMYR1 parasites that
MYR1 appears to be dispensable for the formation and persistence of latent
Toxoplasma stages *per se*. As MYR1 has been demonstrated to be dispensable for
stage conversion *in vitro* (48), the relatively low number of cysts is likely explained by a
failure of ΔMYR1 parasites to efficiently disseminate and/or persist within the murine
host (34). In alternative, we hypothesise that the absence of CCL2 in ΔMYR1 infections
limits recruitment of host cells that *Toxoplasma* can potentially use as vehicles to reach
the brain.
373
This work also highlights the limitations of restriction-based CRISPR-screens in
capturing the variety of pathogen's mechanisms to survive host clearance. Novel
applications of the CRISPR screens, for example in combination with single-cell RNA
sequencing (49) or with functional assays to explore immunological contexts (43) could
help understand how infected cells affect the neighbouring environment to support
infection, and contribute to parasite survival, dissemination and persistence within the
host. Here, we show that MYR1-dependent proteins play a critical role in promoting a
favourable environment for growth beyond the infected cell in a paracrine manner. This
is different to the injection of rhoptry effector proteins by *Toxoplasma* into cells it does
not invade, which requires parasite-host cell contact (50), and provides a novel angle on
how the parasite can systematically alter the host environment in its favour during
infection.
386
387 Our work draws attention to an understudied aspect of pathogen manipulation in
388 complex multicellular settings which warrants further studies. The limitation we highlight
389 here, that mutant phenotypes can be masked in pooled CRISPR screens, likely extends
390 to other experimental setups where paracrine effects are possible.
391
MATERIAL AND METHODS
Cell culture and parasite strains. Primary human foreskin fibroblasts (HFFs) (ATCC)
were maintained in Dulbecco's modified Eagle's Medium (DMEM) with 4.5 g/L glucose
and GlutaMAX-1 (Gibco Thermo Fisher Scientific) supplemented with 10% foetal bovine
serum (FBS) (Gibco Thermo Fisher Scientific) at 37°C and 5% CO_2. To generate bone
marrow-derived macrophages (BMDMs), bone marrow cells were extracted from femurs
of C57BL/6J mice and differentiated to macrophages for 7 days at 37°C and 5% CO_2.
Briefly, bone marrow cells were seeded in 15 cm^2 Petri dishes (Corning) and
differentiated into BMDMs in RPMI 1640 medium (Gibco Thermo Fisher Scientific)
supplemented with 20% L929 conditioned media (containing the murine macrophage
colony stimulating factor (mM-CSF)), 50 µM *2*-mercaptoethanol, Penicillin/Streptomycin
(Thermo Fisher Scientific), and 10% FBS. For experiments BMDMs were grown in the

above RPMI medium but lacking 2-mercaptoethanol (called working media further on). *Toxoplasma gondii* strains PruΔKU80 (51), PruΔUPRT::mCherry (3), PruΔGRA12::mCherry (2), PruΔMYR1::mCherry (2) and derived strains were maintained in confluent HFFs and passaged by syringe lysis through 23G needles every 2-3 days.

Generation of parasite lines. 10^6 freshly lysed parasites were transfected with 20-25 µg of DNA by electroporation using the 4D-Nucleofector (Lonza) with previously optimised protocols as described in Young *et al* (2). All primers used are listed in SI Appendix Fig. S1a. To generate pSAG1::Cas9sgIST the IST gRNA sequence was inserted into pSAG1::Cas9sgUPRT (52) by inverse PCR using primers 1&2. To generate the IST KO (PruΔIST::mCherry), PruΔKU80 parasites were co-transfected with pSAG1::CAS9sgIST and a Pro-GRA1::mCherry::T2A::HXGPRT::Ter-GRA2 construct amplified with primers 3&4 containing 40 bp homology regions to the 5′- and 3′- untranslated regions of IST (ToxoDB TGME49_240060). 24 h after transfection, 50 µg/ml Mycophenolic acid (Merck) and Xanthine (Sigma) (M/X) were added to select for integration, and a clonal culture was verified by PCR with primers 5-8. To generate Luciferase expressing parasite lines, PruΔKU80 or PruΔMYR1::mCherry were co-transfected with pSAG1::CAS9sgUPRT and PciI digested pUPRT-ffLucHA to insert a HA-tagged Firefly luciferase gene (LucHA) into the *Uprt* locus and establish WT-Luc and ΔMYR1-Luc strains respectively. To generate pUPRT-ffLucHA, the GRA1 promoter region and firefly Luciferase sequences were amplified from pGRA (53) and pDHFR-Luc (54) plasmids respectively using primers 9&10 and 11&12 and combined with BamHI/Xma digested pUPRT-HA in a Gibson reaction. 24 h post transfection 20 µg/ml FUDR (Sigma) was added to select for disruption of the *Uprt* locus and clones verified by PCR with primers 13&14.

Parasite growth in BMDMs. $5x10^4$ BMDMs were seeded per well in 96-well Ibitreat black µ-plates (Ibidi GmbH) in working media. The following day cells were either treated with IFN-γ (100 U/mL, Peprotech) or left untreated. 24 h later cells were infected with the strains in triplicate at a multiplicity of infection (MOI) of 0.3, and the plate was centrifuged at 210 *g* for 3 min. At 3 h post infection (p.i.) the medium was replaced to remove non-invaded parasites. Cells were fixed in 4% paraformaldehyde (PFA) at 24 h p.i., washed in PBS and stained with Far Red Cell Mask (1:2,000, Thermofisher Scientific). Plates were imaged on an Opera Phenix High-Content Screening System (PerkinElmer) with a 40x NA1.1 water immersion objective. 38 fields of view with 10 planes were imaged per well. Analysis was performed on a maximum projection of the planes and the percentage of infected cells was quantified over triplicate samples similarly to previously established protocols (3, 55).

Competition assay. $5x10^5$ BMDMs were seeded in 12-well plates in working media. The following day cells were either treated with 100 U/ml IFN-γ or left untreated. 24 h post treatment BMDMs were infected with a 1:1 mix of either WT (PruΔKU80) and PruΔGRA12::mCherry parasites or WT and PruΔMYR1::mCherry parasites at a MOI of 0.3 ($1.5x10^5$ parasites per well). Confluent HFFs in 12-well plates were similarly infected to evaluate defects in growth of the KO parasites. A sample of input parasites were fixed in 4% PFA to verify the starting ratio. 48 h p.i. cells were scraped and passed through a 27G needle, and the parasites inoculated onto HFF monolayers and grown for a further 3-4 days. For flow cytometry analysis, cells were lysed with 23G needles and the parasites passed through a 5 μm filter before fixation in 4% PFA and staining with Hoechst 33342 (Thermofisher Scientific). Samples were analysed on a BD LSR Fortessa flow cytometer and with FlowJo software v10. Hoechst 33342 was excited by a 355 nm laser and detected by a 450/50 band pass filter. mCherry was excited by a 561 nm laser and detected by a 600 long pass filter and a 610/20 band pass filter. To eliminate debris from the analysis, events were gated on forward scatter, side scatter and Hoechst 33342 fluorescence. Parasites were identified by their nuclear staining and KO were discriminated by their mCherry signal. The ratios of KO/WT parasites (mCherry$^+$/ mCherry$^-$) from two independent experiments, each in technical triplicates for each condition, were calculated and normalised by dividing by the input ratio, to allow comparison between strains and biological replicates.

Plaque assay. HFF were grown to confluency in T25 flasks and infected with 200 parasites to grow undisturbed for 10 days. Cells were fixed and stained in a solution with 0.5% (w/v) crystal violet (Sigma), 0.9% (w/v) ammonium oxalate (Sigma), 20% (v/v) methanol in distilled water, then washed with tap water. Plaques were imaged on a ChemiDoc imaging system (BioRad) and measured in FIJI (56).

Animal Ethics Statement. C57BL/6J (Jackson Laboratories), C57BL/6NTac and Rag2 N12 C57BL/6N (Rag2-deficient; *Rag2*$^{-/-}$, Taconic) mice were bred and housed under specific pathogen-free conditions in the biological research facility at the Francis Crick Institute. Mice maintenance and handling adhered to the Home Office UK Animals Scientific Procedures Act 1986. All work and procedures performed were approved by the UK Home Office and performed in accordance with the granted Project License (P1A20E3F9), the Francis Crick Institute Ethical Review Panel, and conforms to European Union directive 2010/63/EU.

***In vivo* infections.** Male and female mice, aged 6-12 weeks were used in this study. For experiments, animals were sex- and age-matched. Mice were injected intraperitoneally (i.p.) with a total of 25,000 *Toxoplasma gondii* tachyzoites in 200 μL PBS either as a single-strain inoculum (100% PruΔUPRT::LucHA or

PruΔMYR1::LucHA), or as a mixed strain inoculum including 5,000 tachyzoites of
luciferase-expressing strain PruΔMYR1::LucHA::mCherry (20%) and 20,000 tachyzoites
of PruΔKU80 or PruΔMYR1::mCherry strains (80%). Mice were monitored and weighed
regularly throughout the experiments. Parasite *in vivo* growth was monitored by
intravital imaging (IVIS) at days 3, 5 and 7 p.i. Mice were injected i.p. with 100 µL of 30
mg/ml luciferin (PerkinElmer) in PBS and were anaesthetised (isoflurane 5% for
induction and 2.5% afterwards) 15 min prior to bioluminescent imaging on an IVIS
Spectrum CT (Perkin-Elmer). Animals were euthanised at day 7 p.i. or at humane
endpoints, blood was collected by cardiac puncture and peritoneal exudate cells were
harvested by peritoneal lavage through injection of 1 mL PBS i.p. Blood was transferred
to serum separating tubes and centrifuged at 10,000 rpm for 10 min at 4°C to isolate
serum, and peritoneal exudate was spun at 500 g for 5 min to remove the cellular
component. IL-12p40 (#88-7120-88), TNF (#88-7324-88), IFN-γ (#88-7314-88) and
CCL2/MCP-1 (#88-7391-88) levels were detected on serum and/or in suspension at
peritoneal exudates of infected mice by ELISA, following manufacturer's protocol
(Thermofisher). To confirm cyst formation, the brain of mice surviving infection with
PruΔMYR1::LucHA::mCherry tachyzoites were homogenised in 1 mL PBS and 300 µL
of the sample were stained with FITC-conjugated Dolichos Biflorus Agglutinin (DBA; 1:
200; Vector Laboratories #RL-1031) for 1 h at room temperature. Fluorescently labelled
cysts were counted using a Ti-E Nikon microscope.

Data analysis. Data was analysed in GraphPad Prism v10. All data shown is presented
as means ± SEM, except for violin plots, where median and quartiles are presented.
Two-tailed unpaired Welch's t-test (Gaussian-distributed data) or Mann-Whitney test
(non-Gaussian-distributed data) were used for statistical analysis of data with only two
experimental groups. For analysis of data with three or more experimental groups, One-
way or Two-way ANOVA were performed. When datasets did not follow a Gaussian
distribution, data was transformed to a logarithmic scale and parametric statistical
analysis was performed on transformed datasets (57). If logarithmic-transformed data
still did not follow a Gaussian distribution, untransformed data was analysed by non-
parametric tests (standard or multiple Mann-Whitney tests). Statistical significance was
set as: ns – not statistically significant, * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$, **** $p <$
0.0001.

Author Contributions. Conceptualisation: F.T., D.F., J.C.Y., M.T.; investigation: F.T.,
D.F., J.C.Y., S.B.; writing, review, editing: all authors; supervision: M.T.; funding
acquisition: F.T., M.T.

Acknowledgments. We thank the High-Throughput Screening Science Technology
Platform (STP), the Flow Cytometry STP, the Biological Research Facility and the

Imaging STP for support. We thank Jeanette Wagener and Aïcha Stierlen for their help in experiment performance and setup. We thank Andreas Wack's lab for providing protocols and support. We want to thank all members of the Treeck lab for critical and continuous discussion. This work was supported by an award to M.T. from the Wellcome Trust (223192/Z/21/Z), by funding to M.T. from the Francis Crick Institute, which receives its core funding from Cancer Research UK (CC2132), the UK Medical Research Council (CC2132), and the Wellcome Trust (CC2132). F.T. is supported by the Deutsche Forschungsgemeinschaft (TO 1349/1-1). J.C.Y. is funded by an MRC Career Development award (MR/V03314X/1). We thank VEuPathDB (58) for providing access to the *Toxoplasma* databases.

Competing interests. The authors declare no competing interests.

REFERENCES

1. C. Bock, *et al.*, High-content CRISPR screening. *Nature reviews. Methods primers* **2** (2022).

2. J. Young, *et al.*, A CRISPR platform for targeted in vivo screens identifies Toxoplasma gondii virulence factors in mice. *Nat Commun* **10**, 3963 (2019).

3. S. Butterworth, *et al.*, Toxoplasma gondii virulence factor ROP1 reduces parasite susceptibility to murine and human innate immune restriction. *PLoS Pathog* **18**, e1011021 (2022).

4. G. Pappas, N. Roussos, M. E. Falagas, Toxoplasmosis snapshots: global status of Toxoplasma gondii seroprevalence and implications for pregnancy and congenital toxoplasmosis. *Int J Parasitol* **39**, 1385–94 (2009).

5. J. P. Dubey, D. S. Lindsay, C. A. Speer, Structures of Toxoplasma gondii tachyzoites, bradyzoites, and sporozoites and biology and development of tissue cysts. *Clin Microbiol Rev* **11**, 267–99 (1998).

6. M. Sasai, M. Yamamoto, Innate, adaptive, and cell-autonomous immunity against Toxoplasma gondii infection. *Exp Mol Med* **51**, 1–10 (2019).

7. R. T. Gazzinelli, *et al.*, Parasite-induced IL-12 stimulates early IFN-gamma synthesis and resistance during acute infection with Toxoplasma gondii. *J Immunol* **153**, 2533–43 (1994).

8. J. P. Saeij, E.-M. Frickel, Exposing Toxoplasma gondii hiding inside the vacuole: a role for GBPs, autophagy and host cell death. *Curr Opin Microbiol* **40**, 72–80 (2017).

9.   J. D. MacMicking, Interferon-inducible effector mechanisms in cell-autonomous
immunity. *Nat Rev Immunol* **12**, 367–82 (2012).

10. J. P. Hunn, C. G. Feng, A. Sher, J. C. Howard, The immunity-related GTPases in
mammals: a fast-evolving cell-autonomous resistance system against intracellular
pathogens. *Mamm Genome* **22**, 43–54 (2011).

11. L. D. Sibley, L. B. Adams, Y. Fukutomi, J. L. Krahenbuhl, Tumor necrosis factor-
alpha triggers antitoxoplasmal activity of IFN-gamma primed macrophages. *J
Immunol* **147**, 2340–5 (1991).

12. J. A. Langermans, *et al.*, IFN-gamma-induced L-arginine-dependent
toxoplasmastatic activity in murine peritoneal macrophages is mediated by
endogenous tumor necrosis factor-alpha. *J Immunol* **148**, 568–74 (1992).

13. A. Sher, I. P. Oswald, S. Hieny, R. T. Gazzinelli, Toxoplasma gondii induces a T-
independent IFN-gamma response in natural killer cells that requires both
adherent accessory cells and tumor necrosis factor-alpha. *J Immunol* **150**, 3982–
9 (1993).

14. P. M. Robben, M. LaRegina, W. A. Kuziel, L. D. Sibley, Recruitment of Gr-1+
monocytes is essential for control of acute toxoplasmosis. *J Exp Med* **201**, 1761–
9 (2005).

15. I. R. Dunay, *et al.*, Gr1(+) inflammatory monocytes are required for mucosal
resistance to the pathogen Toxoplasma gondii. *Immunity* **29**, 306–17 (2008).

16. J. de D. Ruiz-Rosado, *et al.*, MIF Promotes Classical Activation and Conversion of
Inflammatory Ly6C(high) Monocytes into TipDCs during Murine Toxoplasmosis.
Mediators Inflamm **2016**, 9101762 (2016).

17. R. S. Goldszmid, *et al.*, NK cell-derived interferon-γ orchestrates cellular
dynamics and the differentiation of monocytes into dendritic cells at the site of
infection. *Immunity* **36**, 1047–59 (2012).

18. K. Barylyuk, *et al.*, A Comprehensive Subcellular Atlas of the Toxoplasma
Proteome via hyperLOPIT Provides Spatial Context for Protein Functions. *Cell
Host Microbe* **28**, 752-766.e9 (2020).

19. M.-A. Hakimi, P. Olias, L. D. Sibley, Toxoplasma Effectors Targeting Host
Signalling and Transcription. *Clin Microbiol Rev* **30**, 615–645 (2017).

20. M. Franco, *et al.*, A Novel Secreted Protein, MYR1, Is Central to Toxoplasma's
Manipulation of Host Cells. *mBio* **7** (2016).

21. M. W. Panas, J. C. Boothroyd, Toxoplasma Uses GRA16 To Upregulate Host c-
Myc. *mSphere* **5** (2020).

22. N. D. Marino, *et al.*, Identification of a novel protein complex essential for effector
translocation across the parasitophorous vacuole membrane of Toxoplasma
gondii. *PLoS Pathog* **14**, e1006828 (2018).

23. M. W. Panas, *et al.*, Translocation of Dense Granule Effectors across the
Parasitophorous Vacuole Membrane in Toxoplasma-Infected Cells Requires the
Activity of ROP17, a Rhoptry Protein Kinase. *mSphere* **4** (2019).

24. A. M. Cygan, *et al.*, Coimmunoprecipitation with MYR1 Identifies Three Additional
Proteins within the Toxoplasma gondii Parasitophorous Vacuole Required for
Translocation of Dense Granule Effectors into Host Cells. *mSphere* **5** (2020).

25. L. O. Sangaré, *et al.*, In Vivo CRISPR Screen Identifies TgWIP as a Toxoplasma
Modulator of Dendritic Cell Migration. *Cell Host Microbe* **26**, 478-492.e8 (2019).

26. P. Olias, R. D. Etheridge, Y. Zhang, M. J. Holtzman, L. D. Sibley, Toxoplasma
Effector Recruits the Mi-2/NuRD Complex to Repress STAT1 Transcription and
Block IFN-γ-Dependent Gene Expression. *Cell Host Microbe* **20**, 72–82 (2016).

27. G. Gay, *et al.*, Toxoplasma gondii TgIST co-opts host chromatin repressors
dampening STAT1-dependent gene regulation and IFN-γ–mediated host
defenses. *Journal of Experimental Medicine* **213**, 1779–1798 (2016).

28. M. W. Panas, A. Naor, A. M. Cygan, J. C. Boothroyd, Toxoplasma Controls Host
Cyclin E Expression through the Use of a Novel MYR1-Dependent Effector
Protein, HCE1. *mBio* **10** (2019).

29. L. Braun, *et al.*, The Toxoplasma effector TEEGR promotes parasite persistence
by modulating NF-κB signalling via EZH2. *Nat Microbiol* **4**, 1208–1220 (2019).

30. A. Rosenberg, L. D. Sibley, Toxoplasma gondii secreted effectors co-opt host
repressor complexes to inhibit necroptosis. *Cell Host Microbe* **29**, 1186-1198.e8
(2021).

31. A. Bougdour, *et al.*, Host cell subversion by Toxoplasma GRA16, an exported
dense granule protein that targets the host cell nucleus and alters gene
expression. *Cell Host Microbe* **13**, 489–500 (2013).

32. H. He, *et al.*, Characterization of a Toxoplasma effector uncovers an alternative
GSK3/β-catenin-regulatory pathway of inflammation. *Elife* **7** (2018).

33.   L. Braun, *et al.*, A Toxoplasma dense granule protein, GRA24, modulates the early
immune response to infection by promoting a direct and sustained host p38
MAPK activation. *Journal of Experimental Medicine* **210**, 2071–2086 (2013).

34.   A. L. Ten Hoeve, *et al.*, The Toxoplasma effector GRA28 promotes parasite
dissemination by inducing dendritic cell-like migratory properties in infected
macrophages. *Cell Host Microbe* **30**, 1570-1588.e7 (2022).

35.   A. Naor, *et al.*, MYR1-Dependent Effectors Are the Major Drivers of a Host Cell's
Early Response to Toxoplasma, Including Counteracting MYR1-Independent
Effects. *mBio* **9** (2018).

36.   Y. Tachibana, E. Hashizaki, M. Sasai, M. Yamamoto, Host genetics highlights IFN-
γ-dependent Toxoplasma genes encoding secreted and non-secreted virulence
factors in in vivo CRISPR screens. *Cell Rep* **42**, 112592 (2023).

37.   C. J. Giuliano, *et al.*, CRISPR-based functional profiling of the Toxoplasma gondii
genome during acute murine infection. *Nat Microbiol* (2024).
https://doi.org/10.1038/s41564-024-01754-2.

38.   K. D. C. Jensen, *et al.*, Toxoplasma Polymorphic Effectors Determine Macrophage
Polarization and Intestinal Inflammation. *Cell Host Microbe* **9**, 472–483 (2011).

39.   B. A. Fox, *et al.*, Toxoplasma gondii Parasitophorous Vacuole Membrane-
Associated Dense Granule Proteins Orchestrate Chronic Infection and GRA12
Underpins Resistance to Host Gamma Interferon. *mBio* **10** (2019).

40.   Y. Wang, *et al.*, Genome-wide screens identify Toxoplasma gondii determinants of
parasite fitness in IFNγ-activated murine macrophages. *Nat Commun* **11**, 5258
(2020).

41.   S. Nishiyama, A. Pradipta, J. S. Ma, M. Sasai, M. Yamamoto, T cell-derived
interferon-γ is required for host defense to Toxoplasma gondii. *Parasitol Int* **75**,
102049 (2020).

42.   Y. Shinkai, *et al.*, RAG-2-deficient mice lack mature lymphocytes owing to inability
to initiate V(D)J rearrangement. *Cell* **68**, 855–67 (1992).

43.   H. Shi, J. G. Doench, H. Chi, CRISPR screens for functional interrogation of
immunity. *Nat Rev Immunol* **23**, 363–380 (2023).

44.   E. A. Holcomb, *et al.*, High-content CRISPR screening in tumor immunology. *Front*
Immunol **13**, 1041451 (2022).

45.   D. G. Mordue, L. D. Sibley, A novel population of Gr-1+-activated macrophages
induced during acute toxoplasmosis. *J Leukoc Biol* **74**, 1015–25 (2003).

46. E. N. Rudzki, *et al.*, Toxoplasma gondii GRA28 Is Required for Placenta-Specific
Induction of the Regulatory Chemokine CCL22 in Human and Mouse. *mBio* **12**,
e0159121 (2021).

47. S. H. Seo, S. G. Kim, J. H. Shin, D. W. Ham, E. H. Shin, Toxoplasma GRA16
Inhibits NF-κB Activation through PP2A-B55 Upregulation in Non-Small-Cell Lung
Carcinoma Cells. *Int J Mol Sci* **21**, 6642 (2020).

48. S. Seizova, *et al.*, Transcriptional modification of host cells harboring Toxoplasma
gondii bradyzoites prevents IFN gamma-mediated cell death. *Cell Host Microbe*
30, 232-247.e6 (2022).

49. S. Butterworth, *et al.*, High-throughput identification of Toxoplasma gondii effector
proteins that target host cell transcription. *Cell Host Microbe* **31**, 1748-1762.e8
(2023).

50. A. A. Koshy, *et al.*, Toxoplasma co-opts host cells it does not invade. *PLoS Pathog*
8, e1002825 (2012).

51. B. A. Fox, *et al.*, Type II Toxoplasma gondii KU80 Knockout Strains Enable
Functional Analysis of Genes Required for Cyst Development and Latent
Infection. *Eukaryot Cell* **10**, 1193–1206 (2011).

52. B. Shen, K. M. Brown, T. D. Lee, L. D. Sibley, Efficient Gene Disruption in Diverse
Strains of Toxoplasma gondii Using CRISPR/CAS9. *mBio* **5** (2014).

53. I. Coppens, *et al.*, Toxoplasma gondii sequesters lysosomes from mammalian
hosts in the vacuolar space. *Cell* **125**, 261–74 (2006).

54. J. P. J. Saeij, J. P. Boyle, M. E. Grigg, G. Arrizabalaga, J. C. Boothroyd,
Bioluminescence Imaging of Toxoplasma gondii Infection in Living Mice Reveals
Dramatic Differences between Strains. *Infect Immun* **73**, 695–702 (2005).

55. E. J. Lockyer, *et al.*, A heterotrimeric complex of Toxoplasma proteins promotes
parasite survival in interferon gamma-stimulated human cells. *PLoS Biol* **21**,
e3002202 (2023).

56. J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis.
Nat Methods **9**, 676–82 (2012).

57. J. H. MacDonald, *Handbook of Biological Statistics*, 3rd Ed. (Sparky House
Publishing, 2014).

58. B. Amos, *et al.*, VEuPathDB: the eukaryotic pathogen, vector and host
bioinformatics resource center. *Nucleic Acids Res* **50**, D898–D911 (2022).

SI Appendix Fig. S1. (a) List of primers used in the study. (b) PCR validation of the PruΔIST strain compared to parental PruΔKu80 (WT). (c) PCR validation of the luciferase-expressing WT-Luc and ΔMYR-Luc strains compared to their respective parental strains PruΔKu80 (WT) and PruΔMYR1 (ΔMYR1). (d, e and f) IFN-γ at peritoneal lavage (d) and CCL2 at both peritoneal lavage (e) and serum (f), were detected by ELISA at day 7 p.i. with mixed *Toxoplasma* inocula (20:80 ratio) of luciferase-expressing ΔMYR1-Luc to WT or ΔMYR1 strains that do not express luciferase. The violin-plots display data shown in Fig. 3d converted to a logarithmic scale, with the median (continued black line) and quartile values (dotted grey lines) highlighted. Significance was tested with the Mann-Whitney test for IFN-γ, and by Welch's t-test for CCL2 data. The number of biological replicates (n) in each graph is the same as mentioned in the original graphs on Fig. 3d. *** p<0.001, **** p<0.0001.