Title: BEHAV3D Tumor Profiler to map heterogeneous cancer cell behavior in the tumor microenvironment

Emilio Rios-Jimenez*[1], Anoek Zomer*[2,3], Raphael Collot[2,3], Mario Barrera Román[2,3], Hendrikus Ariese[2,3], Ravian L. van Ineveld[2,3], Michiel Kleinnijenhuis[2,3], Nils Bessler[2,3], Hannah Johnson[2,3], Anne Rios[2,3] #, Maria Alieva[1] #

* Equal contribution

Shared last authorship

Affiliations:

[1]Institute for Biomedical Research Sols-Morreale (IIBM), Spanish National Research Council-Universidad Autónoma de Madrid, Madrid, Spain.

[2]Princess Máxima Center for Pediatric Oncology, Utrecht, the Netherlands.

[3]Oncode Institute, Utrecht, the Netherlands.

Abstract

Intravital microscopy (IVM) enables live imaging of animals at single-cell level, offering
essential insights into cancer progression. This technique allows for the observation of single-
cell behaviors within their natural 3D tissue environments, shedding light on how genetic and
microenvironmental changes influence the complex dynamics of tumors. The complexity of
data generated by IVM often surpasses the capabilities of conventional analyses accessible to
biomedical scientists, thereby neglecting single-cell heterogeneity and limiting the exploration
of microenvironmental influences on cellular behavior without bias. To address this challenge,
here we introduce BEHAV3D Tumor Profiler (BEHAV3D-TP), a user-friendly computational
framework designed for the comprehensive analysis of single tumor cell behaviors and their
interactions with the tumor microenvironment (TME). BEHAV3D-TP facilitates unbiased
profiling of cancer cell dynamics without requiring advanced computational expertise. Here,
we apply BEHAV3D-TP to study diffuse midline glioma (DMG), a highly aggressive pediatric
brain tumor characterized by invasive growth. Our analysis reveals that distinct migratory
behaviors of DMG cells correlate with specific TME components such as tumor-associated
macrophages and vasculature. This approach, initially aimed at uncovering tumor invasive
patterns and their interactions with the TME, holds promise for understanding additional cancer
cell behaviors like intravasation and metastasis. BEHAV3D-TP represents a significant
advancement in democratizing the analysis of heterogeneous cancer cell behaviors and their
TME interactions, providing accessible computational insights into tumor dynamics.

INTRODUCTION

Cancer is a complex disease driven by a sequence of genetic and microenvironmental changes. As a result, tumors comprise many different cell populations that are in constant evolution[1] . This tumor heterogeneity has important biological and clinical implications as it differentially promotes key characteristics of individual cancer cells, including proliferation, invasion, and drug resistance [2]. Yet, we are far from fully understanding all the dynamic and bi-directional interactions in heterogeneous tumors, and how these interactions influence cancer cell behavior at the single-cell level. Emerging single-cell technologies and analysis tools provide a new opportunity to profile individual cells within tumors. For example, recent spatial transcriptomic and proteomic profiling provides evidence for the existence of multiple distinct microenvironmental niches within one tumor [3]. However, these techniques can only provide static information and make use of thin tissue sections that are only one cell layer thick. To investigate the real-time behavior of individual cells intravital microscopy (IVM) has been developed to study tumor cell behavior evolution over time but also in their three-dimensional native tissue environment.

IVM is a technique used to capture both spatial and temporal information at a single-cell level in living animals. Over the past years, it has provided many new insights into the dynamics of tumor heterogeneity and the development of therapy resistance [4–14 6–10,12–15]. For example, intravital imaging of mouse breast cancer models revealed that only a small portion of tumor cells is motile [16], and that fast- and slow-migrating cells are present in distinct regions of the tumor [7,17]. Moreover, additional heterogeneity exists among migrating tumor cells that can, for instance, move randomly or directionally relative to other tumor cells and/or surrounding healthy tissue [7]. However, traditional IVM analyses have considered cancer cells or the tumor microenvironment (TME) as a homogeneous population, thereby ignoring heterogeneity at the single-cell level and the influence of the local microenvironment, where interactions between neighboring cells may affect cellular behavior. To address these complexities, recent studies have used unbiased approaches using morpho-kinetic parameters derived from live *in vitro*[18] or *in vivo* imaging to classify cells and explore the single-cell behavioral landscape as an additional omic layer [19–21]. In line with this concept of characterizing cellular dynamic properties for cell classification, we have previously developed an analytical platform termed BEHAV3D [20,22] allowing to perform behavioral phenotyping of engineered T cells targeting cancer. This approach revealed substantial behavioral heterogeneity among the engineered T cells and

identified, based on behavioral characteristics, a subpopulation of T cells with increased killing
capacity.

However, the implementation of unbiased methods for cellular dynamic analysis, such as
BEHAV3D, often demands computational expertise beyond the reach of many biomedical
researchers conducting IVM experiments. To democratize the thorough understanding of
heterogeneous cancer cell behavior and the microenvironmental components influencing it,
there is a need for comprehensive, user-friendly, zero-code analysis tools. Such tools would
facilitate unbiased analysis of diverse tumor dynamics, making advanced computational
techniques accessible to a broader range of researchers.

Building upon our BEHAV3D pipeline for analyzing engineered T cell behavior[22], we
introduce BEHAV3D Tumor Profiler (BEHAV3D-TP) (Fig. 1), a new zero-code computational
framework tailored to map the behavioral patterns of single tumor cells and their interplay with
the TME. We used this novel tool to get more insights into the *in vivo* migration behavior of
diffuse midline glioma (DMG), a highly aggressive pediatric brain tumor characterized by
invasive growth and its capacity to spread to other regions of the brain and the spinal cord [23].
For the first time, by visualizing TME components either during or after IVM, we were able to
show that DMG cell migration behavior correlates with specific microenvironmental factors,
including tumor-associated macrophages and vasculature. In our study, we specifically focused
on cancer cell migration as a specific feature of DMG, but our methodology can easily be
extended to other cancer cell behaviors such as intravasation or metastatic colonization. Overall,
BEHAV3D-TP provides an accessible computational approach for a better understanding of
tumor-TME crosstalk and the functional implications of this communication in heterogeneous
tumors.

RESULTS

BEHAV3D Tumor Profiler: an accessible tool for mapping heterogeneity in tumor dynamics

To comprehensively map and analyze the dynamic heterogeneity of tumor cells within their TME we developed BEHAV3D-TP [https://github.com/imAIgene-Dream3D/BEHAV3D_Tumor_Profiler]. Aiming at enhancing accessibility and widespread adoption across scientists studying single tumor dynamics in live imaging, and inspired by previous work on democratizing advanced image analysis[24–26], we designed BEHAV3D-TP as a modular Jupyter Notebook featuring an intuitive graphical user interface (GUI) (Figure 1 and Supplementary Video 1). Operating directly within a web browser, the platform only requires a Google account for access. BEHAV3D-TP implementation within the Google Colaboratory (Colab) framework not only ensures efficient cloud-based computation with free and scalable resources but also enhances tool accessibility by eliminating the need for local software installation, a common barrier for inexperienced users. Regardless of programming proficiency, users can effortlessly navigate the computational pipeline to extract comprehensive insights into tumor dynamics (Figure 1). After the time-lapse image file processing and preparation (Figure 1a), BEHAV3D-TP includes a central module called the *heterogeneity module* (Figure 1b), enabling researchers to uncover intricate patterns of tumor behavior. Additionally, we provide two optional modules—the *large-scale phenotyping module* (Figure 1c) and the *small-scale phenotyping module* (Figure 1d)—which, combined with TME visualization during or after IVM, allow for correlating different tumor behavioral patterns with the composition of the TME. To demonstrate the application of BEHAV3D-TP, we utilized single-cell dynamic measurements obtained from intravital imaging of DMG-bearing mice.

Identification of highly heterogeneous *in vivo* cell behavior displayed by DMGs

To get insights into the behavioral heterogeneity among individual DMG cells, we performed IVM on mice implanted with a cranial imaging window to visualize cancer cells expressing H2BmNeonGreen marker for up to 5.4 hours (Figure supplement 1a). We tracked intravitally imaged DMG cells from different mice and in each imaged position we determined how each DMG cell's position changes relative to the tumor edge (Figure 2a). Next, to untangle the complexity of tumor cell dynamics in an unbiased manner, we used the BEHAV3D-TP *heterogeneity module,* that implements multiparametric single-cell time-series classification allowing to identify distinct single-cell behavioral patterns (Figure 2 a, b). For this, we used

kinetic parameters, including displacement, speed, persistence, and movement direction relative

to the tumor edge, to measure similitudes between cellular behavior using the dynamic time

warping algorithm[22], followed by dimensionality reduction to classify cells. The BEHAV3D-

TP *heterogeneity module* identified 7 DMG behavioral clusters: *retreating*, *slow retreating*,

erratic, *static*, *slow*, *slow invading,* and *invading* (Figure 2b, c, d, Supplementary video 2). To

accurately describe each of the clusters, we analyzed their most predominant features (Figure

2d) and back-projected the cluster information into the original time-lapse to visually inspect

cell behavior in relation to the tumor (Figure 2a, c, Figure supplement 1b, Supplementary video

2). Of particular interest were cells from the *invading* and *retreating* clusters, characterized by

fast movement away from and towards the tumor edge, respectively (Figure 2c-f, and Figure

supplement 1c). In fact, in this particular DMG model, approximately 10% of the cancer cells

were displaying *invasive* behavior (Figure supplement 1d, e), in line with the intrinsic

infiltrative nature of DMG[27] . Interestingly, about another 10% of DMG cells were moving back

to the tumor edge (*retreating*, *slow retreating* clusters) (Figure supplement 1d, e), behavior that

has been previously observed for glioblastoma and suggested to be regulated by differential

chemotactic signaling by the tumor edge and the surrounding brain parenchyma [7]. We also

observed *erratic* cells that display fast non-directed migration behavior, and *static* cells that are

non-migratory (Figure 2c-f). Crucially, the distinct behavioral patterns were detected in all

intravitally imaged mice (Figure supplement 1d, e), ruling out any sample bias that might lead

to clusters consisting solely of cells from a single tumor or mouse. This underscores the

effectiveness of the BEHAV3D-TP in revealing representative heterogeneous tumor behavioral

patterns.

Mapping *in vivo* tumor cell migratory properties to distinct TME regions identified

through *ex vivo* large-scale 3D imaging

To investigate whether the TME influences behavioral heterogeneity among cancer cells, we

developed two complementary approaches to correlate the identified behavioral clusters (Figure

2) to the composition of the TME (Figures 1c, 3a and 4a).

The first approach consists of performing large-scale TME phenotyping and identifying regions

with a specific cellular composition and architecture within the TME of intravitally imaged

tumors that have been intravitally imaged and subsequently fixed. This is accomplished by

using multispectral large-scale single-cell resolution 3D (mLSR-3D) imaging data [28,29] and a

spatial analysis framework called CytoMAP [30] (Figure 3a). We selected microenvironmental

components of interest, such as perivascular niches and glial cells, as they have been previously

reported to play an important role in the invasive behavior of glioblastoma cells [7,31–35].
Specifically, intravitally imaged brains were collected immediately after the IVM imaging
sessions and were fixed and stained for blood vessels (CD31), oligodendrocytes (Olig2), and
tumor-associated microglia/macrophages (TAMM) (Iba1) (Figure 3b). Subsequent CytoMAP[30]
spatial phenotyping analysis identified three different environmental niches: *Void* regions (less
vascularized and with low glial infiltration), *TAMM/oligo* regions (enriched in
oligodendrocytes, TAMM, and tumor cells), and *TAMM/vascularized* regions (highly
vascularized regions enriched in TAMM) (Figure 3b, and Figure supplement 2a-c). To
investigate the heterogeneity of tumor cell behavior in relation to the identified
microenvironmental niches, BEHAV3D-TP *large-scale phenotyping module* mapped the
different niches (*Void*, *TAMM/Oligo*, and *TAMM/vascularized*) identified by CytoMAP onto
3D intravital imaging data (Figures 1c, 3c). In the TME-defined IVM-imaged positions we
compared the frequencies of the different behavioral clusters that we identified by with the
BEHAV3D-TP *heterogeneity* module (Figure 3d). Interestingly, *invading* cells were more
abundant in *TAMM/vascularized* regions compared to *Void* regions that contain a higher
proportion of *static* cells, and *TAMM/Oligo* microenvironments contain more *slow retreating*
cells compared to the other regions (Figure 3d). Finally, we compared the results obtained with
BEHAV3D-TP to a more classical IVM analysis approach that relies on assessing single
dynamic parameters. Interestingly, single parameters showed a restricted ability to identify
differences in cellular behavior among various environments (Figure supplement 2e-g),
underscoring the importance of multiparametric behavioral mapping in revealing more nuanced
cellular behavior that cannot be captured solely by individual dynamic parameters.

Linkage of *in vivo* tumor cell behavior to the composition of the microenvironment at a
189
single-cell level
190

With the goal of further refining TME phenotyping to better understand tumor cell behavior
determinants, we implemented an alternative approach using the BEHAV3D-TP *small-scale*
phenotyping module (Figure 1d, 4a). This module utilizes the detection of TME components
during IVM to offer insights into their abundance and spatial relationship with DMG cells at a
single-cell level. Together with the *heterogeneity module* that profiles the single-cell behavior
of tumor cells (Figure 1), this additional module allows linking different DMG behavioral
profiles with DMG cell small-scale niches (Figures 1d, 4a). For *in vivo* TME labeling, we
injected mice with an anti-CD31-AF647 antibody to label blood vessels, and with either SR101,

a fluorescent marker known to label oligodendrocytes and astrocytes [36–39], or CD20r, a novel

molecule described to label microglial cells [40] (Figure supplement 3a, b). This allowed us to

measure spatial TME characteristics of each single DMG cell, such as minimal distance to a

certain TME component, the number of these cell types in a certain radius, or the cell density

for this particular cell (Figure 1d). Interestingly, the BEHAV3D-TP *small-scale phenotyping*

module revealed substantial TME heterogeneity across the different behavioral DMG clusters

(Figure 4b). Firstly, compared to *slow* and *slow-retreating* cells, *invading* cells were usually

found to be less densely positioned relative to other DMG cells (Figure 4b, and Figure

supplement 3c), correlating with their invasive potential into the brain parenchyma. Second,

and in line with the data from the large-scale TME phenotyping (Figure 3), our data suggests

that *invading* DMG cells reside in a local environment that is enriched in TAMMs (Figure 4b).

Indeed, *invading* cells are closer to TAMMs compared to *erratic* and *slow* migrating cells

(Figure 4c), and move directionally towards TAMMs (Figure 4d, Supplementary video 3),

suggesting that migration of this behavioral cluster is at least partially mediated by chemotaxis.

Third, all fast-migrating behavioral clusters (*invading*, *erratic,* and *retreating*) were found to

be generally closer to blood vessels (Figure 4b and e), suggesting that perivascular invasion is

a more efficient movement route for these cells (Figure 4f), a mode of migration that has been

previously observed for adult glioblastoma cells as well [7,31,33–35]. Lastly, particularly *slow-*

retreating DMG cells were found in regions enriched for SR101[+] cells (Figure 4b and Figure

supplement 3d), perhaps reflecting oligodendrocyte-like (OC-like) tumor DMG cells that have

been found near oligodendrocytes [41]. Altogether, these data demonstrate the power of our

BEHAV3D-TP pipeline in correlating single-cell tumor cell behavior to its local environment,

with an important opportunity to identify novel environmental regulators of invasion.

DISCUSSION

Here, we provide a new user-friendly platform tailored for IVM single tumor cell dynamics

analysis, termed BEHAV3D-TP, aiming to better understand the mechanisms underlying tumor

heterogeneity. In contrast to traditional pipelines that rely on the analysis of single parameters

such as cell speed or displacement[42–44] or that use an artificial threshold to assign a specific

behavior to cells[10], BEHAV3D-TP is an unbiased approach based on the integration of multiple

parameters measured by IVM. Using the *heterogeneity module* of our BEHAV3D-TP pipeline,

we uncovered behavioral phenotypes that have not been described before and that could not

have been identified using single parameter analysis. For example, our analysis revealed

invading and *retreating* clusters, composed of cells with a similar cell speed but with an

opposite migration direction, an important biological and clinical difference. Likely,
bidirectional migration is mediated by different chemotactic cues and by the composition of the
local TME[45]. It is of vital importance for DMG and other cancer patients to better understand
the biology underlying the different behavioral phenotypes, especially concerning *invading*
cells that are difficult to therapeutically target [46].

Previous studies suggest a major role for the microenvironment in promoting tumor cell spread
into the healthy brain [47–49]. Indeed, using the complementary *large-* and *small-scale TME*
phenotyping modules of BEHAV3D-TP, we identified substantial differences in the TME
composition of the different behavioral DMG clusters. Our results also show that the small-
scale TME analysis has a higher predictive value compared to the large-scale TME analysis,
suggesting that a cell's direct microenvironment is a more important regulator of its behavior
compared to its macroenvironment. Future experiments directed towards identifying more TME
components, for example using multispectral LSR-3D imaging[28], could therefore further refine
the *large-scale phenotyping module* by detecting subregion heterogeneity.

IVM imaging allows for simultaneous labeling of various TME components and can be used to
study live their association with tumor cell behavior. Indeed, we found that TAMMs and blood
vessels were associated with specific behavioral phenotypes, showing the power of
BEHAV3D-TP *small-scale phenotyping module* in identifying novel regulators of cellular
behavior. This is exemplified by our finding that *slow-retreating* DMG cells were found in
environments enriched for SR101$^+$ cells, an observation that has not been made before. To take
further advantage of BEHAV3D-TP, future work should be directed toward identifying more
TME components, including cells of the adaptive immune system. For example, it would be
interesting to use our pipeline on recently developed immunocompetent mouse models of DMG
[50,51], or in mice with a humanized immune system[52].

To enhance the accessibility of BEHAV3D-TP for researchers keen on studying tumor cell
behavioral profiles through live imaging, we have incorporated our software in a Google Colab
notebook, allowing researchers with minimal programming skills to employ the tool online
without the requirement to install specific software environments nor the need to directly
interact with the code. The user can easily install the necessary dependencies, set up the working
environment, process datasets and adjust parameters without requiring deep coding skills.
Google Colab provides the necessary computational resources to be able to run the pipeline
without demanding a powerful local machine. However, there are RAM usage and runtime
limitations that could affect the analysis, given a big enough dataset in the free version of

Google Colab. Given this and data privacy concerns, the workflow can also be run locally via

a Jupyter Notebook, that can be downloaded from the Google Colab notebook. Aligned with

the goals of BEHAV3D-TP, other methods utilizing the Google Colab framework are emerging,

such as CellTracksColab[24], which is a general tool for tracking data analysis. While

CellTracksColab offers valuable insights into tracking data, BEHAV3D-TP is specifically

designed for IVM data and includes additional features. These features include directionality

metrics crucial for invasion studies, detailed analysis of identified behavioral populations, data

back-projection and correlation of these behaviors with both large-scale and small-scale TME

features. This comprehensive approach provides a deeper understanding of the TME drivers

behind intratumoral heterogeneity, making BEHAV3D-TP a robust tool for researchers

investigating the complex behaviors of tumor cells in their natural environments.

In summary, BEHAV3D-TP allows to get insights into the heterogeneity of tumor cell

migration behavior in an accessible way; however, it would be exciting to extend the application

to other behaviors, such as therapy response to study TME-mediated drug resistance. Such an

endeavor may benefit from including additional parameters beyond kinetic features. For

example, others have additionally used morphological parameters [19,21], and we envision that a

multiparametric analysis based on a combination of morpho-kinetic features may further refine

behavioral phenotyping.

METHODS

Animal material

NOD.Cg-PrkdcscidIl2rgtm1Wjl/SzJ (NSG) mice were purchased from Charles River Laboratories (France). Experiments were conducted per Institutional Guidelines under acquired permission from the local Ethical Committee and as per current Dutch laws on Animal Experimentation. Mice were housed under sterile conditions using an individually ventilated cage (IVC) system and were fed with sterile food and water.

Human DMG sphere-forming culture

DMG007 (HSJD-DIPG-007) cells established from primary DMG material were grown in a base medium (TSM) consisting of 48% DMEM/F12 and 48% Neurobasal medium (Thermo Fisher) supplemented with GlutaMAX, 100mM Sodium Pyruvate, MEM non-essential amino acids and 1M Hepes buffer, at 1% each. Primocin (InvivoGen) at 50mg/ml was additionally added to the TSM. Working medium was prepared by adding 20ng/ml of human EGF and human basic FGF, 10ng/ml of PDGF-AA and PDGF-BB (Peprotech) and 2ug/ml of Heparin (StemCell Technologies). DMG cells were cultured at 37 °C in a humidified atmosphere with 5% CO_2.

Fluorescent reporter cloning and lentiviral transduction

Fluorescent reporter construct was based on the combination of Histone2B-mNeonGreen, P2A and mScarlet I-CAAX gifted from Dr. Hugo Snippert (UMC Utrecht, The Netherlands), and inserted by In-Fusion HD Cloning Plus (Takarabio) into a lentiviral vector including cPPT/CTS, WPRE, an EF1a promoter and IRES-puromycin-resistance cassette[11]. Subsequently the above-mentioned elements were amplified by PCR and the amplicon was assembled in the final lentiviral backbone using Gibson Assembly (NEB). Sequences were validated using Sanger sequencing. DMG007 cells were transduced using a standard lentiviral transduction protocol and selected using puromycin to achieve a stable cell line.

Cranial imaging window (CIW) surgery and tumor cell injection

CIW surgery and tumor cell injection were performed on the same day as previously described [7,10]. Briefly, mice were sedated with Hypnorm (Fluanison [neuroleptic] + Fentanyl [opioid]) (0.4 ml/kg) + Midazolam [benzodiazepine sedative] (2 mg/kg) at a dose of 1:1:2 in sterile water and mounted in a stereotactic frame. The head was secured using a nose clamp and two custom-made 3D printed ear bars, made out of Polylactic Acid (PLA, Geeetech, 700-001-043). The

head was shaved, and the skin was cut circularly. The mouse was placed under a stereo-microscope to ensure precise surgical manipulation. The periosteum was scraped, and a circular groove of 5 mm diameter was drilled over the right parietal bone. The bone flap was lifted under a drop of cortex buffer (125 mM NaCl, 5 mM KCl, 10 mM glucose, 10 mM HEPES buffer, 2 mM MgSO$_4$, and 2 mM CaCl$_2$, pH 7.4), and the dura mater was removed. Gelfoam sponge (Pfizer) was used to stop potential bleedings. Next, 1×10^5 DMG007-H2B-mNeonGreen cells resuspended in 3 μl phosphate-buffered saline (PBS) were injected stereotactically using a 5 μl Hamilton syringe with a 2 pt style in the middle of the craniotomy at a depth of 0.5 mm. The exposed brain was sealed with silicone oil and a 6 mm coverslip was glued on top. Dental acrylic cement (Vertex) was applied on the skull surface, covering the edge of the coverslip, and a custom-made 3D printed ring made of biocompatible Polylactide acid (PLA, Geeetech, 700-001-0433) was glued around the coverslip to provide fixation during intravital imaging and to serve as a reservoir for a water drop required for the objective of the microscope. A single dose of 100 μg/kg of buprenorphine (Temgesic, BD pharmaceutical limited) was administered for post-surgical pain relief. Mice were closely monitored twice per week to assess behavior, reactivity, and appearance.

Intravital imaging

Mice were intravitally imaged as previously described [7,10]. In short, were sedated with isoflurane and placed face-down in a stereotactic frame. The Polylactide ring was used to fix the mouse's head to the frame with custom-made Polylactide bars. Time-lapse images of the entire tumor volume were acquired for a maximum of 5.4 hours. The minimal time interval between serial images was set to 20 minutes. For tile scans, images of the complete z-stack of the tumor were acquired, with a step size of 2 μm. In a group of 3 mice, oligodendrocytes and astrocytes were imaged by intravenous injection of 50 μL SR101 (Thermofisher) at a concentration of 5 mM dissolved in PBS. In a different group of 3 mice, TAMMs were imaged by intravenous injection of 50 μL 100 μM CDr20 [40]. Simultaneously with the injection of either SR101 or CDr20, all mice received an intravenous injection of 10 ul CD31-AF647 antibody (ThermoFisher, A14716) to visualize the blood vessels.

Intravital imaging was performed on an upright FVMPE-RS multiphoton microscope (Olympus) equipped with a MaiTai DeepSee and an InSight DeepSee laser (Spectra-Physics), a 25x/1.05 numerical aperture water objective, and two GaAsP and two Multialkali photomultiplier tubes (Olympus). Images were acquired with simultaneous imaging with the MaiTai (960 nm) and Insight (1100 nm) lasers. Green, red, and far-red fluorescence was

separated by 570 nm and 650 nm dichroic mirrors, and 610/70 nm and 705/90 nm filters
(Semrock) and collected by GaAsP or Multialkali photomultiplier tubes. Scanning was
performed in a bidirectional mode with a resonant scanner at 400 Hz and 12 bit, with a zoom of
1x, 512×512 pixels. For optimal detection of blood vessels (CD31-AF647), a single tile scan
was made at the last timepoint with Insight laser tuned at 1250 nm. At the postprocessing step,
this image was co-registered to the time-lapse movie.

Co-registration of IVM imaged blood vessel imaging

CD31-AF647 tile scan featuring optimal visualization of blood vessels during live imaging was
co-registered to the last timepoint of the time-lapse movie. Both images (last timepoint of time-
lapse tile-scan and blood vessel tile-scan) were visually inspected to determine and annotate the
internal landmarks using Imaris (Oxford Instruments), versions 9.5 to 9.6. These landmarks
served as a reference for the co-registration software to overlap both images. To perform the
co-registration, a modification of *elastix*[53] was used; a python-based open-source software
based on Insight Segmentation and Registration Toolkit (ITK) plus SimpleElastix[54] : a
collection of different algorithms used on medical image registration. Due to the extensive
number of potential parameter combinations, specific parameters were manually chosen. This
selection was conducted using the Slicer3D visualization software, version 5.2.2.

Immunohistochemistry

3D immunohistochemistry was performed as described before [29]. After the live imaging session,
mouse brains were immersed in 5 ml 4% paraformaldehyde (PFA) pH 7.4 overnight on ice.
After fixation, brains were blocked in 5-10 ml Wash Buffer 1 (2 ml Tween-20, 2 ml Triton X-
100, 2 ml 10% SDS, and 2 g BSA in 1 l PBS) for 5 hours. For multiplex immunolabeling, a
two- or three-round staining protocol was used as previously described [28,29]. Washing and
incubation steps were performed in Wash Buffer 2 (1 ml Triton X-100, 2 ml 10% SDS, and 2
g BSA in 1 l PBS). A thick (1 mm) slice of the cortex part that was under the cranial imaging
window was dissected with a scalpel before proceeding to immunolabeling. The following
primary antibodies were used: rabbit anti-Olig2 1:100 (AB9610, Merk), goat anti-Iba1 1:200
(NB100-1028SS, NovusBio) and CD31-AF647 1:250 (A14716, ThermoFisher). As secondary
antibodies, donkey anti-rabbit AF405 1:250 (ab175651, Abcam) and donkey anti-goat AF633
1:250 (A21082, ThermoFisher) were used. After the last washing step, tissues were optically
cleared by three stepwise incubations of an increasing concentration of FUnGI [55] clearing agent

(25%/50%/75%, diluted in PBS) for 1h at room temperature [28]. The final incubation step with
100% FUnGI was performed overnight at 4°C.

mLSR3D imaging

mLSR3D imaging [28,29] of thick slices of fixed mouse cortex was performed using a Zeiss
LSM880 system equipped with a 32-channel spectral detector. The signal from all fluorophores
present in the samples was collected in a single acquisition and linear unmixing was used to
obtain separate channels. Unmixing was carried out by the Online Fingerprinting mode of the
microscope. Pre-acquired references of each single fluorophore were used to enable
multispectral on-the-fly unmixing. Imaging was performed using a 25x multi-immersion
objective 0.8 NA (Zeiss 420852-9871) with a working distance of 570µm. Tile scans were
acquired using a voxel size of 0.332x0.332x1.200 µm, a pixel dwell of 2µs, and a 10% tile
overlap.

Image processing

391 For 3D visualization, shift correction, object rendering, and cell tracking, time-lapse movies
392 were processed with Imaris (Oxford Instruments), versions 9.5 to 10.1.

Cell tracking

The *Channel Arithmetics* Xtension was used for channel unmixing of overlapping signals. The
Spots ImarisTrack module was used for object detection and semi-automated tracking of tumor
cells (autoregressive motion). Tumor cell tracks were manually corrected to ensure accurate
tracking. Around 50-100 cells per position were manually corrected and labeled as such for
posterior selection of accurate tracks. To determine the direction of tumor cell movement
relative to the tumor core, at the image edge closer to the tumor core a "Tumor edge" surface
object was manually rendered using the contour tool. The *Distance Transformation* Xtension
was used to measure the distance between tumor cells and the "Tumor edge". For tracked tumor
cells, time-lapse data containing the coordinates of each cell, the values of cell speed, mean
square displacement, displacement delta length, displacement length, and distance to "Tumor
edge" were exported. See https://qbi.uq.edu.au/files/40820/ImarisManual.pdf for statistical
details.

Microenvironmental factors rendering

407 For rendering the microenvironmental factors detected during IVM or by post-IVM LSR-3D
408 imaging, the Surface and Spots modules of Imaris were used. Surfaces were used for the SR101,

409 CDr20, Olig2, DAPI, and Iba1 stainings, and Spots were used for CD31+ protruding or
410 elongated structures, as previously described [30]. For rendered structures, positional data
411 containing the coordinates of objects were exported.

Tumor large-scale spatial phenotyping with Cytomap

LSR-3D imaging data was processed for large-scale spatial phenotyping using the Cytomap
Spatial Analysis Toolbox [30]. Briefly, neighborhoods with a 100 µm radius were generated.
Neighborhoods were then clustered into 2 regions based on DMG cell positioning using the
David Balwin algorithm. Using the gating tool, the tumor regions (containing DMG objects)
were defined. Next, the neighborhoods were clustered again into 3 different regions based on
the distribution of Iba1, Olig2, and CD31: *Void, TAMM/Oligo,* and *TAMM/vascularized*. By
using the *Manually Defined Regions* option, only the tumor region was classified. Finally, for
each region, fold-change values were calculated and exported using the *Region Statistics* tab.
A 3D image of the cluster distribution was exported for each sample (Figure supplement 2a)
and then used to visually assign phenotypical regions in the IVM movies (Figure 3c). For each
mouse, one or two positions for each type of region were cropped and analyzed for tumor cell
behavior as described above (*Cell tracking*).

BEHAV3D Tumor Profiler framework

BEHAV3D-TP was developed using the R Studio version 4.3.3. It features three modules: 1)
Heterogeneity module with optional 2) *Large-scale and/or* 3) *Small-scale phenotyping*
modules. All of the modules are included in the same script available in Github
[https://github.com/imAIgene-Dream3D/BEHAV3D_Tumor_Profiler], but can be run
independently from one another. To acquire this independency, the optional modules are
directly combined to the necessary sections of the *Heterogeneity module*. Consequently, the
user can use whichever module is best suited for their purposes. In order to further facilitate
public access to the BEHAV3D-TP, a Google Colab notebook has also been created accessible
through the same Github page [https://github.com/imAIgene-
Dream3D/BEHAV3D_Tumor_Profiler]. This environment is friendly to anyone regardless of
their coding expertise as it has a step-by-step follow-through execution through the pipeline.
Nevertheless, a Wiki demo run and a video tutorial (Supplementary Video 1) are also provided
through the GitHub page [https://github.com/imAIgene-
Dream3D/BEHAV3D_Tumor_Profiler] for further clarification, if necessary. In this Colab
notebook, the user can easily import data, modify analysis parameters, execute the pipeline and

export the obtained results without any computational power demands or computational
knowledge. Additionally, the original R version and R Markdown demo scripts are also
provided.

Heterogeneity module.

To account for missing time points and to create a time series with the same time interval for
each tumor cell time series, linear interpolation was used to estimate the values of missing time
points. To compare time series independently, tracks were cropped to the minimal common
length among all tracks. In our case, to a 2.6 hour duration. For tracked tumor cells, time-lapse
data containing the coordinates of each cell, the values of cell speed, mean square displacement,
displacement delta length, displacement length, and relative distance to "Tumor edge" were
used to calculate a cross-distance matrix between time series. First, we performed a principal
component (PC) analysis and selected the number of PCs that explained at least 90% of data
variance (3).

The previously calculated PCs were then used to compute the cross-distance matrix between
each tumor cell multivariate time series using the dynamic time warping algorithm from the
package "dtwclust". To visualize heterogenous tumor cell behaviors in 2 dimensions,
dimensionality reduction on the cross-distance matrix was performed by the Uniform Manifold
Approximation and Projection method ("umap" package). Clustering was performed using the
k-means clustering algorithm. For each cluster, the average values of distinct behavioral and
environmental components were calculated and plotted as a heatmap ("pheatmap").

Large-scale phenotyping module.

462　The information obtained from mLSR3D and Cytomap about large-scale TME regions were
463　combined with the DMG cells behavioral information to examine the large-scale features of
464　tumor cells. For tumor cells, the mean speed, square displacement and raw movement were
465　analyzed.

Small-scale phenotyping module.

The positional coordinates of tracked tumor cells and detected microenvironmental factors were
used to determine the small-scale landscape of tumor cells. For tumor cells, the average distance
to the closest 10 neighbors was measured to determine tumor cell density. For each tumor cell
and each environmental component, the number of microenvironmental objects in a 30-μm
radius and the distance to the closest object were computed. For each behavioral cluster, the

values of closest neighbors, minimal distance to each environmental factor, and number of
environmental objects in the 30 µm radius were plotted and compared.

Statistical analyses

Statistical analyses were performed using R. Results are represented as mean ± s.e.m. unless
indicated otherwise; n represents independent positions in 6 biological replicates. Two-tailed
unpaired t-tests were performed between two groups unless indicated otherwise. To compare
frequencies of different behavioral signatures between large scale regions identified, a Tukey
Honestly Significant Difference (HSD) adjusted Estimated marginal means (emmeans)
pairwise comparison test was applied. To compare the distribution of behavioral clusters
between the environmental clusters, a mixed linear regression followed by a one-way ANOVA
was performed. To compare the distribution of different behavioral and environmental features
between the behavioral clusters, a one-way ANOVA followed by Tukey HSD correction was
performed.

Data availability

486 Data will be made publicly available after publication.

Code availability

We provide the BEHAV3D-TP framework in a Google Colab user-friendly environment
[https://colab.research.google.com/drive/1JI7ysqFf3tvdi6Df4YUsSZ8RbuXw8wba?usp=shari
ng], as well as in an R script format in Github [https://github.com/imAIgene-
Dream3D/BEHAV3D_Tumor_Profiler]. Additionally, a Wiki [https://github.com/imAIgene-
Dream3D/BEHAV3D_Tumor_Profiler/wiki] to follow through a demo dataset as well as a
video tutorial (Supplementary Video 1) has been developed for further assistance.

Acknowledgements

We thank the Princess Máxima Center for Pediatric Oncology for technical support and the
Hubrecht Institute and Zeiss for imaging support and collaboration. All imaging was performed
at the Princess Máxima Imaging Center. We thank D. van Vuurden (Princess Máxima Center
for Pediatric Oncology) and Dr. Ángel Montero Carcaboso (Sant Joan de Déu Barcelona
Hospital) for providing primary DMG cultures; Dr Young-Tae Chang (Postech, Korea) for
providing us the CD20r dye; Sandra Fernandez Archidona for her assistance with IVM data
visualization; and members of the Dream3DLAB (Rios group) and imAIgene-lab (Alieva
group) for offering critical feedback on the project. A.Z. was supported by a Veni fellowship

from the Netherlands Organization for Scientific Research (09150161910076). R.C. and A.C.R
were supported by an ERC-starting grant 2018 project (no. 804412). This work was produced
with the support of a 2023 Leonardo Grant for Scientific Research and Cultural Creation,
BBVA Foundation (LEO23-2-10305-BBM-BAS-144). M.A was supported by the Comunidad
de Madrid (2022-T1/BMD-24021).

FIGURE LEGENDS

Fig. 1: BEHAV3D Tumor Profiler pipeline. Schematic representation of the workflow showing data preparation, input and outputs in each of the three modules. The programmatic processing is represented with a yellow background **a,** Time-lapse image processing and preparation, based on segmentation and tracking of tumor and labelled TME cells. Upload to Google Drive is optional but recommended to work in the Google Colab framework. **b,** Heterogeneity Module provides distinct behavioral clusters for DMG cells, and relates them to dynamic features and provides a back-projection of the tracked cells. **c, d,** Optional modules, additional to the Heterogeneity module. **c,** Large-scale phenotyping module combines TME large-scale information with DMG behavioral profiling to further depict population frequencies in distinct TME regions. **d,** Small-scale phenotyping module combines TME small-scale information with DMG behavioral profiling to depict how the cellular environment can influence tumor cell behavior.

Fig. 2: Behavioral profiling of DMG cells with BEHAV3D-TP heterogeneity module. a, Representative intravitally imaged position showing DMG cells tracked relative to tumor Edge (left panel). Tracks were classified according to DMG behavior (right, color coded by cluster). Representative of $n = 18$ independent positions. **b,** UMAP plot showing seven color-coded DMG cell behavioral clusters identified by BEHAV3D-TP. Each datapoint represents one T cell track. **c,** Representative intravitally imaged positions and enlarged sections for showcasing distinct DMG behavioral clusters *Invading* (yellow), *Retreating* (green), *Erratic* (dark purple) and *Static* (light purple) through the time series. $n = 18$ independent experiments. Scale bars 10μm. **d,** Heatmap depicting relative values of DMG cell features indicated for each cluster, named according to their most distinct characteristics. Arbitrary units in respect to maximal and minimal values for each feature. Bubble size according to significance levels, all features represented have a significance level of P < 0.001 (***): *mean_displ2, mean squared displacement; speed; delta displacement; displacement length; persistence; invasion*. P-valuves represent the significance of differences across clusters tested through ANOVA for each feature. **e, f,** UMAP representation of DMG cells' velocity relative to the tumor edge(μm)

(**e**) and speed (µm/h) (**f**). Each datapoint represents one T cell track and is colored following a

color gradient. For each feature colors represent the mean DMG track values.

Fig. 3: Mapping distinct DMG behavioral patterns to distinct TME regions. a, Overview

of the large-scale TME region phenotyping workflow. After IVM imaging session, the tissue is

fixed and stained and then analyzed using mLSR-3D. CytoMAP Spatial Analysis Toolbox is

used to identify distinct large-scale regions, that are subsequently matched to IVM imaging

dataset. **b,** Representative fixed multispectral image of a mLSR3D imaged DMG tumor and

zoomed-in images of TME large-scale regions: *Void* (blue outline), *TAMM/vascularized* (red

outline) and *TAMM/Oligo* (yellow outline). DMG nuclei (blue), CD31 (pink), Olig2 (purple),

and Iba1 (yellow) are represented on the imaged. Scale bars, 100 µm (overview), 30 µm

(zoomed-in regions). **c,** 3D multispectral image DMG cell rendering color-coded per large TME

region: *Void* (blue), *TAMM/Oligo* (yellow) and *TAMM/vascularized* (red) (left). Regions were

mapped on intravital imaging data (right). **d,** DMG single-cell behavioral cluster frequencies

among the TME large-scale regions identified with CytoMAP. *Slow retreating,* P < 0.05; *Static,*

P < 0.05; *Invading,* P < 0.05. P-values represent the significance of differences across TME

regions tested through ANOVA for each behavioral cluster.

Fig. 4: The BEHAV3D-TP small-scale phenotyping correlates DMG behavioral profiles

with microenvironmental composition at a single-cell resolution. a, Overview of the small-

scale TME phenotyping module. During IVM imaging, information about spatial distribution

of SR101+ (purple), CD20r+ (purple) and CD31+ (red) cells is collected and processed by the

BEHAV3D-TP *small-scale phenotyping module*. Single DMG cell spatial TME features like

distance to its neighbors, number of cells in a radius or minimal distance to a cell type were

measured. **b,** Heatmap depicting DMG cell small-scale niches' features across distinct DMG

behavioral clusters. Arbitrary units in respect to maximal and minimal values for each feature.

Bubble size according to significance levels, from smallest (P < 0.05 (**)) to medium (P < 0.01

(**)) to biggest (P < 0.001 (***)): *dist_10_cell*, Distance to nearest 10 cells, P < 0.01 (**);

n_SR101, number of SR101 cells, P < 0.05 (*); *min_SR101,* minimal distance to a SR101 cell,

P < 0.05 (*); *n_CD20r,* number of CD20r cells, P < 0.05 (*); *min_Cd20r,* minimal distance to

a CD20r cell, P < 0.01 (**); *min_BV_dist,* minimal distance to a Blood vessel, P < 0.001 (***).

P-values represent the significance of differences across clusters tested through ANOVA for

each feature. **c, e,** Boxplot depicting the *Minimal distance to a CD20r+ TAMM (µm)* (**c**) and

the *Minimal distance to a blood vessel (µm)* (**e**) across DMG behavioral clusters. **c,** *Invading*

versus *Erratic,* P < 0.01 (**); *Invading* versus *Slow*, P < 0.05 (*). **e,** *Static* versus *Erratic*, P <

0.01 (**); *Static* versus *Retreating*, P < 0.001 (***); *Static* versus *Invading*, P < 0.001 (***);

Slow versus *Retreating*, P < 0.1 (·); *Slow invading* versus *Retreating*, P < 0.1 (·); *Slow* versus

Invading, P < 0.1 (·); *Slow invading* versus *Invading*, P < 0.1 (·). **d,** Fixed multispectral image

showing the movement of DMG cells (grey) relative to CD20r+ TAMM (blue) and the Tumor

core. The trajectories show a path from the initial (blue) to the final position (red). Scale bar,

10 µm. **f,** Time lapse multispectral images (left to right) showing the DMG cells (blue)

displacement along CD31 blood vessel (purple) away from the tumor edge. Scale bars, 30 µm.

Figure supplement 1: DMG cells heterogeneity captured by intravital imaging. a,

Schematic representation of an intravital imaging experimental design, capturing timelapses up

to 5.4 hours of DMG cells expressing H2BmNeonGreen and mScarlet proteins, followed by

single-cell tracking. **b,** Representative intravitally imaged position showing DMG cell tracks

colored by behavioral cluster over the last timepoint of the original IVM time-lapse. **c,**

Quantification of average DMG cell speed (µm/h) in each identified behavioral cluster. **d,**

Behavioral cluster frequency distribution across IVM imaged mice. **e,** Behavioral cluster

average distribution in IVM imaged mice, shown in percentage across mice. Representative of

n=18 independent positions.

Figure supplement 2: Cell type and behavioral cluster frequency distribution in identified

TME large-scale regions. a, Representative CytoMap output showcasing**,** 3D spatial

distribution of DMG cells in distinct TME large-scale regions: *Void* (blue), *TAMM/Oligo*

(yellow) and *TAMM/vascularized* (red). Each datapoint represent a unique DMG cell. **b, c,** Cell

count fold change of DMG, Iba1+, Olig2+ and CD31+ cell types relative to the mean cell count

in each TME large scale region (*Void, TAMM/Oligo, TAMM/vascularized)*, both in boxplot (**b**)

and table (**c**) formats. In DMG cell count: *TAMM/Oligo* versus *Void,* P < 0.01 (**);

TAMM/Oligo versus *TAMM/vascularized*, P < 0.05 (*). In Iba1 cell count: *TAMM/Oligo* versus

Void, P < 0.001 (***); *TAMM/vascularized* versus *Void*, P < 0.001 (***). In Olig2 cell count:

TAMM/Oligo versus *Void,* P < 0.001 (***); *TAMM/Oligo* versus *TAMM/vascularized*, P <

0.001 (***). In CD31 cell count: *TAMM/Oligo* versus *Void,* P < 0.001 (***);

TAMM/vascularized versus *Void,* P < 0.001 (***); *TAMM/Oligo* versus *TAMM/vascularized*,

P < 0.001 (***). **d,** Boxplots showing the frequencies of *Invading* (left), *Slow retreating*

(middle), and *Static* (right) DMG behavioral clusters in *Void, TAMM/Oligo and*

TAMM/vascularized TME large-scale regions. The cell proportion was normalized per mouse.

In *Invading* cluster: *TAMM/vascularized* versus *Void*, P < 0.01 (**). In *Slow retreating* cluster:

TAMM/Oligo versus *Void*, P < 0.01 (**); *TAMM/Oligo* versus *TAMM/vascularized,* P < 0.01

(**). In *Static* cluster: *Void* versus *TAMM/vascularized,* P < 0.05 (*). Each point represents a

different position imaged. **e, f, g,** Violin plots showing DMG single cell features (*Tumor cell*

speed (µm/hour) (**e**), *Average speed displacement* (**f**) *and Tumor cell velocity* (µm/hour)(**g**))

differences among *Void, TAMM/Oligo and TAMM/vascularized* TME large-scale regions. **e,**

TAMM/vascularized versus *Void*, P < 0.01 (**); *TAMM/vascularized* versus *TAMM/Oligo*, P <

0.01 (**). **g,** Tumor cell velocity can be either towards brain core (down, negative values) or

towards brain parenchyma (up, positive values).

Figure supplement 3: Cell distribution and behavioral clusters in small-scale TME. a, b,

Representative 3D IVM rendering of *DMG nuclei* (cyan), CD31+ cells (grey) and SR101+ (**a**,

red) or CD20r+ cells (**b**, red). Scale bars, 100 µm. **c, d,** Boxplot showing *Average distance to*

the 10 closest DMG neighbors (µm) (**c**) and *Number of SR101+ cells in a 30 µm radius* (**d**)

across distinct DMG single-cell behavioral clusters. **c,** *Invading* versus *Slow*, P < 0.05 (*);

Invading versus *Slow retreating*, P < 0.01 (**). **d,** *Slow retreating* versus *Static*, P < 0.01 (**);

Slow retreating versus *Slow*, P < 0.1 (·); *Slow retreating* versus *Slow Invading*, P < 0.1 (·); *Slow*

retreating versus *Invading*, P < 0.1 (·).

Supplementary video 1: BEHAV3D Tumor Profiler pipeline main characteristics and

walkthrough. Summary video of BEHAV3D TP pipeline.

Supplementary video 2: Representative intravital time-lapse video of DMG cells. Raw

DMG cells time-lapse (right panel). Cell tracks were classified according to DMG behavior

(right panel, color coded by cluster). Representative movie from n=18 independent positions

imaged in 6 mice. Scale bars 40 µm.

Supplementary video 3: Dynamic multispectral representative time-lapse video of DMG

cells and CD20r+ TAMM. Time-lapse showing the movement of DMG cells (grey) relative

to CD20r+ TAMM (blue). The trajectories towards the TAMM of two DMG cells show a path

from the initial (blue) to the final position (red). Left: Tumor edge; right: tumor core.

Representative movie from n=18 independent positions imaged in 6 mice. Scale bars 10 µm, 7

µm after zoom in.

REFERENCES

1. Vitale, I., Shema, E., Loi, S. & Galluzzi, L. Intratumoral heterogeneity in cancer progression and response to immunotherapy. *Nat Med* **27**, 212–224 (2021).

2. Lawson, D. A., Kessenbrock, K., Davis, R. T., Pervolarakis, N. & Werb, Z. Tumour heterogeneity and metastasis at single-cell resolution. *Nat. Cell Biol.* **20**, 1349–1360 (2018).

3. Elhanani, O., Ben-Uri, R. & Keren, L. Spatial profiling technologies illuminate the tumor microenvironment. *Cancer Cell* **41**, 404–420 (2023).

4. Scheele, C. L. G. J., Maynard, C. & Rheenen, J. van. Intravital Insights into Heterogeneity, Metastasis, and Therapy Responses. *Trends Cancer* **2**, 205–216 (2016).

5. Entenberg, D., Oktay, M. H. & Condeelis, J. S. Intravital imaging to study cancer progression and metastasis. *Nat. Rev. Cancer* **23**, 25–42 (2023).

6. Alieva, M., Ritsma, L., Giedt, R. J., Weissleder, R. & Rheenen, J. van. Imaging windows for long-term intravital imaging. *IntraVital* **3**, e29917 (2014).

7. Alieva, M. *et al.* Intravital imaging of glioma border morphology reveals distinctive cellular dynamics and contribution to tumor cell invasion. *Sci. Rep.* **9**, 2054 (2019).

8. Alieva, M. & Rios, A. C. Longitudinal Intravital Imaging of Brain Tumor Cell Behavior in Response to an Invasive Surgical Biopsy. *J. Vis. Exp.* (2019) doi:10.3791/59278.

9. Zomer, A., Croci, D., Kowal, J., Gurp, L. van & Joyce, J. A. Multimodal imaging of the dynamic brain tumor microenvironment during glioblastoma progression and in response to treatment. *iScience* **25**, 104570 (2022).

10. Alieva, M. *et al.* Preventing inflammation inhibits biopsy-mediated changes in tumor cell behavior. *Sci. Rep.* **7**, 7529 (2017).

11. Janssen, A., Beerling, E., Medema, R. & Rheenen, J. van. Intravital FRET Imaging of Tumor Cell Viability and Mitosis during Chemotherapy. *PLoS ONE* **8**, e64029 (2013).

12. Croci, D., Zomer, A., Kowal, J. & Joyce, J. A. Cranial imaging window implantation technique for longitudinal multimodal imaging of the brain environment in live mice. *STAR Protoc.* **4**, 102197 (2023).

13. Zomer, A. *et al.* Brief Report: Intravital Imaging of Cancer Stem Cell Plasticity in Mammary Tumors. *STEM CELLS* **31**, 602–606 (2013).

14. Zomer, A. *et al.* In Vivo Imaging Reveals Extracellular Vesicle-Mediated Phenocopying of Metastatic Behavior. *Cell* **161**, 1046–1057 (2015).

15. Chen, N. *et al.* Neutrophils Promote Glioblastoma Tumor Cell Migration after Biopsy. *Cells* **11**, 2196 (2022).

16. Chen, N., Ritsma, L. M. A. & Vrisekoop, N. In vivo characteristics of human and mouse breast tumor cell lines. *Exp. Cell Res.* **381**, 86–93 (2019).

17. Bayarmagnai, B., Perrin, L., Pourfarhangi, K. E. & Gligorijevic, B. Cell Migration, Methods and
Protocols. *Methods Mol. Biol.* **1749**, 175–193 (2018).

18. Shannon, M. J., Eisman, S. E., Lowe, A. R., Sloan, T. & Mace, E. M. cellPLATO: an unsupervised
method for identifying cell behaviour in heterogeneous cell trajectory data. *J. Cell Sci.* **137**, jcs261887.

19. Crainiciuc, G. *et al.* Behavioural immune landscapes of inflammation. *Nature* **601**, 415–421 (2022).

20. Dekkers, J. F. *et al.* Uncovering the mode of action of engineered T cells in patient cancer organoids.
Nat. Biotechnol. **41**, 60–69 (2023).

21. Freckmann, E. C. *et al.* Traject3d allows label-free identification of distinct co-occurring phenotypes
within 3D culture by live imaging. *Nat. Commun.* **13**, 5317 (2022).

22. Alieva, M. *et al.* BEHAV3D: a 3D live imaging platform for comprehensive analysis of engineered
T cell behavior and tumor response. *Nat. Protoc.* 1–33 (2024) doi:10.1038/s41596-024-00972-6.

23. Buczkowicz, P., Bartels, U., Bouffet, E., Becher, O. & Hawkins, C. Histopathological spectrum of
paediatric diffuse intrinsic pontine glioma: diagnostic and therapeutic implications. *Acta Neuropathol.*
128, 573–581 (2014).

24. Gómez-de-Mariscal, E. *et al.* CellTracksColab — A platform for compiling, analyzing, and
exploring tracking data. *bioRxiv* 2023.10.20.563252 (2024) doi:10.1101/2023.10.20.563252.

25. Hidalgo-Cenalmor, I. *et al.* DL4MicEverywhere: deep learning for microscopy made flexible,
shareable and reproducible. *Nat. Methods* **21**, 925–927 (2024).

26. Chamier, L. von *et al.* Democratising deep learning for microscopy with ZeroCostDL4Mic. *Nat*
Commun **12**, 2276 (2021).

27. Persson, M. L. *et al.* The intrinsic and microenvironmental features of diffuse midline glioma:
Implications for the development of effective immunotherapeutic treatment strategies. *Neuro-Oncol.* **24**,
1408–1422 (2022).

28. Ineveld, R. L. van *et al.* Multispectral confocal 3D imaging of intact healthy and tumor tissue using
mLSR-3D. *Nat. Protoc.* **17**, 3028–3055 (2022).

29. Ineveld, R. L. van *et al.* Revealing the spatio-phenotypic patterning of cells in healthy and tumor
tissues with mLSR-3D and STAPL-3D. *Nat. Biotechnol.* **39**, 1239–1245 (2021).

30. Stoltzfus, C. R. *et al.* CytoMAP: A Spatial Analysis Toolbox Reveals Features of Myeloid Cell
Organization in Lymphoid Tissues. *Cell Rep.* **31**, 107523 (2020).

31. Winkler, F. *et al.* Imaging glioma cell invasion in vivo reveals mechanisms of dissemination and
peritumoral angiogenesis. *Glia* **57**, 1306–1315 (2009).

32. Wallmann, T. *et al.* Microglia Induce PDGFRB Expression in Glioma Cells to Enhance Their
Migratory Capacity. *iScience* **9**, 71–83 (2018).

33. Seano, G. & Jain, R. K. Vessel co-option in glioblastoma: emerging insights and opportunities.
Angiogenesis **23**, 9–16 (2020).

34. Ravin, R. *et al.* Perivascular invasion of primary human glioblastoma cells in organotypic human
brain slices: human cells migrating in human brain. *J. Neuro-Oncol.* **164**, 43–54 (2023).

35. Gupta, R. K. *et al.* Tumor-specific migration routes of xenotransplanted human glioblastoma cells
in mouse brain. *Sci. Rep.* **14**, 864 (2024).

36. Nimmerjahn, A., Kirchhoff, F., Kerr, J. N. D. & Helmchen, F. Sulforhodamine 101 as a specific
marker of astroglia in the neocortex in vivo. *Nat. Methods* **1**, 31–37 (2004).

37. Appaix, F. *et al.* Specific In Vivo Staining of Astrocytes in the Whole Brain after Intravenous
Injection of Sulforhodamine Dyes. *PLoS ONE* **7**, e35169 (2012).

38. Rasmussen, R., Nedergaard, M. & Petersen, N. C. Sulforhodamine 101, a widely used astrocyte
marker, can induce cortical seizure-like activity at concentrations commonly used. *Sci. Rep.* **6**, 30433
(2016).

39. Hagos, L. & Hülsmann, S. Unspecific labelling of oligodendrocytes by sulforhodamine 101 depends
on astrocytic uptake via the thyroid hormone transporter OATP1C1 (SLCO1C1). *Neurosci. Lett.* **631**,
13–18 (2016).

40. Kim, B. *et al.* Visualizing Microglia with a Fluorescence Turn-On Ugt1a7c Substrate. *Angew. Chem.*
Int. Ed. **58**, 7972–7976 (2019).

41. Liu, I. *et al.* The landscape of tumor cell states and spatial organization in H3-K27M mutant diffuse
midline glioma across age and location. *Nat. Genet.* **54**, 1881–1894 (2022).

42. Xin, T. *et al.* Oncogenic Kras induces spatiotemporally specific tissue deformation through
converting pulsatile into sustained ERK activation. *Nat. Cell Biol.* **26**, 859–867 (2024).

43. Venkataramani, V. *et al.* Glioblastoma hijacks neuronal mechanisms for brain invasion. *Cell* **185**,
2899-2917.e31 (2022).

44. Bera, K. *et al.* Extracellular fluid viscosity enhances cell migration and cancer dissemination. *Nature*
611, 365–373 (2022).

45. Gertler, F. & Condeelis, J. Metastasis: tumor cells becoming MENAcing. *Trends Cell Biol.* **21**, 81–
90 (2011).

46. Kluiver, T. A., Alieva, M., Vuurden, D. G. van, Wehrens, E. J. & Rios, A. C. Invaders Exposed:
Understanding and Targeting Tumor Cell Invasion in Diffuse Intrinsic Pontine Glioma. *Front. Oncol.*
10, 92 (2020).

47. Quail, D. F. & Joyce, J. A. The Microenvironmental Landscape of Brain Tumors. *Cancer Cell* **31**,
326–341 (2017).

48. Taylor, K. R. & Monje, M. Invasive glioma cells: The malignant pioneers that follow the current.
Cell **185**, 2846–2848 (2022).

49. Erices, J. I. *et al.* Glioblastoma Microenvironment and Invasiveness: New Insights and Therapeutic
Targets. *Int. J. Mol. Sci.* **24**, 7047 (2023).

50. Patel, S. K. *et al.* Generation of diffuse intrinsic pontine glioma mouse models by brainstem-targeted
in utero electroporation. *Neuro-Oncol.* **22**, 381–392 (2020).

51. Chatinier, A. du *et al.* Generation of immunocompetent syngeneic allograft mouse models for
pediatric diffuse midline glioma. *Neuro-Oncol. Adv.* **4**, vdac079 (2022).

52. Srivastava, R. *et al.* Development of a human glioblastoma model using humanized DRAG mice for
immunotherapy. *Antib. Ther.* **6**, 253–264 (2023).

53. Klein, S., Staring, M., Murphy, K., Viergever, M. A. & Pluim, J. P. W. elastix: A Toolbox for
Intensity-Based Medical Image Registration. *IEEE Trans. Méd. Imaging* **29**, 196–205 (2010).

54. Marstal, K., Berendsen, F., Staring, M. & Klein, S. SimpleElastix: A User-Friendly, Multi-Lingual
Library for Medical Image Registration. *2016 IEEE Conf. Comput. Vis. Pattern Recognit. Work.*
(CVPRW) 574–582 (2016) doi:10.1109/cvprw.2016.78.

55. Rios, A. C. *et al.* Intraclonal Plasticity in Mammary Tumors Revealed through Large-Scale Single-
Cell Resolution 3D Imaging. *Cancer Cell* **35**, 618-632.e6 (2019).

Figure 1. BEHAV3D Tumor Profiler pipeline. Schematic representation of the workflow showing data preparation, input and outputs in each of the three modules. The programmatic processing is represented with a yellow background **a,** Time-lapse image processing and preparation, based on segmentation and tracking of tumor and labelled TME cells. Upload to Google Drive is optional, but recommended to work in the Google Colab framework. **b,** Heterogeneity Module provides distinct behavioral clusters for DMG cells, and relates them to dynamic features and provides a back-projection of the tracked cells. **c, d,** Optional modules, additional to the Heterogeneity module. **c,** Large-scale phenotyping module combines TME large-scale information with DMG behavioral profiling to further depict population frequencies in distinct TME regions. **d,** Small-scale phenotyping module combines TME small-scale information with DMG behavioral profiling to depict how the cellular environment can influence tumor cell behavior.

Figure 2. Behavioral profiling of DMG cells with BEHAVED-TP heterogeneity module. a, Representative intravitally imaged position showing DMG cells tracked relative to tumor Edge (left panel). Tracks were classified according to DMG behavior (right, color coded by cluster). Representative of *n = 18* independent positions. **b,** UMAP plot showing seven color-coded DMG cell behavioral clusters identified by BEHAV3D-TP. Each datapoint represents one T cell track. **c,** Representative intravitally imaged positions and enlarged sections for showcasing distinct DMG behavioral clusters *Invading* (yellow), *Retreating* (green), *Erratic* (dark purple) and *Static* (light purple) through the time series. n = 18 independent experiments. Scale bars 10μm. **d,** Heatmap depicting relative values of DMG cell features indicated for each cluster, named according to their most distinct characteristics. Arbitrary units in respect to maximal and minimal values for each feature. Bubble size according to significance levels, all features represented have a significance level of P < 0.001 (***): *mean_displ2, mean squared displacement; speed; delta displacement; displacement length; persistence; invasion*. P-values represent the significance of differences across clusters tested through ANOVA for each feature. **e, f,** UMAP representation of DMG cells' velocity relative to the tumor edge(μm) (**e**) and speed (μm/h) (**f**). Each datapoint represents one T cell track and is colored following a color gradient. For each feature colors represent the mean DMG track values.

Figure 3. Mapping distinct DMG behavioral patterns to distinct TME regions. a, Overview of the large-scale TME region phenotyping workflow. After IVM imaging session, the tissue is fixed and stained and then analyzed using mLSR-3D. CytoMAP Spatial Analysis Toolbox is used to identify distinct large-scale regions, that are subsequently matched to IVM imaging dataset. **b,** Representative fixed multispectral image of a mLSR3D imaged DMG tumor and zoomed-in images of TME large-scale regions: *Void* (blue outline), *TAMM/vascularized* (red outline) and *TAMM/Oligo* (yellow outline). DMG nuclei (blue), CD31 (pink), Olig2 (purple), and Iba1 (yellow) are represented on the imaged. Scale bars, 100 µm (overview), 30 µm (zoomed-in regions). **c,** 3D multispectral image DMG cell rendering color-coded per large TME region: *Void* (blue), *TAMM/Oligo* (yellow) and *TAMM/vascularized* (red) (left). Regions were mapped on intravital imaging data (right). **d,** DMG single-cell behavioral cluster frequencies among the TME large-scale regions identified with CytoMAP. *Slow retreating*, P < 0.05; *Static*, P < 0.05; *Invading*, P < 0.05. P-values represent the significance of differences across TME regions tested through ANOVA for each behavioral cluster.

Figure 4. The BEHAV3D-TP small-scale phenotyping correlates DMG behavioral profiles with micro-environmental composition at a single-cell resolution. a, Overview of the small-scale TME phenotyping module. During IVM imaging, information about spatial distribution of SR101+ (purple), CD20r+ (purple) and CD31+ (red) cells is collected and processed by the BEHAV3D-TP *small-scale phenotyping module*. Single DMG cell spatial TME features like distance to its neighbors, number of cells in a radius or minimal distance to a cell type were measured. **b,** Heatmap depicting DMG cell small-scale niches' features across distinct DMG behavioral clusters. Arbitrary units in respect to maximal and minimal values for each feature. Bubble size according to significance levels, from smallest (P < 0.05 (**)) to medium (P < 0.01 (**)) to biggest (P < 0.001 (***)): *dist_10_cell*, Distance to nearest 10 cells, P < 0.01 (**); *n_SR101*, number of SR101 cells, P < 0.05 (*); *min_SR101*, minimal distance to a SR101 cell, P < 0.05 (*); *n_CD20r*, number of CD20r cells, P < 0.05 (*); *min_Cd20r*, minimal distance to a CD20r cell, P < 0.01 (**); *min_BV_dist*, minimal distance to a Blood vessel, P < 0.001 (***). P-values represent the significance of differences across clusters tested through ANOVA for each feature. **c, e,** Boxplot depicting the *Minimal distance to a CD20r+ TAMM (µm)* (**c**) and the *Minimal distance to a blood vessel (µm)* (**e**) across DMG behavioral clusters. **c,** *Invading* versus *Erratic*, P < 0.01 (**); *Invading* versus *Slow*, P < 0.05 (*). **e,** *Static* versus *Erratic*, P < 0.01 (**); *Static* versus *Retreating*, P < 0.001 (***); *Static* versus *Invading*, P < 0.001 (***); *Slow* versus *Retreating*, P < 0.1 (·); *Slow invading* versus *Retreating*, P < 0.1 (·); *Slow* versus *Invading*, P < 0.1 (·); *Slow invading* versus *Invading*, P < 0.1 (·). **d,** Fixed multispectral image showing the movement of DMG cells (grey) relative to CD20r+ TAMM (blue) and the Tumor core. The trajectories show a path from the initial (blue) to the final position (red). Scale bar, 10 µm. **f,** Time lapse multispectral images (left to right) showing the DMG cells (blue) displacement along CD31 blood vessel (purple) away from the tumor edge. Scale bars, 30 µm..