Immunogenicity and protection efficacy of self-amplifying and circular mRNA vaccines for SARS-CoV-2

Oinam Ningthemmani Singh[1], Umang Berry[1], Garima Joshi[2], Tejeswara Rao Asuru[2], Kannan Chandrasekar[1], Sriram Narayanan[3], Puneet Srivastava[1], Mahima Tiwari[1], Souvick Chattopadhyay[4], Farha Mehdi[4], Bhisma Narayan Panda[5], Debasis Nayak[6], Shailendra Mani[1], Tripti Shrivastava[1], Gaurav Batra[4], C.T. Ranjith-Kumar[3], Prasenjit Guchhait[2], Milan Surjit[1*]

[1]Centre for Virus research, therapeutics and vaccines, Translational Health Science and Technology Institute, National Capital Region Biotech Science Cluster, Faridabad, Haryana, India.

[2]Regional Centre for Biotechnology, National Capital Region Biotech Science Cluster, Faridabad, Haryana, India.

[3]University School of Biotechnology, Guru Gobind Singh Indraprastha University, New Delhi, India.

[4]Centre for Biodesign, Translational Health Science and Technology Institute, National Capital Region Biotech Science Cluster, Faridabad, Haryana, India.

[5]Experimental animal facility, Translational Health Science and Technology Institute, National Capital Region Biotech Science Cluster, Faridabad, Haryana, India.

[6]Department of Biological Science, Indian Institute of Science Education and Research (IISER) Bhopal IISER, Bhopal, India.

Correspondence: milan@thsti.res.in

Running title: SARS-CoV-2 SAM & circ-RNA vaccine comparison

Abstract

Recent studies have demonstrated safety and reliability of messenger RNA (mRNA) vaccines for human use. Conventionally, mRNA vaccines use linear or self-amplifying mRNA (SAM), the latter considered to be superior. However, COVID-19 SAM vaccines showed limited success. Further, Circular mRNA (Circ-RNA) vaccines against the SARS-CoV-2, Ebola and monkey pox proved its efficacy. Circ-RNAs are highly stable, do not induce innate immune pathway RNA sensors nor require any extracellular protein for their function. Here, we compared the efficacy of SAM- and Circ-RNA vaccines using the SARS-CoV-2-RBD (receptor binding domain) antigen. Both SAM-RBD and Circ-RBD induced a comparable anti-RBD IgG titer and virus-neutralization titer. However, the latter induced a significantly higher memory T-cell response. Circ-RBD formulation was stable in refrigerator, induced a durable neutralizing antibody response, protected mice from COVID-19-like pathology and a bivalent vaccine containing the Circ-RNA-RBD of SARS-CoV-2-Delta and Omicron variants potently neutralized both viruses. These finding demonstrate the overall superiority of the Circ-RNA over the SAM vaccine technology and provides the proof of concept for developing bivalent Circ-RNA vaccines against emerging variants of SARS-CoV-2.

Introduction

Although messenger RNA (mRNA) vaccines have been researched for more than two decades, its proof of success in real-life was witnessed for the first time during the COVID-19 pandemic. Comirnaty, the first FDA (United States food and drug administration)-approved COVID-19 vaccine (developed by Pfizer-BioNTech) and Spikevax (COVID-19 vaccine developed by Moderna), both mRNA vaccines, were pivotal in controlling the COVID-19. Since then, there is considerable momentum to develop mRNA vaccines against multiple pathogens, allergic diseases and cancer[1-6]. Advances in mRNA delivery technology, *in vitro* transcription technology and reduction in the cost of production have improved the overall yield, stability, efficacy and economic viability of the mRNA vaccine, thereby positioning it as the preferred option for vaccine development[7]. Both linear mRNA (such as Comirnaty and Spikevax) and self-amplifying mRNA (SAM) (such as Gemcovac)-based vaccines have been approved for use in human[1,2,8]. More recently, several independent studies have demonstrated the efficacy of circular mRNA (Circ-RNA) vaccines[9-12].

Linear mRNA-encoding the vaccine antigen is generally small in size (compared to the SAM), contain a methyl-guanosine cap analog at the 5'-end, poly A-tailed at the 3'-end and the antigen-encoding sequence is flanked by untranslated regions (UTR) at both sides. A major challenge of linear mRNA-based vaccines is low stability of the input mRNA. Stability of the linear mRNAs are increased by incorporating pseudo-uridine or N6-Methyladenosine (m6A) into the sequence[13,14]. SAM vaccines are generally based on engineered genome of alphaviruses such as the Venezuela Equine Encephalitis virus (VEEV), Semliki Forest Virus (SFV) or Sindbis virus (SINV)[15]. Here, the viral structural protein-encoding sequence is replaced with the sequence of the vaccine antigen so that non-structural proteins of the virus enables replication the viral genome without production of infectious progeny virions. Thus, SAM vaccines should produce the required quantity of antigen for a prolonged period with less amount of the input RNA (compared to the linear mRNA), leading to a stronger and more durable immune response[16]. However, use of viral replication machinery for amplifying the input RNA has certain disadvantages such as: size of the SAM RNA is considerably longer as it contains the sequence to express viral non-structural proteins; viral RNA-dependent RNA polymerase, which is required for replication of the SAM RNA has low fidelity, increasing the possibility of introducing mutation(s) in the antigen-encoding sequence; pre-existing anti-vector immunity remains a concern; generation of dsRNA during the replication of SAM might activate host defence machinery leading to degradation of SAM; possibility of uncontrolled and excessive replication of SAM due to mutations in the non-structural proteins cannot be ruled out. Notably, one of the SAM vaccines (COVAC1) failed to induce 100% seroconversion while another SAM vaccine (ARCT154) showed a very good response in phase I clinical

trials[17,18]. Gemcovac (Gennova), a commercially available SAM vaccine, also shows very good

protection efficacy in clinical trials[8].

Circular RNAs (Circ-RNA) are covalently closed single-stranded RNA without free 5'- and 3'-

ends. It is produced naturally through back-splicing, which involves the connection of the 3'-

end of an exon to the 5'- end of another exon, creating a close loop RNA. The circular structure

provides resistance to most of the RNases except RNase A, RNase T1 and RNase T2, thus

significantly increasing their half-life compared to linear mRNAs. Endogenous Circ-RNAs act

as miRNA sponges to regulate various cellular processes by binding to nucleic acids and

proteins, thereby modulating transcription and cellular signalling pathways[19]. Circ-RNAs do

not induce the cellular innate immune pathway RNA sensors[20,21]. Recent reports have

demonstrated the ability of engineered Circ-RNAs to express the vaccine antigens using

internal ribosomal entry site (IRES)-mediated translation, thereby eliminating the requirement

of capping of the 5'-end of the mRNA[22,9]. Circ-RNA-based vaccines against the Severe acute

respiratory syndrome-coronavirus-2 (SARS-CoV-2) and Monkey pox has been shown to

induce potent immune response and protect the animals from challenge with the respective

viruses[9,12]. Notably, in the case of SARS-CoV-2, Circ-RNA immunization produced a higher

T_H1 biased humoral immune response and a higher neutralizing to binding antibody ratio than

its mRNA counterpart, supporting its protection efficacy and superior potential for evading the

antibody-dependent enhancement (ADE)[9].

SARS-CoV-2, the causative agent of the coronavirus-induced disease-19 (COVID-19), is

responsible for the worst global pandemic of the 21st century. So far approximately 771 million

COVID-19 cases and 7 million deaths have been reported globally

(https://data.who.int/dashboards/covid19). The viral genome shows high mutation rate,

leading to emergence of multiple variants, some of which, such as the B.1.617.2 (Delta) and

the Omicron B.1.1.529 (BA.1) variants pose serious health issues[23,24]. Omicron B.1.1.529

(BA.1) became the most dominant circulating strain with 22 subvariants accounting for 87%

of infections. Omicron strain dominated the previous variant of concern (VOC) in 2022 and

carries the most significant number of mutations than the previous VOC, with 37 mutations

within its spike protein. 15 mutations at the RBD region covering almost all the critical mutation

sites, such as K417N, S477N, T478K, E484A and N501Y make more transmissible and

immune evasion to pre-existing immunity. Previous non-Omicron infection or vaccination has

little protection against the Omicron variant[25-27]. The Omicron sub-variants, including the

BA.4/5 and XBB, continue to emerge, with total immune escaping potential to the previous

vaccination. Few studies have shown the broader protective efficacy of bivalent mRNA and

nanoparticle vaccines composed of a mixture of antigens from the SARS-CoV-2-Beta and

Delta or Delta and Omicron variants[28-30]. Despite development of several vaccines against the

SARS-CoV-2, there is limited access to vaccines globally and there is scope for improving the efficacy and duration of protection offered by the vaccines.

Being the viral outer surface protein, spike protein of the SARS-CoV-2 is the main target for vaccine development. The receptor binding domain (RBD) of the spike protein, which accounts for one-fifth of the total protein, produced over 90% of the neutralizing antibody in infected patients[31]. There are no reports of antibody dependent enhancement (ADE) by vaccines based on the full length spike protein of the SARS-CoV-2. However, possibility of ADE induction by the spike protein of emerging SARS-CoV-2 variants cannot be ruled out. Therefore, RBD would be the best target for developing SARS-CoV-2 vaccines.

In the present study, we compared the immunogenicity and virus neutralization potential of the Circ-RNA and SAM-RNA vaccine candidates using the SARS-CoV-2-RBD as the vaccine antigen. Although both Circ-RBD (Circ-RNA backbone) and SAM-RBD (SAM-RNA backbone) elicited comparable humoral and neutralizing antibody response, the former induced a significantly higher T_H1-biased cellular immune response and protected mice from challenge with the SARS-CoV-2. Further, Circ-RBD vaccine was stable in refrigerator, induced a durable neutralizing antibody response and a bivalent vaccine formulation encoding the RBD of Delta (Circ-RBD[Del]) and Omicron (Circ-RBD[Omi]) variants induced high cross-neutralizing antibody response against both variants, confirming the protective efficacy of the Circ-RNA vaccines. Significance of these findings in designing better mRNA vaccines based on circular RNA technology is discussed.

Results

Production and characterization of the SARS-CoV-2 SAM-RBD and Circ-RBD vaccine formulations

Self-splicing property of the Anabaena intron (Group 1) was utilized to generate the Circ-RNA encoding the SARS-CoV-2-RBD (Wuhan isolate) fused to the CD5 signal sequence (SS) and the sequence encoding the trimerization motif of bacteriophage T4 fibritin (foldon) at the N- and C- terminus, respectively. The fusion protein was translated under control of a Coxsackie virus B3 (CVB3) internal ribosome entry site, as reported earlier[22]. A bacteriophage T7 promoter sequence was placed upstream of the anabaena intron sequence at the 5'-end to enable *in-vitro* transcription-mediated synthesis of Circ-RBD. A Semliki Forest virus (SFV) replicon was engineered to produce the self-amplifying RNA encoding the SARS-CoV-2-RBD (SAM-RBD). RBD-encoding sequence fused to the CD5 signal sequence and T4 fibritin foldon sequence at its 5'- and 3'-ends was inserted into the SFV replicon. *In-vitro* transcription was carried out

using the SP6 polymerase, followed by Cap1-capping of the RNA. Sized integrity of the RNAs

were verified by spectrophotometry and formaldehyde-agarose gel electrophoresis. Expected

sized of the RNAs were observed (Figure 1A). An aliquot of the RNAs were treated with the

RNase R to check their circularization status. SAM-RBD was completely digested by RNase

R treatment whereas majority of Circ-RBD was resistant, confirming its circularization

(Figure1A). Next, RNA was encapsulated into the lipid-nanoparticle (LNP) by microfluidic

mixing and the diameter of the resulting RNA-LNP formulation was measured. The average

diameter of the SAM-RBD-LNP and the Circ-RBD-LNP was found to be 140 and 136nm,

respectively (Figure 1B). Average polydispersity index (PDI) of the SAM-RBD-LNP and the

Circ-RBD-LNP was 0.06 and 0.04, respectively (Figure 1C). LNP encapsulation and stability

of the SAM-RBD and the Circ-RBD RNA in the LNP complex was verified by resolving the

RNA-LNP complex with or without treatment with Triton X-100 (disrupts the LNP). No RNA

was visible in the SAM-RBD-LNP and the Circ-RBD-LNP complex-containing samples (Figure

1D). Treatment of the sample with Triton X-100 results in detection of the SAM-RBD and Circ-

RBD (Figure 1D). No difference was observed in the quality of the SAM-RBD and Circ-RBD

RNA before and after LNP encapsulation (Figure 1D). Percentage LNP encapsulation of the

SAM-RBD and Circ-RBD RNA was quantified using Ribogreen. 98.3% and 95.9% of SAM-

RBD and Circ-RBD RNA, respectively, was LNP-encapsulated (Figure 1E). Scanning electron

microscopy analysis of the SAM-RBD-LNP and Circ-RBD-LNP complexes revealed spherical

structure with average diameter of 136nm and 149nm respectively (Figure 1F). Next, SAM-

RBD-LNP and Circ-RBD-LNP complexes were transfected into HEK293T cells to detect the

expression of RBD. Western Blot analysis of the whole cell extract using the anti-RBD antibody

confirmed expression of the RBD protein (Figure 1G, upper panel). An aliquot of the whole

cell extract was probed using the anti-GAPDH antibody to check equal loading of the sample

(Figure 1G, lower panel). Secretion of RBD protein into the culture medium was checked by

anti-RBD western blot of the culture medium collected from the cells used in the above

experiment (shown in Figure 1G). RBD was detected in the culture medium as 2 distinct

bands, likely representing the glycosylated variants of the same protein (Figure 1H). Aliquots

of the same sample were resolved in non-reduced state to check trimerization status of the

RBD protein. As expected, anti-RBD western blot detected a band of approximately 100kDa

in the non-reduced samples, indicative of RBD-trimer (Figure 1I). Possible activation of the

innate immune pathways by the SAM-RBD and Circ-RBD RNAs were checked by RT-qPCR

analysis of IFN-β, RIG-I, OASL, PKR and OAS1 genes. None of these genes were induced

by either of the RNAs (Figure 1J). As a positive control, cells were transfected with poly I:C in

parallel, which is known to induce the cellular antiviral response pathway (Figure 1J).

Collectively, these results confirmed LNP encapsulation of SAM-RBD and Circ-RBD RNA and

expression of RBD protein trimer by both the RNAs. Importantly, western blot analysis also

demonstrated that approximately same amount of RBD was produced by both the SAM-RBD
and the Circ-RBD. Therefore, our experimental condition was suitable to compare the immune
response induced by the SAM-RBD and the Circ-RBD.
218

SAM-RBD and Circ-RBD induced significant T_H1-biased humoral and cellular immune
response in mice
To select the injection route and the optimal RNA dose, Balb/c mice were immunized via
intradermal (ID) or intramuscular (IM) route with freshly prepared vaccine formulation
containing 25μg Circ-RBD RNA. Two weeks later, immunization was repeated and blood was
collected after four weeks (day 28). Both ID and IM routes showed similar anti-RBD IgG titer
(average values: $6X10^5$ and $1X10^6$, respectively) and virus neutralization titer (average NT_{50}=
$1.7X10^3$ and $1.9X10^3$, respectively) (Figures S1A and B). ID immunization route was selected
for all subsequent experiments. Next, mice were injected with 5, 10 and 25μg of Circ-RBD-
RNA containing vaccine formulations in a two-dose immunization schedule as described
above. Measurement of anti-RBD IgG titer revealed average values to be $2.2X10^4$, $2.8X10^5$,
$6X10^5$ and average virus neutralization titer (NT_{50}) was found to be $0.5X10^3$, $1.5X10^3$, $0.8 X10^3$,
in 5, 10 and 20μg RNA containing formulations, respectively (Figures S1C and S1D). There
was no significant difference in values obtained between all 3 doses, hence 10μg RNA
containing formulation was selected for all subsequent experiments.
In order to evaluate the immune response generated by the Circ-RBD and SAM-RBD, 10μg
of each RNA containing vaccine formulation was intradermally injected into Balb/c mice in two
dose immunization schedules at two-week interval, followed by collection of blood and spleen
samples for analysis of different immune parameters (Figure 2A). Significant level of anti-RBD
IgM was detected in the sera of mice immunized with both SAM-RBD and Circ-RBD RNA
(Figure 2B). Importantly, no significant difference was observed in the level of IgM between
SAM-RBD and Circ-RBD immunized mice (Figure 2B). Measurement of anti-RBD IgG titer
and virus neutralization titer (NT_{50}) in the sera collected on day 28 from SAM-RBD and Circ-
RBD immunized mice revealed the average values to be $2.8X10^5$ and $3.2X10^6$ (anti-RBD-IgG
titer) and $1X10^3$ and $1.5X10^3$ (NT_{50}), respectively (Figures 2C and 2D). IgG-isotyping of
aliquots of the same sera showed comparable values between SAM-RBD and Circ-RBD
groups (Figures 2E-2G). The ratio of anti-RBD (IgG2a+IgG2b) to IgG1 was found to be 1.9
and 1.8, respectively, indicating a T_H1 response (Figure 2H). Next, splenocytes from these
mice were cultured *in vitro* and induced with overlapping peptide pool corresponding to the
RBD region, followed by selection of CD4$^+$ and CD8$^+$ T cells and quantification of intracellular
T_H1 (IFN-γ, TNF-α) and T_H2 (IL-4) cytokines. Both CD4$^+$ and CD8$^+$ T cells of mice immunized
with Circ-RBD induced significant IFN-γ and TNF-α response (Figures 2I-2L). A significant

IFN-γ and TNF-α response was observed in CD4$^+$ and CD8$^+$ T cells, respectively, in SAM-

RBD immunized mice (Figures 2I and 2L). An increased trend was observed in IFN-γ and

TNF-α level in CD8$^+$ and CD4$^+$ T cells, respectively, in SAM-RBD immunized mice although it

was not statistically significant (Figures 2J and 2K). There were no significant changes in IL-4

response in both SAM-RBD and Circ-RBD immunized mice, indicating that both vaccine

formulations induced T_H1-biased T-cell immune response (Figures 2M and 2N).

Representative FACS plots are shown (Figure S2). Collectively, these results demonstrate

that both SAM-RBD and Circ-RBD induced a robust and comparable humoral immune

response, however, the latter induced a better T_H1-biased cell-mediated immune response.

Circ-RBD vaccine formulation is stable at four degrees centigrade and induces a

durable immune response in both female and male mice

We evaluated the stability of the Circ-RBD vaccine formulation upon prolonged storage (up to

4 weeks) in the refrigerator (4^{0}C). The Circ-RBD vaccine formulations were stored in aliquots

in the refrigerator for 2 or 4 weeks, followed by evaluation of their physical and functional

characteristics. No change in particle diameter, polydispersity index and percent RNA

encapsulation was observed between the samples stored for 2 and 4 weeks, compared to the

freshly prepared sample (0 week) (Figures 3A-3C). Mice were injected with aliquots of the

same samples in a two-dose immunization schedule (as described above), followed by

collection of blood on the 28th day. There was no difference in the anti-RBD IgG titer and the

virus neutralization titer (NT$_{50}$) between freshly prepared and two weeks-stored samples (IgG

titer=2.8X10^5 and 4X10^5, NT$_{50}$=1.7X10^3 and 1.3X10^3 respectively) (Figures 3D and 3E). There

was approximately one log drop in the anti-RBD IgG titer and a 1.5 log drop in the virus

neutralization titer in the samples stored for 4 weeks at 4^{0}C (IgG titer=0.6X10^5,

NT$_{50}$=0.04X10^3) (Figures 3D and 3E).

Next, durability of the immune response and influence of gender on the immune response

elicited by the Circ-RBD was evaluated by immunizing male and female mice with the Circ-

RBD vaccine formulation in a two-dose immunization schedule (day 1 and day 14), followed

by measurement of the anti-RBD IgG titer and the virus neutralization titer (NT$_{50}$) up to 9

months. Both the anti-RBD IgG titer and the virus neutralization titer (NT$_{50}$) were measured in

the mice sera at 1, 3 and 9 months post-first immunization. There was no drop in the anti-RBD

IgG titer or the virus neutralization titer (NT$_{50}$) after 9 months of immunization, with reference

to the samples collected after one month of the first immunization (Figures 3G and 3H). Both

male and female mice induced a similar level of the anti-RBD IgG titer and the virus

neutralization titer (NT$_{50}$) (Figures 3G and 3H). Collectively, these results demonstrated that

Circ-RBD vaccine formulation is stable in standard refrigeration condition and induced a highly

durable neutralizing antibody response in both male and female mice.

Immunization with the Circ-RBD vaccine protects mice from the SARS-CoV-2 infection and prevents COVID-19 associated pathological changes

A mouse model of intranasal SAR-CoV-2 infection was established as described[32]. The experimental plan is illustrated (Figure 4A). Mice were injected with freshly prepared Circ-RBD vaccine formulation on day 0 and day 14, intranasally transduced with replication deficient adenovirus expressing the human angiotensin converting enzyme 2 (Ad5CMV-hACE2, denoted as hACE2 hereafter) on day 28, followed by infection with 10^5 PFU of SARS-CoV-2 (Wuhan isolate) on day 33. Body weight of the animals were measured every day, blood was collected prior to infection and 4 days after infection (day 37), animals were sacrificed on day 37 and bronchoalveolar lavage (BAL) and lung tissue were collected for analysis. As expected, SARS-CoV-2 infection led to loss of body weight in the placebo group (Figure 4B). There was equal loss of weight in the Circ-RBD immunized group on the day one post-infection (Figure 4B). However, no further weight loss was observed in the immunized group on second and third days and animals starting gaining weight by fourth day post-infection, suggesting that vaccinated mice were more resistant to SARS-CoV-2 infection induced weight loss (Figure 4B). Anti-RBD IgG titer was much higher in the Circ-RBD immunized SARS-CoV-2 infected mice compared to the placebo group infected with SARS-CoV-2 on day 37 (10^6 vs 10^3) (Figure 4C). No virus or viral RNA was detected in the lungs and BAL of Circ-RBD immunized mice whereas significant quantity of virus and viral RNA was detected in the placebo group on day 37 (the 5^{th} day of infection) (Figures 4D, 4E, 4F). Further, morphology and histopathology analysis of lung tissue of placebo and Circ-RBD immunized showed a striking difference between the two groups. Multiple lesions were visible in the placebo group lungs whereas Circ-RBD lungs appeared normal (Figure 4G). Histopathology analysis of the hematoxylin and eosin-stained lung tissue sections showed pneumonia and inflammatory cells infiltration (Figure 4H). Quantification of lung infiltration score and percent hematoxylin-stained area in the slides from placebo and Circ-RBD immunized mice revealed high lung infiltration score and higher hematoxylin-stained area in the placebo group (Figures 4I, 4J). Further, expression of the proinflammatory cytokines and IFN-γ were significantly higher in the lungs of the placebo group compared to the Circ-RBD immunized group (Figure 4K). Taken together, these results confirmed that immunization with Circ-RBD protected the mice against the SARS-CoV-2.

In order to further evaluate the protection efficacy of the Circ-RBD vaccine, we wanted to establish another mouse model of the SARS-CoV-2 infection that produces a more severe disease phenotype. Intra-tracheal administration of the hACE2 has been shown to express the hACE2 in mouse lungs within 5 days of transduction[33]. We hypothesized that increased hACE2 expression in the lungs will permit more pronounced SARS-CoV-2 infection in the lungs, leading to a more severe disease phenotype. To test the hypothesis, mice were

transduced with the hACE2 via intra-tracheal administration, followed by intranasal infection with the SARS-CoV-2. This model displayed a more severe pathogenic effect, including lethality. The experimental plan is summarized (Figure 5A). Next, the animals were immunized twice at 2 weeks intervals followed by intratracheal transduction of the hACE2 (on day 28). 10^5 PFU of the SARS-CoV-2 (Wuhan) was intranasally administered after 5 days and disease parameters were followed up for next three weeks (day 54). In the placebo group, there was a marked reduction in the body weight from the 2^{nd} day of infection up to the 10^{th} day, after which there was a gradual increase in the body weight up to the 21^{st} day (Figure 5B). In the case of Circ-RBD immunized group, weight loss was observed between the 2^{nd} to 5^{th} day post-infection after which the animals gained weight, indicating that Circ-RBD vaccine protected the animals from progressing to severe disease pathology (Figure 5B). The weight profile of each mouse is represented (Figure S3A). Placebo group animals appeared sick, displayed hunchback posture, barely moved and importantly, three animals died during the observation period (one animal each died on day 36, 38 and 40) (Figure 5C). The remaining four animals started improving from day 47 (two weeks post-infection) onwards. Notably, Circ-RBD immunized animals were almost normal throughout the observation period.

Anti-RBD IgG titer was measured in the placebo and Circ-RBD immunized group on day 40 and 47 (one- and two-weeks post-infection, respectively). There was no significant anti-RBD IgG in the placebo group after one week of infection and an average titer of 10^5 was observed after two weeks of infection (Figure 5D). Circ-RBD immunized group showed anti-RBD IgG titer of 10^6 in the samples collected after one- and two-weeks of infection (Figure 5D). Estimation of viral load and viral RNA levels in the lung revealed higher viral load (10^3 PFU/mg of lung) and higher viral RNA levels (approximately ten-fold higher) in the lungs of placebo group (samples collected on day 40) compared to the Circ-RBD immunized group (Figures 5E, 5F). Both viral load and viral RNA levels were significantly less in the placebo group samples collected on day 47, which inversely correlated with the anti-RBD IgG titer observed at the same time point (Figures 5E, 5F). Placebo group also showed higher expression of proinflammatory cytokines and IFN-γ in the lung compared to that of the Circ-RBD immunized group (Figure 5G). Morphology and histopathology analysis of lung tissue of placebo group showed multiple lesions, pneumonia and infiltration of inflammatory cells whereas Circ-RBD immunized lungs appeared closed to normal in the samples collected on day 54 (Figures 5H, 5I). A similar profile was observed in the lung samples collected on day 47 (Figures S3B, S3C). Lung morphology and histopathology of both placebo and Circ-RBD immunized mice, collected on day 40 showed few lesions and inflammatory cells (Figures S3D, S3E). Quantification of lung inflammatory score and percentage of hematoxylin-stained area in the slides further confirmed the above observation (Figures 5J, 5K). Collectively, SARS-CoV-2

360   challenged studies in both the mice models confirmed the functional ability of the Circ-RBD

361   vaccine in protecting the animals against COVID-19.

363   A bivalent SARS-CoV-2 Circ-RBD vaccine induces robust immune response and

364   efficiently neutralizes both the Delta and the Omicron BA.1 variant viruses

Recent reports on bivalent mRNA vaccines and nanoparticle vaccine demonstrate their

efficacy in conferring broad spectrum immunity against the SARS-CoV-2 variants of concern[29-

31]. Feasibility of using the Circ-RNA vaccine technology for developing a bivalent vaccine was

tested by evaluating the immunogenicity and virus neutralization potential of a vaccine

formulation containing 1:1 mixture of the Circ-RNAs encoding the RBD of the Delta (Circ-

RBD^{Del}) and the Omicron BA.1 (Circ-RBD^{Omi}) variants of the SARS-CoV-2. Sequence

encoding the RBD region of the Delta and the Omicron variants of the SARS-CoV-2 were

inserted into the Circ-RNA backbone vector in place of the RBD (Wuhan) sequence so that

both RBD^{Del} and RBD^{Omi} are fused in-frame with the CD5 secretory signal and foldon

sequence at their N- and C- terminus, respectively. Expression and secretion of the RBD^{Del}

and RBD^{Omi} were verified by western blot analysis of the whole cell lysate and the culture

medium of the HEK239T cells transfected with the Circ-RBD^{Del}-LNP and Circ-RBD^{Omi}-LNP

formulations. Anti-RBD western blot reveled that both Circ-RBD^{Del} and Circ-RBD^{Omi} were

expressed in the HEK239T cells and secreted to the culture medium (Figures 6A, upper panel,

6B, upper and middle panel). Secretion of the Circ-RBD^{Omi} was not as efficient as that of the

Delta and the Wuhan variants as the ratio of protein in intracellular to culture medium was

much higher in the RBD^{Omi} (Figure 6B, middle panel). GAPDH level was measured in aliquots

of the same samples to monitor equal loading (Figures 6A, 6B, lower panel). Mice were

injected with vaccine formulations containing 10μg of Circ-RBD^{Wuhan}, Circ-RBD^{Del}, Circ-RBD^{Omi}

or 1:1 mixture of 5μg Circ-RBD^{Del} and 5ug Circ-RBD^{Omi} in a two-dose immunization schedule

at two weeks intervals, as described earlier, followed by collection of blood on day 28 and

estimation of the anti-RBD IgG titer and the virus neutralization titer. Mice immunized with the

vaccines containing the Circ-RBD^{Wuhan}, Circ-RBD^{Del} and Circ-RBD$^{Del+Omi}$ produced an anti-

RBD IgG titer of approximately 10^6 against the SARS-CoV-2 Wuhan and Delta variants

whereas it was 0.3x10^5, 0.6x10^5 and 0.6x10^5, respectively in case of the Omicron variant

(Figures 6C, 6D, 6E). In contrast, the Circ-RBD^{Omi} vaccine induced an average anti-RBD IgG

titer of 1.2x10^5 against the SARS-CoV-2 Wuhan and Delta variants and 1.8x10^5 against the

Omicron variant (Figures 6C, 6D, 6E). The virus neutralization titer (NT$_{50}$) of the sera showed

a pattern broadly similar to that of the anti-RBD IgG titer. As observed earlier, NT$_{50}$ of sera

from the Circ-RBD^{Wuhan} immunized group against the Wuhan variant was 3.5x10^3 (Figure 6F).

Sera from the Circ-RBD^{Del} and Circ-RBD$^{Del+Omi}$ showed approximately similar NT$_{50}$ as that of

the Circ-RBD^{Wuhan} (Figure 6F). However, NT$_{50}$ of sera from the Circ-RBD^{Omi} immunized mice

against the Wuhan variant was found to be 0.2x10^3, which is in line with the data on its anti-

RBD IgG titer against the same variant (Figure 6F). NT$_{50}$ of sera from the Circ-RBD^{Del} and the

Circ-RBD$^{Del+Omi}$ immunized mice against the SARS-CoV-2 Delta variant was found to be

8.1x10^3 and 8.8x10^3, respectively, indicating that the virus neutralization efficacy of the

bivalent vaccine was similar to its monovalent counterparts (Figure 6G). Sera from the Circ-

RBD^{Wuhan} immunized mice was almost as efficient as sera from the Circ-RBD^{Del} immunized

mice in neutralizing the SARS-CoV-2 Delta variant (NT$_{50}$= 1.7x10^3) (Figure 6G). However,

sera from the Circ-RBD^{Omi} immunized mice were significantly less efficient in neutralizing the

SARS-CoV-2 Delta variant than that of the Circ-RBD^{Wuhan}, Circ-RBD^{Del} and Circ-RBD$^{Del+Omi}$

immunized mice (Figure 6G). Sera from the Circ-RBD^{Omi} and Circ-RBD$^{Del+Omi}$ immunized mice

showed an NT$_{50}$ of 8.1x10^3 and 2.3x10^3, respectively, against the SARS-CoV-2 Omicron BA.1

variant, indicating that bivalent vaccine was equally effective as its monovalent (Circ-RBD^{Omi})

counterpart (Figure 6H). NT$_{50}$ of sera from the Circ-RBD^{Del} and the Circ-RBD^{Wuhan} immunized

mice against the Omicron BA.1 variant were found to be 0.3x10^3 and 0.06x10^3 respectively,

suggesting that they were less efficient in neutralizing the Omicron BA.1 virus (Figure 6H).

Collectively, these data demonstrate that the bivalent Circ-RBD vaccine produces potent

neutralizing antibodies against the Delta and the Omicron BA.1 variants of the SARS-CoV-2.

Discussion

With the success of the COVID-19 mRNA vaccines, immunogenicity and protective efficacy

of a number of vaccine candidates based on linear mRNAs, self-amplifying mRNAs and

circular mRNAs have been evaluated in multiple independent studies. Given their ability to

replicate inside the host, SAM RNAs express the antigen for a longer duration with lesser

amount of the input RNA. Analogous to the SAM-RNA, Circ-RNAs also offer the advantage of

expression of the antigen for a longer duration with lesser amount of the input RNA, owing to

their higher stability. On top of that, Circ-RNA vaccines do not produce any additional

exogenous protein in the target cells, lack open 5'- and 3'-ends and do not produce double

strand RNA (dsRNA) in the target cell, all of which are potential unwanted features in the SAM

vaccine. Therefore, Circ-RNA vaccines appear to be more promising than the linear mRNA

and SAM vaccines. Comparison between linear mRNA and Circ-RNA vaccines have

demonstrated the superiority of the later[9]. However, there are no report on comparison of the

protective efficacy between the Circ-RNA and the SAM vaccines. Here, we compared the

efficacy of the Circ-RNA and the SAM vaccines by measuring their immunogenicity and virus

neutralization potential using the SARS-CoV-2 RBD as the vaccine antigen. After confirming

that Circ-RBD vaccine induced a better overall response, we further evaluated its protective

efficacy, longevity of the immune response and compatibility for production of bivalent vaccines using mouse model of the SARS-CoV-2 infection.

SAM-RNA was produced using the SFV replication machinery. SFV replicon was opted as there are very few reports of SFV infection in human, with very mild symptoms and no instances of mortality. SFV replicon has been used in earlier studies for heterologous protein expression and a recently it has been used to produce a vaccine candidate against the Ebola virus[34,35]. Also note that, multiple outbreaks of VEEV has been reported in North and South American countries with cases of human death. Hence chances of anti-vector immunity is higher in the case of VEEV[36].

The Anabaena permuted intron-exon sequence was used to produce the Circ-RNA and CVB3 IRES-mediated translation was used to produce the secretory signal and foldon fused-RBD protein in the target cells. Circ-RNA vaccine production using the above vector has been reported and well characterized[9]. Only the RBD region was used instead of the entire Spike protein of the SARS-CoV-2 as it contains the majority of neutralizing epitopes, avoids induction of antibody-dependent enhancement (ADE) and permits insertion of multiple elements due to its smaller size. The trimeric structure of RBD was maintained using the foldon-motif to improve its immunogenicity and virus neutralization potential, as reported earlier[37].

In order to compare the SAM-RBD vaccine with the Circ-RBD vaccine, it was important that both vaccine antigens were expressed at comparable levels and did not induce the host innate immune response. Western blot analysis using anti-RBD antibody and RT-qPCR analysis of the ISGs confirmed that both vaccine candidates expressed similar levels of the vaccine antigen and evaded the host innate immune pathway RNA sensors, respectively.

Although both SAM-RBD and Circ-RBD induced a similar humoral response, CD4$^+$ and CD8$^+$ memory T cell response was significantly increased in the Circ-RBD immunized mice. There was a trend of increased production of IFNγ and TNFα in CD4+ and CD8+ memory T cells in SAM-RBD immunized mice, however, it was not statistically significant, indicating that Circ-RBD vaccine induced a more potent cell-mediated immune response. SFV replicon system has been used to develop a dual antigen vaccine candidate against the Ebola virus by co-expressing the viral glycoprotein and nucleoprotein where it was observed that high intradermal dose of the Ebola virus glycoprotein induced IFNγ in CD4+ T cells and high intradermal dose of nucleoprotein induced very low level of IFNγ in CD8+ T cells[35]. In the case of VEEV-replicon based SARS-CoV-2-RBD and nucleocapsid (N)- expressing dual antigen vaccine immunized outbred mice, IFNγ level was moderately increased in the CD4+ and CD8+ T cells stimulated with either the S- or N-peptide pool[38]. Other studies on evaluation of a SAM vaccine candidate in mice based on the SARS-CoV-2-RBD or the Spike protein expressed from an alpha virus (VEEV/ VEEV+Sindbis) replicon reported a significant T- cell response in

splenocytes of the immunized mice, upon stimulation with the S-peptide pool[39-42]. However,

SAM-Spike immunized pigtail macaques did not induce high level of IFNγ production in CD4+

and CD8+ T cells upon stimulation of splenocyte with S-peptide pool[40]. On the other hand, the

Circ-RBD vaccine of the SARS-CoV-2 induced similar level of T cell response as that of the

corresponding linear mRNA vaccine and a higher neutralizing antibody response than that of

the linear mRNA[9]. Given that different assays were conducted in different laboratories to

measure the T cell response, it is possible that lower IFNγ and TNFα production in CD4+ and

CD8+ T cells (splenocytes stimulated with RBD peptide pool) may be attributed to the overall

lesser sensitivity of our assay protocol. However, the relative difference in the level of IFNγ

and TNFα between SAM-RBD and Circ-RBD immunized sample would remain constant as

the samples were processed in parallel. Therefore, Circ-RBD vaccine appears to be a better

alternative to a SAM-RBD vaccine. Whether a VEEV replicon-based SAM vaccine shows a

similar profile as the SFV replicon used in our study, remains to be tested. A recent study also

showed potent adjuvant function of the circular RNA, irrespective of the antigen sequence and

robust CD8+ T-cell response induced by a circular RNA vaccine encoding the nucleoprotein

of the influenza virus, further supporting practical utility of the circular RNA vaccine

technology[10].

Functional efficacy of the Circ-RBD vaccine was characterized by evaluating its

thermostability, durability of the neutralizing antibody response, ability to protect the mice upon

SARS-CoV-2 challenge and compatibility as a bivalent vaccine formulation. Circ-RBD

immunization induced peak anti-RBD IgG titer at 4 weeks, which protected the mice from

COVID-19 like disease pathology when SARS-CoV-2 infection was initiated in the upper

respiratory tract (mild disease) or lower respiratory tract (severe disease, often leading to

lethality). Nine month follow up study of the anti-RBD IgG titer and the virus neutralization titer

(NT_{50}) did not show any decline in both. Moreover, neutralizing antibody titer was similar in

both male and female mice. With regard to stability of the Circ-RBD vaccine in normal

refrigeration conditions, there was no change in the characteristic of the Circ-RBD-LNP

formulation after 4 weeks of storage, however neutralizing antibody titer was stable for up to

2 weeks of storage. Finally, we evaluated the functional efficacy of a bivalent Circ-RBD

vaccine. Failure of monovalent vaccines in preventing COVID-19 for a longer period is

ascribed to emergence of newer variants of the SARS-CoV-2 containing escape mutations in

the RBD region. Bivalent vaccines based on linear mRNA platform or nanoparticle platform

expressing the spike protein or RBD region of the spike protein of the SARS-CoV-2-Delta and

Omicron variants have shown cross-protection against SARS-CoV-2 variants[28-30]. On similar

line, our data on the bivalent Circ-RBD vaccine expressing the RBD of Delta and Omicron

variants demonstrate cross-protection against the Wuhan, Delta and Omicron variants.

In summary, the current study compared the immunogenicity and virus neutralization efficacy

of the SAM and Circ-RNA vaccine candidates encoding the SARS-CoV-2-RBD antigen, which

revealed similar humoral immune response and virus neutralization potential of both vaccines

and a superior cell-mediated immune response induced by the later. Our data also showed

reasonable stability of the Circ-RNA vaccine at four degrees centigrade, high durability of the

neutralizing antibody response induced by it and high virus neutralization efficacy of the

bivalent Circ-RNA vaccine, all of which support the use of Circ-RNA technology in vaccine

development. An earlier study showed the superiority of the SARS-CoV-2 Circ-RNA vaccine

over the corresponding linear mRNA vaccine. Data obtained from this study further paves the

way for use of the Circ-RNA technology for vaccine development.

Materials and Methods:

Plasmids and reagents

Plasmid encoding the noninfectious Semliki Forest virus replicon (pSFV3) was a gift from

Henrik   Garoff   (Addgene   plasmid   #   92072   ;   http://n2t.net/addgene:92072   ;

RRID:Addgene_92072)[34]. 988 to 1578 nucleotides encoding the RBD region (330 to 526

amino acids) of spike protein of the SARS-CoV-2 Wuhan isolate (hCoV-19/USA-WA1/2020)

fused in-frame with the CD5 secretory sequence at the N-terminal and the Bacteriophage T4

fibritin trimerization motif sequence at the C-terminus was synthesized and inserted into the

pSFV3 vector at the BamH1 restriction site. The resulting plasmid was named pSFV3-RBD.

Transcription of SFV-RBD hybrid RNA in the pSFV3-RBD vector was under control of the SP6

promoter. A circular RNA expression cassette containing the sequence of the T7 promoter,

Anabaena Group 1 intron 3' sequence, CVB3 IRES sequence, CD5 secretory sequence, RBD

region (988 to 1578 nucleotides) of spike protein of the SARS-CoV-2 Wuhan isolate (hCoV-

19/USA-WA1/2020), bacteriophage T4 fibritin trimerization motif and anabaena group 1 intron

5' sequence was chemically synthesised in the given order and cloned into the pUC57 vector.

The resulting plasmid was named pUC57 cRNA-RBD^{Wuhan}. RBD corresponding to the SARS-

CoV-2-Delta   (hCoV-19/USA/MD-HP05285/2021)   and   Omicron   (hCov-19/USA/MD-

HP2087/2021) variants were cloned similarly and the resulting plasmids were named pUC57

cRNA-RBD^{Del} and pUC57 cRNA-RBD^{Omi}, respectively. The goat anti-Mouse IgG-HRP

(Catalog No. 1030-05) was from Southern Biotech (Alabama, USA). Megascript SP6

transcription kit (Catalog no. AM1330), MegaScript T7 transcription kit (Catalog no. AM1334)

lipofectamine 3000 (Catalog No. L3000075), anti-SARS-CoV-2 Spike S1 (Catalog no. PA5-

81795), goat anti-Rabbit IgG-HRP (Catalog No. 31460), Anti-mouse IgG (H+L) Alexa Flour

488 (Catalog no. A28175) and Ribogreen (Catalog no. R11490) were from Thermo Fisher

Scientific (Massachusetts, USA). Scriptcap Cap 1 capping system (Catalog no. sccs1710) was

from CellScript (Wisconsin, USA). RNAse R (Catalog no. ab286929) was from Abcam

(Cambridge, United Kingdom). RNeasy Kit was from Qiagen ((Catalog no. 74104, Hilden,

Germany). SM102 (Catalog no. 33474) was from Cayman Chemical company (Michigan,

USA). DSPC (Catalog no. 850365P), Cholesterol (Catalog no. 700000P), and DMG-PEG

2000 (catalog no. 880151P), was from Sigma (Missouri, USA). Amicon ultra centrifugal filter

(Catalog no. UFC703008) was from Merck Millipore (Darmstadt, Germany). anti-GAPDH

antibody (Catalog no. SC-25778) was from Santa Cruz Biotechnology (Texas, USA). TRI-

reagent (Catalog no. TR118) was from MRC (Massachusetts, USA). Firescript cDNA

synthesis kit (Catalog no. 06-15-00050) and HOT FIREPol SolisGreen qPCR Mix (Catalog

no.08-47-0000S) from Solis Biodyne (Estonia). Ad5-hACE (catalog no. VVC-McCray-7723)

was obtained from the Viral Vector Core (University of Iowa, Iowa, USA). Vetbond (catalog

no. 1469Sb) was from 3M (Minnesota, USA). Mouse monoclonal antibody isotyping reagent

(Cat no. ISO2) was from Sigma Aldrich (Missouri, USA). SARS-CoV-2 spike antibody (catalog

no. GTX632604) was from GeneTex (California, USA). SARS-CoV-2 RBD peptide pool

(Catalog no: PM-WCPV-S_RBD-1) was from JPT innovative peptide solution (Berlin,

Germany). GolgiStop (Catalog no. 554724) was from BD bioscience (New Jersey, USA). FC

(Catalog no.14-9161-73), IC fixation buffer (Catalog no. FB001), permeabilizing buffer

(Catalog no. 00-8333-56), anti-mouse CD3 (Catalog no.100204), anti-mouse CD4 (Catalog

no.100408), anti-mouse CD8 (Catalog no.100714), anti-mouse CD44 (Catalog no.103030),

anti-mouse IFNγ (Catalog no. 505810), TNFα (Catalog no. 506328), and IL-4 (Catalog no.

504126) are from Thermo Fisher Scientific (Massachusetts, USA).

In vitro transcription, capping, circularization, RNase R treatment and purification

pSFV3-RBD, pUC57 cRNA-RBD[Wuhan], pUC57 cRNA-RBD[Del] and pUC57 cRNA-RBD[Omi]

plasmids were linearized and purified DNA was used for the _in vitro_ transcription using the

Megascript SP6 or T7 transcription kit, following the manufacturer's instruction. SAM-RNA was

capped using the Scriptcap Cap 1 capping system, following the manufacturer's instructions.

In the case of Circ-RNA, _in vitro_ transcribed RNA was incubated at 55^0C for 15 min in the

circularization buffer (10mM MgCl$_2$, 50mM Tris-HCl, 100mM NaCl, 1mM DTT, pH 7.9; New

England Biolabs, USA) and 2μM rGTP to circularize the RNA, as described earlier[16].

Circularized RNA was incubated at 37^0C for 7 min with 0.1units RNase R/μg RNA, followed

by addition of 0.1 units RNase R/μg RNA and incubation for another 7 min. RNA was purified

using the RNeasy kit, following the manufacturer's protocol (Qiagen, Germany).

Preparation and characterization of the RNA-LNP vaccine formulation,

SM102, DSPC, Cholesterol and DMG-PEG 2000 were mixed at a molar ratio of 50:10:38.5:1.5

in absolute ethanol at 1/5 of the final LNP amount. An aqueous solution of RNA was prepared

by diluting it in 50mM acetate buffer (1.2mM sodium acetate, 4mM acetic acid, pH 4.0) at 4/5

of the final LNP volume. The two solutions were mixed using a microfluidics system (ElveFlow

Elvesys, France) at a flow rate ratio of 1:5. The resulting RNA-LNP complex was dialyzed with

phosphate buffered saline (137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4,

pH 6.8) and concentrated using a 30kDa MWCO Amicon ultra centrifugal filter to a volume of

0.1μg/μl RNA concentration. Further, the RNA-LNP vaccine formulation was passed through

a 0.2μm filter and aliquoted into glass vials and stored at 4^0C. Size and polydispersity (PDI) of

the RNA-LNP vaccine formulation was evaluated by dynamic light scattering (ZetaSizer lab,

Malvern panalytical, United Kingdom). For transmission electron microscopy (TEM), RNA-

LNP vaccine formulation was diluted ten times in H_2O, followed by staining with 2% uranyl

acetate and image acquisition using JEM-1400 Flash TEM equipped with tungsten filament

and sCMOS camera (JEOL Ltd, Japan). RNA encapsulation into LNP was measured by

fluorescent-based RNA quantification system using Ribogreen, following manufacturer's

instruction. The encapsulated RNA was released from the LNP by treating the RNA-LNP

vaccine formulation with 2% Triton X-100, followed by measurement of RNA quantity by

Ribogreen assay and evaluation of RNA quality by agarose gel electrophoresis.

Mammalian cell culture, transfection, western blot, RT-qPCR, SARS-CoV-2 propagation

HEK293T cells were maintained in Dulbecco's modified Eagle medium (DMEM)

supplemented with 10% FBS (fetal bovine serum) and Penicillin, Streptomycin during

maintenance. Cells were seeded on 6 well plates (8×10^5 cell/well) and transfected with 6μg

of RNA-LNP complex. Only LNP was added in the mock. For western blot analysis under

reducing condition, samples (cells or culture medium) were incubated at 95^0C for 5 minutes

in Laemlli buffer (2% SDS, 13% glycerol, 2.5% β-mercaptoethanol, 0.005% bromo phenol

blue and 63 mM Tris HCl, pH 6.8). For non-reducing condition, β-merceptoethanol was

omitted from the Laemlli buffer. Total protein in the samples was quantified by Bicinchoninic

acid assay and equal amount of protein were resolved in SDS-PAGE, followed by transfer to

Polyvinylidene fluoride (PVDF) membrane. The membrane was blocked with 5% skimmed

milk, incubated with anti-Spike S1 or anti-GAPDH antibody at 1:5000 dilution. Anti-Rabbit

secondary antibodies (Santa Cruz Biotechnology) were used at 1:5000 dilution. Protein

bands were developed by enhanced chemiluminescence reagent, using a commercially

available kit (BioRad, USA) and visualized in a gel Documentation system (Chemidoc MP,

Bio Rad, USA).

For RT-qPCR analysis of genes involved in the antiviral response pathway, HEK293T cells

were transfected with 1μg of SAM-RBD or Circ-RBD RNA or Poly I:C was using lipofectamine

3000, following manufacturer's instruction (Thermo Fisher Scientific, Massachusetts, USA).

24 hours post-transfection, total RNA was isolated using the TRI-reagent (MRC, Massachusetts, USA), followed by reverse transcription (RT) using the Firescript cDNA synthesis kit (Solis Biodyne, Estonia). The relative transcript levels of different genes were determined by SYBR green based RT-qPCR, as described earlier[43]. Values obtained for GAPDH RNA was used as an internal control for normalization of the data. Primer sequences are as follows: hGAPDH FP: 5' GAGTCAACGGATTTGGTCGT 3', hGAPDH RP: 5' TTGATTTTGGAGGGATCTCG 3', hIFNβ FP: 5' ACGCCGCATTGACCATCTAT 3', hIFNβ RP: 5' TTGGCCTTCAGGTAATGCAGA 3', hRIG-I FP: 5' TGTGGGCAATGTCATCAAAA 3', hRIG-I RP: 5' GAAGCACTTGCTACCTCTTGC 3', hOASL FP; 5' AGGGTACAGATGGGACATCG 3', hOASL RP: 5' AAGGGTTCACGATGAGGTTG 3', hPKR FP: 5' TCTTCATGTATGTGACACTGC 3', hPKR RP: 5' CACACAGTCAAGGTCCTT 3', hOAS1 FP: 5' GCTCCTACCCTGTGTGTGTGT 3', hOAS1 RP: 5' TGGTGAGAGTACTGAGGAAGA 3'.

For RT-qPCR analysis in the mouse lung tissue, total RNA was isolated from 10mg lung tissue using TRI reagent, following manufacturer's instruction. cDNA synthesis and RT-qPCR was done as mentioned above. HPRT RNA level was quantified for normalization of data. Primer sequences are as follows: SCoV2 FP: 5' TGGACCCCAAAATCAGCGAA 3', SCoV2 RP: 5' TCGTCTGGTAGCTCTTCGGT 3', mHPRT FP: 5' GTTGGATACAGGCCAGACTTTGTTG 3', mHPRT RP: 5' GATTCAACTTGCGCTCATCTTAGGC 3', mIL-6 FP: 5' GAGGATACCACTCCCAACAGACC 3', mIL-6 RP: 5' AAGTGCATCATCGTTGTTCATACA 3', mIL-1β FP: 5' GCCCATCCTCTGTGACTCAT 3', mIL-1β RP: 5' AGGCCACAGGTATTTTGTCG 3', mTNFα FP: 5' CATCTTCTCAAAATTCGAGTGACAA 3', mTNFα RP: 5' TGGGAGTAGACAAGGTACAACCC 3', mIFNγ FP: 5' AACGCTACACACTGCATCTTGG 3', mIFNγ RP: 5' GACTTCAAAGAGTCTGAGG 3'.

SARS-CoV-2 Wuhan isolate (NR-52281, SARS-related coronavirus-2 isolate ISA-WA1/2020) and Omicron Variants (B.1.1.529, NR56461, SARS-related coronavirus-2 isolate hCoV-19/USA/MD-HP20874/2021) were obtained from BEI Resources, amplified in Vero E6 cells in the BSL3 facility of THSTI, India, titrated and stored frozen in aliquots. SARS-CoV-2-Delta (SARS-CoV-2/human/IND/THSTI_287/2021) was isolated from clinical sample at THSTI and has been reported (GenBank: MZ356566.1). SARS-CoV-2 infection was done as described[43].

Mice handling, immunization, SARS-CoV-2 challenge study

Mice were housed in individually ventilated cages at the experimental animal facility of THSTI. For SARS-CoV-2 infection experiments, animals were housed in the animal biosafety level-3 (ABSL-3) laboratory of THSTI. Institutional animal ethics committee and institutional biosafety committee approval was obtained for the experiments. Male and Female Balb/c mice of 6 to 8 weeks old were intradermally injected on the back of the animal. In the case of intramuscular

injection, mice were injected in the hind leg muscle. All injections were in $100\mu l$ volume.

Placebo animal received $100\mu l$ PBS. Blood was collected at the indicated time points through

retro-orbital sinus using a sterile hematocrit capillary tube. Wherever indicated, mice were

euthanized by anesthetic overdose (Ketamine 300 mg/kg and xylazine 30 mg/kg) and blood,

spleen and lungs were collected. Serum was isolated from the blood by centrifugation at

10,000g for 10 min at $4°C$. Lung tissue was cut into parts and used for RNA, protein or

histopathology analysis.

SARS-CoV-2 challenge experiments were conducted in inbred Balb/c mice. Six to eight-week-

old mice were immunized with $10\mu g$ of Circ-RBD twice at two-weeks interval. Blood was

collected through retro-orbital sinus and anti-RBD IgG was measured. Replication deficient

adenovirus expressing human ACE2 (Ad5CMV-hACE2, denoted as hACE2) was transduced

through intranasal or intratracheal route[32,33]. For Intranasal transduction, mice were lightly

anaesthetized with isoflurane and $2x10^8$ pfu of hACE2 was administered, as described[32].

Intratracheal transduction of hACE2 was performed as described[33]. Mice were anaesthetised

with ketamine xylazine injection (Ketamine 100 mg/kg and xylazine 10 mg/kg), hair was

shaved from the neck region and a 5mm incision was made in the skin to expose the trachea.

$2x10^8$ pfu of hACE2 was administered in the trachea using a 24-gauge I.V. cannula. The

wound opening was glued with tissue adhesive (Vetbond, 3M Corp, USA). Animals did not

show any sign of illness or inflammation at the site of glue. Mice were infected with 10^5 pfu

SARS-CoV-2 intranasally after 5 days of hACE2 administration. Weight of the animals were

measured before infection and every day after the infection. In the intranasal transduction

model, mice were sacrificed 5 days post-SARS-CoV-2 infection and lung and BAL fluid were

harvested. In the case of intratracheal transduction model, weight was measured every day

after infection and survival was checked till 21 days of post-infection. When an animal showed

critical illness symptoms after SARS-CoV-2 infection, it was euthanized and samples were

collected. An animal from the immunized group was euthanized for comparison when mice

from the placebo group were euthanized.

ELISA, Flow cytometry, quantification of viral load, virus Neutralization assay,

The Enzyme-Linked Immunosorbent Assay (ELISA) was performed as described, with below

mentioned modifications[44]. The receptor-binding domains (RBDs) of the three SARS-CoV-2

variants-Wuhan, Delta, and Omicron, were expressed in-house in the Expi293 cells (Thermo

Fisher Scientific, Massachusetts, USA) and used in the ELISA. The assay samples (mice sera)

were diluted in a 3-fold series from 1:500 to 1:1,093,500. Horse radish peroxidase (HRP)-

labelled Goat anti-Mouse IgG (H+L) (Jackson ImmunoResearch, USA) was used at a dilution

of 1:5000. The endpoint anti-RBD IgG titer (denoted as anti-RBD IgG titer in the text and

figures) refers to the dilution factor at which the absorbance (obtained in the ELISA) of the

test samples exceeded three times that of the background (secondary Ab alone). Anti-RBD

IgM and IgG isotype (IgG1, IgG2a and IgG2b) ELISA was performed using a commercially

available kit, following the manufacturer protocol (Sigma Aldrich) at a serum dilution of 1:500.

For FACS study, splenocytes were prepared as described and cultured in RPMI-10

supplemented with 10%fetal bovine serum and 1% penicillin-streptomycin antibiotic[45]. After 3

hours, RBD peptide pool (2 μg/ml each peptide) and GolgiStop (BD) was added to the culture

medium for 6 hours. Cells were blocked with FC at room temperature for 15 min, stained with

anti-CD3, anti-CD4, anti-CD8 and anti-CD44 antibodies followed by fixation and

permeabilization. Intracellular staining of IFNγ, TNFα and IL-4 was performed using respective

antibodies. Cells were sorted using the BD FACS-aria (BD Biosciences) and data was

analyzed using the FlowJo software (FlowJo LLC, Oregon, USA).

For quantification of SARS-CoV-2 level in the lungs, 5mg of lung tissue was homogenised in

1ml serum free DMEM and sterilized using 0.2μm filter. The homogenate was serially diluted

10-fold and added to 2×10^5 VeroE6 cells/well in 12 well plate for 1 hour. For quantification of

viral load in the BAL, BAL fluid were collected twice in PBS (500μl each), followed by passing

through 0.2μm filter. Two-fold serially diluted solution was added to 2×10^5 VeroE6 cells/well in

12 well plate for 1 hour. Infection medium was removed, 1% carboxymethyl cellulose (CMC)

was added and plates were incubated for 48 hours at 37^0C, with 5% CO_2 supplementation.

Cells were fixed in 3.7% formaldehyde for overnight, plaques stained with 0.05% crystal violet

and number of plaques in each well was counted.

50% virus neutralization efficiency (NT_{50}) of the sera from mice immunized with the SAM-

RBD[Wuhan] and Circ-RBD[Wuhan] vaccine formulations (shown in Figs. 2D, 3E, 3F, 3H, 3J, S1B

and S1D) were measured by the Plaque reduction neutralization test (PRNT), as

described[46]. Briefly, mouse sera were heated at 55^0C for 30 min, 2-fold serially diluted

starting from 1:10 in serum-free DMEM, mixed with an equal volume of virus [virus stock was

appropriately diluted so that each well produces approximately 50 plaque forming units (pfu)]

and incubated at 37^0C for 1 hour. 2×10^5 VeroE6 cells/well in 12 well plates were incubated

with the serum-virus mix for 1 hour. Infection medium was removed and cells were overlaid

with 1% CMC in 2% FBS supplemented DMEM and incubated at 37^0C with 5%CO_2

supplementation for 48 hours. Cells were fixed in 3.7% formaldehyde for overnight and

stained with 0.2% crystal violet. The number of plaques were counted and the serum dilution

leading to 50% neutralization of the virus (NT_{50}) was calculated using the GraphPad Prism

9.0 (Massachusetts, USA). NT_{50} of the Wuhan, Delta and Omicron variants (shown in

Figures 6F-6H) were measured by the Focus reduction neutralization test (FRNT). Heat

inactivated serum sample were 2- fold serially diluted, starting from 1:20 and mixed with an

equal volume of the virus [virus stock was appropriately diluted so that each well produces

approximately 100 focus forming units (ffu)] and incubated at 37^0C for 1 hour. $2x10^4$ VeroE6

cells/well in 96 well plates were incubated with the serum-virus mix for 1 hour. Infection

medium was removed and cells were overlaid with 0.5% CMC in 2% FBS supplemented

DMEM and incubated at 37^0C with 5%CO_2 supplementation for 24 hours. Cells were fixed in

formaldehyde and permeabilized with 0.2% Triton X-100. Next, the cells were stained with

anti-Spike antibody, following manufacturer's protocol followed by staining with alexa fluor-

488-conjugated anti-mouse IgG antibody. The number of foci was counted using an

ELISPOT reader (iSPOT, AID GmBh, Strassberg, Germany) and the serum dilution leading

to 50% neutralization of the virus was calculated using GraphPad Prism 9.0.

Histopathology

Histopathology analysis of the lung tissue samples were performed by the histology service of

the experimental animal facility of THSTI, following established protocol. Briefly, lung

tissues were kept in 10% neutral buffered formalin solution immediately after collection. After

ensuring proper fixation, samples were subjected to processing and paraffin embedded blocks

were prepared. 4μm thick sections were cut using a semi-automatic microtome (Histocore

Multicut- Leica Biosystem, Germany) and mounted on glass slides. The sections were stained

with haematoxylin and eosin, following standard operating procedures of the facility. Slides

were analysed by a qualified histopathologist, following blind evaluation method. Semi-

quantitative scoring system was adopted to score the pathological conditions, 0 indicating

absence of a pathological condition while 5 indicating its highest severity. Pathological

conditions such as; bronchiolitis, vasculitis, interstitial pneumonia, alveolitis and haemorrhage

were analysed and scored. Images were visualized and captured using a Bright field upright

microscope (CX43RF- Olympus Life science Solutions, Japan) equipped with a colour camera

(DP28- Olympus Life science Solutions, Japan) and imaging software (cellSens Entry-

Olympus Life science Solutions, Japan) at 200X magnification. Haematoxylin[+] area as an

index of cellularity was quantified from four different fields of 200X resolution using the ImageJ

software, as described[47].

Statistical analysis

750   All cell line data are derived from 3 independent experiments and represented as mean±SD

751   of 3 independent experiments. For mouse study, 6 or more animals were used in each group,

752   as indicated in the figures. Average values are shown in the bar and individual values are

753   shown as dots in each bar in the graphs. Two-tailed unpaired t-test (Mann-Whitney) and One-

754   way ANOVA (Kruskal-Wallis test) followed by Dunn's multiple comparisons was used for

comparison between groups, as indicated in the figures. Logistic regression was used to determine the NT_{50} titre of the mice sera. All statistical analysis was performed using the GraphPad Prism 9. A *p*-value <0.05 with 95% confidence interval was considered statistically significant.

Acknowledgement

This study was partly supported by a BIRAC (Biotechnology Industry Research Assistance Council, Department of Biotechnology, Government of India) grant to MS (BT/COVID0081/01/20) and THSTI core grant to MS. ONS and UB are supported by senior research fellowships from the Council of scientific and industrial research and the department of Biotechnology, Government of India, respectively. We are thankful to the experimental animal facility and biosafety level three (BSL3) facility of THSTI for helping with the experiments. The following reagent was deposited by the Centre for Disease Control and Prevention (USA) and obtained through BEI Resources, NIAID, NIH (USA): SARS-Related Coronavirus 2, isolate hCoV-19/USA-WA1/2020, NR-52281 and SARS-Related Coronavirus 2, isolate hCoV-19/USA/MD-HP20874/2021 (Lineage B.1.1.529; Omicron Variant), NR-56461, contributed by Andrew S. Pekosz. We are thankful to Henrik Garoff for the pSFV3 plasmid.

Author contribution

Conceptualization: MS; Investigation: ONS, UB, GJ, TRA, KC, SN, PS, SC, FM; Data analysis: ONS, GJ, TRA, BNP, SM, GB, CTRK, PG, MS; Resources: TS, SM, DN, GB, CTRK, PG, MS; manuscript writing: ONS, MS, GB, SM, CTRK. All authors have reviewed, edited and agreed to the final version of the manuscript.

Declaration of interest

The authors have no conflict of interest to declare.

Keywords: SARS-CoV-2, mRNA vaccine, self-amplifying mRNA, Circular mRNA, COVID-19, SARS-CoV-2 vaccine.

Reference

1. Thomas, S.J., Moreira Jr, E.D., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Perez, J.L., Pérez Marc, G., Polack, F.P., Zerbini, C., Bailey, R., et al. (2021). Safety

791   and efficacy of the BNT162b2 mRNA Covid-19 vaccine through 6 months. N. Engl. J.
792   Med. 385, 1761-1773.
793

794 2. Baden, L.R., El Sahly, H.M., Essink, B., Kotloff, K., Frey, S., Novak, R., Diemert, D.,
795   Spector, S.A., Rouphael, N., Creech, C.B., et al. (2021). Efficacy and safety of the
796   mRNA-1273 SARS-CoV-2 vaccine. N. Engl. J. Med. 384, 403-416.
797

798 3. Arevalo, C.P., Bolton, M.J., Le Sage, V., Ye, N., Furey, C., Muramatsu, H., Alameh,
799   M.G., Pardi, N., Drapeau, E.M., Parkhouse, K., et al. (2022). A multivalent
800   nucleoside-modified mRNA vaccine against all known influenza virus subtypes.
801   Science 378, 899-904.
802

803 4. Krienke, C., Kolb, L., Diken, E., Streuber, M., Kirchhoff, S., Bukur, T., Akilli-Öztürk,
804   Ö., Kranz, L.M., Berger, H., Petschenka, J., et al.(2021). A noninflammatory mRNA
805   vaccine for treatment of experimental autoimmune encephalomyelitis. Science 371,
806   145-153.
807

808 5. Heine, A., Juranek, S., Brossart, P. (2021).Clinical and immunological effects of
809   mRNA vaccines in malignant diseases. Mol. Cancer 20, 52.
810

811 6. Parhiz, H., Atochina-Vasserman, E.N., Weissman, D. (2024). mRNA-based
812   therapeutics: looking beyond COVID-19 vaccines. The Lancet 403, 1192-1204.
813

814 7. Chaudhary, N., Weissman, D., Whitehead, K.A. (2021). mRNA vaccines for
815   infectious diseases: principles, delivery and clinical translation. Nat. Rev. Drug
816   Discov. 20, 817-838.
817

818 8. Saraf, A., Gurjar, R., Kaviraj, S., Kulkarni, A., Kumar, D., Kulkarni, R., Virkar, R.,
819   Krishnan, J., Yadav, A., Baranwal, E., et al. (2024). An Omicron-specific, self-
820   amplifying mRNA booster vaccine for COVID-19: a phase 2/3 randomized trial. Nat.
821   Med. 30, 1363-1372.
822

823 9. Qu, L., Yi, Z., Shen, Y., Lin, L., Chen, F., Xu, Y., Wu, Z., Tang, H., Zhang, X., Tian,
824   F., et al. (2022). Circular RNA vaccines against SARS-CoV-2 and emerging variants.
825   Cell 185, 1728-1744 e1716.
826

827 10. Amaya, L., Grigoryan, L., Li, Z., Lee, A., Wender, P.A., Pulendran, B., Chang, H.Y.
828   (2023). Circular RNA vaccine induces potent T cell responses. Proc. Natl. Acad. Sci.
829   USA 120, e2302191120.
830

831 11. Seephetdee, C., Bhukhai, K., Buasri, N., Leelukkanaveera, P., Lerdwattanasombat,
832   P., Manopwisedjaroen, S., Phueakphud, N., Kuhaudomlarp, S., Olmedillas, E.,
833   Saphire, E.O., et al.(2022). A circular mRNA vaccine prototype producing VFLIP-X
834   spike confers a broad neutralization of SARS-CoV-2 variants by mouse sera.
835   Antiviral Res. 204, 105370.
836

837 12. Zhou, J., Ye, T., Yang, Y., Li, E., Zhang, K., Wang, Y., Chen, S., Hu, J., Zhang, K.,
838   Liu, F., et al. (2024). Circular RNA vaccines against monkeypox virus provide potent
839   protection against vaccinia virus infection in mice. Mol. Ther. 32, 1779–1789.
840

841 13. Karikó, K., Muramatsu, H., Welsh, F.A., Ludwig, J., Kato, H., Akira, S., Weissman,
842   D., (2008). Incorporation of pseudouridine into mRNA yields superior
843   nonimmunogenic vector with increased translational capacity and biological stability.
844   Mol. Ther. 16, 1833-1840.
845

14. Wang, X., Lu, Z., Gomez, A., Hon, G.C., Yue, Y., Han, D., Fu, Y., Parisien, M., Dai, Q., Jia, G., et al. (2014). N6-methyladenosine-dependent regulation of messenger RNA stability. Nature 505, 117-120.

15. Bloom, K., van den Berg, F., Arbuthnot, P. (2021). Self-amplifying RNA vaccines for infectious diseases. Gene therapy 28, 117-129.

16. Schmidt, C., Schnierle, B.S. (2023). Self-Amplifying RNA Vaccine Candidates: Alternative Platforms for mRNA Vaccine Development. Pathogens 12.

17. Pollock, K.M., Cheeseman, H.M., Szubert, A.J., Libri, V., Boffito, M., Owen, D., Bern, H., McFarlane, L.R., O'Hara, J., Lemm, N.M., et al. (2022). Safety and immunogenicity of a self-amplifying RNA vaccine against COVID-19: COVAC1, a phase I, dose-ranging trial. EClinicalMedicine 44, 101262.

18. Oda, Y., Kumagai, Y., Kanai, M., Iwama, Y., Okura, I., Minamida, T., Yagi, Y., Kurosawa, T., Greener, B., Zhang, Y., et al. (2024) Immunogenicity and safety of a booster dose of a self-amplifying RNA COVID-19 vaccine (ARCT-154) versus BNT162b2 mRNA COVID-19 vaccine: a double-blind, multicentre, randomised, controlled, phase 3, non-inferiority trial. Lancet Infect. Dis. 24, 351-360.

19. Liu, C.X., Chen, L.L. (2022). Circular RNAs: Characterization, cellular roles, and applications. Cell 185, 2016-2034.

20. Breuer, J., Barth, P., Noe, Y., Shalamova, L., Goesmann, A., Weber, F. and Rossbach, O. (2022). What goes around comes around: Artificial circular RNAs bypass cellular antiviral responses. Mol. Ther. Nucleic Acids 28, 623-635.

21. Wesselhoeft, R.A., Kowalski, P.S., Parker-Hale, F.C., Huang, Y., Bisaria, N., Anderson, D.G. (2019). RNA circularization diminishes immunogenicity and can extend translation duration in vivo. Mol. cell 74, 508-520. e504.

22. Wesselhoeft, R.A., Kowalski, P.S., Anderson, D.G. (2018). Engineering circular RNA for potent and stable translation in eukaryotic cells. Nat. Commun. 9, 2629.

23. Carabelli, A.M., Peacock, T.P., Thorne, L.G., Harvey, W.T., Hughes, J., Peacock, S.J., Barclay, W.S., De Silva, T.I., Towers, G.J., et al. (2023). SARS-CoV-2 variant biology: immune escape, transmission and fitness. Nat. Rev. Microbio. 21, 162-177.

24. Tao, K., Tzou, P.L., Nouhin, J., Gupta, R.K., de Oliveira, T., Kosakovsky Pond, S.L., Fera, D., Shafer, R.W. (2021). The biological and clinical significance of emerging SARS-CoV-2 variants. Nat. Rev. Genet. 22, 757-773.

25. Planas, D., Saunders, N., Maes, P., Guivel-Benhassine, F., Planchais, C., Buchrieser, J., Bolland, W.H., Porrot, F., Staropoli, I., Lemoine, F., et al. (2022). Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. Nature 602, 671-675.

26. Liu, L., Iketani, S., Guo, Y., Chan, J.F.W., Wang, M., Liu, L., Luo, Y., Chu, H., Huang, Y., Nair, M.S., et al. (2022). Striking antibody evasion manifested by the Omicron variant of SARS-CoV-2. Nature 602, 676-681.

27. Cameroni, E., Bowen, J.E., Rosen, L.E., Saliba, C., Zepeda, S.K., Culap, K., Pinto, D., VanBlargan, L.A., De Marco, A., di Iulio, J., et al. (2022) Broadly neutralizing antibodies overcome SARS-CoV-2 Omicron antigenic shift. Nature 602, 664-670.

28. Corleis, B., Hoffmann, D., Rauch, S., Fricke, C., Roth, N., Gergen, J., Kovacikova, K., Schlottau, K., Halwe, N.J., Ulrich, L., et al. (2023) Efficacy of an unmodified bivalent mRNA vaccine against SARS-CoV-2 variants in female small animal models. Nat. Commun. 14, 816.

29. Li, J., Liu, Q., Liu, J., Fang, Z., Luo, L., Li, S., Lei, Y., Li, Z., Jin, J., Xie, R., et al. (2022) Development of Bivalent mRNA Vaccines against SARS-CoV-2 Variants. Vaccines (Basel) 10.

30. Yuan, Y., Zhang, X., Chen, R., Li, Y., Wu, B., Li, R., Zou, F., Ma, X., Wang, X., Chen, Q., et al. (2022). A bivalent nanoparticle vaccine exhibits potent cross-protection against the variants of SARS-CoV-2. Cell Rep. 38, 110256.

31. Piccoli, L., Park, Y.J., Tortorici, M.A., Czudnochowski, N., Walls, A.C., Beltramello, M., Silacci-Fregni, C., Pinto, D., Rosen, L.E., Bowen, J.E., et al. (2020). Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike Receptor-Binding Domain by Structure-Guided High-Resolution Serology. Cell 183, 1024-1042.

32. Hassan, A.O., Case, J.B., Winkler, E.S., Thackray, L.B., Kafai, N.M., Bailey, A.L., McCune, B.T., Fox, J.M., Chen, R.E., Alsoussi, W.B., et al. (2020). A SARS-CoV-2 Infection Model in Mice Demonstrates Protection by Neutralizing Antibodies. Cell 182, 744-753.

33. Gu, T., Zhao, S., Jin, G., Song, M., Zhi, Y., Zhao, R., Ma, F., Zheng, Y., Wang, K., Liu, H., et al. (2020). Cytokine Signature Induced by SARS-CoV-2 Spike Protein in a Mouse Model. Front. Immunol. 11, 621441.

34. Liljeström, P., Garoff, H. (1991). A new generation of animal cell expression vectors based on the Semliki Forest virus replicon. Nat. Biotech. 9, 1356-1361.

35. Krähling, V., Erbar, S., Kupke, A., Nogueira, S.S., Walzer, K.C., Berger, H., Dietzel, E., Halwe, S., Rohde, C., Sauerhering L., et al. (2023). Self-amplifying RNA vaccine protects mice against lethal Ebola virus infection. Mol. Ther. 31, 374-386.

36. Weaver, S.C., Salas, R., Rico-Hesse, R., Ludwig, G.V., Oberste, M.S., Boshell, J., Tesh, R.B. (1996). Re-emergence of epidemic Venezuelan equine encephalomyelitis in South America. Lancet 348, 436-440.

37. Routhu, N.K., Cheedarla, N., Bollimpelli, V.S., Gangadhara, S., Edara, V.V., Lai, L., Sahoo, A., Shiferaw, A., Styles, T.M., Floyd, K., et al. (2021). SARS-CoV-2 RBD trimer protein adjuvanted with Alum-3M-052 protects from SARS-CoV-2 infection and immune pathology in the lung. Nat. Commun. 12, 3587.

38. McCafferty, S., Haque, A.A., Vandierendonck, A., Weidensee, B., Plovyt, M., Stuchlíková, M., François, N., Valembois, S., Heyndrickx, L., Michiels, J., et al. (2022). A dual-antigen self-amplifying RNA SARS-CoV-2 vaccine induces potent humoral and cellular immune responses and protects against SARS-CoV-2 variants through T cell-mediated immunity. Mol. Ther. 30, 2968-2983.

39. McKay, P.F., Hu, K., Blakney, A.K., Samnuan, K., Brown, J.C., Penn, R., Zhou, J., Bouton, C.R., Rogers, P., Polra, K., et al. (2020). Self-amplifying RNA SARS-CoV-2 lipid nanoparticle vaccine candidate induces high neutralizing antibody titers in mice. Nat. Commun. 11, 3523.

956 40. Erasmus, J.H., Khandhar, A.P., O'Connor, M.A., Walls, A.C., Hemann, E.A., Murapa,
957 P., Archer, J., Leventhal, S., Fuller, J.T., Lewis, T.B., et al. (2020). An Alphavirus-
958 derived replicon RNA vaccine induces SARS-CoV-2 neutralizing antibody and T cell
959 responses in mice and nonhuman primates. Sci. Transl. Med. 12, eabc9396.
960

961 41. De Alwis, R., Gan, E.S., Chen, S., Leong, Y.S., Tan, H.C., Zhang, S.L., Yau, C., Low,
962 J.G., Kalimuddin, S., Matsuda, D., et al. (2021). A single dose of self-transcribing and
963 replicating RNA-based SARS-CoV-2 vaccine produces protective adaptive immunity
964 in mice. Mol. Ther. 29, 1970-1983.
965

966 42. Maruggi, G., Mallett, C.P., Westerbeck, J.W., Chen, T., Lofano, G., Friedrich, K., Qu,
967 L., Sun, J.T., McAuliffe, J., Kanitkar, A., et al. (2022). A self-amplifying mRNA SARS-
968 CoV-2 vaccine candidate induces safe and robust protective immunity in preclinical
969 models. Mol. Ther. 30, 1897-1912.
970

971 43. Verma, R., Saha, S., Kumar, S., Mani, S., Maiti, T.K., Surjit, M. (2021). RNA-protein
972 interaction analysis of SARS-CoV-2 5′ and 3′ untranslated regions reveals a role of
973 lysosome-associated membrane protein-2a during viral infection. mSystems 6,
974 00643-00621.
975

976 44. Mehdi, F., Chattopadhyay, S., Thiruvengadam, R., Yadav, S., Kumar, M., Sinha,
977 S.K., Goswami, S., Kshetrapal, P., Wadhwa, N., Chandramouli Natchu, U., et al.
978 (2021). Development of a fast SARS-CoV-2 IgG ELISA, based on receptor-binding
979 domain, and its comparative evaluation using temporally segregated samples from
980 RT-PCR positive individuals. Front. Microbio. 11, 618097.
981

982 45. Agarwal, S., Kaur, S., Asuru, T.R., Joshi, G., Shrimali, N.M., Singh, A., Singh, O.N.,
983 Srivastva, P., Shrivastava, T., Vrati, S., et al. (2022). Dietary alpha-ketoglutarate
984 inhibits SARS CoV-2 infection and rescues inflamed lungs to restore O(2) saturation
985 by inhibiting pAkt. Clin. Transl. Med. 12, e1041.
986

987 46. Bewley, K.R., Coombes, N.S., Gagnon, L., McInroy, L., Baker, N., Shaik, I., St-Jean,
988 J.R., St-Amant, N., Buttigieg, K.R., Humphries, H.E., et al. (2021). Quantification of
989 SARS-CoV-2 neutralizing antibody by wild-type plaque reduction neutralization,
990 microneutralization and pseudotyped virus neutralization assays. Nat. Protoc. 16,
991 3114-3140.
992

993 47. Chong, Z., Karl, C.E., Halfmann, P.J., Kawaoka, Y., Winkler, E.S., Keeler, S.P.,
994 Holtzman, M.J., Yu, J., Diamond, M.S. (2022). Nasally delivered interferon-lambda
995 protects mice against infection by SARS-CoV-2 variants including Omicron. Cell Rep.
996 39, 110799.

 Figure Legends

 Figure 1. Production and characterization of the mRNA vaccine candidates.

(A) Agarose gel electrophoresis analysis of the *Invitro* transcribed and purified SAM-RBD and

Circ-RBD RNA with (+) and without (-) treatment with RNase R.

(B) Average particle size of SAM-RBD-RNA-LNP and Circ-RBD-RNA-LNP complex.

(C) Average polydispersity Index of SAM-RBD-RNA-LNP and Circ-RBD-RNA-LNP complex.

(D) Agarose gel electrophoresis analysis of the SAM-RBD and the Circ-RBD-RNA

incorporation into the LNP. RNA-LNP formulations were treated with Triton-x-100, as

indicated.

(E) Average percent SAM-RBD and Circ-RBD RNA incorporation into the LNP. RNA-LNP

formulations were treated with Triton-x-100, as indicated.

(F) Scanning electron micrograph of SAM-RBD-LNP and Circ-RBD-LNP. Scale:100nm.

(G) Western blot analysis of the RBD protein in HEK239T cells transfected with the SAM-

RBD-LNP and Circ-RBD-LNP (upper panel). Mock: Lysate from HEK293T cells treated with

LNP. Lower panel: Anti GAPDH western blot of aliquots of the cell lysate used in the upper

panel.

(H) Anti-RBD western blot of RBD protein in the cell culture medium of HEK239T cells

mentioned in (G).

(I) Anti-RBD western blot of aliquots of sample used in (H) under reducing and non-reducing

conditions, as indicated.

(J) RT-qPCR analysis of the indicated genes in total RNA isolated from HEK239T cells

transfected with the SAM-RBD, Circ-RBD-RNA or poly I:C, as indicated. Mock: lipofectamine

3000 transfected cells. PKR and OAS1 are shown in the right panel for clear representation.

Dotted line indicates the background level. Data are mean±SD of three independent

experiments. *P*-values were calculated by the non-parametric student's t-test, using the

Mann-Whitney test.

1033

1034

1035

1036

1037

1038

1039

1040

1041

Figure 2. Immunogenicity assessment of the SAM-RBD and Circ-RBD-RNA vaccine

formulations.

(A) Schematic of the immunization study. Balb/c mice were intradermally injected with $10\mu g$

SAM-RBD or Circ-RBD-RNA-LNP on day 1 and 14. Blood was collected via retro-orbital route

prior to the first immunization and subsequently on day 7 and day 28. Animals were euthanized

on day 28, blood and spleen were collected.

(B) Anti-RBD IgM level at 1 week post-immunization in the sera (1:500 dilution) of mice

injected with the placebo (PBS), SAM-RBD and Circ-RBD-LNP, evaluated by ELISA.

(C) Anti-RBD IgG titre at 4 weeks post-immunization in the sera of mice described in (B),

evaluated by ELISA.

(D) 50% virus neutralization titre (NT_{50}) at 4 weeks post-immunization in the sera of mice

described in (B), evaluated by PRNT assay.

(E) Anti-RBD IgG1 level (1:500 dilution) at 4 weeks post-immunization in the sera of mice

described in (B), evaluated by ELISA.

(F) Anti-RBD IgG2a level (1:500 dilution) at 4 weeks post-immunization in the sera of mice

described in (B), evaluated by ELISA.

(G) Anti-RBD IgG2b level (1:500 dilution) at 4 weeks post-immunization in the sera of mice

described in (B), evaluated by ELISA.

(H) Ratio of anti-RBD T_H1 antibody (IgG2a and IgG2b) to T_H2 (IgG1) antibody in the sera of

mice injected with the SAM-RBD and Circ-RBD at 4 weeks, evaluated by ELISA.

(I) Percentage of IFNγ and CD4 positive T-cells in the splenocytes, following stimulation with

the SARS-CoV-2-Spike-RBD peptide pool, evaluated by FACS.

(J) Percentage of IFNγ and CD8 positive T-cells in the splenocytes, following stimulation with

the SARS-CoV-2-Spike-RBD peptide pool, evaluated by FACS.

(K) Percentage of TNFα and CD4 positive T-cells in the splenocytes, following stimulation with

the SARS-CoV-2-Spike-RBD peptide pool, evaluated by FACS.

(L) Percentage of TNFα and CD8 positive T-cells in the splenocytes, following stimulation with

the SARS-CoV-2-Spike-RBD peptide pool, evaluated by FACS.

(M) Percentage of IL-4 and CD4 positive T-cells in the splenocytes, following stimulation with

the SARS-CoV-2-Spike-RBD peptide pool, evaluated by FACS.

(N) Percentage of IL-4 and CD8 positive T-cells in the splenocytes, following stimulation with

the SARS-CoV-2- Spike-RBD peptide pool, evaluated by FACS.

In all images, bars represent the average and dots represent the value obtained from each

animal. Data are mean$\pm$SD, *p*-values were calculated by the non-parametric student's t-test,

using the Mann-Whitney test (B-H) and the non-parametric one-way ANOVA, using the

Kruskal-wallis test (I-N). Values showing distinct difference from the majority in the group were

not considered for calculating the p-value in the panels I-L.

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

Figure 3. Stability and durability of the immune response induced by the Circ-RBD-LNP
vaccine formulation.

(A) Average size of the Circ-RBD-LNP stored at 4^0C for 0, 2 and 4 weeks, evaluated by
dynamic light scattering.

(B) Average polydispersity index of the Circ-RBD-LNP, stored as indicated in (A).

(C) Percentage of RNA encapsulated into the LNP in the Circ-RBD-LNP formulation, stored
as indicated in (A). Equal aliquots of the samples were treated with the Triton X-100.

(D) Anti-RBD IgG titre in the sera of mice immunized for 4 weeks with the Circ-RBD-LNP,
stored as indicated in (A).

(E) 50% virus neutralization titre (NT_{50}) in aliquots of the sera used in (D), evaluated by PRNT
assay.

(F) Anti-RBD IgG titre at 0, 1, 3 and 9 months post-immunization in the sera of mice immunized
with the Circ-RBD-LNP, evaluated by ELISA.

(G) 50% virus neutralization titre (NT_{50}) in aliquots of the sera described in (G), evaluated by
PRNT assay.

(H) Anti-RBD IgG titre at 9 months post-immunization with the Circ-RBD-LNP, in male and
female mice, evaluated by ELISA.

(I) 50% virus neutralization titre (NT_{50}) in aliquots of the sera described in (I), evaluated by
PRNT assay.

Data are mean±SD of triplicate samples (A-C) and six mice/group (D-I), *p*-values were
determined by the unpaired student's t-test, using the Mann-Whitney test.

Figure 4. Immunization with the Circ-RBD protects mice from intranasal SARS-CoV-2

challenge-induced disease symptoms.

(A) Schematic of the Circ-RBD-LNP immunization and the SARS-CoV-2 challenge study in

mice. Blood was collected before immunization on day 1 and day 14. Mice were injected with

the Circ-RBD-LNP or PBS (placebo group) on day 1 and day 14. Intranasal transduction of

the hACE2 was done on day 28, followed by intranasal administration of the SARS-CoV-2

(Wuhan) on day 33. Mouse weight was measured every day from day 33 to day 37. Mice were

euthanized on day 37 and blood, lung tissue, bronchoalveolar lavage (BAL) were collected.

(B) Effect of the SARS-CoV-2 (Wuhan) infection on mice weight. Weight of each mouse in the

placebo and the Circ-RBD immunization group was measured (N=6) every day from day 33-

37 and the average of percent weight change with respect to the weight on day 33 (considered

100%) is plotted in the graph.

(C) Anti-RBD IgG titre in sera of the placebo and the Circ-RBD immunized, SARS-CoV-2

(Wuhan) infected mice, as indicated in (A). Sera collected on day 33 and 37 were evaluated

by ELISA.

(D) Measurement of viral load (SARS-CoV-2,Wuhan) in the lung tissue of the placebo and the

Circ-RBD immunized mice. Samples were collected on day 37, as indicated in (A) and viral

load was measured by plaque assay.

(E) Measurement of viral RNA level in aliquots of the lung tissue described in (D), determined

by RT-qPCR.

(F) Measurement of viral load in the BAL of the placebo and the Circ-RBD immunized, SARS-

CoV-2 (Wuhan) infected mice, collected on day 37 and evaluated by plaque assay.

(G) Lung morphology of the placebo and the Circ-RBD immunized, SARS-CoV-2 (Wuhan)

infected mice, collected on day 37.

(H) Hematoxylin and eosin-stained image of lung tissue section prepared from the placebo

and Circ-RBD immunized, SARS-CoV-2 (Wuhan) infected mice, collected on day 37. Images

were acquired at 40X (left panel) and 200X (right panel) magnification. Black and red arrows

indicate peribronchiolar and perialveolar infiltration, respectively, black arrow-head indicates

neutrophils, double headed black arrow indicates alveolar septa thickening and dotted region

indicates epithelial damage.

(I) Lung infiltration score of the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan)

infected mice, evaluated by analysis of the hematoxylin and eosin-stained images of tissue

sections prepared from the lungs collected on day 37.

(J) Percentage of hematoxylin-stained area in the hematoxylin and eosin-stained images of

lung tissue section of the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan) infected

mice, shown in (H).

(K) RT-qPCR analysis of the indicated cytokines in the total RNA isolated from lung tissue of

the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan) infected mice.

Data are shown as the mean±SD, *p*-values were determined by the unpaired student's t-test,

using the Mann-Whitney test.

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

Figure 5

Figure 5. Immunization with the Circ-RBD protects mice from the intratracheal SARS-
CoV-2 challenge-induced disease symptoms.

(A) Schematic of the Circ-RBD-LNP immunization and the SARS-CoV-2 challenge study in

mice. Blood was collected before immunization on day 1 and day 14. Mice were injected with

the Circ-RBD-LNP or PBS (placebo group) on day 1 and day 14. Intra-tracheal transduction

of the hACE2 was done on day 28, followed by intranasal infection with the SARS-CoV-2

(Wuhan) on day 33. Mouse weight was measured every day from day 33-54. Mice were

euthanized on day 40, 47 and 54 and blood and lung tissue were collected.

(B) Effect of the SARS-CoV-2 (Wuhan) infection on mice weight. Weight of each mouse in the

placebo and the Circ-RBD immunization group was measured (initial no. of mice in both

groups = 7) on the indicated days between day 33 to day 54 and the average of percent weight

change with respect to the weight on day 33 (considered 100%) is plotted in the graph. Few

animals in the placebo group died during the experimental period, as indicated in (D). Average

weight of the remaining animals is plotted.

(C) Survival of mice in the placebo and the Circ-RBD immunized group was monitored every

day from day 33 to day 54.

(D) Anti-RBD IgG titre in the sera of the placebo and Circ-RBD immunized, SARS-CoV-2

(Wuhan) infected mice as indicated in (A). Sera collected on day 40 and day 47 were evaluated

by ELISA.

(E) Measurement of viral load (SARS-CoV-2,Wuhan) in lung tissue of the placebo and Circ-

RBD immunized, SARS-CoV-2 (Wuhan) infected mice, evaluated by plaque assay.

(F) Measurement of the viral RNA level in aliquots of the lung tissue described in (E),

determined by RT-qPCR.

(G) RT-qPCR analysis of the indicated cytokines in the total RNA isolated from lung tissue of

the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan) infected mice, collected on day

54.

(H) Lung morphology of the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan)

infected mice, collected on day 54.

(I) Representative hematoxylin and eosin-stained image of lung tissue section prepared from

the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan) infected mice, collected on day

54. Images were acquired at 40X (left panel) and 200X magnification (right panel). Black and

red arrow indicate peribronchiolar and perialveolar infiltration, respectively; black arrow-head

indicate neutrophils, double headed black arrow indicates alveolar septa thickening and dotted

region indicates epithelial damage.

(J) Lung infiltration score of the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan)

infected mice, evaluated by analysis of the hematoxylin and eosin-stained images of tissue

sections prepared from the lung tissue collected on day 54.

(K) Percentage of hematoxylin-stained area in the hematoxylin and eosin-stained images of

lung tissue section of the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan) infected

mice, collected on day 54.

In (B) and (D), data are shown as the mean$\pm$SD and p-values were determined by the unpaired

student's t-test, using the Mann-Whitney test.

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

Figure 6. Assessment of immunogenicity and virus neutralization potential of the Circ-

RBD^{Del} and Circ-RBD^{Omi} bivalent vaccine formulation.

(A) Western blot analysis of the RBD protein in the HEK239T cells transfected with the Circ-

RBD^{Del}-LNP (upper panel). Mock: HEK293T cells treated with LNP. Lower panel: Anti GAPDH

western blot of aliquots of the samples used in the upper panel.

(B) Western blot analysis of the RBD protein in the HEK239T cells transfected with the Circ-

RBD^{Omi}-LNP (upper panel-lower exposure, middle panel-higher exposure of the same blot).

Mock: HEK293T cells treated with LNP. Lower panel: anti GAPDH western blot of aliquots of

the samples used in the upper panel.

(C) Anti-RBD^{Wuhan} IgG titre at 4 weeks post-immunization in the sera of mice injected with the

placebo (PBS), Circ-RBD^{Wuhan}, Circ-RBD^{Del}, Circ-RBD^{Omi} and bivalent Circ-RBD^{Del}+Circ-

RBD^{Omi} vaccine formulations, evaluated by ELISA.

(D) Anti-RBD^{Del} IgG titre at 4 weeks post-immunization in the sera of mice described in (C),

evaluated by ELISA.

(E) Anti-RBD^{Omi} IgG titre at 4 weeks post-immunization in the sera of mice described in (C),

evaluated by ELISA.

(F) 50% virus neutralization titre (NT$_{50}$) against the SARS-CoV-2 (Wuhan) at 4 weeks post-

immunization in the sera of mice described in (C), evaluated by FRNT assay.

(G) 50% virus neutralization titre (NT$_{50}$) against the SARS-CoV-2 (Delta) at 4 weeks post-

immunization in the sera of mice described in (C), evaluated by FRNT assay.

(H) 50% virus neutralization titre (NT$_{50}$) against the SARS-CoV-2 (Omicron BA.1) at 4 weeks

post-immunization in the sera of mice described in (C), evaluated by FRNT assay.

In (C)-(E), data are shown as the mean±SD of pooled sera of 6 animals in each group, assayed

in triplicate. In (F)-(H), data are shown as the mean±SD of 6 animals in each group, assayed

individually. *p*-values were determined by the non-parametric one-way ANOVA, using the

Kruskal-wallis test.

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

 Supplementary Figure legends

Figure S1. Selection of immunization route and dose of RNA-LNP formulation that

elicits significant immune response in mice.

(A) Anti-RBD IgG titre at 4 weeks post-immunization in the sera of mice injected with the

placebo (PBS) and the Circ-RBD-LNP via intramuscular (IM) or intradermal (ID) route,

evaluated by ELISA.

(B) 50% virus neutralization titre (NT_{50}) at 4 weeks post-immunization in the sera of mice

described in (B), evaluated by PRNT assay.

(C) Anti-RBD IgG titre at 4 weeks post-immunization in the sera of mice intradermally injected

with the placebo (PBS) or 5, 10, 25µg of Circ-RBD-LNP, evaluated by ELISA.

(D) 50% virus neutralization titre (NT_{50}) at 4 weeks post-immunization in the sera of mice

described in (C), evaluated by PRNT assay.

Data are shown as the mean±SD of pooled sera from five animals, assayed in triplicate; *p*-

values were determined by the unpaired student's t-test, using the Mann-Whitney test.

1344

1345

1346

1347

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

1363

Figure S2. Analysis of intracellular IFNγ, TNFα and IL-4 levels in the RBD peptide pool-

stimulated CD4, CD8 positive T-cells derived from the splenocytes of Circ-RBD

immunized mice.

(A)-(D) Representative FACS plot showing the gating strategy of lymphocytes (A), single cells

(B), CD3$^+$ (C) and CD4$^+$CD8$^+$(D) T-cells.

(E)-(H) Representative FACS plot showing the gating strategy of CD4$^+$ memory T-cells (E) and

quantification of IFNγ (F), IL-4 (G) and TNFα (H) in CD4$^+$ memory T-cells.

(I)-(L) Representative FACS plot showing the gating strategy of CD8$^+$ memory T-cell (J) and

quantification of IFNγ (K), IL-4 (K) and TNFα (L) in CD8$^+$ memory T-cells.

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

Figure S3. Immunization with the Circ-RBD protects mice from intra-tracheal SARS-

CoV-2 challenge.

(A) Effect of the SARS-CoV-2 (Wuhan) infection on mice weight. Weight of each mouse in the

placebo and the Circ-RBD immunization group was measured (initial no. of mice in both

groups = 7) on the indicated days between day 33 to day 54 and the percent weight change

with respect to the weight on day 33 (considered 100%) is plotted in the graph. Few animals

in the placebo group died during the experimental period.

(B) Lung morphology of the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan)

infected mice, collected on day 47.

(C) Representative hematoxylin and eosin-stained image of lung tissue section prepared from

the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan) infected mice, collected on day

47. Images were acquired at 40X (left panel) and 200X magnification (right panel).

(D) Lung morphology of the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan)

infected mice, collected on day 40.

(E) Representative hematoxylin and eosin-stained image of lung tissue section prepared from

the placebo and Circ-RBD immunized, SARS-CoV-2 (Wuhan) infected mice, collected on day

40. Images were acquired at 40X (left panel) and 200X magnification (right panel).