1 **Growth patterns of theoretical bite force and jaw musculature in southern sea otters**

2 (***Enhydra lutris nereis***)

4 Chris J. Law[1,2]

5 [1]Burke Museum and Department of Biology, University of Washington

6 [2]Department of Integrative Biology, University of California Berkeley

Abstract

The transition from milk to solid food requires drastic changes in the morphology of the feeding

apparatus and its performance. As durophagous mammals, southern sea otters exhibit significant

ontogenetic changes in cranial and mandibular morphology to presumably enable them to feed

on a variety of hard-shelled invertebrate prey. Juvenile sea otters begin feeding independently by

6–8 months of age, but how quickly they reach sufficient maturity in biting performances

remains unknown. Here, I found that theoretical bite force of southern sea otters does not reach

full maturation until during the adult stage at 3.6 and 5.0 years of age in females and males,

respectively. The slow maturation of biting performance can be directly attributed to the slow

growth and development of the cranium and the primary jaw adductor muscle (i.e., the

temporalis) and may ultimately impact the survival of newly weaned juveniles by limiting their

ability to process certain hard-shelled prey. Alterative foraging behaviors such as tool use,

however, may mitigate the disadvantages of delayed maturation of biting performance. In

analyses of sexual dimorphism, I found that female otters reached bite force maturation earlier

whereas male otters exhibit initial rapid growth in bite force–to quickly reach sufficient biting

performances needed to process prey early in life–followed by a slower growth phase towards

bite force maturation that coincides with sexual maturity. This biphasic growth in bite force

suggests that male to male competition for resources and mates exhibits strong selection in the

growth and development of skull form and function in male otters. Overall, this study

demonstrates how the analysis of anatomical data can provide insight on the foraging ecologies

and life histories of sea otters across ontogeny.

Key words: Carnivora, functional morphology, jaw biomechanics, ontogeny, performance,

sexual dimorphism, skull morphology

Introduction

The growth and development of the vertebrate feeding apparatus often correspond to

major ontogenetic shifts in diet and resource use (e.g., Herrel and O'Reilly 2006; Baliga and

Mehta 2014; Gignac and Erickson 2014; Santana and Miller 2016; Higgins et al. 2018; Chatar et

al. 2024). In mammals, weaning is a major ontogenetic dietary shift, as juveniles must transition

from a liquid food source (i.e., milk) during the nursing stage to a diet consisting of solid foods

that must be obtained and processed for survival. Unsurprisingly, the mammalian feeding

apparatus undergoes tremendous ontogenetic changes in skull morphology and musculature to

accommodate these dietary shifts. Ontogenetic changes in skull size (growth) and skull shape

(development) facilitate the enlargement of jaw muscles and their attachment areas, increase in

osteological robustness, and stronger bite force (Tanner et al. 2010; La Croix et al. 2011b;

Segura 2015; Forbes☐Harper et al. 2017; Law et al. 2017; Law 2020; Stanchak et al. 2023).

Furthermore, selection is expected to favor individuals that exhibit fast rates and/or short

durations of growth and development, and thus more quickly become efficient foragers on adult

diets (Herrel and Gibb 2006; Gignac and Santana 2016). The need for quick transitions to adult

morphologies and performance may be particularly important for species with highly specialized

diets such as carnivory or durophagy, a diet that consists of hard or tough material including

bamboo, bones, exoskeletons, or shells. In carnivoran mammals, the growth and development of

the carnivoran skull have been investigated by an increasing number of researchers. These

ontogenetic changes (e.g. the flattening of the dorsal cranial profile, development of a more

pronounced sagittal crest, broadening of the zygomatic arches, and increase in size) all contribute

to stronger bite performance (Tanner et al. 2010; La Croix et al. 2011a; Segura and Prevosti

2012; Tarnawski et al. 2014; Forbes☐Harper et al. 2017; Chatar et al. 2024). Although adults of

durophagous species exhibit strong biting performances to specialize on hard material (Binder

and Van Valkenburgh 2000; Christiansen and Wroe 2007; Figueirido et al. 2014; Tseng et al.

2017), these traits are often not fully developed in the skulls of juveniles and may constrain their

ability to break into their hard, tough food items. Thus, how quickly durophagous juveniles reach

sufficient maturity in biting performances requires further investigation.

Southern sea otters (*Enhydra lutris nereis*) are a great model species to examine the

growth and development of biting performance and the underlying jaw musculature. This

durophagous marine mustelid feeds on a variety of hard-shelled invertebrates such as clams,

crabs, mussels, snails, and urchins (Riedman and Estes 1990). As one of the smallest marine

mammals, sea otters exhibit the highest known mass-specific metabolic rates of marine

mammals (Morrison et al. 1974; Costa and Kooyman 1984; Yeates et al. 2007) and consequently

must spend 20–50% of the day foraging to consume 20–30% of their body mass in food per day

(Morrison et al. 1974; Costa and Kooyman 1982; Estes et al. 1986; Ralls and Siniff 1990;

Thometz et al. 2016). Thus, as adults, sea otters exhibit several cranial adaptations that facilitate

durophagy and enable them to feed effectively including short, blunt skulls (Riley 1985); taller

and wider mandibular rami (Timm-Davis et al. 2015); and bunodont dentition with fracture-

resistant dental enamel (Constantino et al. 2011; Ziscovici et al. 2014). Southern sea otters

exhibit dramatic changes in cranial and mandibular shape and size over ontogeny to reach these

adult morphologies including the widening of the zygomatic arches and the mastoid processes,

dorsoventral compression of the cranium, shortening of the jaw length, and anterodorsal

expansion of the coronoid process (Law et al. 2017). These ontogenetic changes enable southern

sea otters to exhibit positive allometric increases in theoretical bite force and underlying jaw

adductor musculature with respect to body size and skull size (Law et al. 2016b). Southern sea

otters obtain most of their nourishment from nursing during the first few weeks of life (Payne &

Jameson, 1984). However, by 6 weeks of age, pups begin soliciting small pieces of food from

their mothers, and by 8.9 weeks of age, pups begin to catch and consume some of their own prey

(Staedler 2011). Pups can break open some hard-shelled prey items by 14–20 weeks of age and

begin to display proto-tool use behaviors (Staedler 2011). Pups are weaned and completely

independent by 6–8 months of age (Payne and Jameson 1984; Jameson and Johnson 1993).

Although significant ontogenetic changes in skull morphology occur during these first 6–8

months, the full growth and development of the skull does not occur until well after weaning

(Law et al. 2017). The mandible does not reach adult maturity until the subadult stage at 1–3

years old, and most aspects of cranial maturity do not occur until after mature body size is

obtained during the adult stage at 4–5.5 years old (Law et al. 2017). Law et al. (2017) postulated

that the slower maturation of the skull may hamper the ability of newly weaned juveniles to

successfully process hard-shelled prey by constraining jaw adductor muscle size, and thus biting

performance. Although positive allometry in bite force suggests fast growth to overcome the

smaller, less developed morphology of the juvenile stage (Law et al. 2016), the rate and duration

of growth and development of biting performance remains to be quantified.

In this study, I quantified the age at which juvenile southern sea otters reach maturity in

biting performance. To accomplish this goal, I conducted dissection-based estimations of

theoretical bite forces (Davis et al. 2010; Santana et al. 2010) across an ontogenetic series of

southern sea otters and generated a growth curve model of theoretical bite force. I also generated

growth curves for the underlying jaw adductor musculature and lever mechanics known to

contribute to bite force generation in mammals. Lastly, because male sea otters attain maturation

of skull morphologies earlier than female otters (Law et al. 2017), I tested whether male otters

also exhibit faster maturation of theoretical bite force and underlying jaw musculature.

Methods

Specimens and gross dissections

Osteological and muscular data were gathered from 74 southern sea otter specimens. 55

of these specimens were obtained from Law et al. (2016b) and 19 specimens were added in this

study. All 74 specimens originated from naturally deceased southern sea otters obtained from the

California Department of Fish and Wildlife (CDFW) Marine Wildlife Veterinary Care and

Research Center between October 2013 and June 2017. The age of each specimen was estimated

by CDFW using a suite of morphological characteristics including total body length, tooth wear,

and closure of cranial sutures (Nicholson et al. 2020). Ages of studied specimens ranged from 1

day to 11 years (Fig. 1). Following CDFW, I also classified each individual into one of five age

classes based on their estimated age: pups (0–6 months), immatures (6–12 months), subadults

(1–3.5 years), adults (3.5–10 years), and aged adults (>10 years). All specimens stranded along

the central California coast, within and throughout the current southern sea otter range from

Pigeon Point in the north to Gaviota in the south. All specimens were deposited at the California

Academy of Sciences. Overall, my dataset consisted of 35 female otters (nine pups, six

immatures, eight subadults, 10 adults, and two aged adults) and 39 male otters (seven pups, six

immatures, seven subadults, 15 adults, and four aged adults).

125   I dissected the major jaw adductor muscles (superficial temporalis, deep temporalis,

superficial masseter, deep masseter, and zygomaticomandibularis). While there was distinct

separation between the superficial and deep temporalis, there was inadequate separation between

the superficial masseter, deep masseter, and zygomaticomandibularis muscles (Scapino 1968;

Law et al. 2016b). I did not dissect the medial and lateral pterygoids because these muscles are

positioned deep along the medial side of the mandible and cannot be excised intact fully.

Muscles were removed from the left side of the skull, blotted dry, and weighed to the nearest 0.1

g using a digital scale. I measured the length and pennation angle of muscle fibers by digesting

and separating the muscles in a solution of 15% aqueous nitric acid for 3–7 d depending on

muscle size (Biewener and Full 1992). Muscle fiber lengths were measured to the nearest 0.01

mm using a digital caliper. Skulls were subsequently cleaned by a dermestid beetle colony at the

California Academy of Sciences or with a maceration tank at CDFW. I then photographed and

digitally measured the condylobasal length (i.e., distance from the anteriormost point on the

premaxillae to the plane of the posterior surface of the occipital condyles), zygomatic breadth

(i.e., greatest distance across the zygomatic arches), and cranial height (i.e., distance

perpendicular to the palate plane from the lateralmost point of the mastoid process to the point of

the sagittal crest directly superior to the mastoid process) of each cranium using ImageJ

(Schneider et al. 2012). I then calculated the geometric mean of these three measurements as a

proxy for overall skull size of each specimen. The geometric mean was derived from the Nth

root of the product of N (N = 3) linear measurements.

Estimating theoretical bite force

I calculated a theoretical bite force for each specimen by modeling the mandible as a

static third-class lever (Fig. 2). In this model, an axis passing through the temporomandibular

joints (TMJs) serves as a fulcrum and muscle forces generated by contractions of the temporalis

and masseter adductor muscles create rotation of the lower jaw about this fulcrum (Davis et al.

2010). Under static lever equilibrium, the force of biting balances the rotation of the lower jaw

created by these muscle forces. To simplify the bite force model, I treated the superficial and

deep temporalis subdivisions of the temporalis as one muscle (i.e., the temporalis) and the

superficial masseter, deep masseter, and zygomaticomandibularis as another muscle (i.e., the

masseter). I first calculated the physiological cross sectional area (PCSA) for the temporalis and

masseter muscles based on their masses (m), mean fiber lengths (f), fiber pennation angles (θ),

and muscle density as a constant (ρ = 1.06 g/cm (Mendez and Keys 1960)), (Sacks et al. 1982):

$$PCSA = \frac{m}{\rho} \frac{\cos(\theta)}{f}$$

Pennation angle remains difficult to measure, particularly in muscle complexes that consist of

multiple muscles with different lines of action. Thus, as part of my two-muscle bite force model

simplification, I used a pennation angle of 0° in estimations of both muscle PCSAs.

I then estimated muscle forces of the temporalis and masseter by multiplying the PCSA

with a muscle stress value of 30 N/cm^2 (Davis et al. 2010). I modeled each muscle force as a

single force vector (F_T and F_M) perpendicular to the muscle in-levers and applied them to the

insertion points of the temporalis (the top of the coronoid process of the mandible) and the

masseter (the mid-point between the anteriormost edge of the masseteric fossa and the angular

process of the mandible) (Fig. 2). The temporalis in-lever (MAT) is the in-lever length measured

from the insertion point of the temporalis to the TMJ, and the masseter in-lever (MAM) is the in-

lever length from the insertion point of the masseter to the TMJ. Estimations of muscle forces

assumed that all jaw muscles were maximally activated.

I calculated theoretical bite forces in between the first and second molars of the lower jaw

because the molars are used to crush items before consumption (Riedman and Estes 1990).

Maximum theoretical bite force (BF) was estimated by adding the moment (product of the force

vector and in-lever length) of each jaw adductor muscle, dividing by the out-lever ,and

multiplying by 2 to account for bilateral biting:

$$BF = 2\frac{F_T * MAT + F_M * MAM}{OL}$$

where F_T and F_M are the force vectors of the temporalis and masseter, respectively; MAT is the

temporalis in-lever length; MAM is the masseter in-lever length; and OL is the out-lever

measured from the bite point and the TMJ (Fig. 2). All in-lever and out-lever measurements were

taken from photographs of the lateral view of the mandible in ImageJ v. 1.48 (Schneider et al.

2012). Because all muscles were dissected and measured at occlusion in all specimens, muscle

force vectors were also estimated based on gape angles at occlusion (i.e., 0º). Lastly, I calculated

the mechanical advantage (MA) of jaw closing based on the temporalis (MAT/OL) and masseter

(MAM/OL) muscles. MA is the ratio of the length of the in-lever (MAT or MAM) divided by the

length of the out-lever (OL). A relatively higher mechanical advantage suggests a jaw more

optimized for stronger bite force.

Statistical Analyses

I determined the rate and the age at which skull size, theoretical bite force, underlying

muscle parameters (i.e., PCSA, mass, and fiber length), and mechanical advantage of jaw closing

reached maturity using Brody growth models:

$$x(t) = A(1 - Be^{-kt})$$

where *x(t)* is the trait of interest at time *t*, *A* is the asymptotic maturity, *k* is the exponential decay

of the growth rate and *B* is the scaling constant signified by the ratio between the initial growth

rate at t = 0 and *k* (Brody 1945). Thus, the initial growth rate (*L*) can be calculated as the product

between *B* and *k*. I used Brody models in this study to keep the parameters consistent with

previous work on sea otter skull growth that found that Brody models were the best fitting

models compared to a suite of other growth models (Law et al. 2017). I fit Brody models to each

trait separately by sex because southern sea otters exhibit sexual dimorphism in the skull (Law et

al. 2016a, 2017). I then determined whether the parameter estimates significantly differed

between the sexes by comparing the mean values of one sex with the 95% confidence intervals

of the mean with the other sex. I also reported the age of maturity of each trait as the estimated

age at which the trait reached the lower limit of the 95% confidence interval for its asymptotic

value (Zelditch et al. 2003). All growth models were fit using the nls function in R 4.3.1

(R Core Team 2023).

Results

205  Asymptotic values of theoretical bite force significantly differed between male and

206 female southern sea otters; adult males exhibited 142% greater theoretical bite forces ($BF_{\female}$ =

207 437.7 N; $BF_{\male}$ = 620.7 N) than adult females (Table 1; Fig. 3). Although females displayed faster

208 initial growth rates, males displayed slower rates of growth decay to reach their theoretical bite

209 force, leading to later maturation (i.e., 95% of the asymptotic values) in theoretical bite force (5.0

210 years) compared to females (3.6 years). However, males reached the asymptotic value of

211 theoretical bite force of females in 1.8 years.

212  Asymptotic values of the temporalis PCSA, mass, and fiber lengths also significantly

213 differed between males and females (Table 1; Fig. 4). Adult male otters exhibited greater

214 temporalis PCSA, larger temporalis mass, and longer temporalis fiber lengths but displayed

215 slower initial growth rates (significantly lower decay rates with similar scaling constants) and

216 thus later maturation in these traits compared to adult female otters (Table 1). Adult male otters

217 also exhibited greater masseter PCSA and larger masseter mass than adult female otters;

218 however, neither scaling constants nor rates of growth decay in these masseter traits do not

219 significantly differ between the sexes, suggesting that increased growth duration rather than

220 slower growth rate contributed to the larger PCSA and mass found in males. In contrast,

221 asymptotic values of masseter fiber lengths did not significantly differ between the sexes.

222  Asymptotic values and rates of growth decay of the mechanical advantage of jaw closing

223 for either the temporalis or masseter did not differ between males and females (Table 1; Fig. 5).

224 Both female and male otters reached mature mechanical advantage during weaning (6–12

225 months of age). The 95% confidence interval of the k parameter in the model for temporalis

226 mechanical advantage in males includes 0, suggesting that the Brody model may be a poor fit to

227 this data. Asymptotic values of temporalis in-lever (MAT) and out-lever to the molars (O)

differed between the sexes; adult male otters exhibited significantly longer MAT and OL than

females (Table 1; Fig. 5). Interestingly, the temporalis in-lever of males reached maturation

earlier than the out-lever of males by over a year, which may contribute to the poor fit of the k

parameter. In contrast, rates of growth decay of masseter in-lever (MAM) differed between the

sexes but not their asymptotic values (Table 1; Fig. 5).

Discussion

The transition from milk to a diet consisting of hard-shelled invertebrate prey presumably

requires equally drastic changes in the morphology of the feeding apparatus and biting

performance. In this study, I found that bite force does not reach full maturity until the adult

stage (3.6 years old in females and 5.0 years old in males). The slow maturation of biting

performance may limit newly weaned juveniles from processing certain hard-shelled prey and

may therefore impact their survival. The first year post-weaning age class often shows depressed

survivorship (Monson et al. 2000; Tinker et al. 2006), as newly weaned otters exhibit the highest

daily energy demands of all juvenile age classes (Thometz et al. 2014) but do not have the

benefit of obtaining food from their mothers and must compete with conspecific adults for the

same food resources (Riedman and Estes 1990) in already resource-poor environments (Tinker et

al. 2008; Layman et al. 2015; Tinker and Hatfield 2015). Together, these results suggest that

newly weaned southern sea otters are at a disadvantage in foraging until the early stages of

adulthood when they reach adult skull morphologies and biting performances. Below, I discuss

how alternative feeding behaviors may help mitigate delayed maturation of biting performance

and how craniomandibular asynchronization and sexual dimorphism influence bite performance

growth. Lastly, I compare how sea otter bite force growth compares to other carnivorans and

discuss caveats to my bite force model.

Tool use may mitigate delayed maturation of biting performance

Alternative feeding behaviors can potentially provide pathways to foraging success even

with suboptimal morphologies or performances such as a less developed feeding apparatus. One

such feeding behavior found in southern sea otters is the use of rocks, shells, or other hard

objects as hammers or anvils to crack open their hard-shelled prey (Riedman and Estes 1990).

The use of tools varies greatly among individual otters and is associated with individual variation

in diet specialization; for instance, adult otters specializing on snails or clams tend to use tools

frequently whereas adult otters specializing on urchins, crabs, and small bivalves tend not to use

tools (Fujii et al. 2015, 2017). Overall, individuals that used tools frequently processed harder,

larger prey items that would be typically inaccessible with biting alone (Law et al. 2024b). These

benefits may be enhanced in juvenile otters as they exhibit weaker bite forces compared to their

adult conspecifics. All sea otter pups exhibit proto-tool use behaviors (such as repetitive

pounding of objects on their chests) and begin using tools at the age of 14 weeks (Staedler 2011).

During this stage, an average otter's bite force is only 88 N, far lower than the amount of force it

takes to break open even the least hard prey such as urchins (mean fracture force = 185 N) and

crabs (mean fracture force = 247 N) (Law et al. 2016b, 2024b). Tool use behavior does not vary

with the mother's diet specialization (Staedler 2011). Thus, although urchin and crab specialists

tend not to use tools as adults, juvenile otters use tools to process these prey types (Staedler

2011), suggesting that tool use may contribute to their foraging success and compensate for their

weaker bite forces. Furthermore, because larger prey items contain more calories but are harder

to break open (Law et al. 2024b), tool use may also enable juveniles to gain access to additional

calories that would have been inaccessible with their immature bite forces. These tool use

benefits may be particularly important for female otters as they used tools with greater

frequencies than male otters, which enabled them to consume prey items that were 21–35%

harder compared with males despite their smaller crania and weaker bite forces (Law et al.

2024b). Therefore, it is tempting to postulate that tool use may serve as an essential behavior in

facilitating prey access for individuals with weak bite forces that are still growing/developing or

are smaller. Nevertheless, the majority of tool use research in sea otters has focused on adults

and additional work is required to further understand the importance of tool use on foraging

success and survival in juvenile otters.

Effects of craniomandibular asynchronization on biting performance growth

Asynchronized growth and development between the cranium and mandible found in

southern sea otters may explain the underlying factors that are affecting the maturation rate of

biting performance. Compared to the mandible, the sea otter cranium exhibits slower growth and

development of its morphology (Fig. 6) (Law et al. 2017). Because the cranium serves as the

underlying structure that supports the primary jaw muscle (i.e., the temporalis muscle) used to

generate bite force, the slow maturation of cranial size and shape directly contributes to the slow

maturation of biting performance (3.6 years of age for females and 5.0 years of age for males).

Specifically, broadening of the zygomatic arches, lengthening along the anteroposterior axis, and

development of a more pronounced sagittal crest all provide additional attachment sites for the

temporalis muscle (Scapino 1968; Turnbull 1970; Tanner et al. 2010; La Croix et al. 2011b; Law

and Mehta 2019; Law 2020; Stanchak et al. 2023). Cranial size maturity is not reached until 4.3

years of age for females and 3.3 years of age for males, cranial shape maturity is not reached

until 5.5 years of age for both sexes (Fig. 6; Table 1; Law et al. 2017), and correspondingly,

physiological cross-sectional area of the temporalis does not reach full maturity until 3.8 years of

age for females and 4.8 years of age for males (Fig. 4; Fig. 6; Table 1). In contrast, mandibular

size and shape reach full maturity during the subadult class between 1.5 to 2.8 years of age,

much earlier than the cranium (Fig. 6; Table 1; Law et al. 2017). Ontogenetic changes in

mandibular shape of southern sea otters are associated with increasing jaw adductor muscle in-

levers and the out-lever (Law et al. 2017); thus, these lever measurements along with mechanical

advantage of jaw closure also reach full maturity during the subadult class (Fig. 5; Fig. 6; Table

1). The mandible also serves as the underlying structure that supports the masseter muscle.

Unsurprisingly, physiological cross-sectional area of the masseter also reaches full maturity at

2.1 years of age for females and 3.3 years of age for males, on par with the mandibular

morphologies (Fig. 4; Table 1). The earlier maturation of mandibular morphology and

musculature in juvenile otters may contribute to positive allometric increases in bite force early

in ontogeny (Law et al. 2016b) and provide a boost in bite force during a period in which the

cranium remains smaller and less developed and the temporalis has not yet reached full

development. Similar patterns (i.e., earlier maturation of mechanical advantage and mandible

compared to the cranium) are also found in the bone-eating spotted hyena (Tanner et al. 2010),

providing evidence that adaptations to durophagy suggest fast maturation of jaw mechanics

while waiting for the rest the skull and jaw musculature to reach full size and shape.

The different functional roles of the cranium and mandible may also influence the

asynchrony between their growth and/or development. The cranium is the more complex

structure of the two, serving as a multifunctional structure that takes part in sensory functions,

respiration, and brain protection in addition to feeding, whereas the mandible is involved

primarily in feeding. Thus, the cranium may require slow, prolonged growth and/or development

of its morphologies to accommodate its multiple functions. Additional studies are needed to

reveal the maturation ages of growth and development among the cranium, brain, and other

sensory organs in sea otters and other mammals.

Although asynchrony has been quantified in only a handful of mammals, primarily

carnivorans (Tanner et al. 2010; La Croix et al. 2011b; Segura et al. 2013; Segura 2015; Law et

al. 2017), this phenomenon has also been uncovered at the macroevolutionary level. Decoupled

evolutionary modes between the cranium and mandible are found in various mammal clades

(McLean et al. 2018; Arbour et al. 2019; Michaud et al. 2020; Cassini and Toledo 2021; Law et

al. 2022, 2024a) despite strong integration between the cranium and mandible (Hautier et al.

2012; Figueirido et al. 2013; McLean et al. 2018; Michaud et al. 2020; Law et al. 2024a). A

possible explanation is that the cranium exhibits more structural, functional, or phylogenetic

constraints on its evolution, whereas the mandible may evolve more directly in response to

dietary changes. Across carnivorans, cranial shape follows clade-based evolutionary shifts,

suggesting that cranial diversity is phylogenetically constrained in carnivorans. In contrast,

mandibular shape evolution is linked to broad dietary regimes, suggesting greater evolutionary

lability of the mandible (Figueirido et al. 2011; Law et al. 2022, 2024a). Similarly, in bats,

sensory functions (i.e., echolocation and vision) and phylogenetic structure are the most

influential factors shaping cranial evolution, whereas diet has a stronger influence on mandibular

evolution (Arbour et al. 2019). The comparison between asynchronous growth/development at

the species level and decoupled evolution at the macroevolutionary level presents itself as an

exciting avenue of future investigation in the factors influencing cranial and mandibular

variation.

Effects of sexual dimorphism on biting performance growth

Like many species within the order Carnivora (Lindenfors et al. 2007; Law 2019), a

combination of sexual selection and niche divergence is hypothesized to drive sexual

dimorphism in sea otters in which males are bigger. Despite their larger size, male southern sea

otters displayed faster initial growth and developmental rates in skull size and some cranial shape

elements compared to females (Law et al. 2017). Presumably, reaching maturation of

craniomandibular morphology earlier enables males to reach maximum biting performance

faster, which may lead to competitive advantages in processing hard-shelled prey and in male–

male agonistic interactions. However, the growth curves of theoretical bite force provide a more

nuanced pattern that displayed both faster and slower growth (Fig. 3; Table 1). Within their first

two years, male otters reached the asymptotic value of theoretical bite force of females (437.7 N)

in just 1.8 years compared to 3.6 years for females. Because of their lower rate of growth decay,

subsequent growth of biting performance slows down and takes an additional >3 years for males

to reach their asymptotic value of theoretical bite force (620.7 N). Biphasic growth of biting

performance facilitated by lower rate of growth decay may be advantageous for males, which

must compete to establish reproductive territories, aggressively exclude other mature males, and

maintain exclusive access to estrous females (Ames et al. 1983; Jameson 1989; Ralls et al. 1996;

Pearson and Davis 2005). In the first phase, quickly reaching biting performances on par with

female bite forces enables males to compete with conspecifics in obtaining and processing the

energy required to further grow and develop. In the second phase, continual growth of biting

performance is slowed, likely because muscle tissue is energetically costly to grow (Millward et

al. 1976). Consistently, the mass of the temporalis muscles is the last element associated with the

feeding apparatus to reach full maturation (Fig. 4; Table 1). During this period, males do not

have territories and instead congregate in 'bachelor areas' where they take part in extensive

male–male interactions ranging from playful to aggressive that may be important for the

development of fighting skills and possibly even the establishment of social hierarchies (Jameson

1989; Lafferty and Tinker 2014). Because biting is the primary weapon used in territorial

conflicts between males, males that attain stronger biting performances faster may rise through

the hierarchical ranks and begin to venture out of bachelor groups to establish their own

reproductive territories. The ability to maintain these territories has lifetime reproductive

consequences as males that maintained their territories for the longest sired the most pups

(Tarjan and Tinker 2016). That full maturation of theoretical bite force and the onset of sexual

maturity and body size maturity occur at the same time at five years of age in males suggest

biting performance has an important influence on reproductive success.

Female otters, in contrast, exhibit a faster rate of growth decay, leading to weaker bite

forces compared to male otters. A possible explanation for this pattern is that selection for larger

bite forces is relaxed once female otters attained sufficient bite force to process hard prey as they

do not need stronger bites and larger body sizes for territorial conflicts. Rather than for growth,

energetic resources may be used to maintain their extremely high daily total energetic demands

associated with raising a pup (Thometz et al. 2014, 2016). Furthermore, these lower bite forces

may be sufficient to process hard-shelled prey because female otters tend to use tools more

frequently than male otters, providing them an alternative method to forage effectively (Law et

al. 2024b).

Comparisons of bite force growth to other carnivorans

Despite a large body of work examining the growth and development of the carnivoran

skull (Tanner et al. 2010; La Croix et al. 2011a; Segura and Prevosti 2012; Tarnawski et al.

2014; Forbes□Harper et al. 2017; Chatar et al. 2024), the growth of bite force and underlying

masticatory muscles is rarely studied in carnivorans. The most likely explanation is that

performance and muscular data are difficult to obtain in carnivorans, much less across an entire

ontogenetic series ranging from day old pups to years old adults. Thus, the few studies

examining performance across ontogeny have relied on osteological specimens to estimate biting

performance across ontogeny such as with calculations of mechanical advantage of jaw closing

or estimations of bite force using the dry skull method (Tanner et al. 2010; La Croix et al. 2011b;

Segura and Prevosti 2012; Forbes□Harper et al. 2017). However, as seen in previous work,

mechanical advantage reaches maturation ages much faster than other metrics of biting

performance and thus may be unreliable to assess performance maturation by itself (Tanner et al.

2010; La Croix et al. 2011b); this study). Similarly, bite forces estimated with the dry skull

method remain to be validated with empirical data (see *Caveats to bite force model* section

below). Dissection-based models may provide better insight on biting performance growth, but

muscular data remains difficult to gather across a full ontogenetic series in carnivorans. For

example, Law (2020) examined ontogenetic patterns of theoretical bite forces and jaw

musculature in American martens and fishers; however, this study was limited to just subadult

and adult age classes and was unable to generate growth curves. Lastly, there have been very few

published successes in measuring empirical bite forces in carnivorans using force transducers.

Binder and Van Valkenburgh's (2000) examination of spotted hyenas remains the only study to

successfully measure empirical bite force across ontogeny in a carnivoran. In contrast to sea

otters, spotted hyena bite forces continued to increase well past skull maturity; spotted hyena

skulls reached maturity around 3 years of age and their bite forces reached maturity later around

5–6.5 years of age (Binder and Van Valkenburgh 2000; Tanner et al. 2010), whereas sea otter

bite forces reached maturity 3.6–5 years of age (depending on sex) prior to skull maturation at

5.5 years of age (Fig. 6). Comparison of growth rates of the underlying masticatory muscles may

elucidate why bite force maturation ages differ between these two durophagous species, but this

data remains to be collected in hyenas. La Croix et al. (La Croix et al. 2011b) also attempted to

measure bite forces in coyotes but found that the juveniles were unmotivated biters, thus

highlighting the difficulties of measuring empirical bite force across ontogeny. Overall, this

study emphasizes the clear need for additional anatomical work to investigate ontogenetic

relationships among skull morphology, bite performance, and the underlying masticatory

musculature in carnivorans.

Caveats to bite force model

I acknowledge that my approach of using a dissection-based 2D model with single points for

each muscle force vector may be an oversimplification of bite force estimation. First, my model

does not account for how different degrees of gape can influence the length and orientation of

moment arms, which in turn can influence how the muscle force vectors are oriented. Increased

gape is typically linked to reduced bite force (Herring and Herring 1974; Dumont and Herrel

2003; Santana 2015), suggesting that my theoretical bite force estimations may be at the upper

limits. Second, the use of 3D methods may provide better estimations of bite force than 2D

models. In an analysis across various bat species, Davis et al. (2010) found that 3D distributed

traction models that distribute forces over each muscle's respective attachment region on the

skull is the best predictor of empirical bite force. Interestingly, the authors also found that 2D

models using estimated muscle PCSAs (i.e., Thomason's dry skull method (Thomason 1991)) do

a pretty good job of estimating empirical bite forces in bats, whereas theoretical bite forces

estimated using dissection-based 2D models that incorporate measured muscle PCSAs (as done

in this current study) tend to be the worst of the three models as it underestimated empirical bite

forces (Davis et al. 2010). These differences in bite force models may be restricted to bats, as

Law and Mehta (2019) found an opposite pattern in which bite forces estimated from

Thomason's dry skull method underestimated bite forces derived from 2D dissection-based

models. However, the dry skull method remains valuable in estimating muscle mass and PCSA

when gross muscle is unavailable, at least for carnivorans. Dickinson et al. ((Dickinson et al.

2021)) found that the dry skull method is the best technique to estimate muscle mass and PCSA

across carnivorans compared to other 2D or 3D approaches that estimate muscle attachment

areas. Therefore, a 3D distributed traction model may not vastly outperform the 2D dissection-

based model or Thomason's dry skull method in sea otters; nevertheless, how these three models

compare in sea otters requires additional investigation. Overall, the validity of bite force models

in sea otters remains tricky to elucidate without empirical bite force measurements from wild

southern sea otters. Unfortunately, direct measurements of bite forces are difficult to obtain from

mammals outside of small rodents and bats (Herrel et al. 2008; Santana et al. 2010; Becerra et al.

2014), and especially from mammals protected under the US Marine Mammal Protection Act

and the U.S. Endangered Species Act, as the author of this paper discovered after five years of

trying.

Conclusion

The ability to feed is paramount to survival; therefore, it is critical to understand the

growth and development of the feeding apparatus and its impact on the maturation of biting

performance. In this study, I found that southern sea otters exhibit slow maturation of theoretical

bite force due to slow growth and development of the cranium and the temporalis muscle.

Although slow bite force maturation may limit newly weaned juveniles from processing certain

hard-shelled prey and may thus impact their survival, tool using behavior may help mitigate

these limitations. In addition, faster growth and development of the mandible and mechanical

advantage may also provide a boost in biting performance during a period in which the cranium

and temporalis has not yet reached full development. Furthermore, I found that bite force

maturation ages differed between the sexes: female otters reached bite force maturation earlier

whereas male otters exhibit biphasic growth in bite force in which they quickly reach sufficient

bite forces needed to process prey early in life followed by a slower growth phase towards bite

force maturation that coincides with sexual maturity. These patterns suggest that male to male

competition for resources and mates exhibits a strong selective force in the growth and

development of skull form and function in male otters. Overall, this study demonstrates how the

analysis of anatomical data can provide insight on the foraging ecologies of sea otters across

ontogeny. Future work integrating the functional morphology of the feeding apparatus,

biomechanics of tool using behavior, and changes in prey characteristics will elucidate the

processes that improve feeding performances through ontogeny.

Acknowledgements

I am grateful to the staff of the California Department of Fish & Wildlife (Colleen

Young, Erin Dodd, and Francesca Batac) and California Academy of Sciences (Moe Flannery

and Sue Pemberton) for access to specimens; Lilian Carswell (US Fish and Wildlife Service) for

permitting logistics; and Tim Tinker (US Geological Survey) and Rita Mehta (University of

California Santa Cruz) for helpful discussions on sea otter life history. Funding was provided

partly by a Grant-in-Aid of Research from the American Society of Mammalogists, a Lerner-

Gray Fund through the American Museum of Natural History, a Rebecca and Steve Sooy

Graduate Research Fellowship for Marine Mammals, and the National Science Foundation (NSF

DBI-2128146).

Figures

Fig. 1. Frequency plots of sea otter specimens by estimated age (years).

498   **Fig. 2.** Diagram of the bite force model. Forces of the temporalis and masseter muscles (F_T

499   and F_M) are applied at given distances (MAT and MAM, respectively) from the TMJ (orange

500   circle), creating rotation of the lower jaw, and bite force at a given distance (OL) balances these

501   muscle forces: $BF = 2[(F_T \times MAT + F_M \times MAM)/OL]$.

510   **Fig. 3.** Growth curve for theoretical bite force. Red circles and growth curves represent female

511   otters and blue diamonds and growth curves represent male otters. Red and blue dashed lines

512   indicate the times at which traits reach 95% of their asymptotic values for females and males,

513   respectively.

Fig. 4. Growth curves for PCSA, mass, and fiber length of the temporalis and masseter muscles.

Red circles and growth curves represent female otters, and blue diamonds and growth curves

represent male otters. Red and blue dashed lines indicate the times at which traits reach 95% of

their asymptotic values for females and males, respectively.

Fig. 5. Growth curves for mechanical advantage of jaw closure from the temporalis and masseter

muscles. Red circles and growth curves represent female otters and blue diamonds and growth

curves represent male otters. Red and blue dashed lines indicate the times at which traits reach

95% of their asymptotic values for females and males, respectively.

Fig. 6. Timeline illustrating the maturation ages of southern sea otter skull morphology and theoretical bite force in relation to major life-history events. Maturation ages of cranial and mandibular size and shape were obtained from Law et al. (2017). Cranial and mandibular size and shape were calculated using geometric morphometrics. Maturation age was defined as the age at which the trait reached 95% of their asymptotic values. Estimated age (in years) are in circles. Imm = immature pup.

Table 1. Parameter estimates of growth curves generated with a Brody model. *A* is the
asymptotic maturity, *k* is the exponential decay of the growth rate, *B* is the scaling constant
signified by the ratio between the initial growth rate at t = 0 and *k*, and *L* is the initial growth rate
calculated as the product between *B* and *k*. 95% confidence intervals are given in parentheses.
Bold values indicate significant differences between male and female parameters. Age of
maturity indicates the ages (in years) at which element reach 95% of their respective asymptotic
values.
555

	sex	*A*	*B*	*k*	*L*	Age of maturity
bite force	F	**437.7 (410.1, 465.2) N**	0.88 (0.80, 0.96)	**0.80 (0.57, 1.02)**	0.70	3.60
	M	**620.7 (580.4, 660.9) N**	0.93 (0.84, 1.02)	**0.54 (0.37, 0.71)**	0.50	5.00
temporalis PCSA	F	**8.32 (7.83, 8.81)**	0.84 (0.77, 0.92)	**0.75 (0.54, 0.95)**	0.63	3.80
	M	**12.07 (11.13, 13.01)**	0.92 (0.82, 1.01)	**0.51 (0.32, 0.70)**	0.47	4.80
masseter PCSA	F	**2.36 (2.17, 2.55)**	0.83 (0.70, 0.96)	**1.15 (0.67, 1.62)**	0.95	2.10
	M	**3.08 (2.88, 3.27)**	0.92 (0.81, 1.03)	0.82 (0.52, 1.12)	0.75	3.30
temporalis mass	F	**36.77 (34.25, 39.29)**	0.93 (0.85, 1.01)	**0.64 (0.46, 0.82)**	0.60	4.20
	M	**63.47 (55.91, 71.04)**	0.98 (0.87, 1.09)	**0.38 (0.21, 0.54)**	0.37	5.80
masseter mass	F	**8.34 (7.77, 8.92)**	0.90 (0.81, 1.00)	0.85 (0.59, 1.11)	0.77	3.20
	M	**10.85 (10.14, 11.56)**	0.99 (0.88, 1.09)	0.70 (0.46, 0.94)	0.69	3.90
temporalis length	F	**4.07 (3.83, 4.30)**	0.53 (0.42, 0.64)	**1.76 (0.81, 2.70)**	0.93	1.30
	M	**4.74 (4.50, 4.97)**	0.55 (0.47, 0.62)	**0.69 (0.38, 1.00)**	0.38	3.50
masseter length	F	3.29 (3.12, 3.45)	0.43 (0.35, 0.52)	1.21 (0.60, 1.82)	0.52	1.90
	M	3.33 (3.15, 3.50)	**0.53 (0.42, 0.64)**	1.30 (0.53, 2.08)	0.69	1.90
temporalis MA	F	0.73 (0.70, 0.75)	**0.29 (0.21, 0.36)**	3.32 (1.15, 5.50)	0.96	0.90
	M	0.72 (0.71, 0.74)	**0.14 (0.07, 020)**	3.72 (-2.09, 9.54)	0.52	0.60
masseter MA	F	0.54 (0.52, 0.56)	0.42 (0.31, 0.53)	4.70 (1.69, 7.72)	1.97	0.50
	M	0.52 (0.50, 0.54)	0.40 (0.29, 0.50)	3.66 (0.55, 6.77)	1.46	0.70
temporalis in-lever	F	**3.07 (2.97, 3.16)**	0.56 (0.51, 0.62)	1.36 (1.03, 1.70)	0.76	2.30
	M	**3.22 (3.13, 3.31)**	0.59 (0.53, 0.64)	1.07 (0.77, 1.36)	0.63	3.10
masseter in-lever	F	2.22 (2.13, 2.31)	0.64 (0.56, 0.74)	**2.21 (1.44, 2.98)**	1.41	1.60
	M	2.29 (2.21, 2.38)	0.71 (0.63, 0.78)	**1.27 (0.88, 1.66)**	0.90	2.20
out-lever	F	**4.23 (4.04, 4.42)**	**0.39 (0.32, 0.47)**	1.09 (0.56, 1.61)	0.43	2.10
	M	**4.43 (4.33, 4.54)**	**0.52 (0.48, 0.57)**	1.03 (0.76, 1.30)	0.54	3.20

559   **References**
560
561

Ames, J. A., R. A. Hardy, and F. E. Wendell. 1983. Tagging materials and methods for sea
otters, Enhydra lutris. California Fish and Game 69:243–252.

Arbour, J. H., A. A. Curtis, and S. E. Santana. 2019. Signatures of echolocation and dietary
ecology in the adaptive evolution of skull shape in bats. Nat Commun 10:2036.

Baliga, V. B., and R. S. Mehta. 2014. Scaling patterns inform ontogenetic transitions away from
cleaning in Thalassoma wrasses. J. Exp. Biol. 217:3597–3606.

Becerra, F., A. I. Echeverría, A. Casinos, and A. I. Vassallo. 2014. Another one bites the dust:
Bite force and ecology in three caviomorph rodents (Rodentia, Hystricognathi). J. Exp. Zool.
321:220–232.

Biewener, A. A., and R. J. Full. 1992. Force platform and kinematic analysis. Pp. 45–73 *in* A. A.
Biewener, ed. Biomechanics Structures and Systems. A Practical Approach. Biomechanics
Structures and Systems. A Practical Approach.

Binder, W. J., and B. Van Valkenburgh. 2000. Development of bite strength and feeding
behaviour in juvenile spotted hyenas (Crocuta crocuta). J Zool 252:273–283.

Brody, S. 1945. Chapter 16 Time relations of growth of individuals and populations. Pp. 484–
574 *in* Bioenergetics and growth. Hafner Publishing Company, Inc., Neew York, NY.

Cassini, G. H., and N. Toledo. 2021. An Ecomorphological Approach to Craniomandibular
Integration in Neotropical Deer. J Mamm Evol 28:111–123.

Chatar, N., R. Boman, V. Fischer, V. Segura, C. Julémont, and Z. J. Tseng. 2024. Growing
sabers: Mandibular shape and biomechanical performance trajectories during the ontogeny of
Smilodon fatalis. Anat. Rec., doi: 10.1002/ar.25504.

Christiansen, P., and S. Wroe. 2007. Bite forces and evolutionary adaptations to feeding ecology
in carnivores. Ecology 88:347–358.

Constantino, P. J., J. J. W. Lee, D. Morris, P. W. Lucas, A. Hartstone-Rose, W.-K. Lee, N. J.
Dominy, A. Cunningham, M. Wagner, and B. R. Lawn. 2011. Adaptation to hard-object
feeding in sea otters and hominins. J. Hum. Evol. 61:89–96.

Costa, D. P., and G. L. Kooyman. 1984. Contribution of specific dynamic action to heat-balance
and thermoregulation in the sea otter Enhydra lutris. Physiol. Zool. 57:199–203.

Costa, D. P., and G. L. Kooyman. 1982. Oxygen consumption, thermoregulation, and the effect
of fur oiling and washing on the sea otter, Enhydra lutris. Can. J. Zool. 60:2761–2767.

Davis, J. L., S. E. Santana, E. R. Dumont, and I. R. Grosse. 2010. Predicting bite force in
mammals: two-dimensional versus three-dimensional lever models. J. Exp. Biol. 213:1844–
1851.

Dickinson, E., J. S. Davis, A. R. Deutsch, D. Patel, A. Nijhawan, M. Patel, A. Blume, J. L.
Gannon, C. M. Turcotte, C. S. Walker, and A. Hartstone-Rose. 2021. Evaluating bony
predictors of bite force across the order Carnivora. J Morpho 282:1499–1513.

Dumont, E. R., and A. Herrel. 2003. The effects of gape angle and bite point on bite force in
bats. J. Exp. Biol. 206:2117–2123.

Estes, J. A., K. E. Underwood, and M. J. Karmann. 1986. Activity-time budgets of sea otters in
California. J. Wildl. Manage. 50:626–636.

Figueirido, B., N. MacLeod, J. Krieger, M. D. Renzi, J. A. Pérez-Claros, and P. Palmqvist. 2011.
Constraint and adaptation in the evolution of carnivoran skull shape. Paleobiology 37:490–
518.

Figueirido, B., Z. J. Tseng, and A. Martín-Serra. 2013. Skull shape evolution in durophagous
carnivorans. Evolution 67:1975–1993.

Figueirido, B., Z. J. Tseng, F. J. Serrano-Alarcón, A. Martín-Serra, and J. F. Pastor. 2014. Three-
dimensional computer simulations of feeding behaviour in red and giant pandas relate skull
biomechanics with dietary niche partitioning. Biol. Lett. 10:20140196–20140196.

Forbes☐Harper, J. L., H. M. Crawford, S. J. Dundas, N. M. Warburton, P. J. Adams, P. W.
Bateman, M. C. Calver, and P. A. Fleming. 2017. Diet and bite force in red foxes: ontogenetic
and sex differences in an invasive carnivore. J. Zoo☐l. 303:54–63.

Fujii, J. A., K. Ralls, and M. T. Tinker. 2017. Food abundance, prey morphology, and diet
specialization influence individual sea otter tool use. Behav Ecol 28:1206–1216.

Fujii, J. A., K. Ralls, M. T. Tinker, and M. T. Tinker. 2015. Ecological drivers of variation in
tool-use frequency across sea otter populations. Behav Ecol 26:519–526.

Gignac, P. M., and G. M. Erickson. 2014. Ontogenetic changes in dental form and tooth
pressures facilitate developmental niche shifts in American alligators. J Zool 295:132–142.

Gignac, P. M., and S. E. Santana. 2016. A bigger picture: organismal function at the nexus of
development, ecology, and evolution: an introduction to the symposium. Integr Comp Biol
56:369–372.

Hautier, L., R. Lebrun, and P. G. Cox. 2012. Patterns of Covariation in the Masticatory
Apparatus of Hystricognathous Rodents: Implications for Evolution and Diversification. J
Morpho 273:1319–1337.

Herrel, A., and A. C. Gibb. 2006. Ontogeny of Performance in Vertebrates*. Physiol Biochem
Zool 79:1–6.

Herrel, A., and J. C. O'Reilly. 2006. Ontogenetic scaling of bite force in lizards and turtles.
Physiol Biochem Zool 79:31–42.

Herrel, A., A. D. Smet, L. F. Aguirre, and P. Aerts. 2008. Morphological and mechanical
determinants of bite force in bats: do muscles matter? J Exp Biol 211:86–91.

Herring, S. W., and S. E. Herring. 1974. The Superficial Masseter and Gape in Mammals. Am.
Nat. 108:561–576.

Higgins, B. A., C. J. Law, and R. S. Mehta. 2018. Eat whole and less often: ontogenetic shift
reveals size specialization on kelp bass by the California moray eel, Gymnothorax mordax.
Oecologia 101:1–13.

Jameson, R. J. 1989. Movements, home range and territories of male sea otters off central
California. Mar Mammal Sci 5:159–172.

Jameson, R. J., and A. M. Johnson. 1993. Reproductive characteristics of female sea otters. Mar
Mammal Sci 9:156–167.

La Croix, S., K. E. Holekamp, J. A. Shivik, B. L. Lundrigan, and M. L. Zelditch. 2011a.
Ontogenetic relationships between cranium and mandible in coyotes and hyenas. J Morpho
272:662–674.

La Croix, S., M. L. Zelditch, J. A. Shivik, B. L. Lundrigan, and K. E. Holekamp. 2011b.
Ontogeny of feeding performance and biomechanics in coyotes. J Zool 285:301–315.

Lafferty, K. D., and M. T. Tinker. 2014. Sea otters are recolonizing southern California in fits
and starts. Ecosphere 5:1–11.

Law, C. J. 2020. Sex☐specific ontogenetic patterns of cranial morphology, theoretical bite force,
and underlying jaw musculature in fishers and American martens. J Anat 237:727–740.

Law, C. J. 2019. Solitary meat-eaters: solitary, carnivorous carnivorans exhibit the highest
degree of sexual size dimorphism. Sci Rep-uk 9:15344.

Law, C. J., V. B. Baliga, M. T. Tinker, and R. S. Mehta. 2017. Asynchrony in craniomandibular
development and growth in Enhydra lutris nereis (Carnivora: Mustelidae): are southern sea
otters born to bite? Biol J Linn Soc 121:420–438.

Law, C. J., E. A. Blackwell, A. A. Curtis, E. Dickinson, A. Hartstone☐Rose, and S. E. Santana.
2022. Decoupled evolution of the cranium and mandible in carnivoran mammals. Evolution
76:2959–2974.

Law, C. J., L. J. Hlusko, and Z. J. Tseng. 2024a. Uncovering the mosaic evolution of the
carnivoran skeletal system. Biol. Lett. 20:20230526.

Law, C. J., and R. S. Mehta. 2019. Dry versus wet and gross: Comparisons between the dry skull
method and gross dissection in estimations of jaw muscle cross☐sectional area and bite forces
in sea otters. J Morphol 280:1706–1713.

Law, C. J., M. T. Tinker, J. A. Fujii, T. Nicholson, M. Staedler, J. A. Tomoleoni, C. Young, and
R. S. Mehta. 2024b. Tool use increases mechanical foraging success and tooth health in
southern sea otters (Enhydra lutris nereis). Science 384:798–802.

Law, C. J., V. Venkatram, and R. S. Mehta. 2016a. Sexual dimorphism in craniomandibular
morphology of southern sea otters (Enhydra lutris nereis). J Mammal 97:1764–1773.

Law, C. J., C. Young, and R. S. Mehta. 2016b. Ontogenetic scaling of theoretical bite force in
southern sea otters (Enhydra lutris nereis). Physiol Biochem Zool 89:347–363.

Layman, C. A., S. D. Newsome, and T. G. Crawford. 2015. Individual-level niche specialization
within populations: emerging areas of study. Oecologia 178:1–4.

Lindenfors, P., J. L. Gittleman, and K. E. Jones. 2007. Sexual size dimorphism in mammals. Pp.
16–26 *in* Sex, size and gender roles: evolutionary studies of sexual size dimorphism.

McLean, B. S., K. M. Helgen, H. T. Goodwin, and J. A. Cook. 2018. Trait-specific processes of
convergence and conservatism shape ecomorphological evolution in ground-dwelling
squirrels. Evolution 72:473–489.

Mendez, J., and A. Keys. 1960. Density and composition of mammalian muscle. Metab. Clin.
Exp. 9:184–188.

Michaud, M., G. Veron, and A. Fabre. 2020. Phenotypic integration in feliform carnivores:
Covariation patterns and disparity in hypercarnivores versus generalists. Evolution 74:2681–
2702.

Millward, D. J., P. J. Garlick, and P. J. Reeds. 1976. The energy cost of growth. Proc. Nutr. Soc.
35:339–349.

Monson, D. H., J. A. Estes, J. L. Bodkin, and D. B. Siniff. 2000. Life history plasticity and
population regulation in sea otters. Oikos 90:457–468.

Morrison, P., M. Rosenmann, and J. A. Estes. 1974. Metabolism and thermoregulation in the sea
otter. Physiol. Zool. 47:218-229.

Nicholson, T. E., K. A. Mayer, M. M. Staedler, T. O. Gagné, M. J. Murray, M. A. Young, J. A.
Tomoleoni, M. T. Tinker, and K. S. V. Houtan. 2020. Robust age estimation of southern sea
otters from multiple morphometrics. Ecol Evol 10:8592–8609.

Payne, S. F., and R. J. Jameson. 1984. Early behavioral development of the sea otter, Enhydra
lutris. J Mammal 65:527–531.

Pearson, H. C., and R. W. Davis. 2005. Behavior of territorial male sea otters Enhydra lutris in
Prince William Sound, Alaska. aquatic mammals 31:226–233.

Ralls, K., T. C. Eagle, and D. B. Siniff. 1996. Movement and spatial use patterns of California
sea otters. Can J Zool 74:1841–1849.

Ralls, K., and D. B. Siniff. 1990. Time budgets and activity patterns in California sea otters. J.
Wildl. Manage. 54:251–259.

R Core Team. 2023. R: A language and environment for statistical computing.

Riedman, M., and J. A. Estes. 1990. The sea otter (Enhydra lutris): behavior, ecology, and
natural history. Biological Report 90:1–117.

Riley, M. A. 1985. An analysis of masticatory form and function in three mustelids (Martes
americana, Lutra canadensis, Enhydra lutris). J Mammal 66:519–528.

Sacks, R. D., R. D. Sacks, R. R. Roy, and R. R. Roy. 1982. Architecture of the hind limb
muscles of cats: Functional significance. J Morpho 173:185–195.

Santana, S. E. 2015. Quantifying the effect of gape and morphology on bite force: biomechanical
modelling and in vivomeasurements in bats. Funct. Ecol. 30:557–565.

Santana, S. E., E. R. Dumont, E. Dumont, and J. L. Davis. 2010. Mechanics of bite force
production and its relationship to diet in bats. Funct. Ecol. 24:776–784.

Santana, S. E., and K. E. Miller. 2016. Extreme postnatal scaling in bat feeding performance: a
view of ecomorphology from ontogenetic and macroevolutionary perspectives. Integr Comp
Biol icw075-10.

Scapino, R. P. 1968. Biomechanics of Feeding in Carnivora.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of
image analysis. Nat Meth 9:671–675.

Segura, V. 2015. A three-dimensional skull ontogeny in the bobcat (Lynx rufus) (Carnivora:
Felidae): a comparison with other carnivores. Canadian Journal of Zoology 93:225–237.

Segura, V., and F. Prevosti. 2012. A quantitative approach to the cranial ontogeny of Lycalopex
culpaeus (Carnivora: Canidae). Zoomorphology 131:79–92.

Segura, V., F. Prevosti, and G. Cassini. 2013. Cranial ontogeny in the Puma lineage, Puma
concolor, Herpailurus yagouaroundi, and Acinonyx jubatus(Carnivora: Felidae): a three-
dimensional geometric morphometric approach. Zool J Linn Soc 169:235–250.

Staedler, M. M. 2011. Maternal care and provisioning in the southern sea otter (Enhydra lutris
nereis): reproductive consequences of diet specialization in an apex predator.

Stanchak, K. E., P. A. Faure, and S. E. Santana. 2023. Ontogeny of cranial musculoskeletal
anatomy and its relationship to allometric increase in bite force in an insectivorous bat
(Eptesicus fuscus). Anat. Rec. 306:2842–2852.

Tanner, J. B., M. L. Zelditch, and B. L. Lundrigan. 2010. Ontogenetic change in skull
morphology and mechanical advantage in the spotted hyena (Crocuta crocuta). J Morpho
271:353–365.

Tarjan, L. M., and M. T. Tinker. 2016. Permissible home range estimation (PHRE) in restricted
habitats: a new algorithm and an evaluation for sea otters. PLoS ONE 11:e0150547-20.

Tarnawski, B. A., G. H. Cassini, and D. A. Flores. 2014. Skull allometry and sexual dimorphism
in the ontogeny of the southern elephant seal (Miroungaleonina leonina). Can. J. Zool. 92:19–
31.

Thomason, J. J. 1991. Cranial strength in relation to estimated biting forces in some mammals.
Can J Zool 69:2326–2333.

Thometz, N. M., T. L. Kendall, B. P. Richter, and T. M. Williams. 2016. The high cost of
reproduction in sea otters necessitates unique physiological adaptations. J. Exp. Biol.
219:2260–2264.

Thometz, N. M., M. T. Tinker, M. T. Tinker, M. M. Staedler, M. Staedler, K. A. Mayer, T. M.
Williams, and T. M. Williams. 2014. Energetic demands of immature sea otters from birth to
weaning: implications for maternal costs, reproductive behavior and population-level trends. J
Exp Biol 217:2053–2061.

Timm-Davis, L. L., T. J. DeWitt, and C. D. Marshall. 2015. Divergent skull morphology
supports two trophic specializations in otters (Lutrinae). PLoS ONE 10:e0143236-18.

Tinker, M. T., G. Bentall, and J. A. Estes. 2008. Food limitation leads to behavioral
diversification and dietary specialization in sea otters. Proc Natl Acad Sci USA 105:560–565.

Tinker, M. T., D. F. Doak, J. A. Estes, B. B. Hatfield, M. M. Staedler, and J. L. Bodkin. 2006.
Incorporating diverse data and realistic complexity into demographic estimation procedures
for sea otters. Ecological Applications 16:2293–2312.

Tinker, M. T., and B. B. Hatfield. 2015. Southwest U.S. Southern sea otter annual range-wide
census results: U.S. Geological Survey data release.

Tseng, Z. J., D. F. Su, X. Wang, S. C. White, and X. Ji. 2017. Feeding capability in the extinct
giant Siamogale melilutra and comparative mandibular biomechanics of living Lutrinae.
Scientific Reports 7:1–10.

Turnbull, W. D. 1970. Mammalian masticatory apparatus.

Yeates, L. C., T. M. Williams, and T. L. Fink. 2007. Diving and foraging energetics of the
smallest marine mammal, the sea otter (Enhydra lutris). J Exp Biol 210:1960–1970.

Zelditch, M. L., B. L. Lundrigan, H. D. Sheets, and T. Garland. 2003. Do precocial mammals
develop at a faster rate? A comparison of rates of skull development in Sigmodon fulviventer
and Mus musculus domesticus. Journal of Evolutionary Biology 16:708–720.

Ziscovici, C., P. W. Lucas, P. J. Constantino, T. G. Bromage, and A. van Casteren. 2014. Sea
otter dental enamel is highly resistant to chipping due to its microstructure. Biol. Lett.
10:20140484–20140484.

765