Intra-individual variability in the effects of transcranial direct current stimulation on free choice saccade behaviour

Brandon Caie[1] and Gunnar Blohm[1]

[1]Centre for Neuroscience Studies, Queen's University

Abstract

Transcranial direct current stimulation (tDCS) is used as a tool to causally influence neural activity in humans non-invasively. Although most studies recruit a large number of participants in order to uncover population-level effects, growing evidence suggests that tDCS may be expected to induce different effects in different individuals, leading to large inter-individual variability and confounds in population-level testing. Alternatively, this variability may arise from intra-individual sources that are difficult to assess in standard designs. Here, we performed between 8 and 10 sessions of tDCS within individuals to understand how intra-individual variability impacts the inference of tDCS effects. We recruited 5 participants who underwent functional MRI to localize the human frontal eye field (FEF) homologue. An HD-tDCS montage was used to stimulate the target location based on individual MRI localizations, alternating the polarity between anodal or cathodal current over 8-10 repeated sessions during a 5 week period. Participants performed a free choice task before and after stimulation while recording EEG activity. We then developed a difference-in-difference method based on permutation testing to assess the likelihood of a causal effect of tDCS at different levels of abstraction: group-level, inter-individual, and intra-individual. At the group-level, we found evidence for an influence of tDCS on choice reaction times, which followed a reaction-time dependent change in alpha-band activity, and on how choices depended on recent history. However, individuals showed heterogeneous, and often contradictory, effects. We then analyzed the distribution of session permutations at the intra-individual level, and found a discrepancy between the inter-individual effects that survived significance testing and the intra-individual effects that correlated on a session-session basis. We argue that, while the

observed variability may have arisen from a combination of inter and intra-individual differences relevant to tDCS-dependent mechanisms of action, it may be equally well explained by spurious effects arising from history-dependence between repeated measures that are typically assumed to be independent. In light of this, we assess the counterfactuals that must be evaluated in order to make data-driven inferences about the causal effects of tDCS on free choice behaviour and its neural correlates.

Introduction

It has long been known that externally applied electric fields over the cortical surface can influence neural activity (Bindman et al., 1964; Purpura and McMurty, 1965). Transcranial direct current stimulation (tDCS) was subsequently developed as a method to causally influence neural activity non-invasively (Merton and Morton, 1980; Rothwell, 2018). tDCS has several advantages: it is non-invasive, cost-effective, well-tolerated and easy to use. Early application of tDCS to human study suggested that cortical activity could be modulated according to the polarity of the applied current (Nitsche and Paulus, 2000, 2001; Pellicciari et al., 2013; Priori et al., 1998). While this finding generated much interest in the applications of directly intervening on neural activity in humans, future studies using tDCS were marred by confounding sources of variability and failures to replicate findings (Horvath et al., 2014; B. Krause and Cohen Kadosh, 2014).

The use of tDCS to influence human neural activity was motivated by studies in animal models, where slice preparations or invasive in vivo measurements isolated mechanisms of action at a resolution inaccessible to human study (Bindman et al., 1964; Pelletier and Cicchetti, 2014; Purpura and McMurty, 1965). It is now generally supported that anodal stimulation can increase, while cathodal stimulation can decrease, neural firing rate through a net polarization along a somatodendritic axis (Kabakov et al., 2012; Liu et al., 2018), although it is unlikely that this is the sole physiological mechanism at play in vivo (Lefaucheur and Wendling, 2019; Rahman et al., 2013; Stagg and Nitsche, 2011). Thus, studies using tDCS were often motivated with this mechanism of action as an *a priori* that could be used to dissociate experimental or clinical hypotheses (Jackson et al., 2016; Lefaucheur and Wendling, 2019). However, computational models of current flow have shown that individual anatomical differences – such as skull thickness and composition, CSF volume, and gyri-sulcal morphology – can lead to different electric field distributions across individuals when the 'same' brain region is targeted (Bikson et al., 2012; Datta et al., 2012; Mikkonen et al., 2020; Opitz et al., 2015; Rawji et al., 2018; Truong et al., 2013). Given these known sources of variability, even if tDCS could be reduced to a singular mechanism of action (sometimes called the 'somatic doctrine (Jackson et al., 2016)), predictions about the physiological outcome in absence of detailed knowledge of how individual neuroanatomy relates to the stimulation protocol may be expected to yield mixed and/or contradictory effects at the population level (Filmer et al., 2019; Vergallito et al., 2022).

With the recognition that tDCS may not predict consistent effects across individuals comes the need for causal assessments at the individual level. However, this introduces methodological challenges. For

one, individualized stimulation protocols that predict current distributions across the cortical tissue require imaging technology that negates the cost-effectiveness of tDCS. But perhaps more fundamental to the experimentalist, where effects are often *inferred* through statistical changes to behaviour and/or neural measurements, is the introduction of statistical dependencies that can be abstracted over when testing hypotheses at the population-level (Caie and Blohm, 2024; Molenaar, 2004; Molenaar and Campbell, 2009). While population-level studies can treat sessions from different individuals as statistically independent within reason, individual assessments collect repeated measurements that come from an obvious common source. While it is well known that many behaviours and neural processes within individuals are history-dependent(Caie et al., 2023; Chaudhuri et al., 2015; Codling et al., 2008; Deco and Romo, 2008; Fischer and Whitney, 2014; Linkenkaer-Hansen et al., 2001; Medaglia et al., 2011; Murray et al., 2014) – that is to say, they are statistically dependent on past measurements – the consequences of this mixture of different sources of intra-individual variability on the ability to infer causal effects from repeated measures (cf Caie and Blohm, 2024; Granger and Newbold, 1974; Macke and Nienborg, 2019) has not been considered as a source of variability in the interpretation of individual responses to tDCS.

Here, we report the results of 10 repeated sessions of a HD-tDCS experiment testing the effect of an externally applied current over the right frontal eye fields (rFEF) in human participants performing a free choice saccade task (Caie et al., 2023; Noudoost and Moore, 2011; Soltani et al., 2013). Individual fMRI localizations of the human homologue of the right frontal eye field (rFEF) were performed prior to the delivery of twice-weekly alternating sessions of anodal or cathodal current using high-definition transcranial direct current stimulation (HD-tDCS) for 5 participants over a 5 week period. Using a data-driven statistical method based on permutation testing, we assessed the likelihood that pre-post differences to psychometric measures and EEG recordings were influenced by the polarity of the stimulation at different levels of analysis: group, inter-individual, and intra-individual. At each level, we found evidence for causal effects of tDCS on choice psychometrics, and corresponding changes in EEG activity. However, at both the group and inter-individual level, the variance in its constituent repeated measures was large and often contradictory. We argue that this is sufficient to call into question the validity of either group or individual-level testing as a privileged level of analysis for the influence of tDCS on repeated measures where statistical independence cannot be assumed *a priori*. We then discuss the counterfactuals that should be evaluated in order to make inferences about the influence of tDCS on choice behaviour and its neural correlates at the individual level.

Methods

Participants were recruited to undergo an fMRI-guided HD-tDCS study on the role of the frontal eye field in free choice saccade behaviour. Prior to the stimulation sessions, a structural and functional MRI was obtained from each participant in order to localize a region of the fronto-parietal cortex active during saccadic planning and execution. Then, an individualized stimulation setup was designed using 3D models of the neuroanatomy, which in turn guided electrode placement prior to each session. Participants performed 10 sessions of a free choice saccade task over an approximately 5 week period, in a pre/post design alternating between anodal and cathodal stimulation.

Participants

Five participants took part in the study. All had normal or corrected-to-normal vision. The participants gave written consent prior to undergoing the procedure. The study was conducted with an experimental protocol approved by the Queen's University General Research Ethics Board, which adheres to the principles of the Canadian Tri-council Policy Statement on Ethical Conduct for Research Involving Humans and the principles of the Declaration of Helsinki (2013).

Free Choice Task

The participants began by fixating on a central target (*Fig 1A*) projected onto a screen (20" Mitsubishi Diamond Pro [16x12inches] 1280x1024 pixels, 60Hz, contrast 75.2%, brightness 0%). Two targets spaced at 6 degrees visual angle from central fixation in each visual field, were then presented with randomly assigned temporal asynchrony values (0 ms, 16ms, 33ms, 66ms, 99ms) and onset times drawn from a uniform distribution between 750 and 1250ms. The participants were instructed to look at either target, so long as it was done as fast as possible, without anticipating. The participants were informed prior to the beginning of the experiment that the targets would come up at different times relative to each other. The participants were instructed to blink in between trials in order to minimize disruption of stimulus perception. The script for the task was written in MATLAB using Psychtoolbox.

Behavioural data acquisition and filtering

During each trial, eye movements were measured via an EyeLink 1000 Tower Mount (SR Research, Mississauga, Canada) with a 1000Hz infrared camera that tracked retinal position. Retinal position was calibrated prior to each session. The EyeLink apparatus was placed 60cm from the screen containing the saccade targets. Saccades were calculated offline using a saccade detection algorithm with a velocity criterion of 50°/s, and were individually verified through a custom visual marking program. Trials where the tracker lost the eye (either through tracking error or participant lapse) were excluded. Trials were also excluded if the eye position deviated more than 15 degrees from the horizontal. The first saccade following target onset, or the first saccade within 3 degrees of the future target position prior to target onset, were taken to calculate the saccadic reaction time and endpoint error. Saccades with a positive endpoint horizontal position were given a choice value of 1, and saccades with a negative endpoint horizontal position a choice value of 0.

fMRI-guided HD-tDCS

To target the site of stimulation, we localized the right frontal eye field using a center-out saccade task during a functional MRI scan. Structural MRI was first acquired on a 3-Tesla Magnetom Trio scanner (Siemens Medical Systems, Erlangen, Germany). Participants rested supine in the scanner and wore a 32-channel head coil. At the beginning of the localizer session, a 176 slice, high resolution T1- weighted anatomic reference volume was acquired using a 3D MP-RAGE sequence (single shot, ascending sequence in the sagittal plane with TR = 1760 ms, TE = 2.2 ms, FoV = 256 mm, flip angle = 9°, and voxel size = 1 mm3). Participants performed center-out saccades, where targets were organized in a ring around a center fixation point. Participants were then tasked to perform saccades from the center point to each target in clockwise succession, pausing for approximately 1 second between each saccade.

Individual structural MRI scans were registered to Talaraich space by manually finding the anterior commisure and translating the cerebrum into AC-PC space. Functional MRI scans taken during the center-out saccade task were collected using a gradient echo planar imaging sequence, and pre-processed using a sinc slice scan time correction interpolation and high-pass filtering before co-registration to the anatomical scan with an initial and fine alignment in Brainvoyager QX. A general linear model was fit to the aligned BOLD activation map to find the voxel most correlated with the saccade period in the vacinity of the pre-central sulcus (Vernet et al., 2014).

HD-tDCS was chosen to maximize current delivery at the target location while minimizing current spread to other brain regions (citation). Briefly, HD-tDCS is a method to improve focalization of current by positioning 4 return electrodes in a concentric ring close to a central electrode. The position of the central electrode was chosen individually for each participant by finding the position on the scalp closest to the voxel chosen as the right FEF using Brainsight (Rogue Research Inc, Montreal Quebec). This mapping was done by performing a 3D reconstruction of the surface of the scalp, and coregistering it to the functional MRI data. A skin reconstruction was performed to identify anatomical landmarks for performing the subject-image registration. Then, a 3D brain reconstruction was overlayed to select a point on the scalp corresponding to the FEF ROI. Prior to each session, the central electrode position on the scalp was then found using a stereotactic neuronavigation system. Participants wore an optical tracking device fixed over the head that was tracked by a position sensory camera located overhead. The position on the scalp corresponding with the target location was then found using an infrared pointer that mapped the position of the pointer to the position on the scalp corresponding to the 3D model .

This scalp position was then recorded with a marker, then a custom HD-tDCS cap was fitted with the central electrode placed over the marked scalp location. The cap was designed so that the central electrode would be surrounded by four pickup electrodes at a radius of 4cm. To lower impedance, hair was dried and separated under the scalp using cotton swabs. Electrodes were 1.5cm diameter Ag/AgCl ring electrodes (UltraStim, Axelgaard Manufacturing, Denmark). Once the center and surround electrodes were placed in the plastic holder, the holder was filled with conducting gel and the stimulation cap was fixed with a chin velcro strap. Continuous direct current was then generated during the stimulation period using a microprocessor-controlled constant current source (neuroConn DC-STIMULATOR MC, Brainbox) . Current was delivered at 2mA for 20 minutes, with a 30 second ramp on and ramp off period.

Current Modelling

Individualized modelling of the resulting electric field induced by HD-tDCS over the target site was performed in order to predict the strength and orientation of the induced electric field at the target ROI across participants. A tetrahedral head mesh was constructed using the T1-weighted structural MR images using *MRI2Mesh* in SimNIBS (Saturnino et al., 2015). The position of the central electrode was then transformed from BrainVoyager Internal Coordinates to World Coordinates in the SimNIBS GUI. The resulting mesh was

used to generate a 4cm ring in the tetrahedral mesh from which the surround electrode positions were selected, then projected on to the scalp in the SimNibs GUI. Electric field simulations were then performed for each subject individually. The central electrode was set to 2mA current, and each surround electrode -.5mA for the anodal current simulation. Cathodal electric fields were taken as the negative sign of the anodal simulations. Electrode parameters in SimNIBS were set to elliptical 1.5cm x 1.5cm, with type "Electrode plus Gel" and electrode thickness of 5mm. Simulation results were visualized using Gmsh 4. Tissue conductivity values were taken from the standard values in SimNibs (Wagner, 2004, Opitz, 2015). A table of the central and surround electrode positions in SimNIBS World Coordinates are shown for each participant in table xx.

Session Design

Each individual session was organized into a pre-stimulation period, where EEG and behavioural measures were taken simultaneously, followed by a 20 minute stimulation period where either a net anodal or cathodal current was introduced to the site of the central electrode, and finally a post-stimulation EEG-behavioural recording. In each of the pre/post periods, 5 blocks of 90 trials of the free choice task were recorded, lasting about 20 minutes and totalling 450 trials per session per condition. Although it was initially planned to standardize the timing of each session across a 5-week period, in order to accommodate participant's schedules, there was some variability in the timing and number of sessions performed. Sessions were scheduled for twice-weekly, with alternating anodal and cathodal stimulation. **3(double check)** sessions were excluded due to a trigger failure aligning the EEG data to the task timestamps, or due to excessive noise in the EEG signal.

EEG Pre-Processing

EEG data was recorded in the pre and post stimulation periods using the same center-surround electrode setup as for current delivery. EEG data was recorded using a V-Amp 16 channel EEG amplifier (Brain Products, Gilching, Germany) at a sampling frequency of 512 Hz, acquired using a custom protocol in OpenVibe. Reference and ground electrodes were placed on the back of the neck and the right elbow respectively. For each block of trials, EEG channels were independently notch-filtered to remove 60 Hz powerline noise, and then bandpass-filtered between 5 and 30 Hz. The current source density at the central electrode was then estimated using the surface laplacian spatial filter of the central and surround electrodes.

Time-frequency responses of the surface laplacian were computed for each trial individually. EEG data

from each trial was aligned to the onset of the fixation cross using a trigger protocol in OpenVibe. The surface laplacian was first detrended to remove slow drifts. Then, a continuous wavelet transform was performed across 25 frequency bands evenly spaced between 5 and 30 Hz. The wavelet transform was performed with a Morlet wavelet of width 5. The average baseline power for the first 50ms of the trial was then subtracted within each frequency band.

Psychometric Analysis

We quantified choice probability by fitting psychometric functions and extracting parameters corresponding to choice bias and sensitivity. For a given dataset, the proportion of rightward responses for each TOA bin were computed, and the results were fit to a cumulative distribution function of the standard normal distribution using *psignifit* (Schütt et al., 2016). Default parameterization in *psignifit* was used, with a lapse rate set to 1%. Parameters for the point of equal selection (PES) and width (W) of the psychometric function were then extracted to quantify the choice bias and sensitivity respectively. The PES corresponded to the TOA at which the probability of right target responses was 0.5, while W corresponds to the interval over which the psychometric function rises between the 5th and 95th percentile of right target choices. Additionally, we measured the reaction time (RT) of each saccade relative to the onset of the first choice target in the trial sequence, and computed the mean μ_{RT} and standard deviation σ_{RT} of the resulting distribution.

In addition to trial-average analyses, we performed future statistical tests of psychometrics conditioned on the outcome of the previous trial, in order to i) assess whether the underlying behaviour was nonstationary and ii) assess whether our inferences about causal effects of tDCS were specific to a particular trial-type. We analyzed three different choice history effects: choice repetition, choice direction, and reaction time. For choice repetition testing, all trials where the choice direction from the past trial corresponded with the current trial were compared with trials where the choice direction alternated, thus this was agnostic to the direction of the choice. In contrast, the previous choice direction grouping contrasted trials where the previous choice was left vs. right. Finally, the previous reaction time contrast was done by performing a median split on the previous trial reaction time.

Difference-In-Difference Permutation Testing

We used a statistical method based on permutation testing (Blair and Karniski, 1993; Maris, 2004) of a difference-in-difference design (Angrist and Pischke, 2009; Marinescu et al., 2018) to assess whether the polarity of stimulation influenced choice behaviour and EEG time-frequency responses at different levels of analysis: group-level, individual-level, and session-level. This method takes into account potential confounds arising from nonstationarities between pre and post stimulation periods by estimating the null distribution of a difference-in-difference test. For the group-level analysis, data across all participants and all sessions was pooled together. Four trial groups were then constructed corresponding to the stimulation polarity $P = \{Anodal, Cathodal\}$ and stimulation condition $C = \{Pre, Post\}$. The PSE, W, μ_{RT}, σ_{RT}, and EEG time-frequency responses were computed for each polarity and condition, and a difference in difference D for a given metric m was computed as

$$D_m = [m^{Pre}_{Anodal} - m^{Post}_{Anodal}] - [m^{Pre}_{Cathodal} - m^{Post}_{Cathodal}] \tag{1}$$

This acted as the test statistic for the hypothesis that, for a given metric m, the polarity of the stimulation had an effect given any relative change between pre and post stimulation that was independent of stimulation polarity. To estimate the null distribution of the difference-in-difference, we took the combined matrix E and randomly assigned the trial indices to evenly-sized matrices $n = \{n_1, ..n_{P \times C = 4}\}$, and for a given metric computed the difference-in-difference

$$N(i) = [\mu_{n_1} - \mu_{n_2}] - [\mu_{n_3} - \mu_{n_4}] \tag{2}$$

where i is the permutation number. Because testing all possible permutations is computationally intractable, we selected 500 permutations for each statistical test, and used the resulting distribution to estimate the probability that the true difference in difference D_m could have resulted from random combinations of the data under the null assumption that there is no difference between either pre and post blocks or anodal and cathodal sessions. For a given metric m, a p-value was computed by comparing the number of permutations for which the true difference-in-difference was as extreme as the permuted test statistic.

$$p(N \geq D) = \frac{1}{i} \min(\sum_{j=1}^{i} \mathbb{1}(N(j) \geq D(j)), \sum_{j=1}^{i} \mathbb{1}(N(j) \geq D(j))) \tag{3}$$

where $\mathbb{1}$ is the indicator function (1 if true, 0 of false).

This procedure was then performed by combining all data within a single participant, and combining data within single session combinations. For the participant-level analysis, the procedure was identical as outlined above. For the session combination, we took all possible pairs of anodal and cathodal sessions in order to assess how the participant-average findings correspond to how paired sessions were distributed. Therefore, the same procedure was performed for all possible sessions.

We then modified the difference-in-difference method to assess whether polarity-dependent effects were specific to certain types of behavioural contexts at the three different levels of analysis. This was motivated by the idea that polarity-dependent changes to, for example the influence of a previous left and previous right choice, may be averaged out when not taken into account. In order to account for a behavioural context $b = b1, b2$ as an intervening factor between a given metric m and the stimulation polarity, we performed a triple-difference estimator. For psychometric responses of the current trial, we computed this triple-difference with respect to three choice history contexts (previous RT, previous choice direction, repetition). For EEG time-frequency responses, we computed the triple difference with respect to both the previous and current trial (previous RT, previous choice direction, repetition, future choice, future RT). The resulting test statistic for a given combination of M and b

$$D_m = [[m_{A,b_1 l}^{Pre} - m_{A,b_2 l}^{Pre}] - [_{A,b_1}^{Post} - _{A,b_2}^{Post}]] - [[m_{C,b_1 l}^{Pre} - m_{C,b_2 l}^{Pre}] - [_{C,b_1}^{Post} - _{C,b_2}^{Post}]] \tag{4}$$

and the resulting null distribution constructed from 500 permutations of 8 evenly sized null matrices $n = \{n_1, .. n_{P \times C \times B}\}$. The triple-difference test statistic and p-value was then computed for each psychometric group-level, participant-level, and all session combinations.

Results

The study was composed of 5 participants, who underwent between 8-10 repeated fMRI-guided HD-tDCS sessions in a pre-post design, alternating between anodal and cathodal stimulation over an approximately 5 week period. The free choice task was performed in 5 blocks of 90 trials while EEG activity over the right frontal eye fields was recorded during the pre and post-stimulation period. By recording multiple sessions from the same participants, we were able to evaluate the data at different levels of abstraction: Group, inter-individual, and intra-individual. *Fig 1* illustrates the relationship between higher and lower

levels of abstraction, i.e. higher-level relationships are constructed by deriving summary statistics, such as averages, of the constituent relationships of a lower-level. At the group-level, studies aim to collect enough participants such that an average response within a sample generalizes to the average response of a population. Inter-individual studies in turn assess whether these group-level averages are consistent with effects on individual participants, while intra-individual variability studies assess consistency of effects across repeated measures.

Figure 1: Levels of Abstraction. **A**: Causal diagram depicting the influence of tDCS on EEG activity, and EEG activity at different levels of abstraction. The validity of higher-level abstractions relies on the statistical independence of repeated measures of lower-level abstraction. **B**: Levels of abstraction analyzed in this paper. Group-level analysis are composed of repeated measures of individual participants, in turn are composed of repeated measures of individual sessions, are in turn composed of different ways we categorize different trials.

For a given level of analysis, a summary statistic is typically chosen to represent the influence of an intervention on the aggregate of many causal processes that comprise its lower level. How a summary statistic derived from one level of abstraction translates to understanding a component of a lower-level abstraction depends on the statistical properties of the system. In systems that have a property known as ergodicity, the average of an ensemble of measurements taken at one time is equivalent to the average of the long-run trajectory of a single component of that ensemble. Ergodicity is an important property of a physical system to deduce, because it allows for measurements at different levels of abstraction (cf. longitudinal measure-

ments of single processes & ensemble measurements over short periods) to be interchangeable. However, in non-ergodic systems the ensemble behaviour at a given time is statistically distinct from the behaviour of a single component across time. One way to confer non-ergodicity in a system is to add history-dependence, with the canonical example being the distinction between a Bernoulli process – a coin flip– and a random walk. While a Bernoulli process is ergodic, integrating the outcome of Bernoulli processes over time results in a statistical distinction between the outcomes of a single realization of the process and a collection of realizations at one time point. In such a process, even though the ensemble average of the system is zero for all points in time, the probability of a single trajectory diverging from the ensemble average grows as time extends – because of this, the variance of the ensemble increases over time, and measuring the long-run trajectory single process is not equivalent to measuring the ensemble at a single point in time.

The translational value of many studies using causal interventions to understand brain-behaviour relationships at least loosely relies on an assumption of ergodicity (Fisher et al., 2018; Mangalam and Kelty-Stephen, 2021; Molenaar, 2004) . If one could indeed anticipate that gathering outcome measures from a large group over a short period of time has a translational equivalency to outcome mesaures of a single outcome over a long period of time, this would provide a great service. But the exchangeability of cross-sectional and longitudinal perspectives is often hypothetical. In non-invasive brain stimulation approaches like tDCS, much effort has gone into understanding population effects, but it is still largely unclear if we have achieved a level of understanding that allows any degree of confidence in its ability to reliably intervene on pre-specified physiological processes at the individual level (either for establishing additional causal relationships in research or for clinical applications). Although the last decade of tDCS research has cast aspersions on the the exchangeability of group-level and individual inferences, it is largely unknown if these individual inferences (derived from repeated measures of trials within single-session designs) are sufficient to provide causal evidence for the effect of tDCS while taking into account the potential for tDCS-dependent and tDCS-independent sources of intra-individual variability.

Given this, we feel it is important to make serious inquiry of the way in which we combine data into the independent statistical units required for conventional inference methods – does this represent an useful summary of the causal relationships that comprise it, or is it an aggregation of a diversity of potential causal chains whose average is not an unbiased estimator? We therefore assess the data presented using different assumptions about statistical independence that may, or may not, appear justifiable, with the intent not to

make a specific ontological commitment to the appropriate level of abstraction for tDCS research *per se*, but to illuminate how different levels of abstraction translate. The manuscript is organized into the deconstruction of the same dataset at three levels of abstraction: group-level, inter-individual, and intra-individual.

Free Choice Task

We assessed the influence of tDCS at the three different levels of abstraction with a free choice saccade task (Caie et al., 2023; Noudoost and Moore, 2011; Soltani et al., 2013). In this task, and indeed in many sequential designs, trials are randomized such that there is no inherent utility in using information from previous trials. Given this normative definition of the task, a researcher may assume that trials can be treated as independent statistical units. However (although this is a somewhat thorny inference to derive, such that we would not commit to a specific ontology *vis a vis* generative sources of sequential biases in humans) (cite some counterfactuals)) it is often stated that humans are poor randomizers; indeed in this task and many others there is evidence for history-dependence that defies the normative independent-and-identically-distributed description of the task, as well as the assumptions of statistical analyses that follow the reification of trials as a behavioural "natural kind".

While the reasons for this may be diverse and overlapping, the importance with respect to the task motivation is that the most fundamental unit of analysis specified within the task design, the trial, can either be conceived as an independent measure suitable for aggregation, or a dependent measure that carries a unique sequence. As we saw noted conceptually by distinguishing Bernoulli processes and random walks, this interpretational duality carries important consequences for the translational value of ensemble measurements to process-level statements. It is also one that we will see reiterated at levels of abstraction beyond the single-trial, so it is important to emphasize that the ambiguity surrounding statistical dependence at the levels of abstraction we attend to in this manuscript is unlikely to terminate at the trial.

Each trial consisted of a delay interval, followed by two targets presented asynchronously to each visual hemifield (*Fig 2, Methods: Free Choice Saccade Task*). Participants directed a saccade to either target, as fast as possible, without anticipating. The temporal onset asynchrony (TOA) of the targets was randomized such that the preceding trial carried no additional information about the current trial. We quantified choice behaviour for each trial by recording the outcome of the choice (left or right) with respect to the TOA, that

Figure 2: Free Choice Task. **A**: Free choice task schematic. Each trial begins with a randomized delay period, followed by the appearance of two choice targets. The temporal onset asynchrony (TOA) denotes the delay between the onset of the left and right target, with negative asynchrony values denoting the precedence of the left target. **B**: Eye trace of two trials. Positive horizontal eye position is right of the fixation cross. Slid red lines indicate leftward target, dashed red lines indicate rightward target. **C**: Group-average choice probability. Proportion of right target choices are plotted against the temporal onset asynchrony. Circles indicated average choice probability for a given TOA. **D** Histogram (grey bars) and cumulative response function (dashed line) of grouped reaction times. The horizontal red line indicates the appearance of the first target.

is the difference in time between the two target onsets, and the reaction time of the saccade relative to the onset of the first choice target. We then fit psychometric curves to the choice direction probability as a function of the TOA (*Fig 2C*). From this, we derived two measures that were used to parameterize how choice direction depended on the TOA: the point of equal selection (PES), or the TOA at which there was an equal selection probability between left and right choices, and the slope of the psychometric function W, which denoted how fast the psychometric function rose between the 25th and 75th right choice probability percentile. These two parameters were used as aggregate measures of choice bias and sensitivity respectively; the PES denotes a directional bias in the data, while the slope denotes how much a change in the TOA around the midpoint corresponds with a change in the choice probability. Psychometric functions were fit with psignifit in python (*Methods: Psychometric Analysis*). The group-average psychometric function indicates that average choice probability for the rightward target reliably increased as a function of the TOA (*Fig 2C*), thus the TOA influenced the choice probability.

We also measured the reaction time of each trial , that is how long it took from the target onset to trigger a saccade. Model based decompositions of decision processes predict that reaction times are determined by the time it takes to integrate or amplify sequentially-sampled sensory information from an initial baseline evidence level to an action-triggering threshold (Carpenter, 1999; Carpenter and Williams, 1995; Cisek et al., 2009; Ratcliff and McKoon, 2008). In a bounded-accumulator framework, the mean reaction times (μ_{RT}) directly correspond to the rate at which evidence is amplified or accumulated – therefore, we used mean reaction times as a proxy measurement for a component of the choice process that may not be directly accessible from measurements of choice probability. Additionally, we recorded the standard deviation of the reaction times (σ_{RT}) as a proxy measurement for components of the choice process that may not change the mean reaction time, such as anticipatory responses, baseline evidence updating, and sensory noise (Caie et al., 2023; Kim et al., 2017; Nakahara et al., 2006). We stress that these are merely proxy measures for a model-based abstraction of the choice process, and are not intended to provide evidence for anything akin to a cognitive natural kind. Rather, they are instruments to decompose the inference process – we remain agnostic as to their functional interpretation beyond this utility herein. *Fig 2D* shows the empirical histograms (grey bars) and overlapped cumulative distribution (dashed lines) for the combined data. The same choice psychometrics and reaction time plots for the individual participants are then shown in *Fig 3*.

Figure 3: Individual Participant Psychometrics. **A**: Individual choice probabilities are plotted for the 5 participants. **B**: Reaction time histograms and cumulative response functions. Conventions as before in *Fig 2*.

Group-Average

We first analyzed the full dataset by combining trials across all participants into separate groups comprising the pre and post-stimulation periods for anodal and cathodal sessions separately. At this level of analysis, no distinction between participants are made. Although this method of combining data has obvious interpretational issues for any group-level conclusions drawn from a dataset with 5 participants (despite the repeated measures within participants), we perform this analysis not to support the generality of its outcome so much as to assess how they correspond to within-participants variability, whatever the result may be. In doing so, the claim may be voiced that we risk introducing a straw man for how group-level studies are conducted in the first place. In a sense this is true, however we contend that even if group-level variability is not always combined across participants in such a blind fashion, data within participants often is. To be explicit, the intent is to assess whether that the methodological issues presented by combining data from different sources, obvious that they may be to some at the group-level, are recapitulated at the intra-individual level.

Figure 4: Permutation Testing. **A**: Time series representation of a single tDCS session. Depicted is an arbitrary time series for anodal (top) and cathodal (bottom) sessions, demarcated by the dashed horizontal line. Pre and post-stimulation epochs are demarcated by the vertical horizontal line. Sessions were organized into 5 blocks of 90 trials for both pre and post-stimulation. Pre and post stimulation were separated by a 20 minute interval (not shown). **B**: Cartoon depiction of permutation testing, Top: for a given metric, the true difference-in-difference (DID) subtraction is obtained by subtracting the across-trial means from pre and post-stimulation for anodal (top left), subtracting the across-trial means from pre and post-stimulation for cathodal (top right), and then subtracting the difference of the two differences. The null DID distribution (bottom) is then obtained by randomly assigning trial indices to 4 evenly sized groups, performing the same subtraction, and iterating over a chosen number of permutations.

First we briefly justify how the analysis proceeded at the group-level. In a pre-post design, statistical

changes to outcome metrics may arise from a causal effect of the intervention, or from a confounding variable that influenced how the measured processes changed over time. It is well established that many behaviours (Fründ et al., 2014; Gilden, 2001; Kello et al., 2010; Mahmoodi et al., 2023; Urai et al., 2019), and the statistics of neural processes (Fox et al., 2007; Linkenkaer-Hansen et al., 2001; Marcos et al., 2013 are autocorrelated in time – because of this, treating pre and post-stimulation trials independently for a given session may yield a statistical change between pre and post, simply due to autocorrrelations in independent signals (Caie and Blohm, 2024; Granger and Newbold, 1974; Phillips, 1986). By alternating between anodal and cathodal stimulation, we sought to factor out this influence of temporal variability within-sessions in order to isolate the causal influence of tDCS as precisely as possible. However, given this we had to factor in the probability that differences could arise from random combinations of sessions in absence of a causal effect of tDCS. To this end, we developed a permutation-testing approach based on a difference-in-difference design (*Methods: Permutation Testing*). Briefly, this method estimates the null distribution of a given metric in the case where there is no difference between anodal and cathodal sessions, and compares it to the true difference-in-difference. Permutation testing therefore takes into account variability arising from the difference between pre and post blocks by providing a data-driven likelihood that the true difference between anodal and cathodal sessions would have arisen from random combinations of sessions. For each metric, we computed a difference-in-difference between anodal and cathodal sessions by subtracting pre-post difference in anodal sessions from the pre-post difference in cathodal sessions, and compared that result to the null distribution we computed from randomly permuted sets of trials. The procedure is explained visually in *Fig 4.*

Fig 5 shows the outcome of permutation testing for the psychometric (*Fig 5A*) and EEG time-frequency responses (*Fig 5B*) at the group-average level, where data across all participants and sessions were pooled together. *Fig xA* shows the empirical null distributions for the PES (top left), W (top right), μ_{RT} (bottom left) and σ_{RT} (bottom right) in grey, with the true difference-in-difference denoted by the solid vertical black line. Here, it can be seen that it was statistically likely that the difference-in-difference for both metrics used to quantify choice probability, the PES and the width, could have arisen from random permutations of trials (PES $p = .264$, W $p = .392$) . Thus, there is no evidence to suggest that the choice outcome was influenced by the polarity of the stimulation at the group-level. However, both metrics for reaction times were statistically unlikely to have arisen from random permutations of trials (μ_{RT} p < .001, σ_{RT} < .001), suggesting that the polarity of stimulation explained a significant portion of the variance in the data with

Figure 5: Group Permutation Testing. **A**: True DID (solid black bar) and empirical null distribution (grey bars) are plotted for the PSE (top left), width (top right), mean reaction time (bottom left) and RT standard deviation (bottom right). Significant p values are reported in bold. **B**: EEG time-frequency subtractions for left vs right choice (top) and slow vs fast RT (bottom). Depicted are the mean power subtractions (pre-post) for anodal (left) and cathodal (middle) sessions, and the polarity DID subtraction (right).

respect to how reaction times changed between pre and post-stimulation epochs at the group level.

We then modified the difference-in-difference calculations in the psychometrics, where a single metric was chosen to perform a double subtraction (anodal/cathodal, pre/post), to assess whether a change in behaviour corresponded with a change in the EEG time-frequency response. To do this, we computed a triple-difference estimator (Olden and Møen, 2022) by including a median split between a given type of behavioural trial as an additional way to divide the data. We split behavioural trials in three ways here based on the outcome of the trial following the EEG activity: left vs right choice and a median split between slow and fast RT . For a given split, we had a difference-in-difference for each time-frequency point for anodal sessions that first subtracted the median split in pre-stimulation, then the median split in post-stimulation, and then subtracted these two differences to yield a single difference-in-difference for the anodal sessions. This was then performed again for the cathodal sessions, and the final triple-difference estimator was calculated as the subtraction between the two. The triple-difference was then compared statistically to an empirical null distribution that was calculated as before, where instead of dividing the data in permutations of 4 groups it is now 8.

Fig 5 shows the EEG time-frequency triple-difference subtractions with respect to the upcoming choice direction (top row) and reaction time (bottom row) for the two session conditions (pre-post) for anodal (left)

and cathodal (middle) sessions. We then computed the triple-difference subtraction, plotted in the right column. For the triple-difference subtraction, we plotted the true difference first in grayscale, with statistically significant time-frequency pixels overlayed in colour (sidak corrected for multiple comparisons, p values < .0001 reported). Here, it can be seen that, just as with the behavioural psychometrics, there was no evidence for a polarity-dependent effect on how the EEG time-frequency response changed with respect to the future choice direction. However, this subtraction performed relative to the upcoming reaction time revealed a significant relationship in alpha and beta-band activity in the early portion of the trial, and alpha-band activity in the late portion of the trial. Therefore, when combining data from all participants and all sessions together as a single functional unit of analysis, we found evidence for a behavioural effect of tDCS on reaction times that followed a change in the recorded EEG response.

Free choice saccadic behaviour has been demonstrated to depend on information from the previous trial (Caie et al., 2023; Soltani et al., 2013). In the case of a tDCS effect that was dependent on choice history, it is conceivable that choice history-dependent effects may be averaged together, and thus obscured, in trial-averaged analyses. For example, if anodal and cathodal stimulation differentially influenced the choice process depending on whether the previous choice was in the left or right direction, combining these different trials together could obscure the tDCS-dependence. We therefore assessed whether the influence of tDCS on choice psychometrics and EEG time-frequency responses was dependent on information from the previous trial (*Methods: Permutation Testing*) We used three different metrics for the previous trial: the direction of the previous choice, whether the current choice corresponded to a repetition or an alternation of the previous choice, and whether the previous trial resulted in a slow or a fast reaction time (based on a median split). These outcome measures reflect three distinct abstractions of feedback that may occur following a trial: the previous direction split indexes whether tDCS influenced the trial-trial buildup of a directional bias (Dorris et al., n.d.; Fecteau et al., 2004, the repetition split indexes a trial-trial buildup of sequence effects that may not be specifically lateralized (Cho and Cohen, 2002; Gao et al., 2009), and the reaction time split indexes a metric of task performance analogous to post-error slowing (Dutilh et al., 2012; Laming, 1979), given the requirement of participants to choose as fast as possible.

We compared the true triple-difference estimator to the permuted null distribution for these three outcome measures. *Fig 6* shows the results for the repetition/alternation difference (top row), previous choice difference (middle row) and previous reaction time difference (bottom row). With respect to the psychomet-

rics, we found evidence for an effect of the previous choice direction on the PES ($p < .001$), and an effect of the previous reaction time on the W ($p = .016$) and σ_{RT} ($p = .05$). In contrast to the group-averaged choice and reaction time subtractions, statistical testing of the EEG time-frequency responses did not perfectly correspond with the psychometrics. While the direction of the previous choice was a better predictor of the influence of polarity on psychometrics, the identity of the previous choice *relative to the current choice* (repetition vs alternation) was a better predictor of changes to the EEG time frequency response, with a modulation in both the alpha and beta band surviving statistical significance. However, we did observe a modulation of beta-band activity with respect to the previous reaction time that co-occurred with the change to the width and RT variance. Taken together, we observe that the outcome of the previous trial was predictive of how choice psychometrics and EEG activity depended on tDCS polarity, however only when conditioning on the previous reaction time did we find a correspondence between statistical changes to the psychometrics and the EEG. From one perspective this provides evidence supporting the contextual importance of the choice behaviour with respect to the tDCS effect at the group level. From another perspective, this provides evidence to undermine any deconstruction of the correlational structure between the intervention and repeated measures of single trials that requires statistical independence in order to retain validity.

Figure 6: Choice History Permutation Testing. **A-B**: Permutation testing for triple difference of repetition vs alternation trials across all participants. **A**: depicts the psychometric true TD and null distribution as in *Fig 6*. **B**: EEG time-frequency response triple-difference. Shown are the repetition/alternation difference-in-difference (DID) for anodal (top left) and cathodal (top middle), from which the polarity TD is computed (top right). Significant TFR pixels are in colour. Bottom left: Time-series of alpha band (7Hz band, mean power triple-difference in black, std power triple-difference in grey, significant time-points shaded in blue). Bottom right: The same, for the beta band (15 Hz). **C-D**: Previous left vs right choice triple difference. Plots are otherwise the same. **E-F**: Previous reaction time (fast vs slow median split)

Inter-individual

Group-level analyses of tDCS studies have previously been shown to be comprised of largely heterogeneous results at the inter-individual level. This may result from fixed anatomical-physiological differences that result in variable effects being averaged together at the group level, or unexplained variability within participants that is erroneously interpreted as relevant to a causal effect of tDCS. Previously, it has been shown that the induced electric field for a given target site can vary dramatically across participants depending on individual morphological differences. Our stimulation protocol (*Methods: fMRI-guided HD-tDCS*) was designed so as to minimize confounding variables by a) minimizing current spread with a 4x1 HD-tDCS electrode montage (cite sources on that) and b) selecting a region of interest that corresponded to the individually-specific neural activation close to the putative human homologue of the right frontal eye field (frontal eye field where art thou), as measured through an fMRI saccade localizer task.

The resulting electrode montages for each participant are overlayed on individual 3D models of the scalp in *Fig 7*, with the induced electric fields for an anodal stimulation session (center electrode 2mA, surround electrodes -.5 mA) overlayed on both the surface of the scalp (top row) and the gray matter (bottom row). Visually, variation in the location, intensity, and spread of the current can be observed. We depicted the distribution of current intensity for a single participant as a histogram of the electric field normal vector (measured in V/m) for each triangle in the mesh composing the gray matter surface, with the average and standard deviation of each electric field normal shown at the target ROI for each participant in *Fig 7C*. Here, it can be seen that the intensity of the current was different for each participant. This confirms that inter-individual anatomical differences should be considered as a source of variation in outcome metrics, however with the number of participants in this study little systematic can be said about how current intensity and spread relates to the effect variance across individuals. Coordinates for the region on the gray-matter and pial surface for the center electrode, and the positions of the surround electrodes on the pial surface for each participant are shown in *Fig 8*.

We now describe the results of permutation testing for psychometrics and EEG time-frequency responses at the inter-individual level. *Fig 9* plots the z-score of each difference-in-difference for all trials (left), and the previous-trial triple-difference conditioned on repetitions vs. alternations (middle-left), previous left vs. right choice (middle-right), and previous slow vs. fast RT (right), with z-scores corresponding to a p-value below 0.05 coloured in black. First we assessed whether the null effect of choice probability and positive effect of

reaction times was recapitulated consistently at the individual level. Here we found that within individuals, contrary to the group-level findings, choice probability (as indicated by the PES and width parameters in the leftmost columns) was statistically distinguishable in all participants, however in contrasting directions (PES *subtraction:* increased in 2/5 participants, decreased in 2/5, null change in 1/5; W *subtraction:* increased in 1/5, decreased in 1/5, null in 3/5). Thus, the null effect observed at the group-level appeared to be composed of a mixture of null and contrasting positive results that averaged as a significant effect. In the case of reaction times, where we observed a group-level significance for both the mean and standard deviations, we again found hetereogeneity in the individual data. For the standard deviation, there was some consistency, however only two participants showed significant interactions with the tDCS polarity. For μ_{RT}, again only two participants (1 and 4) showed significant changes, however now in opposite directions.

Figure 7: Individual current modelling **A**: Electric field simulations for each participant (left to right) are overlayed on the scalp (top row), showing the center-surround HD-tDCS ring setup, and the gray matter surface (bottom row). Electric fields are plotted in v/M. **B**: Distribution of normal component of electric field for an example participant. Histogram is shown for all triangles in the 3D mesh corresponding to the gray matter surface. **C**: Normal component of the electric field at the ROI for each participant. The mean and standard deviation are shown for mesh triangles corresponding to less than a 2cm radial distance from the ROI.

At the group-level, we did not observe a significant change in the choice-dependent EEG activity. How-

Table 1: Table of ROI coordinates in SimNibs for each participant. $rFEF$ corresponds to location of ROI projected onto the gray matter surface. $rFEF_p$ corresponds to the location of the center electrode, as projected onto the skin surface. Surround electrodes are also projected onto the skin surface.

Participant	rFEF	rFEF$_p$	Surround 1	Surround 2	Surround 3	Surround 4
1	[57.57, 28.85, 40.44]	[58.95, 37.60, 41.19]	[35.24, 29.91, 70.47]	[71.19, -1.03, 41.43]	[72.06, 40.96, 4.56]	[39.84, 70.91, 37.96]
2	[61.00, 28.41, 38.58]	[68.30, 18.81, 32.54]	[46.06, 13.38, 64.42]	[78.89, -19.60, 34.07]	[78.81, 21.72, -4.33]	[56.01, 55.63, 32.34
3	[55.86, 32.96, 34.98]	[60.19, 36.63, 24.77]	[40.43, 31.74, 58.12]	[71.66, -1.07, 32.67]	[72.59, 36.31, -16.10]	[43.49, 70.44, 15.54]
4	[54.90, 16.85, 43.62]	[61.90, 17.00, 33.62]	[39.51, 9.59, 64.98]	[78.03, -19.24, 32.79]	[72.90, 20.35, -4.48]	[46.26, 51.88, 36.18]
5	[57.28, 27.02, 46.49]	[58.90, 37.60, 30.99]	[47.40, 28.37, 67.89]	[77.14, 1.59, 31.11]	[66.45, 45.27, -8.85]	[38.53, 73.04, 38.40]

ever, within participants we found a significant change in the choice-dependent alpha-band activity in 2 of 5 participants, and a change in the choice-dependent beta-band activity in 1 of 5 participants. *Fig 9B* plot the mean and standard deviation of the choice (left) and RT-dependent (middle left) alpha (top) and beta-band (bottom) power subtractions, with significant time-points (multiple comparison corrected p < .001) overlayed in blue. In contrast, at the group-level we did observe a reaction-time dependent modulation of alpha and beta-band activity, although a significant relationship at the individual participant level was only found in a single participant, suggesting that the detection of this effect resulted from an averaging effect that poorly explained the inter-individual variation. As such, we cannot conclude that the group-level statistics we reported previously are supported by a consistent relationship with inter-individual psychometrics.

We then performed the same analysis on choice history, using the same variables (repetition/alternation, previous choice direction, previous RT median split) as with the group-average. While the group-level averaging only resulted in a joint effect of history on psychometrics and EEG activity for the previous RT split, we report a diversity of effects reaching significance within participants here. For the repetition/alternation split, we see a change in the choice psychometrics for two participants, but alpha and beta-band changes surviving the significance threshold for all participants. Notably, these modulations occurred with distinct profiles across participants. Repetition-dependence on stimulation polarity resulted in a diversity of temporal profiles of alpha and beta modulation such that a clean distinction between an increase and decrease in band-power was difficult to assess – for example, in participant 2 (Fig 9B), there was an observed power difference in alpha activity that increased in the early portion of the trial, and decreased in the late portion.

In the previous choice split, we find additional evidence for heterogeneity in the psychometric effects,

with 3 of 5 participants showing an effect of PES in one direction, and 1 participant showing an effect in the opposite direction. Similarly, 2 of 5 participants showed an effect of σ_{RT}, again in opposing directions, with no corresponding consistency observable in either EEG bands analyzed here. Notably, in the previous RT split, having previously observed a significant group-level relationship in the W and σ_{RT}, only a single participant survived significance testing for either of these metrics (participant 4, $\sigma_{RT}\, p = xx$. The group-level change in W appears to have been driven by individual participant effects that were in the same direction for 4/5 participants, but none surviving significance testing in isolation. For σ_{RT}, the positive result seen in the group-level was composed of a single positive result at the group-level (participant 4), plus 3 participants with statistically discarded trends in this direction, and one participant with a statistically discarded trend in the other. As with the other findings, EEG alpha and beta-band activity was likewise heterogeneous. Although no group-level alpha-band relationship was observed, an individual alpha-band relationship was found in 2 of 5 participants, however in opposite directions with respect to changes in band-power. Conversely, a group-level beta-band relationship was observed, but again we found a mixture of previous RT-dependent increases and decreases across participants, again indicating that a reductionist approach to the correlational structure of tDCS-behaviour-EEG in this dataset at the level of frequency band-power over time may belie the complexity of the generative process.

Figure 8: Inter-Individual Permutation Testing. **A-B**: Psychometric and EEG permutation testing for participant 1. **A**: Z-scores for the true DID relative to the permuted null distribution for PSE, W, μ_{RT}, and σ_{RT}. Significant tests are denoted by black bars and stars (left). TD permutation z-scores for repetition vs alternation (second from left), previous left vs right (second from right) and previous slow vs fast RT (right) are also reported. **B**: Top: mean alpha-band DID for choice subtraction, RT subtraction, and mean alpha-band TD for repetition, previous choice, and RT subtractions. Bottom: The same, for beta band. **C-D**: Participant 2. **E-F**: Participant 3. **G-H**: Participant 4. **I-J**: Participant 5.

Intra-Individual

The previous analyses of inter-individual psychometrics and EEG time-frequency responses aggregated all of the tDCS sessions together, and split them by the polarity of the stimulation on that day only. We now appraise these findings in light of the variability observed across-sessions. Our method of assessing intra-individual variability was to treat every possible anodal and cathodal session pairing within our dataset as a possible permutation that could be used in isolation to provide evidence for the underlying relationship of tDCS polarity at the individual level. In our permutation analysis, we preserved the order of the polarities such that the subtraction logic would be consistent by only including the session permutations with the anodal stimulation session as the first entry (*Methods: Session Permutations*). For a participant with 10 sessions, 5 anodal and 5 cathodal, this resulted in 25 possible session pairings that could be used to contrast the effects of anodal and cathodal tDCS. Our general analysis strategy was to see if previously significant findings at the inter-individual level corresponded to consistent relationships across different session pairings, and if this relationship between the psychometric subtractions at the individual session level predicted differences in the EEG response.

Fig 10 shows the permutation z-scores for each session pairing within participants, plotted along rows, where the columns are separated by the choice psychometric (left: PSE, middle left: W, middle right: μ_{RT}, right: σ_{RT}). Session pairings that survived statistical significance (p < .05) are indicated with black bars, whereas session pairings that did not are indicated with grey. For comparative purposes, psychometric subtractions that survived significance testing at the inter-individual level are shaded in blue. The first observation that should be stated is that for every possible participant-psychometric combination, at least one possible pairing of anodal and cathodal sessions would have resulted in a statistically significant difference. This in itself is notable, because it calls in to question the notion that variability seen in inter-individual tDCS studies across participants reflects a veritable inter-individual difference, rather than a sample of a larger variability space than is measurable within the contexts of single paired-session studies. Naturally, such questions are specific to the variability imposed by the combination of the task and behaviour, the temporal sequence of the paired-sessions, and the sources of variability the participant imposes; because of this, these conclusions do not provide a refutation of any specific paired-session design *per se*, but offers a singular example where caution in the evidential value of single paired-sessions should be exercised.

The second observation that should be noted is a disconnect between psychometric subtractions that

Figure 9: Intra-individual variability in psychometric subtractions at the session-pairing level. **A**: Participant 1 Z-scores for each possible session-pairing permutation are plotted for $PSE, W, \mu_{RT}, \sigma_{RT}$, left to right. Black bars denote z-scores corresponding to a significant relationship (p<.05), grey bars denote p > .05. Plots shaded in blue denote where significance was reached for the corresponding psychometric subtraction when combining data across sessions. **B**: Participant 2. **C**: Participant 3. **D**: Participant 4. **E**: Participant 5

survived significance at the inter-individual level, and the consistency of the session pairings that constitute it. While some psychometric subtractions that survived inter-individual significance testing were remarkably consistent across session-pairings (see Participant 1 PSE, μ_{RT}, Participant 2 W, Participant 4 W), other psychometric subtractions surviving significance were composed of sessions that, when paired together, showed

highly variable effects (see Participant 1 σ_{RT}, Participant 2 σ_{RT}, Participant 4 PSE, μ_{RT}). Thus, it can be seen that for psychometric effects detected by combining repeated measures of anodal and cathodal sessions together, individual combinations of these sessions may yield strong effects in either the detected direction, or its opposite. Taken together, these assessments could be used to suggest that only psychometric subtractions with a consistent relationship at the session-level should be viewed as evidence, due to averaging confounds arising from the likelihood of detecting effects at the single paired-session level. Alternatively, we may take each session-pairing as a veridical representation of the relative effect of tDCS on these specific occassions, and as such interpret intra-individual differences in how tDCS influences behaviour.

Next, we turned to the correlational structure between the session-pairing psychometric effect, and the EEG activity in the alpha and beta bands. To compress this analysis, we averaged the band-power across the delay epoch. Although this is a concession, in the sense that we previously observed complex band-power changes that did not necessarily translate to a time-averaged directional change with a suitable summary statistic, we stress that this an analysis tool to interrogate how a correlation may be drawn through a common means of abstraction, which necessarily cleaves away some of the complexity we previously observed. *Fig 11* plots, for each participant and psychometric subtraction, the correlation between the mean triple-difference alpha band-power change across the delay epoch, and the z-score of the psychometric difference-in-difference subtraction for each session pairing.

To contrast the correlations we report here with the inter-individual level of analysis, we report the consistency between tests at the inter-individual level (where both the psychometric and EEG time-frequency subtractions reached significance), and tests at the intra-individual level (where they significantly correlated). The distinction between what these test is important to state up front. While the former reports whether the these separate significance tests coincided when aggregating all sessions, the latter reports whether there was a relationship between the measures on a session-session basis. For data limitations within-sessions, we neglect to include the metrics of choice history that we previously reported at the inter-individual level. In the inter-individual tests (fig xx), only three non-choice history tests coincided with significance in both the psychometrics and a corresponding change in the reported EEG band-power activity, all with respect to the PSE: participant 1, where a increase in the PSE subtraction coincided with an early-trial increase in alpha power, participant 3, where a increase in the PSE subtraction coincided with an early-trial reduction in beta-power, and participant 5, where a decrease in the PSE subtraction corresponded to a reduction in

Figure 10: Session-wise correlations between psychometrics and EEG alpha-band activity. **A**: Double-subtraction z-scores are plotted against mean alpha-band double subtractions (averaged across trial timepoints) for $PSE, W, \mu_{RT}, \sigma_{RT}$ for participant 1. Grey boxes denote correlations where there was significance in both the psychometrics and EEG measures at the session-average level. Blue boxes denote significant session-wise correlations. **B**: Participant 2. **C**: Participant 3. **D**: Participant 4. **E**: Participant 5.

middle-trial alpha power.

Beginning with the alpha-band correlations in *Fig 11*, we see that for both $P1_{PSE-->Alpha}$, $p = .312$ and $P5_{PSE-->Alpha}$, $p = .7$, despite both psychometric and EEG permutations surviving significance when

Figure 11: Session-wise correlations between psychoemtrics and EEG beta-band activity. Conventions the same as *Fig 11*.

averaged across sessions, we fail to find evidence for a relationship between the psychometric subtractions, as denoted by the z-score on the abscissa, and the corresponding mean alpha power-subtraction. In contrast, we find evidence for several relationships between the psychometric Z-score and the power subtractions ($P4_{PSE}$, $P1_{\mu_{RT}}$, $P3_{\mu_{RT}}$, $P5_{\mu_{RT}}$, $P1_{\sigma_{RT}}$, $P3_{\sigma_{RT}}$) that did not correspond with a coinciding change in both the average psychometric and EEG subtractions. In *Fig 12*, the same analysis is shown

for beta-band activity averaged across trials. Here we see that seven significant session-wise correlations ($P1_{PSE-->Beta}$, $P1_{\mu_{RT}-->Beta}$, $P1_{\sigma_{RT}-->Beta}$, $P2_{\mu_{RT}-->Beta}$, $P2_{\sigma_{RT}-->Beta}$, $P4_{PSE-->Beta}$, $P5_{\mu_{RT}-->Beta}$, none aligning with the single consistent change in psychometric and EEG beta power at the session-average ($P3_{PSE-->Beta}$, $p = .056$).

Discussion

In this work, we performed a test-retest study of the influence of HD-tDCS on free choice behaviour and its neural correlates (via pre-post EEG and behavioural measures) at multiple levels of abstraction: group-level, inter-individual, and intra-individual. A growing body of evidence has suggested that the exchangeability of results derived from these levels of abstraction are subject to an array of confounds (Horvath et al., 2015; B. Krause and Cohen Kadosh, 2014; Li et al., 2015). Our experimental setup went to lengths to minimize the influence of some of these confounds by individualizing the selection of a brain region involved in the planning and execution of saccadic eye movements with fMRI, using the individual MRIs to guide electrode participant prior to every session, and selecting an electrode montage that minimizes current spread to non-targeted brain regions while maintaining focality (Mikkonen et al., 2020; Villamar et al., 2013). We then developed a permutation testing procedure that allowed us to estimate in a data-driven fashion the probability that changes to psychometric parameters and EEG time-frequency data were related to the tDCS polarity. We observed a general inequivalency between the statistical relationships uncovered at one level, and the statistical relationships that comprised it at a lower level of abstraction.

While our approach may be criticized as a *reductio ad absurdem* with respect to group-level analysis, as the consequences of inter-individual sources of variability are established enough such that blindly combining participants into common statistical units is recognized by many to be flawed (Gallistel, 2012), we argue that the recapitulation of this inconsistency at the intra-individual level of abstraction provides evidence for the interpretation that averaging repeated measures across sessions within-individuals is similarly confounded. Because of the intra-individual variability we report, caution is warranted in interpreting the variability seen in inter-individual studies as evidence of inter-individual variability in tDCS-mediated effects *per se*, in absence of a principled explanation for the intra-individual variability intrinsic that should be expected under repeated measures of a given experimental context. Without such an explanation, we cannot at this point recommend a single privileged level of analysis (group, inter-individual, intra-individual)

at which the effects of tDCS *should* be expected to be suitably generalizable in the context of our task and paradigm, such that repeated measures could reliably separate tDCS-mediated effects from the unknown statistical properties of unmeasured supervening processes.

The early studies evaluating tDCS in humans tested how TMS-induced motor evoked potentials changed over time as a function of the polarity of stimulation (Nitsche and Paulus, 2000, 2001). It was found that anodal stimulation potentiated the size of the motor evoked potential following stimulation, while following cathodal stimulation it was depressed. Although responsible for an influx of human tDCS research, this work have undergone a re-evaluation in recent years. Numerous studies have reported substantial inter and intra-subject variability in this paradigm (Chew et al., 2015; Dyke et al., 2016; Horvath et al., 2016; Vergallito et al., 2022; Zappasodi et al., 2018). Some of this variability may be attributed to the specifics of the TMS-MEP protocol (ex Herwig et al., 2001) which are beyond the scope of our work, however other sources of variability are suitably general such that they warrant discussion here.

Most of the narrative surrounding inter-individual variability in response to tDCS concerns differential anatomical and physiological features that may generate individually specific effects subject to masking by group-averaging. With appropriate electrode montages, it is uncontroversial that electric fields can be induced in humans using tDCS beyond the skull, and this can influence neural activity (Chhatbar et al., 2018; Huang et al., 2017; Opitz et al., 2016; Ruhnau et al., 2018; Vöröslakos et al., 2018; Wiethoff et al., 2014). However individual differences in anatomical features are substantial enough to predict different strengths and orientations of current flow at targeted sites, and casts doubt on the anatomical specificity of tDCS (Bikson and Rahman, 2013) – this may be predicted to have markedly different influence on aggregate neural measures, such as EEG, that largely sample activity arising from dipole moments generated by the temporal synchrony of neuronal activity along the columnar architecture of cortical microcircuitry (Buzsáki et al., 2012).

With respect to our paradigm, individual differences in tDCS-induced electric fields can be viewed from at least two vantages. The first is based on predictions derived from a circuit-level reductionist approach (Douglas et al., 1989) to the relationship between neural activity and free choice saccadic behaviour (Heinzle et al., 2007; Soltani et al., 2013). In this view, a dissociation between the predictions of modifying activity in superficial and deep layers of a given brain region may be made, such that a dissociation between the

effects of a tangential and radial current intervention may be predicted (for similar logic see Evans et al., 2022; Rawji et al., 2018, but see Joshi et al., 2023; Laakso et al., 2024). Similarly, variability in the functional definition of a brain region, and its consistency across individuals, may contribute – indeed the frontal eye field in humans is a somewhat nebulous concept that largely inherits etiology from non-human primate studies, and has not converged on a singular definition in humans (Vernet et al., 2014). Accordingly, across our 5 participants there was variability in the localization procedure when aligned to a common reference frame. Alternatively, the variance associated with the induced electric field may be interpreted from a more holistic perspective with respect to global neural activity. In our study, the intensity and the variability of the induced current was inconsistent across participants. The degree to which this may cause differential changes to distal activity in the network (either causally through the target region or indirectly through current spread) is challenging to assess, but these unknowns should not be neglected when condensing the results into a mechanistic-reductionist interpretation of intervening on a single brain region with tDCS (Bergmann and Hartwigsen, 2021; M. R. Krause et al., 2017; Louviot et al., 2022; Mitchell and Potter, 2024; Polanía et al., 2011; Silberstein and Chemero, 2013).

We show that inter-individual changes to psychometric measures and EEG time-frequency correlates, when averaged across sessions, were composed of a mixture of confirmatory and contradictory results. In some cases, individual session-pairings in our permutation testing analysis largely followed the session-averages, while others session-pairings yielded polarizing effects. All we can say from this definitively is that single session- pairings contrasting the effects of anodal and cathodal stimulation are insufficient to provide a representative statistic of an inter-individual effect, as they may diverge substantially from session-averages across repeated measures. However, this intra-individual variability may have arisen from a number of distinct causes that can be roughly divided into two classes. The first would posit that there are intra-individual variables within the causal chain between the measured EEG signal and free choice saccadic behaviour that modify how tDCS impacts the measurements, and that these intra-individual variables change over suitable timescales to confound the independence of sessions. This would essentially propose that i) the causal effect of tDCS on the brain-behaviour relationship is state-dependent, and ii) the underlying states that determine its causal effect are non-stationary across repeated-measures. Although causal assessments at this level can be challenging, and of course depend on how an all-encompassing term like 'brain-state' is actually operationalized (Greene et al., 2023), there is some experimental evidence for state-dependent effects of tDCS at different levels of abstraction – from cellular mechanisms to functional networks (Antal et al., 2007; Paulus

Figure 12: Sequential and I.I.D Perspectives of Statistical Variations in Free Choice Behaviour. Depicted is a hypothetical generative process that supervenes over the correlational structure of free choice saccadic behaviour and EEG activity at the group-level for two intervention conditions (ex: pre-post anodal and pre-post cathodal subtractions). These groups of measurements are composed of a set of individuals each with a set of anodal and cathodal sessions. These sessions in turn can be conceived as independent and identically distributed measurements (I.I.D Perspective), or history-dependent measures arising from a unique sequence (depicted as an ensemble of random walks here, although this choice is arbitrary). For each session measurement, individual trial outcomes can arise from a sequential (depicted as a choice sequence tree with unique historicities) or an ensemble statistical model (a generic bounded integration process here).

and Rothwell, 2016; Vergallito et al., 2023).

A second class of intra-individual explanans are somewhat agnostic to specific mechanistic predictions of tDCS, but instead are concerned with the expectations of the null model –what should be expected of the statistical structure of intra-individual variability in absence of a causal effect? In this manuscript, we developed a permutation-testing approach to assess in a data-driven manner the probability that a change in psychometrics or EEG time-frequency measures were statistically distinguishable by the polarity of the stimulation at three different levels of abstraction. Although this provides us with a useful tool to appraise statistical variations as a function of the intervention we used, the way in which these results are interpreted

is far from straightforward, and ultimately depends on the degree to which the measured processes are taken to be independent statistical units (Caie and Blohm, 2024; Mangalam and Kelty-Stephen, 2022; Molenaar, 2004). In the task analyzed in this manuscript, it is fairly straightforward to conceive of different generative models that may confirm or contradict the independence of these statistical units at different levels of abstraction (this division is depicted in *Fig 13*). While our analysis of sequence-dependence at the group and inter-individual level, and prior work on this task, suggest that the outcome of past trials can provide predictive value, we caution against this being used as proof positive of a sequentially-dependent generative process underlying the observations – it would be perhaps trivial to construct a null proposition wherein ensemble models of sequences could be viewed as independent functional units over a longer timescale, such that any historicity we observe falls out of an ensemble of sequences that are I.I.D with respect to the timescale of the entire experiment [1]. However, this should not discourage us in discussing potential generative models, if only to clarify how the predictions they generate differ.

The atomization of behavioural measures into discrete functional units is practically axiomatic to the cognitive neuroscience approach (Cisek, 2019; Huk et al., 2018). Under this view, repeated measurements of a certain psychological classification of behaviour are recorded, and variance associated with repeated measures provide evidence for process-level variation that can be used to guide model-based decompositions of behaviour, and help separate expected sources of process-level variance from causal changes in the presence of interventions. Although there is plenty of counterargumentation suggesting that these measurements, and the neural processes they are often correlated with (), are not independent – aphorisms about the inability for humans to appear algorithmically random abound – the consequences for this on statistical inference are often not explicated. It is well known in econometrics, a field where the inability to uncover generative models with predictive power is more accepted (and perhaps more obvious), that autocorrelations in statistical models produce spurious correlations (Granger and Newbold, 1974; Phillips, 1986). The consequence of this for single paired measurements of sequentially-dependent behaviours, such as might be expected in the case of our pre-post experiment, is a high rate of false-positive assessments when averaging across trials, and a general inability to perform inference of meaningful statistical variations with single session pairings.

Eschewing the atomization of trials is not a death knell for statistical testing in general. While we may

---

[1] this argumentation relies on arguments from computability theory, where it has been shown repeatedly that no algorithm could be constructed to determine whether a given input measurement was a product of a random event or a deterministic machine totalling the sum of the observations. This is related to incompleteness theorems that state a formal system cannot demonstrate its own consistency

have sufficient evidence to cast doubt on the statistical independence of trials in tasks such as ours (despite attempts at 'randomization'), it may be found that sessions, themselves composed of sequentially-dependent measures, are independent (or *independent enough*) to be used as repeated measurements for inferential studies. Our assessment of intra-individual variability is inconclusive in this regard – while we can say that variance in session-pairings may arise from either an IID perspective combined with tDCS-dependent intra-individual variance, or a null model incorporating statistical dependencies between sessions, it is unclear to what degree these contrasting views are separable. Future work assessing invariances at timescales relevant to session-wise test-retesting are likely needed to provide useful answers to these questions, although this is in a sense a recursive logic that will in turn necessitate new counterfactuals about timescales yet longer. What we can conclude is that the competing hypotheses and sources of variability we observe in free choice behaviour and its EEG correlates are such that a privileged level of abstraction for inferring causal effects of tDCS remains elusive in our paradigm.

References

Angrist, J. D., & Pischke, J.-S. (2009). *Mostly Harmless Econometrics: An Empiricist's Companion*. Princeton University Press.

Antal, A., Terney, D., Poreisz, C., & Paulus, W. (2007). Towards unravelling task-related modulations of neuroplastic changes induced in the human motor cortex. *The European Journal of Neuroscience*, *26*(9).

Bergmann, T. O., & Hartwigsen, G. (2021). Inferring Causality from Noninvasive Brain Stimulation in Cognitive Neuroscience. *Journal of Cognitive Neuroscience*, *33*(2).

Bikson, M., & Rahman, A. (2013). Origins of specificity during tDCS: Anatomical, activity-selective, and input-bias mechanisms. *Frontiers in Human Neuroscience*, *7*.

Bikson, M., Rahman, A., & Datta, A. (2012). Computational models of transcranial direct current stimulation. *Clinical EEG and neuroscience*, *43*(3).

Bindman, L. J., Lippold, O. C. J., & Redfearn, J. W. T. (1964). The action of brief polarizing currents on the cerebral cortex of the rat (1) during current flow and (2) in the production of long-lasting after-effects. *The Journal of Physiology*, *172*(3).

Blair, R. C., & Karniski, W. (1993). An alternative method for significance testing of waveform difference potentials. *Psychophysiology*, *30*(5).

Buzsáki, G., Anastassiou, C. A., & Koch, C. (2012). The origin of extracellular fields and currents — EEG, ECoG, LFP and spikes.

Caie, B., & Blohm, G. (2024, March). History-dependence shapes causal inference of brain-behaviour relationships [arXiv:2403.00947 [q-bio]].

Caie, B., Endres, D., Khan, A., & Blohm, G. (2023, December). Choice anticipation as gated accumulation of sensory expectations.

Carpenter, R. H. S. (1999). A neural mechanism that randomises behaviour. *Journal of Consciousness Studies*, *6*(1).

Carpenter, R. H. S., & Williams, M. L. L. (1995). Neural computation of log likelihood in control of saccadic eye movements. *Nature*, *377*(6544).

Chaudhuri, R., Knoblauch, K., Gariel, M.-A., Kennedy, H., & Wang, X.-J. (2015). A Large-Scale Circuit Mechanism for Hierarchical Dynamical Processing in the Primate Cortex. *Neuron*, *88*(2).

Chew, T., Ho, K.-A., & Loo, C. K. (2015). Inter- and Intra-individual Variability in Response to Transcranial Direct Current Stimulation (tDCS) at Varying Current Intensities. *Brain Stimulation*, *8*(6).

Chhatbar, P. Y., Kautz, S. A., Takacs, I., Rowland, N. C., Revuelta, G. J., George, M. S., Bikson, M., & Feng, W. (2018). Evidence of transcranial direct current stimulation-generated electric fields at subthalamic level in human brain *in vivo. Brain Stimulation*, *11*(4).

Cho, R., & Cohen, J. D. (2002). Mechanisms underlying dependencies of performance on stimulus history in a two-alternative forced-choice task. *Cognitive, Affective, & Behavioral Neuroscience*, *2*(4).

Cisek, P. (2019). Resynthesizing behavior through phylogenetic refinement. *Attention, Perception & Psychophysics*, *81*(7).

Cisek, P., Puskas, G. A., & El-Murr, S. (2009). Decisions in changing conditions: The urgency-gating model. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *29*(37).

Codling, E. A., Plank, M. J., & Benhamou, S. (2008). Random walk models in biology. *Journal of The Royal Society Interface*, *5*(25).

Datta, A., Truong, D., Minhas, P., Parra, L. C., & Bikson, M. (2012). Inter-Individual Variation during Transcranial Direct Current Stimulation and Normalization of Dose Using MRI-Derived Computational Models. *Frontiers in Psychiatry*, *3*.

Deco, G., & Romo, R. (2008). The role of fluctuations in perception. *Trends in Neurosciences*, *31*(11).

Dorris, M. C., Taylor, T. L., Klein, R. M., & Munoz, D. P. (n.d.). Influence of Previous Visual Stimulus or Saccade on Saccadic Reaction Times in Monkey.

Douglas, R. J., Martin, K. A., & Whitteridge, D. (1989). A Canonical Microcircuit for Neocortex. *Neural Computation*, *1*(4).

Dutilh, G., Vandekerckhove, J., Forstmann, B. U., Keuleers, E., Brysbaert, M., & Wagenmakers, E.-J. (2012). Testing theories of post-error slowing. *Attention, Perception, & Psychophysics*, *74*(2).

Dyke, K., Kim, S., Jackson, G. M., & Jackson, S. R. (2016). Intra-Subject Consistency and Reliability of Response Following 2mA Transcranial Direct Current Stimulation. *Brain Stimulation*, *9*(6).

Evans, C., Zich, C., Lee, J. S. A., Ward, N., & Bestmann, S. (2022). Inter-individual variability in current direction for common tDCS montages. *NeuroImage*, *260*.

Fecteau, J. H., Au, C., Armstrong, I. T., & Munoz, D. P. (2004). Sensory biases produce alternation advantage found in sequential saccadic eye movement tasks. *Experimental Brain Research*, *159*(1).

Filmer, H. L., Ehrhardt, S. E., Shaw, T. B., Mattingley, J. B., & Dux, P. E. (2019). The efficacy of transcranial direct current stimulation to prefrontal areas is related to underlying cortical morphology. *NeuroImage*, *196*.

Fischer, J., & Whitney, D. (2014). Serial dependence in visual perception. *Nature Neuroscience*, *17*(5).

Fisher, A. J., Medaglia, J. D., & Jeronimus, B. F. (2018). Lack of group-to-individual generalizability is a threat to human subjects research. *Proceedings of the National Academy of Sciences, 115*(27).

Fox, M. D., Snyder, A. Z., Vincent, J. L., & Raichle, M. E. (2007). Intrinsic Fluctuations within Cortical Systems Account for Intertrial Variability in Human Behavior. *Neuron, 56*(1).

Fründ, I., Wichmann, F. A., & Macke, J. H. (2014). Quantifying the effect of intertrial dependence on perceptual decisions. *Journal of Vision, 14*(7).

Gallistel, C. R. (2012). On the evils of group averaging: Commentary on Nevin's "Resistance to extinction and behavioral momentum". *Behavioural Processes, 90*(1).

Gao, J., Wong-Lin, K., Holmes, P., Simen, P., & Cohen, J. D. (2009). Sequential Effects in Two-Choice Reaction Time Tasks: Decomposition and Synthesis of Mechanisms. *Neural Computation, 21*(9).

Gilden, D. L. (2001). Cognitive emissions of 1/f noise. *Psychological Review, 108*(1).

Granger, C. W. J., & Newbold, P. (1974). Spurious regressions in econometrics. *Journal of Econometrics, 2*(2).

Greene, A. S., Horien, C., Barson, D., Scheinost, D., & Constable, R. T. (2023). Why is everyone talking about brain state? *Trends in Neurosciences, 46*(7).

Heinzle, J., Hepp, K., & Martin, K. A. C. (2007). A Microcircuit Model of the Frontal Eye Fields. *The Journal of Neuroscience, 27*(35).

Herwig, U., Padberg, F., Unger, J., Spitzer, M., & Schönfeldt-Lecuona, C. (2001). Transcranial magnetic stimulation in therapy studies: Examination of the reliability of "standard" coil positioning by neuronavigation. *Biological Psychiatry, 50*(1).

Horvath, J. C., Carter, O., & Forte, J. D. (2014). Transcranial direct current stimulation: Five important issues we aren't discussing (but probably should be). *Frontiers in systems neuroscience, 8*.

Horvath, J. C., Forte, J. D., & Carter, O. (2015). Evidence that transcranial direct current stimulation (tDCS) generates little-to-no reliable neurophysiologic effect beyond MEP amplitude modulation in healthy human subjects: A systematic review. *Neuropsychologia, 66*.

Horvath, J. C., Vogrin, S. J., Carter, O., Cook, M. J., & Forte, J. D. (2016). Effects of a common transcranial direct current stimulation (tDCS) protocol on motor evoked potentials found to be highly variable within individuals over 9 testing sessions. *Experimental Brain Research, 234*(9).

Huang, Y., Liu, A. A., Lafon, B., Friedman, D., Dayan, M., Wang, X., Bikson, M., Doyle, W. K., Devinsky, O., & Parra, L. C. (2017). Measurements and models of electric fields in the in vivo human brain during transcranial electric stimulation. *eLife, 6*.

Huk, A., Bonnen, K., & He, B. J. (2018). Beyond Trial-Based Paradigms: Continuous Behavior, Ongoing Neural Activity, and Natural Stimuli. *The Journal of Neuroscience*, *38*(35).

Jackson, M. P., Rahman, A., Lafon, B., Kronberg, G., Ling, D., Parra, L. C., & Bikson, M. (2016). Animal Models of transcranial Direct Current Stimulation: Methods and Mechanisms. *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology*, *127*(11).

Joshi, R., Murali, S., & Thirugnanasambandam, N. (2023). Behavioral Validation of Individualized Low-Intensity Transcranial Electrical Stimulation (tES) Protocols. *eNeuro*, *10*(12).

Kabakov, A. Y., Muller, P. A., Pascual-Leone, A., Jensen, F. E., & Rotenberg, A. (2012). Contribution of axonal orientation to pathway-dependent modulation of excitatory transmission by direct current stimulation in isolated rat hippocampus. *Journal of Neurophysiology*, *107*(7).

Kello, C. T., Brown, G. D. A., Ferrer-i-Cancho, R., Holden, J. G., Linkenkaer-Hansen, K., Rhodes, T., & Van Orden, G. C. (2010). Scaling laws in cognitive sciences. *Trends in Cognitive Sciences*, *14*(5).

Kim, T. D., Kabir, M., & Gold, J. I. (2017). Coupled Decision Processes Update and Maintain Saccadic Priors in a Dynamic Environment. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *37*(13).

Krause, B., & Cohen Kadosh, R. (2014). Not all brains are created equal: The relevance of individual differences in responsiveness to transcranial electrical stimulation. *Frontiers in Systems Neuroscience*, *8*.

Krause, M. R., Zanos, T. P., Csorba, B. A., Phillips, M. E., Datta, A., Pack, C. C., Pilly, P. K., & Choe, J. (2017). Transcranial Direct Current Stimulation Facilitates Associative Learning and Alters Functional Connectivity in the Primate Brain.

Laakso, I., Tani, K., Gomez-Tames, J., Hirata, A., & Tanaka, S. (2024). Small effects of electric field on motor cortical excitability following anodal tDCS. *iScience*, *27*(2).

Laming, D. (1979). Autocorrelation of choice-reaction times. *Acta Psychologica*, *43*(5).

Lefaucheur, J.-P., & Wendling, F. (2019). Mechanisms of action of tDCS: A brief and practical overview. *Neurophysiologie Clinique*, *49*(4).

Li, L. M., Uehara, K., & Hanakawa, T. (2015). The contribution of interindividual factors to variability of response in transcranial direct current stimulation studies. *Frontiers in Cellular Neuroscience*, *9*.

Linkenkaer-Hansen, K., Nikouline, V. V., Matias Palva, J., & Ilmoniemi, R. J. (2001). *Long-Range Temporal Correlations and Scaling Behavior in Human Brain Oscillations* (tech. rep.).

Liu, A., Vöröslakos, M., Kronberg, G., Henin, S., Krause, M. R., Huang, Y., Opitz, A., Mehta, A., Pack, C. C., Krekelberg, B., Berényi, A., Parra, L. C., Melloni, L., Devinsky, O., & Buzsáki, G. (2018). Immediate neurophysiological effects of transcranial electrical stimulation [Publisher: Nature Publishing Group]. *Nature Communications*, *9*(1).

Louviot, S., Tyvaert, L., Maillard, L. G., Colnat-Coulbois, S., Dmochowski, J., & Koessler, L. (2022). Transcranial Electrical Stimulation generates electric fields in deep human brain structures. *Brain Stimulation*, *15*(1).

Macke, J. H., & Nienborg, H. (2019). Choice (-history) correlations in sensory cortex: Cause or consequence? *Current Opinion in Neurobiology*, *58*.

Mahmoodi, K., Kerick, S. E., Grigolini, P., Franaszczuk, P. J., & West, B. J. (2023). Temporal complexity measure of reaction time series: Operational versus event time. *Brain and Behavior*, *13*(7).

Mangalam, M., & Kelty-Stephen, D. G. (2021). Point estimates, Simpson's paradox, and nonergodicity in biological sciences. *Neuroscience & Biobehavioral Reviews*, *125*.

Mangalam, M., & Kelty-Stephen, D. G. (2022). Ergodic descriptors of non-ergodic stochastic processes. *Journal of The Royal Society Interface*, *19*(189).

Marcos, E., Pani, P., Brunamonti, E., Deco, G., Ferraina, S., & Verschure, P. (2013). Neural Variability in Premotor Cortex Is Modulated by Trial History and Predicts Behavioral Performance. *Neuron*, *78*(2).

Marinescu, I. E., Lawlor, P. N., & Kording, K. P. (2018). Quasi-experimental causality in neuroscience and behavioural research. *Nature Human Behaviour*.

Maris, E. (2004). Randomization tests for ERP topographies and whole spatiotemporal data matrices. *Psychophysiology*, *41*(1).

Medaglia, J. D., Ramanathan, D. M., Venkatesan, U. M., & Hillary, F. G. (2011). The challenge of non-ergodicity in network neuroscience. *Network: Computation in Neural Systems*, *22*(1).

Merton, P. A., & Morton, H. B. (1980). Stimulation of the cerebral cortex in the intact human subject [Publisher: Nature Publishing Group]. *Nature*, *285*(5762).

Mikkonen, M., Laakso, I., Tanaka, S., & Hirata, A. (2020). Cost of focality in TDCS: Interindividual variability in electric fields. *Brain Stimulation*, *13*(1).

Mitchell, K. J., & Potter, H. D. (2024, June). Beyond mechanism – extending our concepts of causation in neuroscience.

Molenaar, P. C. M. (2004). A Manifesto on Psychology as Idiographic Science: Bringing the Person Back Into Scientific Psychology, This Time Forever. *Measurement: Interdisciplinary Research & Perspective*, *2*(4).

Molenaar, P. C., & Campbell, C. G. (2009). The New Person-Specific Paradigm in Psychology. *Current Directions in Psychological Science*, *18*(2).

Murray, J. D., Bernacchia, A., Freedman, D. J., Romo, R., Wallis, J. D., Cai, X., Padoa-Schioppa, C., Pasternak, T., Seo, H., Lee, D., & Wang, X.-J. (2014). A hierarchy of intrinsic timescales across primate cortex. *Nature Neuroscience*, *17*(12).

Nakahara, H., Nakamura, K., & Hikosaka, O. (2006). Extended LATER model can account for trial-by-trial variability of both pre-and post-processes. *Neural Networks*, *19*.

Nitsche, M. A., & Paulus, W. (2000). Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *The Journal of Physiology*, *527 Pt 3*.

Nitsche, M. A., & Paulus, W. (2001). Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans. *Neurology*, *57*(10).

Noudoost, B., & Moore, T. (2011). Control of visual cortical signals by prefrontal dopamine. *Nature*, *474*(7351).

Olden, A., & Møen, J. (2022). The triple difference estimator. *The Econometrics Journal*, *25*(3).

Opitz, A., Falchier, A., Yan, C.-G., Yeagle, E. M., Linn, G. S., Megevand, P., Thielscher, A., Deborah A, R., Milham, M. P., Mehta, A. D., & Schroeder, C. E. (2016). Spatiotemporal structure of intracranial electric fields induced by transcranial electric stimulation in humans and nonhuman primates. *Scientific Reports*, *6*.

Opitz, A., Paulus, W., Will, S., Antunes, A., & Thielscher, A. (2015). Determinants of the electric field during transcranial direct current stimulation. *NeuroImage*, *109*.

Paulus, W., & Rothwell, J. C. (2016). Membrane resistance and shunting inhibition: Where biophysics meets state-dependent human neurophysiology. *The Journal of Physiology*, *594*(10).

Pelletier, S. J., & Cicchetti, F. (2014). Cellular and molecular mechanisms of action of transcranial direct current stimulation: Evidence from in vitro and in vivo models. *The International Journal of Neuropsychopharmacology*, *18*(2).

Pellicciari, M. C., Brignani, D., & Miniussi, C. (2013). Excitability modulation of the motor system induced by transcranial direct current stimulation: A multimodal approach. *NeuroImage*, *83*.

Phillips, P. C. B. (1986). Understanding spurious regressions in econometrics. *Journal of Econometrics*, *33*(3).

Polanía, R., Paulus, W., & Nitsche, M. A. (2011). Modulating cortico-striatal and thalamo-cortical functional connectivity with transcranial direct current stimulation. *Human Brain Mapping*, *33*(10).

Priori, A., Berardelli, A., Rona, S., Accornero, N., & Manfredi, M. (1998). Polarization of the human motor cortex through the scalp. *Neuroreport*, *9*(10).

Purpura, D. P., & McMurty, J. G. (1965). Intracellular Activities and Evoked Potential Changes During Polarization of the Motor Cortex. *Journal of neurophysiology*, *28*.

Rahman, A., Reato, D., Arlotti, M., Gasca, F., Datta, A., Parra, L. C., & Bikson, M. (2013). Cellular effects of acute direct current stimulation: Somatic and synaptic terminal effects. *The Journal of Physiology*, *591*(Pt 10).

Ratcliff, R., & McKoon, G. (2008). The Diffusion Decision Model: Theory and Data for Two-Choice Decision Tasks. *Neural computation*, *20*(4).

Rawji, V., Ciocca, M., Zacharia, A., Soares, D., Truong, D., Bikson, M., Rothwell, J., & Bestmann, S. (2018). tDCS changes in motor excitability are specific to orientation of current flow. *Brain Stimulation*, *11*(2).

Rothwell, J. (2018). Transcranial brain stimulation: Past and future. *Brain and Neuroscience Advances*, *2*.

Ruhnau, P., Rufener, K. S., Heinze, H. .-., & Zaehle, T. (2018). Sailing in a sea of disbelief: In vivo measurements of transcranial electric stimulation in human subcortical structures. *Brain Stimulation*, *11*(1).

Saturnino, G., Antunes, A., Stelzer, J., & Thielscher, A. (2015). SimNIBS: A versatile toolbox for simulating fields generated by transcranial brain stimulation.

Schütt, H. H., Harmeling, S., Macke, J. H., & Wichmann, F. A. (2016). Painfree and accurate Bayesian estimation of psychometric functions for (potentially) overdispersed data. *Vision Research*, *122*.

Silberstein, M., & Chemero, A. (2013). Constraints on Localization and Decomposition as Explanatory Strategies in the Biological Sciences. *Philosophy of Science*, *80*(5).

Soltani, A., Noudoost, B., & Moore, T. (2013). Dissociable dopaminergic control of saccadic target selection and its implications for reward modulation. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(9).

Stagg, C. J., & Nitsche, M. A. (2011). Physiological Basis of Transcranial Direct Current Stimulation [Publisher: SAGE Publications Inc STM]. *The Neuroscientist*, *17*(1).

Truong, D. Q., Magerowski, G., Blackburn, G. L., Bikson, M., & Alonso-Alonso, M. (2013). Computational modeling of transcranial direct current stimulation (tDCS) in obesity: Impact of head fat and dose guidelines. *NeuroImage. Clinical, 2.*

Urai, A. E., de Gee, J. W., Tsetsos, K., & Donner, T. H. (2019). Choice history biases subsequent evidence accumulation. *eLife, 8.*

Vergallito, A., Feroldi, S., Pisoni, A., & Romero Lauro, L. J. (2022). Inter-Individual Variability in tDCS Effects: A Narrative Review on the Contribution of Stable, Variable, and Contextual Factors. *Brain Sciences, 12*(5).

Vergallito, A., Varoli, E., Pisoni, A., Mattavelli, G., Del Mauro, L., Feroldi, S., Vallar, G., & Romero Lauro, L. J. (2023). State-dependent effectiveness of cathodal transcranial direct current stimulation on cortical excitability. *NeuroImage, 277.*

Vernet, M., Quentin, R., Chanes, L., Mitsumasu, A., Valero-Cabré, A., & Oliveira-Maia, A. J. (2014). Frontal eye field, where art thou? Anatomy, function, and non-invasive manipulation of frontal regions involved in eye movements and associated cognitive operations INTRODUCTION: FEF, A CROSS-ROADS FOR EYE MOVEMENTS AND VISUO-SPATIAL COGNITION.

Villamar, M. F., Volz, M. S., Bikson, M., Datta, A., Dasilva, A. F., & Fregni, F. (2013). Technique and considerations in the use of 4x1 ring high-definition transcranial direct current stimulation (HD-tDCS). *Journal of Visualized Experiments: JoVE,* (77).

Vöröslakos, M., Takeuchi, Y., Brinyiczki, K., Zombori, T., Oliva, A., Fernández-Ruiz, A., Kozák, G., Kincses, Z. T., Iványi, B., Buzsáki, G., & Berényi, A. (2018). Direct effects of transcranial electric stimulation on brain circuits in rats and humans. *Nature Communications, 9*(1).

Wiethoff, S., Hamada, M., & Rothwell, J. C. (2014). Variability in response to transcranial direct current stimulation of the motor cortex. *Brain Stimulation, 7*(3).

Zappasodi, F., Musumeci, G., Navarra, R., Di Lazzaro, V., Caulo, M., & Uncini, A. (2018). Safety and effects on motor cortex excitability of five cathodal transcranial direct current stimulation sessions in 25hours. *Neurophysiologie Clinique = Clinical Neurophysiology, 48*(2).