Evidence of play behavior in captive California Two-Spot Octopuses, *Octopus bimaculoides*

Katarina Jarmoluk[1] and Galit Pelled[1,2*]

[1] Department of Mechanical Engineering, Michigan State University, East Lansing, Michigan,
United States of America

[2] Department of Radiology, Michigan State University, East Lansing, Michigan, United States of
America

* Corresponding author

E-mail: pelledga@msu.edu (GP)

Abstract

Play is considered to be an essential part of development that supports learning, memory, and the development of flexible behavioral strategies. It may also serve as an informative factor in assessing an animal's welfare state and in improving care and husbandry practices. An increasing number of non-mammalian species have been discovered to engage in play behavior, including several cephalopod species. Here, we characterized play behavior in wild-caught, laboratory-housed California Two-Spot Octopuses, *Octopus bimaculoides*, a species with growing relevance as a model in biomedical research, with the goal of establishing a behavioral repertoire and encouraging further research into the behavior and welfare of this species.

Introduction

Play is an essential part of animal development, as it is known to engage neurobiological mechanisms that support learning, memory, and the development of flexible behavioral strategies. It is also important in adulthood, and both human and animal research suggests that playful behavior and internal motivation are linked to mental health [1].

When play behavior first started growing in popularity as a field of scientific study, the only non-human animals thought to be capable of it were other mammals and possibly birds [2, 3], but over the past few decades there has been growing appreciation of the importance of play in development, socialization and communication, as well as an increasing amount of evidence showing that play behavior is also found in reptiles [4, 5], fish [5-7], and even insects [8].

Octopods are soft-bodied marine mollusks of over 300 known species within the class Cephalopoda, containing two major groups: the finless incirrate octopuses and the finned cirrate octopuses. Within the taxon of incirrate octopuses is the family Octopodidae, containing the majority of octopods at over 200 species. Octopodidae are benthic creatures and are characterized by their eight arms, each possessing one to two rows of suckers [9]. The majority of an octopus' neurons are located within the arms [10, 11], giving it a decentralized nervous system. They are also capable of changing the color and texture of their skin, which they commonly use as a method of camouflage or startle display [12-14].

With their large brains, unique nervous system, and wide behavioral repertoire, octopuses are ideal models in a broad variety of fields and subjects, including aquaculture, neurobiology, climate change, and more [15]. In particular, they are becoming increasingly common as a model in biomedical and neuroscience fields, as well as in robotics [16-19].

The discussion of play in octopuses is not a new occurrence. Play or play-like behaviors have been observed and recorded in two species: the common octopus, *Octopus vulgaris* [20], and the Giant Pacific Octopuses, *Octopus dofleini* [21]; it has been anecdotally described in the Caribbean Reef Octopus, *Octopus briareus* [22]; and briefly mentioned in regards to the California Two-Spot Octopus, *Octopus bimaculoides* [23]. *O. bimaculoides* (Pickford and McConnaughey, 1949) is a species endemic to the coasts of California and Mexico's Baja Peninsula. This species is one of the most common species of "pet" octopuses and is also growing in popularity as a model for biomedical research [24]. Our current work marks the first time, to our knowledge, that

play behavior has been recorded and quantified in *O. bimaculoides*, and that video and photo
evidence have been used to characterize play in any octopus species.

Understanding octopus play may be valuable for assessing an animal's welfare state. It is
well-understood that an animal's welfare state affects the quality of scientific research. An
animal's body is affected by its state of mind [25] and inadequate welfare can cause abnormal
behavior, physiology, and immunology [26]. The most important measure for ensuring the
protection of cephalopod welfare in both laboratory and field settings is the refinement of methods
of maintenance, care, and culture [27, 28]. According to Sykes & Gestal [29], "The practice of
good (positive) or bad (negative) welfare in [cephalopod] research, maintenance, rearing or
culture conditions will determine the existence of pathologies."

Existing research shows that stress negatively affects both *O. bimaculoides* physiology
and behavior [30, 31] and may be related to stereotypic behaviors such as autophagy [32]. For
captive cephalopods, enrichment is an important management tool that can be used to overcome
the inherent stress of being held in captivity [33]. Ahloy-Dallaire et. al. [34] suggested that there
is a consistent relationship between animal welfare and most types of play. Evidence suggests
that positive affect increases play, while negative affect can significantly suppress many types of
play in both human and non-human animals. Since there have been very few deliberate records
and measurements of octopus play behavior, the association between play and welfare in the
Octopodidae family, and in *O. bimaculoides* specifically, remains to be determined. As it grows in
popularity as a research model, the importance of establishing a detailed behavioral repertoire for
O. bimaculoides increases as well.

In this paper, the behavior of *O. bimaculoides* in a lab setting was analyzed in order to
characterize play behavior within this species and to demonstrate how a deeper understanding
of play in captive octopuses may aid in improving and maintaining octopus welfare. Although the
presented findings are preliminary in nature, the authors hope that the presence of such findings
may encourage further research into the behavior and welfare of this increasingly biomedically
relevant species.

87 Methods

All procedures were approved by the Institutional Animal Care and Use Committee at
Michigan State University. Nine adult *O. bimaculoides* were obtained off the coast of southern
California by Aquatic Research Consultants then shipped overnight to the lab at Michigan State
University. Octopuses were housed separately in the lab for an average of 93 ± 43 days. Welfare
was regularly assessed using a slightly modified version of the Giant Pacific Octopus health and
welfare assessment tool developed by Holst and Miller-Morgan [35].

Each octopus was housed individually in a 40-gallon tank with a separate sump
compartment for filtration. Each tank system was isolated from the others to prevent possible
water contamination. Water parameters were maintained within the ranges recommended by
Hanlon & Forsythe, 1985, as follows: temperature: 15-20°C; salinity: 34-36 ppt; pH: 7.7-8.2;
ammonia: <0.2 mg/L; nitrite: <0.2 mg/L; and nitrate: <100.0 mg/L [36]. A reverse
osmosis/deionization filtration system provided automatic top-off water to each tank, and saltwater

was made by mixing Tropic Marin Classic Sea Salt with the RO/DI water. More detailed tank and
environment set-up and protocols are described by Van Buren et. al., 2021 [19].

All octopus tanks contained at least one den in the form of a small flowerpot or ceramic
rock cave, and some octopuses rearranged live rock or other tank items to construct their own
dens. Plastic aquarium plants, live rock, and seashells were also frequently used by the octopuses
to further conceal dens. Octopuses were either fed live fiddler crabs or frozen-thawed shrimp.
Novel enrichment items such as Legos, shells, and other live foods (such live snails, shrimp, and
other species of crabs) were provided on a rotating basis to prevent habituation and boredom.
Feeding the octopus shrimp in a test tube was introduced as a method of enrichment. To train the
octopuses to open the test tube by unscrewing the cap, octopuses were first presented with the
test tube containing a piece of shrimp without the cap. If the octopuses removed the shrimp from
the uncapped test tube successfully, at the next feeding the cap would be screwed on loosely. As
they succeeded in opening the loosely capped tube, the cap would be gradually tightened in
subsequent feedings to increase the difficulty of opening it. This process continued with each
feeding until the octopuses could remove a cap that was fully screwed on.

To monitor the octopuses, Wyze Cam V2 and V3 cameras were set up facing the front of
each tank. The cameras operated 24 hours a day and recorded all motion events to a micro-SD
card. In addition, the Wyze phone app allowed for manual recording of live videos to the user's
phone using a start/stop recording function. Videos and pictures taken with portable video
cameras were also analyzed.

Burghardt's [7] five criteria for play were used to determine whether observed behaviors
constituted play. Briefly, these criteria are as follows: 1) there is no immediate function to the
behavior in the context in which it is performed; 2) the behavior is voluntary, spontaneous, or
autotelic (done for its own sake); 3) there is a structural or temporal difference from other typical
behavior; 4) the behavior is repeated but not stereotyped; and 5) the animal is healthy and free
of stress and competition.

Additional criteria was used to determine the intensity of play behavior based on the "Play
Intensity Scale" described by Kuba et al., 2006 [20]. Holding the object close to the mouth was
classified as level 0 behavior, exploring the object with the arms as level 1 behavior, and
completing one release-grasp sequence (comparative to the push/pull sequence described by
Kuba et al., 2006) was classified as level 2 behavior. Level 3 behavior, which was characterized
by 2-5 release-grasp sequences, was considered "play-like," and level 4 "play" behavior consisted
of the release-grasp sequence being repeated over 5 times. Release-grasp sequences were only
counted when they were performed continually.

Results

136   Of the nine octopuses, only three (octopuses E, F, and H) learned to unscrew the test tube
137   cap to access the shrimp within. The characteristics of all nine octopuses are shown in Table 1.
138   Because the test tube cap was not presented to octopuses that had not learned to unscrew it,
139   only these three octopuses had access to the free-floating test tube cap.

Octopus ID	Sex	Average Weight (g)	Opened Test Tube Cap
A	F	229.5	no
B	M	172.0	no
C	F	213.0	no
D	F	153.3	no
E	M	128.2	yes
F	M	220.8	yes
G	M	137.5	no
H	F	170.5	yes
I	F	147.8	no

Table 1. Characteristics of the nine octopuses housed in the lab. F = female; M = male.

In accordance with the Play Intensity Scale discussed in the methods, all three octopuses
who had access to the free-floating cap engaged in level 4 object play with it. None of the
octopuses, including those that were only given the test tube without the cap, chose to play with
the test tube itself despite its similar floating properties. For all three octopuses, play consisted of
the octopus releasing the cap from its grasp so that it floated upwards into the water current of
the tank. As it began to float past, the octopus would reach out to grab it again, pull it in close to
their body, then release it again (Figure 1). A video capturing one instance of octopus H playing
can be found in the supplemental date (S1 Video).

Fig 1. Octopus play sequence with test tube cap. An adult *O. bimaculoides* (Octopus F) grabs a test tube cap with his arms (a), pulls the cap towards his body (b), then releases it into the tank current (c) and reaches to grab it again as it moves away (d, e). The blue cap is indicated by a green arrow in pictures a-d, and is not visible in picture e. This release-grasp sequence was observed in this individual octopus on five separate occasions throughout the four months it was housed in the lab.

The arms used to grasp and release the test tube cap were primarily the front two arms on either side (R1, L1, R2, L2). In some cases, such as when an octopus was using those arms to hold onto the test tube still, other arms were used (R3 and R4). In all octopuses, at least two of the rear arms were used as an anchoring point to a surface in the tank during play, such as the substrate, live rock, or the inside of the den.

Octopus ID	Number of Times Observed Playing	Skin Patterns During Play	Arms Used
E	1	Acute mottle	L1, L2, R1, R2
F	5	Acute mottle	L1, L2, L3, R1, R2, R3
H	2	Uniform pale brown phase	L1, R1, R2, R3, R4

Table 2. Characteristics of play behavior with test tube caps in octopuses E, F, and H.

Discussion

This work demonstrates that *O. bimaculoides* engages in play behavior, a behavior that appears to satisfy intellectual and emotional needs. Elucidating this behavior is important to understand the role of play in development and as a tool to assess animals' wellbeing. These findings also invite further research into how play behavior may vary between octopus species, or into what factors may have contributed to the occurrence of play in individual octopuses, such as life stage, sex, or personality. Mather and Anderson [37] determined that octopuses (*Octopus rubescens*) have individual personalities, characterized by three factors: activity, reactivity, and avoidance. These personality traits may influence the likelihood of an octopus to engage in play activity. The octopuses that displayed play behavior were often active during daytime hours and sought to engage handlers during daily tank maintenance. Comparatively, the octopuses that frequently withdrew from handlers and remained in their dens did not show any signs of play behavior. There is a possibility that the difference in personality between the octopuses housed in the lab could have influenced the likelihood of an individual octopus both to learn how to open the test tube, and to engage in object play with the plastic cap.

In conclusion, the behavior of these *O. bimaculoides* is consistent with previously established criteria for animal play. These findings are also aligned with previously established object play behavior in other octopus species. While there is not enough data here to make any generalized conclusions about the overall correlation between octopus play and welfare, the three animals presented here were determined to be in positive welfare states, and this paper provides preliminary evidence for play behavior in *O. bimaculoides* that may inspire future research. The

full characterization of behavior in a species that is increasingly popular in captive settings is an essential aspect to be understood in order to ensure proper care and husbandry methods. A controlled experiment analyzing the play behavior of *O.* bimaculoides, what husbandry factors may impact the occurrence of play, and how the presence or absence thereof correlates with the individual's welfare state could assist greatly in the development of more detailed and comprehensive standards of care for these animals.

References

1.   Blois-Heulin C, Rochais C, Camus S, Fureix C, Lemasson A, Lunel C, et al. Animal Welfare: Could Adult Play be a False Friend? 2015;2(2):156-85.

2.   Fagen R. Animal play behavior. Oxford, England: Oxford University Press; 1981. 684 p.

3.   Beach FA. Current Concepts of Play in Animals. The American Naturalist. 1945;79(785):523-41.

4.   Barabanov V, Gulimova V, Berdiev R, Saveliev S. Object play in thick-toed geckos during a space experiment. Journal of Ethology. 2015;33(2):109-15. doi: 10.1007/s10164-015-0426-8.

5.   Dinets V. Play behavior in ectothermic vertebrates. Neuroscience & Biobehavioral Reviews. 2023;155:105428. doi: https://doi.org/10.1016/j.neubiorev.2023.105428.

6.   Burghardt GM, Dinets V, Murphy JB. Highly Repetitive Object Play in a Cichlid Fish (Tropheus duboisi). Ethology. 2015;121(1):38-44. doi: https://doi.org/10.1111/eth.12312.

7.   Burghardt GM. The genesis of animal play : testing the limits. Cambridge, Mass.: MIT Press; 2005. xvi, 501 p. p.

8.   Dapporto L, Turillazzi S, Palagi E. Dominance Interactions in Young Adult Paper Wasp (Polistes dominulus) Foundresses: A Playlike Behavior? Journal of comparative psychology (Washington, DC : 1983). 2006;120:394-400. doi: 10.1037/0735-7036.120.4.394.

9.   Jereb P, Roper C, Norman M, Finn J. Cephalopods of the world. An annotated and illustrated catalogue of cephalopod species known to date. Volume 3. Octopods and Vampire Squids. FAO species catalogue for fishery purposes. 2014;4(3):370.

10.   Young JZ. THE NUMBER AND SIZES OF NERVE CELLS IN OCTOPUS. Proceedings of the Zoological Society of London. 1963;140(2):229-54.

11.   Young JZ. The anatomy of the nervous system of Octopus vulgaris. Oxford: Clarendon Press; 1971.

12.   Packard A, Hochberg FG, Nixon M, Messenger JB, editors. Skin patterning in Octopus and other genera1977.

13.   How MJ, Norman MD, Finn J, Chung W-S, Marshall NJ. Dynamic Skin Patterns in Cephalopods. Frontiers in Physiology. 2017;8. doi: 10.3389/fphys.2017.00393.

14.   Forsythe JW, Hanlon RT. Behavior, Body Patterning and Reproductive-Biology of Octopus-Bimaculoides from California. Malacologia. 1988;29(1):41-55. PubMed PMID: WOS:A1988P041200006.

15.   O'Brien CE, Roumbedakis K, Winkelmann IE. The Current State of Cephalopod Science and Perspectives on the Most Critical Challenges Ahead From Three Early-Career Researchers. Front Physiol. 2018;9:700. Epub 20180606. doi: 10.3389/fphys.2018.00700. PubMed PMID: 29962956; PubMed Central PMCID: PMCPMC6014164.

16.   Di Cosmo A, Pinelli C, Scandurra A, Aria M, D'Aniello B. Research Trends in Octopus Biological Studies. Animals (Basel). 2021;11(6). Epub 20210617. doi: 10.3390/ani11061808. PubMed PMID: 34204419; PubMed Central PMCID: PMCPMC8233767.

17.   Sumbre G, Gutfreund Y, Fiorito G, Flash T, Hochner B. Control of Octopus Arm Extension
by a Peripheral Motor Program. Science. 2001;293(5536):1845-8. doi: 10.1126/science.1060976.
18.   Gedela NSS, Radawiec RD, Salim S, Richie J, Chestek C, Draelos A, et al. In vivo
electrophysiology recordings and computational modeling can predict octopus arm movement.
Bioelectron Med. 2025;11(1):4. Epub 20250214. doi: 10.1186/s42234-025-00166-9. PubMed
PMID: 39948616; PubMed Central PMCID: PMCPMC11827351.
19.   VanBuren T, Cywiak C, Telgkamp P, Mallett CL, Pelled G. Establishing an Octopus
Ecosystem for Biomedical and Bioengineering Research. J Vis Exp. 2021;(175). Epub 2021/10/12.
doi: 10.3791/62705. PubMed PMID: 34633382.
20.   Kuba MJ, Byrne RA, Meisel DV, Mather JA. When do octopuses play? Effects of repeated
testing, object type, age, and food deprivation on object play in Octopus vulgaris. J Comp Psychol.
2006;120(3):184-90. doi: 10.1037/0735-7036.120.3.184. PubMed PMID: 16893255.
21.   Mather JA, Anderson RC. Exploration, play and habituation in octopuses (Octopus dofleini).
Journal of Comparative Psychology. 1999;113(3):333-8. doi: 10.1037/0735-7036.113.3.333.
22.   Wood JB, Wood D. Enrichment for an advanced invertebrate. The shape of Enrichment.
1999;8(3):1-5.
23.   Zylinski S. Fun and play in invertebrates. Curr Biol. 2015;25(1):R10-2. doi:
10.1016/j.cub.2014.09.068. PubMed PMID: 25562291.
24.   VanBuren T, Cywiak C, Telgkamp P, Mallett CL, Pelled G. Establishing an Octopus
Ecosystem for Biomedical and Bioengineering Research. JoVE. 2021;(175):e62705. doi:
doi:10.3791/62705.
25.   Poole T. Happy animals make good science. Lab Anim. 1997;31(2):116-24. doi:
10.1258/002367797780600198. PubMed PMID: 9175008.
26.   Prescott MJ, Lidster K. Improving quality of science through better animal welfare: the
NC3Rs strategy. Lab Anim (NY). 2017;46(4):152-6. doi: 10.1038/laban.1217. PubMed PMID:
28328893.
27.   Moltschaniwskyj NA, Hall K, Lipinski MR, Marian JEAR, Nishiguchi M, Sakai M, et al. Ethical
and welfare considerations when using cephalopods as experimental animals. 2007;17(2):455-76.
PubMed PMID: Moltschaniwskyj2007.
28.   Mather JA, Anderson RC. Ethics and invertebrates: a cephalopod perspective. Diseases of
aquatic organisms. 2007;75(2):119-29.
29.   Sykes A, Gestal C. Welfare and Diseases Under Culture Conditions. 2014. p. 97-112.
30.   Bennett H, Toll RB. Intramantle inking: a stress behavior in Octopus bimaculoides
(Mollusca: Cephalopoda). J Am Assoc Lab Anim Sci. 2011;50(6):943-5. PubMed PMID:
22330791; PubMed Central PMCID: PMCPMC3228935.
31.   Chancellor S, Grasse B, Sakmar T, Scheel D, Brown JS, Santymire RM. Exploring the Effect
of Age on the Reproductive and Stress Physiology of Octopus bimaculoides Using Dermal
Hormones. Animals (Basel). 2023;13(19). Epub 20231006. doi: 10.3390/ani13193115. PubMed
PMID: 37835721; PubMed Central PMCID: PMCPMC10571824.
32.   Reimschuessel R, Stoskopf MK. Octopus automutilation syndrome. J Invertebr Pathol.
1990;55(3):394-400. doi: 10.1016/0022-2011(90)90083-i. PubMed PMID: 2351844.
33.   Vidal EAG, Villanueva R, Andrade JP, Gleadall IG, Iglesias J, Koueta N, et al. Chapter One -
Cephalopod Culture: Current Status of Main Biological Models and Research Priorities. Advances
in Cephalopod Science: Biology, Ecology, Cultivation and Fisheries. 67: Academic Press; 2014. p.
1-98.
34.   Ahloy-Dallaire J, Espinosa J, Mason G. Play and optimal welfare: Does play indicate the
presence of positive affective states? Behavioural Processes. 2018;156:3-15. doi:
https://doi.org/10.1016/j.beproc.2017.11.011.

35.   Holst MM, Miller-Morgan T. The Use of a Species-Specific Health and Welfare Assessment
Tool for the Giant Pacific Octopus,. J Appl Anim Welf Sci. 2021;24(3):272-91. Epub 20200916. doi:
10.1080/10888705.2020.1809412. PubMed PMID: 32937082.
36.   Hanlon RT, Forsythe JW. Advances in the laboratory culture of octopuses for biomedical
research. Lab Anim Sci. 1985;35(1):33-40. PubMed PMID: 3981958.
37.   Mather J, Anderson RC. Personalities of octopuses (Octopus rubescens). Journal of
Comparative Psychology. 1993;107:336-40.

S1 Video. *O. bimaculoides* playing with test tube cap.