Increased Postprandial Metabolic Flexibility is Associated with Higher Body Fat Percentages in Healthy Young Adults

Nicholas A. Foreman[1], Sahil Rajwade[1], Jaiden Bluth[1], Lauren C. Skoglund[1], Audrey M. Letts[1], Loretta DiPietro[1], Adam Ciarleglio[2], Matthew D. Barberio[1].

[1]Department of Exercise and Nutrition Sciences, The Milken Institute School of Public Health, George Washington University, Washington D.C.

[2]Department of Biostatistics, The Milken Institute School of Public Health, George Washington University, Washington D.C.

Corresponding Author

Matthew D. Barberio
Assistant Professor
Department of Exercise and Nutrition Sciences
The Milken Institute School of Public Health
George Washington University
Washington, D.C.
Email: mbarberio@gwu.edu
Phone: 202-994-1408

Abstract

PURPOSE: High levels of adiposity are associated with an increased risk for chronic cardiometabolic disease. Adipose tissue is a critical metabolic tissue during the postprandial (feeding) period. We sought to assess the relationship between body composition (body fat percentage) and measures of postprandial metabolic flexibility in otherwise healthy young adults. **METHODS:** Young adults (n = 27, n = 15 females) of varying body compositions (BMI = 27.1 ± 4.5; Body Fat (BF)% = 30.4 ± 8.7) but without overt pathology completed a 100g oral glucose tolerance test (OGTT). Indirect calorimetry was conducted before (fasting) and following (30, 60, 90, 120 min) consumption of the glucose beverage. Serum and plasma were collected at corresponding time points and analyzed for blood glucose, insulin, non-esterified fatty acids, and inflammatory markers. Linear mixed models were used to test for the effect of body composition on postprandial metabolism over time while controlling for sex, age, fasting metabolism, measures of insulin sensitivity, visceral adipose tissue volume, and the repeated measures design. **RESULTS:** As expected, there was a significant ($p < 0.05$) increase in blood glucose, serum insulin, RER, and CHO with concomitant decreases in plasma NEFAs and whole-body FOX during the OGTT. Using mixed effects models, BF% modified the relationship between postprandial RER and time (β = -0.133, p = 0.019), as well as postprandial CHO and time (β = -0.134, p = 0.023) such that individuals with higher BF% increase their RER and CHO at a greater rate and to a greater extent than those with lower BF%. There was no evidence that BF% modifies the relationship between FOX and time. **CONCLUSION:** We demonstrate that a higher body fat percentage is associated with greater postprandial metabolic flexibility (RER and CHO) during an OGTT in young adults without overt metabolic pathology. Metabolic flexibility in this sample may be preserved and represents a compensatory adaptation to decreased glucose storage in the anabolic, postprandial period.

Introduction

Metabolic flexibility is described as the appropriate shift in substrate oxidation based upon substrate availability (e.g. fasting to fed) or metabolic demand (e.g. exercise). It was originally described as changes in the respiratory quotient (RQ) across the leg limb from rest to insulin stimulation during the euglycemic-hyper-insulinemic clamp (EHC).[1, 2] In the two decades since the concept of metabolic flexibility was described in humans, several methodological approaches to assess metabolic flexibility have been utilized, including indirect calorimetry during the EHC, during oral glucose tolerance tests (OGTT), and during mixed-meal feeding challenges (MMC).[1, 3] Individuals with overt metabolic pathologies, such as type 2 diabetes and insulin resistance, as well as individuals classified as obese based on BMI, have been described as metabolically inflexible.[1] That is, these individuals showed blunted (reduced) changes in the respiratory exchange ratio (RER) during metabolic challenges such as EHC and feeding.

In the absence of overt metabolic pathology or metabolic dysfunction, transitioning from the fasted to the fed state involves shifting away from aerobic fatty acid oxidation to aerobic glucose oxidation and, to a lesser degree, glycolytic energy production.[4] This shift results in an anabolic environment, in which exogenous energy is stored, as compared to the catabolic environment of fasting. Elevated insulin, which is the primary driver of the metabolic shift following feeding, suppresses adipose tissue lipolysis and promotes the uptake, storage, and oxidation of exogenous macronutrients.[4] Adipose tissue serves a critical role in buffering fatty acid flux in the circulation during this postprandial period; it is also a site of significant glucose uptake for lipogenesis and has been identified as a central regulator of metabolic flexibility.[2, 4]

The presence of obesity and increased fat mass is believed to negatively impact metabolic flexibility.[5, 6] Individuals with obesity have higher resting RER indicating a preference for higher carbohydrate oxidation in the fasting state while adipose tissue lipolysis is more resistant to insulin-induced suppression in feeding. However, studies have not thoroughly

analyzed the effect of adiposity on postprandial metabolic flexibility. By dividing participants into

high vs. low metabolic flexibility, Sparks et al. (2009) demonstrated that healthy young males

with low metabolic flexibility during insulin stimulation (EHC) had significantly higher body fat

percentage and larger fat cell size.[6] Follow-up analysis in this cohort demonstrated that females

were more metabolically flexible than their male counterparts despite significantly higher body

fat percentages.[5] Conversely, individuals classified as having high metabolic flexibility to an

OGTT had similar body fat percentages but a larger subcutaneous adipose tissue (SAT) as

compared to those described as having low metabolic flexibility.[7] This discrepancy was

highlighted in a recent systematic review by Glaves et al. (2021), who found inconsistent and

unclear relationships between adipose tissue volume, function, and characteristics in studies

using EHC, OGTTs, and MMCs.[3]

The primary objective of this study was to thoroughly assess the relationship between

adiposity and postprandial metabolic flexibility during an OGTT. We recruited 27 healthy male

and female young adults (18 - 40 years old) with body mass index (BMI) between 18.5-39.9.

Using indirect calorimetry, we assessed postprandial metabolic flexibility (RER, Carbohydrate

Oxidation [CHO], and Fat Oxidation [FOX]) during an OGTT. We hypothesized that individuals

with higher body fat percentages would demonstrate significantly reduced postprandial

metabolic flexibility (respiratory exchange ratio and carbohydrate oxidation) throughout the

OGTT.

METHODS

Participants

49
We performed a cross-sectional, experimental study from January 2023 to January 2024.

Potential participants were recruited and screened based on sex, body mass index (BMI)

classification, and lack of known clinical pathology from the George Washington University

campus and surrounding metropolitan area. Participants were required to be between 18 and 40

years old, have a BMI between 18.5 and 39.9, be free of clinically diagnosed cardiometabolic

disease, and not be on prescription medications for hypertension, blood lipids, or blood glucose

homeostasis. Participants who were pregnant or taking medications for cardiometabolic disease

were excluded. Recruitment was stratified by BMI classification to yield a across the BMI

continium. In response to flyers and emails, fifty-two participants were screened for

participation. Of these, thirty-four were enrolled in this study; seven participants did not

complete the study (3 due to scheduling, 3 due to loss to follow-up, 1 due to equipment

malfunction), giving complete data for 27 participants. Participant characteristics are reported in

Table 1. All study procedures were approved by the George Washington University IRB (IRB

#202737). Prior to enrollment, participants read and signed an IRB-approved informed consent

and all study procedures were conducted in accordance with the Declaration of Helsinki.

Experimental Design

66
67   Participants reported to the Metabolic and Exercise Testing Laboratories in the morning

68   between 0800 and 0900 on two separate days. Each visit was separated by at least 48 hours.

69   At the first visit, participants completed written questionnaires for medical and physical activity

70   history, assessment of body composition, and cardiorespiratory fitness. Participants returned to

71   the lab for assessment of postprandial substrate metabolism following an OGTT.

72   For each visit, participants were instructed to arrive after 10 hours of fasting and to

73   abstain from caffeine consumption for 12 hours and from any recreational drug use or structured

74   exercise for 24 hours. Upon arrival, hydration was measured using a portable refractometer. If

75   urine specific gravity was above 1.02, participants were given eight ounces of water. Body

76   weight was measured using a mechanical column scale (Seca 700; Hamburg, Germany).

Table 1. Participant Anthropometric Characteristics

Variable	All (n = 27)	Female (n = 15)	Male (n = 12)	p-value (Sex)	Association with BF%
NIH Race/Ethnicity (%)				0.902	---
Black	7 (25.9%)	4 (26.7%)	3 (25.0%)	---	---
Hispanic or Latino	1 (3.7%)	0 (0.0%)	1 (8.3%)	---	---
Unknown	1 (3.7%)	1 (6.7%)	0 (0.0%)	---	---
White	18 (66.7%)	10 (66.7%)	8 (66.7%)	---	---
Age (years)	22 [19, 27]	22 [19, 24]	25 [21, 28]	0.138	r=0.081; p=0.688
Height (m)	1.7 (0.09)	1.6 (0.05)	1.8 (0.05)	**<0.001**	**r=-0.415; p=0.032**
Weight (kg)	73.0 [66.6, 90.0]	72.4 [63.5, 87.7]	76.8 [71.1, 100.0]	0.075	r=0.257; p=0.195
BMI (kg/m^2)	27.1 (4.5)	27.7 (4.7)	26.4 (4.4)	0.474	**r=0.582; p=0.001**
Fat Mass (kg)	24.2 (9.6)	26.3 (8.6)	21.7 (10.6)	0.221	**r=0.854; p<0.001**
Fat Free Mass (kg)	51.7 (10.9)	45.2 (6.3)	59.7 (10)	**<0.001**	r=-0.351; p=0.072
Body Fat (%)	30.4 (8.7)	34.9 (6.3)	24.7 (8.1)	**0.001**	---
VAT Volume (in^3)	2.7 (1.3)	2.2 (1.3)	3.3 (1.0)	**0.035**	r=0.301; p=0.127
RMR (kcal/d)	1709.8 (287.9)	1538.2 (222.3)	1924.2 (206.8)	**<0.001**	r=-0.25; p=0.208
RER	0.80 (0.06)	0.78 (0.05)	0.82 (0.06)	0.056	r=-0.129; p=0.520
Carbohydrate Oxidation (g/min)	0.10 (0.07)	0.07 (0.04)	0.14 (0.08)	**0.005**	r=-0.16; p=0.424
Lipid Oxidation (g/min)	0.08 (0.02)	0.08 (0.03)	0.08 (0.02)	0.91	r=-0.031; p=0.878
VO$_{2PEAK}$ (mL/kg/min)	34.1 (8.8)	31.4 (7.3)	37.5 (9.7)	0.075	**r=-0.784; p<0.001**
VO$_{2PEAK}$ (L/min)	2.6 (0.6)	2.3 (0.37)	3.1 (0.53)	**<0.001**	**r=-0.66; p<0.001**

Race/ethnicity is reported as count (percent). Numerical data are mean (SD) or median [IQR] based on normality. RMR = Resting metabolic rate, RER = Respiratory Exchange Ratio

81
82
83 *Body mass and body composition*
84
Height was measured using a wall-mounted stadiometer (Seca 216; Hamburg,

Germany) to the nearest 0.1 centimeters. Weight was measured using the digital scale as part

of the bioelectrical impedance analysis (InBody 770; Seoul, South Korea). Dual X-ray

absorptiometry (GE Lunar iDXA; GE Healthcare, Chicago, IL, USA) with manufacturer software

(enCORE Forma version 15) was used to measure body fat percentage, fat-free mass (FFM),

fat mass, and visceral adipose tissue (VAT) volume. Before the measurement, participants

removed all jewelry and were instructed to wear clothing free of metal zippers or reflective

surfaces.

Exercise Testing
94
Following body composition assessment, participants completed a graded exercise test

on a cycle ergometer (Monark LC6 novo; Varberg, Sweden) to volitional exhaustion. During

cycling, expired gases were measured using a breath-by-breath metabolic cart (Cosmed Quark

CPET; Rome, Italy), and heart rate was monitored using a chest strap (Polar H10; Kempele,

Finland). All participants completed three 180-second stages of 25, 50, and 75W each before

resistance increased by 25W per minute. The test continued until the participant was unable to

continue cycling or the investigator observed a plateau in oxygen consumption (VO_2). Breath-

by-breath data were averaged into five-second bins. Because not every participant satisfied the

criteria for VO2max,[8] the highest 15-second value is reported as VO2peak.

104
105   *Oral Glucose Tolerance Test with Indirect Calorimetry and Blood Sampling*
106
On a separate day, participants arrived following the same pre-testing guidelines

outlined above. Assessment of body weight and hydration were performed, and the participant

rested in a reclined chair in a dimly lit room for at least 10-15 minutes. A venous catheter was

placed in an antecubital vein and fasting whole blood was sampled for plasma (K-EDTA) and

serum (Serum Separator Tube); saline was used to keep the catheter patent between blood

draws. After the resting blood draw, indirect calorimetry was conducted via the canopy method

(Parvo Medics TrueOne 2400; Salt Lake City, Utah) while the participant lay supine on a

padded table for determination of fasting metabolic measures. Following this, subjects

consumed a 100g glucose beverage (GlucoCrush Cardinal Health; Dublin, OH) and were

instructed to finish consumption of the drink within 5 minutes. After finishing the drink, subjects

returned to the supine position for the remainder of the test. Indirect calorimetry was used to

assess postprandial substrate oxidation at the following postprandial intervals (min): 15 – 30, 45

– 60, 75 – 90, 105 – 120.

Determination of substrate oxidation and metabolic rate
121
For each 15-minute indirect calorimetry measurement, expired gases were collected from a

mixing chamber at 5-second intervals, and the final 8 minutes were used for analysis.

Respiratory exchange ratio (RER) was calculated as the ratio between exhaled carbon dioxide

(VCO_2) and VO_2. Metabolic rate and rates of carbohydrate (CHO) and fat oxidation (FOX) were

calculated using published equations for fat oxidation = [VO_2 (L/min) x 1.67) – (VCO_2 (L/min) x

1.67)] and carbohydrate oxidation = [VCO_2 (L/min) x 4.55) – (VO_2 (L/min) x 3.21)].[9] For

metabolic rate calculations, urinary nitrogen excretion was assumed to be zero.

Laboratory measurements and calculations

Fasting and postprandial blood glucose were determined in fresh whole blood

(OneTouch Ultra, Lifescan, Inc.; Milpitas, CA). Fasting and postprandial serum samples were

used to measure insulin (Insulin Human ELISA; Invitrogen Inc. CA), and non-esterified fatty

acids (NEFA; LabAssay NEFA; FujiFilm; Richmond, VA). Fasting serum samples were used to

determine C-reactive protein (CRP; CRP Human ELISA; Invitrogen Inc., CA), leptin (Leptin

Human ELISA; Invitrogen Inc., CA), and adiponectin (Adiponectin human ELISA; Adipogen Life

Science; San Diego, CA). Manufacturer instructions were followed for conducting and analyzing

all assays. Manufacturer internal controls were assessed along with coefficients of variations for

biological replicates. Samples whose performance was assessed to be outside of manufacturer

recommendations were re-analyzed. Fresh whole blood (fasting) was used to determine

HbA1C% (PTS Diagnostics; Whitestown, IN) and blood lipid profiles (CardioChek; PTS

Diagnostics; Whitestown, IN) via point-of-care analyzers

Matsuda Index was calculated as follows: 10,000/√ [fasting glucose (mg/dl) × fasting

insulin (μIU/mL)] × [mean glucose (mg/dl) × mean insulin (μIU/mL) during OGTT before

normalizing to the 100g dose.[10] Adipose tissue insulin resistance (Adip-IR) was calculated using

fasting values ADIP-IR (fasting) = fasting insulin (μIU/mL) * fasting NEFA (mEq/L) and

postprandial (30, 60, 90, 120) values ADIP-IR (OGTT) = mean plasma NEFA (mEq/L) x mean

serum insulin (μIU/mL).[11] Fasting insulin and glucose measurements were used to calculate the

homeostatic model assessment of insulin resistance (HOMA-IR) = [fasting insulin (μIU/mL) -

(fasting glucose (mg/dl)/18.156)] / 22.5.[12]

Statistical Analyses

151
We report mean (sd) or median (interquartile range, IQR) both overall and by sex for

anthropometric (**Table 1**) and clinical (**Table 2**) measures and report frequency (percent) for

race. We conducted two-sample t-tests or Mann-Whitney tests to compare numerical variables

between the sexes for normally and non-normally distributed variables, respectively. We

compared race categories between the sexes using Fisher's exact test. We computed and

tested the association between each of the clinical variables and percent body fat (BF%) using

Pearson correlation coefficients. To describe the change in each postprandial outcome variable

during the OGTT (i.e., RER, CHO, FOX, glucose, insulin, and NEFA), we used one-way,

repeated measures analysis of variance (ANOVA) with time modeled as a categorical variable.

Contrasts were set *a priori* as the comparison from baseline to each postprandial timepoint (i.e.,

0 vs. 30, 0 vs. 60, etc.) with a Sidak adjustment for multiple comparisons. Separately, linear

mixed-effects models were used to determine the effect of body composition on postprandial

metabolism. For each postprandial outcome variable, we included fixed effects for time as a

continuous variable, percent body fat (BF%), time x BF% interaction, sex, age, Matsuda index,

AdipIR, visceral adipose tissue volume, and the fasted outcome measure. A subject-specific

random intercept term was included to account for repeated measures. To account for the

possible nonlinear effect of time on postprandial metabolism (i.e., a nonlinear change in

metabolism), models with an additional quadratic time term and quadratic time x BF% were

considered. Models with and without the quadratic time terms were compared using likelihood

ratio tests to determine if the quadratic terms should be retained. Variables in the models were

tested for normality with the Shapiro-Wilk test and transformed. As a result, insulin and NEFA

concentrations were transformed via natural log, and VAT volume was transformed via cube

root. Each variable included in the models was standardized to a mean of 0 and standard

deviation of 1, and time was divided by 30 to correct scaling. Fasting and postprandial variables

were standardized separately due to the inclusion of the fasting state as a covariate in the

models. All analyses were conducted in R (version 4.2.2). The lmerTest[13] and performance[14]

packages were used for model fitting and checking of relevant assumptions, and the emmeans

package[15] was used to visualize continuous interactions. For the interpretation of linear mixed-

effects models, we assess significance using both the 95% confidence interval (e.g., does it

cross 0) and the p-value (p < 0.05). Due to ambiguity in the determination of p-values for these

models[16], we deferred to the confidence interval when there was disagreement between the

confidence intervals and p-values.

RESULTS

There were 27 young adults (15 (55.6%) female) with no previously diagnosed

metabolic pathology. Body composition was heterogeneous across the sample, as indicated by

the wide interquartile range, suggesting our sample is sufficient to assess the relationship

between body composition and postprandial metabolic flexibility. Compared to females, males

had higher resting carbohydrate oxidation (p = 0.005) and resting metabolic rate (p < 0.001)

without differences in resting lipid oxidation (**Table 1**). Absolute VO_{2PEAK} was higher (p <0.001)

in males than females with no significant differences in relative VO_{2PEAK} (p = 0.075).

Triglycerides were higher in males than females (p = 0.026), and leptin was higher in females

than males (p<0.001) (**Table 2**). All other analyzed clinical measures were similar between the

sexes.

Table 2. Participant Clinical Data

Variable	All (n = 27)	Female (n = 15)	Male (n = 12)	p-value (Sex)	Association with BF%
Blood Glucose (mg/dL)	100 (8)	101 (10)	99 (7)	0.580	r=0.036; p=0.858
Serum Insulin (µIU/mL)	2.5 (0.68)	2.6 (0.56)	2.2 (0.79)	0.173	**r=0.427; p=0.037**
NEFA (mEq/L)	-1.01 (0.52)	-0.93 (0.6)	-1.12 (0.4)	0.383	r=0.371; p=0.068
HbA1c (%)	5.1 [4.8, 5.25]	5.0 [4.8, 5.2]	5.15 [5.03, 5.3]	0.432	r=-0.221; p=0.269
Total Cholesterol (mg/dL)	160.4 (33.9)	156.2 (30.1)	166.2 (39.5)	0.489	r=0.198; p=0.353
HDL (mg/dL)	51.0 [47.8, 59.5]	50.0 [45.2, 67.0]	51.5 [50.0, 56.5]	0.725	r=0.086; p=0.691
LDL (mg/dL)	92.3 (29.6)	88.6 (22.7)	97.3 (37.9)	0.517	r=0.085; p=0.715
Triglycerides (mg/dL)	74.5 [66.2, 93.0]	70.5 [64.8, 76.0]	94.5 [77.2, 119.5]	**0.026**	r=-0.016; p=0.940
Matsuda Index	4.24 [2.58, 6.00]	3.00 [2.48, 4.70]	5.96 [4.14, 6.67]	0.074	**r=-0.616; p=0.001**
HOMA-IR	3.4 (2.0)	3.8 (1.8)	2.9 (2.3)	0.277	r=0.398; p=0.054
AdipIR (OGTT)	7.7 [4.4, 12.2]	8.8 [6.2, 13.8]	6.5 [3.5, 8.8]	0.259	**r=0.481; p=0.017**
AdipIR (fasting)	5.2 [3.0, 6.9]	5.8 [5.2, 12.4]	3.4 [2.4, 4.0]	0.056	**r=0.443; p=0.030**
Adiponectin (ng/mL)	4613 [3955, 6013]	4798 [4145, 6847]	4175 [3639, 5433]	0.291	r=-0.151; p=0.470
Leptin (pg/mL)	118.1 [53.8, 148.5]	143.5 [114.7, 254.0]	45.2 [30.6, 88.1]	**<0.001**	**r=0.817; p<0.001**
CRP (mg/L)	0.48 [0.26, 1.20]	0.36 [0.27, 0.68]	0.64 [0.32, 1.60]	0.291	r=0.091; p=0.664

Data are mean (SD) or median [IQR], depending on normality. VAT volume was adjusted with a cube root transformation. Insulin and NEFA concentrations were transformed via natural log.

Substrate Response to the OGTT

Individual and aggregate data from the OGTT is presented by sex for outcome

measures in the blood (**Figure 1**) and whole-body substrate oxidation (**Figure 2**); all aggregate

values are available in **Supplementary Table 1**. During the OGTT, one-way repeated

measures ANOVAs indicated increases in blood glucose, serum insulin, RER, and CHO with

concomitant decreases in plasma NEFAs and whole-body FOX. Blood glucose was elevated

above fasting levels (99.7 ± 8.5 mg/dL) at 30 (139.9 ± 21.0; $p<0.001$), 60 (124.9 ± 29.6;

$p<0.001$), and 90 (115.3 ± 25.5; $p = 0.007$) minutes but not 120 minutes (109 ± 21.7; $p = 0.214$).

Compared to fasting insulin (14.2 ± 8.1 µIU/mL), postprandial insulin was elevated at 30 (79.9 ±

75.1), 60 (77.9 ± 54.2), 90 (63.2 ± 47.7), and 120 (55.0 ± 36.4) minutes (all $p<0.001$). NEFAs

were suppressed compared to fasting (0.41 ± 0.19 mEq/L) at 30 (0.28 ± 0.13), 60 (0.15 ± 0.08),

90 (0.15 ± 0.08), and 120 (0.09 ± 0.04) minutes (all $p<0.001$). RER was not different from

fasting (0.795 ± 0.055) at 30 minutes (0.806 ± 0.052; $p = 0.493$) but was significantly higher at

60 (0.832 ± 0.051; $p<0.001$), 90 (0.839 ± 0.044; $p<0.001$), and 120 (0.845 ± 0.048; $p<0.001$)

minutes. Similarly, CHO was not different from fasting (0.102 ± 0.071 g/min) at 30 minutes

(0.125 ± 0.071; $p = 0.083$) but was significantly higher at 60 (0.157 ± 0.063; $p<0.001$), 90 (0.160

± 0.049; $p<0.001$), and 120 (0.166 ± 0.061; $p<0.001$) minutes. FOX was unchanged from fasting

(0.084 ± 0.024) at 30 (0.086 ± 0.024; $p = 0.927$) and 60 (0.078 ± 0.027; $p = 0.383$) minutes. At

90 (0.073 ± 0.026; $p = 0.011$) and 120 minutes (0.068 ± 0.023; $p<0.001$), FOX was suppressed

below fasting levels.

Figure 1. Substrate (Glucose: Left column; NEFA: right column) and hormonal (Insulin: middle column) response to a 100g oral glucose tolerance test in young adults (Top; All Subjects, Middle; Males, Bottom; Females) with no overt metabolic pathology. Black lines are mean (SD) while lighter grey lines are individual responses. Data for the means and SDs of each postprandial measure is supplied in Supplementary Table 1.

Figure 2. Metabolic response (Respiratory Exchange Ratio: left column; Carbohydrate Oxidation: Right column) measured by Indirect calorimetry, to a 100g oral glucose tolerance test in young adults (Top; All Subjects, Middle; Males, Bottom; Females) with no overt metabolic pathology. Black lines are mean (SD) while lighter grey lines are individual responses. Data for the means and SDs of each measure is supplied in Supplementary Table 1.

Effect of body composition on substrate response

Using mixed-effects models, we tested the effect of BF% on glucose, insulin, and NEFAs

over time and time^2 to account for a potential nonlinear response. To interpret our results,

obtained beta values were multiplied by 8.7%, the standard deviation for body fat percentage in

our sample, to demonstrate the expected changes in blood glucose, serum insulin, and serum

NEFAs based on the standard deviations for those measures (**Table 3**). The addition of a time^2

main effect and a time^2 x BF% interaction did not improve the model fit for the effect of BF% on

postprandial blood glucose (p = 0.237) or insulin (p = 0.545), suggesting that a linear model has

adequate fit relative to a quadratic model. In the linear models without time^2, BF% did not modify

the relationship between time and glucose (p = 0.294) or insulin (p = 0.178). In the subsequent

main effects models, a 6.9% increase in body fat percentage would result in a significant (β =

0.802; 95% CI: 0.155, 1.45; p = 0.055) 19.1 mg/dl increase in postprandial blood glucose at

each time point. A 2.0% change in body fat is associated with a non-significant increase in

serum insulin (β = 0.232; 95% CI: -0.175, 0.638; p = 0.355). For NEFAs, model fit was improved

by the addition of a time^2 main effect and a time^2 x BF% interaction (p < 0.001) compared to the

model with only linear time terms. In this quadratic model, BF% did not modify the relationship

between NEFAs and time (p = 0.082) or time^2 (p = 0.313). In the main effects model without the

interaction term, a 3.4% body fat percent increase resulted in a non-significant (β = 0.39; 95%

CI: -0.115, 0.894; p = 0.216) increase postprandial plasma NEFA 1.19 (mEq/L). Thus, body

composition affects postprandial blood glucose but not serum insulin or NEFAs throughout the

OGTT (**Table 3**).

Table 3. Effect of body composition on glucose, insulin, and NEFAs during an OGTT

	Glucose			Insulin			NEFA		
Characteristic[1]	Beta[2]	95% CI	p-value	Beta	95% CI	p-value	Beta	95% CI	p-value
Time	-0.55	-0.723, -0.382	**<0.001**	-0.17	-0.286, -0.046	**0.008**	-2.13	-2.83, -1.44	**<0.001**
Time[2]	---	---	---	---	---	---	0.26	0.123, 0.397	**<0.001**
BF%	0.802	0.155, 1.45	0.055	0.232	-0.175, 0.638	0.355	0.39	-0.115, 0.894	0.216
Fasting Measure	0.231	-0.061, 0.523	0.206	-0.24	-0.728, 0.239	0.412	0.211	-0.067, 0.488	0.224
Sex (Male)	0.585	-0.676, 1.84	0.451	0.058	-0.681, 0.794	0.898	1.33	0.360, 2.31	**0.036**
Age	-0.1	-0.464, 0.267	0.66	0.016	-0.190, 0.224	0.897	0.198	-0.075, 0.472	0.244
Matsuda Index	0.188	-0.247, 0.623	0.482	-0.6	-1.17, -0.018	0.105	0.08	-0.242, 0.403	0.683
AdipIR (OGTT)	0.137	-0.280, 0.554	0.591	0.438	0.205, 0.671	**0.006**	0.797	0.452, 1.14	**0.001**
VAT volume[1/3]	-0.01	-0.624, 0.595	0.969	0.056	-0.285, 0.396	0.789	-0.47	-0.934, 0.001	0.116

[1]Fasting Measure denotes the fasting concentration of the outcome variable.
[2]A one-unit increase in time represents 30 minutes. A 1-SD increase in each variable corresponds to a 1-SD increase in glucose (19.1), insulin (0.59), or NEFA (0.45). Insulin and NEFA were transformed via natural log for normality. Variable SDs: BF%: 8.7; Fasting glucose: 8.5; Fasting insulin: 0.7; Fasting NEFA: 0.5; Age: 4.97; Matsuda: 2.9; AdipIR (OGTT): 13.3; VAT: 1.29.

Effect of body composition on postprandial metabolic flexibility (RER, CHO, and FOX)

We used mixed effects models to test for an effect of BF% on postprandial metabolic flexibility (**Table 4**). In these models, a quadratic model with a main effect for time^2 and a time^2 x BF% interaction term was a better fit to the postprandial data for RER (p = 0.002) and CHO (p = 0.002) than the linear model. In the quadratic models, BF% modified the relationship between postprandial RER and time (p = 0.011) and time^2 (p = 0.019), as well as postprandial CHO and time (p = 0.014) and time^2 (p = 0.023), indicating a nonlinear change in RER and CHO throughout the OGTT that depends on body composition. There was no evidence that BF% modifies the relationship between FOX and time (**Supplementary Table 2 and Supplementary Figure 1**). We also considered the effect of fat mass on postprandial RER and CHO. However, there was no evidence that fat mass modifies the relationship between postprandial RER and time (p = 0.364) or time^2 (p = 0.409), as well as postprandial CHO and time (p = 0.364) or time^2 (p = 0.409; see **Supplementary Table 3)**.

To assess this interaction of time and BF% as a measure of postprandial metabolic flexibility, we visualized the effect of time and BF% as continuous variables. We display (**Figure 3**) the modeled RER and CHO for three representative values of BF%: the mean BF%, the mean BF% + 1 SD, and the mean BF% - 1 SD. The interaction of time and BF% displayed in these plots indicates that higher BF% is associated with a greater rate of increase in

postprandial RER (**Figure 3a**) and CHO (**Figure 3b**). This demonstrates that higher BF% is

associated with greater postprandial metabolic flexibility in our sample of young adults. The

resulting visualization is adjusted for all covariates in **Table 4**.

Further, greater insulin sensitivity (Matsuda Index) was associated with a higher

postprandial RER or CHO (**Table 4**). Because skeletal muscle is a primary site of postprandial

glucose disposal[17] and people with obesity often have higher fat-free mass,[18] we tested the

effect of adding fat-free mass as an additional covariate. This did not improve the model fit (p =

0.204) over the original CHO model. Like the original CHO model, the addition of a $time^2$ main

effect and a $time^2$ x BF% interaction term improved the model fit (p = 0.002), and percent body

fat modified the relationship between CHO and time (p = 0.014) and $time^2$ (p = 0.023) (**Table 4**).

Like RER and CHO, higher BF% was associated with greater CHO adjusted for FFM (**Figure**

3c), indicating FFM does not affect the interaction between time and BF%.

Table 4. Effect of body composition (body fat percentage) on postprandial metabolic flexibility

Characteristic[1]	RER			CHO (g/min)			CHO (g/min/FFM)		
	Beta[2]	95% CI	p-value	Beta	95% CI	p-value	Beta	95% CI	p-value
BF% x $Time^2$	-0.133	-0.240, -0.026	**0.019**	-0.134	-0.245, -0.023	**0.023**	-0.134	-0.245, -0.023	**0.023**
BF% x Time	0.732	0.189, 1.28	**0.011**	0.74	0.175, 1.31	**0.014**	0.74	0.175, 1.31	**0.014**
BF%	-0.346	-1.07, 0.374	0.39	-0.51	-1.26, 0.239	0.225	-0.311	-1.11, 0.492	0.502
$Time^2$	-0.156	-0.263, -0.050	**0.006**	-0.166	-0.277, -0.056	**0.005**	-0.166	-0.277, -0.056	**0.005**
Time	1.08	0.537, 1.61	**<0.001**	1.06	0.498, 1.62	**<0.001**	1.06	0.498, 1.62	**<0.001**
Fasting Measure	0.803	0.549, 1.06	**<0.001**	0.793	0.474, 1.11	**<0.001**	0.7	0.358, 1.04	**0.005**
Sex (Male)	0.556	-0.310, 1.42	0.3	0.421	-0.487, 1.33	0.451	0.458	-0.422, 1.34	0.414
Age	-0.095	-0.345, 0.155	0.536	-0.07	-0.332, 0.192	0.661	-0.139	-0.414, 0.137	0.429
Matsuda Index	0.505	0.207, 0.802	**0.012**	0.558	0.246, 0.869	**0.009**	0.59	0.285, 0.895	**0.007**
AdipIR (OGTT)	0.201	-0.083, 0.486	0.255	0.123	-0.176, 0.421	0.504	0.069	-0.232, 0.370	0.715
VAT volume$^{1/3}$	-0.504	-0.909, -0.098	0.055	-0.224	-0.648, 0.199	0.39	-0.368	-0.837, 0.100	0.224
FFM	---	---	---	---	---	---	0.305	-0.177, 0.786	0.323

[1]Fasting Measure denotes the fasting measurement of the outcome variable. See main text for other abbreviations.

[2]A one-unit increase in time represents 30 minutes. A 1-SD increase in each covariate corresponds to a 1-SD increase in postprandial RER (0.044) or CHO (0.055 g/min). Covariate SDs: Fasting RER: 0.055; Fasting CHO: 0.071; FFM: 10.9; See Table 3 for other relevant SDs.

Figure 3. Visualization of the interaction between body fat percentage and postprandial metabolic metabolism as a measure of metabolic flexibility. Respiratory Exchange Ratio (RER) and carbohydrate oxidation (CHO) were plotted as continuous variables across time for representative body fat percentages (BF%) from our sample. The mean BF% of our sample was 30.4 (circles) while 21.7 (squares) and 39.0 (triangles) represent the mean ± 1 SD of BF% in our sample. Error bars represent the 95% confidence interval for each timepoint estimate for a given BF%. The data presented represents modeled data from the interaction of BF% x Time and BF% xTime^2 adjusted for covariates.

DISCUSSION

Metabolic flexibility is the ability to adjust substrate oxidation based on substrate

availability or demand.[1] The relationship between adiposity and measures of metabolic

flexibility has been poorly analyzed and lacks clarity.[3, 19-24] The majority of studies to

date have categorized individuals based on BMI (an indicator of adiposity) and made

weak, indirect associations between adiposity and measures of metabolic flexibility.[3] In

this analysis, we analyzed the interaction between body fat percentage and postprandial

metabolism (RER, CHO, and FOX) and demonstrate that higher body fat percentage is

associated with elevated postprandial metabolic flexibility (RER and CHO) during an

OGTT in young adults without overt metabolic pathology. These findings may seem

paradoxical since individuals with higher BMI (e.g., obesity) are most often reported to

have lower, or blunted, metabolic flexibility. However, this finding is not inconsistent with

studies that show females have greater metabolic flexibility than males despite higher

body fat[5] and individuals with high metabolic flexibility had greater subcutaneous

adipose tissue volumes.[7] Furthermore, loss of subcutaneous adipose tissue during

prolonged bed rest is accompanied by decreased metabolic flexibility.[20] Thus, our work

expands on previously published reports by directly analyzes the relationship between

body fat and postprandial metabolic flexibility while adjusting for known covariates that

significantly affect substrate oxidation.

Adipose tissue function is proposed to be central to the regulation of metabolic

flexibility[2] by buffering fatty acid flux during the postprandial period.[4] The obese state is

associated with increased adipose tissue volume,[25] elevated adipose tissue insulin

resistance, adipose tissue inflammation, and altered adipokine release;[26] all of which

are associated with altered postprandial metabolism.[27] In our healthy young adults,

higher body fat percentages did not alter suppression of lipolysis, as measured by

plasma NEFAs, following ingestion of the glucose beverage. Furthermore, measures of

adipose tissue insulin resistance (ADIP-IR) did not have a significant effect on

postprandial RER or CHO. However, we cannot fully rule out reduced adipose tissue

function in our findings. A critical role of adipose tissue in the postprandial state is

glucose uptake and lipogenesis which cannot be accounted for here.[28-30] Importantly,

our approach demonstrates that an increase in body fat percentage of ~6.9% (i.e., 20%

to 26.9%) would result in an increase in approximately 19.1 mg/dl of blood glucose at

each time point during the OGTT (**Table 3**). This suggests the capacity for adipose

tissue to buffer the blood glucose flux over time may be reduced in people with a higher

body fat percentage, which would be consistent with observational studies in individuals

with obesity, diabetes, and animal models.[31-34] This function is an important part of

adipose tissue's role in reducing hyperglycemia in the postprandial period while

subsequently storing nutrients for later oxidation.

In the context of greater hyperglycemia following the OGTT, the elevated

postprandial metabolic flexibility (postprandial RER and CHO) associated with higher

body fat percentage in our sample would not be paradoxical. The postprandial period

should induce an anabolic state (e.g. storage of nutrients) as compared to the catabolic

state of fasting (e.g. breakdown of substrate).[4] Metabolic flexibility is an critical

metabolic adaptation to handle the complex integration of metabolism during caloric

excess and restriction.[35] Furthermore, it is essential for regular maintenance of blood

glucose homeostasis following a feeding. Our observation of elevated CHO and RER in

the setting of greater hyperglycemia, suggesting greater metabolic flexibility in people

with higher body fat percentages, would fit the concept of metabolic flexibility precisely.

That is, they are adjusting substrate oxidation based on availability of glucose in the

blood for longer periods of time. Individuals with lower body fat percentages not

elevating CHO and RER during this period are likely experiencing the appropriate

catabolic (storage) response to feeding by taking up and storing glucose as glycogen

(liver and skeletal muscle) or fat (adipose tissue). As a result, they have no further need

to increase CHO to maintain blood glucose homeostasis.

This raises the question if our sample of young adults across the BMI spectrum

but with no overt metabolic pathology are demonstrating a resilient metabolically flexible

phenotype.[36] Metabolic flexibility is diminished following hypercaloric, high-fat feeding

interventions and improved following significant weight loss, indicating it is a plastic

phenotype.[37, 38] No study has successfully tracked the disappearance of metabolic

flexibility, but we suspect that long-term elevated levels of adiposity, such as that seen

in chronic obesity, eventually lead to the loss of postprandial metabolic flexibility. In our

sample, higher body fat percentage is inversely associated with insulin sensitivity (r =-

0.616; p=0.001) while being positively associated with fasting values for serum insulin

(r=0.427; p=0.037), plasma leptin (r=0.817; p<0.001), and adipose tissue insulin

resistance (r=0.443; p=0.030). These relationships may be indicative or worsening

metabolic health with higher body fat percentage, but the ability to elevate CHO

oxidation during the OGTT (e.g. metabolic flexibility) demonstrates a resilient,

metabolically flexibility phenotype in these individuals. The eventual loss of metabolic

flexibility would therefore play a significant role in the progression of glucose dysfunction

towards overt metabolic disease. That process and the mechanisms involved are still

poorly understood and require both experimental and observational longitudinal studies.

Our approach to analyzing metabolic flexibility during an OGTT as a continuous

variable across time is a novel aspect of our approach. Most studies assessing

adiposity and metabolic flexibility categorize subjects based on BMI, a surrogate for

adiposity, and define metabolic flexibility as the change in RER or CHO from fasting to a

defined timepoint (e.g. 60 min in an OGTT) or the peak change (max – min) in the

measure.[3] That approach largely demonstrates that individuals with overweight and

obesity have reduced (blunted) metabolic flexibility. Our data, when analyzed in that

way, produces similar results (**Supplementary Figure 2**). However, these indirect

associations do not clarify the relationship between adiposity and metabolic flexibility

and oversimplify the temporal postprandial metabolic response into a single timepoint,

leading to inconclusive results. Defining postprandial metabolic flexibility based on a

single timepoint during a dynamic and prolonged response, such as feeding, is likely a

carryover from the euglycemic clamp method, which applies indirect calorimetry to the

steady-state phase of insulin infusion. Because postprandial metabolism is dynamic, a

feeding challenge such as an OGTT has no equivalent to the steady-state period.[39, 40]

Using a continuous approach has allowed us to demonstrate an unexpected, but

explainable, relationship in which higher body fat percentage is associated with

postprandial metabolic flexibility in young adults without overt metabolic pathology.

In conclusion, we demonstrate that higher body fat percentage is associated

with greater postprandial metabolic flexibility (RER and CHO) during an oral glucose

tolerance test in young adults without overt metabolic pathology. We believe this finding

is indicative of a preserved metabolically flexible phenotype where elevated

postprandial blood glucose is resulting in elevated CHO with higher body fat

percentage. Our approach of analyzing data in a continuous manner using linear mixed

models, accounting for covariates known to modify the postprandial substrate oxidation

response, adds clarity to the previously poorly defined relationship between adiposity

and postprandial metabolic flexibility. While we cannot account for some functional (e.g.

adipocyte size, lipogenesis, and molecular) or anatomical (e.g. distribution) aspects of

adiposity, we directly demonstrate what has long been ambiguous in published reports.

These findings raise several new questions about adiposity and the continuum of

metabolic health.

Conflicts of Interest: The authors declare no conflicts of interest.

Acknowledgements: The authors would like to acknowledge the Milken Institute

School of Public Health Office of Research Excellence which provided funding for this

project. The authors would also like to acknowledge the participants for their

volunteered time and effort.

Funding: This study was funded by a Research Innovation Award (PI Barberio) from

the Office of Research Excellence in the Milken Institute School of Public Health at The

George Washington University.

Contributions: **NF:** Conceptualization, methodology, formal analysis, investigation,
data curation, writing – original draft, visualization. **SR:** Investigation. JB: Investigation.
LS: Investigation. **AL:** Investigation. **LD:** Conceptualization, writing – review & editing.
AC: Formal analysis, writing – review & editing. **MB:** Conceptualization, methodology,
investigation, writing – original draft, project administration, funding acquisition.

All authors have reviewed this manuscript and agree to its publication as written.

References

1.   Goodpaster BH, Sparks LM. Metabolic Flexibility in Health and Disease. Cell Metab.

2017;25(5):1027-36. doi: 10.1016/j.cmet.2017.04.015. PubMed PMID: 28467922; PMCID:

PMC5513193.

2.   Storlien L, Oakes ND, Kelley DE. Metabolic flexibility. Proc Nutr Soc. 2004;63(2):363-8.

doi: 10.1079/PNS2004349. PubMed PMID: 15294056.

3.   Glaves A, Diaz-Castro F, Farias J, Ramirez-Romero R, Galgani JE, Fernandez-Verdejo

R. Association Between Adipose Tissue Characteristics and Metabolic Flexibility in Humans: A

Systematic Review. Front Nutr. 2021;8:744187. Epub 20211203. doi:

10.3389/fnut.2021.744187. PubMed PMID: 34926544; PMCID: PMC8678067.

4.   Frayn KN. Adipose tissue as a buffer for daily lipid flux. Diabetologia. 2002;45(9):1201-

10. Epub 20020724. doi: 10.1007/s00125-002-0873-y. PubMed PMID: 12242452.

5.   Sparks LM, Pasarica M, Sereda O, deJonge L, Thomas S, Loggins H, Xie H, Miles JM,

Smith SR. Effect of adipose tissue on the sexual dimorphism in metabolic flexibility. Metabolism.

2009;58(11):1564-71. doi: 10.1016/j.metabol.2009.05.008. PubMed PMID: 19595383.

6.   Sparks LM, Ukropcova B, Smith J, Pasarica M, Hymel D, Xie H, Bray GA, Miles JM,

Smith SR. Relation of adipose tissue to metabolic flexibility. Diabetes Res Clin Pract.

2009;83(1):32-43. Epub 20081126. doi: 10.1016/j.diabres.2008.09.052. PubMed PMID:

19038471; PMCID: PMC2749984.

7.   Fernandez-Verdejo R, Malo-Vintimilla L, Gutierrez-Pino J, Lopez-Fuenzalida A, Olmos

P, Irarrazaval P, Galgani JE. Similar Metabolic Health in Overweight/Obese Individuals With

Contrasting Metabolic Flexibility to an Oral Glucose Tolerance Test. Front Nutr. 2021;8:745907.

Epub 20211116. doi: 10.3389/fnut.2021.745907. PubMed PMID: 34869522; PMCID:

PMC8637191.

8.   Taylor HL, Buskirk E, Henschel A. Maximal oxygen intake as an objective measure of

cardio-respiratory performance. J Appl Physiol. 1955;8(1):73-80. doi: 10.1152/jappl.1955.8.1.73.

PubMed PMID: 13242493.

9.   Frayn KN. Calculation of substrate oxidation rates in vivo from gaseous exchange. J

Appl Physiol Respir Environ Exerc Physiol. 1983;55(2):628-34. doi:

10.1152/jappl.1983.55.2.628. PubMed PMID: 6618956.

10.   Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose

tolerance testing: comparison with the euglycemic insulin clamp. Diabetes Care.

1999;22(9):1462-70. doi: 10.2337/diacare.22.9.1462. PubMed PMID: 10480510.

11.   Gastaldelli A, Gaggini M, DeFronzo RA. Role of Adipose Tissue Insulin Resistance in

the Natural History of Type 2 Diabetes: Results From the San Antonio Metabolism Study.

Diabetes. 2017;66(4):815-22. Epub 20170104. doi: 10.2337/db16-1167. PubMed PMID:

28052966.

12.   Shashaj B, Luciano R, Contoli B, Morino GS, Spreghini MR, Rustico C, Sforza RW,

Dallapiccola B, Manco M. Reference ranges of HOMA-IR in normal-weight and obese young

Caucasians. Acta Diabetol. 2016;53(2):251-60. Epub 20150613. doi: 10.1007/s00592-015-

0782-4. PubMed PMID: 26070771.

13.   Kuznetsova A, Brockhoff, P.B., & Christensen R.H.B. lmerTest Package: Tests in Linear

Mixed Effects Models. Journal of Statistical Software. 2017;82(13):1-26. doi:

10.18637/jss.v082.i13.

14.   Lüdecke D, Ben-Shachar, M.S., Patil, I., Waggoner, P., Makowski, D. performance: An R

Package for Assessment, Comparison

and Testing of Statistical Models. The Journal of Open Source Software. 2021;6(60). doi:

10.21105/joss.03139.

15.   Lenth R. emmeans: Estimated Margin2l Means, aka Least-Squares Means. R package

version 1.10.3-090006. 2024.

16.   Baayen RH, Davidson DJ, Bates DM. Mixed-effects modeling with crossed random

effects for subjects and items. Journal of Memory and Language. 2008;59(4):390-412. doi:

10.1016/j.jml.2007.12.005. PubMed PMID: WOS:000261651700003.

17.   Kelley D, Mitrakou A, Marsh H, Schwenk F, Benn J, Sonnenberg G, Arcangeli M, Aoki T,

Sorensen J, Berger M, et al. Skeletal muscle glycolysis, oxidation, and storage of an oral

glucose load. J Clin Invest. 1988;81(5):1563-71. doi: 10.1172/JCI113489. PubMed PMID:

3130396; PMCID: PMC442590.

18.   Dulloo AG, Jacquet J, Miles-Chan JL, Schutz Y. Passive and active roles of fat-free

mass in the control of energy intake and body composition regulation. Eur J Clin Nutr.

2017;71(3):353-7. Epub 20161214. doi: 10.1038/ejcn.2016.256. PubMed PMID: 27966570.

19.   Lightowler H, Schweitzer L, Theis S, Henry CJ. Changes in Weight and Substrate

Oxidation in Overweight Adults Following Isomaltulose Intake During a 12-Week Weight Loss

Intervention: A Randomized, Double-Blind, Controlled Trial. Nutrients. 2019;11(10). Epub

20191004. doi: 10.3390/nu11102367. PubMed PMID: 31590285; PMCID: PMC6836138.

20.   Rudwill F, O'Gorman D, Lefai E, Chery I, Zahariev A, Normand S, Pagano AF, Chopard

A, Damiot A, Laurens C, Hodson L, Canet-Soulas E, Heer M, Meuthen PF, Buehlmeier J,

Baecker N, Meiller L, Gauquelin-Koch G, Blanc S, Simon C, Bergouignan A. Metabolic

Inflexibility Is an Early Marker of Bed-Rest-Induced Glucose Intolerance Even When Fat Mass Is

Stable. J Clin Endocrinol Metab. 2018;103(5):1910-20. doi: 10.1210/jc.2017-02267. PubMed

PMID: 29546280; PMCID: PMC7263792.

21.   Kahlhofer J, Lagerpusch M, Enderle J, Eggeling B, Braun W, Pape D, Muller MJ, Bosy-

Westphal A. Carbohydrate intake and glycemic index affect substrate oxidation during a

controlled weight cycle in healthy men. Eur J Clin Nutr. 2014;68(9):1060-6. Epub 20140709. doi:

10.1038/ejcn.2014.132. PubMed PMID: 25005676.

22.   Bergouignan A, Antoun E, Momken I, Schoeller DA, Gauquelin-Koch G, Simon C, Blanc

S. Effect of contrasted levels of habitual physical activity on metabolic flexibility. J Appl Physiol

(1985). 2013;114(3):371-9. Epub 20121213. doi: 10.1152/japplphysiol.00458.2012. PubMed

PMID: 23239872.

23.   Assaad M, El Mallah C, Obeid O. Phosphorus ingestion with a high-carbohydrate meal

increased the postprandial energy expenditure of obese and lean individuals. Nutrition.

2019;57:59-62. Epub 20180619. doi: 10.1016/j.nut.2018.05.019. PubMed PMID: 30153580.

24.   Huda MS, Dovey T, Wong SP, English PJ, Halford J, McCulloch P, Cleator J, Martin B,

Cashen J, Hayden K, Wilding JP, Pinkney J. Ghrelin restores 'lean-type' hunger and energy

expenditure profiles in morbidly obese subjects but has no effect on postgastrectomy subjects.

Int J Obes (Lond). 2009;33(3):317-25. Epub 20090203. doi: 10.1038/ijo.2008.270. PubMed

PMID: 19188925.

25.   Turer AT, Khera A, Ayers CR, Turer CB, Grundy SM, Vega GL, Scherer PE. Adipose

tissue mass and location affect circulating adiponectin levels. Diabetologia. 2011;54(10):2515-

24. Epub 20110722. doi: 10.1007/s00125-011-2252-z. PubMed PMID: 21779869; PMCID:

PMC4090928.

26.   Shafiei MS, Shetty S, Scherer PE, Rockey DC. Adiponectin regulation of stellate cell

activation via PPARgamma-dependent and -independent mechanisms. Am J Pathol.

2011;178(6):2690-9. doi: 10.1016/j.ajpath.2011.02.035. PubMed PMID: 21641391; PMCID:

PMC3124230.

27.   Sun K, Kusminski CM, Scherer PE. Adipose tissue remodeling and obesity. J Clin

Invest. 2011;121(6):2094-101. Epub 20110601. doi: 10.1172/JCI45887. PubMed PMID:

21633177; PMCID: PMC3104761.

28.   Chadt A, Al-Hasani H. Glucose transporters in adipose tissue, liver, and skeletal muscle

in metabolic health and disease. Pflugers Arch. 2020;472(9):1273-98. Epub 20200626. doi:

10.1007/s00424-020-02417-x. PubMed PMID: 32591906; PMCID: PMC7462924.

29.   Luo L, Liu M. Adipose tissue in control of metabolism. J Endocrinol. 2016;231(3):R77-

R99. doi: 10.1530/JOE-16-0211. PubMed PMID: 27935822; PMCID: PMC7928204.

30.   Roberts R, Hodson L, Dennis AL, Neville MJ, Humphreys SM, Harnden KE, Micklem KJ,

Frayn KN. Markers of de novo lipogenesis in adipose tissue: associations with small adipocytes

and insulin sensitivity in humans. Diabetologia. 2009;52(5):882-90. Epub 20090228. doi:

10.1007/s00125-009-1300-4. PubMed PMID: 19252892.

31.   Fabbrini E, Yoshino J, Yoshino M, Magkos F, Tiemann Luecking C, Samovski D,

Fraterrigo G, Okunade AL, Patterson BW, Klein S. Metabolically normal obese people are

protected from adverse effects following weight gain. J Clin Invest. 2015;125(2):787-95. Epub

20150102. doi: 10.1172/JCI78425. PubMed PMID: 25555214; PMCID: PMC4319438.

32.   Graham TE, Kahn BB. Tissue-specific alterations of glucose transport and molecular

mechanisms of intertissue communication in obesity and type 2 diabetes. Horm Metab Res.

2007;39(10):717-21. doi: 10.1055/s-2007-985879. PubMed PMID: 17952832.

33.   Herman MA, Peroni OD, Villoria J, Schon MR, Abumrad NA, Bluher M, Klein S, Kahn

BB. A novel ChREBP isoform in adipose tissue regulates systemic glucose metabolism. Nature.

2012;484(7394):333-8. doi: 10.1038/nature10986. PubMed PMID: 22466288; PMCID:

PMC3341994.

34.   Hoffstedt J, Forster D, Lofgren P. Impaired subcutaneous adipocyte lipogenesis is

associated with systemic insulin resistance and increased apolipoprotein B/AI ratio in men and

women. J Intern Med. 2007;262(1):131-9. doi: 10.1111/j.1365-2796.2007.01811.x. PubMed

PMID: 17598821.

35.   Smith RL, Soeters MR, Wust RCI, Houtkooper RH. Metabolic Flexibility as an Adaptation

to Energy Resources and Requirements in Health and Disease. Endocr Rev. 2018;39(4):489-

517. doi: 10.1210/er.2017-00211. PubMed PMID: 29697773; PMCID: PMC6093334.

36.   Mietus-Snyder M, Perak AM, Cheng S, Hayman LL, Haynes N, Meikle PJ, Shah SH,

Suglia SF, American Heart Association Atherosclerosis H, Obesity in the Young Committee of

the Council on Lifelong Congenital Heart D, Heart Health in the Y, Council on L,

Cardiometabolic H, Council on Arteriosclerosis T, Vascular B, Council on C, Stroke N. Next

Generation, Modifiable Cardiometabolic Biomarkers: Mitochondrial Adaptation and Metabolic

Resilience: A Scientific Statement From the American Heart Association. Circulation.

2023;148(22):1827-45. Epub 20231030. doi: 10.1161/CIR.0000000000001185. PubMed PMID:

37902008.

37.   Berk ES, Kovera AJ, Boozer CN, Pi-Sunyer FX, Albu JB. Metabolic inflexibility in

substrate use is present in African-American but not Caucasian healthy, premenopausal,

nondiabetic women. J Clin Endocrinol Metab. 2006;91(10):4099-106. Epub 20060725. doi:

10.1210/jc.2005-2411. PubMed PMID: 16868062; PMCID: PMC2670464.

38.   Kelley DE, Goodpaster B, Wing RR, Simoneau JA. Skeletal muscle fatty acid

metabolism in association with insulin resistance, obesity, and weight loss. Am J Physiol.

1999;277(6):E1130-41. doi: 10.1152/ajpendo.1999.277.6.E1130. PubMed PMID: 10600804.

39.   Alcantara JMA, Galgani JE. Association of metabolic flexibility indexes after an oral

glucose tolerance test with cardiometabolic risk factors. Eur J Clin Nutr. 2024;78(3):180-6. Epub

20231218. doi: 10.1038/s41430-023-01373-w. PubMed PMID: 38110728.

40.   Alcantara JMA, Sanchez-Delgado G, Jurado-Fasoli L, Galgani JE, Labayen I, Ruiz JR.

Reproducibility of the energy metabolism response to an oral glucose tolerance test: influence

of a postcalorimetric correction procedure. Eur J Nutr. 2023;62(1):351-61. Epub 20220825. doi:

10.1007/s00394-022-02986-w. PubMed PMID: 36006468; PMCID: PMC9899729.

564