Increased Postprandial Metabolic Flexibility is Associated with Higher Body Fat Percentages in Healthy Young Adults

Nicholas A. Foreman[1], Sahil Rajwade[1], Jaiden Bluth[1], Lauren C. Skoglund[1], Audrey M. Letts[1], Loretta DiPietro[1], Adam Ciarleglio[2], Matthew D. Barberio[1].

[1]Department of Exercise and Nutrition Sciences, The Milken Institute School of Public Health, George Washington University, Washington D.C.

[2]Department of Biostatistics, The Milken Institute School of Public Health, George Washington University, Washington D.C.

Corresponding Author

Matthew D. Barberio
Assistant Professor
Department of Exercise and Nutrition Sciences
The Milken Institute School of Public Health
George Washington University
Washington, D.C.
Email: mbarberio@gwu.edu
Phone: 202-994-1408
ORCID: 0000-0003-0468-7842

Keywords: Substrate oxidation, Postprandial, Metabolic Health, Glucose Tolerance

Abstract

PURPOSE: Because higher adiposity is associated with cardiometabolic disease, we assessed the relationship between body composition (body fat percentage; BF%) and postprandial metabolic flexibility. **METHODS:** Young adults (n = 27, n = 15 females, BMI = 27.1 ± 4.5; BF% = 30.4 ± 8.7) without overt pathology completed a 100g oral glucose tolerance test (OGTT). Indirect calorimetry before (fasting) and following (30, 60, 90, 120 min) consumption was used to calculate respiratory exchange ratio (RER) and oxidation of carbohydrates (CHO) and fats (FOX). Serum and plasma were collected at corresponding time points and analyzed for blood glucose, insulin, and NEFAs. Data from individuals with normal weight were compared to those with overweight/obesity by two-way repeated measures ANVOA. The effect of BF% on postprandial metabolism was tested via linear mixed models while adjusting for potential confounders. **RESULTS:** During the OGTT, blood glucose, serum insulin, plasma lactate, RER, and CHO all significantly increased while plasma NEFAs and whole-body FOX decreased (all $p<0.05$). BF% modified the relationship between postprandial RER and time ($p = 0.019$) as well as postprandial CHO and time ($p = 0.023$) without an effect of BF% on FOX; individuals with higher BF% increase their RER and CHO faster and to a greater extent than those with lower BF%. **CONCLUSION:** Body fat percentage is associated with greater postprandial metabolic flexibility during an OGTT in young adults. Despite increased adiposity, metabolic flexibility may be preserved, representing a compensatory adaptation to decreased glucose storage in the postprandial period.

1 **Introduction**
2
Metabolic flexibility is described as the appropriate shift in substrate oxidation based

upon substrate availability (e.g. fasting to fed) or metabolic demand (e.g. exercise). It was

originally described as the change in the respiratory quotient (RQ) across the leg limb from rest

to insulin stimulation during the euglycemic-hyper-insulinemic clamp (EHC).[1, 2] In the two

decades since the concept of metabolic flexibility was described in humans, several

methodological approaches to assess metabolic flexibility have been utilized. These include

indirect calorimetry during the EHC, during oral glucose tolerance tests (OGTT), and during

mixed-meal feeding challenges (MMC).[1, 3] Individuals with overt metabolic pathologies, such as

type 2 diabetes and insulin resistance, as well as individuals with obesity based on BMI, have

been described as metabolically inflexible.[1] That is, these individuals showed blunted (reduced)

changes in the respiratory exchange ratio (RER) during metabolic challenges such as EHC and

feeding.

In the absence of overt metabolic pathology or metabolic dysfunction, transitioning from

the fasted to the fed state involves shifting away from aerobic fatty acid oxidation to aerobic

glucose oxidation and, to a lesser degree, glycolytic energy production.[4] This shift results in an

anabolic environment, in which exogenous energy is stored, as compared to the catabolic

environment of fasting. Elevated insulin, which is a primary driver of the metabolic shift following

feeding, suppresses adipose tissue lipolysis and promotes the uptake, storage, and oxidation of

exogenous macronutrients.[4] Adipose tissue serves a critical role in buffering fatty acid flux in the

circulation during this postprandial period; it is also a site of significant glucose uptake for

lipogenesis and has been identified as a central regulator of metabolic flexibility.[2, 4]

The presence of obesity and increased fat mass is believed to negatively impact

metabolic flexibility.[5, 6] Individuals with obesity have higher resting RER indicating a preference

for higher carbohydrate oxidation in the fasting state. In obesity, adipose tissue lipolysis is also

more resistant to insulin-induced suppression in feeding. However, studies have not thoroughly

analyzed the effect of adiposity on postprandial metabolic flexibility. By dividing participants into

high vs. low metabolic flexibility, Sparks et al. (2009) demonstrated that healthy young males

with low metabolic flexibility during insulin stimulation (EHC) had significantly higher body fat

percentage and larger fat cell size.[6] Follow-up analysis in this cohort demonstrated that females

were more metabolically flexible than their male counterparts despite significantly higher body

fat percentages.[5] Conversely, individuals classified as having high metabolic flexibility to an

OGTT had similar body fat percentages but a larger subcutaneous adipose tissue (SAT) as

compared to those described as having low metabolic flexibility.[7] This discrepancy was

highlighted in a recent systematic review by Glaves et al. (2021), who found inconsistent and

unclear relationships between adipose tissue volume, function, and characteristics in studies

using EHC, OGTTs, and MMCs.[3]

The primary objective of this study was to thoroughly assess the relationship between

adiposity and postprandial metabolic flexibility during an OGTT. We recruited 27 healthy male

and female young adults (18 - 40 years old) with body mass index (BMI) between 18.5-39.9.

Using indirect calorimetry, we assessed postprandial metabolic flexibility (RER, Carbohydrate

Oxidation [CHO], and Fat Oxidation [FOX]) during an OGTT. We hypothesized that individuals

with higher body fat percentages would demonstrate significantly reduced postprandial

metabolic flexibility (RER and CHO) throughout the OGTT.

METHODS

Participants

48
We performed a cross-sectional, experimental study from January 2023 to January 2024.

Potential participants were recruited and screened based on sex, body mass index (BMI)

classification, and lack of known clinical pathology from the George Washington University

campus and surrounding metropolitan area. Participants were required to be between 18 and 40

years old, have a BMI between 18.5 and 39.9, be free of clinically diagnosed cardiometabolic

disease, and not be on prescription medications for hypertension, blood lipids, or blood glucose

homeostasis. Participants who were pregnant or taking medications for cardiometabolic disease

were excluded. Recruitment was stratified by BMI classification to yield a representative sample

across the BMI continuum. In response to flyers and emails, fifty-two participants were screened

for participation. Of these, thirty-four were enrolled in this study; seven participants did not

complete the study (3 due to scheduling, 3 due to loss to follow-up, 1 due to equipment

malfunction), giving complete data for 27 participants. Participant characteristics are reported in

Table 1. All study procedures were approved by the George Washington University IRB (IRB

#202737). Prior to enrollment, participants read and signed an IRB-approved informed consent,

and all study procedures were conducted in accordance with the Declaration of Helsinki.

Experimental Design

65
Participants reported to the Metabolic and Exercise Testing Laboratories in the morning

between 0800 and 0900 on two separate days. Each visit was separated by at least 48 hours.

At the first visit, participants completed written questionnaires for medical and physical activity

history, assessment of body composition, and cardiorespiratory fitness. Participants returned to

the lab for assessment of postprandial substrate metabolism following an OGTT.

For each visit, participants were instructed to arrive after 10 hours of fasting and to

abstain from caffeine consumption for 12 hours and from any recreational drug use or structured

exercise for 24 hours. Upon arrival, hydration was measured using a portable refractometer. If

urine specific gravity was above 1.02, participants were given eight ounces of water. Body

weight was measured using a mechanical column scale (Seca 700; Hamburg, Germany).

Table 1: Participant Characteristics

Variable	All (n = 27)	NW (n = 12)	OW/OB (n = 15)	p	Association with BF%
Sex, % Female	12 (56%)	6 (50%)	6 (40%)	0.897	---
NIH Race/Ethnicity, %				0.328	---
Black	7 (25.9)	2 (16.7)	5 (33.3)		---
Hispanic or Latino	1 (3.7)	0 (0.0)	1 (6.7)		---
Unknown	1 (3.7)	0 (0.0)	1 (6.7)		---
white	18 (66.7)	10 (83.3)	8 (53.3)		---
Age, yrs	22 [19, 27]	19 [19, 22]	25 [23, 29.5]	0.002	r=0.081; p=0.688
Height, m	1.71 (0.09)	1.7 (0.08)	1.71 (0.1)	0.641	r=-0.415; p=0.032
Weight, kg	73.0 [66.6, 90.0]	66.6 [61.5, 71.4]	89.9 [75.5, 96.9]	<0.001	r=0.257; p=0.195
BMI, kg/m^2	26.2 [24.1, 31.4]	23.7 [21.8, 24.4]	31.2 [27.2, 32.6]	<0.001	r=0.582; p=0.001
Fat Mass, kg	24.2 (9.6)	17 (4.6)	30 (8.5)	<0.001	r=0.854; p<0.001
Fat Free Mass, kg	51.7 (10.9)	46.5 (8.3)	55.8 (11.2)	0.025	r=-0.351; p=0.072
Body Composition, % Fat	30.4 (8.7)	26 (7.8)	33.8 (8)	0.017	NA
VAT Volume, in^3	2.7 (1.29)	1.9 (1.06)	3.3 (1.12)	0.003	r=0.301; p=0.127
Resting Metabolic Rate, kcal/d	1709.8 (287.9)	1627.3 (260.9)	1775.7 (299.9)	0.188	r=-0.25; p=0.208
Respiratory Exchange Ratio (RER)	0.8 (0.06)	0.78 (0.06)	0.8 (0.05)	0.413	r=-0.129; p=0.520
Carbohydrate Oxidation, g/min	0.1 (0.07)	0.08 (0.06)	0.12 (0.07)	0.198	r=-0.16; p=0.424
Lipid Oxidation, g/min	0.08 (0.02)	0.09 (0.03)	0.08 (0.02)	0.733	r=-0.031; p=0.878
VO2peak, mL/kg/min	34.1 (8.8)	40 (8.3)	29.5 (6.2)	0.001	r=-0.784; p<0.001
VO2peak, L/min	2.6 (0.6)	2.7 (0.68)	2.6 (0.54)	0.736	r=-0.66; p<0.001

Data are mean ± SD or median [IQR], depending on normality.

Body mass and body composition

Height was measured using a wall-mounted stadiometer (Seca 216; Hamburg,

Germany) to the nearest 0.1 centimeters. Weight was measured using the digital scale as part

of the bioelectrical impedance analysis (InBody 770; Seoul, South Korea). Dual X-ray

absorptiometry (GE Lunar iDXA; GE Healthcare, Chicago, IL, USA) with manufacturer software

(enCORE Forma version 15) was used to measure body fat percentage, fat-free mass (FFM),

fat mass, and visceral adipose tissue (VAT) volume. Before the measurement, participants

removed all jewelry and were instructed to wear clothing free of metal zippers or reflective

surfaces.

Exercise Testing

89
Following body composition assessment, participants completed a graded exercise test

on a cycle ergometer (Monark LC6 novo; Varberg, Sweden) to volitional exhaustion. During

cycling, expired gases were measured using a breath-by-breath metabolic cart (Cosmed Quark

CPET; Rome, Italy), and heart rate was monitored using a chest strap (Polar H10; Kempele,

Finland). All participants completed three 180-second stages of 25, 50, and 75W each before

resistance increased by 25W per minute. The test continued until the participant was unable to

continue cycling or the investigator observed a plateau in oxygen consumption (VO_2). Breath-

by-breath data were averaged into five-second bins. Because not every participant satisfied the

criteria for VO2max,[8] the highest 15-second value is reported as VO2peak.

99
100   *Oral Glucose Tolerance Test with Indirect Calorimetry and Blood Sampling*

101
On a separate day, participants arrived following the same pre-testing guidelines

outlined above. Assessment of body weight and hydration were performed, and the participant

rested in a reclined chair in a dimly lit room for at least 10-15 minutes. A venous catheter was

placed in an antecubital vein and fasting whole blood was sampled for plasma (K-EDTA) and

serum (Serum Separator Tube); saline was used to keep the catheter patent between blood

draws. After the resting blood draw, indirect calorimetry was conducted via the canopy method

(Parvo Medics TrueOne 2400; Salt Lake City, Utah) while the participant lay supine on a

padded table for determination of fasting metabolic measures. Following this, subjects

consumed a 100g glucose beverage (GlucoCrush Cardinal Health; Dublin, OH) and were

instructed to finish consumption of the drink within 5 minutes. After finishing the drink, subjects

returned to the supine position for the remainder of the test. Indirect calorimetry was used to

assess postprandial substrate oxidation at the following postprandial intervals (min): 15 – 30, 45

– 60, 75 – 90, and 105 – 120.

Determination of substrate oxidation and metabolic rate

116
For each 15-minute indirect calorimetry measurement, expired gases were collected from a

mixing chamber at 5-second intervals, and the final 8 minutes were used for analysis.

Respiratory exchange ratio (RER) was calculated as the ratio between exhaled carbon dioxide

(VCO_2) and VO_2. Metabolic rate and rates of carbohydrate (CHO) and fat oxidation (FOX) were

calculated using published equations for fat oxidation as [VO_2 (L/min) x 1.67) – (VCO_2 (L/min) x

1.67)] and carbohydrate oxidation as [VCO_2 (L/min) x 4.55) – (VO_2 (L/min) x 3.21)].[9] For

metabolic rate calculations, urinary nitrogen excretion was assumed to be zero.

Laboratory measurements and calculations

Fasting and postprandial blood glucose were determined in fresh whole blood

(OneTouch Ultra, Lifescan, Inc.; Milpitas, CA). Fasting and postprandial serum samples were

used to measure insulin (Insulin Human ELISA; Invitrogen Inc. CA), and non-esterified fatty

acids (NEFA; LabAssay NEFA; FujiFilm; Richmond, VA). Fasting serum samples were used to

determine C-reactive protein (CRP; CRP Human ELISA; Invitrogen Inc., CA), leptin (Leptin

Human ELISA; Invitrogen Inc., CA), and adiponectin (Adiponectin human ELISA; Adipogen Life

Science; San Diego, CA). Manufacturer instructions were followed for conducting and analyzing

all assays. Manufacturer internal controls were assessed along with coefficients of variations for

biological replicates. Samples whose performance was assessed to be outside of manufacturer

recommendations were re-analyzed. Fresh whole blood (fasting) was used to determine

HbA1C% (PTS Diagnostics; Whitestown, IN) and blood lipid profiles (CardioChek; PTS

Diagnostics; Whitestown, IN) via point-of-care analyzers.

Matsuda Index was calculated as follows: 10,000/√ [fasting glucose (mg/dl) × fasting

insulin (μIU/mL)] × [mean glucose (mg/dl) × mean insulin (μIU/mL) during the OGTT before

normalizing to the 100g dose.[10] Adipose tissue insulin resistance (Adip-IR) was calculated using

fasting values as fasting insulin (µIU/mL) * fasting NEFA (mEq/L) and using postprandial OGTT

values (30, 60, 90, 120) as mean plasma NEFA (mEq/L) x mean serum insulin (µIU/mL).[11]

Fasting insulin and glucose measurements were used to calculate the homeostatic model

assessment of insulin resistance (HOMA-IR) as [fasting insulin (µIU/mL) - (fasting glucose

(mg/dl)/18.156)] / 22.5.[12]

Statistical Analyses

146
We report data as mean (SD) or median (interquartile range, IQR) both overall and by

BMI status for participant characteristics (**Table 1**) and clinical measures (**Table 2**). We

compared race categories between the sexes using Fisher's exact test and report them as

frequency (percent). We computed and tested the association between each of the clinical

variables and percent body fat (BF%) using Pearson correlation coefficients. To describe the

change in each postprandial outcome variables during the OGTT (i.e., RER, CHO, FOX,

glucose, insulin, lactate, and NEFA) between individuals with normal weight (NW) and

individuals with overweight and obesity (OW/OB) as defined by the BMI, we used two-way,

repeated measures analysis of variance (ANOVA) with time modeled as a categorical variable.

Contrasts were set *a priori* as the comparison from baseline to each postprandial timepoint (i.e.,

0 vs. 30, 0 vs. 60, etc.). Analysis and figures were conducted in OriginPro 2020.

To determine the effect of body composition on postprandial metabolism, linear mixed-

effects models were used. For each postprandial outcome variable, we included fixed effects for

time as a continuous variable, percent body fat (BF%), time x BF% interaction, sex, age,

Matsuda index, AdipIR, visceral adipose tissue volume, and the fasted outcome measure. A

subject-specific random intercept term was included to account for repeated measures. To

account for the possible nonlinear effect of time on postprandial metabolism (i.e., a nonlinear

change in metabolism), models with an additional quadratic time term and quadratic time x BF%

were considered. Models with and without the quadratic time terms were compared using

likelihood ratio tests to determine if the quadratic terms should be retained. Variables in the

models were tested for normality with the Shapiro-Wilk test and transformed. As a result, insulin

and NEFA concentrations were transformed via natural log, and VAT volume was transformed

via cube root. Each variable included in the models was standardized to a mean of 0 and

standard deviation of 1, and time was divided by 30 to correct scaling. Fasting and postprandial

variables were standardized separately due to the inclusion of the fasting state as a covariate in

the models. All data transformation and subsequent mixed-effects models were conducted in R

(version 4.2.2) using the lmerTest[13], performance[14], and emmeans[15] packages. For the

interpretation of linear mixed-effects models, we assess significance using both the 95%

confidence interval (e.g., does it cross 0) and the p-value ($p < 0.05$). Due to ambiguity in the

determination of p-values for these models[16], we deferred to the confidence interval when there

was disagreement between the confidence intervals and p-values.

RESULTS

There were 27 young adults (n=15 (56%) female) with no previously diagnosed

metabolic pathology (**Table 1)**. As expected, individuals with OW/OB had significantly ($p <$

0.001) higher body mass, volume of body fat, visceral fat area, and fat free mass as compared

to individuals with NW. There were no significant differences in resting metabolic rate, RER, or

substrate oxidation between individuals with OW/OB and NW (**Table 1**). Comparisons of

participant characteristics based on biological sex are presented in **Supplementary Table 1**.

Table 2: Clinical Data

Variable	All (n = 27)	NW (n = 12)	OWOB (n = 15)	p-value	Association with BF%
Blood Glucose (mg/dL)	100 (8)	98 (12)	101 (5)	0.315	r=0.036; p=0.858
Serum Insulin (microIU/mL)	14.2 (8.1)	11.7 (5.8)	16.3 (9.4)	0.17	r=0.404; p=0.050
NEFA (mEq/L)	0.42 [0.28, 0.49]	0.34 [0.25, 0.47]	0.43 [0.32, 0.49]	0.344	r=0.339; p=0.098
Serum Lactate (mM)	1.58 (0.42)	1.43 (0.33)	1.72 (0.45)	0.09	r=0.237; p=0.266
HbA1c (%)	5.1 [4.8, 5.25]	5.05 [4.8, 5.12]	5.2 [4.85, 5.4]	0.154	r=-0.221; p=0.269
Total Cholesterol (mg/dL)	160.4 (33.9)	139 (22.8)	175.6 (32.7)	0.006	r=0.198; p=0.353
HDL (mg/dL)	51.0 [47.8, 59.5]	52.0 [46.2, 64.8]	50.5 [48.2, 56.5]	0.769	r=0.086; p=0.691
LDL (mg/dL)	92.3 (29.6)	71.1 (20.8)	102.9 (28)	0.016	r=0.085; p=0.715
Triglycerides (mg/dL)	74.5 [66.2, 93.0]	73.0 [44.5, 76.0]	82.0 [67.8, 96.8]	0.135	r=-0.016; p=0.940
Matsuda Index	4.24 [2.58, 6.00]	5.35 [4.03, 7.48]	3.06 [2.46, 4.78]	0.082	r=-0.616; p=0.001
HOMA-IR	3.1 [2.2, 4.3]	2.5 [1.9, 3.5]	3.7 [2.9, 4.6]	0.207	r=0.398; p=0.054
AdipIR (OGTT)	7.7 [4.4, 12.2]	4.2 [3.0, 10.1]	8.9 [6.7, 30.4]	0.035	r=0.481; p=0.017
AdipIR (fasting)	5.2 [3.0, 6.9]	3.5 [2.4, 5.9]	5.3 [4.1, 9.3]	0.167	r=0.443; p=0.030
Adiponectin (ng/mL)	4613 [3955, 6013]	4613 [4048, 6874]	4540 [3307, 5426]	0.467	r=-0.151; p=0.470
Leptin (pg/mL)	118.1 [53.8, 148.5]	89.8 [36.9, 124.1]	134.0 [84.3, 198.2]	0.085	r=0.817; p<0.001
CRP (mg/L)	0.48 [0.26, 1.2]	0.48 [0.23, 1.1]	0.49 [0.29, 1.1]	0.727	r=0.091; p=0.664

Data are mean ± SD or median [IQR], depending on normality. VAT volume was adjusted with a cube root transformation. Insulin and NEFA concentrations were log transformed.

Substrate Response to OGTT

Blood [glucose], serum [insulin], and plasma [lactate] significantly increased while

plasma [NEFAs] significantly decreased following consumption of 100g glucose beverage

(**Figure 1A-D**). Blood [glucose] (**Figure 1A**) was elevated above fasting levels (99.7 ± 8.5

mg/dL) at 30 (139.9 ± 21.0; p<0.001), 60 (124.9 ± 29.6; p<0.001), and 90 (115.3 ± 25.5; p =

0.007) minutes but not 120 minutes (109 ± 21.7; p = 0.214). [Glucose] area under the curve was

not significantly different between individuals with obesity and individuals with normal weight

BMI (15,076 ± 1912 mg/dl*min vs. 13,852 ± 2126 mg/d*minl; p = 0.13) Compared to fasting

[insulin] (14.2 ± 8.1 µIU/mL), postprandial [insulin] was elevated at 30 (79.9 ± 75.1 µIU/mL; p <

0.001), 60 (77.9 ± 54.2 µIU/mL; p < 0.001), 90 (63.2 ± 47.7 µIU/mL; p = 0.006), and 120 (55.0 ±

36.4 µIU/mL; p = 0.03) minutes (all p<0.001; **Figure 1B**). [Insulin] area under the curve was not

significantly different between individuals with obesity and individuals with normal weight BMI

(9454 ± 5295 µIU/mL*min vs. 5929 ± 4284 µIU/mL*min; p = 0.09) Plasma [lactate] (**Figure 1C**)

did not change from fasting (1.6 ± 0.4 mM) to 30 (1.5 ± 0.4 mM; p = 0.79), but significantly

increased at 60 (1.9 ± 0.3 mM; P < 0.001), and 90 (1.8 ± 0.3 mM; p < 0.001) before returning to

near baseline at 120 (1.7 ± 0.3; p = 0.06) minutes. [Lactate] area under the curve was not

significantly different between individuals with obesity and individuals with normal weight BMI

(205.7 ± 37.3 mM*minl vs. 208.9 ± 24.7 mM*min; p = 0.88) Plasma [NEFAs] (**Figure 1D**) were

suppressed compared to fasting (0.41 ± 0.19 mEq/L) at 30 (0.28 ± 0.13 mEq/L), 60 (0.15 ± 0.08

mEq/L), 90 (0.15 ± 0.08 mEq/L), and 120 (0.09 ± 0.04 mEq/L) minutes (all p < 0.001). [NEFA]

area under the curve was significantly higher for individuals with obesity than individuals with

normal weight BMI (27.5 ± 10.4 mEq/L*min vs. 19.1 ± 7.2 mEq/L*min; p = 0.03).

[Insert Figure 1 here]

CHO (**Figure 2A**) was significantly elevated over time (time: p < 0.001), but there was no

significant interaction between time and obesity status (time*BMI: p = 0.14). CHO was not

different from fasting (0.102 ± 0.071 g/min) at 30 minutes (0.125 ± 0.071; p = 0.62) but was

significantly higher at 60 (0.157 ± 0.063; p = 0.009), 90 (0.160 ± 0.049; p = 0.004), and 120

(0.166 ± 0.061; p = 0.002) minutes. There was a significant interaction between time and

obesity status (time*BMI: p = 0.03) for FOX which indicated higher FOX throughout the

postprandial period for individuals with OW/OB. Overall, FOX was unchanged from fasting

(0.084 ± 0.024) at 30 (0.086 ± 0.024; p = 0.99) and 60 (0.078 ± 0.027; p = 0.83) minutes.

Compared to fasting, FOX was suppressed at 90 (0.073 ± 0.026; p = 0.011) and 120 minutes

(0.068 ± 0.023; p<0.001). RER (**Figure 2C**) was not different from fasting (0.795 ± 0.055) at 30

minutes (0.806 ± 0.052; p = 0.493) but was significantly higher at 60 (0.832 ± 0.051; p<0.001),

90 (0.839 ± 0.044; p<0.001), and 120 (0.845 ± 0.048; p<0.001) minutes. There was no

significant time*BMI interaction (p = 0.08).

[Insert Figure 2 here]

Effect of body composition on postprandial metabolic flexibility (RER, CHO, and FOX)

We used mixed effects models to test for an effect of BF% on postprandial metabolic

flexibility (**Table 4**). In these models, a quadratic model with a main effect for time^2 and a time^2 x

BF% interaction term was a better fit to the postprandial data for RER (p = 0.002) and CHO (p =

0.002) than the linear model. In the quadratic models, BF% modified the relationship between

postprandial RER and time (p = 0.011) and time^2 (p = 0.019), as well as postprandial CHO and

time (p = 0.014) and time^2 (p = 0.023), indicating a nonlinear change in RER and CHO

throughout the OGTT that depends on body composition. There was no evidence that BF%

modifies the relationship between FOX and time (**Supplementary Table 2**). We also considered

the effect of fat mass on postprandial RER and CHO. However, there was no evidence that fat

mass modifies the relationship between postprandial RER and time (p = 0.364) or time^2 (p =

0.409), as well as postprandial CHO and time (p = 0.364) or time^2 (p = 0.409; see

Supplementary Table 3).

Table 3. Effect of body composition on OGTT response

Characteristic[1]	Glucose			Insulin			NEFA		
	Beta[2]	95% CI	p-value	Beta	95% CI	p-value	Beta	95% CI	p-valu
Time	-0.55	-0.723, -0.382	**<0.001**	-0.17	-0.286, -0.046	**0.008**	-2.13	-2.83, -1.44	**<0.00**
Time^2	---	---	---	---	---	---	0.26	0.123, 0.397	**<0.00**
BF%	0.802	0.155, 1.45	0.055	0.232	-0.175, 0.638	0.355	0.39	-0.115, 0.894	0.21(
Fasting Measure	0.231	-0.061, 0.523	0.206	-0.24	-0.728, 0.239	0.412	0.211	-0.067, 0.488	0.22₄
Sex (Male)	0.585	-0.676, 1.84	0.451	0.058	-0.681, 0.794	0.898	1.33	0.360, 2.31	**0.03(**
Age	-0.1	-0.464, 0.267	0.66	0.016	-0.190, 0.224	0.897	0.198	-0.075, 0.472	0.24₄
Matsuda	0.188	-0.247, 0.623	0.482	-0.6	-1.17, -0.018	0.105	0.08	-0.242, 0.403	0.68₃
AdipIR (OGTT)	0.137	-0.280, 0.554	0.591	0.438	0.205, 0.671	**0.006**	0.797	0.452, 1.14	**0.00·**
VAT volume$^{1/3}$	-0.01	-0.624, 0.595	0.969	0.056	-0.285, 0.396	0.789	-0.47	-0.934, 0.001	0.11(

[1]Fasting Measure denotes the fasting concentration of the outcome variable.
[2]A one-unit increase in time represents 30 minutes. A 1-SD increase in each variable corresponds to a 1-SD increase in glucose (19.1), insulin (0.59), or NEFA (0.45). Insulin and NEFA were transformed via natural log for normality. Variable SDs: BF%: 8.7; Fasting glucose: 8.5; Fasting insulin: 0.7; Fasting NEFA: 0.5; Age: 4.97; Matsuda: 2.9; AdipIR (OGTT): 13.3; VAT: 1.29.

To assess the interaction of time and BF% and their effects on postprandial substrate

oxidation as a measure of postprandial metabolic flexibility, we visualized the effect of time and

BF% as continuous variables. We display (**Figure 3A-B**) the modeled RER and CHO for three

representative values of BF%: the mean BF% of our study population, the mean BF% + 1 SD,

and the mean BF% - 1 SD. The interaction of time and BF% displayed in these plots indicates

that higher BF% is associated with higher postprandial RER (**Figure 3A**) and CHO (**Figure 3B**)

demonstrating that higher BF% is associated with greater postprandial metabolic flexibility in our

sample of young adults. The resulting visualization is adjusted for all covariates in **Table 4**.

Further, greater insulin sensitivity (Matsuda Index) was associated with a higher

postprandial RER or CHO (**Table 4**). Because skeletal muscle is a primary site of postprandial

glucose disposal[17] and people with obesity often have higher fat-free mass,[18] we tested the

effect of adding fat-free mass as an additional covariate. This did not improve the model fit (p =

0.204) over the original CHO model. Like the original CHO model, the addition of a $time^2$ main

effect and a $time^2$ x BF% interaction term improved the model fit (p = 0.002), and percent body

fat modified the relationship between CHO and time (p = 0.014) and $time^2$ (p = 0.023) (**Table 4;**

Supplementary Results). Like RER and CHO, higher BF% was associated with greater CHO

adjusted for FFM, indicating differences in FFM across our sample do not affect the interaction

between time and BF%.

Table 4. Effect of body composition (body fat percentage) on postprandial metabolic flexibility

Characteristic[1]	RER			CHO			CHO with FFM		
	Beta[2]	95% CI	p-value	Beta	95% CI	p-value	Beta	95% CI	p-value
BF% x Time[2]	-0.133	-0.240, -0.026	**0.019**	-0.134	-0.245, -0.023	**0.023**	-0.134	-0.245, -0.023	**0.023**
BF% x Time	0.732	0.189, 1.28	**0.011**	0.74	0.175, 1.31	**0.014**	0.74	0.175, 1.31	**0.014**
BF%	-0.346	-1.07, 0.374	0.39	-0.51	-1.26, 0.239	0.225	-0.311	-1.11, 0.492	0.502
Time[2]	-0.156	-0.263, -0.050	**0.006**	-0.166	-0.277, -0.056	**0.005**	-0.166	-0.277, -0.056	**0.005**
Time	1.08	0.537, 1.61	**<0.001**	1.06	0.498, 1.62	**<0.001**	1.06	0.498, 1.62	**<0.001**
Fasting Measure	0.803	0.549, 1.06	**<0.001**	0.793	0.474, 1.11	**<0.001**	0.7	0.358, 1.04	**0.005**
Sex (Male)	0.556	-0.310, 1.42	0.3	0.421	-0.487, 1.33	0.451	0.458	-0.422, 1.34	0.414
Age	-0.095	-0.345, 0.155	0.536	-0.07	-0.332, 0.192	0.661	-0.139	-0.414, 0.137	0.429
Matsuda Index	0.505	0.207, 0.802	**0.012**	0.558	0.246, 0.869	**0.009**	0.59	0.285, 0.895	**0.007**
AdipIR (OGTT)	0.201	-0.083, 0.486	0.255	0.123	-0.176, 0.421	0.504	0.069	-0.232, 0.370	0.715
VAT volume[1/3]	-0.504	-0.909, -0.098	0.055	-0.224	-0.648, 0.199	0.39	-0.368	-0.837, 0.100	0.224
FFM	---	---	---	---	---	---	0.305	-0.177, 0.786	0.323

[1]Fasting Measure denotes the fasting measurement of the outcome variable. See main text for other abbreviations.

[2]A one-unit increase in time represents 30 minutes. A 1-SD increase in each covariate corresponds to a 1-SD increase in postprandial RER (0.044) or CHO (0.055 g/min). Covariate SDs: Fasting RER: 0.055; Fasting CHO: 0.071; FFM: 10.9; See Table 3 for other relevant SDs.

[Insert Figure 3 here]

DISCUSSION

Metabolic flexibility is the ability to adjust substrate oxidation based on substrate

availability or demand.[1] The relationship between adiposity and measures of metabolic flexibility

has been poorly analyzed and lacks clarity.[3, 19-24] The majority of studies to date have

categorized individuals based on BMI (an indicator of adiposity) and made inconsistent and

inconclusive associations between adiposity and measures of metabolic flexibility.[3] In this

analysis, we analyzed the interaction between body fat percentage and postprandial metabolism

(RER, CHO, and FOX) and demonstrate that higher body fat percentage is associated with

elevated postprandial metabolic flexibility (RER and CHO) during an OGTT in young adults

without overt metabolic pathology. These findings may seem paradoxical since individuals with

higher BMI (e.g., obesity) are most often reported to have lower, or blunted, metabolic flexibility.

However, this finding is not inconsistent with studies that show females have greater metabolic

flexibility than males despite higher body fat[5] and individuals with high metabolic flexibility had

greater subcutaneous adipose tissue volumes.[7] Furthermore, loss of subcutaneous adipose

tissue during prolonged bed rest is accompanied by decreased metabolic flexibility.[20] This study

expands on previously published reports by directly analyzing the relationship between body fat

and postprandial metabolic flexibility while adjusting for known covariates (i.e., age, sex, VAT,

insulin sensitivity, etc.) that significantly affect substrate oxidation.

Adipose tissue function is proposed as a central regulator of metabolic flexibility[2] by

buffering fatty acid flux during the postprandial period.[4] Obesity is associated with increased

adipose tissue volume,[25] elevated adipose tissue insulin resistance, adipose tissue

inflammation, and altered adipokine release;[26] all of which are associated with altered

postprandial metabolism.[27] In our sample of healthy young adults, higher body fat percentages

did not alter suppression of lipolysis, as measured by plasma NEFAs, following ingestion of the

glucose beverage. Furthermore, measures of adipose tissue insulin resistance (ADIP-IR) did not

have a significant effect on postprandial RER or CHO. However, we cannot fully rule out

reduced adipose tissue function in our findings. A critical role of adipose tissue in the

postprandial state is glucose uptake and lipogenesis, which we cannot account for here.[28-30]

Importantly, our approach demonstrates that an increase in body fat percentage of ~6.9% (i.e.,

20.0% to 26.9%) would result in an average increase of approximately 19.1 mg/dl of blood

glucose at each time point during the OGTT (**Table 3**). This suggests the capacity for adipose

tissue to buffer the blood glucose flux over time may be reduced in people with a higher body fat

percentage, which would be consistent with observational studies in individuals with obesity,

diabetes, and animal models.[31-34] This function is an important part of adipose tissue's role in

reducing hyperglycemia in the postprandial period while subsequently storing nutrients for later

oxidation.

In the context of greater hyperglycemia following the OGTT, the elevated postprandial

metabolic flexibility (postprandial RER and CHO) associated with higher body fat percentage in

our sample would not be paradoxical. The postprandial period should induce an anabolic state

(e.g. storage of nutrients) as compared to the catabolic state of fasting (e.g. breakdown of

substrate).[4] Metabolic flexibility is a critical metabolic adaptation to handle the complex

integration of metabolism during caloric excess and restriction.[35] Furthermore, it is essential for

regular maintenance of blood glucose homeostasis following a feeding. Our observation of

elevated CHO and RER in the setting of greater hyperglycemia suggests greater metabolic

flexibility in people with higher body fat percentages, which would fit the concept of metabolic

flexibility precisely; these individuals with higher body fat percentages are adjusting substrate

oxidation based on the increased availability of glucose in the blood for longer periods of time.

Individuals with lower body fat percentages who do not elevate CHO and RER during this period

are likely experiencing the appropriate catabolic (storage) response to feeding by taking up and

storing glucose as glycogen (liver and skeletal muscle) or fat (adipose tissue). As a result, they

have no further need to increase CHO to maintain blood glucose homeostasis. A recent report

by Curl et al.[36] demonstrated altered glucose kinetics in older individuals as compared to

younger adults without differences in metabolic flexibility, which they defined as the difference

between peak and fasting RER. We did not assess glucose kinetics, but our findings on the

impacts of body composition on postprandial blood glucose would be consistent with their

findings.

This raises the question if our sample of young adults across the BMI spectrum but with

no overt metabolic pathology are demonstrating a resilient, metabolically flexible phenotype.[37]

Metabolic flexibility is diminished following hypercaloric, high-fat feeding interventions and

improved following significant weight loss, indicating it is a plastic phenotype.[38, 39] No study has

successfully tracked the disappearance of metabolic flexibility, but we suspect that long-term

elevated levels of adiposity, such as that seen in chronic obesity, eventually lead to the loss of

postprandial metabolic flexibility. In our sample, higher body fat percentage is inversely

associated with insulin sensitivity (r =-0.616; p=0.001) while being positively associated with

fasting values for serum insulin (r=0.427; p=0.037), plasma leptin (r=0.817; p<0.001), and

adipose tissue insulin resistance (r=0.443; p=0.030). These relationships may be indicative or

worsening metabolic health with higher body fat percentage, but the ability to elevate CHO

oxidation during the OGTT (e.g. metabolic flexibility) demonstrates a resilient, metabolically

flexibility phenotype in these individuals. The eventual loss of metabolic flexibility would

therefore play a significant role in the progression of glucose dysfunction towards overt

metabolic disease. That process and the mechanisms involved are still poorly understood and

require both experimental and observational longitudinal studies.

Studies assessing the relationship between metabolic flexibility and adiposity primarily

note inconsistent and inconclusive results which include positive associations, inverse

associations, and no associations between indices of metabolic flexibility and measures of

adiposity (e.g. total fat mas, visceral adipose volume).[3] Some studies also rely on surrogates of

adiposity such as BMI, waist circumference, or waist:hip ratio.[3, 22] Though not the primary goal of

our study, we note no significant differences in postprandial RER or CHO between individuals

with normal weight and individuals with obesity as determined by the BMI; this is consistent with

many studies. Directly analyzing the effect of body composition, while adjusting for covariates

known to influence substrate oxidation, has allowed us to more clearly demonstrate an

unexpected, but explainable, relationship in which higher body fat percentage is associated with

postprandial metabolic flexibility in young adults without overt metabolic pathology. Assessing

the relationship between body composition, instead of absolute fat mass or volume, and

metabolic flexibility also accounts for the expected body mass differences between subjects

and, more specifically, between biological sexes. By recruiting similarly from both sexes and

including biological sex in our models, we account for potential sex differences independent of

differences in body composition between the sexes. While our study is one of the larger

studiesy analyzing metabolic flexibility to an OGTT in both males and females to date,[3] we did

not power our study to detect sex differences, which should be considered in future work.

Recent reports demonstrate that total body mass as well as fat free mass are inversely

associated with metabolic flexibility during an OGTT, highlighting the necessity of taking

anthropometric differences into account.[40]

Furthermore, our approach to analyzing metabolic flexibility during an OGTT as a

continuous variable across time is a novel aspect of our approach. Studies assessing metabolic

flexibility during feeding challenges define metabolic flexibility as the change in RER or CHO

from fasting to a defined timepoint (e.g. 60 min in an OGTT) or the peak change (max – min) in

the measure.[3] Our data, when analyzed in that way, produces similar results (**Supplementary**

Figure 1). Defining postprandial metabolic flexibility based on a single timepoint during a

dynamic and prolonged response, such as feeding, is likely a carryover from the EHC method,

which applies indirect calorimetry to the steady-state phase of insulin infusion. Because

postprandial metabolism is dynamic, a feeding challenge such as an OGTT has no equivalent to

the steady-state period.[40, 41] Curl et al.[36] recently argued against postprandial RER as measure

of metabolic flexibility following a glucose challenge when comparing younger vs. older adults,

instead suggesting blood glucose kinetics as a better measure. We agree that the postprandial

period should not be defined as a single measure and should be analyzed in a continuous

manner across time. In fact, Curl et al.[36] note significantly higher RER in their younger adults at

60-min postprandial but not at 120-min postprandial, highlighting the dynamic nature of this

physiological process and the necessity to analyze it continuously.

We demonstrate that higher body fat percentage is associated with greater postprandial

metabolic flexibility (RER and CHO) during an oral glucose tolerance test in young adults

without overt metabolic pathology. We believe this finding is indicative of a preserved

metabolically flexible phenotype where elevated postprandial blood glucose is resulting in

elevated CHO with higher body fat percentage. Our approach of analyzing data in a continuous

manner using linear mixed models, accounting for covariates known to modify the postprandial

substrate oxidation response, adds clarity to the previously poorly defined relationship between

adiposity and postprandial metabolic flexibility. While we cannot account for some functional

(e.g. adipocyte size, lipogenesis, and molecular) or anatomical (e.g. distribution) aspects of

adiposity, we more directly demonstrate what has long been ambiguous in published reports.

These findings raise several new questions about adiposity and the continuum of metabolic

health.

Conflicts of Interest: The authors declare no conflicts of interest.

Acknowledgements: The authors would like to acknowledge the Milken Institute School of

Public Health Office of Research Excellence which provided funding for this project. The authors

would also like to acknowledge the participants for their volunteered time and effort.

Funding: This study was funded by a Research Innovation Award (PI Barberio) from the Office

of Research Excellence in the Milken Institute School of Public Health at The George

Washington University.

Contributions: **NF:** Conceptualization, methodology, formal analysis, investigation, data
curation, writing – original draft, visualization. **SR:** Investigation. **JB**: Investigation.
LS: Investigation. **AL:** Investigation. **LD:** Conceptualization, writing – review & editing. **AC:**
Formal analysis, writing – review & editing. **MB:** Conceptualization, methodology, investigation,
writing – original draft, project administration, funding acquisition.

All authors have reviewed this manuscript and agree to its publication as written.

References

1.   Goodpaster BH, Sparks LM. Metabolic Flexibility in Health and Disease. Cell Metab.

2017;25(5):1027-36. doi: 10.1016/j.cmet.2017.04.015. PubMed PMID: 28467922; PMCID:

PMC5513193.

2.   Storlien L, Oakes ND, Kelley DE. Metabolic flexibility. Proc Nutr Soc. 2004;63(2):363-8.

doi: 10.1079/PNS2004349. PubMed PMID: 15294056.

3.   Glaves A, Diaz-Castro F, Farias J, Ramirez-Romero R, Galgani JE, Fernandez-Verdejo

R. Association Between Adipose Tissue Characteristics and Metabolic Flexibility in Humans: A

Systematic Review. Front Nutr. 2021;8:744187. Epub 20211203. doi:

10.3389/fnut.2021.744187. PubMed PMID: 34926544; PMCID: PMC8678067.

4.   Frayn KN. Adipose tissue as a buffer for daily lipid flux. Diabetologia. 2002;45(9):1201-

10. Epub 20020724. doi: 10.1007/s00125-002-0873-y. PubMed PMID: 12242452.

5.   Sparks LM, Pasarica M, Sereda O, deJonge L, Thomas S, Loggins H, Xie H, Miles JM,

Smith SR. Effect of adipose tissue on the sexual dimorphism in metabolic flexibility. Metabolism.

2009;58(11):1564-71. doi: 10.1016/j.metabol.2009.05.008. PubMed PMID: 19595383.

6.   Sparks LM, Ukropcova B, Smith J, Pasarica M, Hymel D, Xie H, Bray GA, Miles JM,

Smith SR. Relation of adipose tissue to metabolic flexibility. Diabetes Res Clin Pract.

2009;83(1):32-43. Epub 20081126. doi: 10.1016/j.diabres.2008.09.052. PubMed PMID:

19038471; PMCID: PMC2749984.

7.   Fernandez-Verdejo R, Malo-Vintimilla L, Gutierrez-Pino J, Lopez-Fuenzalida A, Olmos

P, Irarrazaval P, Galgani JE. Similar Metabolic Health in Overweight/Obese Individuals With

Contrasting Metabolic Flexibility to an Oral Glucose Tolerance Test. Front Nutr. 2021;8:745907.

Epub 20211116. doi: 10.3389/fnut.2021.745907. PubMed PMID: 34869522; PMCID:

PMC8637191.

8.   Taylor HL, Buskirk E, Henschel A. Maximal oxygen intake as an objective measure of

cardio-respiratory performance. J Appl Physiol. 1955;8(1):73-80. doi: 10.1152/jappl.1955.8.1.73.

PubMed PMID: 13242493.

9.   Frayn KN. Calculation of substrate oxidation rates in vivo from gaseous exchange. J

Appl Physiol Respir Environ Exerc Physiol. 1983;55(2):628-34. doi:

10.1152/jappl.1983.55.2.628. PubMed PMID: 6618956.

10.   Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose

tolerance testing: comparison with the euglycemic insulin clamp. Diabetes Care.

1999;22(9):1462-70. doi: 10.2337/diacare.22.9.1462. PubMed PMID: 10480510.

11.   Gastaldelli A, Gaggini M, DeFronzo RA. Role of Adipose Tissue Insulin Resistance in

the Natural History of Type 2 Diabetes: Results From the San Antonio Metabolism Study.

Diabetes. 2017;66(4):815-22. Epub 20170104. doi: 10.2337/db16-1167. PubMed PMID:

28052966.

12.   Shashaj B, Luciano R, Contoli B, Morino GS, Spreghini MR, Rustico C, Sforza RW,

Dallapiccola B, Manco M. Reference ranges of HOMA-IR in normal-weight and obese young

Caucasians. Acta Diabetol. 2016;53(2):251-60. Epub 20150613. doi: 10.1007/s00592-015-

0782-4. PubMed PMID: 26070771.

13.   Kuznetsova A, Brockhoff, P.B., & Christensen R.H.B. lmerTest Package: Tests in Linear

Mixed Effects Models. Journal of Statistical Software. 2017;82(13):1-26. doi:

10.18637/jss.v082.i13.

14.   Lüdecke D, Ben-Shachar, M.S., Patil, I., Waggoner, P., Makowski, D. performance: An R

Package for Assessment, Comparison

and Testing of Statistical Models. The Journal of Open Source Software. 2021;6(60). doi:

10.21105/joss.03139.

15.   Lenth R. emmeans: Estimated Margin2l Means, aka Least-Squares Means. R package

version 1.10.3-090006. 2024.

16.   Baayen RH, Davidson DJ, Bates DM. Mixed-effects modeling with crossed random

effects for subjects and items. J Mem Lang. 2008;59(4):390-412. doi:

10.1016/j.jml.2007.12.005. PubMed PMID: WOS:000261651700003.

17.   Kelley D, Mitrakou A, Marsh H, Schwenk F, Benn J, Sonnenberg G, Arcangeli M, Aoki T,

Sorensen J, Berger M, et al. Skeletal muscle glycolysis, oxidation, and storage of an oral

glucose load. J Clin Invest. 1988;81(5):1563-71. doi: 10.1172/JCI113489. PubMed PMID:

3130396; PMCID: PMC442590.

18.   Dulloo AG, Jacquet J, Miles-Chan JL, Schutz Y. Passive and active roles of fat-free

mass in the control of energy intake and body composition regulation. Eur J Clin Nutr.

2017;71(3):353-7. Epub 20161214. doi: 10.1038/ejcn.2016.256. PubMed PMID: 27966570.

19.   Lightowler H, Schweitzer L, Theis S, Henry CJ. Changes in Weight and Substrate

Oxidation in Overweight Adults Following Isomaltulose Intake During a 12-Week Weight Loss

Intervention: A Randomized, Double-Blind, Controlled Trial. Nutrients. 2019;11(10). Epub

20191004. doi: 10.3390/nu11102367. PubMed PMID: 31590285; PMCID: PMC6836138.

20.   Rudwill F, O'Gorman D, Lefai E, Chery I, Zahariev A, Normand S, Pagano AF, Chopard

A, Damiot A, Laurens C, Hodson L, Canet-Soulas E, Heer M, Meuthen PF, Buehlmeier J,

Baecker N, Meiller L, Gauquelin-Koch G, Blanc S, Simon C, Bergouignan A. Metabolic

Inflexibility Is an Early Marker of Bed-Rest-Induced Glucose Intolerance Even When Fat Mass Is

Stable. J Clin Endocrinol Metab. 2018;103(5):1910-20. doi: 10.1210/jc.2017-02267. PubMed

PMID: 29546280; PMCID: PMC7263792.

21.   Kahlhofer J, Lagerpusch M, Enderle J, Eggeling B, Braun W, Pape D, Muller MJ, Bosy-

Westphal A. Carbohydrate intake and glycemic index affect substrate oxidation during a

controlled weight cycle in healthy men. Eur J Clin Nutr. 2014;68(9):1060-6. Epub 20140709. doi:

10.1038/ejcn.2014.132. PubMed PMID: 25005676.

22.   Bergouignan A, Antoun E, Momken I, Schoeller DA, Gauquelin-Koch G, Simon C, Blanc

S. Effect of contrasted levels of habitual physical activity on metabolic flexibility. J Appl Physiol

(1985). 2013;114(3):371-9. Epub 20121213. doi: 10.1152/japplphysiol.00458.2012. PubMed

PMID: 23239872.

23.   Assaad M, El Mallah C, Obeid O. Phosphorus ingestion with a high-carbohydrate meal

increased the postprandial energy expenditure of obese and lean individuals. Nutrition.

2019;57:59-62. Epub 20180619. doi: 10.1016/j.nut.2018.05.019. PubMed PMID: 30153580.

24.   Huda MS, Dovey T, Wong SP, English PJ, Halford J, McCulloch P, Cleator J, Martin B,

Cashen J, Hayden K, Wilding JP, Pinkney J. Ghrelin restores 'lean-type' hunger and energy

expenditure profiles in morbidly obese subjects but has no effect on postgastrectomy subjects.

Int J Obes (Lond). 2009;33(3):317-25. Epub 20090203. doi: 10.1038/ijo.2008.270. PubMed

PMID: 19188925.

25.   Turer AT, Khera A, Ayers CR, Turer CB, Grundy SM, Vega GL, Scherer PE. Adipose

tissue mass and location affect circulating adiponectin levels. Diabetologia. 2011;54(10):2515-

24. Epub 20110722. doi: 10.1007/s00125-011-2252-z. PubMed PMID: 21779869; PMCID:

PMC4090928.

26.   Shafiei MS, Shetty S, Scherer PE, Rockey DC. Adiponectin regulation of stellate cell

activation via PPARgamma-dependent and -independent mechanisms. Am J Pathol.

2011;178(6):2690-9. doi: 10.1016/j.ajpath.2011.02.035. PubMed PMID: 21641391; PMCID:

PMC3124230.

27.   Sun K, Kusminski CM, Scherer PE. Adipose tissue remodeling and obesity. J Clin

Invest. 2011;121(6):2094-101. Epub 20110601. doi: 10.1172/JCI45887. PubMed PMID:

21633177; PMCID: PMC3104761.

28.   Chadt A, Al-Hasani H. Glucose transporters in adipose tissue, liver, and skeletal muscle

in metabolic health and disease. Pflugers Arch. 2020;472(9):1273-98. Epub 20200626. doi:

10.1007/s00424-020-02417-x. PubMed PMID: 32591906; PMCID: PMC7462924.

29.   Luo L, Liu M. Adipose tissue in control of metabolism. J Endocrinol. 2016;231(3):R77-

R99. doi: 10.1530/JOE-16-0211. PubMed PMID: 27935822; PMCID: PMC7928204.

30.   Roberts R, Hodson L, Dennis AL, Neville MJ, Humphreys SM, Harnden KE, Micklem KJ,

Frayn KN. Markers of de novo lipogenesis in adipose tissue: associations with small adipocytes

and insulin sensitivity in humans. Diabetologia. 2009;52(5):882-90. Epub 20090228. doi:

10.1007/s00125-009-1300-4. PubMed PMID: 19252892.

31.   Fabbrini E, Yoshino J, Yoshino M, Magkos F, Tiemann Luecking C, Samovski D,

Fraterrigo G, Okunade AL, Patterson BW, Klein S. Metabolically normal obese people are

protected from adverse effects following weight gain. J Clin Invest. 2015;125(2):787-95. Epub

20150102. doi: 10.1172/JCI78425. PubMed PMID: 25555214; PMCID: PMC4319438.

32.   Graham TE, Kahn BB. Tissue-specific alterations of glucose transport and molecular

mechanisms of intertissue communication in obesity and type 2 diabetes. Horm Metab Res.

2007;39(10):717-21. doi: 10.1055/s-2007-985879. PubMed PMID: 17952832.

33.   Herman MA, Peroni OD, Villoria J, Schon MR, Abumrad NA, Bluher M, Klein S, Kahn

BB. A novel ChREBP isoform in adipose tissue regulates systemic glucose metabolism. Nature.

2012;484(7394):333-8. doi: 10.1038/nature10986. PubMed PMID: 22466288; PMCID:

PMC3341994.

34.   Hoffstedt J, Forster D, Lofgren P. Impaired subcutaneous adipocyte lipogenesis is

associated with systemic insulin resistance and increased apolipoprotein B/AI ratio in men and

women. J Intern Med. 2007;262(1):131-9. doi: 10.1111/j.1365-2796.2007.01811.x. PubMed

PMID: 17598821.

35.   Smith RL, Soeters MR, Wust RCI, Houtkooper RH. Metabolic Flexibility as an Adaptation

to Energy Resources and Requirements in Health and Disease. Endocr Rev. 2018;39(4):489-

517. doi: 10.1210/er.2017-00211. PubMed PMID: 29697773; PMCID: PMC6093334.

36.   Curl CC, Leija RG, Arevalo JA, Osmond AD, Duong JJ, Huie MJ, Masharani U, Horning

MA, Brooks GA. Altered glucose kinetics occurs with aging: a new outlook on metabolic

flexibility. Am J Physiol Endocrinol Metab. 2024;327(2):E217-E28. Epub 20240619. doi:

10.1152/ajpendo.00091.2024. PubMed PMID: 38895979; PMCID: PMC11427093.

37.   Mietus-Snyder M, Perak AM, Cheng S, Hayman LL, Haynes N, Meikle PJ, Shah SH,

Suglia SF, American Heart Association Atherosclerosis H, Obesity in the Young Committee of

the Council on Lifelong Congenital Heart D, Heart Health in the Y, Council on L,

Cardiometabolic H, Council on Arteriosclerosis T, Vascular B, Council on C, Stroke N. Next

Generation, Modifiable Cardiometabolic Biomarkers: Mitochondrial Adaptation and Metabolic

Resilience: A Scientific Statement From the American Heart Association. Circulation.

2023;148(22):1827-45. Epub 20231030. doi: 10.1161/CIR.0000000000001185. PubMed PMID:

37902008.

38.   Berk ES, Kovera AJ, Boozer CN, Pi-Sunyer FX, Albu JB. Metabolic inflexibility in

substrate use is present in African-American but not Caucasian healthy, premenopausal,

nondiabetic women. J Clin Endocrinol Metab. 2006;91(10):4099-106. Epub 20060725. doi:

10.1210/jc.2005-2411. PubMed PMID: 16868062; PMCID: PMC2670464.

39.   Kelley DE, Goodpaster B, Wing RR, Simoneau JA. Skeletal muscle fatty acid

metabolism in association with insulin resistance, obesity, and weight loss. Am J Physiol.

1999;277(6):E1130-41. doi: 10.1152/ajpendo.1999.277.6.E1130. PubMed PMID: 10600804.

40.   Alcantara JMA, Galgani JE. Association of metabolic flexibility indexes after an oral

glucose tolerance test with cardiometabolic risk factors. Eur J Clin Nutr. 2024;78(3):180-6. Epub

20231218. doi: 10.1038/s41430-023-01373-w. PubMed PMID: 38110728.

41.   Alcantara JMA, Sanchez-Delgado G, Jurado-Fasoli L, Galgani JE, Labayen I, Ruiz JR.

Reproducibility of the energy metabolism response to an oral glucose tolerance test: influence

of a postcalorimetric correction procedure. Eur J Nutr. 2023;62(1):351-61. Epub 20220825. doi:

10.1007/s00394-022-02986-w. PubMed PMID: 36006468; PMCID: PMC9899729.

Normal Weight (NW) ● Overweight/Obesity (OW/OB)